Population genomics of *Bacillus anthracis* from an anthrax hyperendemic area

reveals transmission processes across spatial scales and unexpected within-host

diversity

Taya L. Forde[1][*], Tristan P. W. Dennis[1], O. Rhoda Aminu[1], William T. Harvey[1], Ayesha

Hassim[2], Ireen Kiwelu[3], Matej Medvecky[1], Deogratius Mshanga[4], Henriette Van Heerden[2],

Adeline Vogel[1], Ruth N. Zadoks[1,#], Blandina T. Mmbaga[3,5], Tiziana Lembo[1][¶], Roman Biek[1][¶]

[1] Institute of Biodiversity, Animal Health & Comparative Medicine, University of Glasgow,

Glasgow, United Kingdom

[2] Department of Veterinary Tropical Diseases, Faculty of Veterinary Science, University of

Pretoria, Onderstepoort, South Africa

[3] Kilimanjaro Clinical Research Institute, Kilimanjaro Christian Medical Centre, Moshi,

Tanzania

[4] Tanzania Veterinary Laboratory Agency, Northern Zone, Arusha, Tanzania

[5] Kilimanjaro Christian Medical University College, Moshi, Tanzania

[#] Current address: Sydney School of Veterinary Science, University of Sydney, Sydney,

Australia

* Corresponding author

E-mail: taya.forde@glasgow.ac.uk (TLF)

¶ These authors contributed equally to this work.

Abstract

Genomic sequencing has revolutionized our understanding of bacterial disease epidemiology, but remains underutilized for zoonotic pathogens in remote endemic settings. Anthrax, caused by the spore-forming bacterium *Bacillus anthracis*, remains a threat to human and animal health and rural livelihoods in low- and middle-income countries. While the global genomic diversity of *B. anthracis* has been well-characterized, there is limited information on how its populations are genetically structured at the scale at which transmission occurs, critical for understanding the pathogen's evolution and transmission dynamics. Using a uniquely rich dataset, we quantified genome-wide single nucleotide polymorphisms (SNPs) among 73 *B. anthracis* isolates derived from 33 livestock carcasses sampled over one year throughout the Ngorongoro Conservation Area, Tanzania, an area hyperendemic for anthrax. Genome-wide SNPs distinguished 22 unique *B. anthracis* genotypes within the study area. However, phylogeographic structure was lacking, as identical SNP profiles were found throughout the study area, likely the result of the long and variable periods of spore dormancy and long-distance livestock movements. Significantly, divergent genotypes were obtained from spatio-temporally linked cases and even individual carcasses. The high number of SNPs distinguishing isolates from the same host is unlikely to have arisen during infection, as supported by our simulation models. This points to an unexpectedly wide transmission bottleneck for *B. anthracis*, with an inoculum comprising multiple variants being the norm. Our work highlights that inferring transmission patterns of *B. anthracis* from genomic data will require analytical approaches that account for extended and variable environmental persistence as well as co-infection.

Importance

Pathogens transmitted between animals and people affect the health and livelihoods of farmers, particularly in developing countries dependent on livestock. Understanding over what distances these pathogens are transmitted and how they evolve is important to inform control strategies towards reducing disease impacts. Information on the circulation of *Bacillus anthracis*, which causes the often-lethal disease anthrax, is lacking for settings where the disease is commonplace. Consequently, we examined its genetic variability in an area in Tanzania where anthrax is widespread. We found no clear link between how closely cases were sampled and their genetic similarity. We suspect this lack of congruence is primarily driven by large-scale livestock movements, which control efforts should take into consideration. Another significant finding was the co-occurrence of multiple *B. anthracis* types within individual hosts, suggesting animals are commonly infected with a mixture of variants. This needs to be accounted for when investigating possible connections between cases.

Introduction

Genomic data have the potential to transform our understanding of the evolution and epidemiology of pathogens of public health importance (1). However, this potential has yet to be fully harnessed for many zoonotic diseases that occur in hard-to-reach areas. Anthrax remains endemic in many low- and middle-income countries (LMICs) worldwide (2). It is a disease characterized by sudden deaths in herbivorous livestock and wildlife, and can also cause serious, potentially fatal disease in people (3). Anthrax is classified among the neglected zoonoses: a group of diseases shared by animals and people that, due to their occurrence in remote, disadvantaged communities, collectively receive less than 0.1% of international global health assistance (4). As for many neglected zoonoses, there is limited genomic data for *Bacillus anthracis*, the bacterium that causes anthrax, from endemic LMIC settings where surveillance tends to be limited. Such data could help to improve our understanding of transmission processes, such as how *B. anthracis* is spread within and between outbreaks, and ultimately contribute to more informed disease management.

The genomic diversity of *B. anthracis* has been well-described at a global scale. Isolates can be broadly divided into three major clades (A, B, C), of which A clade is the most widespread and globally dominant (5–7). Isolates from most *B. anthracis* lineages have been found across geographically widespread areas, often spanning multiple continents (7). While particular variants often predominate regionally, high lineage diversity has also been reported, including co-circulation of strains from multiple linages (8, 9). How *B. anthracis* diversity is structured at smaller scales is less well-defined. The pathogen has limited genomic diversity compared to other bacterial species (i.e. is genetically monomorphic), rendering standard genotyping methods such as multi-locus sequence typing insufficiently discriminatory (10). A hierarchical genotyping

scheme known as PHRANA was therefore developed specifically for *B. anthracis,* based on

quickly evolving repetitive regions nested within more phylogenetically stable markers –

canSNPs – that distinguish among the major lineages (5, 11). Variants of this scheme have been

used to examine the diversity of *B. anthracis* in several endemic settings globally, including in a

few African countries (8, 12–14). Genome-wide SNP data would offer higher resolution for

discriminating among closely-related isolates. However, whole genome sequencing (WGS)

features in only a few studies of local *B. anthracis* diversity (15–18), and has rarely been

conducted outside Europe (19), so the potential for phylogenomic data to be used to understand

transmission patterns within hyperendemic areas has yet to be explored.

Transmission of *B. anthracis* occurs primarily through the environment. After causing the death

of the animal host, vegetative bacteria are released into the environment via bodily fluids. Here,

upon exposure to oxygen and cues related to a lack of nutrients, these bacteria sporulate and can

persist in a dormant yet infectious state for several decades (3, 20). While the viability of spores

decreases over time, new *B. anthracis* infections could theoretically arise from recent cases or

cases that occurred several years or even decades previously. How this environmental

persistence shapes the spatio-temporal diversity of *B. anthracis* in endemic settings has never

been investigated.

In molecular epidemiological studies of bacterial pathogens, a single isolate is typically

sequenced from each individual case. However, this approach fails to recognize the bacterial

population diversity that may exist within the host (21, 22). Such diversity can either result from

mutations that arise during infection, or from heterogeneity (multiple variants) in the inoculum,

either through co-infection (exposure to multiple variants simultaneously) or superinfection

(multiple exposures) (23, 24). Under those scenarios, a single isolate is unlikely to represent the

overall diversity of the pathogen within the host, and as a result this approach can lead to

erroneous inferences about transmission pathways (25). The importance of capturing within-host

diversity is therefore increasingly recognized (26, 27). In the case of anthrax, multiple genotypes

of *B. anthracis* have been previously found within individual hosts (28–30), but it remains

unclear whether this represents a more widespread phenomenon.

The objective of this study was to quantify the genomic diversity of *B. anthracis* at hierarchical

spatial scales within the livestock population of a hyperendemic setting. This was accomplished

using a unique dataset including 1) isolates collected throughout a large (~8,300 km^2) area of

northern Tanzania where anthrax is widespread; 2) among spatio-temporally linked cases; and 3)

within individual hosts, assessing multiple isolates from the same and different sample types

associated with a case (e.g. tissue, blood, soil).

Results

Genomic sequencing of *B. anthracis* isolates enabled assessment of diversity at hierarchical spatial scales

Bacillus anthracis isolates were recovered from livestock carcasses in the Ngorongoro

Conservation Area (NCA), part of the Serengeti ecosystem of northern Tanzania. The 73 isolates

from which WGS data were obtained were from a total of 33 carcass sites sampled throughout

the NCA (Table S1). Carcasses were of the following species: sheep (n = 18), cattle (n = 7),

goats (n = 4), donkey (n = 1) or unknown host (n = 3). The diversity of *B. anthracis* was assessed

at multiple hierarchical spatial scales (Fig. 1). Carcasses for which detailed sampling location

data were available (n = 32) could be grouped into four distinct areas within the NCA (central,

north, south, east), referred to herein as 'geographical groups'. To assess the genomic relatedness

among spatio-temporally-linked cases, on four occasions, samples were collected from two

carcasses either from the same or neighboring households on the same or consecutive days,

which we refer to as 'epidemiological clusters'. Three of these clusters were in the central

geographical group, while the fourth was in the southern group of carcasses sampled. Finally,

multiple isolates (n = 2-4) were sequenced from a single carcass for 21 carcasses. These were

either i) isolates from multiple sample types (i.e. tissue, blood, swabs, soil and/or insects) from a

single carcass (n = 16 carcasses) and/or ii) multiple isolates from the same sample (n = 15

carcasses). Isolates contributing to investigations of diversity at each hierarchical level are

detailed in Table S1, along with individual sequence quality metrics. Mapped reads had an

average depth of coverage of 85X across isolates, ranging from 24X – 245X (median 72X).

B. anthracis within the NCA is limited to a single subgroup lacking clear

phylogeographic signal

All _B. anthracis_ isolates from the NCA were found to belong to the Ancient A subgroup of Clade

A (Fig. 2). Within this subgroup, the NCA isolates formed a monophyletic clade within Cluster

3.2 as defined by Bruce et al. (7), which also contained the isolate A2075 (GenBank accession

SRR2968187), isolated in 1999 from a baboon in Muhesi Game Reserve, central Tanzania.

A total of 125 SNPs polymorphic among the NCA isolates and A2075 were retained for analysis,

of which 13 were unique to A2075. Twenty-two unique genotypes (SNP profiles) were found

among the 73 NCA isolates (Fig. 3). Based on a rarefaction analysis, the observed genotypic

diversity was close to that present throughout the study area (i.e. further sampling would have

been unlikely to reveal additional genotypes; Fig. S1). The maximum pairwise nucleotide

difference between any two NCA-derived isolates was 49 SNPs [median = 24, interquartile range

(IQR) = 10-35] (Fig. S2). There was no clear relationship between the pairwise nucleotide

differences and the geographical distance between sampling locations (Fig. S3), as confirmed by

a test for isolation by distance (r = 0.04, p = 0.214). Isolates from the central, eastern and

southern geographical groups were observed throughout the phylogenetic tree, while all isolates

from the northern sampling area were restricted to a single clade that contained the majority of

NCA isolates (Fig. 3). In eight instances, identical *B. anthracis* genotypes were found in

carcasses from different geographical groups, all of which involved isolates from the central

geographical group and one of the other areas, with all areas implicated (Table S3). Identical or

nearly identical SNP profiles (1 SNP difference) were obtained from carcasses sampled 3-5

months apart on six occasions, and 10 months apart on one occasion (Table S4).

B. anthracis isolates from spatio-temporally linked cases are rarely genetically

related

Between one and four isolates were sequenced from each carcass sampled as part of an

epidemiological cluster (i.e. pair of spatio-temporally linked carcasses), resulting in between 4-8

isolates per cluster. In most cases, isolates deriving from the same epidemiological cluster were

phylogenetically unrelated. Only in one of the four clusters examined (C2) did both carcasses

contain isolates with identical core SNP profiles (Fig. 3B, S6 and S7). Overall, isolates from

different carcasses within the same epidemiological cluster had similar numbers of SNP

differences when compared with randomly selected carcasses (linked cases: median = 21, IQR =

0-33; unlinked cases: median = 23, IQR = 10-35) (Fig. 4).

Within-host diversity of _B. anthracis_ is similar to between-host diversity in the

NCA

Overall, the number of pairwise SNP differences between isolates from the same carcass was

lower than that found between isolates from different carcasses [same: median = 11, interquartile

range = 0-34; different: median = 23, interquartile range = 10-35] (Fig. 4). However, isolates

with multiple distinct genotypes were obtained on 15 of 21 occasions wherein multiple isolates

were sequenced from the same carcass, with as many as 43 SNP differences between isolates

(Fig. 5). A high level of divergence was seen, regardless of whether isolates were from the same

or different sample type (e.g. multiple isolates from a tissue sample vs. isolates from tissue and

soil samples, Fig. 5A). Divergent genotypes were observed in all carcasses from which three or

more isolates were sequenced (12/12), and in a third (3/9) of carcasses for which two isolates

were sequenced. A single SNP difference separated one isolate from two others within one

carcass (AN16-83) and two SNP differences separated two isolates from a single soil sample

(LNA). All other pairwise within-host SNP differences were eight or above. All of the within-

host SNP profiles detected were shared with those from other carcasses.

Observed within-host diversity is unlikely to have arisen during the course of

infection

To assess the likelihood of different levels of within-host diversity arising during the course of

infection, we performed simulation modelling of *B. anthracis* infection with a homogenous

inoculum of varying size (1, 2, 5, 10, 20, 50 and 100 bacterial genomes), and across a range of

replication cycles (up to 25 generations) using previously established mutation rates (29). With

an infectious dose d, discrete generations and no die-off, the population size in generation i is

therefore $d \times 2^{(i-1)}$. Averaging across simulations with varying initial doses, in the 25[th]

generation, 90.0% of the population was identical to the infecting dose, 9.53% differed by 1

SNP, and 0.489% by 2 SNPs, with higher numbers of mutations very rare (<0.01%). While

stochasticity was present in early generations and particularly at low inoculum doses (Fig. S8-

10), the mean proportion of genomes with various numbers of SNPs changed predictably after

the initial generations (Fig. 6A-B). Extrapolating to the 40th generation, then ~84% of the

population would be expected to be identical to the infecting dose, ~15% differing by 1 SNP and

~0.9% by 2 SNPs.

To illustrate how this within-host generated diversity would be captured in our sampling, pairs of

genomes were repeatedly sampled from simulated populations and pairwise differences

calculated (Fig 6C and 6D, File S1). Overall, pairwise differences greater than 2 SNPs occurred

in less than 0.2% of all simulated populations (Table S5). Higher SNP distances such as those

observed (Fig. 5B) are therefore unlikely to arise within-host following infection with a

homogenous inoculum.

Discussion

Genomic data for understanding the population structure and transmission patterns of bacterial

zoonoses has been limited for LMIC settings where these diseases tend to have the greatest

impact. Despite *B. anthracis* having limited genomic diversity in comparison with other bacterial

species, WGS provided sufficient resolution to discriminate among isolates collected from within

a relatively small geographic area of a few thousand square kilometers. The way in which the

genomic diversity was partitioned across hierarchical spatial scales within this area has a series

of novel implications for our understanding of how the pathogen is transmitted and evolves

during endemic circulation.

B. anthracis diversity within the hyperendemic NCA region is limited to a single Clade A sub-group

The NCA is an area where anthrax has likely been endemic for decades, if not centuries. Local

community members claim that it has been an issue for their health and that of their livestock

throughout living memory. Despite WGS providing sufficient discriminatory power to

differentiate among individual *B. anthracis* isolates in this setting, fewer than 50 SNP differences

were found among NCA isolates across the 5.2 MB chromosomal genome, highlighting the

degree to which this bacterial pathogen is monomorphic. All 73 sequenced isolates formed a

monophyletic cluster within the Ancient A lineage – also known as canSNP group A.Br.005/006

or A.Br.034 (6) – a subgroup of Clade A comprised mostly of isolates from south-eastern Africa.

This contrasts with some previous studies of anthrax diversity in chronically endemic areas

reporting co-circulation of isolates from multiple lineages using lower resolution markers (8, 9).

Our study, which only included isolates collected over one year, can be considered a snapshot of

recent diversity only. That being said, Tanzanian isolates genotyped in a previous study (n = 17)

were also all found to belong to Ancient A clade (5), demonstrating that this particular lineage is

dominant and well-established in this country. The most closely related publicly available isolate

(A2075), sampled in central Tanzania approximately 300km away from our study area, differed

from some NCA isolates by only 13 SNPs, which illustrates that highly related *B. anthracis*

isolates can be geographically widespread. Broader, longitudinal WGS studies of *B. anthracis*

across different regions of Africa will be needed to assess the actual range of individual

genotypes over space and time.

No phylogeographic signal observed despite considerable SNP diversity

Despite the considerable diversity observed (i.e. 22 unique genotypes within the sampled

population), a phylogeographic signal was not detected at the scale of this study area. The

finding of identical SNP profiles across distances of tens of kilometers likely reflects the ecology

of anthrax in general and in our study system. First, there are few opportunities for genetic

diversity to arise within the *B. anthracis* lifecycle, which is characterized by long periods of

environmental dormancy in spore form, punctuated by brief interludes of a few days where it

develops into its vegetative state and replicates within an infected host. It is estimated that *B.*

anthracis undergoes only 20-40 replications per infection (11). Given its low mutation rate (5.2 –

8.3 x 10^{-10} mutations/site/generation) (5, 29, 31), novel mutations would likely arise in only a

small proportion of *B. anthracis* infections, as also supported by our simulations. The short time

of active replication within a host also means there is minimal opportunity for horizontal gene

transfer with other bacteria (11), which further restricts the ability of *B. anthracis* to diversify. It

is believed that *B. anthracis* rarely multiplies outside of a host, although there is some evidence

for limited environmental replication (32, 33). Viability of *B. anthracis* decays exponentially

over time, although infectious spores remain detectable at carcass sites at least four years after

the death of the animal (34), and under favorable conditions, spores can remain viable for up to

several decades (20). It is therefore reasonable to expect that spores from a single anthrax carcass

with few or no novel mutations could be the source of subsequent infections over highly variable

time periods, a phenomenon referred to by Sahl et al. (2016) as a 'time capsule'. This situation

violates typical molecular clock assumptions (35), wherein SNPs would be expected to arise at a

relatively steady rate within the pathogen population over time. The observation that

contemporary isolates from the NCA are phylogenetically basal to isolate A2075, collected

nearly two decades earlier, highlights this issue. In order for molecular clock models to be

applied to environmentally-persistent pathogens such as *B. anthracis* – for instance to estimate

how long particular lineages have been in circulation – current analytical frameworks will need

to be extended.

In addition to long and variable environmental persistence, animal movements and spatial

admixture likely contribute to the lack of phylogeographic structure of *B. anthracis* within this

study area. Potential sources of infection are quite spatially restricted, as the highest

concentrations of viable spores are found within only a few meters of anthrax carcasses (36), in

what have been termed 'localized infectious zones' (37). However, given that the incubation

period of anthrax in livestock typically ranges from 1-14 days and potentially longer (3),

extensive movement can occur between the time of infection and the animal's death, meaning

the site of sampling likely does not reflect the site of infection. In parts of rural Africa where

pastoralism is the main form of agriculture, livestock are moved for various reasons, including to

access water, pastures and minerals. Such daily and/or seasonal livestock movements could

contribute to the observation of identical SNP profiles over distances of tens of kilometers in our

study area and the lack of relationship between genetic distance and geographic distance between

sampling locations. While livestock appear to be the primary drivers of *B. anthracis* transmission

in the NCA, movement of infected wildlife and scavengers acting as carriers of *B. anthracis*

spores could represent an additional mechanism contributing to our observations (38, 39).

Further comparative genomic studies across wider areas will be essential for elucidating the

geographic scales at which transmission occurs. This would help to delineate areas across which

coordinated livestock vaccination campaigns should occur to avoid regular re-incursions through

animal movement and trade.

High within-host diversity is the result of simultaneous infection with multiple

variants, not within-host evolution

We observed high *B. anthracis* diversity within individual hosts, essentially indistinguishable

from levels of diversity found throughout the study area. Smaller numbers of SNPs (1-2) could

potentially have arisen during culture; however, based on previous passaging experiments with

B. anthracis (29), we do not expect this to have contributed in our case due to the limited number

of passages performed. Alternatively, small numbers of SNPs could have arisen during the

course of infection within the host. The simulations we performed suggest that isolates with >2

SNP differences between them are unlikely to be the product of within-host evolution. The

regular occurrence of such differences or greater among isolates from the same carcass indicates

that animals are commonly infected with a heterogenous infectious dose (i.e. ingestion of a

mixture of *B. anthracis* genotypes from single or possibly multiple grazing or watering points;

Fig. 7)*. Multiple SNP profiles were present among isolates from the same individual soil

samples, supporting the occurrence of heterogeneity in a single infection source. It is also

noteworthy that all within-host SNP profiles observed in this study were shared with isolates

from at least one other sampled carcass (i.e. none were unique); this strongly suggests that most

of the observed within-host diversity is the result of various genotypes having been present in the

inoculum rather than having been generated *de novo*. This points to a wide transmission

bottleneck, since a small inoculum comprising only a few spores would limit the possible

diversity that could be transmitted.

Our findings make an important contribution to the ongoing debate about the size of the

transmission bottleneck for naturally-occurring anthrax in animals (i.e. the number of spores that

give rise to a case). Recent work has proposed that founding populations may be as small as 1-3

individual spores (30). However, our findings are clearly at odds with such narrow bottlenecks,

since the multiple genotypes we regularly observed within individual hosts could not have been

transmitted within such a small inoculum. While our results provide no information about the

exact size of the infectious dose, they align more closely with earlier suggestions of higher

infectious doses, which are biologically plausible given that animals grazing at carcass sites

might ingest hundreds of thousands of spores with each bite (34).

Spatio-temporally linked anthrax cases are rarely genetically linked

Limiting phylogenetic analyses to a single isolate per host often leads to incorrect inference of

transmission events (22), including the potential to overlook important epidemiological

connections (40). This issue is exemplified in the current study by the spatio-temporally linked

pair of cases in Cluster 2: whereas several of the isolates from both carcasses (n = 3 each) had

identical SNP profiles, supporting a transmission link, both carcasses also harbored non-identical

genotypes (Fig. S5 and S6). Under these circumstances, transmission links may be missed, even

with complete sampling. As noted by Ågren et al. (2014), in the case of multi-clonal infections,

subsequent cases may stem from different genotypes from within the founding population,

masking the fact that these cases stemmed from a common source, regardless of the number of

isolates sequenced among the subsequent cases. This could be the case for the other three spatio-

temporally linked pairs of anthrax carcasses investigated in this study (Fig. S6), in which we

only detected isolates with distinct SNP profiles. Alternatively, cases without a genetic link

could be temporally linked for reasons other than exposure to a common source. For instance,

animals might be more susceptible or at greater risk of exposure to infection at particular times

of year, e.g. due to lower immune function related to their nutritional status and/or associated

with weather extremes including prolonged rains or droughts (8). More extensive sampling of

within-carcass diversity would be necessary to investigate these hypotheses and to determine

whether co-occurrence of genotypes could be used to track transmission patterns.

Conclusions

In this study, the genomic diversity of *B. anthracis* was quantified at various spatial scales within a hyperendemic setting. While WGS could discriminate among isolates within a relatively small geographic area, there was a lack of phylogeographic signal and limited genetic relatedness was observed among isolates from spatio-temporally linked cases. We hypothesize that this lack of spatial structure reflects the long-term persistence of *B. anthracis* spores in the environment, combined with extensive livestock movements related to local pastoralist practices. Based on simulations, the high within-host heterogeneity we observed points to an inoculum comprised of diverse genotypes, suggestive of a wide transmission bottleneck. Our work paves the way for studying *B. anthracis* genomic diversity and evolution within anthrax-endemic areas more broadly and to confirm the temporal and spatial scales over which genomic data are most informative for inferring transmission dynamics.

Methods

Study area

This study was conducted in the NCA of northern Tanzania, which covers 8,292 km^2. Located to the south-east of Serengeti National Park, this multiple-land use area is inhabited by roughly 87,000 people (41) and one million livestock (sheep, goats and cattle) (Veterinary Officer for Ngorongoro District, pers comm). Northern Tanzania remains hyperendemic for anthrax (42), and prior to this study the NCA was recognized as a potential hotspot for this disease (43).

Research and ethical approval

This study received ethical approval from the Kilimanjaro Christian Medical University College

Ethics Review Committee (certificate No. 2050); the National Institute for Medical Research,

Tanzania (NIMR/HQ/R.8a/Vol. IX/2660); Tanzanian Commission for Science and Technology

(2016-95-NA-2016-45); and the College of Medical Veterinary and Life Sciences ethics

committee at the University of Glasgow (200150152). It also received permission under Section

20 of the Animal Diseases Act 35 (1984) at the University of Pretoria, South Africa (Ref

12/11/1/1/6).

Sample collection

Samples were collected between May 2016 and April 2017 inclusive through active surveillance

by a dedicated field team. Sudden deaths in animals reported by community members throughout

the NCA were investigated and samples were collected when anthrax was suspected (File S1).

When available, the following samples were collected: a piece of skin tissue (tip of the ear if the

carcass was still intact, or a piece of hide if the carcass had already been opened); whole blood;

swab of blood or body fluid at natural orifices; blood- or body fluid-soaked soil from below the

carcass; and insects found on or around the carcass. Various metadata were recorded, including

the species of animal affected and the location of sampling (Table S1, File S1). All samples were

stored at ambient temperature for up to six months at local veterinary facilities until transport to

the Kilimanjaro Clinical Research Institute (KCRI) in Moshi, Tanzania for molecular

diagnostics, as previously described (44), with aliquots shipped to the University of Pretoria,

South Africa, for selective culture and DNA extraction from *B. anthracis* isolates.

Selective culture, DNA extraction and sequencing

Sample pretreatment (i.e. to inhibit competition from heat-sensitive bacteria) is described in File

S1. Sample homogenates (100 µL) were plated onto both polymyxin-EDTA thallous acetate

(PET) selective media and 5% sheep blood agar (SBA). These were incubated at 37 °C overnight

and the plates inspected for growth after 15 – 24 hours incubation. The PET was then further

incubated and inspected at 48 hours. Suspect *B. anthracis* colonies based on typical

morphological characteristics were sub-cultured onto SBA for purification and identification

(File S1). In parallel, a single colony was streaked onto a new purity plate for nucleic acid

extraction. In some instances multiple isolates were selected from the same sample where the

colonies demonstrated differences in morphology but were identified on the same plate and met

the selection criteria (File S1).

DNA extracts from 75 isolates from 33 carcasses were submitted for library preparation and

sequencing at MicrobesNG (Birmingham, UK). Libraries were prepared using the Nextera XT

v2 kit (Illumina, San Diego, USA) and sequenced on the Illumina HiSeq platform, generating

250 base pair paired-end reads.

Bioinformatics and genomic analyses

Reads were adapter trimmed by MicrobesNG using Trimmomatic v0.30 (45), and basic statistics

determined using QUAST (46) (Table S1). Bacterial species identification was confirmed using

Kraken (47). Based on these quality metrics, sequences from two isolates were excluded from

further analyses: one due to a low number of reads (<40,000), and another that was identified as

B. cereus. There were some further indications that not all cultures were pure *B. anthracis,*

despite multiple rounds of sub-culture (File S1, Table S1). A reference-based mapping approach

and strict variant filtering criteria were implemented to minimize the issues associated with the

sequence quality while making use of as much of the data as possible.

Read mapping and variant calling were performed on the CLIMB computing platform for

microbial genomics (48). Trimmed reads were aligned to the chromosome of the Ames Ancestor

reference genome (NC_007530) using bwa-mem (version 0.7.17). Picard was used to mark and

remove duplicate reads, add read group information, and index the bam files (49). Quality

metrics for read mapping were obtained using Qualimap (50) (Table S1). SNPs were detected in

individual isolates by VarScan v2.4.4 (51) with parameters set as follows: minimum read depth

of 4; minimum base quality of 20; variant allele frequency ≥ 0.95. Subsequent SNP curation

steps are described in File S1. Custom python scripts for the assessment of read mapping SNP

metrics data, variant site filtering and generation of variant call and alignment files (source codes

with description of their functionality and usage), along with the final variant call and multiple

sequence alignment files are available on GitHub

(https://github.com/matejmedvecky/anthraxdiversityscripts).

The alignment of concatenated SNPs was analyzed using ModelFinder to determine the most

appropriate model of nucleotide substitution (52). Subsequently, a maximum-likelihood

phylogeny was estimated in IQ-Tree (53) under the Kimura-3-parameters (K3P) model, using

1000 ultrafast bootstrap replicates (54). A distance matrix detailing SNP differences between

isolates was constructed using snp-dists v0.6 (https://github.com/tseemann/snp-dists). The

distance between GPS points was calculated using the pointDistance command in the R package

raster (55). Isolation by distance was tested using Mantel test to assess the correlation between

SNP distance and Euclidean geographic distance within the R package *adegenet* (56). All

program versions and commands used, the distance matrix, as well as small custom scripts are

available on GitHub (https://github.com/tristanpwdennis/anthrax_diversity).

To place the newly sequenced isolates within the global phylogeny of *B. anthracis*, 80 WGS

from GenBank were accessed (Table S2), and a core genome alignment generated using Parsnp

v1.1.2 (57). The resulting phylogeny indicated that NCA isolates belong to the Ancient A

lineage. To further resolve the diversity among NCA isolates compared with other publicly

available isolates from the same lineage, reads from eight additional isolates available on SRA

from a study by Bruce et al. 2019 were accessed using fastq-dump from the SRA-toolkit: all

isolates (n=4) belonging to the 3.2 linage, and two arbitrarily selected isolates from each the 3.1

and 3.3 lineages (Table S2). These were run through our SNP-calling pipeline as described

above, resulting in a sequence alignment file free of -/N characters, which was used to infer a

phylogeny in RAxML v8.2.11 (58) using a GTR model of nucleotide substitution, and using the

Ames Ancestor reference genome as an outgroup.

Simulation modelling

The initial number of genomes in the inoculum was varied between one and 100 genomes to

reflect uncertainty in anthrax infectious dose. In each bacterial generation, genomes underwent a

round of replication followed by cell division and accordingly population size doubled in each

generation. The number of mutations occurring in the replication of each genome was drawn

from a Poisson distribution parameterized to reflect the estimated *B. anthracis* mutation rate (i.e.

$\lambda = 0.004316$). This genome-level mutation rate is based on the genome size of 5.2 million base

pairs and a mutation rate of 8.3×10^{-10} mutations per site (29); this represents an upper estimate

of the mutation rate and was chosen as we were interested in estimating the upper limits on

reasonable expectation of diversity emerging during the course of an infection initiated by a

homogenous dose. Simulations were carried out for up to 30 generations (replication cycles), and

results extrapolated to 40 generations, proposed to be the upper limit on the number of

replication cycles during an infection (11). Details of simulations run are provided in File S1.

Pairs of genomes were repeatedly sampled from these simulated populations and the count of

mutations separating each pair was calculated. Sampling was performed 100 times per

generation for each simulation. Simulations, sampling of simulated populations, and linear

models summarizing trends in the outcome of these processes were performed in R (59).

Data availability

440 Raw sequencing reads are available on European Nucleotide Archive SRA under accession

441 number PRJEB45684.

Acknowledgements

We are grateful for all the support received for this research, particularly from the NCA

community and authorities. We thank the Ngorongoro District Council, Ngorongoro

Conservation Area Authority, District Veterinary Office, Tanzania Wildlife Research Institute

and members of our field team – Sabore Ole Moko, Sironga Nanjicho, Kadogo Lerimba and

Godwin Mshumba – for assistance with this study. We also thank the Directorate of Veterinary

Services, Ministry of Agriculture, Livestock and Fisheries, and Ministry of Health, Community

Development, Gender, Elderly and Children for their support. Finally, we thank Yi Xuan Chew

and Nichith Kollanandi Ratheesh for initial assessments of the publicly available genomic data.

Establishment and maintenance of the Zoonoses laboratory at KCRI (BTM) was supported by

the BBSRC (BB/J010367/1), BBSRC Zoonoses and Emerging Livestock Systems (BB/L017679,

BB/L018926, BB/L018845), and the Wellcome Trust (096400/Z/11/Z).

References

1.   Kao RR, Haydon DT, Lycett SJ, Murcia PR. 2014. Supersize me: how whole-genome

sequencing and big data are transforming epidemiology. Trends Microbiol 22:282–91.

2.   Carlson CJ, Kracalik IT, Ross N, Alexander KA, Hugh-Jones ME, Fegan M, Elkin BT, Epp

T, Shury TK, Zhang W, Bagirova M, Getz WM, Blackburn JK. 2019. The global

distribution of *Bacillus anthracis* and associated anthrax risk to humans, livestock and

wildlife. Nat Microbiol 4:1337–1343.

3.   WHO. 2008. Anthrax in Humans and Animals. 4th Ed.

4.   WHO. 2011. The control of neglected zoonotic diseases. Community-based interventions

for prevention and control. Report of the third conference organized with ICONZ, DFID-

RIU, Gates Foundation, SOS, EU, TDR and FAO with the participation of ILRI and OIE.,

Geneva, Switzerland.

5.   Van Ert MN, Easterday WR, Huynh LY, Okinaka RT, Hugh-Jones ME, Ravel J, Zanecki

SR, Pearson T, Simonson TS, U'Ren JM, Kachur SM, Leadem-Dougherty RR, Rhoton SD,

Zinser G, Farlow J, Coker PR, Smith KL, Wang B, Kenefic LJ, Fraser-Liggett CM, Wagner

DM, Keim P. 2007. Global genetic population structure of *Bacillus anthracis*. PLoS ONE

2:e461.

6.   Sahl JW, Pearson T, Okinaka R, Schupp JM, Gillece JD, Heaton H, Birdsell D, Hepp C,

Fofanov V, Noseda R, Fasanella A, Hoffmaster A, Wagner DM, Keim P. 2016. A *Bacillus*

anthracis genome sequence from the Sverdlovsk 1979 autopsy specimens. mBio 7: e01501-

16.

7.   Bruce SA, Schiraldi NJ, Kamath PL, Easterday WR, Turner WC. 2020. A classification

476   framework for *Bacillus anthracis* defined by global genomic structure. Evol Appl 13:935–

477   944.

478   8.   Smith KL, DeVos V, Bryden H, Price LB, Hugh-Jones ME, Keim P. 2000. *Bacillus*

479   *anthracis* diversity in Kruger National Park. J Clin Microbiol 38:3780–3784.

480   9.   Simonson TS, Okinaka RT, Wang B, Easterday WR, Huynh L, U'Ren JM, Dukerich M,

481   Zanecki SR, Kenefic LJ, Beaudry J, Schupp JM, Pearson T, Wagner DM, Hoffmaster A,

482   Ravel J, Keim P. 2009. *Bacillus anthracis* in China and its relationship to worldwide

483   lineages. BMC Microbiol 9:71.

484   10.   Achtman M. 2008. Evolution, population structure, and phylogeography of genetically

485   monomorphic bacterial pathogens. Annu Rev Microbiol 62:53–70.

486   11.   Keim P, Van Ert MN, Pearson T, Vogler AJ, Huynh LY, Wagner DM. 2004. Anthrax

487   molecular epidemiology and forensics: using the appropriate marker for different

488   evolutionary scales. Infect Genet Evol 4:205–213.

489   12.   Fasanella A, Serrecchia L, Chiaverini A, Garofolo G, Muuka GM, Mwambazi L. 2018. Use

490   of Canonical Single Nucleotide Polymorphism (CanSNPs) to characterize *Bacillus*

491   *anthracis* outbreak strains in Zambia between 1990 and 2014. PeerJ 6:e5270.

492   13.   Beyer W, Bellan S, Eberle G, Ganz HH, Getz WM, Haumacher R, Hilss KA, Kilian W,

493   Lazak J, Turner WC, Turnbull PCB. 2012. Distribution and molecular evolution of *Bacillus*

494   *anthracis* genotypes in Namibia. PLoS Negl Trop Dis 6:e1534.

14. Maho A, Rossano A, Hächler H, Holzer A, Schelling E, Zinsstag J, Hassane MH,

Toguebaye BS, Akakpo AJ, Van Ert M, Keim P, Kenefic L, Frey J, Perreten V. 2006.

Antibiotic susceptibility and molecular diversity of *Bacillus anthracis* strains in Chad:

detection of a new phylogenetic subgroup. J Clin Microbiol 44:3422–3425.

15. Girault G, Blouin Y, Vergnaud G, Derzelle S. 2014. High-throughput sequencing of

Bacillus anthracis in France: investigating genome diversity and population structure using

whole-genome SNP discovery. BMC Genomics 15:288.

16. Derzelle S, Girault G, Roest HIJ, Koene M. 2015. Molecular diversity of *Bacillus anthracis*

in the Netherlands: investigating the relationship to the worldwide population using whole-

genome SNP discovery. Infect Genet Evol 32:370–376.

17. Derzelle S, Girault G, Kokotovic B, Angen Ø. 2015. Whole genome-sequencing and

phylogenetic analysis of a historical collection of *Bacillus anthracis* strains from Danish

cattle. PLoS ONE 10:e0134699.

18. Lienemann T, Beyer W, Pelkola K, Rossow H, Rehn A, Antwerpen M, Grass G. 2018.

Genotyping and phylogenetic placement of *Bacillus anthracis* isolates from Finland, a

country with rare anthrax cases. BMC Microbiology 18:102.

19. Lekota KE, Hassim A, Madoroba E, Hefer CA, van Heerden H. 2020. Phylogenomic

structure of *Bacillus anthracis* isolates in the Northern Cape Province, South Africa

revealed novel single nucleotide polymorphisms. Infection, Genetics and Evolution

80:104146.

20. Carlson CJ, Getz WM, Kausrud KL, Cizauskas CA, Blackburn JK, Bustos Carrillo FA,

Colwell R, Easterday WR, Ganz HH, Kamath PL, Økstad OA, Turner WC, Kolstø A-B,

Stenseth NC. 2018. Spores and soil from six sides: interdisciplinarity and the environmental

biology of anthrax (*Bacillus anthracis*). Biol Rev Camb Philos Soc 93:1813–1831.

21. Döpfer D, Buist W, Soyer Y, Munoz MA, Zadoks RN, Geue L, Engel B. 2008. Assessing

genetic heterogeneity within bacterial species isolated from gastrointestinal and

environmental samples: how many isolates does it take? Appl Environ Microbiol 74:3490–

3496.

22. Worby CJ, Lipsitch M, Hanage WP. 2014. Within-host bacterial diversity hinders accurate

reconstruction of transmission networks from genomic distance data. PLoS Comput Biol

10:e1003549.

23. Futse JE, Brayton KA, Dark MJ, Knowles DP, Palmer GH. 2008. Superinfection as a driver

of genomic diversification in antigenically variant pathogens. Proc Natl Acad Sci USA

105:2123–2127.

24. Sintchenko V, Holmes EC. 2015. The role of pathogen genomics in assessing disease

transmission. BMJ 350:h1314.

25. Didelot X, Walker AS, Peto TE, Crook DW, Wilson DJ. 2016. Within-host evolution of

bacterial pathogens. Nat Rev Microbiol 14:150–162.

26. Paterson GK, Harrison EM, Murray GGR, Welch JJ, Warland JH, Holden MTG, Morgan

FJE, Ba X, Koop G, Harris SR, Maskell DJ, Peacock SJ, Herrtage ME, Parkhill J, Holmes

MA. 2015. Capturing the cloud of diversity reveals complexity and heterogeneity of MRSA

carriage, infection and transmission. Nat Commun 6:6560.

27. Alamil M., Hughes J., Berthier K., Desbiez C., Thébaud G., Soubeyrand S. 2019. Inferring

epidemiological links from deep sequencing data: a statistical learning approach for human,

animal and plant diseases. Philosophical Transactions of the Royal Society B: Biological

Sciences 374:20180258.

28. Beyer W, Turnbull PCB. 2013. Co-infection of an animal with more than one genotype can

occur in anthrax. Lett Appl Microbiol 57:380–384.

29. Ågren J, Finn M, Bengtsson B, Segerman B. 2014. Microevolution during an anthrax

outbreak leading to clonal heterogeneity and penicillin resistance. PLoS ONE 9:e89112.

30. Easterday WR, Ponciano JM, Gomez JP, Van Ert MN, Hadfield T, Bagamian K, Blackburn

JK, Stenseth NC, Turner WC. 2020. Coalescence modeling of intrainfection *Bacillus*

anthracis populations allows estimation of infection parameters in wild populations. Proc

Natl Acad Sci USA 117:4273-4280.

31. Vogler AJ, Busch JD, Percy-Fine S, Tipton-Hunton C, Smith KL, Keim P. 2002. Molecular

analysis of rifampin resistance in *Bacillus anthracis* and *Bacillus cereus*. Antimicrob

Agents Chemother 46:511–513.

32. Saile E, Koehler TM. 2006. *Bacillus anthracis* multiplication, persistence, and genetic

exchange in the rhizosphere of grass plants. Appl Environ Microbiol 72:3168–3174.

33. Braun P, Grass G, Aceti A, Serrecchia L, Affuso A, Marino L, Grimaldi S, Pagano S,

Hanczaruk M, Georgi E, Northoff B, Schöler A, Schloter M, Antwerpen M, Fasanella A.

2015. Microevolution of Anthrax from a Young Ancestor (M.A.Y.A.) suggests a soil-borne

life cycle of *Bacillus anthracis*. PLoS ONE 10:e0135346.

34. Turner WC, Kausrud KL, Beyer W, Easterday WR, Barandongo ZR, Blaschke E, Cloete

CC, Lazak J, Van Ert MN, Ganz HH, Turnbull PCB, Stenseth NC, Getz WM. 2016. Lethal

exposure: An integrated approach to pathogen transmission via environmental reservoirs.

Sci Rep 6:27311.

35. Biek R, Pybus OG, Lloyd-Smith JO, Didelot X. 2015. Measurably evolving pathogens in

the genomic era. Trends in Ecology & Evolution 30:306–313.

36. Dragon DC, Bader DE, Mitchell J, Woollen N. 2005. Natural dissemination of *Bacillus*

anthracis spores in northern Canada. Appl Environ Microbiol 71:1610–1615.

37. Blackburn JK, Ganz HH, Ponciano JM, Turner WC, Ryan SJ, Kamath P, Cizauskas C,

Kausrud K, Holt RD, Stenseth NC, Getz WM. 2019. Modeling R_0 for pathogens with

environmental transmission: Animal movements, pathogen populations, and local infectious

zones. Int J Environ Res Public Health 16:954.

38.   1979. Vultures as carriers of anthrax. J S Afr Vet Assoc 50:35.

39. Stears K, Schmitt MH, Turner WC, McCauley DJ, Muse EA, Kiwango H, Mathayo D,

Mutayoba BM. 2021. Hippopotamus movements structure the spatiotemporal dynamics of

an active anthrax outbreak. Ecosphere 12:e03540.

40. Forde TL, Orsel K, Zadoks RN, Biek R, Adams LG, Checkley SL, Davison T, De Buck J,

Dumond M, Elkin BT, Finnegan L, Macbeth BJ, Nelson C, Niptanatiak A, Sather S,

Schwantje HM, Van Der Meer F, Kutz SJ. 2016. Bacterial genomics reveal the complex

epidemiology of an emerging pathogen in arctic and boreal ungulates. Front Microbiol

7:1759.

41. National Bureau of Statistics, Office of Chief Government Statistician. 2013. 2012

Population and Housing Census. The United Republic of Tanzania.

42. Mwakapeje ER, Høgset S, Fyumagwa R, Nonga HE, Mdegela RH, Skjerve E. 2018.

Anthrax outbreaks in the humans - livestock and wildlife interface areas of Northern

Tanzania: a retrospective record review 2006–2016. BMC Public Health 18:106.

43. Lembo T, Hampson K, Auty H, Beesley CA, Bessell P, Packer C, Halliday J, Fyumagwa R,

Hoare R, Ernest E, Mentzel C, Mlengeya T, Stamey K, Wilkins PP, Cleaveland S. 2011.

Serologic surveillance of anthrax in the Serengeti ecosystem, Tanzania, 1996-2009.

Emerging Infect Dis 17:387–394.

44. Aminu OR, Lembo T, Zadoks RN, Biek R, Lewis S, Kiwelu I, Mmbaga BT, Mshanga D,

Shirima G, Denwood M, Forde TL. 2020. Practical and effective diagnosis of animal

anthrax in endemic low-resource settings. PLOS Neglected Tropical Diseases 14:e0008655.

45. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114–2120.

46. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for

genome assemblies. Bioinformatics 29:1072–1075.

47. Wood DE, Salzberg SL. 2014. Kraken: ultrafast metagenomic sequence classification using

exact alignments. Genome Biol 15:R46.

48. Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, Bull MJ,

Richardson E, Ismail M, Elwood-Thompson S, Kitchen C, Guest M, Bakke M, Sheppard

SK, Pallen MJ. 2016. CLIMB (the Cloud Infrastructure for Microbial Bioinformatics): an

online resource for the medical microbiology community. Microb Genom 2:e000086.

49. Broad Institute. Picard Tools. Available: http://broadinstitute.github.io/picard/

50. García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, Dopazo J,

Meyer TF, Conesa A. 2012. Qualimap: evaluating next-generation sequencing alignment

data. Bioinformatics 28:2678–2679.

51. Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER,

Ding L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration

discovery in cancer by exome sequencing. Genome Res 22:568–576.

52. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nature Methods

14:587–589.

53. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol

Evol 32:268–274.

54.   Minh BQ, Nguyen MAT, von Haeseler A. 2013. Ultrafast approximation for phylogenetic

615   bootstrap. Mol Biol Evol 30:1188–1195.

616   55.   raster: Geographic Data Analysis and Modeling version 3.0-12 from CRAN.

617   56.   Jombart T. 2008. adegenet: a R package for the multivariate analysis of genetic markers.

618   Bioinformatics 24:1403–1405.

619   57.   Treangen TJ, Ondov BD, Koren S, Phillippy AM. 2014. The Harvest suite for rapid core-

620   genome alignment and visualization of thousands of intraspecific microbial genomes.

621   Genome Biol 15:524.

622   58.   Stamatakis A. 2014. RAxML Version 8: A tool for phylogenetic analysis and post-analysis

623   of large phylogenies. Bioinformatics 30:1312-1313.

624   59.   R Core Team. 2018. R: A language and environment for statistical computing. R

625   Foundation for Statistical Computing, Vienna, Austria.

626   60.   UNEP-WCMC. 2020. Protected Area Profile for Ngorongoro Conservation Area. World

627   Database of Protected Areas. Available: www.protectedplanet.net

628   61.   Wickham H. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New

629   York.

630   62.   Pebesma E. 2018. Simple features for R: Standardized support for spatial vector data. The R

631   Journal 10:439–446.

63. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. 2017. ggtree: an R package for visualization

and annotation of phylogenetic trees with their covariates and other associated data.

Methods Ecol Evol 8:28–36.

64. Ciccarelli FD, Doerks T, Mering C von, Creevey CJ, Snel B, Bork P. 2006. Toward

automatic reconstruction of a highly resolved Tree of Life. Science 311:1283–1287.

65. Hsieh TC, Ma KH, Chao A. 2016. iNEXT: an R package for rarefaction and extrapolation

of species diversity (Hill numbers). Methods Ecol Evol 7:1451–1456.

Figures

1. Geographical Groups   2. Epidemiological Clusters   3. Within Infected Host

Fig 1. Hierarchical levels at which the genomic diversity of *Bacillus anthracis* was studied.

1. Sequenced isolates originating from livestock carcasses sampled throughout the anthrax

hyperendemic Ngorongoro Conservation Area (NCA), northern Tanzania, (shown as grey dots in

panel 1) were categorized into four geographical groups (colored circles); 2. A subset of these

isolates were from spatio-temporally linked pairs of carcasses (n = 4, represented by grey shapes

in panel 1), referred to as 'epidemiological clusters'; 3. Multiple isolates (n = 2-4; represented by

asterixis) were sequenced from individual carcasses, either originating from multiple sample

types (e.g. tissue and blood), and/or multiple isolates sequenced from a single sample. The shape

file for the NCA was provided by Tanzania National Parks (TANAPA) (60).

Fig 2. Phylogenetic position of *Bacillus anthracis* isolated from the Ngorongoro

Conservation Area within the global population. A) Global phylogeny of *B. anthracis*,

showing the major clades (A, B, and C) and sub-lineages. This tree was estimated based on a

core single nucleotide polymorphism (SNP) phylogeny of 80 publicly available genomes (Table

S2). B) Maximum likelihood phylogenetic tree of the Ancient A lineage (Cluster 3 based on

Bruce et al., 2019). All isolates from the Ngorongoro Conservation Area (NCA) form a

monophyletic lineage within Cluster 3.2, along with the publicly available isolate A2075

(SRR2968187) isolated in 1999 from a baboon in central Tanzania. This tree was inferred using

the general time reversible (GTR) model of nucleotide substitution, using the Ames Ancestor

reference genome (NC_007530) as an outgroup. Tree scales reflect number of substitutions per

site.

Fig 3. Phylogeography of *Bacillus anthracis* in the hyperendemic area of the Ngorongoro

Conservation Area, Tanzania. A) Spatial distribution of carcasses from which *B. anthracis*

isolates were obtained. The map outlines the Ngorongoro Conservation Area (NCA) and shows

its location in northern Tanzania (outlined in red in inset). Carcasses were assigned to four

geographical groups within the NCA based on spatial proximity, shown by colored circles. B)

Maximum likelihood tree estimating the phylogenetic relationship among *B. anthracis* isolates

from the NCA. This tree is based on an un-gapped alignment of 125 high quality core single

nucleotide polymorphisms (SNPs) across the whole chromosome, rooted to the Ames Ancestor

reference sequence (NC_005730) and including the publicly available isolate A2075

(SRR2968187) originating from central Tanzania. Using the more closely related isolates from

Cluster 3.2 as an outgroup produced the same root position. Isolates are colored on the tree based

on their collection site (geographical group) within the NCA. Epidemiological clusters of cases

(pairs of carcasses sampled from the same or neighboring households on the same or consecutive

days) are distinguished by symbol shape. Letters distinguish the 22 unique genotypes (SNP

profiles) detected. Isolate-labelled versions of these figures can be found as Fig. S4 and S5 in the

Supporting Information. The base earth, river and lake data for the map were downloaded from

Natural Earth (https://www.naturalearthdata.com/). Figure was plotted in R v.3.6.1 with *ggplot2*

(61), with the addition of the *sf* (62) and *ggtree* (63) packages.

Fig 4. Comparative numbers of single nucleotide differences among *Bacillus anthracis*

isolates from hierarchical spatial scales. Violin plots comparing pairwise nucleotide (SNP)

differences between all *B. anthracis* isolates from the Ngorongoro Conservation Area versus

SNP differences between isolates from i) within the same geographical group; ii) within the same

epidemiological cluster (but not from the same carcass); and iii) within a single carcass. Central

boxplot shows median and interquartile range, with whiskers showing minimum and maximum

values.

Fig 5. Within-host diversity of *Bacillus anthracis* among livestock in the Ngorongoro

Conservation Area. A) Circularized maximum likelihood tree – based on high quality core

single nucleotide polymorphisms (SNPs) – displayed as a cladogram (branch-lengths ignored),

rooted to Ames Ancestor reference genome (NC_005730). Isolates from the same carcass are

shown in the same color and are linked by inner connecting lines. Isolates without labels are

singletons (i.e. only one isolate sequenced per carcass site). Sample type is shown by the

different symbol shapes indicated in the legend. The figure was prepared using iTOL (64). For

labeled taxa, see Fig. S7. B) Histogram showing the relative frequency of pairwise SNP

differences among *B. anthracis* isolates collected from the same carcass. Red line shows 99%

upper limit of nucleotide differences observed among sampled pairs of genomes based on

simulation of within host evolution. Results suggest that almost all diversity observed within the

same infected host is the result of a heterogenous inoculum.

Fig 6. Simulation of within-host populations and sampling of resulting genetic diversity.

Box and whiskers plots show the proportion of genomes across simulated populations that differ

from the founding genome by either one nucleotide (A) or by two nucleotides (B). Boxes

represent the interquartile range with a line showing the median and with outliers shown as

points. Both boxes and outliers are colored by inoculum size (dose) according to the legend.

Simulations were run for 25 generations, or 20 generations for larger inoculum sizes. Greater

stochastic heterogeneity was observed for lower starting doses. Note that in (B), only outliers are

visible as the proportions observed were very low. C) Bar-plot showing the relationship between

inoculum size (dose) and single nucleotide polymorphism (SNP) distances between pairs of

sampled genomes. Bars represent average proportions of pairwise SNP differences between

genomes sampled from simulated within-host populations. Populations were simulated from

various inoculum sizes and were sampled in the 20th generation after 19 replication cycles. Fifty

simulations were run from an initial dose of 20, and 100 simulations were run from each of the

other inoculum sizes, with each simulated population sampled 100 times in each generation. D)

Bar-plot showing the relationship between the number of replication cycles and SNP distances

between pairs of sampled genomes. Populations were simulated for 25 generations and samples

taken in generations 20-25 are represented in shades of grey according to the legend. Proportions

are averaged across simulations initiated with inoculum sizes of 1, 2, 5, 10 and 20 and sampled

repeatedly.

Fig 7. Conceptual framework for the acquisition of within-host diversity of *Bacillus*

***anthracis*.** Colors represent different genotypes of *B. anthracis* (with multiple nucleotide

differences), while different shades (i.e. light green) represent single nucleotide variants arising

during the course of infection. Vertical black lines represent the environment (e.g. soil, water)

from which subsequent cases of anthrax arise through exposure to *B. anthracis* spores. A)

Example of infections stemming from small transmission bottleneck (inoculum of few spores),

resulting in primarily clonal transmission. Where the bottleneck is narrow, a limited number of

genotypes could comprise the inoculum, regardless of the number of variants present in the

environment. A small number of SNPs may arise during the course of infection (see Figs 5B and

6), but are unlikely to be transmitted unless they arise early during the course of infection. These

variants rarely differ from the founding genotype by > 1-2 SNPs (Table S5). B) Example of

infections stemming from a wide bottleneck (infectious dose with multiple spores), wherein

sufficient numbers of spores comprise the inoculum such that multiple genotypes present in the

environmental source may seed initial and subsequent infections. Our results strongly support

heterogenous transmission (B), either from single or multiple carcass sites, and a large

transmission bottleneck. Figure adapted from Ågren et al., 2014.

Supporting Information

Fig S1. Rarefaction curve of *Bacillus anthracis* genotypic diversity within the study area.

Based on inclusion of all isolates sequenced (n = 73, left) and on all isolates that were not

sampled as part of an epidemiological cluster (n = 51, right), which might be expected to be non-

independent. Results suggest genotypes within this population have been exhaustively sampled

(i.e. that further sampling would not be expected to reveal additional genotypes). Figure

generated in R package iNEXT (65).

Fig S2. Histogram showing the relative frequency of pairwise nucleotide (SNP) differences

among *B. anthracis* isolates. Includes 73 isolates from the Ngorongoro Conservation Area,

northern Tanzania.

Fig S3. Scatter plots showing the number of nucleotide differences as a function of

geographic difference between the sampling locations. Geographic distance (in meters) is

shown on the x-axis versus nucleotide differences on the y-axis, with each point representing a

pair of isolates. A) All *Bacillus anthracis* isolates from the study area. B) The same relationship

is observed when limited to isolates from the dominant clade; this was done in order to account

for deeper divergences potentially obscuring patterns.

Fig S4. Locations of carcasses sampled within the Ngorongoro Conservation Area,

northern Tanzania. Each carcass is assigned a unique identifier (Table S1). Epidemiological

clusters are the same as those explained for Fig 3 and Fig S5.

Fig S5. Phylogenetic relationship among *Bacillus anthracis* isolates from the Ngorongoro

Conservation Area. Estimated through maximum likelihood, based on high quality core SNPs.

Geographic groups and epidemiological clusters are the same as those explained for Fig 3. Each

isolate is attributed to a carcass ID, with the sample type indicated following the underscore (B =

blood, D = soil, I = insect, S = swab, T = tissue). A number follows the sample type if more than

one isolate was sequenced from the same sample.

Fig S6. Genotypes of *Bacillus anthracis* observed in isolates from within and between pairs

of carcasses from the same epidemiological clusters (C1-4). Genotype letters correspond to

those in Fig 3 and Fig S5. Individual carcasses are numbered /1 or /2. In cluster C3, two isolates

were from a soil sample (C3/1) collected at the same household as the two cases (C3/2 and

C3/3); genotype T from this soil sample was shared with an isolate from C3/2. Otherwise, only

in C2 was there evidence of a shared genotype between pairs of carcasses (genotype A). Thus,

the level of sampling conducted here (1-4 isolates per carcass) did not produce evidence for the

same combinations of genotypes being found among linked carcasses.

Fig S7. Within-host diversity of *Bacillus anthracis* isolated from livestock in the

Ngorongoro Conservation Area of northern Tanzania. This circularized maximum likelihood

tree – based on high quality core single nucleotide polymorphisms – is displayed as a cladogram

(branch-lengths ignored). Isolates from the same carcass are shown in the same colour and linked

by inner connecting lines. Isolates in black are singletons (i.e. only one isolate sequenced per

carcass site). The figure was prepared using ITOL (64).

Fig S8. Proportion of simulated within-host populations identical to the inoculating genome

over 25 generations. Populations were simulated from homogenous inoculating doses of

varying size (A-G) and each line tracks a single simulated population through generations.

Represented are 100 simulations run for 25 generations from doses 1, 2, 5 and 10; 50 simulations

run for 25 generations (dose 20); 100 simulations run for 20 generations from doses 50 and 100

and 7 simulations run for 25 generations from inoculum dose of 50.

Fig S9. Proportion of simulated within-host populations differing from the inoculating

genome by one nucleotide (SNP). Populations were simulated from homogenous inoculating

doses of varying size (A-G) and each line tracks a single simulated population through

generations. Represented are 100 simulations run for 25 generations from doses 1, 2, 5 and 10;

50 simulations run for 25 generations (dose 20); 100 simulations run for 20 generations from

doses 50 and 100 and 7 simulations run for 25 generations from inoculum dose of 50.

Fig S10. Proportion of simulated within-host populations differing from the inoculating

genome by two nucleotides (SNPs). Populations were simulated from homogenous inoculating

doses of varying size (A-G) and each line tracks a single simulated population through

generations. Represented are 100 simulations run for 25 generations from doses 1, 2, 5 and 10;

50 simulations run for 25 generations (dose 20); 100 simulations run for 20 generations from

doses 50 and 100 and 7 simulations run for 25 generations from inoculum dose of 50.

Table S1. Metadata and sequence quality metrics of *Bacillus anthracis* isolates included in

this study. Each isolate is attributed to a carcass ID, with the sample type indicated following the

underscore (B = blood, D = soil, I = insect, S = swab, T = tissue). A number follows the sample

type if more than one isolate was sequenced from the same sample.

Table S2. Publicly available *Bacillus anthracis* isolates included to contextualize those from

the Ngorongoro Conservation Area. Sheet 1: Publicly available global collection of assembled

B. anthracis genomes. Sheet 2: Publicly available sequence data for Ancient A isolates, available

on SRA.

Table S3. Identical isolates found from different geographical groups.

Table S4. Identical or nearly identical isolates from carcasses sampled several months

apart.

Table S5. Single nucleotide polymorphism (SNP) distances among pairs of isolates sampled

from simulated within-host populations of *Bacillus anthracis*. Proportion of pairs of isolates

in evolved populations with different SNP distances across varying initial inoculum size (dose),

sampled in generations 20 and 25, and mean SNP differences across sampled pairs.

File S1. Supplementary methods and results.