Circadian patterns of energy metabolism are only partially reconstituted in gnotobiotic mice

Authors:

- Hoces, Daniel[1]

- Lan, Jiayi[2]

- Wenfei, Sun[3]

- Geiser, Tobias[1]

- Arnoldini, Markus[1]

- Nowok, Sven[4]

- Macpherson, Andrew J.[5]

- Stecher, Bärbel[6,7]

- Zenobi, Renato[2]

- Hardt, Wolf-Dietrich[8]

- Wolfrum, Christian[3]

- Slack, Emma[1,9]

1. Laboratory for Food Immunology, Institute of Food, Nutrition and Health, Department of Health Sciences and Technology, ETH Zürich, 8093 Zürich, Switzerland

2. Laboratory of Organic Chemistry, Department of Chemistry and Applied Biosciences, ETH Zürich, 8093 Zürich, Switzerland

3. Laboratory of Translational Nutrition Biology, Institute of Food, Nutrition and Health, Department of Health Sciences and Technology, ETH Zürich, 8603 Schwerzenbach, Switzerland

4. ETH Phenomics Center, Department of Biology, ETH Zürich, 8093 Zürich, Switzerland

5. Department of Biomedical Research, Gastroenterology & Mucosal Immunology, University of Bern, 3010 Bern, Switzerland

6. Max-von-Pettenkofer Institute, LMU Munich, Munich, Germany

7. German Center for Infection Research (DZIF), partner site LMU Munich, 80539 Munich, Germany

8. Institute of Microbiology, Department of Biology, ETH Zürich, 8093 Zürich, Switzerland

9. Corresponding author. Email: emma.slack@hest.ethz.ch

SUMMARY

Microbiota composition correlates with host metabolic health in both humans and mice. Here we

investigated the extent to which a gnotobiotic microbiota can mimic the influence of a complete

microbiota on mouse metabolism using an isolator-housed TSE-PhenoMaster® system. We found

that energy expenditure was equivalent between germ-free (GF) and specific-opportunistic-

pathogen free line (SPF) mice, and we extend this observation to the OligoMM12 microbiota (12

species across 5 phyla that are naturally abundant in a murine gut microbiota). Moreover,

microbiota-released calories were well compensated by food intake in all groups. However, each

group of mice have a distinctive circadian metabolic profile. OligoMM12 clustered with GF during

the light phase, but with SPF during the dark phase for both RER and metabolome. Therefore,

interactions between the host and a reductionist microbiota are non-uniform over the circadian

cycle, revealing a promising tool to identify key microbiota species/functions that modify host

metabolism.

KEYWORDS

INTRODUCTION

The gut microbiota is currently considered a key regulator of host energy metabolism (Sonnenburg and Bäckhed, 2016). In the absence of a microbiota, mice accumulated less fat (Bäckhed et al., 2004) and were protected from obesity induced by certain types of high-fat diets (Bäckhed et al., 2007; Fleissner et al., 2010; Kübeck et al., 2016). Several mechanisms have been proposed to explain this phenomenon and its relationship to metabolic imbalances (Cani et al., 2019). These include endocrine regulation of food intake (Goswami et al., 2018; Lin et al., 2012), additional energy liberated by the microbiota from dietary fibers (Turnbaugh et al., 2006), alterations in bile acid profiles (Sayin et al., 2013; Yao et al., 2018), inflammatory responses induced by some members of the microbiota (Caesar et al., 2015) and induction of thermogenesis in adipose tissue (Krisko et al., 2020; Li et al., 2019, 2021). However, given the complexity of a complete microbiota and its interactions with the host, validating any of these theories and identifying causal relationships remains a major experimental challenge (Harley and Karp, 2012; Walter et al., 2020).

Circadian variations in microbiota function adds an extra layer of complexity to metabolic interactions between the host and the microbiota. Circadian feeding is a major driver of microbiota composition (Thaiss et al., 2014; Zarrinpar et al., 2014). The luminal concentration of fermentation products such as short-chain fatty acids (SCFA) shows a dramatic circadian oscillation linked both to food intake and to intestinal motility (Tahara et al., 2018). Microbiota-derived molecules are known to influence host nutrient absorption (Wang et al., 2017) and host metabolic gene expression (Kuang et al., 2019; Thaiss et al., 2016). However, obtaining resolution on how these circadian variations affect host energy metabolism remains challenging and much of our current knowledge is derived from indirect calorimetry measurements made over a time period shorter than 24h (Bäckhed et al., 2004; Halatchev et al., 2019; Kübeck et al., 2016; Wostmann et al., 1968). Measurements of the same host-microbiota system, if taken at different timepoints in the circadian cycle of metabolism, could therefore be wrongly interpreted as qualitative shifts in

microbiota function. Consequently, to understand the influence of the microbiota on host energy

metabolism, it is key to register variation over the circadian cycle.

Gnotobiotic mice, colonized with a simplified microbiota made up of defined species, have become

a major tool to identify potential mechanisms of interaction between the microbiota and host (Koh

and Bäckhed, 2020; Mallapaty, 2017; Steimle et al., 2021). Simulations and mathematical analysis

can be applied with higher certainty to a defined consortium and the use of cultivable strains also

allows for *in vitro* analysis of metabolism and behavior of the strains. Such approaches can

generate a mechanistic understanding of how external factors (i.e. diet, infection) act on the

different microbiota members individually and at a community level (Faith et al., 2014; Kovatcheva-

Datchary et al., 2019). A widely used example, the OligoMM12, is a gnotobiotic consortium of 12

cultivable mouse-derived strains representing the major five bacterial phyla in the murine gut

(Brugiroux et al., 2016). It is reproducible between facilities (Eberl et al., 2020) and extensive data

now exists on the metabolism of individual species and their interactions with each other and the

host (Streidl et al., 2021; Weiss et al., 2021; Wotzka et al., 2019; Yilmaz et al., 2021). However,

working with these mice involves hygiene conditions similar to those required to work with germ-

free mice, which poses a technical challenge for the long-term analysis of mouse metabolism. In

particular, standard metabolic cage systems do not permit maintenance of an axenic environment

and moving mice between the isolator systems where such animals are normally bred to IVC-

cage-like systems can be associated with stress and behavioral abnormalities, which also

influence metabolism (Rabasa and Dickson, 2016). This explains why metabolic characterizations

of gnotobiotic mice are so scarce (Halatchev et al., 2019; Yao et al., 2018).

To address the challenge of long-term energy metabolism measurements in gnotobiotic mice, we

have built an isolator-housed metabolic cage system. Based on the TSE PhenoMaster® system,

we can monitor levels of O_2, CO_2 and hydrogen every 24min for up to 8 cages, across two separate

isolators, while maintaining a strict hygienic barrier. This way, longitudinal monitoring of

metabolism can be carried out over periods of several weeks in germ-free and gnotobiotic mice.

In this study, we explored how well a gnotobiotic microbiota replicates the influence of a complex

microbiota on host metabolism. We compared the metabolic profile of germ-free (GF), gnotobiotic

(OligoMM12) and conventionally raised mice (specific-opportunistic-pathogen free, SPF) fed *ad*

libitum with standard chow, using isolator-based indirect calorimetry. Similar to what has been

described before (Krisko et al., 2020; Kübeck et al., 2016), we found no significant differences in

energy expenditure among GF and SPF mice. These results are in contrast to other work

(Bäckhed et al., 2004; Levenson et al., 1969; Li et al., 2019; Wostmann et al., 1968), but the

discrepancies can potentially be explained by the methods applied for normalizing energy-

expenditure data. These methods consider (or not) that germ-free and gnotobiotic mice exhibit

extensive water retention in the cecal lumen which can contribute up to 10% of the total body

weight. When accounting for cecal inert mass, we also observed no significant difference in energy

expenditure in the gnotobiotic OligoMM12 mouse line (Brugiroux et al., 2016). Interestingly,

despite indistinguishable energy expenditure, and indistinguishable energy absorption from food

each day, OligoMM12 mice showed increased fat mass compared to both, germ-free and SPF

mice. Moreover, their circadian patterns of respiratory exchange ratio (RER) and certain

metabolites in liver and plasma, phenocopied SPF mice during the dark phase, but germ-free mice

during the light phase. Our study shows how a reductionist/synthetic microbiota can specifically

recover microbiota function in the dark (active) phase, but not in the light (resting/fasting) phase

of the circadian cycle. This represents a powerful tool for identifying critical microbiota species and

functions needed to support healthy host metabolism throughout the day.

RESULTS

To describe the influence of the microbiota on the metabolic profile of mice, and to be able to compare to published literature, we compared male, adult age-matched (12-14wks old) germ-free (GF), gnotobiotic (OligoMM12) and conventionally raised (SPF) mice, all bred and raised in flexible-film isolators and with a C57BL/6J genetic background. As these data are inherently noisy, and data quality can be improved considerably by pooling data of multiple days and animals, we needed to carry out indirect calorimetry in a system where we could be certain that the risk of microbial contamination is very close to zero. We therefore adapted flexible-film surgical isolators to accommodate a TSE PhenoMaster® system (schematic view in Fig. 1A, picture Fig. 1B) to perform indirect calorimetry measurements of eight single housed mice. Mice were adapted for between 24-36h to the single-housing condition inside isolator-based metabolic chambers before data collection. Variations on O_2, CO_2 and hydrogen, along with food and water consumption, were recorded every 24 min on each metabolic cage.

Body composition in GF, OligoMM12 and SPF mice

After 6-12 days of data recording mice were fasted for 4-5 hours and euthanized (approximately at ZT6 ± 1 hour), and body mass and body composition were measured. As cecal mass (cecal tissue plus its content) is affected by the colonization status (Wostmann et al., 1968), we first assessed the cecal mass in GF, OligoMM12 and SPF and its impact on body mass. We found that cecal mass was inversely correlated to the microbiota complexity, starting at approximately 0.5 g in SPF mice, increasing to around 1.5 g in OligoMM12 mice and reaching 3 g on average in GF mice (Fig.1C).

Given that the cecal mass represents a major proportion of total body mass, accounting for approximately 10% in GF mice (Suppl. Fig.1A), we compared the body mass of GF, OligoMM12

and SPF mice before and after the removal of the cecal mass. Although we found no significant

difference between these mice, we observed a trend in total body mass with GF mice showing

higher mass than SPF and OligoMM12 mice (Fig, 1D). This trend was completely reverted after

removal of the cecum (Fig. 1E).

Measurements of body composition in mice are typically performed using EchoMRI, which yields

data on lean, fat and water mass. As we observed that the cecum represented such a large and

variable fraction of body mass, it was essential to compare EchoMRI read-outs of "lean" and "fat"

body mass before and after removal of the cecum (Suppl. Fig.1B-G). We found a strong correlation

between the total lean mass measured by EchoMRI with and without the cecum (Suppl. Fig 1B),

with a decrease on the lean mass readout by EchoMRI after cecum removal ranging from around

1.3 g in SPF mice up to approximately 2 g in GF mice (Suppl. Fig.1C), i.e., 5 to 10% of the original

lean mass readout (Suppl. Fig. 1D). We concluded that a relatively consistent mass of cecum

seems to be scored as lean mass by EchoMRI, regardless of total cecum size, and all groups of

mice are similarly affected. Therefore, for ease of comparison to published work, we decided to

report and to use lean mass obtained by EchoMRI *before dissection* for definitive energy

expenditure calculations. We observed a trend in lean body mass with GF having a lower lean

body mass than SPF mice, and OligoMM12 mice showing an intermediate phenotype (Fig. 1F).

However, we were underpowered to detect a significant difference between groups, and we

estimate that at least 22 mice per group would be needed to achieve statistical significance by

one-way ANOVA if the current group differences are real – a number that was beyond the scope

of the current study.

Interestingly, EchoMRI fat mass measurements pre- and post-cecum dissection were poorly

correlated in GF mice (Suppl. Fig. 1E) attributable to a rather variable scoring of cecal content as

fat or water. In GF mice, fat mass readout by EchoMRI decreased from 0.2 g to 2.3 g (Suppl. Fig.

1F) when comparing values with and without the cecum. This represented between 5 to 48% of

the original fat mass readout. We also observed a shift towards *higher* fat mass readings in SPF

mice after cecum removal (Suppl. Fig. 1F and Suppl. Fig. 1G); further highlighting the need for

caution in interpreting EchoMRI readouts for fat mass in mice with major differences in intestinal

colonization. Therefore, we proceeded to directly weigh the fat depots accessible to dissection

(interscapular brown adipose tissue, iBAT; and, inguinal and visceral white adipose tissue, iWAT

and vWAT) as a more reliable method. There was no significant difference between GF and SPF

mice in size of the explored fat depots; however, OligoMM12 mice accumulated more fat in all

explored depots than GF mice, and more iBAT and vWAT compared to SPF mice (Fig 1G).

Energy metabolism and energy balance in GF, OligoMM12 and SPF mice

In order to explain the body composition differences observed in the GF, OligoMM12 and SPF

mice, we explored the daily energy expenditure of these mice. Energy expenditure was estimated

using VO_2 and VCO_2 readouts (Meyer et al., 2015) and normalized as described before (Mina et

al., 2018; Speakman, 2013; Tschop et al., 2011) using lean body mass and dissected fat mass.

We could confirm that germ-free mice maintain their germ-free status over at least 10 days of

accommodation in these cages, via culture-dependent and culture-independent techniques (see

Methods).

In contrast to some previous reports (Bäckhed et al., 2004; Levenson, 1978; Li et al., 2019;

Wostmann et al., 1968), but aligned with others (Krisko et al., 2020; Kübeck et al., 2016), we found

no significant difference in daily energy expenditure (Fig. 2A and Fig. 2B) or VO_2 (Suppl. Fig. 2A

and Suppl. Fig. 2B) between GF, OligoMM12 and SPF mice after normalization using a regression

model that included lean body mass and total dissected fat mass as predictive variables. This lack

of difference was also observed when light and dark phases were analyzed separately (Fig. 2B

and Suppl. Fig. 2B). "Classical" normalization procedures (dividing by mass) also showed no

difference between groups when lean body mass, or "total body mass after cecum dissection" was

used for normalization of energy expenditure (Fig. 2C and D) or VO_2 (Suppl. Fig. 2C and D).

Unsurprisingly, we did find a significant difference during the dark phase in energy expenditure

(Fig. 2E) and VO_2 (Suppl. Fig. 2E) between GF and SPF mice if "total body mass" (i.e., including

cecum mass) was used for normalization, as on average 10% of total body mass in GF mice

comprises the non-metabolically active sterile cecal mass (compared to the 2.5% in SPF mice,

Suppl. Fig.1A). Therefore, at least when comparing to the SPF microbiota used in this study,

absence of a microbiota does not result in altered daily energy expenditure in metabolically active

tissues.

We hypothesized that adiposity in the OligoMM12 mice can be due to increased calorie absorption

from food. This was assessed by comparing the daily energy ingestion from food and calorie

excretion in feces, in GF, OligoMM12 and SPF mice. The difference between these two values

estimates the absorbed calories. As reported previously (Wostmann et al., 1983), GF animals

ingested on average between 10-20% more standard chow compared to OligoMM12 and SPF

mice (Fig. 2F). Correspondingly, GF animals also excreted a much larger dry mass of feces, while

OligoMM12 mice produced an intermediate fecal mass and SPF mice excreted the least (Fig. 2G).

Remarkably, energy density of dry feces was lower in GF mice (3.7 Kcal/g) compared to colonized

mice (OligoMM12 and SPF, 4.0 Kcal/g); with the latter showing no difference among them (Fig.

2H). This gap between GF and mice with microbiota can likely be explained by the fact that

although fecal bacteria improve energy release from food, a considerable fraction of that energy

remains stored in the bacteria present in the feces. Assuming certain averaged parameters (dry

mass of a bacterium = 2.26×10^{-13}g / bacteria cell (Dennis and Bremer, 2008), density of bacteria

in feces = 5×10^{11} bacteria cells / g of feces (Barlow et al., 2020), and energy stored in bacteria =

4.58 Kcal/g of dry bacteria mass (Popovic, 2019)); we estimated that the fecal microbiota of

colonized mice can contribute approx. 0.52 Kcal/g of dry fecal mass – slightly more than the energy

density difference between fecal energy density in colonized and germ-free mice.

We then used these values for food intake, fecal dry mass output, and fecal energy density to

estimate energy absorbed from the feces. We found that the higher food consumption in GF mice

(Fig. 2I) almost perfectly counterbalance their corresponding higher energy excretion in feces (Fig.

2J), such that all mice extract around 9 Kcal per day from their food (Fig. 2K). This is consistent

with our measurements of daily energy expenditure by indirect calorimetry (Fig. 2B), although it

fails to explain the observed adiposity in the OligoMM12 mice (Fig. 1G). Unexpectedly, the

OligoMM12 microbiota does not fully recapitulate the ability of a complex microbiota to improve

energy availability from food, as the fraction of all calories extracted from consumed food remains

similar between germ-free and OligoMM12 mice. Given that the gut content of both OligoMM12

and SPF mice is densely colonized, and the fecal energy density is similar; it should be noted that

the lower amount of energy extracted by the OligoMM12 is not so much due to a poorer digestive

capacity of the gnotobiotic gut microbes, but rather that compared to the SPF microbiota, the

calories extracted by the OligoMM12 microbiota are either retained within the microbes or not

converted into compounds that can be taken up by their host (Fig. 2L).

We therefore concluded that daily energy expenditure and daily energy absorption from food

varies only within the range of experimental error intrinsic to indirect calorimetry experiments. At

a fundamental level, food intake therefore seems to be well regulated by microbiota-released

calories. Despite this, OligoMM12 mice have an elevated fat mass. It remains a distinct possibility

that gain of fat mass depends on the cumulative effect of minute differences in energy intake and

energy expenditure that are simply not resolvable in our system. Nevertheless, in order to gain a

deeper insight into underlying mechanisms we carried out a series of more detailed analyses of

metabolism.

Circadian changes in RER and microbiota-derived hydrogen and short-chain fatty acids

(SCFA)

The tight control of eating behavior based on absorbed calories with or without a microbiota is even more impressive when one considers the difference in substrates being used for oxidative metabolism and the timing of their arrival in the circulation with relation to eating. Respiratory exchange ratio (RER, the ratio of CO_2 produced per O_2 consumed) is widely used as an informative proxy for substrate utilization (i.e., glucose or fatty acids) for oxidation in tissues. In general, GF mice have a lower RER compared to SPF in both light and dark phases, potentially explained by increased fat/decreased glucose metabolism in GF mice (Fig 3A). Therefore, appetite regulation functions despite very different metabolic substrate availability in GF and SPF mice. Intriguingly, OligoMM12 mice displayed an RER at the GF level during the light phase (when mice typically sleep and fast) but raised up to SPF levels during the dark phase (i.e., when mice are active and eating). It therefore appears that the OligoMM12 microbiota better recapitulates the microbiome effects on the host during the dark (active) phase when food-derived carbohydrates are abundant in the large intestine, but not when mainly host-derived carbon sources are available in the large intestine (Fig. 3A). These changes in RER are not related to differences in feeding patterns as all mice have a similar food intake during the periods in which their RERs differ the most (Fig. 3B).

Differences in RER provided a clue that there could be differences in energy storage in mice with different microbiota status. Microbial fermentation products, including short-chain fatty acids (SCFA) and lactate, can be directly used as energy and carbon-sources by the murine host, and are generated by the microbiota via processes that liberate molecular hydrogen. We therefore quantified hepatic concentrations of glycogen, and cecal concentrations SCFA, at Zeitgeber 5 (ZT5, 5h into the light phase) and 16 (ZT16, 4h into the dark phase). Hydrogen was measured continuously during the circadian cycle.

Hepatic glycogen levels show a circadian rhythm, which usually peaks early during the transition between dark to light phase (ZT2-4), and drops to its minimum during the early hours of the dark

phase (ZT14-16) in nocturnal rodents (Doi et al., 2010; Ishikawa and Shimazu, 1976). We found

similar accumulation of hepatic glycogen in GF, OligoMM12 and SPF mice at ZT5; however, GF

and OligoMM12 mice drop to lower levels of hepatic glycogen compared to SPF mice at ZT16 (Fig

3C). This differential pattern in GF/OligoMM12 compared to SPF mice may indicate that, although

they can equally fill up hepatic glycogen storages at the end of the dark phase, GF and OligoMM12

show faster depletion of hepatic glycogen during the light phase.

Hydrogen, a byproduct of fiber fermentation by the microbiota, was also measured in the exhaust

air of the metabolic cages. We found a clear circadian pattern in hydrogen production between

OligoMM12 and SPF mice (Fig. 3D). Hydrogen levels in OligoMM12 and SPF mice decreased

down to the limit of blank (GF level as reference) during the light phase, to later peak after food

intake resumes during the dark phase. In addition, OligoMM12 mice showed a higher production

of hydrogen than SPF mice during the dark phase even after regression-based normalization by

cecal mass, which is larger in OligoMM12 mice (Fig. 3D). Therefore, the higher dark phase

hydrogen measurements in the OligoMM12 mice are not just the result of a larger cecal mass, but

of the OligoMM12 microbiota producing hydrogen at a higher rate per cecal content mass than

the SPF microbiota.

SCFA are the other major output of bacterial fermentation in the large intestine, as well as being

key bioactive compounds produced by the large intestinal microbiota. SPF mice showed the

highest cecal concentrations of all SCFA during both the light phase (butyrate and propionate)

and dark phase (acetate, butyrate, and propionate). Interestingly, OligoMM12 mice showed only

20-50% of the SCFA concentrations observed in SPF mice but instead showed high production of

lactate during the dark phase (Fig. 3E). In germ-free mice, all analyzed metabolites had levels

below the limit of blank except for lactate, which could correspond to host-produced L-lactate

(Zarrinpar et al., 2018) (our assay is not able to differentiate the enantiomers). As the total mass

of cecum content is widely different among GF, OligoMM12 and SPF mice, we also estimated the

total quantity of each compound in the cecal content by multiplying the concentration (Fig. 3E) by

the cecal mass for each group (Fig. 1C) while propagating the uncertainty of each measurement,

as a way of representing the total availability of SCFA for the different hosts. This transformation

has quite a major impact on how these data can be interpreted: when taking cecal mass into

account, OligoMM12 mice have considerably higher levels of acetate during the light and dark

phase and of propionate during the light phase than SPF mice, while butyrate levels remain low.

There is also an increased abundance of lactate and succinate in the OligoMM12 cecum content

(Fig. 3F). Although we cannot directly link these microbial metabolites to the phenotype of the

OligoMM12 mice, this underlines the major differences in metabolite profiles in the large intestine

when comparing germ-free, gnotobiotic and SPF mice. High lactate production by the microbiome

certainly warrants further study for potential metabolic effects on the host.

Circadian changes in liver and plasma metabolites in GF, OligoMM12 and SPF mice

Finally, to increase our metabolic resolution, we applied UPLC-MS to perform untargeted

metabolomics in the liver and plasma during the light (Zeitgeber 5) and dark phase (Zeitgeber 16)

in GF, OligoMM12 and SPF mice. Hierarchical clustering (Fig. 4A and Fig. 4C) and PCA analysis

(Fig. 4B and Fig. 4D) separate the SPF, OligoMM12 and GF groups during both light and dark

phases. Correlating to what we observed in the RER during the light phase, GF and OligoMM12

cluster closer and are clearly separated from the SPF in the light phase PCA analysis for both liver

(Fig. 4B) and plasma samples (Fig. 4D). During the dark phase, although OligoMM12 mice

differentiate from GF, they do not recapitulate the metabolome of SPF mice neither in liver (Fig.

4B) or plasma samples (Fig. 4D).

We used the package MetaboAnalystR (Chong and Xia, 2018) to identify compounds that are

significantly different in pair comparisons between OligoMM12 mice and their GF and SPF

counterparts by untargeted peak extraction and map them into metabolic pathways using the

KEGG database. We found several pathways differentially enriched when OligoMM12 mice were

compared to GF or SPF counterparts during the light and dark phase in liver (Fig. 4E) and plasma

(Fig. 4F), including amino acid, bile acids, and fatty acid metabolism. Additionally, we selected

compounds that belong to these differentially enriched pathways or have been previously

identified to have circadian changes in obese patients (Nowak, 2021), confirmed their structure by

using chemical standards, and performed a targeted peak extraction for a more precise

comparison among groups (Suppl. Table 1). For certain metabolites the OligoMM12 microbiota

produce an intermediate phenotype between GF and SPF mice, e.g., a subset of bile acids, in

liver (Suppl. Fig 3A) and plasma (Suppl. Fig 3B).

DISCUSSION

Since the early days of nutritional studies, there has been a clear interest to understand the role of microbiota in host morphology, physiology and nutrition (Gordon and Pesti, 1971; Levenson, 1978). Pioneering work comparing germ-free animals with conventionally raised or gnotobiotic counterparts already described differences in food intake, energy extraction from diet and energy expenditure by indirect calorimetry (Levenson et al., 1969; Wostmann et al., 1983). More recently, researchers have explored the effect of specific microbiota communities and how they influence energy metabolism and body composition in the host (Ridaura et al., 2013; Suárez-Zamorano et al., 2015; Turnbaugh et al., 2006).

Here, by carefully checking the validity of different measurement types, we found no significant difference in lean body mass among germ-free (GF), gnotobiotic (OligoMM12) and conventionally raised (SPF) mice. Interestingly, there was a significant increase in fat depots in OligoMM12 mice compared to GF and SPF animals. Previous studies diverged on the effect of microbiota on fat mass accumulation during conventional/low-fat diet feeding; reporting either increased fat mass in SPF mice (Bäckhed et al., 2004) or no difference compared to GF mice (Kübeck et al., 2016). However, it should be noted that there can be huge differences between SPF microbiota within and between animal facilities. GF mice transplanted with microbiota derived from obese donors accumulated more fat mass compared to those transplanted with microbiota derived from lean donors (Halatchev et al., 2019; Ridaura et al., 2013; Turnbaugh et al., 2006), with correlates identified to individual species/strain abundance (Woting et al., 2014, 2015). SPF microbiota matching more closely to those from obese donors could therefore be expected to give differing results to ours. In contrast, minimal microbiota communities can be perfectly replicated across sites, and can help to clarify the complex processes linking microbiota and host metabolism (Becker et al., 2011). Further exploration of the metabolic effects of the OligoMM12 microbiota community, and extended versions thereof, has potential to clarify if specific strains, species or

functional classes (Schmidt et al., 2018) are sufficient and necessary to drive the development of increased fat depots in these mice.

Our analysis compared energy expenditure in GF, OligoMM12 and SPF mice and found no significant difference among these mice. Normalization of mass-dependent variables by a per-mass (or allometric transformation) ratio has been recognized as a common source of controversy (Packard and Boardman, 1999; Tanner, 1949; White and Seymour, 2005), especially with large changes in body mass composition (Butler and Kozak, 2010; Kaiyala and Schwartz, 2011), and there have been several publications calling for the use of better statistical methods (Arch et al., 2006; Fernández-Verdejo et al., 2019; Tschop et al., 2011). We found that, when the cecal content mass is very different among groups, using total body mass for normalization introduces a considerable bias in normalized energy expenditure estimation. Interestingly, it was long-ago observed that surgical removal of the cecum equalized the oxygen consumption between germ-free and conventional rats; as well as other measurements normalized by total body mass (Wostmann et al., 1968). With normalization using linear regression models based on lean-mass and fat-mass (Mina et al., 2018), we and others found no significant differences in energy expenditure by indirect calorimetry between GF and SPF mice under standard chow diet conditions (Krisko et al., 2020; Kübeck et al., 2016; Li et al., 2021). Importantly, we suggest cautious interpretation of lean and fat mass readouts obtained by EchoMRI when comparing mice with major differences in intestinal content composition. We are also keen to point out the more general limitations of our observations: only one gnotobiotic microbiota and one SPF microbiota were analyzed, and our conclusions pertain exclusively to these. We in no way exclude the possibility that some microbiota constituents or conformations *can* influence host energy expenditure (Halatchev et al., 2019) and/or body composition (Ridaura et al., 2013; von Schwartzenberg et al., 2021; Turnbaugh et al., 2006). In addition, it should be noted that indirect

calorimetry is an inherently noisy data type, and small differences in daily energy expenditure are

impossible to resolve via this technique (Corrigan et al., 2020; Fernández-Verdejo et al., 2019).

Nevertheless, the lack of measurable difference in energy expenditure between GF, OligoMM12

and SPF mice is aligned with our finding that the amount of energy obtained by *ad libitum* food

intake was also remarkably similar among the groups. GF mice seem to accurately compensate

the lower capacity of energy extraction from diet by increasing food intake. While this seems

generally to be in agreement with models that described the regulation of appetite (and therefore

energy intake) by the basal energy requirement of the individual (MacLean et al., 2017; Stubbs et

al., 2018), it remains surprising given the discrepancy in the types of substrates available for

oxidative metabolism in colonized and germ-free mice. Although germ-free mice have a longer

total gastrointestinal transit time than SPF mice (Touw et al., 2017), very little calorie absorption

from food can occur after ingested food reaches the cecum of a germ-free mouse, whilst an SPF

mouse will release usable energy from their food via microbial fermentation for several more hours

in the cecum and colon, generating a major time-difference in the absorption of calories after

eating in germ-free and SPF animals. This compensation seems also to function in mice colonized

with the OligoMM12 microbiota, where despite robust microbial fermentation (read out as

hydrogen and fermentation product production) and identical fecal energy density to SPF mice,

energy recovery from ingested food is poor due to the volume of feces shed. A clear conclusion

from these observations is that microbiota-dependent changes in metabolic substrates, and timing

of calorie absorption, are well integrated in the murine central regulation of appetite over the

course of a day (Fetissov, 2017).

Despite this broadly successful regulation of food intake and energy expenditure, at the molecular

level, major differences were observed between the mice with different microbiota. In cecum

content, the OligoMM12 microbiota was associated with a predominance of succinate and lactate,

at the expense of propionate and butyrate. In complex microbiotas, lactate is typically further

metabolized to butyrate by specific firmicutes (Belenguer et al., 2011; Duncan et al., 2004; Flint et

al., 2015), which may be lacking or insufficiently abundant in the OligoMM12 mice. As lactate can

inhibit lipolysis in adipocytes (Cai et al., 2008; Liu et al., 2009), this raises an interesting theme for

follow-up studies to define the role of microbiota-derived lactate in host metabolism. In line with

the RER data, we also observed that the liver and plasma metabolite profiles of OligoMM12 mice

clustered closer to GF mice than to SPF mice. Although certain compounds were differentially

represented between OligoMM12 and GF or SPF mice, we could not replicate earlier observations

(Claus et al., 2008; Mardinoglu et al., 2015). This discrepancy may be explained by the location

of the sample collection (portal vein vs. liver/peripheral plasma) or the lack of power of our sample

size to detect significant differences having both circadian phase and microbiota as covariables.

Interestingly, OligoMM12 had a bile acid profile closer to GF than SPF mice, for example showing

GF-levels of β-murocholic acid, the predominant bile acid in the liver of GF mice (Sayin et al.,

2013). Follow up studies with manipulation of the OligoMM12 microbiota or metabolic interventions

are a promising tool to pull apart the circadian effects on RER, the influence of an unusual

fermentation product profile, and other more subtle metabolic changes on overall metabolic health

of the murine host.

In conclusion, our study showed that isolator-based indirect calorimetry is possible and allows

detailed analysis of the metabolism of germ-free and gnotobiotic mice in real-time. Data generated

with this system demonstrated that microbiota-released calories are well integrated in host energy

balance. Intriguingly, the OligoMM12 gnotobiotic microbiota accumulate fat mass, and displays a

GF-like RER during the light phase but an SPF-like RER during the dark phase. The OligoMM12

mice have a skewed profile of cecal fermentation products during both light and dark phases,

including very high concentrations of lactate, as well as alterations in bile acid, fatty acid and amino

acid metabolism in blood and liver. This reveals the full potential for gnotobiotic microbiota

communities to investigate the mechanisms underlying the influence of microbiota composition

and function on host metabolic health. As microbial dysbiosis is associated with a range of human

diseases, circadian analysis of energy balance represents a crucial tool in the mining of

microbiome data for therapeutic and diagnostic purposes.

ACKNOWLEDGEMENTS

We would like to thanks to Sven Nowok, Thomas Fehr, Andre Galhano and Susanne Freedrich for their support in the establishment of the gnotobiotic metabolic phenotype facility in the ETH Phenomic Center. Also, we thank Maria L. Balmer for her comments and suggestions for the manuscript. The gnotobiotic metabolic phenotype facility was initially funded by the FreeNovation grant (Novartis). E.S. and W-D.H. are supported by the NCCR Microbiomes, a research consortium financed by the SNF. E.S. acknowledges the support of the Swiss National Science Foundation (40B2-0_180953, 310030_185128), European Research Council Consolidator Grant, and Gebert Rüf Microbials (GR073_17) and the Botnar Research Centre for Child Health Multi-Invesitigator Project 2020. This project is part of "Zurich Exhalomics", a flagship project of "Hochschulmedizin Zürich".

Author Contributions

Conceptualization, D.H., W-D.H., C.W. and E.S.; Methodology, D.H., J.L., W.S., R.Z. C.W. and E.S.; Formal Analysis, D.H., J.L, W.S., T.G. and M.A.; Investigation, D.H., J.L., W.S. and S.N.; Resources, B.S., R.Z., W-D.H., C.W. and E.S.; Writing – Original draft, D.H. and E.S.; Writing – Review and Editing D.H., J.L., W.S, T.G, M.A, S.N., A.J.M., B.S., R.Z., W-D.H., C.W and E.S.; Visualization, D.H., W-D.H., C.W. and E.S.; Supervision, W-D.H., C.W. and E.S.; Funding acquisition, W-D.H., C.W., A.J.M, and E.S.

Declaration of Interest

The authors declare no competing interests.

FIGURES TITLES AND LEGENDS

Figure 1: OligoMM12 mice have increase fat mass compared to GF mice and SPF C57B6/J mice. (A) Schematic representation of isolator-based indirect calorimetry system, with a TSE PhenoMaster® calorimeter connected to two flexible surgical isolators with four metabolic cages each. (B) Pictures of isolator-based indirect calorimetry system inside the facility. (C) Cecal mass (tissue including luminal content). (D) Total body mass at the end of the experiment and before cecum removal. (E) Total body mass after cecum removal. (F) Lean body mass acquired by EchoMRI before cecum removal (N of mice per group with EchoMRI and indirect calorimetry measurements: GF = 12, OligoMM12 = 8, SPF = 11). (G) Fat mass from interscapular brown adipose tissue (iBAT), inguinal white adipose tissue (iWAT) and visceral white adipose tissue (vWAT). Number of mice per group in all figures unless otherwise specified: GF = 16, OligoMM12 = 12, SPF = 11. p-values obtained by Tukey's honest significance test.

Figure 2: Energy metabolism in GF, OligoMM12 and SPF C57B6/J mice. (A) Linear regression of energy expenditure and lean body mass based on EchoMRI during light and dark phase. Each colored vertical line represents energy expenditure measurements during the experiment for one mouse. (B) Energy expenditure during 24h period, or during the 12h light or dark phase. Values represent area-under-curve normalized by regression-based analysis using lean body mass obtained by EchoMRI and dissected fat mass. (C, D, E) Energy expenditure values obtained by "classical" ratio-based normalization methods (dividing energy expenditure values per phase by mass). (C) Area-under-curve after normalization by total mass after cecal dissection. (D) Area-under-curve after normalization by lean body mass (EchoMRI). (E) Area-under-curve after normalization by total body mass before cecal dissection. (F) Average daily food intake per mouse. Mice represented in this figure include those that underwent long-term indirect calorimetry (Fig. 3) and mice that only contribute to daily fecal pellet quantification/bomb calorimetry. (N of mice per group: GF = 24, OligoMM12 = 19, SPF = 10) (G) Dry fecal output per mouse collected during a

24h period. (N of mice per group: GF = 12, OligoMM12 = 8, SPF = 4) (H) Energy content of dry

fecal output by bomb calorimetry. (N of mice per group: GF = 21, OligoMM12 = 11, SPF = 11). (I,

J, K, L) Estimation energy metabolism parameters. Number represented estimate mean value ±

1.96*combined standard uncertainty from measurements used for calculations. (I) Estimated daily

energy input (food intake* 3.94 Kcal/g). (J) Estimated daily energy excretion (daily fecal dry

mass*fecal energy content). (K) Estimated daily energy extraction (daily energy input – daily

energy excretion). (L) Estimated energy extraction from food as percentage of energy input ((daily

energy input - daily energy excretion)/daily energy input*100). Note that calculations in L, N and

M are per mouse and are not normalized to body mass. Number of mice per group in all figures

unless otherwise specified: GF = 9, OligoMM12 = 8, SPF = 10. p-values obtained by Tukey's

honest significance test.

Figure 3: Circadian changes in Respiratory Exchange Ratio (RER), microbiota-derived

hydrogen and short-chain fatty acids (SCFAs). (A) Comparison of circadian changes in RER

among GF, OligoMM12 and SPF C57B6/J mice. RER curves obtained by smoothing function of

data obtained every 24min per mouse over 10 days. Mean RER during the light phase (Zeitgeber

0-12) and dark phase (Zeitgeber 12-24). (B) Cumulative food intake during described ZT periods.

Mice included in this analysis are those that underwent long-term indirect calorimetry, and they

are a subset of the mice represented in Fig. 2F (C) Hepatic glycogen and triglyceride concentration

in samples obtained at Zeitgeber 5 and 16 (N=3 per group). (D) Hydrogen production, curves

obtained by smoothing function of data obtained every 24min per mouse. Area-under-curve after

regression-based normalization by cecal mass during the light and dark phase (N of mice per

group: OligoMM12 = 11, SPF = 10). (E) Concentration of short-chain fatty acids (acetate, butyrate,

propionate) and intermediate metabolites (lactate, succinate) products in cecal content (N=3 per

group). (F) Estimation total amount of short-chain fatty acids and intermediate metabolites by

multiplying measured concentration values by the cecal mass of the group. Number represented

estimate mean value ± combined standard uncertainty from measurements used for calculations.

Number of mice per group in all figures unless otherwise specified: GF = 13, OligoMM12 = 12,

SPF = 10. p-values obtained by Tukey's honest significance test.

Figure 4. Metabolic profile comparison of GF, OligoMM12 and SPF C57B6/J mice by

UPLC/MS. (A and C) Heatmaps of metabolites identified by untargeted UPLC/MS during the light

phase (Zeitgeber 5) and dark phase (Zeitgeber 16) in (A) liver and (C) plasma. (B and D) PCA

analysis of metabolites identified by untargeted UPLC/MS during the light phase (Zeitgeber 5) and

dark phase (Zeitgeber 16) in (B) liver and (D) plasma. (E and F) Metabolic pathways identified in

the KEGG PATHWAY database; red dots represent pathways containing compounds differentially

enriched in (*top*) OligoMM12 vs. GF and (*bottom*) OligoMM12 vs. SPF comparisons. Samples

obtained during the light phase (Zeitgeber 5) and dark phase (Zeitgeber 16) in (E) liver and (F)

plasma. Number of mice per group in all figures: GF = 3, OligoMM12 = 3, SPF = 3.

METHODS

Animals

We used C57B6/J male mice aged between 12-14 weeks. We compare germ-free mice (GF), with a 12-strain gnotobiotic microbiota (Brugiroux et al., 2016) (OligoMM12) and specific-pathogen free mice (SPF). GF and OligoMM12 mouse lines are bred and maintained in open-top cages within flexible-film isolators, supplied with HEPA-filtered air, and autoclaved food and water ad libitum. As we are aware that housing conditions may influence behavior and potentially metabolism, we also bred and maintained a SPF colony under identical conditions inside a flexible-film isolator specifically for this study, such that all mice experienced identical living conditions, food, and water. Mice were adapted for between 24-36h after transfer from the breeding isolators to the isolator-based metabolic chambers. For long term experiments, mice were periodically re-housed in couples for short periods of times to avoid stress of extended single-housing conditions. In all cases, animals were maintained with standard chow (diet 3807, Kliba-Nafag, Switzerland) and autoclaved water. Germ-free status was confirmed at the end of the long-term experiments by culturing cecal content in sterile BHIS media in aerobic and anaerobic conditions for a week. In addition, cecal content was frozen at -20°C for a week, then stained with SYBR Gold and assessed by bacterial flow cytometry (Moor et al., 2016) using similarly processed SPF mice cecal content as positive control for the presence of bacteria. All experiments were conducted in accordance with the ethical permission of the Zürich Cantonal Authority.

Indirect calorimetry

The isolator-housed TSE PhenoMaster® system allows instantaneous measurements of oxygen, carbon dioxide and hydrogen levels as well as total feed and water consumption while keeping a strict hygiene level of control. The metabolic isolator system consists of an adapted set of two

flexible-film surgical isolators, each of them housing four metabolic cages from the TSE

PhenoMaster® system (TSE Systems, Germany). Room air is pulled into the isolator by a vacuum

pump passing through a double set of HEPA filters. Then, each cage is connected via a second

HEPA filter through the back wall of the isolator to the CaloSys setup, which pulls sterile air from

the isolator into the cages using negative pressure. Air coming from the cages is dehumidified at

4°C and sequentially passed by a Sensepoint XCD Hydrogen gas analyzer (Honeywell Analytics,

Hegnau, Switzerland) and standard oxygen and carbon dioxide censors provided in the TSE

PhenoMaster® system. A two-point calibration of all analyzers using reference gases was

performed within 24 h before each animal experiment. Data was recorded using a customized

version of the TSE PhenoMaster® software modified to integrate hydrogen measurements.

For indirect calorimetry measurements, the animals were transported in pre-autoclaved, sealed

transport cages from the breeding isolators into the metabolic isolator system. Mice were single

housed and adapted for between 24-36h before starting recording measurements to ensure

proper access to food and water as well as account for initial exploratory behavior. Mice were kept

up to 10 days at a stable temperature (21-22°C) with *ad libitum* availability of standard chow and

water. The days were divided into a dark and light period of 12 hours each. In this study, we kept

the air flow of 0.4 L/min and recorded individual cage data (gases production and food/water

consumption) every 24min (time set per cage for measurement stabilization 2.5min). In long

experiments, mice were periodically pair-housed for 24h to prevent stress due to prolonged single

housing.

Body composition measurements

At the end of the experiment, mice were fasted for 4 hours (Zeitgeber 1 till 5) before for body

composition measurements. We used magnetic-resonance whole-body composition analyzer

(EchoMRI, Houston, USA) to analyze mice body composition (lean and fat mass). Then, mice

were euthanized using CO_2 according to approved protocols. Total body mass was obtained by

weighing the full carcass and cecum was dissected and weighed by one investigator (DH). For a

set of mice, we remeasured body composition by EchoMRI after cecum removal. Finally, fat

depots were dissected from all mice by one investigator (W.S.) that was blinded to the hygiene

status of the mice. Interscapular brown adipose tissue (iBAT), inguinal white adipose tissue (iWAT)

and visceral white adipose tissue (vWAT) were sampled and weighted.

Food intake, fecal samples and bomb calorimetry

Daily food intake was obtained as the mean value of food intake recorded by the TSE

PhenoMaster® system during the course of the experiment. In addition to the mice reported in the

indirect calorimetry experiments, we also collected food intake data from a set of selected

experiments in which we collected fecal pellets produced during 24h. For daily fecal excretion

measurements, we cleaned up the bedding in the cage and replaced it for a clean and reduced

amount of bedding. After 24h, we collected the mix of bedding and fecal pellets. Fecal pellets were

manually collected from the bedding, transferred to 15ml tubes and stored at -20°C until bomb

calorimetry. Before bomb calorimetry, fecal samples were freeze dried in a lyophilizer overnight

(ALPHA 2-4 LDplus, Christ, Germany) and dry mass recorded. We used a C1 static jacket oxygen

bomb calorimeter (IKA, Germany) to quantify the residual energy present in these dry fecal pellets,

using approximately 0.2-0.5g of material. Energy content was normalized to grams of dry fecal

pellets.

Metabolomics by UPLC/MS

Sample obtention and preparation

Approximately at Zeitgeber 5 and 16, three mice of each group were euthanized, and liver and

plasma samples collected. To minimize variations among mice, individual mice were euthanized

with CO_2 and sampled as fast as possible. Blood was obtained by cardiac puncture, collected in

lithium heparin coated tubes, and kept on ice for further processing. Mice were perfused with PBS

and liver samples were obtained by dissection of the lower right lobe, collected on an 2ml

Eppendorf tube and flash frozen in liquid nitrogen. Finally, between 60-80 mg of cecal content was

collected in a 2ml Eppendorf tube and flash frozen in liquid nitrogen. After samples all samples

were obtained, blood samples were centrifuged 8000rcf for 5min, supernatant collected, and flash

frozen in liquid nitrogen. Samples were kept at -80°C until preparation for UPLC/MS.

Short chain fatty acid quantification by UPLC/MS

Samples were first homogenized in 70%-isopropanol (1 mL per 10 mg sample), centrifuged.

Supernatants were used for SCFA quantification using a protocol similar to previously described

(Liebisch et al., 2019). Briefly, a 7-points calibration curve was prepared. Calibrators and samples

were spiked with mixture of isotope-labeled internal standards, derivatized to 3-

nitrophenylhydrazones, and the derivatization reaction was quenched by mixing with 0.1% formic

acid. Four µL of the reaction mixture was then injected into a UPLC-MS system, [M-H]- peaks of

the derivatized SCFAs were fragmented and characteristic MS2 peaks were used for

quantification.

Untargeted UPLC/MS

Samples were thawed on ice. Serum samples were diluted with 90% methanol in water with a

volumetric ratio of 1:7, incubated for 10 min on ice for allowing protein to precipitate. Liver samples

were mixed with 75% methanol in water (2 mL/ 100 mg liver), homogenized using a TissueLyser

(Qiagen, Germany) at 25 Hz for 5 min. The result mixtures were centrifuged at 15,800 g, 4 °C for

15 min. 100 μL of the supernatants were filtered with 0.2 μm reversed cellulose membrane filter

and transferred to sample vials and used for UPLC/MS analysis with an ACQUITY UPLC BEH

AMIDE column (1.7 μm, 2.1 × 150 mm, Waters). Another 400 μL of the supernatants were then

lyophilized and resuspended in 80 μL 5% methanol in water, sonicated, filtered, and used for

UPLC/MS analysis with an ACQUITY UPLC BEH C18 column (1.7 μm, 2.1 × 150 mm, Waters,

RP column).

An ACQUITY UPLC system (I-Class, Waters, MA, USA) coupled with an Orbitrap Q-Exactive Plus

mass spectrometer (Thermo Scientific, San Jose, CA) were used for UPLC/MS analysis. For the

AMIDE column a flow rate of 400 μL/min was used with a binary mixture of solvent A (water with

0.1% formic acid) and solvent B (acetonitrile with 0.1% formic acid). The gradient starts from 1%

of A, then gradually increases to 70% of A within 7 min. Then a 1% of A is kept for 3 min. The

column was kept at 45 °C and the autosampler at 5° C.

For the RP column, the flow rate was set to 240 μL/min using a binary mixture of solvent A (water

with 5 % methanol and 0.1 % formic acid) and solvent B (methanol with 0.1 % formic acid). The

gradient starts from 95% of A, then gradually decreases to 5% of A within 10 min. A 100% solvent

of B is kept for 2 min, then a 100% of A is kept for 2min to restore the gradient. The column was

kept at 30 °C and the autosampler at 5 °C.

The MS was operated at a resolution of 140,000 at m/z = 200, with automatic gain control target

of $2x10^5$ and maximum injection time was set to 100ms. The range of detection was set to m/z 50

to 750. Untargeted MS data was extracted from raw MS files by using XCMS (Smith et al., 2006)

in R (v3.6.1), and then subject to pathway enrichment by using MetaboAnalystR (Chong and Xia,

2018).

Compound identification and targeted peak extraction

Chemical standards of selected compounds were diluted to 10 µg/mL and were analyzed using

the UPLC/MS methods described before. Identification was done by comparing retention time and

MS2 in liver/plasma samples with the chemical standards (Nowak, 2021). After confirming the

chemical identities of the compounds, targeted peak extraction was done using Skyline (v21.1)

(Adams et al., 2020).

Data Analysis

Data quality control

To facilitate analysis across different experimental runs, all times were converted into Zeitgeber

time (ZT; [h]), where 0-12 represents the light phase and 12-24 represents the dark phase. Any

datapoint taken before the start of the first occurrence of ZT=0 was discarded. To account for

faulty measurements caused by measurement imprecision, equipment malfunction or other

disruptive events, datapoints were removed from the raw datasets according to criteria based on

statistical and biological arguments. Food consumption values of 0.01 g during the 24min intervals

were considered as measurement noise and discarded. Negative values for food and water

consumption, as well as oxygen (dO_2) and carbon dioxide (dCO_2) differentials between the

measurement chambers and the reference chamber were also considered as measurement noise

and discarded. For the remaining subsets of measurements from the individual mice, we cleaned

up outlier measurements in food and water intake by eliminating values greater than 75[th] percentile

+ 1.5 times interquartile range. Potential sources for outlier measurements in food and water

consumption observed included leaky water bottles and loss of food pellets during mice husbandry

procedures. A similar approach was used to eliminate outliers from dO_2 and dCO_2 values below

25[th] percentile - 1.5 times interquartile range. Potential sources for outlier measurements in gas

differentials included inappropriate sealing of individual metabolic cages or clogging of pre-

analyzer filters. Oxygen consumption (VO_2) and CO_2 production (VCO_2) was calculated using dO_2

and dCO_2 and the Haldane transformation as described before (Arch et al., 2006). Energy

expenditure was estimated from dO_2 and dCO_2 using Weir's approximation (Weir, 1949). As one

of the study objectives is to explore circadian patterns, if more than 20% of datapoints had to be

removed from a particular day for a particular mouse, all other datapoints from that subset were

discarded as well. After the cleanup process described above, the data from all different

experiment runs were pooled together for further analysis. The above processes lead to a

reduction in dataset size from 10472 to 9453 entries.

Statistical analysis

From the resulting dataset, energy expenditure over a certain period was calculated as the area

under the curve (trapezoid interpolation) of instantaneous values obtained during the 24min

measurements intervals. Food intake values calculated over a certain time are always cumulative.

To compare different mice in the above variables, variations in body mass and composition

between individuals need to be accounted for. As suggested in several publications (Fernández-

Verdejo et al., 2019; Speakman, 2013; Tschop et al., 2011), this was done by regression-based

analysis of covariance (ANCOVA). As such, a linear regression is performed on energy

expenditure as a function of lean body mass and fat depots mass, with the microbiota group as a

qualitative covariate. Then, each individual value is replaced by the sum of the corresponding

residual and the energy expenditure predicted by the linear model using the average lean body

and fat depot mass (calculated over all groups). Hydrogen production (difference in hydrogen

concentration between the measurement chambers and the reference chamber) was adjusted in

analogous fashion, using cecal mass (as a proxy for total gut microbiota mass) as a predictor.

For variables where the continuous evolution during the circadian cycle is of interest (RER, gross

hydrogen production), values were averaged at each time point for each individual. A generalized

additive model was used to fit a smooth line to these averages using a cubic penalized regression

spline (using R function mgcv::gam with formula y ~ s(x; bs = "cs")).

For estimating derived variables (i.e., daily energy excretion) we used the R package "errors"(Ucar

et al., 2019). This package links uncertainty metadata to quantity values (i.e., mean "daily fecal

dry mass excretion", mean "fecal energy content") and this uncertainty is automatically propagated

when calculating derived variables (i.e., "daily energy excretion" = "daily fecal dry mass excretion"

x "fecal energy content"). Uncertainty is treated as coming from Gaussian and linear sources and

propagates them using the first-order Taylor series method for propagation of uncertainty.

Hierarchical clustering and heatmap visualization were produced using the R package "pheatmap"

using Pearson correlation as distance measure for clustering and Ward's minimum variance

method using an algorithm that includes Ward's criterion (Murtagh and Legendre, 2014). For the

Principal Component Analysis, we used the *prcomp* function which is present in built-in R stats

and the R package "factoextra" for visualization.

All group comparisons were analyzed by ANOVA and Tukey's honest significance test. For

comparisons of metabolites identified by targeted peak extraction among groups, area values

were log2 transformed before the statistical test.

Resource availability

Lead contact

Any further communication, including those related to resource sharing, may be directed to and

fulfilled by the lead contact Emma Slack (emma.slack@hest.ethz.ch).

Materials availability

657 This study did not generate new unique reagents.

658 *Data and code availability*

659 Source data for Fig. 1, 2 and 3, and Suppl. Fig. 1 and 2 are available in the Supplementary

660 Information. Source data for Fig. 4 and Suppl. Fig. 3 and the datasets and code used for all figures

661 in this publication are made available in a curated data archive at ETH

662 Zurich (https://www.research-collection.ethz.ch/handle/20.500.11850/501168) under the DOI

663 https://doi.org/10.3929/ethz-b-000501168.

	SUPPLEMENTAL INFORMATION TITLES AND LEGENDS

Supplementary Figure 1: Cecal mass interferes with fat mass estimation by EchoMRI. (A)

Cecal mass (tissue including luminal content) as percentage of total body mass (N of mice per

group: GF = 16, OligoMM12 = 12, SPF = 11) (B) Lean body mass estimated by EchoMRI with and

without cecum. Equations show simple linear regression for estimating lean mass without cecum

based on lean mass with cecum; in brackets adjusted-R squared. (C) Lean mass variation after

cecum removal. (D) Lean mass variation after cecum removal as percentage of lean mass before

cecum removal. (E) Fat body mass estimated by EchoMRI with and without cecum. Equations

show simple linear regression for estimating fat mass without cecum based on fat mass with

cecum; in brackets adjusted-R squared. (F) Fat mass variation after cecum removal. (G) Fat mass

variation after cecum removal as percentage of lean mass before cecum removal. Number of mice

per group in all figures unless otherwise specified: GF = 13, OligoMM12 = 11, SPF = 15. p-values

obtained by Tukey's honest significance test.

Supplementary Figure 2: Cecal mass interferes with normalization of VO$_2$. (A) Linear

regression of VO$_2$ and lean body mass (EchoMRI) during light and dark phase. Each colored

vertical line represents energy expenditure measurements during the experiment per mouse. (B)

VO$_2$ during 24h period, or during the 12h light or dark phase. Values represent area-under-curve

normalized by regression-based analysis using lean body mass obtained by EchoMRI and

dissected fat mass (C, D, E) VO$_2$ values obtained by "classical" ratio-based normalization methods

(dividing energy expenditure values per phase by mass). (C) Area-under-curve after normalization

by total mass after cecal dissection. (D) Area-under-curve after normalization by lean body mass.

(E) Area-under-curve after normalization by total body mass before cecal dissection. Number of

mice per group in all figures unless otherwise specified: GF = 9, OligoMM12 = 8, SPF = 10. p-

values obtained by Tukey's honest significance test.

Supplementary Figure 3. Metabolic profile comparison of GF, OligoMM12 and SPF C57B6/J

mice by UPLC/MS. (A and B) Manually-curated list of compounds obtained by targeted peak

extraction from differentially expressed pathways in (A) liver and (B) plasma samples. p-values

obtained by Tukey's honest significance test after log2 transformation of area value.

Supplementary Table 1. List of metabolites identified by targeted peak extraction in the

UPLC/MS data. Table indicates compound name, KEGG Entry number, type of column was

used for UPLC and if the peak matched the retention time and MS2 spectra identified with the

chemical standard in liver and plasma samples.

REFERENCES

Adams, K.J., Pratt, B., Bose, N., Dubois, L.G., St John-Williams, L., Perrott, K.M., Ky, K., Kapahi, P., Sharma, V., MacCoss, M.J., et al. (2020). Skyline for Small Molecules: A Unifying Software Package for Quantitative Metabolomics. J. Proteome Res. *19*, 1447–1458, 10.1021/acs.jproteome.9b00640.

Arch, J.R.S., Hislop, D., Wang, S.J.Y., and Speakman, J.R. (2006). Some mathematical and technical issues in the measurement and interpretation of open-circuit indirect calorimetry in small animals. Int. J. Obes. (Lond). *30*, 1322–1331, 10.1038/sj.ijo.0803280.

Bäckhed, F., Ding, H., Wang, T., Hooper, L. V, Koh, G.Y., Nagy, A., Semenkovich, C.F., and Gordon, J.I. (2004). The gut microbiota as an environmental factor that regulates fat storage. Proc. Natl. Acad. Sci. U. S. A. *101*, 15718–15723, 10.1073/pnas.0407076101.

Bäckhed, F., Manchester, J.K., Semenkovich, C.F., and Gordon, J.I. (2007). Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. Proc. Natl. Acad. Sci. U. S. A. *104*, 979–984, 10.1073/pnas.0605374104.

Barlow, J.T., Bogatyrev, S.R., and Ismagilov, R.F. (2020). A quantitative sequencing framework for absolute abundance measurements of mucosal and lumenal microbial communities. Nat. Commun. *11*, 1–13, 10.1038/s41467-020-16224-6.

Becker, N., Kunath, J., Loh, G., and Blaut, M. (2011). Human intestinal microbiota: Characterization of a simplified and stable gnotobiotic rat model. Gut Microbes *2*, 25–33, 10.4161/gmic.2.1.14651.

Belenguer, A., Holtrop, G., Duncan, S.H., Anderson, S.E., Calder, A.G., Flint, H.J., and Lobley,

G.E. (2011). Rates of productionand utilization of lactate by microbial communities fromthe human colon. FEMS Microbiol. Ecol. *77*, 107–119, 10.1111/j.1574-6941.2011.01086.x.

Brugiroux, S., Beutler, M., Pfann, C., Garzetti, D., Ruscheweyh, H.J., Ring, D., Diehl, M., Herp, S., Lotscher, Y., Hussain, S., et al. (2016). Genome-guided design of a defined mouse microbiota that confers colonization resistance against Salmonella enterica serovar Typhimurium. Nat Microbiol *2*, 16215, 10.1038/nmicrobiol.2016.215.

Butler, A.A., and Kozak, L.P. (2010). A recurring problem with the analysis of energy expenditure in genetic models expressing lean and obese phenotypes. Diabetes *59*, 323–329, 10.2337/db09-1471.

Caesar, R., Tremaroli, V., Kovatcheva-Datchary, P., Cani, P.D., and Bäckhed, F. (2015). Crosstalk between Gut Microbiota and Dietary Lipids Aggravates WAT Inflammation through TLR Signaling. Cell Metab. *22*, 658–668, 10.1016/J.CMET.2015.07.026.

Cai, T.Q., Ren, N., Jin, L., Cheng, K., Kash, S., Chen, R., Wright, S.D., Taggart, A.K.P., and Waters, M.G. (2008). Role of GPR81 in lactate-mediated reduction of adipose lipolysis. Biochem. Biophys. Res. Commun. *377*, 987–991, 10.1016/j.bbrc.2008.10.088.

Cani, P.D., Van Hul, M., Lefort, C., Depommier, C., Rastelli, M., and Everard, A. (2019). Microbial regulation of organismal energy homeostasis. Nat. Metab. *1*, 34–46, 10.1038/s42255-018-0017-4.

Chong, J., and Xia, J. (2018). MetaboAnalystR: an R package for flexible and reproducible analysis of metabolomics data. Bioinformatics *34*, 4313–4314, 10.1093/bioinformatics/bty528.

Claus, S.P., Tsang, T.M., Wang, Y., Cloarec, O., Skordi, E., Martin, F.-P., Rezzi, S., Ross, A.,

Kochhar, S., Holmes, E., et al. (2008). Systemic multicompartmental effects of the gut microbiome on mouse metabolic phenotypes. Mol. Syst. Biol. *4*, 219, 10.1038/msb.2008.56.

Corrigan, J.K., Ramachandran, D., He, Y., Palmer, C.J., Jurczak, M.J., Chen, R., Li, B., Friedline, R.H., Kim, J.K., Ramsey, J.J., et al. (2020). A big-data approach to understanding metabolic rate and response to obesity in laboratory mice. Elife *9*, 10.7554/elife.53560.

Dennis, P.P., and Bremer, H. (2008). Modulation of Chemical Composition and Other Parameters of the Cell at Different Exponential Growth Rates. EcoSal Plus *3*, 10.1128/ecosal.5.2.3.

Doi, R., Oishi, K., and Ishida, N. (2010). CLOCK regulates circadian rhythms of hepatic glycogen synthesis through transcriptional activation of Gys2. J. Biol. Chem. *285*, 22114–22121, 10.1074/jbc.M110.110361.

Duncan, S.H., Louis, P., and Flint, H.J. (2004). Lactate-utilizing bacteria, isolated from human feces, that produce butyrate as a major fermentation product. Appl. Environ. Microbiol. *70*, 5810–5817, 10.1128/AEM.70.10.5810-5817.2004.

Eberl, C., Ring, D., Münch, P.C., Beutler, M., Basic, M., Slack, E.C., Schwarzer, M., Srutkova, D., Lange, A., Frick, J.S., et al. (2020). Reproducible Colonization of Germ-Free Mice With the Oligo-Mouse-Microbiota in Different Animal Facilities. Front. Microbiol. *10*, 2999, 10.3389/fmicb.2019.02999.

Faith, J.J., Ahern, P.P., Ridaura, V.K., Cheng, J., and Gordon, J.I. (2014). Identifying gut microbe-host phenotype relationships using combinatorial communities in gnotobiotic mice. Sci. Transl. Med. *6*, 220ra11-220ra11, 10.1126/scitranslmed.3008051.

Fernández-Verdejo, R., Ravussin, E., Speakman, J.R., and Galgani, J.E. (2019). Progress and challenges in analyzing rodent energy expenditure. Nat. Methods *16*, 797–799, 10.1038/s41592-019-0513-9.

Fetissov, S.O. (2017). Role of the gut microbiota in host appetite control: Bacterial growth to animal feeding behaviour. Nat. Rev. Endocrinol. *13*, 11–25, 10.1038/nrendo.2016.150.

Fleissner, C.K., Huebel, N., El-Bary, M.M.A., Loh, G., Klaus, S., and Blaut, M. (2010). Absence of intestinal microbiota does not protect mice from diet-induced obesity. Br. J. Nutr. *104*, 919–929, 10.1017/S0007114510001303.

Flint, H.J., Duncan, S.H., Scott, K.P., and Louis, P. (2015). Links between diet, gut microbiota composition and gut metabolism. Proc. Nutr. Soc. *74*, 13–22, 10.1017/S0029665114001463.

Gordon, H.A., and Pesti, L. (1971). The gnotobiotic animal as a tool in the study of host microbial relationships. Bact. Rev. *35*, 390–429, 10.1126/science.173.3992.171.

Goswami, C., Iwasaki, Y., and Yada, T. (2018). Short-chain fatty acids suppress food intake by activating vagal afferent neurons. J. Nutr. Biochem. *57*, 130–135, 10.1016/j.jnutbio.2018.03.009.

Halatchev, I.G., O'Donnell, D., Hibberd, M.C., and Gordon, J.I. (2019). Applying indirect open-circuit calorimetry to study energy expenditure in gnotobiotic mice harboring different human gut microbial communities. Microbiome *7*, 158, 10.1186/s40168-019-0769-4.

Harley, I.T.W., and Karp, C.L. (2012). Obesity and the gut microbiome: Striving for causality. Mol. Metab. *1*, 21–31, 10.1016/j.molmet.2012.07.002.

Ishikawa, K., and Shimazu, T. (1976). Daily rhythms of glycogen synthetase and phosphorylase

activities in rat liver: influence of food and light. Life Sci. *19*, 1873–1878, 10.1016/0024-3205(76)90119-3.

Kaiyala, K.J., and Schwartz, M.W. (2011). Toward a more complete (and less controversial) understanding of energy expenditure and its role in obesity pathogenesis. Diabetes *60*, 17–23, 10.2337/db10-0909.

Koh, A., and Bäckhed, F. (2020). From Association to Causality: the Role of the Gut Microbiota and Its Functional Products on Host Metabolism. Mol. Cell *78*, 584–596, 10.1016/j.molcel.2020.03.005.

Kovatcheva-Datchary, P., Shoaie, S., Lee, S., Wahlström, A., Nookaew, I., Hallen, A., Perkins, R., Nielsen, J., and Bäckhed, F. (2019). Simplified Intestinal Microbiota to Study Microbe-Diet-Host Interactions in a Mouse Model. Cell Rep. *26*, 3772-3783.e6, 10.1016/j.celrep.2019.02.090.

Krisko, T.I., Nicholls, H.T., Bare, C.J., Holman, C.D., Putzel, G.G., Jansen, R.S., Sun, N., Rhee, K.Y., Banks, A.S., and Cohen, D.E. (2020). Dissociation of Adaptive Thermogenesis from Glucose Homeostasis in Microbiome-Deficient Mice. Cell Metab. *31*, 592-604.e9, 10.1016/J.CMET.2020.01.012.

Kuang, Z., Wang, Y., Li, Y., Ye, C., Ruhn, K.A., Behrendt, C.L., Olson, E.N., and Hooper, L. V. (2019). The intestinal microbiota programs diurnal rhythms in host metabolism through histone deacetylase 3. Science *365*, 1428–1434, 10.1126/science.aaw3134.

Kübeck, R., Bonet-Ripoll, C., Hoffmann, C., Walker, A., Müller, V.M., Schüppel, V.L., Lagkouvardos, I., Scholz, B., Engel, K.-H., Daniel, H., et al. (2016). Dietary fat and gut microbiota interactions determine diet-induced obesity in mice. Mol. Metab. *5*, 1162–1174, 10.1016/j.molmet.2016.10.001.

Levenson, S.M. (1978). The influence of the indigenous microflora on mammalian metabolism and nutrition. JPEN. J. Parenter. Enteral Nutr. *2*, 78–107, 10.1177/014860717800200203.

Levenson, S.M., Doft, F., Lev, M., and Kan, D. (1969). Influence of microorganisms on oxygen consumption, carbon dioxide production and colonic temperature of rats. J. Nutr. *97*, 542–552, 10.1093/jn/97.4.542.

Li, B., Li, L., Li, M., Lam, S.M., Wang, G., Wu, Y., Zhang, H., Niu, C., Zhang, X., Liu, X., et al. (2019). Microbiota Depletion Impairs Thermogenesis of Brown Adipose Tissue and Browning of White Adipose Tissue. Cell Rep. *26*, 2720-2737.e5, 10.1016/j.celrep.2019.02.015.

Li, M., Li, L., Li, B., Hambly, C., Wang, G., Wu, Y., Jin, Z., Wang, A., Niu, C., Wolfrum, C., et al. (2021). Brown adipose tissue is the key depot for glucose clearance in microbiota depleted mice. Nat. Commun. *12*, 1–13, 10.1038/s41467-021-24659-8.

Liebisch, G., Ecker, J., Roth, S., Schweizer, S., Öttl, V., Schött, H.F., Yoon, H., Haller, D., Holler, E., Burkhardt, R., et al. (2019). Quantification of fecal short chain fatty acids by liquid chromatography tandem mass spectrometry—investigation of pre-analytic stability. Biomolecules *9*, 121, 10.3390/biom9040121.

Lin, H. V., Frassetto, A., Kowalik Jr, E.J., Nawrocki, A.R., Lu, M.M., Kosinski, J.R., Hubert, J.A., Szeto, D., Yao, X., Forrest, G., et al. (2012). Butyrate and Propionate Protect against Diet-Induced Obesity and Regulate Gut Hormones via Free Fatty Acid Receptor 3-Independent Mechanisms. PLoS One *7*, e35240, 10.1371/journal.pone.0035240.

Liu, C., Wu, J., Zhu, J., Kuei, C., Yu, J., Shelton, J., Sutton, S.W., Li, X., Su, J.Y., Mirzadegan, T., et al. (2009). Lactate inhibits lipolysis in fat cells through activation of an orphan G-protein-coupled receptor, GPR81. J. Biol. Chem. *284*, 2811–2822, 10.1074/jbc.M806409200.

MacLean, P.S., Blundell, J.E., Mennella, J.A., and Batterham, R.L. (2017). Biological control of appetite: A daunting complexity. Obesity 25, S8–S16, 10.1002/oby.21771.

Mallapaty, S. (2017). Gnotobiotics: getting a grip on the microbiome boom. Lab Anim. (NY). *46*, 373–377, 10.1038/laban.1344.

Mardinoglu, A., Shoaie, S., Bergentall, M., Ghaffari, P., Zhang, C., Larsson, E., Bäckhed, F., and Nielsen, J. (2015). The gut microbiota modulates host amino acid and glutathione metabolism in mice. Mol. Syst. Biol. *11*, 834, 10.15252/MSB.20156487.

Meyer, C.W., Reitmeir, P., and Tschop, M.H. (2015). Exploration of Energy Metabolism in the Mouse Using Indirect Calorimetry: Measurement of Daily Energy Expenditure (DEE) and Basal Metabolic Rate (BMR). Curr Protoc Mouse Biol *5*, 205–222, 10.1002/9780470942390.mo140216.

Mina, A.I., LeClair, R.A., LeClair, K.B., Cohen, D.E., Lantier, L., and Banks, A.S. (2018). CalR: A Web-Based Analysis Tool for Indirect Calorimetry Experiments. Cell Metab. *28*, 656–666, 10.1016/j.cmet.2018.06.019.

Moor, K., Fadlallah, J., Toska, A., Sterlin, D., Balmer, M.L., Macpherson, A.J., Gorochov, G., Larsen, M., and Slack, E. (2016). Analysis of bacterial-surface-specific antibodies in body fluids using bacterial flow cytometry. Nat Protoc *11*, 1531–1553, 10.1038/nprot.2016.091.

Murtagh, F., and Legendre, P. (2014). Ward's Hierarchical Agglomerative Clustering Method: Which Algorithms Implement Ward's Criterion? J. Classif. 2014 313 *31*, 274–295, 10.1007/S00357-014-9161-Z.

Nowak, N. (2021). Metabolic Insights Related to Sleep and Circadian Clocks from Mass

Spectrometry-Based Analysis of Blood and Breath. 10.3929/ETHZ-B-000480810.

Packard, G.C., and Boardman, T.J. (1999). The use of percentages and size-specific indices to normalize physiological data for variation in body size: Wasted time, wasted effort? Comp. Biochem. Physiol. - A Mol. Integr. Physiol. *122*, 37–44, 10.1016/S1095-6433(98)10170-8.

Popovic, M. (2019). Thermodynamic properties of microorganisms: determination and analysis of enthalpy, entropy, and Gibbs free energy of biomass, cells and colonies of 32 microorganism species. Heliyon *5*, e01950, 10.1016/j.heliyon.2019.e01950.

Rabasa, C., and Dickson, S.L. (2016). Impact of stress on metabolism and energy balance. Curr. Opin. Behav. Sci. *9*, 71–77, 10.1016/J.COBEHA.2016.01.011.

Ridaura, V.K., Faith, J.J., Rey, F.E., Cheng, J., Duncan, A.E., Kau, A.L., Griffin, N.W., Lombard, V., Henrissat, B., Bain, J.R., et al. (2013). Gut microbiota from twins discordant for obesity modulate metabolism in mice. Science *341*, 1241214, 10.1126/science.1241214.

Sayin, S.I., Wahlström, A., Felin, J., Jäntti, S., Marschall, H.U., Bamberg, K., Angelin, B., Hyötyläinen, T., Orešič, M., and Bäckhed, F. (2013). Gut microbiota regulates bile acid metabolism by reducing the levels of tauro-beta-muricholic acid, a naturally occurring FXR antagonist. Cell Metab. *17*, 225–235, 10.1016/j.cmet.2013.01.003.

Schmidt, T.S.B., Raes, J., and Bork, P. (2018). The Human Gut Microbiome: From Association to Modulation. Cell *172*, 1198–1215, 10.1016/j.cell.2018.02.044.

von Schwartzenberg, R.J., Bisanz, J.E., Lyalina, S., Spanogiannopoulos, P., Ang, Q.Y., Cai, J., Dickmann, S., Friedrich, M., Liu, S.-Y., Collins, S.L., et al. (2021). Caloric restriction disrupts the microbiota and colonization resistance. Nature 1–6, 10.1038/s41586-021-03663-4.

Smith, C.A., Want, E.J., O'Maille, G., Abagyan, R., and Siuzdak, G. (2006). XCMS: processing mass spectrometry data for metabolite profiling using nonlinear peak alignment, matching, and identification. Anal. Chem. *78*, 779–787, 10.1021/ac051437y.

Sonnenburg, J.L., and Bäckhed, F. (2016). Diet-microbiota interactions as moderators of human metabolism. Nature *535*, 56–64, 10.1038/nature18846.

Speakman, J.R. (2013). Measuring energy metabolism in the mouse - theoretical, practical, and analytical considerations. Front Physiol *4*, 34, 10.3389/fphys.2013.00034.

Steimle, A., De Sciscio, A., Neumann, M., Grant, E.T., Pereira, G. V., Ohno, H., Martens, E.C., and Desai, M.S. (2021). Constructing a gnotobiotic mouse model with a synthetic human gut microbiome to study host–microbe cross talk. STAR Protoc. *2*, 100607, 10.1016/J.XPRO.2021.100607.

Streidl, T., Karkossa, I., Segura Muñoz, R.R., Eberl, C., Zaufel, A., Plagge, J., Schmaltz, R., Schubert, K., Basic, M., Schneider, K.M., et al. (2021). The gut bacterium Extibacter muris produces secondary bile acids and influences liver physiology in gnotobiotic mice. Gut Microbes *13*, 1–21, 10.1080/19490976.2020.1854008.

Stubbs, R.J., Hopkins, M., Finlayson, G.S., Duarte, C., Gibbons, C., and Blundell, J.E. (2018). Potential effects of fat mass and fat-free mass on energy intake in different states of energy balance. Eur. J. Clin. Nutr. *72*, 698–709, 10.1038/s41430-018-0146-6.

Suárez-Zamorano, N., Fabbiano, S., Chevalier, C., Stojanović, O., Colin, D.J., Stevanović, A., Veyrat-Durebex, C., Tarallo, V., Rigo, D., Germain, S., et al. (2015). Microbiota depletion promotes browning of white adipose tissue and reduces obesity. Nat. Med. *21*, 1497–1501, 10.1038/nm.3994.

Tahara, Y., Yamazaki, M., Sukigara, H., Motohashi, H., Sasaki, H., Miyakawa, H., Haraguchi, A., Ikeda, Y., Fukuda, S., and Shibata, S. (2018). Gut Microbiota-Derived Short Chain Fatty Acids Induce Circadian Clock Entrainment in Mouse Peripheral Tissue. Sci. Rep. *8*, 1395, 10.1038/s41598-018-19836-7.

Tanner, J.M. (1949). Fallacy of per-weight and per-surface area standards, and their relation to spurious correlation. J. Appl. Physiol. *2*, 1–15, 10.1152/jappl.1949.2.1.1.

Thaiss, C.A., Zeevi, D., Levy, M., Zilberman-Schapira, G., Suez, J., Tengeler, A.C., Abramson, L., Katz, M.N., Korem, T., Zmora, N., et al. (2014). Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. Cell *159*, 514–529, 10.1016/j.cell.2014.09.048.

Thaiss, C.A., Levy, M., Korem, T., Dohnalova, L., Shapiro, H., Jaitin, D.A., David, E., Winter, D.R., Gury-BenAri, M., Tatirovsky, E., et al. (2016). Microbiota Diurnal Rhythmicity Programs Host Transcriptome Oscillations. Cell *167*, 1495-1510 e12, 10.1016/j.cell.2016.11.003.

Touw, K., Ringus, D.L., Hubert, N., Wang, Y., Leone, V.A., Nadimpalli, A., Theriault, B.R., Huang, Y.E., Tune, J.D., Herring, P.B., et al. (2017). Mutual reinforcement of pathophysiological host-microbe interactions in intestinal stasis models. Physiol. Rep. *5*, e13182, 10.14814/PHY2.13182.

Tschop, M.H., Speakman, J.R., Arch, J.R., Auwerx, J., Bruning, J.C., Chan, L., Eckel, R.H., Farese Jr., R. V, Galgani, J.E., Hambly, C., et al. (2011). A guide to analysis of mouse energy metabolism. Nat Methods *9*, 57–63, 10.1038/nmeth.1806.

Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., and Gordon, J.I. (2006). An obesity-associated gut microbiome with increased capacity for energy harvest. Nature *444*, 1027–1031, 10.1038/nature05414.

Ucar, I., Pebesma, E., and Azcorra, A. (2019). Measurement errors in R. R J. *10*, 549–557, 10.32614/RJ-2018-075.

Walter, J., Armet, A.M., Finlay, B.B., and Shanahan, F. (2020). Establishing or Exaggerating Causality for the Gut Microbiome: Lessons from Human Microbiota-Associated Rodents. Cell *180*, 221–232, 10.1016/j.cell.2019.12.025.

Wang, Y., Kuang, Z., Yu, X., Ruhn, K.A., Kubo, M., and Hooper, L. V. (2017). The intestinal microbiota regulates body composition through NFIL3 and the circadian clock. Science *357*, 912–916, 10.1126/science.aan0677.

Weir, J.B.D.B. (1949). New methods for calculating metabolic rate with special reference to protein metabolism. J. Physiol. *109*, 1–9, 10.1113/jphysiol.1949.sp004363.

Weiss, A.S., Burrichter, A.G., Chakravarthy Durai Raj, A., Von, A., Meng, C., Kleigrewe, K., Münch, P.C., Rössler, L., Eisenreich, W., Jochum, L.M., et al. (2021). Exploring the interaction network of a synthetic gut bacterial community. BioRxiv 2021.02.25.432904, 10.1101/2021.02.25.432904.

White, C.R., and Seymour, R.S. (2005). Allometric scaling of mammalian metabolism. J. Exp. Biol. *208*, 1611–1619, 10.1242/jeb.01501.

Wostmann, B.S., Bruckner-Kardoss, E., and Knight, P.L. (1968). Cecal Enlargement, Cardiac Output, and O2 Consumption in Germfree Rats. Exp. Biol. Med. *128*, 137–141, 10.3181/00379727-128-32962.

Wostmann, B.S., Larkin, C., Moriarty, A., and Bruckner-Kardoss, E. (1983). Dietary intake, energy metabolism, and excretory losses of adult male germfree Wistar rats. Lab. Anim. Sci. *33*,

46–50.

Woting, A., Pfeiffer, N., Loh, G., Klaus, S., and Blaut, M. (2014). Clostridium ramosum promotes High-Fat diet-induced obesity in Gnotobiotic Mouse Models. MBio *5*, 1530–1544, 10.1128/mBio.01530-14.

Woting, A., Pfeiffer, N., Hanske, L., Loh, G., Klaus, S., and Blaut, M. (2015). Alleviation of high fat diet-induced obesity by oligofructose in gnotobiotic mice is independent of presence of Bifidobacterium longum. Mol. Nutr. Food Res. *59*, 2267–2278, 10.1002/mnfr.201500249.

Wotzka, S.Y., Kreuzer, M., Maier, L., Arnoldini, M., Nguyen, B.D., Brachmann, A.O., Berthold, D.L., Zünd, M., Hausmann, A., Bakkeren, E., et al. (2019). Escherichia coli limits Salmonella Typhimurium infections after diet shifts and fat-mediated microbiota perturbation in mice. Nat. Microbiol. 10.1038/s41564-019-0568-5.

Yao, L., Seaton, S.C., Ndousse-Fetter, S., Adhikari, A.A., DiBenedetto, N., Mina, A.I., Banks, A.S., Bry, L., and Devlin, A.S. (2018). A selective gut bacterial bile salt hydrolase alters host metabolism. Elife *7*, 10.7554/eLife.37182.

Yilmaz, B., Mooser, C., Keller, I., Li, H., Zimmermann, J., Bosshard, L., Fuhrer, T., Gomez de Agüero, M., Trigo, N.F., Tschanz-Lischer, H., et al. (2021). Long-term evolution and short-term adaptation of microbiota strains and sub-strains in mice. Cell Host Microbe *29*, 650-663.e9, 10.1016/j.chom.2021.02.001.

Zarrinpar, A., Chaix, A., Yooseph, S., and Panda, S. (2014). Diet and feeding pattern affect the diurnal dynamics of the gut microbiome. Cell Metab. *20*, 1006–1017, 10.1016/j.cmet.2014.11.008.

Zarrinpar, A., Chaix, A., Xu, Z.Z., Chang, M.W., Marotz, C.A., Saghatelian, A., Knight, R., and

Panda, S. (2018). Antibiotic-induced microbiome depletion alters metabolic homeostasis by

affecting gut signaling and colonic metabolism. Nat. Commun. 2018 91 *9*, 1–13,

10.1038/s41467-018-05336-9.

SUPPLEMENTARY TABLE

Supplementary Table 1. List of metabolites identified by targeted peak extraction in the UPLC/MS data. Table indicates compound name, KEGG Entry number, type of column was used for UPLC and if the peak ID matched the retention time and MS2 spectra identified with the chemical standard in liver and plasma samples.

Compound	KEGG Entry	Column	Matched peak ID in Liver	Matched peak ID in Plasma
(R)-3-hydroxybutanic acid	C01089	RP Negative	Yes	No
5-Oxoproline	C01879	AMIDE Positive	Yes	Yes
Arachidonic acid	C00219	RP Negative	Yes	Yes
Betaine	C00719	AMIDE Positive	No	Yes
beta-Murocholic acid	C17726	RP Negative	Yes	Yes
Cholic acid	C00695	RP Negative	Yes	Yes
Citrulline	C00327	AMIDE Positive	Yes	Yes
Cortisol	C00735	RP Positive	Yes	No
Creatine	C00300	AMIDE Positive	Yes	Yes

Creatinine	C00791	AMIDE Positive	No	Yes
Cystathionine	C02291	AMIDE Positive	Yes	Yes
Deoxycholic acid	C04483	RP Negative	No	Yes
Docosapentaenoic acid	C16513	RP Negative	Yes	Yes
Glutathione (GSH)	C00051	AMIDE Positive	No	Yes
Glycine	C00037	AMIDE Positive	Yes	Yes
Glycocholic acid	C01921	RP Positive	Yes	No
Hexadecanedioic acid		RP Negative	Yes	Yes
Hippuric acid	C01586	RP Negative	Yes	Yes
L-(+)-Ornithine	C00077	AMIDE Positive	Yes	Yes
L-4-Hydroxyproline	C01157	AMIDE Positive	Yes	Yes
L-Alanine	C00041	AMIDE Positive	Yes	Yes
L-Arginine	C00062	AMIDE Positive	Yes	Yes
Lauroylcarnitine		RP Positive	Yes	Yes

Leucine	C00123	RP Positive	Yes	Yes
L-Glutamic acid	C00025	AMIDE Positive	Yes	Yes
L-Histidine	C00135	AMIDE Positive	Yes	Yes
Linoleic acid	C01595	RP Negative	Yes	Yes
L-Isoleucine	C00407	RP Positive	Yes	Yes
L-Lysine	C00047	AMIDE Positive	Yes	Yes
L-Methionine	C00073	AMIDE Positive	Yes	Yes
L-Proline	C00148	AMIDE Positive	Yes	Yes
L-Serine	C00065	AMIDE Positive	Yes	Yes
L-Threonine	C00188	AMIDE Positive	Yes	Yes
L-Tryptophan	C00078	RP Negative	Yes	Yes
L-Tyrosine	C00082	RP Negative	Yes	Yes
Myristic acid	C06424	RP Negative	Yes	Yes
N,N-dimethylglycine	C01026	AMIDE Positive	Yes	Yes

	C02712	RP Positive/Negative	Yes	Yes
N-Acetylmethionine				
Oxidized glutathione (GSSG)	C00127	AMIDE Negative	Yes	Yes
Pantothenic acid	C00864	RP Positive	Yes	Yes
Pipecolinic acid	C00408	RP Positive	Yes	Yes
Serotonin	C00780	RP Positive	Yes	Yes
Succinic acid	C00042	RP Negative	Yes	Yes
Taurine	C00245	AMIDE Negative	Yes	Yes
Taurine-beta-murocholic acid		AMIDE Negative	Yes	Yes
Taurochenodeoxycholic acid	C05465	AMIDE Negative	Yes	Yes
Tetradecanedioic acid		RP Negative	Yes	Yes

Figure 1: OligoMM12 mice have increase fat mass compared to GF mice and SPF C57B6/J mice.

Figure 2: Energy metabolism in GF, OligoMM12 and SPF C57B6/J mice.

Figure 3: Circadian changes in Respiratory Exchange Ratio (RER), microbiota-derived hydrogen and short-chain fatty acids (SCFAs).

Figure 4. Metabolic profile comparison of GF, OligoMM12 and SPF mice by UPLC/MS.

Supplementary Figure 1: Cecal mass interferes with fat mass estimation by Echo-MRI.

Supplementary Figure 2: Cecal mass interferes with normalization of VO2 and Energy Expenditure.

Supplementary Figure 3. Metabolic profile comparison of GF, OligoMM12 and SPF mice by UPLC/MS (targeted peak extraction of selected list of compounds)