Host- and age-dependent transcriptional changes in *Mycobacterium tuberculosis* cell

envelope biosynthesis genes after exposure to human alveolar lining fluid

Anna Allué-Guardia[1*], Andreu Garcia-Vilanova[1], Angélica M. Olmo-Fontánez[1,2], Jay Peters[3],

Diego J. Maselli[3], Joanne Turner[4], Larry S. Schlesinger[4], and Jordi B. Torrelles[1*]

[1] Population Health Program, Texas Biomedical Research Institute, San Antonio, TX 78227, USA

[2] Integrated Biomedical Sciences Program, University of Texas Health Science Center at San

Antonio, TX 78229, USA

[3] Division of Pulmonary and Critical Care Medicine, School of Medicine, UT Health San Antonio,

San Antonio, TX 78229, USA

[4] Host-Pathogen Interactions Program, Texas Biomedical Research Institute, San Antonio, TX

78227, USA

[*] **Corresponding Authors**: Anna Allué-Guardia, PhD (aallueguardia@txbiomed.org) and Jordi

B. Torrelles, PhD (jtorrelles@txbiomed.org)

Running Title: Human lung mucosa and *M. tuberculosis* infection

Keywords: *Mycobacterium tuberculosis*, alveolar lining fluid (ALF), lung mucosa, cell envelope

biosynthesis, gene expression

Abstract

Tuberculosis (TB) infection, caused by the airborne pathogen *Mycobacterium tuberculosis* (*M.tb*), resulted in almost 1.4 million deaths in 2019 and the number of deaths is predicted to increase by 20% over the next 5 years due to the COVID-19 pandemic. Upon reaching the alveolar space, *M.tb* comes in close contact with the lung mucosa before and after its encounter with host alveolar compartment cells. Our previous studies show that homeostatic innate soluble components of the alveolar lining fluid (ALF) can quickly alter the cell envelope surface of *M.tb* upon contact, defining subsequent *M.tb*-host cell interactions and infection outcomes *in vitro* and *in vivo*. We also demonstrated that ALF from 60+ year old elders (E-ALF) *vs.* healthy 18- to 45-year-old adults (A-ALF) is dysfunctional with loss of homeostatic capacity and impaired innate soluble responses linked to high local oxidative stress. In this study, a targeted transcriptional assay demonstrates that *M.tb* exposure to human ALF alters the expression of its cell envelope genes. Further, our results indicate that A-ALF-exposed *M.tb* upregulates cell envelope genes associated with lipid, carbohydrate, and amino acid metabolism, as well as genes associated with redox homeostasis and transcriptional regulators. Conversely, *M.tb* exposure to E-ALF shows lesser transcriptional response, with most of the *M.tb* genes unchanged or downregulated. Overall, this study indicates that *M.tb* responds and adapts to the lung alveolar environment upon contact, and that the host ALF status determined by factors such as age might play an important role in determining infection outcome.

INTRODUCTION

Mycobacterium tuberculosis (*M.tb*), the causative agent of tuberculosis (TB), is one of the top leading causes of mortality worldwide due to a single infectious agent, with ~1.4 million attributed deaths in 2019 [1]. However, global estimates indicate that worldwide disruptions in the healthcare system during the current COVID-19 pandemic could lead to an additional 6.3 million new TB cases between 2020 and 2025 and an added 1.4 million more TB deaths [2, 3]. Strict lockdowns have prevented patients from having access to TB medications and clinical evaluations, and have led to decreased TB diagnosis rates, since available resources have been redirected to prevent the spread of COVID-19 [4, 5]. These factors are predicted to cause not only an increase in the number of TB cases, but also to promote the development of drug-resistant TB, stressing the need for the development of new anti-TB therapies [6].

Most current drugs target the *M.tb* cell envelope, a highly complex and dynamic structure comprised mainly of carbohydrates and lipids, which provide structural support and resistance to osmotic changes, as well as a critical immunoregulatory role during *M.tb* infection [7-10]. It consists of four main layers: 1] an inner plasma membrane with periplasmic space; 2] a peptidoglycan (PG) core covalently linked to arabinogalactan (AG) and mycolic acids (MAcs); 3] a peripheral layer of non-covalently linked lipids, glycolipids, and lipoglycans [*e.g.* phthiocerol dimycocerosates (PDIMs), trehalose dimycolate (TDM) and monomycolate (TMM), sulfolipids (SLs), phosphatidyl-*myo*-inositol mannosides (PIMs), lipomannan (LM), and mannose-capped lipoarabinomannan (ManLAM), among others], and; 4) the outermost layer or capsule [11, 12]. The pathogenesis of *M.tb* is inherently linked to its heterogeneous and dynamic cell envelope surface, and cell envelope remodeling has been observed during infection and in response to

environmental stresses [10]. Thus, the particular cell envelope composition of a mycobacterial cell

at a given moment will define *M.tb*-host cell interactions and determine the infection outcome.

However, it remains poorly understood how the *M.tb* cell envelope changes and adapts to the host

lung environment during the natural course of pulmonary infection, a critical gap in our knowledge

for defining new drug targets against *M.tb* relevant to bacteria in the lung environment.

After the host inhales droplets containing *M.tb*, the airborne pathogen is deposited in the lung

alveolar space. Here it first comes in contact with soluble components of the alveolar lining fluid

(ALF) for an undefined period of time (from minutes to hours/days) [13-16], before and after its

encounter with host alveolar compartment cells such as alveolar macrophages (AMs) or alveolar

epithelial cells (ATs), and immune cells such as neutrophils [17, 18]. Our previous studies have

demonstrated that homeostatic ALF hydrolytic enzymes, whose function is to promote lung health,

can modify the *M.tb* cell envelope without reducing *M.tb* viability [19, 20], including a reduction

in major *M.tb* virulent factors mannose-capped lipoarabinomannan (ManLAM) and trehalose

dimycolate (TDM), from the *M.tb* cell surface. These ALF-derived *M.tb* cell envelope

modifications have an impact on *M.tb* infection outcomes *in vitro* and *in vivo*, since they allow for

a better recognition by cells of the immune system and improved control of the infection [19-23].

Indeed, exposure to ALF results in decreased *M.tb* association and intracellular growth within

human macrophages, as well as altered intracellular trafficking and increased pro-inflammatory

responses [19]. Neutrophils also possess an enhanced innate ability to recognize and kill

intracellular ALF-exposed *M.tb*, while limiting excessive inflammatory responses [21]. *In vivo*

infections using ALF-exposed *M.tb* demonstrate better control of infection in the mouse model

[22]. Further, *M.tb* fragments released after ALF exposure by the action of the ALF hydrolases,

are capable of priming neutrophils and modulating macrophages in an IL-10 dependent manner to

contribute further to the control of *M.tb* [20, 23].

Importantly, the levels and functionality of ALF soluble components are altered in certain human

populations such as the elderly, with increased pro-oxidation and pro-inflammatory pathways,

altered complement and surfactant levels, and decreased binding capability of surfactant protein

A (SP-A) and D (SP-D) in the aging lung, defining what we call 'dysfunctional ALF'.

Consequently, *M.tb* exposed to elderly dysfunctional ALF, while maintaining its viability, show

increased intracellular growth in macrophages and ATs, as well as increased bacterial burden in

mice with increased lung tissue damage [22, 24, 25]. Altogether, these results indicate that the host

ALF functional status plays a key role in shaping the *M.tb* cell envelope during the initial stages

of infection. However, the overall impact of these different ALF microenvironments on *M.tb*

adaptation to the human host and subsequent infection progression is still largely unknown.

Since hydrolytic enzymes present in functional human ALF (from healthy adult individuals)

modify the *M.tb* cell envelope [19], we hypothesize that *M.tb* will compensate for these ALF-

driven modifications by altering the expression of genes related to cell envelope biogenesis.

Conversely, *M.tb* exposed to ALF with decreased functionality (such as elderly ALF) will show

little to no changes. In addition, our published data [20, 26] demonstrate that 15-minute exposure

to human ALF is enough to alter the *M.tb* cell envelope and its interactions with host cells and that

these modifications are maintained for up to 24 h [20, 26]. In this study, we first aim to determine

if short (15 min) or long (12 h) exposure to human ALF has any effects on the expression of

targeted *M.tb* cell envelope genes, indicating that gene expression is altered at 15 min and that

these changes are maintained up to 12 h. Then, we use a multiplex qPCR assay to compare an

extensive transcriptional profile of *M.tb* cell envelope genes associated with lipid, carbohydrate

and amino acid metabolism, among others, after exposure to functional healthy adult (A)- *vs.*

dysfunctional healthy elderly (E)-ALF. Our results show significant differences in gene

expression, where A-ALF exposed *M.tb* upregulated genes involved in cell envelope remodeling,

thus implicating this remodeling in subsequent *M.tb*-host interactions during the infection process.

MATERIALS AND METHODS

Human subjects and ethics statement - Human ALFs used in this study were previously isolated

from collected bronchoalveolar lavage fluid (BALF) from healthy adult and elderly volunteers, in

strict accordance with the US Code of Federal and approved Local Regulations (The Ohio State

University Human Subjects IRB numbers 2012H0135 & 2008H0119 and Texas Biomedical

Research Institute/UT-Health San Antonio/South Texas Veterans Health Care System Human

Subjects IRB numbers HSC20170667H & HSC20170673H), and Good Clinical Practice as

approved by the National Institutes of Health (NIAID/DMID branch), with written informed

consent from all human subjects. Healthy adults (18-45 years old) and elderly (+60 years) were

recruited from both sexes and no discrimination of race or ethnicity.

Collection of BALF and ALF - BALF was collected from healthy adult or elderly donors in

sterile endotoxin free 0.9% of NaCl, filtered through 0.2 μm filters, and further concentrated 20-

fold using Amicon Ultra Centrifugal Filter Units with a 10-kDa molecular mass cut-off membrane

(Millipore Sigma) at 4°C to obtain ALF with a physiological concentration reported within the

human lung (1 mg/mL of phospholipid), as we previously described [19-24, 27]. ALF was

aliquoted in low-protein binding tubes and stored at -80°C until further use.

Bacterial cultures and ALF exposure - *M.tb* strains GFP-Erdman (kindly provided by Dr.

Marcus Horwitz, UCLA) and $H_{37}R_v$ (ATCC# 25618) were cultured in 7H11 agar plates (BD BBL),

supplemented with oleic acid, albumin, dextrose and catalase (OADC) at 37°C for 14 days. Single

bacterial suspensions ($\sim 1 \times 10^9$ bacteria/ml) were obtained as previously described [19-23].

Bacterial pellets were exposed to individual ALFs (from different donors) for 15 min or ~12 h at

37°C. After exposure, ALF was removed and bacterial pellets were directly incubated in

RNAProtect (Qiagen) for 10 min at room temperature (RT), centrifuged at 13,000 x *g* and stored

at -80°C until further use. For each of the ALFs, corresponding heat-inactivated ALFs (2 h at 80°C)

[21] were used as controls in parallel.

RNA extraction and cDNA synthesis - RNA from ALF- exposed bacterial pellets was extracted

using the Quick-RNA Fungal/Bacterial Miniprep kit (Zymo Research), following the

manufacturer's protocol. Briefly, bacterial pellets were resuspended in Lysis buffer and transferred

to a ZR BashingBead Lysis tube containing 0.1 mm and 0.5 mm ceramic beads. Bead beating

procedure was performed to break the tough-to-lyse mycobacterial cell envelope in a Disruptor

Genie [10 cycles of 1 min at maximum speed with 1 min intervals on ice]. RNA was isolated from

the supernatant using Zymo-spin columns, including an in-column DNAse I treatment, and eluted

in nuclease-free water. To completely remove the genomic DNA, a second DNAse treatment was

performed on the isolated RNA using TURBO DNAse reagent (Thermo Fisher Scientific) for 30

min at 37°C. Final RNA concentration and quality were measured with a Qubit 4 Fluorometer

using the HS RNA kit (Thermo Fisher Scientific), and a Nanodrop One[C], respectively. RNA (500

ng) was used for the synthesis of cDNA using the RevertAid H Minus First Strand cDNA Synthesis

kit (Thermo Fisher Scientific) with random hexamer primers, following the manufacturer's

guidelines.

qPCR analysis of targeted genes - Real-time quantitative PCR (qPCR) was performed to measure

the expression of ten genes associated with the *M.tb* cell wall biosynthesis pathways (*pmmA,*

manB, whiB2, pimA, pimB, pimF, embC, pmmB, pmm1, manA) after exposure of *M.tb* $H_{37}R_v$ to

healthy human ALF (n=2 biological replicates, from two different donors). cDNA and primers

were used in a 20 μl qPCR reaction with PowerUp SYBR Green Master Mix (Applied

Biosystems), following manufacturer's instructions. Reactions were run in an Applied Biosystems

7500 Real-Time PCR instrument with the following settings: reporter SYBR Green, no quencher,

passive reference dye ROX, standard ramp speed, and continuous melt curve ramp increment.

Expression was calculated relative to housekeeping genes *rpoB* or *sigA* using the $2^{-\Delta\Delta CT}$ method

[28].

High-throughput multiplex qPCR - Primer pairs for multiplex qPCR assay were designed using

BatchPrimer3 (https://wheat.pw.usda.gov/demos/BatchPrimer3/) [29] and PrimerQuest (IDT,

https://www.idtdna.com/pages/tools/primerquest), with the following settings: primer size 18-21

ntds, T_m of ~ 54-60°C, and maximum 3' self-complementarity of 3 ntd. Best primers within these

parameters were selected and aligned to the *M.tb* $H_{37}R_v$ reference genome (Genbank accession

number: NC_000962.3) [30-32] to confirm that they uniquely aligned to the targeted gene regions.

A high-throughput multiplex qPCR targeting more than 80 genes associated with *M.tb* metabolism

and cell wall biogenesis was performed in *M.tb* Erdman exposed to ALF using the Biomark 96.96

Dynamic Array IFC for Gene Expression in a Biomark HD instrument (Fluidigm). Briefly, 1.25

µl of cDNA was pre-amplified in a 5 µl reaction with our pool of specific primers (500nM) using

the Fluidigm Preamp Master Mix for a total of 10 cycles. Then, a 1/10 dilution of the pre-amplified

cDNA and 100 µM of combined forward and reverse primers were used to prepare the sample pre-

mix and assay mix, respectively. The 96.96 IFC chip was loaded and the assay run in a Biomark

HD following the manufacturer's instructions for Gene expression using Delta Gene Assays.

Relative expression for each of the genes was calculated in the Fluidigm Real-Time PCR Analysis

software using the $2^{-\Delta\Delta CT}$ method with *rpoB* as the reference gene, and reported as the $\log_2$ fold

change of A-ALF (n=3 biological replicates using ALFs from different donors) or E-ALF (n=3

biological replicates using ALFs from different donors) exposed *M.tb* Erdman compared to control

samples (*M.tb* Erdman exposed to corresponding heat-inactivated ALFs). As a control to compare

both methods (Biomark *vs.* targeted qPCR), we included *M.tb* $H_{37}R_v$ exposed to adult ALF in the

Biomark multiplex qPCR assay.

Statistical analysis - Statistical significance between the two qPCR methods used in this study

was calculated in GraphPad Prism v9.0.1 for each of the genes with a two-way ANOVA using the

Sidak's correction for multiple comparisons test with a 95% confidence interval.

RESULTS

Exposure to human ALF alters the expression of cell envelope PIMs/LM/ManLAM

biosynthesis genes in *M.tb*

Our laboratory has previously demonstrated that ALF hydrolases modify the *M.tb* cell envelope,

and that these cell envelope modifications occur in as little as 15 min of *M.tb* being in contact with

ALF and are maintained up to 24 h, without compromising *M.tb* viability [19]. Importantly, these

ALF-driven alterations of the *M.tb* cell envelope had an effect on *M.tb* infection outcomes *in vitro*

and *in vivo* [19-23]. Based on these results, we sought to determine if *M.tb* alters its cell wall

biosynthetic pathways as a direct consequence of ALF exposure. Because one of the main *M.tb*

cell envelope components, ManLAM, is decreased by the action of ALF hydrolases [19] and PIMs,

LM and ManLAM are thought to be part of the same biosynthetic pathway, we first selected a few

key *M.tb* cell envelope genes associated with biosynthesis of initial mannose donors GDP-Man

[*manA/Rv3255c, manB/Rv3264c* (previously annotated as *manC*)*, pmmA/ Rv3257c* (previously

annotated as *manB*)*, pmmB/ Rv3308*] and polyprenylphosphate-based mannose or PPM

(*pmm1/Rv2051c*)*,* the transcriptional regulator *whiB2/Rv3260c* [33], and mannosyl-

(*pimA/Rv2610c, pimB/Rv2188c, pimF/ Rv1500*) and arabynosyl- (*embC/Rv3793*) transferases (**Fig**

1 and **Table 1**) [34, 35], all involved in this pathway. Expression was determined by RT-qPCR

from *M.tb* previously exposed to human ALF for 15 min and 12 h. Relative expression was

normalized to housekeeping genes *rpoB* (**Fig 2A**) or *sigA* (**Fig 2B**). *M.tb* preparations exposed to

the same heat-inactivated ALFs were used as reference samples.

The majority of targeted *M.tb* cell envelope genes were moderately upregulated after 15 min of

exposure to ALF when compared to *M.tb* exposed to heat-inactivated ALF (depicted as log$_2$ fold

changes), except for *manA* that was slightly downregulated and *manB* and *pimB* that did not show

any changes (**Fig 2**). Indeed, *ppm1* (polyprenol-monophosphomannose synthase), which transfers

mannose from GDP-mannose to endogenous PPM [36, 37] in the PIMs/LM/ManLAM

biosynthesis pathway, showed the highest $\log_2$ fold change (1.84 and 1.89 when using *rpoB* or

sigA as reference genes, respectively) after 15 min of exposure compared to heat-inactivated ALF,

although its expression was too low to be detected in our qPCR assay after 12 h (depicted as ND,

Fig 2). Similarly, *manA* was below the detection limit at 12 h and expression differences could not

be quantified (ND, **Fig 2**). A few of the tested genes increased their expression at 12 h post-ALF

exposure (*manB*, *pimB*, and *embC*) compared to heat-inactivated ALF, while others showed no

major transcriptional changes or were slightly downregulated.

When comparing 15 min *vs.* 12 h post-ALF exposure, some PIMs/LM/ManLAM biosynthesis-

associated genes decreased their expression (*pmmA, whiB2, pimF, pimA*) while others increased

(*manB, pimB, embC*) (**Fig 2** and **Table 2**). This suggests a temporal and dynamic adaptation of

M.tb's cell envelope components in response to ALF, in agreement with previous reports showing

cell envelope remodeling of *M.tb* during infection and in response to different environmental

conditions [33, 34, 38-43]. *pmmB* expression was too low to be detected by our qPCR assay at

either timepoints tested, thus no $\log_2$ fold changes are shown.

Effects of A- *vs.* E-ALF exposure on the expression of cell envelope biosynthesis genes in *M.tb*

Since we observed changes in the expression of key *M.tb* cell envelope genes in response to ALF,

and based on our previous publications showing that ALF status influences *M.tb*-host interactions

via multiple factors, including age [22, 24], we next sought to determine if contact with different

host ALFs would result in different transcriptional profiles of the *M.tb* cell envelope. We exposed

M.tb Erdman to individual A-ALFs or E-ALFs for 12 h as described in our previous work [19],

and calculated the relative expression of 83 genes related to cell envelope biosynthesis using a

multiplex qPCR assay in a Biomark HD platform [44]. Results from some of the genes were

compared to the previous targeted qPCR assay, showing concordance in gene expression between

the two qPCR methods used in this study (**Supplemental Figure 1**). Small differences observed

could be attributed to inherent variability within human ALF, since we used different human

donors.

Genes selected for the multiplex qPCR are related to the biosynthesis of key *M.tb* cell envelope

lipid and carbohydrate components (**Supplemental Table 1**). For lipid metabolism, we targeted

genes from the following pathways: fatty acid metabolism, glycerolipid and glycerophospholipid

metabolism, phosphatidylinositol (PI, precursor for more complex glycolipids such as PIM and

LAM), PIMs/LM/ManLAM biosynthesis, mycolic acid biosynthesis, and linoleic acid metabolism

[9]. For carbohydrate metabolism, we included genes related to: carbohydrate biosynthesis,

glycolysis and gluconeogenesis, mannose and fructose metabolism, galactose metabolism, citrate

cycle or TCA, glyoxylate and dicarboxylate metabolism, and inositol phosphate metabolism [45].

Gene names, with corresponding function, pathway and functional categories, as well as primer

sequences are listed in **Supplemental Table 1**.

As shown in **Fig 3A**, most of the cell envelope genes associated with lipid metabolism from the

different *M.tb*-targeted pathways were significantly upregulated in *M.tb* exposed to A-ALFs (A_1

to A_3, three different adult donors) when compared to the same *M.tb* strain exposed to E-ALFs (E_1

to E_3, three different elderly donors), with A_1-ALF having the highest $\log_2$ fold changes (**Fig 3A**

and **Supplemental Table 2**). Only a few genes were downregulated in *M.tb* exposed to A-ALFs,

including *fabH* and *fadD25* (FA metabolism, the latter involved in lipid degradation), and *adhE1*,

glpK, and *cdsA* (glycerolipid and glycerophospholipid synthesis). ATP-binding cassette

transporter Rv1747, thought to be involved in the export of lipooligosaccharides (LOS) through

the mycobacterial membrane, was also downregulated in A-ALF-exposed *M.tb*, whereas negative

regulator of Rv1747, named Rv2623 [46], showed increased expression (**Fig. 3A**). Interestingly,

M.tb exposed to two of the E-ALFs showed upregulation of Rv1747 (**Fig. 3A**), contrary to A-ALF

exposure, indicating a potential increase in *M.tb* PIM export after exposure to elderly ALF. All

genes involved in mycolic acid synthesis tested in this study were upregulated in *M.tb* exposed to

A-ALF. Finally, exposure to E-ALFs showed lesser effects on the overall expression of *M.tb* lipid

metabolism when compared to A-ALFs, with several genes showing moderate upregulation

(especially in E_1-ALF, which showed the highest upregulation among the elders for most of the

genes), and others with no effects or even decreased expression (**Fig. 3A**).

A similar trend was observed for carbohydrate metabolism genes (**Fig. 3B**), where most of the

genes from the different pathways studied showed significant increased expression in *M.tb*

exposed to A-ALFs when compared to E-ALFs. Particularly, 8 out of 12 genes associated with

fructose metabolism and the mannose donor (GDP-Man/PPM) biosynthesis pathways were highly

expressed in A-ALF-exposed *M.tb*, with only 2 genes downregulated (*pmmA*, which converts D-

mannose 1-phosphate in D-mannose 6-phosphate, and *pfkA*, also a key enzyme involved in

glycolysis) [47], and 2 other genes with variable expression across different A-ALF-exposed *M.tb*

(*ppm1* and *fbA*) (**Fig 3B**). Galactose metabolism genes (*galK, galT, galU*, and *aglA*) essential for

the biosynthesis of the cell envelope galactan core [48], were also upregulated after exposure to

A-ALFs, while no changes or even decreased expression were observed in *M.tb* exposed to E-

ALFs. Interestingly, predicted alpha-glucosidase *aglA* was significantly increased after exposure

to E_3-ALF (**Fig 3B**). Further, all tricarboxylic acid cycle (TCA)-associated genes were upregulated

by A-ALF exposure, with the exception of *sdhC* (membrane-anchored subunit of Sdh2, implicated

in *M.tb* growth linked to the TCA cycle under hypoxia conditions) [49] that was downregulated

upon both A- and E-ALF *M.tb* contact. Glyoxylate and dicarboxylate genes were decreased in E-

ALFs when compared to A-ALFs-exposed *M.tb*, whereas the only inositol phosphate gene tested

ino1, catalyzing the first step in inositol synthesis for the production of major thiols and cell wall

lipoglycans [50], had decreased expression after *M.tb* exposure to all ALFs tested, except for E_2-

ALF (**Fig 3B**).

In addition to lipid and carbohydrate pathways, we studied genes belonging to other categories

such as amino acid metabolism, redox homeostasis, and transcriptional regulators (**Fig. 3C** and

Supplemental Table 1). Results indicate that *M.tb* exposed to A-ALFs have major transcriptional

changes compared to *M.tb* exposed to E-ALFs. Indeed, all genes associated with amino acid

metabolism were highly upregulated in *M.tb* exposed to A-ALFs, except for *serC* (serine

metabolism, **Fig 3C**) [51]. Other cell wall-associated genes were upregulated in *M.tb* exposed to

A-ALFs including the putative membrane protein EccB3, part of the ESX-3 secretion system,

important for zinc and iron uptake and homeostasis [52], and the cutinase precursor Cfp21, a

lipolytic enzyme with immunogenic properties shown to elicit T and B cell responses [53, 54].

Similarly, genes *ahpC* and *sodC*, involved in the oxidative stress response [55, 56], and

transcriptional regulators *dosR* and *phoP* had significantly increased expression in *M.tb* exposed

to A-ALFs, whereas *virS* [57] was not significantly increased. Overall, our results demonstrate that

exposure to functional A-ALF results in broad changes in *M.tb* cell envelope biosynthesis,

suggesting highly dynamic cell envelope changes with constant remodeling within the lung

alveolar environment, whereas *M.tb* exposed to dysfunctional E-ALF [22, 24, 25] showed more

limited effects.

DISCUSSION

The cell envelope of *M.tb* is mainly composed of lipids and carbohydrates, and constitutes a

dynamic structure known to adapt to the changing local host environment, especially during

different stages of infection [8, 10]. Human ALF contains hydrolases whose homeostatic function

is to maintain lung health. We have demonstrated that healthy adult individuals have up to 17

hydrolase activities capable of altering the *M.tb* cell wall. Indeed, exposure of *M.tb* to these adult

ALFs significantly alters the *M.tb* cell wall reducing the content of two major cell envelope

components, ManLAM and TDM, without compromising *M.tb* viability [19]. Importantly, these

hydrolase activities are decreased in ALFs from healthy elders [24]. Thus, here, we demonstrate

for the first time that exposure to human ALF, the first environment encountered by *M.tb* upon

infection, results in transcriptional changes in key *M.tb* cell envelope biogenesis genes. Further,

exposure to healthy A-ALFs resulted in increased expression of most of the *M.tb* cell envelope-

associated genes from lipid, carbohydrate and amino acid metabolic pathways, among others. In

contrast, *M.tb* exposed to E-ALFs from healthy elderly donors, which we demonstrated constitutes

a more oxidized, pro-inflammatory and dysfunctional environment [22, 24], did not show many

significant changes in gene expression.

We first assessed essential genes from the PIMs/LM/ManLAM synthesis and the mannose

metabolism pathway (carbohydrate metabolism) (**Figs 1** and **2**), since our previous studies showed

a decrease of ManLAM in the cell envelope of *M.tb* after ALF exposure [19]. PIM/LM/ManLAM

molecules are essential to regulate *M.tb* recognition, uptake, survival and modulate the host

immune response [58, 59]. Indeed, ManLAM has been shown to block phagosome-lysosome (P-

L) fusion by inhibiting the Ca^{2+}/Calmodulin phosphatidyl-inositol-3-kinase (PI3K) hvps34

pathway, promoting *M.tb* intracellular survival [60-62]. The biosynthesis of these molecules

depends on mannose donors such as GPD-Man and PPM. Previous studies suggest that mannose

donor levels are altered during the course of the infection. Indeed, *M.tb* mannose donor

biosynthesis genes had increased expression levels 2 h after macrophage infection and then

gradually decreased [33]. These were also found upregulated in an *in vitro* granuloma model [34].

In our study, most of the genes involved in the PIM/LM/ManLAM biosynthesis pathway were

upregulated after exposure to A-ALFs (**Figs 3A** and **3B**). We speculate that *M.tb* is trying to

compensate for the loss of ManLAM and other mannose-containing cell envelope surface

components due to the action of A-ALF hydrolases [19], with major implications for disease

progression. In this regard, bacilli exposed to functional A-ALF might get taken up by antigen-

presenting cells (APCs) before they can reconstitute ManLAM and will be cleared, while bacilli

that rapidly upregulate and replenish ManLAM on the cell surface before its encounter with APCs

will be able to block P-L fusion and survive withing the host cells. Conversely, exposure to E-

ALFs did not have major transcriptional effects in genes from both mannose donors and

PIM/LM/ManLAM biosynthesis pathways (**Figs 3A** and **3B)**, likely because E-ALF have less

hydrolase activities [24]. This E-ALF deficiency in hydrolase activities could be directly linked to

E-ALF being a highly oxidative environment [22, 24], therefore impacting the *M.tb* cell envelope

and subsequent remodeling to a much lesser degree during infection. Since removal of surface

lipids in *M.tb* enhances trafficking to acidic compartments [61], fewer ALF-driven alterations after

E-ALF exposure might partially explain why *M.tb* replicates faster in the elderly lung environment

by residing in its protective intracellular niche [22, 25]. Further studies will be necessary to

determine the specific impact of elderly ALF hydrolases on the *M.tb* cell envelope.

Most lipid metabolism genes associated with mycolic acid synthesis, fatty acid metabolism,

glycerolipid and glycerophospholipid synthesis, and linoleic acid metabolism were also

significantly upregulated in A-ALF compared to E-ALF-exposed *M.tb* (**Fig 3A**). A similar trend

was observed for carbohydrate metabolism, amino acid metabolism, and genes involved in redox

homeostasis, stress response, and transcriptional regulation (**Figs 3B** and **3C**). Nitrogen and amino

acid metabolism are important for *M.tb* pathogenesis and host colonization during infection, where

intracellular bacteria exploit host nitrogen sources for growth and replication [63, 64]. Importantly,

amino acids acquired from the host such as Ala and Gly might be directly assimilated for the

synthesis of cell wall biomass and incorporated into the PG layer [65-67]. Transcriptional

regulators DosR and PhoP also showed increased expression upon contact with functional A-ALF.

These proteins are part of two-component systems implicated in a large number of *M.tb* adaptive

responses, such as low oxygen levels during dormancy and persistence in the granuloma

environment [68, 69], but they also play key roles as regulators of *M.tb* virulence [70, 71].

Interestingly, DosR has been shown to play a role in lipid accumulation during oxidative stress

and iron starvation in certain *M.tb* clinical strains [72, 73].

These results indicate that the *M.tb* response to the ALF environment is not only limited to *M.tb*

cell envelope remodeling, but also potentially affects overall *M.tb* metabolism and virulence [19-

25, 27], with major implications in infection progression and TB disease outcome. Further global

transcriptomic studies will be able to provide clues as to which other metabolic pathways are

altered by the human ALF environment, with the potential to decipher key upregulated bacterial

determinants during early stages of infection that can be targeted for the development of new

preventative and therapeutic strategies [74]. Indeed, *M.tb* is exposed to ALF during the first

infection stages upon deposition to the alveolar space, but also when escaping from necrotic cells

or in cavities during active TB episodes leading to transmission [17]. Thus, it is plausible that *M.tb*

adapts its cell envelope to the alveolar environment, upregulating the expression of specific genes

to compensate the changes generated upon contact with ALF hydrolases and thus, determining

interactions with host cells. In this regard, timing is expected to be important, as *M.tb* bacilli not

able to restore its cell wall constitution before encountering antigen presenting cells such as

alveolar macrophages might be cleared [19-21].

Key genes such as Rv1747, thought to participate in the export of PIMs to the cell envelope through

negative regulation by stress protein Rv2623 [46], were highly downregulated in *M.tb* exposed to

A-ALFs. Indeed, an ΔRv1747 mutant showed decreased levels of PIMs and was growth-

attenuated, while ΔRv2623 had enhanced PIM expression and was hypervirulent in mice [75, 76].

Our data suggest that contact with functional A-ALF reduces the expression of PIM transporter

Rv1747 through increased expression of Rv2623, modulating the export of immunomodulatory

PIMs and potentially influencing bacterial growth and virulence.

Taken together, these results suggest that *M.tb* compensates for the loss of cell surface components

(due to the action of ALF hydrolases) by upregulating and activating different cell envelope

biosynthesis pathways to rebuild its cell wall, at the detriment of downregulating some key genes

(*e.g.* Rv1747) involved in the transport of cell envelope components to the surface [19-21, 23, 46].

This shift between ALF innate homeostatic mechanisms and *M.tb* countermeasures in the ALF

microenvironment, dependent on the host ALF status (A-AF *vs.* E-ALF), will likely determine

subsequent interactions between *M.tb* and host cells, as well as intracellular trafficking and

infection outcomes [19-21]. Further studies are needed to provide better insight to why E-ALF-

exposed *M.tb* replicates faster than A-ALF-exposed *M.tb* in both professional and non-professional

phagocytes, and to explain why E-ALF status in old age enhances *M.tb* infection *in vitro* and *in*

vivo [22, 24, 25] and contributes to elders being more susceptible to respiratory infections in

general. Finally, it is important to consider that the cell envelope composition of *M.tb* is strain-

specific, with differences observed in *M.tb* strains from different lineages, and thus, different ALF-

driven alterations in *M.tb* metabolism may drive different infection progression [77].

In summary, our study provides evidence that *M.tb* contact within the ALF shapes the composition

of its cell envelope which, depending on the ALF status ('functional A-ALF' *vs.* 'dysfunctional E-

ALF') [22, 24, 25], is likely to define subsequent *M.tb*-host cell interactions. Indeed, E-ALF-

exposed *M.tb* presented minimal transcriptional changes when compared to A-ALF-exposed *M.tb,*

which we speculate provides a fitness advantage to *M.tb* as its cell wall remains intact, and thus, it

has the energy reserves required to efficiently infect and replicate faster within host cells of elderly

individuals. In contrast, *M.tb* undergoes significant alterations on its cell wall (significant loss of

virulent factors ManLAM and TDM, among others) upon exposure to A-ALF [19]. This triggers

a greater transcriptional change that we interpret as efforts of A-ALF-exposed *M.tb* to reprogram

its metabolism to quickly rebuild its cell wall, specifically upregulating genes involved in the

biosynthesis of *M.tb* virulent factors such as ManLAM, with the ensuing energy requirements.

This can be detrimental for A-ALF-exposed *M.tb* and favor host cells to control infection better in

adult individuals. Future studies will investigate the metabolic status of E-ALF-exposed *M.tb* after

infection of professional and non-professional phagocytes *in vitro* and *in vivo*, and will correlate

bacterial and host determinants associated with increased susceptibility to infection in old age.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHORS CONTRIBUTIONS

AAG performed experiments and analyzed data. AGV analyzed data. AOF processed BAL

samples. JP and DJM performed the bronchoalveolar lavage procedures in humans. AAG and JBT

conceptually developed the study and wrote the manuscript. JT and LSS provided critical analysis

of the data and editing of the manuscript. JT, LSS, and JBT provided funding. All authors read and

approved the final version of this manuscript.

FUNDING

This study was supported by the National Institute on Aging (NIA) of the National Institutes of

Health (NIH) under Award Number P01AG051428 to JT, LSS, and JBT. JBT was partially

supported by the Robert J. Kleberg, Jr. and Helen C. Kleberg Foundation. The content is solely

the responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health.

ACKNOWLEDGEMENTS

We want to thank Sean Vargas from the University of Texas San Antonio (UTSA) Genomics Core

for his assistance with the multiplex qPCR assay (Biomark HD).

TABLES

Table 1. List of primers targeting the PIMs/LM/ManLAM biosynthesis pathways in *M.tb*.

Locus tag H$_{37}$R$_v$	Gene name	Product and Function	Primer sequence (5′ to 3′)	Tm (°C)	Ref.
Rv0667	*rpoB*	Reference gene	rpoB-F: CCTGGAAGAGGTGCTCTACG	60	[33]
			rpoB-R: GGGAAGTCACCCATGAACAC	60	[33]
Rv2703	*sigA*	Reference gene	sigA-F: CTCGGTTCGCGCCTACCTCA	68	[35]
			sigA-R: GCGCTCGCTAAGCTCGGTCA	68	[35]
Rv3255c	*manA*	mannose-6-phosphate isomerase / GDP-Man biosynthesis: formation of Man-6-P	manA-F: GTTCACCACCTGGATTACCG	60	[33]
			manA-R: AACCCTCGGTGCATAACAAG	60	[34]
Rv3264c	*manB*[a]	D-alpha-D-mannose-1-phosphate guanylyltransferase / GDP-Man biosynthesis: formation of GDP-Man	manB-F: ACATCGCCGTTAAACACCAT	60	[34]
			manB-R: GTTCCTCACCCATCTGCTGT	60	[34]
Rv3257c	*pmmA*[b]	phosphomannose mutase / GDP-Man biosynthesis: formation of Man-1-P	pmmA-F: GATCACGTTGTGGATGATGG	60	[33]
			pmmA-R: GTGGATCTGCAGGCCTATGT	60	[34]
Rv3308	*pmmB*	phosphomannose mutase / GDP-Man biosynthesis: formation of Man-1-P	pmmB-F: ATACAGATCACGGCGTCACA	60	[34]
			pmmB-R: CGCTGGATATAACGGTCGAT	60	[34]
Rv2051c	*ppm1*	Polyprenol-monophosphomannose synthase / PPM biosynthesis	pmm1-F: TGGTTGAAGTCGATCCTTCC	60	[33]
			pmm1-R: GCGAACAAGACCAGGCATATG	63	[33]
Rv3260c	*whiB2*	Transcript. regulatory protein	whiB2-F: CCATTCGAGGAACCTCTGC	61	[33]
			whiB2-R: CAGGGCGTACTCCAGACACT	60	[33]
Rv2610c	*pimA*	alpha-(1-2)-phosphatidylinositol mannosyl-transferase / PIM biosynthesis (1st step)	pimA-F: CCGCACTGCCTGATTACTTT	60	[34]
			pimA-R: CGGCTCGTGTAGATGCAGTA	60	[34]

Rv2188c	*pimB*	alpha-(1-6)-phosphatidylinositol mannosyl- transferase / PIM biosynthesis (2nd step)	pimB-F: CTCGGTGGTCAAGGTACTCG	61	[34]
			pimB-R: GTGGTCACCTTTGGGAACAT	60	[34]
Rv1500	*pimF*	glycosyltransferase / LM/ManLAM biosynthesis	pimF-F: CGCCGACGTAGTATTTGGTT	60	[34]
			pimF-R: TGCGTACATAGTCGGCTGTC	60	[34]
Rv3793	*embC*	Arabynosyl-tranferase / ManLAM biosynthesis	embC-F: ATCACCGAGCTGCTGATG	58	[35]
			embC-R: TGCGAGTCACCGTTCCTA	59	[35]

a Previously annotated as *manC*.
b Previously annotated as *manB*.

Table 2. Comparison of log$_2$ FC between 15 min and 12 h of ALF exposure for targeted qPCR

genes. Differences in the expression of targeted *M.tb* cell envelope genes between 15 min and 12

h after ALF exposure were calculated (log$_2$ FC at 12 h – log$_2$ FC at 15 min) for each reference

gene. Average differences for *rpoB* and *sigA* are also shown.

Genes	15 min *vs.* 12 h (*rpoB*)	15 min *vs.* 12 h (*sigA*)	Mean *rpoB* and *sigA*
manB	0.107	0.425	0.266
pmmA	-0.580	-0.421	-0.500
whiB2	-0.461	-0.276	-0.368
pimA	-0.418	-0.224	-0.321
pimB	-0.019	0.252	0.116
pimF	-0.552	-0.406	-0.479
embC	-0.024	0.222	0.099

FIGURE LEGENDS

Fig 1. *M.tb* GDP-Man, PPM and PIM/LM/ManLAM biosynthetic pathways. GDP-Man can

be biosynthesized directly from gluconeogenesis, through formation of Fruc-6-P from Glc-6-P (by

the action of Glc-6-P isomerase), formation of Man-6-P (by the action of Man-6-P isomerase or

ManA/Rv3255c), formation of Man-1-P (by the action of phosphomutases PmmA/Rv3257c,

previously annotated as ManB, and PmmB/Rv3308), and finally formation of GDP-Man (by the

action of Man-1-P guanylyn-transferase or ManB/Rv3264c, previously annotated as ManC).

Further, Man-6-P can also be directly formed from Man by the action of a hexokinase (HK). PPM

is formed from GDP-Man by the action of polyprenyl monophosphomannose synthase or

Ppm1/Ppm2). Further, CDP-DAG together with inositol by the action of PI synthase (Rv2612c)

forms PI, which is further mannosylated by several mannosyltransferases (PimA to PimF) to

generate higher orders PIMs using GDP-Man and PPM as mannose donors. At one point, from

PIM_4 and using PPM as the major mannose donor, PIM_4 is heavily mannosylated by an undisclosed

number of mannosyl-transferases generating LM, and further arabinosylated with arabinosyl-

transferases generating LAM, which can be further mannose-capped by the mannosyl-transferases

action. LAM can also contain methylthio-D-xylose (MTX) capping motifs [78, 79], where MtxT

(Rv0541c) transfers MTX to the mannoside caps of LAM [80]. Note: For simplicity,

acyltransferases (*e.g.* Rv2610c) are not depicted, and neither is the formation of MTX-P-C50 by

MtxS (Rv0539).

Fig 2. Relative expression of selected PIM/LM/ManLAM biosynthesis genes in *M.tb* $H_{37}R_v$

after exposure to ALF. *M.tb* was exposed for 15 min and 12 h to healthy human ALF (n=2

biological replicates, from two independent ALF donors), using *rpoB* (**A**) or *sigA* (**B**) as reference

genes. Expression values are shown as log_2 fold changes, and were calculated using the $2^{-\Delta\Delta CT}$

method (ALF-exposed *M.tb* *vs.* heat-inactivated ALF-exposed *M.tb*) and plotted as the mean ±

SEM using Prism v9. ND: not detected (below limit of detection).

Fig 3. Relative expression of cell envelope biogenesis and metabolism genes in *M.tb* Erdman

exposed to A-ALF and E-ALF. Heatmap showing relative expression of cell wall genes

associated to (**A**) lipid metabolism; (**B**) carbohydrate metabolism; and (**C**) other pathways, in *M.tb*

after being exposed to A-ALFs (n=3 biological replicates, A_1- to A_3-ALFs), or to E-ALFs (n=3

biological replicates, E_1- to E_3-ALFs). Expression was normalized using *rpoB* as reference gene

and calculated using the $2^{-\Delta\Delta CT}$ method (ALF-exposed *M.tb vs.* heat-inactivated ALF-exposed

M.tb). Heatmap was constructed using Prism v9, with downregulated genes in blue (0 to 1 $\log_2$

fold changes) and upregulated genes in yellow (1 to 5 or more $\log_2$ fold changes). Genes are

grouped based on their assigned pathways (see **Supplemental Table 1**). Note that genes *pimB*,

accD3, *adhC* (lipid metabolism), *pmmB*, Rv0794c (carbohydrate metabolism), and *metZ* (others)

were below the limit of detection in one or more of the samples, and have not been included in the

heatmaps.

Fig 1

Fig 2

Fig 3

REFERENCES

1. WHO: **Global Tuberculosis Report 2020**. 2020.
2. Hogan AB, Jewell BL, Sherrard-Smith E, Vesga JF, Watson OJ, Whittaker C, Hamlet A, Smith JA, Winskill P, Verity R *et al*: **Potential impact of the COVID-19 pandemic on HIV, tuberculosis, and malaria in low-income and middle-income countries: a modelling study**. *Lancet Glob Health* 2020, **8**(9):e1132-e1141.
3. Cilloni L, Fu H, Vesga JF, Dowdy D, Pretorius C, Ahmedov S, Nair SA, Mosneaga A, Masini E, Sahu S *et al*: **The potential impact of the COVID-19 pandemic on the tuberculosis epidemic a modelling analysis**. *EClinicalMedicine* 2020, **28**:100603.
4. dos Santos FL, Souza LLL, Bruce ATI, Crispim JD, Arroyo LH, Ramos ACV, Berra TZ, Alves YM, Scholze AR, da Costa FBP *et al*: **Patients' perceptions regarding multidrug-resistant tuberculosis and barriers to seeking care in a priority city in Brazil during COVID-19 pandemic: A qualitative study**. *Plos One* 2021, **16**(4).
5. Marwah V, Peter DK, Ajai Kumar T, Bhati G, Kumar A: **Multidrug-resistant tuberculosis in COVID-19: Double trouble**. *Med J Armed Forces India* 2021, **77**:S479-S482.
6. Tiberi S, Vjecha MJ, Zumla A, Galvin J, Migliori GB, Zumla A: **Accelerating development of new shorter TB treatment regimens in anticipation of a resurgence of multi-drug resistant TB due to the COVID-19 pandemic**. *Int J Infect Dis* 2021.
7. Kalscheuer R, Palacios A, Anso I, Cifuente J, Anguita J, Jacobs WR, Jr., Guerin ME, Prados-Rosales R: **The Mycobacterium tuberculosis capsule: a cell structure with key implications in pathogenesis**. *Biochem J* 2019, **476**(14):1995-2016.
8. Garcia-Vilanova A, Chan J, Torrelles JB: **Underestimated Manipulative Roles of Mycobacterium tuberculosis Cell Envelope Glycolipids During Infection**. *Front Immunol* 2019, **10**:2909.
9. Gago G, Diacovich L, Gramajo H: **Lipid metabolism and its implication in mycobacteria-host interaction**. *Curr Opin Microbiol* 2018, **41**:36-42.
10. Dulberger CL, Rubin EJ, Boutte CC: **The mycobacterial cell envelope - a moving target**. *Nat Rev Microbiol* 2020, **18**(1):47-59.
11. Vincent AT, Nyongesa S, Morneau I, Reed MB, Tocheva EI, Veyrier FJ: **The Mycobacterial Cell Envelope: A Relict From the Past or the Result of Recent Evolution?** *Front Microbiol* 2018, **9**:2341.
12. Chiaradia L, Lefebvre C, Parra J, Marcoux J, Burlet-Schiltz O, Etienne G, Tropis M, Daffe M: **Dissecting the mycobacterial cell envelope and defining the composition of the native mycomembrane**. *Sci Rep* 2017, **7**(1):12807.
13. Nicod LP: **Pulmonary defence mechanisms**. *Respiration* 1999, **66**:2-11.
14. Hiemstra PS, Bals R: **Series introduction: Innate host defense of the respiratory epithelium**. *J Leukoc Biol* 2004, **75**(1):3-4.
15. Williams MC: **Alveolar type I cells: molecular phenotype and development**. *Annu Rev Physiol* 2003, **65**:669-695.
16. Mason RJ: **Biology of alveolar type II cells**. *Respirology* 2006, **11 Suppl**:S12-S15.
17. Torrelles JB, Schlesinger LS: **Integrating Lung Physiology, Immunology, and Tuberculosis**. *Trends Microbiol* 2017, **25**(8):688-697.
18. Liu CH, Liu H, Ge B: **Innate immunity in tuberculosis: host defense vs pathogen evasion**. *Cell Mol Immunol* 2017, **14**(12):963-975.

19. Arcos J, Sasindran SJ, Fujiwara N, Turner J, Schlesinger LS, Torrelles JB: **Human lung hydrolases delineate Mycobacterium tuberculosis-macrophage interactions and the capacity to control infection**. *J Immunol* 2011, **187**(1):372-381.

20. Arcos J, Sasindran SJ, Moliva JI, Scordo JM, Sidiki S, Guo H, Venigalla P, Kelley HV, Lin G, Diangelo L *et al*: **Mycobacterium tuberculosis cell wall released fragments by the action of the human lung mucosa modulate macrophages to control infection in an IL-10-dependent manner**. *Mucosal Immunol* 2017, **10**(5):1248-1258.

21. Arcos J, Diangelo LE, Scordo JM, Sasindran SJ, Moliva JI, Turner J, Torrelles JB: **Lung Mucosa Lining Fluid Modification of Mycobacterium tuberculosis to Reprogram Human Neutrophil Killing Mechanisms**. *J Infect Dis* 2015, **212**(6):948-958.

22. Moliva JI, Duncan MA, Olmo-Fontanez A, Akhter A, Arnett E, Scordo JM, Ault R, Sasindran SJ, Azad AK, Montoya MJ *et al*: **The Lung Mucosa Environment in the Elderly Increases Host Susceptibility to Mycobacterium tuberculosis Infection**. *J Infect Dis* 2019, **220**(3):514-523.

23. Scordo JM, Arcos J, Kelley HV, Diangelo L, Sasindran SJ, Youngmin E, Wewers MD, Wang SH, Balada-Llasat JM, Torrelles JB: **Mycobacterium tuberculosis Cell Wall Fragments Released upon Bacterial Contact with the Human Lung Mucosa Alter the Neutrophil Response to Infection**. *Front Immunol* 2017, **8**:307.

24. Moliva JI, Rajaram MV, Sidiki S, Sasindran SJ, Guirado E, Pan XJ, Wang SH, Ross P, Jr., Lafuse WP, Schlesinger LS *et al*: **Molecular composition of the alveolar lining fluid in the aging lung**. *Age (Dordr)* 2014, **36**(3):9633.

25. Olmo-Fontánez AM, Scordo JM, Garcia-Vilanova A, Maselli DJ, Peters JI, Restrepo BI, Clemens DL, Turner J, Schlesinger LS, Torrelles JB: **Human alveolar lining fluid from the elderly promotes Mycobacterium tuberculosis growth in alveolar epithelial cells and bacterial translocation into the cytosol**. *bioRxiv* 2021:2021.2005.2012.443884.

26. Arcos J, Sasindran SJ, Fujiwara N, Turner J, Schlesinger LS, Torrelles JB: **Human Lung Hydrolases Delineate Mycobacterium tuberculosis-Macrophage Interactions and the Capacity to Control Infection**. *J Immunol* 2011, **187**(1):372-381.

27. Scordo JM, Olmo-Fontanez AM, Kelley HV, Sidiki S, Arcos J, Akhter A, Wewers MD, Torrelles JB: **The human lung mucosa drives differential Mycobacterium tuberculosis infection outcome in the alveolar epithelium**. *Mucosal Immunol* 2019, **12**(3):795-804.

28. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2(T)(-Delta Delta C) method**. *Methods* 2001, **25**(4):402-408.

29. You FM, Huo N, Gu YQ, Luo MC, Ma Y, Hane D, Lazo GR, Dvorak J, Anderson OD: **BatchPrimer3: a high throughput web application for PCR and sequencing primer design**. *BMC Bioinformatics* 2008, **9**:253.

30. Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon SV, Eiglmeier K, Gas S, Barry CE, 3rd *et al*: **Deciphering the biology of Mycobacterium tuberculosis from the complete genome sequence**. *Nature* 1998, **393**(6685):537-544.

31. Camus JC, Pryor MJ, Medigue C, Cole ST: **Re-annotation of the genome sequence of Mycobacterium tuberculosis H37Rv**. *Microbiology (Reading)* 2002, **148**(Pt 10):2967-2973.

32. Lew JM, Kapopoulou A, Jones LM, Cole ST: **TubercuList--10 years after**. *Tuberculosis (Edinb)* 2011, **91**(1):1-7.

33. Keiser TL, Azad AK, Guirado E, Bonacci R, Schlesinger LS: **Comparative**
transcriptional study of the putative mannose donor biosynthesis genes in virulent
Mycobacterium tuberculosis and attenuated Mycobacterium bovis BCG strains.
Infect Immun 2011, **79**(11):4668-4673.
34. Guirado E, Mbawuike U, Keiser TL, Arcos J, Azad AK, Wang SH, Schlesinger LS:
Characterization of host and microbial determinants in individuals with latent
tuberculosis infection using a human granuloma model. *mBio* 2015, **6**(1):e02537-
02514.
35. Papavinasasundaram KG, Chan B, Chung JH, Colston MJ, Davis EO, Av-Gay Y: **Deletion**
of the Mycobacterium tuberculosis pknH gene confers a higher bacillary load during
the chronic phase of infection in BALB/c mice. *J Bacteriol* 2005, **187**(16):5751-5760.
36. Rana AK, Singh A, Gurcha SS, Cox LR, Bhatt A, Besra GS: **Ppm1-encoded polyprenyl**
monophosphomannose synthase activity is essential for lipoglycan synthesis and
survival in mycobacteria. *Plos One* 2012, **7**(10):e48211.
37. Gurcha SS, Baulard AR, Kremer L, Locht C, Moody DB, Muhlecker W, Costello CE,
Crick DC, Brennan PJ, Besra GS: **Ppm1, a novel polyprenol monophosphomannose**
synthase from Mycobacterium tuberculosis. *Biochem J* 2002, **365**(Pt 2):441-450.
38. Lv X, Ma CY, Yan ZF, Deng R, Ai XF, Su T, Xiang XH, Li W: **The Mycobacterium**
tuberculosis protein Rv2387 is involved in cell wall remodeling and susceptibility to
acidic conditions. *Biochem Bioph Res Co* 2018, **503**(2):625-630.
39. Stokas H, Rhodes HL, Purdy GE: **Modulation of the M. tuberculosis cell envelope**
between replicating and non-replicating persistent bacteria. *Tuberculosis* 2020, **125**.
40. Howard NC, Marin ND, Ahmed M, Rosa BA, Martin J, Bambouskova M, Sergushichev
A, Loginicheva E, Kurepina N, Rangel-Moreno J *et al*: **Mycobacterium tuberculosis**
carrying a rifampicin drug resistance mutation reprograms macrophage metabolism
through cell wall lipid changes. *Nat Microbiol* 2018, **3**(10):1099-+.
41. Bacon J, Alderwick LJ, Allnutt JA, Gabasova E, Watson R, Hatch KA, Clark SO, Jeeves
RE, Marriott A, Rayner E *et al*: **Non-Replicating Mycobacterium tuberculosis Elicits a**
Reduced Infectivity Profile with Corresponding Modifications to the Cell Wall and
Extracellular Matrix. *Plos One* 2014, **9**(2).
42. Chang DPS, Guan XL: **Metabolic Versatility of Mycobacterium tuberculosis during**
Infection and Dormancy. *Metabolites* 2021, **11**(2).
43. Yang L, Sinha T, Carlson TK, Keiser TL, Torrelles JB, Schlesinger LS: **Changes in the**
major cell envelope components of Mycobacterium tuberculosis during in vitro
growth. *Glycobiology* 2013, **23**(8):926-934.
44. Spurgeon SL, Jones RC, Ramakrishnan R: **High throughput gene expression**
measurement with real time PCR in a microfluidic dynamic array. *Plos One* 2008,
3(2):e1662.
45. Rhee KY, de Carvalho LP, Bryk R, Ehrt S, Marrero J, Park SW, Schnappinger D,
Venugopal A, Nathan C: **Central carbon metabolism in Mycobacterium tuberculosis:**
an unexpected frontier. *Trends Microbiol* 2011, **19**(7):307-314.
46. Glass LN, Swapna G, Chavadi SS, Tufariello JM, Mi K, Drumm JE, Lam TT, Zhu G, Zhan
C, Vilcheze C *et al*: **Mycobacterium tuberculosis universal stress protein Rv2623**
interacts with the putative ATP binding cassette (ABC) transporter Rv1747 to
regulate mycobacterial growth. *PLoS Pathog* 2017, **13**(7):e1006515.

47. Phong WY, Lin W, Rao SP, Dick T, Alonso S, Pethe K: **Characterization of phosphofructokinase activity in Mycobacterium tuberculosis reveals that a functional glycolytic carbon flow is necessary to limit the accumulation of toxic metabolic intermediates under hypoxia**. *Plos One* 2013, **8**(2):e56037.

48. Konyarikova Z, Savkova K, Kozmon S, Mikusova K: **Biosynthesis of Galactan in Mycobacterium tuberculosis as a Viable TB Drug Target?** *Antibiotics (Basel)* 2020, **9**(1).

49. Hartman T, Weinrick B, Vilcheze C, Berney M, Tufariello J, Cook GM, Jacobs WR, Jr.: **Succinate dehydrogenase is the regulator of respiration in Mycobacterium tuberculosis**. *PLoS Pathog* 2014, **10**(11):e1004510.

50. Movahedzadeh F, Smith DA, Norman RA, Dinadayala P, Murray-Rust J, Russell DG, Kendall SL, Rison SC, McAlister MS, Bancroft GJ *et al*: **The Mycobacterium tuberculosis ino1 gene is essential for growth and virulence**. *Mol Microbiol* 2004, **51**(4):1003-1014.

51. Haufroid M, Wouters J: **Targeting the Serine Pathway: A Promising Approach against Tuberculosis?** *Pharmaceuticals (Basel)* 2019, **12**(2).

52. Famelis N, Rivera-Calzada A, Degliesposti G, Wingender M, Mietrach N, Skehel JM, Fernandez-Leiro R, Bottcher B, Schlosser A, Llorca O *et al*: **Architecture of the mycobacterial type VII secretion system**. *Nature* 2019, **576**(7786):321-325.

53. Wang C, Chen Z, Fu R, Zhang Y, Chen L, Huang L, Li J, Shi C, Fan X: **A DNA vaccine expressing CFP21 and MPT64 fusion protein enhances BCG-induced protective immunity against Mycobacterium tuberculosis infection in mice**. *Med Microbiol Immunol* 2011, **200**(3):165-175.

54. Renier W, Bourdin A, Rubbo PA, Peries M, Dedieu L, Bendriss S, Kremer L, Canaan S, Terru D, Godreuil S *et al*: **B cells response directed against Cut4 and CFP21 lipolytic enzymes in active and latent tuberculosis infections**. *Plos One* 2018, **13**(4):e0196470.

55. Wong CF, Shin J, Subramanian Manimekalai MS, Saw WG, Yin Z, Bhushan S, Kumar A, Ragunathan P, Gruber G: **AhpC of the mycobacterial antioxidant defense system and its interaction with its reducing partner Thioredoxin-C**. *Sci Rep* 2017, **7**(1):5159.

56. Liao D, Fan Q, Bao L: **The role of superoxide dismutase in the survival of Mycobacterium tuberculosis in macrophages**. *Jpn J Infect Dis* 2013, **66**(6):480-488.

57. Singh R, Singh A, Tyagi AK: **Deciphering the genes involved in pathogenesis of Mycobacterium tuberculosis**. *Tuberculosis (Edinb)* 2005, **85**(5-6):325-335.

58. Turner J, Torrelles JB: **Mannose-capped lipoarabinomannan in Mycobacterium tuberculosis pathogenesis**. *Pathog Dis* 2018, **76**(4).

59. Torrelles JB, Schlesinger LS: **Diversity in Mycobacterium tuberculosis mannosylated cell wall determinants impacts adaptation to the host**. *Tuberculosis (Edinb)* 2010, **90**(2):84-93.

60. Fratti RA, Backer JM, Gruenberg J, Corvera S, Deretic V: **Role of phosphatidylinositol 3-kinase and Rab5 effectors in phagosomal biogenesis and mycobacterial phagosome maturation arrest**. *J Cell Biol* 2001, **154**(3):631-644.

61. Rajni, Rao N, Meena LS: **Biosynthesis and Virulent Behavior of Lipids Produced by Mycobacterium tuberculosis: LAM and Cord Factor: An Overview**. *Biotechnol Res Int* 2011, **2011**:274693.

62. Kang PB, Azad AK, Torrelles JB, Kaufman TM, Beharka A, Tibesar E, DesJardin LE, Schlesinger LS: **The human macrophage mannose receptor directs Mycobacterium**

716   **tuberculosis lipoarabinomannan-mediated phagosome biogenesis**. *J Exp Med* 2005,
717   **202**(7):987-999.

63. Wietzerbin-Falszpan J, Das BC, Gros C, Petit JF, Lederer E: **The amino acids of the cell wall of Mycobacterium tuberculosis var. bovis, strain BCG. Presence of a poly(L-glutamic acid)**. *Eur J Biochem* 1973, **32**(3):525-532.

64. Alderwick LJ, Harrison J, Lloyd GS, Birch HL: **The Mycobacterial Cell Wall--Peptidoglycan and Arabinogalactan**. *Cold Spring Harb Perspect Med* 2015, **5**(8):a021113.

65. Cava F, de Pedro MA, Lam H, Davis BM, Waldor MK: **Distinct pathways for modification of the bacterial cell wall by non-canonical D-amino acids**. *Embo J* 2011, **30**(16):3442-3453.

66. Maitra A, Munshi T, Healy J, Martin LT, Vollmer W, Keep NH, Bhakta S: **Cell wall peptidoglycan in Mycobacterium tuberculosis: An Achilles' heel for the TB-causing pathogen**. *Fems Microbiol Rev* 2019, **43**(5):548-575.

67. Garcia-Heredia A, Pohane AA, Melzer ES, Carr CR, Fiolek TJ, Rundell SR, Lim HC, Wagner JC, Morita YS, Swarts BM *et al*: **Peptidoglycan precursor synthesis along the sidewall of pole-growing mycobacteria**. *Elife* 2018, **7**.

68. He Z, De Buck J: **Cell wall proteome analysis of Mycobacterium smegmatis strain MC2 155**. *BMC Microbiol* 2010, **10**:121.

69. Mehra S, Foreman TW, Didier PJ, Ahsan MH, Hudock TA, Kissee R, Golden NA, Gautam US, Johnson AM, Alvarez X *et al*: **The DosR Regulon Modulates Adaptive Immunity and Is Essential for Mycobacterium tuberculosis Persistence**. *Am J Respir Crit Care Med* 2015, **191**(10):1185-1196.

70. Ryndak M, Wang S, Smith I: **PhoP, a key player in Mycobacterium tuberculosis virulence**. *Trends Microbiol* 2008, **16**(11):528-534.

71. Broset E, Martin C, Gonzalo-Asensio J: **Evolutionary landscape of the Mycobacterium tuberculosis complex from the viewpoint of PhoPR: implications for virulence regulation and application to vaccine development**. *mBio* 2015, **6**(5):e01289-01215.

72. Singh A, Crossman DK, Mai D, Guidry L, Voskuil MI, Renfrow MB, Steyn AJ: **Mycobacterium tuberculosis WhiB3 maintains redox homeostasis by regulating virulence lipid anabolism to modulate macrophage response**. *PLoS Pathog* 2009, **5**(8):e1000545.

73. Reed MB, Gagneux S, DeRiemer K, Small PM, Barry CE: **The W-Beijing lineage of Mycobacterium tuberculosis overproduces triglycerides and has the DosR dormancy regulon constitutively upregulated**. *Journal of Bacteriology* 2007, **189**(7):2583-2589.

74. Shaku M, Ealand C, Kana BD: **Cell Surface Biosynthesis and Remodeling Pathways in Mycobacteria Reveal New Drug Targets**. *Front Cell Infect Microbiol* 2020, **10**:603382.

75. Drumm JE, Mi K, Bilder P, Sun M, Lim J, Bielefeldt-Ohmann H, Basaraba R, So M, Zhu G, Tufariello JM *et al*: **Mycobacterium tuberculosis universal stress protein Rv2623 regulates bacillary growth by ATP-Binding: requirement for establishing chronic persistent infection**. *PLoS Pathog* 2009, **5**(5):e1000460.

76. Curry JM, Whalan R, Hunt DM, Gohil K, Strom M, Rickman L, Colston MJ, Smerdon SJ, Buxton RS: **An ABC transporter containing a forkhead-associated domain interacts with a serine-threonine protein kinase and is required for growth of Mycobacterium tuberculosis in mice**. *Infect Immun* 2005, **73**(8):4471-4477.

77. Allue-Guardia A, Garcia JI, Torrelles JB: **Evolution of Drug-Resistant Mycobacterium**
tuberculosis Strains and Their Adaptation to the Human Lung Environment. *Front*
Microbiol 2021, **12**:612675.
78. Treumann A, Xidong F, McDonnell L, Derrick PJ, Ashcroft AE, Chatterjee D, Homans
SW: **5-Methylthiopentose: a new substituent on lipoarabinomannan in**
Mycobacterium tuberculosis. *J Mol Biol* 2002, **316**(1):89-100.
79. Turnbull WB, Shimizu KH, Chatterjee D, Homans SW, Treumann A: **Identification of**
the 5-methylthiopentosyl substituent in Mycobacterium tuberculosis
lipoarabinomannan. *Angew Chem Int Ed Engl* 2004, **43**(30):3918-3922.
80. Angala SK, McNeil MR, Shi L, Joe M, Pham H, Zuberogoitia S, Nigou J, Boot CM,
Lowary TL, Gilleron M *et al*: **Biosynthesis of the Methylthioxylose Capping Motif of**
Lipoarabinomannan in Mycobacterium tuberculosis. *ACS Chem Biol* 2017, **12**(3):682-
691.
774