<u>Copper selects for siderophore-mediated virulence in *Pseudomonas aeruginosa*</u>

Luke Lear[1*], Elze Hesse[2], Angus Buckling[2] and Michiel Vos[1].

[1]European Centre for Environment and Human Health, University of Exeter Medical School,

Penryn, Cornwall, United Kingdom, TR10 9FE

[2]College of Life and Environmental Science, University of Exeter, Penryn, Cornwall, United

Kingdom, TR10 9FE

*Correspondence: ll381@exeter.ac.uk
12

Keywords: evolution of virulence, opportunistic pathogen, metal detoxification, pyoverdine,

coincidental selection.

<u>Abstract</u>

Iron is essential for almost all bacterial pathogens and consequently it is actively withheld by

their hosts. However, the production of extracellular siderophores enables iron sequestration

by pathogens, increasing their virulence. Another function of siderophores is extracellular

detoxification of non-ferrous metals. Here, we experimentally link the detoxification and

virulence roles of siderophores by testing whether the opportunistic pathogen *Pseudomonas*

aeruginosa displays greater virulence after exposure to copper. To do this, we incubated *P.*

aeruginosa under different environmentally relevant copper regimes for either two or twelve

days. Subsequent growth in a copper-free environment removed phenotypic effects, before we

quantified pyoverdine production (the primary siderophore produced by *P. aeruginosa*), and

virulence using the *Galleria mellonella* infection model. Copper selected for increased

pyoverdine production, which was positively correlated with virulence. This effect increased

with time, such that populations incubated with high copper for twelve days were the most

virulent. Replication of the experiment with a non-pyoverdine producing strain of *P.*

aeruginosa demonstrated that pyoverdine production was largely responsible for the change in

virulence. Therefore we here show a direct link between metal stress and bacterial virulence,

highlighting another dimension of the detrimental effects of metal pollution on human health.

Introduction

Iron is essential for the growth of almost all bacteria as it serves as a cofactor for many enzymes (1, 2). However, as iron most commonly exists in the insoluble form Fe^{3+}, it is of relatively low bioavailability in the majority of environments (1-7). It is therefore essential for bacteria to actively sequester Fe^{3+} from the environment (8, 9), with many bacteria producing iron-chelating siderophore compounds to do this (10). Siderophores aid iron recovery by forming extracellular complexes with Fe^{3+} which are then taken up by the cell and reduced to the bioavailable form Fe^{2+} (11). The extracellular nature of siderophores means they can benefit the group around the producer, with only the producer paying the cost of production. As a result, siderophore production can be a cooperative trait subject to social evolution (12, 13). Therefore as increased siderophore production evolves under iron limitation to benefit the producer and its kin, genotypes with reduced production can also evolve to exploit these producers and gain a fitness advantage (12). Due to the importance of iron for microbial growth, one of the first lines of host defence is to withhold iron from invading pathogens (14). This often involves the production of molecules such as transferrin that bind to iron with very high association constants (6, 14-16). Such nutritional immunity can be overcome by pathogens through the production of siderophores, as their very high iron affinity enables 'stealing' iron from the host (16). Siderophores are therefore an important virulence factor in both gram-negative- (17-19) and gram-positive human pathogens (20, 21).

In addition to iron, siderophores are known to bind to a range of potentially toxic metals (22). Crucially, these complexes cannot passively diffuse into the cell (*i.e.* not without active use of transport proteins) and consequently the metal is detoxified extracellularly (23). As a result, siderophore production has been shown to increase in the presence of toxic levels of a wide range of non-ferrous metals (2, 3, 9, 22). As many of the metal ions that can be bound by siderophores are essential in small amounts, siderophore production is adjusted in response to metal bioavailability (24), with either active uptake of metal-siderophore complexes or other metal chelating compounds used to acquire these micronutrients when they are scarce (7, 25). Adjustments in production can be due to both genetic changes and phenotypic plasticity, with genetics dictating the limits of siderophore production and phenotypic plasticity used to regulate production within those limits (26). Although the upper limit of production initially increases with metal toxicity, eventually a threshold is reached as the rising metabolic cost of siderophore production can result in selection for reduced production (8) – analogous to the

evolutionary response to iron deficiency (27). This cost is further exacerbated when competitors of the producer benefit from the siderophore (12). Consequently, when there is strong selection for siderophore production in the community, such as when toxic metals are at high concentrations, per capita siderophore production can be reduced both through phenotypic and genotypic change (8, 27, 28). Phenotypic changes in production occur quickly in response to alterations in the environment, including the abundance of metals and population densities, and allow the cost of production to be minimised (23, 28).

Despite strong evidence both for siderophores playing an important role in metal detoxification (3, 8, 22, 24) and for siderophore production being an important determinant for bacterial virulence (16, 29, 30), whether the presence of toxic metals in non-host environments affects virulence remains untested. Here we aim to experimentally link these dual effects of siderophore production, by testing the effects of copper on the virulence of the opportunistic human pathogen *Pseudomonas aeruginosa*. *P. aeruginosa* is of clinical significance capable of surviving in a range of environments including soil, water and fomites in hospitals (31, 32). In clinical settings it is frequently responsible for severe and fatal infections of patients with cystic fibrosis (CF) (33), burns (34), and immunosuppressive illnesses (35). It is important to note that many CF infections are caused by *P. aeruginosa* acquired from the environment (36), and so an understanding how selection for siderophore production outside of the host could alter virulence is of great importance.

The production of pyoverdine, the main *P. aeruginosa* siderophore, has been shown to rapidly evolve (37). We therefore test virulence after both two days (~7 generations), when little genotypic change is expected to have occurred, and twelve days (~40 generations), where genotypic changes in production are expected (12, 38). To allow us to exclude phenotypic changes to production, at each timepoint (two and twelve days) populations are transferred to a common garden environment (copper free media) for 24 hours prior to assays. By quantifying pyoverdine production in every population before the virulence assay, we are able to test whether its production is associated with virulence. We then further test for an association by repeating the experiment using a *P. aeruginosa* strain that cannot produce either pyoverdine or pyochelin (*P. aeruginosa*'s secondary siderophore (39)), and testing whether copper has the same effect on virulence. Moreover, we test if this association is the same across two environmentally relevant copper concentrations that span those found in agricultural soil (40) and in water (41).

                                  Methods

Experimental design

Pseudomonas aeruginosa PAO1 (42) was grown shaking at 180 rpm for 24 hours at 28°C in

glass microcosms containing 6mL of King's medium B (KB; 10g glycerol, 20g proteose

peptone no. 3, 1.5 g K_2HPO_4,1.5 g $MgSO_4$, per litre). After homogenisation by vortexing, 60μL

of undiluted culture was added to 18 microcosms containing KB mixed with sterilised copper

sulphate ($CuSO_4$; Alfa Aesar, Massachusetts, United States) to final concentrations of either

0.0, 0.1 or 1.0 g/L. Inoculated microcosms were kept static at 28°C for 48 hours before being

thoroughly homogenised and 60μL (1% by volume) transferred into fresh media (Fig.1).

Microcosms were incubated at 28°C, rather than the more common 37°C, in order for our

experiment to be more representative of environmental conditions. Transfers occurred every

two days for twelve days. To control for the physiological effects of copper stress, on days two

and twelve each microcosm was homogenised and 60μL transferred into a common garden

environment (6mL of KB medium without copper). These cultures were grown for 24 hours at

28°C before pyoverdine production was quantified and aliquots were frozen at -80°C in

glycerol at a final concentration of 25% for virulence assays and to quantify density. We then

repeated this experiment with an isogenic mutant strain of *P. aeruginosa* (PAO1Δ*PvdD*Δ*PchEF*

(43)) with the ability to produce pyoverdine, and the secondary siderophore pyochelin, knocked

out (a non-pyoverdine producing strain).

Figure 1 Schematic of the experimental design used to test whether copper selects for siderophore-mediated virulence. Microcosms (n=6 per treatment) containing KB medium at a concentration of either 0.0, 0.1 or 1.0 g/L of copper sulphate ($CuSO_4$) were inoculated with *Pseudomonas aeruginosa* (either the pyoverdine producing or non-pyoverdine producing strain), incubated at 28°C and transferred every two days into fresh media. On days two and twelve, cultures were transferred into copper free medium for 24 hours before being homogenised, their per capita pyoverdine production quantified and frozen in glycerol at a final concentration of 25%. Virulence and density assays were performed using defrosted samples.

Pyoverdine assay

Pyoverdine production was quantified after both two and twelve days. Following 24 hours in a common garden environment, cultures were thoroughly homogenised before 600μL was transferred into a 96 well-plate (200μL into three separate wells). The optical density at 600nm (OD_{600}) and the fluorescence at 460nm following excitation at 400 nm of each of the cultures in the 96-well plate was measured using a BioTek Synergy 2 plate reader (BioTek, Vermont, U.S.A.). Pyoverdine, which fluoresces green, is the only culture component quantified using these excitation and emission parameters, with non-producers giving a reading of zero relative fluorescence units (RFU) (4). Each culture was measured three times (one reading of each of three wells per culture), and the average of three sterile media readings was used as a reference and subtracted from each culture value. The optical density measurements were used to estimate pyoverdine production per cell (*i.e.* per capita pyoverdine production) using: RFU / OD_{600}. The average of the three technical replicates was used in the analysis. We chose to quantify pyoverdine production at the population level (rather than for individual clones) as it represents the most relevant measure of the *Galleria mellonella* inoculum (see *Galleria mellonella virulence assay* below). We repeated this assay with the non-pyoverdine producing strain to confirm it produced negligible quantities of pyoverdine.

Galleria mellonella virulence assay

To quantify virulence of both the pyoverdine producing and non-producing strains, we used the insect infection model *Galleria mellonella* (44). Briefly, defrosted samples were diluted 10^5-fold in M9 salt buffer (3g KH_2PO_4, 6g Na_2HPO_4, 5g NaCl per litre) before 10μL was injected into each of 20 final instar larvae per sample using a 50μL Hamilton syringe (Hamilton, Nevada, USA). Larvae were then incubated at 37°C and mortality checked 18 hours

post-injection. Larvae were classed as dead when mechanical stimulation of the head caused
no response (45). M9-injected and non-injected controls were used to confirm mortality was
not due to injection trauma or background *G. mellonella* mortality; >10% control death was
the threshold for re-injecting (no occurrences). Pyoverdine producing and non-producing
strains were injected into different *G. mellonella* groups.

Quantifying Pseudomonas aeruginosa density
The density of *P. aeruginosa* in the common garden environment was quantified by plating
onto agar. To do this, samples were defrosted, serially diluted with M9 salt buffer and plated
onto KB agar. After 48 hours incubation at $28^{o}C$ colonies were counted and the number of
colony forming units (CFU) standardised to CFUs per mL. Note we only quantified the density
of the pyoverdine producing strain.

Statistical analysis
The effect of copper on per capita pyoverdine production (RFU / OD_{600}) and density (CFU/mL)
of *P. aeruginosa* populations was tested using linear mixed effects models (LMEM) with
copper and time as explanatory variables (both factors), as well as their 2-way interaction. To
determine how copper affected virulence we used a binomial generalised linear mixed model
(GLMM), with number of *G. mellonella* dead versus alive as binomial response variable, and
copper and time as explanatory variables, as well as their two-way interaction. To analyse the
combined effect of pyoverdine production and density on virulence, we used a similar
framework and included pyoverdine production, total CFU and time, along with a two-way
pyoverdine-time interaction, as explanatory variables. In all analyses, pyoverdine production
and population density were $\log_{10}$-transformed to improve normality, and random intercepts
were fitted for individual replicates to account for non-independency of observations over time.
For all analyses, we used the '*DHARMa*' package (46) to check residual behaviour,
after which the most parsimonious model was arrived at by sequentially deleting terms and
comparing model fits using χ -tests. These were followed by Tukey post hoc comparisons
adjusted for multiple comparisons using the Benjamini and Hochberg method. All analyses
were carried out in R version 3.3.3 (47), with the '*lme4*' package used for the LMEMs and
GLMMs (48).

Results

Copper selected for increased pyoverdine production

Here we tested the evolutionary consequences of copper stress on *Pseudomonas aeruginosa* by incubating it with copper for either two or twelve days, and then without copper for one day. All results shown here therefore display the evolutionary consequences of copper stress, as the phenotypic effects are removed by growing populations in copper-free common garden environment. Unless otherwise stated, data from the non-pyoverdine producing strain are only shown in the final section of the results. In line with previous findings (3), pyoverdine production evolved to be significantly greater as a function of copper concentration (copper main effect: χ^2=35.4, d.f.=2, p<0.001). This effect was consistent across the two- and twelve-day treatments; however, production was significantly greater after twelve days compared to two days (time main effect: χ^2=161, d.f.=1, p<0.001; Fig. 2). Pyoverdine production was significantly higher in the high copper treatment compared to both the control (two days: p<0.001; twelve days: p<0.001) and the low copper treatment (two days: p=0.032; twelve days: p=0.032). Likewise, the low copper treatment was always significantly higher than the control (two days: p=0.004; twelve days: p=0.004). The non-independence of observations between day two and day twelve samples accounted for only a small amount of the total variation (SD = 0.029) in this model.

Figure 2 Per capita pyoverdine production ($\log_{10}$-transformed standardised fluorescence units per OD_{600}) by *Pseudomonas aeruginosa* populations after growth in different concentrations of copper for (A) two days + one day in the absence of copper, or (B) twelve days + one day in the absence of copper. Circles show individual replicates (n = 6), with colour indicating total pyoverdine production (relative fluorescence units) and size showing optical density at 600nm. Asterisks indicate significant differences between groups (*** = 0.001, ** = 0.01, * = 0.05, NS = non-significant), with the left value comparing the control to the low copper treatment, the middle value comparing the control to the high copper treatment and the right value comparing the low and high copper treatments.

Virulence is higher in populations exposed to copper and this effect increases with time

To test whether copper stress caused differences in *P. aeruginosa* virulence, populations were assayed using the *Galleria mellonella* virulence model. Copper significantly increased virulence (copper main effect: χ^2=20.8, d.f.=2, p<0.001) and this effect increased with time (copper-time interaction: χ^2=6.90, d.f.=2, p=0.031; Fig. 3). After two days, virulence was significantly higher in the two copper treatments compared to the control (p<0.001 for both

comparisons) but these treatments did not differ themselves (p=0.34). After twelve days
virulence was again significantly higher in the two copper treatments than the control (p<0.001
for both comparisons), however the high copper treatment was found to be significantly more
virulent than the low copper treatment (p=0.027). We note these treatment effects were found
despite the random effect of repeatedly measuring the same population accounting for a large
amount of the variation (SD = 0.42).

Figure 3 The proportion of *Galleria mellonella* dead 18 hours after being injected with
Pseudomonas aeruginosa grown in the presence or absence of copper (A) two days + one day
in the absence of copper, or (B) twelve days + one day in the absence of copper. 20 *G.*
mellonella were injected per replicate (circles show individual replicates; n = 6 per unique
treatment combination). Asterisks indicate significant differences between groups (*** =
0.001, ** = 0.01, * = 0.05, NS = non-significant), with the left value comparing the control to
the low copper treatment, the middle value comparing the control to the high copper treatment
and the right value comparing the low and high copper treatments.

The effect of copper on population density differs over time

The effect of copper on population density differed as function of time (copper-time
interaction: χ^2=20.5, d.f.=2, p<0.001; Fig. 4). After two days, population densities did not

significantly differ between any copper treatment (p=>0.060 for all contrasts), whereas after
twelve days of growth density was significantly lower in the two copper treatments compared
to the control (high copper - control: p=0.013 and low copper - control: p=0.002) but did not
differ themselves (p=0.48). The effect of measuring the same populations at two time points
did not explain any of the overall variation in this model (SD = 0.0), but to account for non-
independency of observations it was retained in the model.

Figure 4 The density (log_{10} CFU mL^{-1}) of *Pseudomonas aeruginosa* populations incubated
with copper for (A) two days + one day in the absence of copper, or (B) twelve days + one day
in the absence of copper. Asterisks indicate significant differences between groups (*** =
0.001, ** = 0.01, * = 0.05, NS = non-significant), with the left value comparing the control to
the low copper treatment, the middle value comparing the control to the high copper treatment
and the right value comparing the low and high copper treatments.

Virulence is associated with increased pyoverdine production rather than pathogen load
Next, we tested whether increased virulence was associated with increased pyoverdine
production by replacing 'copper' as an explanatory variable in our GLMM with *per capita*
pyoverdine production and included density as a covariate in this model. Despite there being

substantial variation across populations in virulence (SD = 0.86), virulence increased as a function of pyoverdine production (pyoverdine main effect: χ^2=13.9, d.f.=1, p<0.001; Fig. 5), and this effect did not change with time (pyoverdine-time interaction: X^2=0.009, d.f.=1, p=0.93) although virulence was higher after twelve days (time main effect: X^2=5.40, d.f.=1, p=0.02). Population density had no significant effect on virulence (main effect of density in GLMM: X^2=1.09, d.f.=2, p=0.30).

Figure 5 The relationship between per capita pyoverdine production ($\log_{10}$-transformed relative fluorescence units per OD_{600}) and virulence of *P. aeruginosa* populations grown with copper for (A) two days + one day in the absence of copper, or (B) twelve days + one day in the absence of copper. Virulence was quantified using the *Galleria mellonella* infection model and expressed as the proportion of *G. mellonella* dead (out of 20) 18 hours after injection. Individual replicates are represented by circles (n = six per treatment); red points containing a △ represent the control (no copper) treatments, blue points containing a + the low copper treatment, and black points containing a × the high copper treatment. The line shows the best model fit, and the shaded area shows the 95% confidence interval.

The high copper treatment decreased the virulence of the non-pyoverdine producing strain

Finally, to confirm at least part of the change in virulence was due to pyoverdine production,

we tested whether a non-pyoverdine producing mutant strain of *P. aeruginosa* evolved under

the same copper regime exhibited similar changes in virulence. The fluorescence assay used to

detect pyoverdine suggested that the mutant strain still produced some pyoverdine (mean = 431

± 91.6 SD RFU), but this was on average 94.5% lower than populations of the wild-type

producer (mean = 7763 ± 2979 SD RFU), and therefore deemed negligible (data not shown).

Once again, we found a significant effect of copper on virulence (copper main effect: χ^2=66.6,

d.f.=2, p<0.001; Fig. 6) however, this time the effect was negative, and decreased with time

(copper-time interaction: χ^2=6.45, d.f.=2, p=0.040; Fig. 6). Despite a slight decrease in the

effect of copper over time, we found the same qualitative pattern in the virulence of the non-

pyoverdine producing strain across copper treatments after two and twelve days: virulence was

not significantly different between the low copper treatment and the control (two days:

p=0.087; twelve days: p=0.48), but was significantly lower in the high copper treatment

compared to both the low copper treatment and the control (p=<0.001 for all contrasts; Fig. 6).

As with the virulence model for the pyoverdine producing strain, the random effect of

repeatedly measuring the same population accounted for a large amount of the variation (SD =

0.44). Note that injections with pyoverdine producing and non-pyoverdine producing strains

were done in different batches of *G. mellonella*, and so statistical comparison between the

results from the wild-type and mutant infections is inappropriate.

Figure 6 The proportion of *Galleria mellonella* dead 18 hours after being injected with an isogenic non-pyoverdine producing strain of *P. aeruginosa* grown in the presence or absence of copper for (A) two days (+ one day in the absence of copper) or (B) twelve days (+ one day in the absence of copper). 20 *G. mellonella* were injected per replicate. Circles show individual replicates (n = 6), with size showing optical density at 600nm and colour indicating total pyoverdine production (standardised fluorescence units) to demonstrate the mutant did indeed not produce any pyoverdine. Asterisks indicate significant differences between groups (*** = 0.001, ** = 0.01, * = 0.05, NS = non-significant), with the left value comparing the control to the low copper treatment, the middle value comparing the control to the high copper treatment and the right value comparing the low and high copper treatments.

Discussion

Here, we experimentally tested whether copper causes the evolution of increased siderophore-mediated virulence in *Pseudomonas aeruginosa*. We found copper to select for increased per capita pyoverdine production, which was associated with greater levels of death in the *Galleria*

mellonella infection assay. Moreover, pyoverdine production was found to increase with time, such that populations exposed to copper for twelve days were more virulent than those exposed for two days. In addition to this overall increase in virulence with time, high copper caused significantly greater virulence than low copper in the twelve-day treatment but not in the two-day treatment. As a result, the twelve-day high copper treatment demonstrated the greatest virulence. Furthermore, we found that when using a non-pyoverdine producing mutant strain of *P. aeruginosa*, high copper reduced virulence, and low copper had little effect on virulence. Our results therefore demonstrate that copper stress increases pyoverdine production and consequently virulence, and that this effect increases with exposure time and copper concentration.

Finding copper-mediated increases in pyoverdine production is consistent with a role of pyoverdine in detoxification of copper in *P. aeruginosa* populations (49). Here, we show that a prolonged need for copper detoxification leads to the continued evolution of greater siderophore production. The resulting cost of these genotypic changes to a metabolically costly trait is a likely explanation for the reduced densities in the twelve-day copper treatments. We note that our finding of increased pyoverdine production with copper differs to that of previous work (8). Likely explanations for this are the fewer observed generations in our study (40 opposed to 200), culturing under static (versus shaken) conditions, which favours public goods production, and transfers occurring every two days compared to daily. In addition to copper, pyoverdine production has been shown to be up-regulated in response to Al^{3+}, Ga^{3+}, Mn^{2+} and Ni^{2+} (49). This suggests that these metals may also increase *P. aeruginosa* virulence, and may cause selection for increased production over longer exposure periods. Furthermore, it is likely that mixtures of metals, which are frequently encountered in polluted environments, might additively select for pyoverdine production and thereby virulence.

The positive association between pyoverdine production and virulence is consistent with previous findings, including in murine models (50). As well as directly aiding pathogen growth within a host by increasing iron uptake, pyoverdine production can increase virulence by causing the upregulation of additional virulence factors such as Exotoxin A and PrpL protease (51, 52). Finding per capita pyoverdine production to be a better predictor of virulence than population density in this system shows the cost of evolving greater production on density is inconsequential in a host. Furthermore, finding the virulence of the non-pyoverdine producing mutant to not be affected by low copper, and to be decreased by high copper, strongly suggests

that virulence has changed as a consequence of the evolutionary change in pyoverdine

production. However, we note that in addition to pyoverdine production, copper stress has been

shown to change the expression of over 300 genes in *P. aeruginosa* (53)*,* and it is plausible

these also have implications for virulence.

In addition to *P. aeruginosa*, toxic metals have also been shown to induce the production of

(non-pyoverdine) siderophores in other species (10), with siderophore production being shown

to increase proportionally with toxic metal pollution in natural microbial communities (9). As

the copper concentrations used in this study were chosen for their environmental relevance, we

suggest our findings could be relevant to natural communities inhabiting polluted

environments. We note that remediation techniques, principally lime addition, are used in some

metal-polluted areas to reduce their toxic effect (54), and this can lower community

siderophore production (55). Extending experiments such as those described here to the level

of natural communities could shed light on the consequences of metal pollution and metal

remediation on bacterial siderophore production and virulence.

In conclusion, we experimentally show that copper can increase the virulence of the pathogen

Pseudomonas aeruginosa by selecting for increased production of a metal-detoxifying

siderophore. We therefore demonstrate a direct link between toxic metal stress and virulence

in an opportunistic pathogen of significant clinical importance. Furthermore, we show that the

effect of metal exposure on virulence increases with exposure time and copper concentration.

This raises further concern for the effect of ever-increasing metal pollution on bacterial

pathogens, and highlights further work is needed to understand the role of metals in bacterial

virulence.

<u>Data accessibility</u>

All data and code will be made publically available upon acceptance. Currently, data can be

found on Mendeley Data at

https://data.mendeley.com/v1/datasets/yypr3yfr86/draft?preview=1, and the R code to

reproduce all analysis and create all data figures at

https://data.mendeley.com/v1/datasets/hbf9ht7d8h/draft?preview=1.

Author contributions

LL conceived the study, carried out experimental work, statistical analysis of the results and drafted the manuscript. EH helped with the analysis of the results. All authors helped with experimental design and critically revised the manuscript.

Acknowledgments

LL would like to thank the NERC FRESH GW4 award no. NE/R011524/1, EH the UKRI Future Leaders Fellowship award MR/V022482/1, AB the NERC award NE/V012347/1, and MV the NERC award NE/T008083/1.

References

1.	Cuív PÓ, Clarke P, O'Connell M. Identification and characterization of an iron-regulated gene, chtA, required for the utilization of the xenosiderophores aerobactin, rhizobactin 1021 and schizokinen by Pseudomonas aeruginosa. Microbiology. 2006;152(4):945-54.

2.	Rajkumar M, Ae N, Prasad MNV, Freitas H. Potential of siderophore-producing bacteria for improving heavy metal phytoextraction. Trends in biotechnology. 2010;28(3):142-9.

3.	Braud A, Geoffroy V, Hoegy F, Mislin G, Schalk I. The siderophores pyoverdine and pyochelin are involved in Pseudomonas aeruginosa resistance against metals: another biological function of these two siderophores. Environ Microbiol Rep. 2010;2:419-25.

4.	Kümmerli R, Jiricny N, Clarke L, West S, Griffin A. Phenotypic plasticity of a cooperative behaviour in bacteria. Journal of evolutionary biology. 2009;22(3):589-98.

5.	Winkelmann G. Ecology of siderophores with special reference to the fungi. Biometals. 2007;20(3-4):379.

6.	Cassat JE, Skaar EP. Iron in infection and immunity. Cell host & microbe. 2013;13(5):509-19.

7.	Palmer LD, Skaar EP. Transition metals and virulence in bacteria. Annual review of genetics. 2016;50:67-91.

8.	O'Brien S, Hodgson DJ, Buckling A. Social evolution of toxic metal bioremediation in Pseudomonas aeruginosa. Proceedings of the Royal Society B: Biological Sciences. 2014;281(1787):20140858.

9.	Hesse E, O'Brien S, Tromas N, Bayer F, Luján AM, van Veen EM, et al. Ecological selection of siderophore-producing microbial taxa in response to heavy metal contamination. Ecology letters. 2018;21(1):117-27.

10.	Saha R, Saha N, Donofrio RS, Bestervelt LL. Microbial siderophores: a mini review. Journal of basic microbiology. 2013;53(4):303-17.

11.	Hider RC, Kong X. Chemistry and biology of siderophores. Natural product reports. 2010;27(5):637-57.

12.   Griffin AS, West SA, Buckling A. Cooperation and competition in pathogenic bacteria.
Nature. 2004;430(7003):1024-7.
13.   Harrison F, McNally A, da Silva AC, Heeb S, Diggle SP. Optimised chronic infection
models demonstrate that siderophore 'cheating'in Pseudomonas aeruginosa is context
specific. The ISME journal. 2017;11(11):2492-509.
14.   Skaar EP. The battle for iron between bacterial pathogens and their vertebrate hosts.
PLoS pathogens. 2010;6(8):e1000949.
15.   Iatsenko I, Marra A, Boquete J-P, Peña J, Lemaitre B. Iron sequestration by
transferrin 1 mediates nutritional immunity in Drosophila melanogaster. Proceedings of the
National Academy of Sciences. 2020;117(13):7317-25.
16.   Golonka R, San Yeoh B, Vijay-Kumar M. The iron tug-of-war between bacterial
siderophores and innate immunity. Journal of innate immunity. 2019;11(3):249-62.
17.   Granato E, Harrison F, Kümmerli R, Ross-Gillespie A. Do Bacterial "Virulence Factors"
Always Increase Virulence. A Meta-Analysis of Pyoverdine Production in Pseudomonas
aeruginosa As a Test Case Front Microbiol. 2016;7:1952.
18.   Lam MM, Wyres KL, Judd LM, Wick RR, Jenney A, Brisse S, et al. Tracking key
virulence loci encoding aerobactin and salmochelin siderophore synthesis in Klebsiella
pneumoniae. Genome medicine. 2018;10(1):77.
19.   Saleem M, Shah M. Detection of Siderophore production in Uropathogenic
Escherichia. coli in patients with Type 2 Diabetes Mellitus. Int J Med Microbiol Trop Dis.
2017;3:176-7.
20.   Abergel RJ, Wilson MK, Arceneaux JE, Hoette TM, Strong RK, Byers BR, et al. Anthrax
pathogen evades the mammalian immune system through stealth siderophore production.
Proceedings of the National Academy of Sciences. 2006;103(49):18499-503.
21.   Perry WJ, Spraggins JM, Sheldon JR, Grunenwald CM, Heinrichs DE, Cassat JE, et al.
Staphylococcus aureus exhibits heterogeneous siderophore production within the
vertebrate host. Proceedings of the National Academy of Sciences. 2019;116(44):21980-2.
22.   Braud A, Geoffroy V, Hoegy F, Mislin GL, Schalk IJ. Presence of the siderophores
pyoverdine and pyochelin in the extracellular medium reduces toxic metal accumulation in
Pseudomonas aeruginosa and increases bacterial metal tolerance. Environmental
Microbiology Reports. 2010;2(3):419-25.
23.   Schalk IJ, Hannauer M, Braud A. New roles for bacterial siderophores in metal
transport and tolerance. Environmental microbiology. 2011;13(11):2844-54.
24.   Höfte M, Buysens S, Koedam N, Cornelis P. Zinc affects siderophore-mediated high
affinity iron uptake systems in the rhizosphere Pseudomonas aeruginosa 7NSK2. Biometals.
1993;6(2):85-91.
25.   Lhospice S, Gomez NO, Ouerdane L, Brutesco C, Ghssein G, Hajjar C, et al.
Pseudomonas aeruginosa zinc uptake in chelating environment is primarily mediated by the
metallophore pseudopaline. Scientific reports. 2017;7(1):1-10.
26.   Harrison F, Paul J, Massey RC, Buckling A. Interspecific competition and siderophore-
mediated cooperation in Pseudomonas aeruginosa. The ISME journal. 2008;2(1):49-55.
27.   Dumas Z, Kümmerli R. Cost of cooperation rules selection for cheats in bacterial
metapopulations. Journal of evolutionary biology. 2012;25(3):473-84.
28.   Kümmerli R, Jiricny N, Clarke LS, West SA, Griffin AS. Phenotypic plasticity of a
cooperative behaviour in bacteria. Journal of evolutionary biology. 2009;22(3):589-98.
29.   Meyer J-M, Neely A, Stintzi A, Georges C, Holder IA. Pyoverdin is essential for
virulence of Pseudomonas aeruginosa. Infection and immunity. 1996;64(2):518-23.

30.   Buckling A, Harrison F, Vos M, Brockhurst MA, Gardner A, West SA, et al.
Siderophore-mediated cooperation and virulence in Pseudomonas aeruginosa. FEMS
microbiology ecology. 2007;62(2):135-41.
31.   Rice LB. Federal funding for the study of antimicrobial resistance in nosocomial
pathogens: no ESKAPE. The University of Chicago Press; 2008.
32.   Rutherford V, Yom K, Ozer EA, Pura O, Hughes A, Murphy KR, et al. Environmental
reservoirs for exoS+ and exoU+ strains of Pseudomonas aeruginosa. Environmental
microbiology reports. 2018;10(4):485-92.
33.   Roy-Burman A, Savel RH, Racine S, Swanson BL, Revadigar NS, Fujimoto J, et al. Type
III protein secretion is associated with death in lower respiratory and systemic
Pseudomonas aeruginosa infections. The Journal of infectious diseases. 2001;183(12):1767-
74.
34.   Japoni A, Farshad S, Alborzi A. Pseudomonas aeruginosa: burn infection, treatment
and antibacterial resistance. Iranian Red Crescent Medical Journal. 2009;11(3):244-53.
35.   Fergie JE, Shema SJ, Lott L, Crawford R, Patrick CC. Pseudomonas aeruginosa
bacteremia in immunocompromised children: analysis of factors associated with a poor
outcome. Clinical infectious diseases. 1994;18(3):390-4.
36.   Rossi E, La Rosa R, Bartell JA, Marvig RL, Haagensen JA, Sommer LM, et al.
Pseudomonas aeruginosa adaptation and evolution in patients with cystic fibrosis. Nature
Reviews Microbiology. 2021;19(5):331-42.
37.   Harrison F, Buckling A. Cooperative production of siderophores by Pseudomonas
aeruginosa. Front Biosci. 2009;14:4113-26.
38.   Stilwell P, O'Brien S, Hesse E, Lowe C, Gardner A, Buckling A. Resource heterogeneity
and the evolution of public goods cooperation. Evolution letters. 2020;4(2):155-63.
39.   Dumas Z, Ross-Gillespie A, Kümmerli R. Switching between apparently redundant
iron-uptake mechanisms benefits bacteria in changeable environments. Proceedings of the
Royal Society B: Biological Sciences. 2013;280(1764):20131055.
40.   Brun L, Maillet J, Richarte J, Herrmann P, Remy J. Relationships between extractable
copper, soil properties and copper uptake by wild plants in vineyard soils. Environmental
pollution. 1998;102(2-3):151-61.
41.   UKWIR. The Chemical Investigations Programme Phase 2, 2015–2020 – Initial
Findings. 2018;1–4.
42.   O'Brien S, Luján AM, Paterson S, Cant MA, Buckling A. Adaptation to public goods
cheats in Pseudomonas aeruginosa. Proceedings of the Royal Society B: Biological Sciences.
2017;284(1859):20171089.
43.   Ghysels B. Thi Min Dieu B, Beatson SA, Pirnay JP, Ochsner UA, Vasil ML, Cornelis P.
2004 FpvB, an alternative type I ferripyoverdine receptor of Pseudomonas aeruginosa.
Microbiology.150:1671-80.
44.   Harrison F, Browning LE, Vos M, Buckling A. Cooperation and virulence in acute
Pseudomonas aeruginosainfections. BMC biology. 2006;4(1):21.
45.   Hernandez RJ, Hesse E, Dowling AJ, Coyle NM, Feil EJ, Gaze WH, et al. Using the wax
moth larva Galleria mellonella infection model to detect emerging bacterial pathogens.
PeerJ. 2019;7:e6150-e.
46.   Hartig F. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. R package version 042 2021.
47.   Team RC. R: A language and environment for statistical computing. R Foundation for
Statistical Computing.

48.   Douglas Bates MM, Ben Bolker, Steve Walker. Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software. 2015;67(1):1-48.
49.   Braud A, Hoegy F, Jezequel K, Lebeau T, Schalk IJ. New insights into the metal
specificity of the Pseudomonas aeruginosa pyoverdine–iron uptake pathway. Environmental
microbiology. 2009;11(5):1079-91.
50.   Kang D, Revtovich AV, Chen Q, Shah KN, Cannon CL, Kirienko NV. Pyoverdine-
dependent virulence of Pseudomonas aeruginosa isolates from cystic fibrosis patients.
Frontiers in microbiology. 2019;10:2048.
51.   Lamont IL, Beare PA, Ochsner U, Vasil AI, Vasil ML. Siderophore-mediated signaling
regulates virulence factor production in Pseudomonas aeruginosa. Proceedings of the
National Academy of Sciences. 2002;99(10):7072-7.
52.   Lopez-Medina E, Fan D, Coughlin LA, Ho EX, Lamont IL, Reimmann C, et al. Candida
albicans inhibits Pseudomonas aeruginosa virulence through suppression of pyochelin and
pyoverdine biosynthesis. PLoS Pathog. 2015;11(8):e1005129.
53.   Teitzel GM, Geddie A, Susan K, Kirisits MJ, Whiteley M, Parsek MR. Survival and
growth in the presence of elevated copper: transcriptional profiling of copper-stressed
Pseudomonas aeruginosa. Journal of bacteriology. 2006;188(20):7242.
54.   Palansooriya KN, Shaheen SM, Chen SS, Tsang DCW, Hashimoto Y, Hou D, et al. Soil
amendments for immobilization of potentially toxic elements in contaminated soils: A
critical review. Environment international. 2020;134:105046.
55.   Hesse E, Padfield D, Bayer F, Van Veen EM, Bryan CG, Buckling A. Anthropogenic
remediation of heavy metals selects against natural microbial remediation. Proceedings of
the Royal Society B. 2019;286(1905):20190804.
546