Feasibility of Administering Human Pancreatic Cancer Chemotherapy in

Spontaneous Pancreatic Cancer Mouse Model

Abagail M. Delahoussaye, B.S.[1*], Joseph Abi Jaoude, M.D.[1*], Tara N. Fujimoto, B.S.[1], Jessica

Molkentine, B.S. [1], Carolina J. Garcia Garcia, B.S.[1], Jason P. Gay, B.S., B.A.[1], Ningping Feng,

Ph. D. [1], Joe Marszalek, Ph.D. [1], Cullen M. Taniguchi, M.D., Ph.D.[1§]

[*]Contributed equally as first authors

Affiliations: [1]The University of Texas MD Anderson Cancer Center, Houston, TX, USA.

[§]Corresponding Author:

Cullen M. Taniguchi, M.D., Ph.D., 1515 Holcombe Blvd Unit 1050, Houston, TX 77030.

Phone: 713-745-5269. Email: ctaniguchi@mdanderson.org

Abstract:

Background: Both modified FOLFIRINOX (mFFX) and gemcitabine/*nab*-paclitaxel

chemotherapy regimens have been shown to improve clinical outcomes in patients with

pancreatic cancer, and are often used interchangeably as the standard of care. Preclinical

studies often do not use these regimens, since administering these multiagent approaches

can be difficult. In this study, we assessed the feasibility of administering these two

chemotherapy regimens in spontaneous pancreatic tumors using KPC mice with the ultimate

goal of advancing preclinical studies.

Methods: KPC mice were created by breeding $Kras^{LSL-G12D/+}$ to $Trp53^{fl/fl};Ptf1\alpha^{Cre/+}$, resulting

in $Kras^{LSL-G12D/+};p53^{fl/+};Ptf1\alpha^{Cre/+}$ mice. At 14 weeks of age, mice were palpated for

spontaneous tumor growth that was verified using ultrasounds. Mice with tumors under 15

mm in diameter were used. The mice were assigned to one of seven treatment regimens: 1

cycle of mFOLFIRINOX (FFX X1), 2 cycles of mFOLFIRINOX (FFX X2), 1 cycle of

mFOLFIRINOX with 40Gy SBRT (FFX SBRT), 1 cycle of gemcitabine/*nab*-paclitaxel

(GEM/AB X1), 2 cycles of gemcitabine/*nab*-paclitaxel (GEM/AB X2), 2 cycles of

gemcitabine/*nab*-paclitaxel with 40Gy SBRT (GEM/AB SBRT), or no chemotherapy treatment

(control).

Results: In total, 92 mice were included. The median OS in the FFX X2 group was slightly

longer that the median OS in the FFX X1 group (15 days vs 11 days, P=0.003). Mice in the

GEM/AB X2 group had longer OS when compared to mice in the GEM/AB X1 group (33.5 vs

13 days, P=0.001). Mice treated with chemotherapy survived longer than untreated control

animals (median OS: 6.5 days, P <0.001). Moreover, in mice treated with chemotherapy,

mice that received 2 cycles of GEM/AB X2 had the longest survival, while the FFX X1 group

had the poorest OS (P <0.001). Lastly, chemotherapy followed by consolidative SBRT

trended towards increased local control and survival.

Conclusions: We demonstrate the utility and feasibility of clinicall relevant mFOLFIRINOX

and gemcitabine/*nab*-paclitaxel in preclinical models of pancreatic cancer.

Keywords:

KPC mice,

mFOLFIRINOX,

gemcitabine,

nab-paclitaxel,

pancreatic cancer,

SBRT

Background

Background

Pancreatic cancer is one of the leading causes of cancer-related death in the world [1,

2]. Owing to the limited role of screening in pancreatic cancer, tumors are often discovered at

advanced stages, which contributes to poor outcomes and severe morbidity and mortality.

While surgery is the main curative option in pancreatic cancer, a large proportion of patients

with pancreatic cancer are initially diagnosed with unresectable disease that would not

benefit from local therapy [1, 3]. Moreover, pancreatic cancer is known to metastasize early

in the disease process [4]. Therefore, chemotherapy remains a major part of treatment for all

stages of pancreatic cancer.

Many chemotherapy regimens have been suggested for the treatment of pancreatic

cancer. Previously, gemcitabine and 5-fluorouracil were used as monotherapy, however

recurrence rates remained high [5, 6]. As such, modern systemic treatment consists \of

powerful combinations of chemotherapy. The two most common regimens used in the

modern treatment of pancreatic cancer are modified FOLFIRINOX (mFFX: oxaliplatin,

irinotecan, leucovorin, fluorouracil) and gemcitabine/*nab*-paclitaxel [7–10]. Many studies and

trials analyzing the role of those regimens showed superior clinical results compared to

conventional monotherapy treatments, and established the role of mFOLFIRINOX and

gemcitabine/*nab*-paclitaxel as standard of care in pancreatic cancer.

While those regimens have been thoroughly studied in humans, limited data on the

use of mFOLFIRINOX and gemcitabine/*nab*-paclitaxel are present in pre-clinical studies. In

particular, the clinical utility and toxicity of those regimens is still not clear in spontaneous

pancreatic tumors in KPC mice. In order to properly mimic human clinical treatment, pre-

clinical models need to implement similar chemotherapy regimens. In this study, we aim to

assess the feasibility of administering mFOLFIRINOX and gemcitabine/*nab*-paclitaxel in

spontaneous pancreatic tumors using a KPC mouse model, as well as their toxicity profiles.

Methods

KPC mice and tumor diagnosis

All mouse work was approved by the Institutional Animal Care and Use

Committee of The University of Texas MD Anderson Cancer Center (IACUC

#00001252-RN02). Both female and male mice were used in this study. Mice were

maintained on a 12-hour light/dark cycle and were provided with sterilized water and

standard rodent chow (Prolab Isopro RMH 3000 irradiated feed). Experiments were

carried out during the light cycle.

KPC mice were created by breeding $Kras^{LSL-G12D/+}$ to $Trp53^{fl/fl};Ptf1\alpha^{Cre/+}$, resulting

in a $Kras^{LSL-G12D/+};Trp53^{fl/+};Ptf1\alpha^{Cre}$ (KPC) mice, which is a validated model of locally

advanced pancreatic cancer [11]. Beginning at 14 weeks of age, mice were screened

weekly by palpation under anesthesia for tumor growth, as previously described [12].

Suspicious intrabdominal masses, were then subjected to small animal ultrasound

imaging (FujiFilm Vevo2100), with a 30MHz transducer, acquired in B-MODE to

estimate volumes from the long and short axes [12, 13].

Chemotherapy

We used pharmaceutical grade mFOLFIRINOX that was provided by The

University of Texas MD Anderson Cancer Center pharmacy. Oxaliplatin was

administered intravenously via the tail vein at 10 mg/kg in a volume of 0.01 mL/g. 5-FU

(Fresenius Kabi, 50 mg/mL) and Irinotecan (TEVA, 20 mg/mL) were combined and

diluted with 0.9% Sodium Chloride (normal saline) and given via intraperitoneal injection

at 50 mg/kg. Both injections account for one cycle delivered in the same day. For mice

that received two cycles of mFOLFIRINOX, the second cycle was given 2 weeks after

completion of the first cycle (Supplementary Figure 1).

For mice treated with gemcitabine/nab-paclitaxel, we used pharmaceutical grade

gemcitabine (Gemzar®, NDC 00002-7501) and *nab*-paclitaxel (Abraxane®, Celgene)

generously provided by The University of Texas MD Anderson Cancer Center Institute

for Applied Cancer Science (IACS). Gemcitabine was diluted with saline to 10 mg/mL

and administered with intraperitoneal injections at 100 mg/kg in a volume of 0.01 mL/g,

and given twice weekly. *Nab*-paclitaxel (Abraxane®, Celgene) was diluted with saline to

30 mg/mL and administered intravenously at a dose of 300 mg/kg (30mgPTX/kg) in a

volume of 0.01 mL/g, and given only once weekly one hour after one of the gemcitabine

doses [14] (Supplementary Figure 1).

Chemotherapy dose finding and efficacy

Mice with detectable tumors were randomly assigned to receive modified

FOLFIRINOX (mFFX) or gemcitabine/*nab*-paclitaxel (GEM/AB), and mice received

therapy within 48 hours of assignment.   Both mFFX and GEM/AB arms were initially

assigned to received one cycle of chemotherapy (FFXx1 and GEM/AB x1). If no

excessive toxicity was observed, then the next batch of mice would receive 2 cycles of

chemotherapy (FFXx1 and GEM/AB x1). No further treatment was given after the last

assigned dose of chemotherapy.   Body weights were measured daily, and tumor

growth monitored twice weekly with ultrasound. Endpoints were reached when mice met

predetermined euthanasia endpoints, including tumors measuring greater than 15 mm

in diameter, lethargy, weight loss exceeding 20% from baseline, or appearing moribund.

At euthanasia, necropsies were conducted to look for macrometastases and determine

a potential cause of death, as we have previously described [13].

Radiation Therapy

For mice that received local treatment with radiation therapy, stereotactic body radiation

therapy (SBRT) was delivered using an XRAD 225Cx irradiator with isoflurane

anesthesia manifold and image guidance. The tumors were palpated and their locations

marked with metallic beads. Then, radiation fields were aligned for each mouse using

cone beam CT image guidance. The metallic beads were removed before radiation.

Beam arrangement was anteroposterior/posteroanterior using a 10 mm diameter

collimator, and mice received 5 fractions of 8Gy with 24 hours between fractions, for a

total of 40Gy [13]. mFOLFIRINOX SBRT mice were irradiated two weeks following one

dose of chemotherapy. Gemcitabine/*nab*-paclitaxel SBRT mice were given two doses of

chemotherapy and then received SBRT two weeks after the final dose. Mice were

allowed to feed and drink water *ad libitum* throughout the duration of treatments.

Additional supportive measures, such as moistened food pellets on the ground, were

given if the mouse appeared dehydrated. Mice were also observed for radiation toxicity

daily. The main symptoms assessed were the development of ruffled fur, persistent

diarrhea, and/or drastic weight loss of 20% or more.

Statistical Analysis

Descriptive statistics were used to assess the baseline characteristics of mice

and tumors in each group of mice. Continuous variables were presented as median and

corresponding interquartile range (IQR). Categorical variables were presented as

frequencies and percentages. We further assessed the average weight of mice in each

group every day post-treatment. Furthermore, survival curves for Overall Survival (OS)

were generated using the Kaplan-Meier method. OS was calculated from time of

diagnosis until death of mice. Local tumor recurrence, distant metastasis, and cause of

death were noted for all mice. Statistical significance was set *a priori* at *P* <0.05. All

statistical analyses were performed using IBM SPSS version 26. Kaplan-Meier curves

were generated using Prism Version 8.

Results

Baseline Mice and Tumor Characteristics

In total, 92 mice were included and analyzed in our study. In particular, 54 mice were

treated with chemotherapy only: 6 mice were treated with one cycle of mFOLFIRINOX (FFX

X1), 30 mice with two cycles of mFOLFIRINOX (FFX X2), 6 mice with one cycle of

gemcitabine/*nab*-paclitaxel (GEM/AB X1), 6 mice with two cycles of gemcitabine/*nab*-

paclitaxel (GEM/AB X2), and 28 mice did not receive any chemotherapy (control).

<Insert Table 1>

Table 1 presents the baseline characteristics of mice and tumors in mice treated with

chemotherapy only. Mice in the FFX X2 group had the oldest age at diagnosis (24.8 weeks,

IQR [21.7-26.9]), and those in the GEM/AB X2 had the youngest age at diagnosis (14.3

weeks, IQR [9.1-23.5]). A similar proportion of males and females were present in the FFX

groups, and a few more males than females were present in the GEM/AB and control groups.

Baseline weights were similar between all cohorts (between median: 23.6g, IQR 20.5-28.1

and median: 25.3g, IQR 22.0-28.4).

The baseline tumor diameter was largest in the FFX X1 group (median: 9.3 mmm IQR

7.1-10.2), smallest in the GEM/AB X2 group (median: 4.8 mm, IQR 4.0-6.9), and similar in

the other 3 groups (between median: 6.7 mm, IQR 6.2-7.4 and median: 7.1 mm, IQR 5.5-

8.0). Most tumors were located in the body of the pancreas for the FFX X1 group (3/6,

50.0%), the tail of the pancreas for the FFX X2 group (13/30, 43.3%), the head of the

pancreas for the GEM/AB X1 group (4/6, 66.7%), the tail of the pancreas for the GEM/AB X2

group (3/6, 50.0%), and the head of the pancreas for the control group (13/28, 46.4%) (Table

1).

Comparative analysis between mFOLFIRINOX and gemcitabine/nab-paclitaxel reveals

similar survival times

We ultimately wanted to treat KPC animals with at least two cycles of chemotherapy,

since that might better approximate the multicycle regimens given to patients [7]. Thirty-six

mice were treated with mFOLFIRINOX (6 mice with one cycle and 30 mice with two cycles).

The median OS in the FFX X2 group was slightly longer that the median OS in the FFX X1

group (15 days vs 11 days, log-rank *P*=0.003) (Fig. 1A). Twelve mice were treated with

gemcitabine/*nab*-paclitaxel (6 mice with one cycle and 6 mice with two cycles). Mice in the

GEM/AB X2 group had longer OS when compared to mice in the GEM/AB X1 group (median

OS, GEM/AB X2: 33.5 days vs GEM/AB X1: 13 days, log-rank *P*=0.001) (Fig. 1B).

We compared OS between the mFFX and the GEM/AB treated mice. In mice that

received one cycle of chemotherapy, the GEM/AB group had a slightly longer median OS

compared to the FFX group, but this difference was not statistically significant (GEM/AB X1:

13 days, FFX X1: 11 days, log-rank P=0.104) (Fig. 2A). In mice that received two cycles of

chemotherapy, the GEM/AB mice had better OS than the FFX mice (median OS, GEM/AB

X2: 33.5 days, FFX X2: 15 days, log-rank P=0.002) (Fig. 2B). All mice treated with

chemotherapy showed a better median OS compared to the control group (median OS: 10.0

days, log-rank P <0.001) (Fig. 2C). Overall, in mice treated with chemotherapy, the GEM/AB

X2 group showed the best OS, and the FFX X1 group had the poorest results (log-rank P

<0.001) (Fig. 2C).

Our KPC animals are a validated model of autochthonous and localized pancreatic

cancer. Thus, not surprisingly, local progression was the main cause of death in both FFX

groups (FFX X1: 5, 83.3%; FFX X2: 20, 66.7%, Table 2), compared to distant metastasis

(FFX X1: 1, 16.7%; FFX X2: 11, 36.7%, Table 2). A similar pattern emerged in GEM/AB

treated mice, where local progression of disease led to death more commonly than distant

metastases (Table 2). Particularly, in the GEM/AB X1 cohort, 4 (66.7%) mice died from local

recurrence, and 1 (16.7%) mouse died from distant metastasis. Moreover, in the GEM/AB X2

cohort 4 (66.7%) mice died from local recurrence, and 2 (33.3%) died from distant

metastasis.   In fact, ascites was observed in 7 (7/36, 14.4%) of mFFX and 1 (1/12, 8.3%) of

GEM/AB mice.

<Insert Table 2>

We noted no major differences in acute toxicity between the treatment regimens, as

assessed by weight changes (Fig. 3). The average weight of mice in the GEM/AB X1 and

FFX X1 decreased over time, while that of the GEM/AB X2 and FFX X2 increased.

Chemoradiation may be superior to chemotherapy alone in KPC mouse models

Radiation therapy is a commonly prescribed to consolidate treatment of the primary

mass in pancreatic cancer, especially if surgery cannot be offered [15].   Stereotactic body

radiation therapy (SBRT) has emerged as a convenient and powerful methods to achieve

local control [16]. We previously demonstrated that SBRT is feasible in murine models [17].

To this end, we treated cohorts of KPC mice that first received mFOLFIRINOX or

gemcitabine/*nab*-paclitaxel with a clinically relevant dose of SBRT (8Gy x 5 fx).   Sixteen

mice were treated with chemoradiation: 4 mice (25%) were treated with one cycle of

mFOLFIRINOX and SBRT, and 12 mice (75%) were treated with two cycles of

gemcitabine/*nab*-paclitaxel and SBRT.

<Insert Table 3>

Table 3 presents the baseline characteristics of mice treated with consolidative SBRT

after induction chemotherapy. The median OS in the FFX + SBRT group was 34 days, and

the median OS in the GEM/AB + SBRT group was 41 days (log-rank *P*=0.214) (Fig. 4). The

addition of SBRT to both mFFX and GEM/AB regimens seems to show a trend towards

improved survival compared to mice treated with chemotherapy only (Fig. 2). In total, 4 (25%)

mice treated with chemoradiation died from local recurrence: 3 in the FFX group (3/4, 75%),

and 1 in the GEM/AB group (1/12, 8.3%) (Table 4). Only one mouse (8.3%) in the GEM/AB

group died from radiation toxicity.

<Insert Table 4>

Discussion

The use of chemotherapy is common in patients with pancreatic cancer since most

tumors present with advanced or metastatic disease where local therapy is of limited clinical

benefit. Two common chemotherapy regimens in the modern treatment of pancreatic cancer

are mFOLFIRINOX and gemcitabine/*nab*-paclitaxel [8, 9, 18, 19]. However, the use of those

regimens is not routinely performed in pre-clinical models that often inform targets for clinical

trials. In our study, we show that treatment with mFOLFIRINOX and gemcitabine/*nab*-

paclitaxel is feasible and prolong survival. Thus, the routine use of clinical relevant

chemotherapy should be considered in studies involving spontaneous pancreatic tumors in

KPC mice.

When analyzing mice that were treated with chemotherapy only, we found that mice

treated with two cycles of gemcitabine/*nab*-paclitaxel had the best survival outcomes (median

OS: 33.5 days, Fig. 2C). A major component of this regimen is the nanoparticle-albumin-bound

paclitaxel which contains albumin, a human derived protein [20]. As such, using *nab*/paclitaxel

in KPC mice could be postulated to lead to significant toxicity due to cross reactivity with the

human component of the drug with repeated doses. However, we show that gemcitabine/*nab*-

paclitaxel was very well tolerated by KPC mice even with repeated dosing, and even led to

improved OS compared to mFOLFIRINOX (Fig. 2A, 2B). This similar effect was observed in

prior studies, but it is unclear whether prolonged administration of GEM/AB could be feasible.

When assessing weight change, mice treated with gemcitabine/*nab*-paclitaxel had

similar weight change trends to those treated with mFOLFIRINOX (Fig. 3). Necropsy analysis

revealed that mice treated with gemcitabine/*nab*-paclitaxel died because of disease

progression, without significant treatment toxicity. We note that mice treated with two cycles

of gemcitabine/*nab*-paclitaxel gained weight between diagnosis and death, which we attribute

to abdominal ascites and tumor progression. Additionally, In that regard, it seems that

gemcitabine/*nab*-paclitaxel could be safely considered in KPC mice with spontaneous

pancreatic tumors.

Despite the improved outcomes in mice treated with gemcitabine/*nab*-paclitaxel, the

mice cohorts treated with mFOLFIRINOX still showed reasonable OS (median OS, FFX X1:

11 days, FFX X2: 15 days) (Fig. 1A). Moreover, mice treated with one or two cycles of

mFOLFIRINOX had longer OS compared to mice in the control cohort (Fig. 2C). As such,

mFOLFIRINOX can also be safely considered in KPC mice. This feasibility study was not

designed to assess potential futility of mFFX compared to gemcitabine/abraxane, but we do

note that the there was a survival trend in the latter combination with a more favorable toxicity

profile, as has been reported in a recent Phase II study [21].

Lastly, we analyzed the use of chemoradiation with SBRT in KPC mice, and we show

that the addition of local treatment with radiation therapy to chemotherapy further improved

OS (Fig. 2C). This goes in line with pancreatic cancer treatment in humans, where local therapy

with surgery and/or radiation therapy has been shown to improve survival and recurrence rates

[22, 23]. Furthermore, KPC mice tolerated SBRT treatment well, with only one mouse dying of

radiation-induced toxicity (Table 4). SBRT treatment has become increasingly common in

clinical practice, and as such, incorporating chemoradiation with SBRT in pre-clinical models

would help to best parallel clinical management. In that regard, our data show that

mFOLFIRINOX, *nab*/paclitaxel, and SBRT can be safely used and combined in KPC mice.

Conclusions

Our data show that the use of mFOLFIRINOX and gemcitabine/*nab*-paclitaxel is feasible in

spontaneous pancreatic tumors in KPC mouse models. Mice receiving two cycles of

chemotherapy had improved survival, and did not show signs of severe toxicity. We further

show that chemoradiation using mFOLFIRINOX or gemcitabine/*nab*-paclitaxel chemotherapy

regimens with SBRT is also feasible, and was also well tolerated in KPC mice. Our results

should encourage future pre-clinical studies to incorporate modern chemotherapy

combinations in order to best mimic the current clinical standard of care.

List of Abbreviations:

FFX: FOLFIRINOX

GEM/AB: gemcitabine/*nab*-paclitaxel

IQR: interquartile range

mFFX: modified FOLFIRINOX

OS: overall survival

SAIF: Small Animal Imaging Facility

SBRT: stereotactic body radiation therapy

Declarations:

Ethical Approval: The University of Texas MD Anderson Cancer Center Institutional Review

Board approved all protocols in this study.

Consent for Publication: Not applicable.

Availability of Data and Materials: Research data are stored in an institutional repository and

will be shared upon reasonable request to the corresponding author.

Competing interests: Dr. CMT is on the medical advisory board of Accuray and is a paid

consultant for Xerient Pharma and Phebra Pty, Ltd. All other authors report no competing

interests.

Funding: CMT is supported by funding from NIH under award R01CA227517-01A1,

Cancer Prevention & Research Institute of Texas (CPRIT) grant RR140012, V

Foundation (V2015-22), the Reaumond Family Foundation, the Childress Family

Foundation, the Mark Foundation, and the McNair Foundation. CJGG was supported by

funding from NIH under awards F31DK121384 and U54CA096300/297. This work was

also supported by the NIH/NCI under award number P30CA016672 for use of the Small

Animal Imaging Facility.

Author's Contributions: AMD, JAJ, and CMT were responsible for the concept and study

study design. AMD performed all the mouse experiments and statistical analysis performed

by JAJ. All authors contributed substantially to the writing of this manuscript, through

performing literature review, data analysis, data interpretation, manuscript drafting, and

providing significant comments and/or edits to the final manuscript.

Acknowledgments: We would like to thank the University of Texas MD Anderson

Cancer Center Institute for Applied Cancer Science (IACS) for generously providing the

chemotherapy used in our study. This research is supported in part by the National

Institutes of Health through MD Anderson's Cancer Center Support Grant CA016672.

This work was also supported by the NIH/NCI under award number P30CA016672 for

use of the Small Animal Imaging Facility.

References

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer Statistics, 2021. CA: A Cancer Journal for Clinicians. 2021;71:7–33.

2. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting Cancer Incidence and Deaths to 2030: The Unexpected Burden of Thyroid, Liver, and Pancreas Cancers in the United States. Cancer Res. 2014;74:2913–21.

3. Chakraborty S, Singh S. Surgical resection improves survival in pancreatic cancer patients without vascular invasion- a population based study. Ann Gastroenterol. 2013;26:346–52.

4. Bayne LJ, Beatty GL, Jhala N, Clark CE, Rhim AD, Stanger BZ, et al. Tumor-derived granulocyte-macrophage colony-stimulating factor regulates myeloid inflammation and T cell immunity in pancreatic cancer. Cancer Cell. 2012;21:822–35.

5. Oettle H, Post S, Neuhaus P, Gellert K, Langrehr J, Ridwelski K, et al. Adjuvant chemotherapy with gemcitabine vs observation in patients undergoing curative-intent resection of pancreatic cancer: a randomized controlled trial. JAMA. 2007;297:267–77.

6. Neoptolemos JP, Stocken DD, Friess H, Bassi C, Dunn JA, Hickey H, et al. A randomized trial of chemoradiotherapy and chemotherapy after resection of pancreatic cancer. N Engl J Med. 2004;350:1200–10.

7. Conroy T, Hammel P, Hebbar M, Ben Abdelghani M, Wei AC, Raoul J-L, et al. FOLFIRINOX or Gemcitabine as Adjuvant Therapy for Pancreatic Cancer. New England Journal of Medicine. 2018;379:2395–406.

8. Conroy T, Desseigne F, Ychou M, Bouché O, Guimbaud R, Bécouarn Y, et al. FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. N Engl J Med. 2011;364:1817–25.

9. Chun JW, Lee SH, Kim JS, Park N, Huh G, Cho IR, et al. Comparison between FOLFIRINOX and gemcitabine plus nab-paclitaxel including sequential treatment for metastatic pancreatic cancer: a propensity score matching approach. BMC Cancer. 2021;21:537.

10. Riedl JM, Posch F, Horvath L, Gantschnigg A, Renneberg F, Schwarzenbacher E, et al. Gemcitabine/nab-Paclitaxel versus FOLFIRINOX for palliative first-line treatment of advanced pancreatic cancer: A propensity score analysis. European Journal of Cancer. 2021;151:3–13.

11. Aguirre AJ, Bardeesy N, Sinha M, Lopez L, Tuveson DA, Horner J, et al. Activated Kras and Ink4a/Arf deficiency cooperate to produce metastatic pancreatic ductal adenocarcinoma. Genes Dev. 2003;17:3112–26.

12. Yang Y, Armour M, Wang KK-H, Gandhi N, Iordachita I, Siewerdsen J, et al.
Evaluation of a Cone Beam Computed Tomography Geometry for Image Guided Small
Animal Irradiation. Phys Med Biol. 2015;60:5163–77.

13. Fujimoto TN, Colbert LE, Huang Y, Molkentine JM, Deorukhkar A, Baseler L, et al.
Selective EGLN Inhibition Enables Ablative Radiotherapy and Improves Survival in
Unresectable Pancreatic Cancer. Cancer Res. 2019;79:2327–38.

14. Frese KK, Neesse A, Cook N, Bapiro TE, Lolkema MP, Jodrell DI, et al. nab-
paclitaxel potentiates gemcitabine activity by reducing cytidine deaminase levels in a
mouse model of pancreatic cancer. Cancer Discov. 2012;2:260–9.

15. Abi Jaoude J, Kouzy R, Nguyen ND, Lin D, Noticewala SS, Ludmir EB, et al.
Radiation therapy for patients with locally advanced pancreatic cancer: Evolving
techniques and treatment strategies. Current Problems in Cancer. 2020;:100607.

16. Abi Jaoude J, Thunshelle CP, Kouzy R, Nguyen ND, Lin D, Prakash L, et al.
Stereotactic Versus Conventional Radiation Therapy for Patients with Pancreatic
Cancer in the Modern Era. Advances in Radiation Oncology. 2021;0.
doi:10.1016/j.adro.2021.100763.

17. Molkentine JM, Fujimoto TN, Horvath TD, Grossberg AJ, Garcia CJG, Deorukhkar
A, et al. Enteral Activation of WR-2721 Mediates Radioprotection and Improved Survival
from Lethal Fractionated Radiation. Sci Rep. 2019;9:1949.

18. Goldstein D, El-Maraghi RH, Hammel P, Heinemann V, Kunzmann V, Sastre J, et
al. nab-Paclitaxel plus gemcitabine for metastatic pancreatic cancer: long-term survival
from a phase III trial. J Natl Cancer Inst. 2015;107:dju413.

19. Blomstrand H, Scheibling U, Bratthäll C, Green H, Elander NO. Real world evidence
on gemcitabine and nab-paclitaxel combination chemotherapy in advanced pancreatic
cancer. BMC Cancer. 2019;19:40.

20. Giordano G, Pancione M, Olivieri N, Parcesepe P, Velocci M, Di Raimo T, et al.
Nano albumin bound-paclitaxel in pancreatic cancer: Current evidences and future
directions. World J Gastroenterol. 2017;23:5875–86.

21. Ozaka M, Ueno M, Ishii H, Mizusawa J, Katayama H, Kataoka T, et al. Randomized
phase II study of modified FOLFIRINOX versus gemcitabine plus nab-paclitaxel
combination therapy for locally advanced pancreatic cancer (JCOG1407). JCO. 2021;39
15_suppl:4017–4017.

22. Katz MHG, Ou F-S, Herman JM, Ahmad SA, Wolpin B, Marsh R, et al. Alliance for
clinical trials in oncology (ALLIANCE) trial A021501: preoperative extended
chemotherapy vs. chemotherapy plus hypofractionated radiation therapy for borderline
resectable adenocarcinoma of the head of the pancreas. BMC Cancer. 2017;17:505.

23. Koong AC, Le QT, Ho A, Fong B, Fisher G, Cho C, et al. Phase I study of
stereotactic radiosurgery in patients with locally advanced pancreatic cancer. Int J
Radiat Oncol Biol Phys. 2004;58:1017–21.

Figures

Figure 1. Kaplan-Meier Curve for Overall Survival in Mice Treated with One or Two

Cycles of mFOLFIRINOX (FFX) (A), and in Mice Treated with One or Two Cycles of

gemcitabine/*nab*-paclitaxel (GEM/AB) (B).

Figure 2. Kaplan-Meier Curve for Overall Survival in Mice Treated with One Cycle of

mFOLFIRINOX (FFX) or gemcitabine/*nab*-paclitaxel (GEM/AB) (A), and in Mice Treated

with Two Cycles of mFOLFIRINOX (FFX) or gemcitabine/*nab*-paclitaxel (GEM/AB) (B),

and Among All Mice (C).

Figure 3. Mice Weight Change After Chemotherapy Treatment.

Survival SBRT

P=0.214
FFX+SBRT
GEM/AB + SBRT

422   **Figure 4.** Kaplan-Meier Curve for Overall Survival in Mice Treated Stereotactic Body

423   Radiation Therapy (SBRT) and mFOLFIRINOX (FFX) or gemcitabine/*nab*-paclitaxel

424   (GEM/AB).