Parallel evolution of urban-rural clines

Bradley J. Cosentino[1][*] and James P. Gibbs[2]

[1]Department of Biology, Hobart and William Smith Colleges, 300 Pulteney Street, Geneva, NY

14456, USA

[2]Department of Environmental Biology, State University of New York College of Environmental

Science and Forestry, 1 Forestry Drive, Syracuse, NY 13210, USA

*Corresponding author. E-mail: cosentino@hws.edu

Abstract

Urbanization is the dominant trend of global land use change. The replicated nature of

environmental change associated with urbanization should drive parallel evolution, yet insight

into the repeatability of evolutionary processes in urban areas has been limited by a lack of

multi-city studies. Here we leverage community science data on coat color in >60,000 eastern

gray squirrels (*Sciurus carolinensis*) across 43 North American cities to test for parallel clines in

melanism, a genetically based trait associated with thermoregulation and crypsis in gray

squirrels. We show the prevalence of melanism in these mammals was positively associated with

urbanization. Urban-rural clines in melanism were strongest in the largest cities with extensive

forest cover and weakest or absent in cities with warm winter temperature, where thermal

selection likely limits the prevalence of melanism. Our results demonstrate that novel traits can

evolve in a highly repeatable manner among urban areas, modified by factors intrinsic to

individual cities, including size, land cover, and climate.

Introduction

Urban land expansion is occurring at a faster rate than urban population growth[1], transforming

land cover, climate, hydrology, and biogeochemical cycling[2]. The dramatic environmental

change associated with urbanization renders cities far more similar to one another in many

abiotic and biotic characteristics than to adjacent rural lands[3,4]. For example, cities have

extensive impervious cover (e.g., buildings, transportation networks), increased air temperature,

and increased air, water, light, and noise pollution[2].

The convergence of environmental conditions in cities makes urban ecosystems an ideal

location to address longstanding questions in evolutionary biology about the degree to which

populations exhibit parallel evolution, that is, when multiple populations evolve very similar

traits[5]. Cities are thought to drive parallel evolution as urban populations adapt to similar

selection pressures[6,7,8,9]. Cities are not homogenous, however, and variation in their

characteristics likely mediates processes that facilitate or inhibit parallel evolution, generating

variation in the shape of genotypic or phenotypic clines along urbanization gradients. Clines may

be strongest in large cities where the environmental gradients driving divergent selection

between urban and rural areas are steepest. Conversely, clines may be weak or nonexistent in

small cities or cities with extensive greenspace connecting urban and rural populations via gene

flow. Environmental conditions at large spatial scales – such as regional climate – may also

modulate the evolution of urban-rural clines at a smaller scale. For example, directional selection

for traits that reduce thermal stress at climate extremes may overwhelm selective pressures

associated with local land use[10]. Advances in our understanding of how landscape-scale

differences among cities facilitate or constrain parallel evolution have been limited by a lack of

studies conducted across multiple, replicate cities[7,8,9]. Addressing this knowledge gap will help

clarify how urbanization, the megatrend of global land use change, will shape biodiversity.

Here we test for parallel evolution along urban-rural clines and examine how landscape

features mediate the shape of clines. Our focus is on melanism in eastern gray squirrels (*Sciurus

carolinensis*) in 43 cities in North America (Fig. 1). Gray squirrels are among the most

ubiquitous of fauna in cities and rural woodlands in eastern North America[11]. Coat color in gray

squirrels is typically gray or black (melanic) and inherited in a simple Mendelian fashion at the

melanocortin 1 receptor gene (*Mc1R*)[12]. Evidence suggests divergent selection associated with

crypsis favors the melanic morph in cities and the gray morph in rural woodlands[13], but urban-

rural divergence in melanism has been documented in only two cities (Syracuse, NY[13] and

Wooster, OH[14]). We used a dataset on coat color from squirrel observations collected via the

community science project *SquirrelMapper* to test for parallel clines in melanism along gradients

of impervious cover. We included city size, forest cover across each urbanization gradient, and

winter temperature in a linear mixed model to test how these landscape-scale characteristics

affect the prevalence of melanism in each city (i.e., main effects of each covariate) and drive

variation in cline shape among cities (i.e., interaction effects with impervious cover). Gray

squirrels can disperse long distances (>10 km)[15], but movements are limited by forest

fragmentation[16], so we expected clines to be weakest in small cities and cities with extensive

forest cover where gene flow between urban and rural areas should be greatest. We included

winter temperature in the model because melanic squirrels are known to have greater

thermogenic capacity than the gray morph at cold temperature[17]. Clines may be weak in cities

with cold winter temperature if thermal selection overwhelms the selection pressures on

melanism associated with urbanization.

Results and Discussion

Our analyses revealed the prevalence of melanism in these mammals was positively related to

the degree of urbanization across multiple cities in North America (main effect of impervious

cover, Figs. 1-2; Supplementary Table 2). Repeated shifts in phenotype along environmental

gradients – such as the general increase in melanism from rural to urban areas documented here –

is a hallmark of parallel evolution. Melanism in gray squirrels is a simple Mendelian trait coded

by a 24-bp deletion at *Mc1R*[12], so clines in melanism very likely occur at phenotypic and

genotypic levels.

The presence of parallel clines in heritable traits is often considered evidence of adaptive

evolution because gene flow should homogenize populations in the absence of natural

selection[18]. Known drivers of parallel adaptive evolution in cities from other systems include the

urban heat island[19] and pollution[20]. Evidence suggests multiple selective mechanisms involving

crypsis help maintain clines in melanism in gray squirrels. Predation – including hunting of

squirrels by humans – is often the main driver of squirrel mortality in rural woodlands[21,22], and

the gray morph is more concealed than the melanic morph in regrown, secondary forests that

dominate rural woodlands in our study area[13].

Conversely, predation pressure on tree squirrels is relaxed in cities, where the primary

driver of mortality tends to be vehicular collisions[22]. The melanic morph is more visible to

vehicle drivers than the gray morph on asphalt roads, which may lead to the melanic morph

having greater survival in areas with high car traffic[13]. An alternative explanation for the urban-

rural clines documented here is that melanic squirrels have been introduced to some cities as

novelty[23]. However, these introductions do not explain the low prevalence of the melanic morph

in adjacent rural woodlands given the dispersal ability of the species. Moreover, melanism was

common in northern old growth forests prior to the 1800s[24,25,26], including regions where it is

now rare except in cities (e.g., upstate New York[13,25]). We hypothesize that the dramatic shift

from old growth to secondary forests following European settlement in North America[27,28,29]

changed the degree to which gray and melanic morphs are concealed from visual predators and

hunters, favoring the gray morph in regrown forests[13]. Additional selective factors may play a

role in driving urban melanism (e.g., parasite pressure, pollution)[30], and ultimately genomic tools

and field experiments are needed to clarify the role of natural selection in generating urban-rural

clines[31].

Although clines in melanism tended to take the same form among cities, the strength of

urban-rural clines varied, supporting the idea that repeated evolution is often imperfect[5]. First,

clines most commonly evolved in northern cities with cold winter temperature (Fig. 1b), whereas

clines in melanism were absent in cities with warm winters where melanism was rare or

nonexistent (main effect of winter temperature, Figs. 1-2; Supplementary Table 2). The melanic

morph has greater capacity for non-shivering thermogenesis and retains more heat than the gray

morph at cold temperature[17,32,33]. Our results suggest thermal selection at a regional scale plays

an important role in maintaining melanism in the north at a level where processes operating at a

finer spatial scale cause the evolution of urban-rural clines. Winter temperature has also been

shown to affect the strength of urban-rural clines in cyanogenesis in white clover (*Trifolium*

repens)[34].

Second, the positive relationship between urbanization and melanism was strongest in

large cities with extensive forest cover (interaction effects of city size and forest cover with

impervious cover, Fig. 2; Supplementary Table 2), where melanism increased by nearly 20%

from rural to urban areas (Fig. 2). Although gray squirrels have substantial capacity for long-

distance dispersal, gene flow between urban and rural areas should be most limited in large cities

due to isolation-by-distance and mortality caused by extensive road networks during dispersal.

Gene flow may be strongest between urban and rural populations in smaller cities, maintaining

maladapted morphs across the urbanization gradient[9]. Selection associated with road mortality in

urban areas may also be weak in small cities with limited traffic, which could explain why

melanism was rare in smaller cities (main effect of city size, Fig. 2).

The finding of strong clines in cities with extensive forest cover (Fig. 2) did not support

our prediction that forest cover facilitates gene flow and weakens clines along the urbanization

gradient. Gray squirrels require trees, and squirrel abundance and distribution along urbanization

gradients is limited by greenspace availability, particularly in large cities[35]. It is possible that

cities with more extensive forest cover support a greater abundance of gray squirrels, increasing

effective population size and thereby making selection more efficient in overcoming the effects

of genetic drift on morph frequencies.

Our work reinforces the idea that ecological homogenization among cities drives parallel

evolution[19,20], and it reveals how cities can be a refuge for unique traits generated via rapid

evolutionary response[36]. Our study also provides insight into how landscape-scale differences

among cities alter evolutionary processes associated with urbanization. We show the degree of

parallelism in trait evolution among cities is imperfect, with environmental heterogeneity among

cities contributing to variation in the shape of urban-rural clines. Greenspace availability in

particular has been a focus in the ecological literature for understanding among-city variation in

the response of species distributions to urbanization[35] – our results show that greenspace

availability in the form of forest cover likely mediates evolutionary responses to urbanization as

well. Finally, our work underscores the value of conducting studies in urban evolution across

multiple cities, for which leveraging community science data has great utility.

Methods

Squirrel datasets. Squirrel records were sourced from the community science program

SquirrelMapper. Starting in 2010, participants used a web platform to submit sightings of *S.*

carolinensis, including the date of observation and color morph of each squirrel ("gray",

"melanic", or "other"). Spatial coordinates for each observation were secured via Google Maps

JavaScript API. A total of 5,899 records were submitted to the original *SquirrelMapper* program

from the native range of *S. carolinensis* in North America. Beginning in 2019, *SquirrelMapper*

was integrated with *iNaturalist*, enabling users to submit georeferenced photos of each squirrel

observation. All images of *S. carolinensis* submitted to *iNaturalist* through 21 December 2021

and classified as "research grade" were included for analysis (n = 75,130).

We used the participatory science portal *Zooniverse* to crowdsource the classification of

coat color of squirrels in images submitted to *iNaturalist*. Participants were asked to classify the

number of squirrels observed in each image ("zero", "one", "two or more") and the coat color of

squirrels present in each image ("gray", "melanic", "other", or "unclear"). Each image was

classified by a minimum of 10 participants (range: 10-291; median = 10). We retained images

with a minimum agreement threshold of 80% for the number of squirrels observed. Of the

72,239 images (96%) that met the agreement threshold, 865 were classified as having no squirrel

present (e.g., track or other sign), 70,258 had a single squirrel, and 1,116 had two or more

squirrels. Of the images classified as having at least one squirrel, we retained 63,506 images that

had a minimum agreement threshold of 80% for coat color of at least one squirrel in the image.

We removed 5,182 images that did not have location data available and 4,035 images taken

outside of North America where *S. carolinensis* is introduced. The final *iNaturalist* dataset

consisted of 54,289 images, of which 54,215 were classified as having a single coat color present

(gray: 46,630, melanic: 6,728, other: 857) and 74 had two color morphs present (gray and

melanic: 68, gray and other: 5, melanic and other: 1). A random sample of 500 images classified

by the authors showed that the classification scheme we used with 80% agreement thresholds

was 100% accurate for the number of squirrels observed in the image and 99.8% accurate for

coat color, the only discrepancy being a single image classified as melanic that the authors

classified as unclear. The final *iNaturalist* dataset was merged with squirrel records from the

original *SquirrelMapper* project, yielding 60,262 squirrel records in total. Multiple coat colors

observed at the same location were treated as separate records in this final dataset. Observations

in the final dataset were made between 1969 and 2020 (median = 2019), with 95% of

observations being reported since 2010.

City selection. We selected cities in the native range of *S. carolinensis* that had at least 100

squirrel observations in the city and surrounding rural area. We defined the spatial extent of each

city's footprint using the Urban Area National Shapefile Record Layout (United States Census

Bureau) for the United States and the Population Center Boundary File (Statistics Canada

Catalogue no. 92-160-G) for Canada. We calculated the distance from the centroid of each city

footprint to its maximum extent in each of the four cardinal directions. We then buffered each

urban area by 25% of the average distance from the city centroid to its extent. This buffer

distance was small enough to include a significant area of rural land surrounding each city but

not so large as to produce excessive overlap between polygons of adjacent cities. We determined

the number of squirrels in each city polygon (encompassing the city footprint and hinterland)

after spatially thinning the squirrel records to a minimum distance of 10 m using the s*pThin*

package (v. 0.2.0)[37] in R[38], which helps minimize spatial sampling bias. If two cities with ≥100

squirrels had polygons that overlapped, we retained the city with a greater sample size or spatial

spread of squirrel observations. The final dataset included 26,924 squirrels among 43 cities (Fig.

1), and the median number of squirrels in each city was 254 (range: 101 – 4,731; Supplementary

Table 1; maps of squirrel observations in each city available for download at

https://github.com/bcosentino/squirrelMapper).

Environmental covariates. We measured percent impervious cover within 1 km around each

squirrel observation using the Global Man-made Impervious Surface Dataset (30-m resolution)[39].

A 1-km buffer was chosen to encompass the typical home range size of *S. carolinensis* (<5 ha)

and long-distance movements typical of males during winter and summer mating seasons[40]. For

each city and its surrounding rural area we measured forest cover and winter temperature. Forest

cover was measured using the Global 2010 Tree Cover database (30-m resolution)[41], and we

used WorldClim (version 2.1) to measure winter temperature as mean temperature of the coldest

quarter (BIO11, 30s resolution)[42]. We also measured the area of each city footprint as a metric of

city size. All spatial analyses were conducted in R using the *raster* (v. 3.3-13)[43] and *exactextractr*

packages (v. 0.5.1)[44].

Statistical analyses. We fit a generalized linear mixed model with a binomial error distribution

to test for effects of impervious cover, city size, winter temperature, and forest cover on the

distribution of melanism. Melanism was the response variable and represented by a binary

indicator, with 1 corresponding to melanic and 0 to other coat colors for each squirrel

observation. Impervious cover within 1 km of each observation was included as a fixed effect to

represent each city's urbanization gradient. Because the range of impervious cover varied among

cities, we rescaled impervious cover within each city to range from 0 (lowest impervious cover)

to 1 (greatest impervious cover)[34]. This ensured the slope term for impervious cover has a

consistent interpretation among cities, allowing us to test for a general urban-to-rural cline in

melanism. City size (log-transformed), forest cover, and winter temperature were included as

fixed effects to test whether these variables predict variation in the prevalence of melanism

among cities. Two-way interaction terms were included between each predictor and impervious

cover to test how city characteristics affect the shape of urban-to-rural clines in melanism. We

standardized each predictor so that each variable had mean = 0 and SD = 1, which aided in

model convergence and allowed us to compare the effect size of each slope term[45]. We also

included city as a random effect to account for the nested observations within each city and to

estimate unexplained variance in the prevalence of melanism among cities (Supplementary Table

3). The model was fit with the *lme4* package in R (v. 1.1-26)[46]. Two-sided Wald tests were

conducted for each parameter in the model. Binned residual plots were generated using the *arm*

package (v. 1.11-2)[47] as recommended for models with a binary response variable[48]. Plots of

binned residuals on fitted values and each predictor suggested adequate model fit and no

substantial deviation from model assumptions.

We used the residual autocovariate approach[49] to account for spatial autocorrelation in

the residuals of the model. We extracted residuals from the global model and created a spatial

autocovariate for the residuals using the *spdep* package (v. 1.1-7)[50]. We set the radius

neighborhood to 31 km (the greatest nearest neighbor distance in the thinned dataset) and used

an inverse distance weighting scheme with style "B" for symmetric weights[51]. Our global model

was then refit with the spatial autocovariate term as an additional predictor, which reduced

spatial autocorrelation in the residuals from a maximum of $r = 0.11$ within 1 km to $r < 0.05$

(Moran's I; Supplementary Fig. 1).

We refit our model using different spatial thinning distances (10 m, 50 m, 100 m) and

buffer distances to measure impervious cover (500 m, 1 km, 10 km). These analyses showed the

model results were not sensitive to spatial thinning distances (Supplementary Table 4) or buffer

distances to quantify impervious cover around squirrel observations (Supplementary Table 5).

References

1. Angel, S. et al. The dimensions of global urban expansion: Estimates and projections for all

countries, 2000–2050. *Prog. Plann.* **75,** 53-107 (2011).

2. Grimm, N. B. *et al*. Global change and the ecology of cities. *Science* **319,** 756-760 (2008).

3. McKinney, M. L. Urbanization as a major cause of biotic homogenization. *Biol. Conserv.*

127, 247-260 (2006).

4. Groffman, P. M. *et al*. Ecological homogenization of urban USA. *Front. Ecol. Environ.* **12,**

74-81 (2014).

5. Bolnick, D. I. *et al*. (Non)Parallel evolution. *Annu. Rev. Ecol. Evol. Syst.* **49,** 303-330 (2018).

6. Donihue, C. M. & Lambert, M. R. Adaptive evolution in urban ecosystems. *Ambio* **44,** 194-

203 (2015).

7. Johnson, M. T. J. & Munshi-South, J. Evolution of life in urban environments. *Science* **358,**

eaam8327 (2017).

8. Rivkin, L. R. *et al*. A roadmap for urban evolutionary ecology. *Evol. Appl.* **12,** 384-398

(2019).

9. Santangelo, J. S. *et al*. Urban environments as a framework to study parallel evolution in

Urban Evolutionary Biology, Szulkin, M., Munshi-South, J. & Charmantier, A. (Eds).

(Oxford University Press, 2020).

10. Cosentino, B. J., Moore, J-D., Karraker, N. E., Ouellet, M. & Gibbs, J. P. Evolutionary

response to global change: climate and land use interact to shape color polymorphism in a

woodland salamander. *Ecol. Evol.* **7,** 5426-5434 (2017).

11. Koprowski, J. L., Munroe, K. E. & Edelman, A. J. Gray not grey: ecology of *Sciurus*

carolinensis in their native range in North America in *Grey Squirrels: Ecology and*

Management of an Invasive Species in Europe, Shuttleworth, C. M., Lurz P. W. W. &

Gurnell, J (Eds). (European Squirrel Initiative, 2016).

12. McRobie, H., Thomas, A. & Kelly, J. The genetic basis of melanism in the gray squirrel

(*Sciurus carolinensis*). *J. Hered.* **100,** 709-714 (2009).

13. Gibbs, J. P., Buff, M. F. & Cosentino, B. J. The biological system – Urban wildlife,

adaptation and evolution: urbanization as a driver of contemporary evolution in gray squirrels

(*Sciurus carolinensis*) in *Understanding Urban Ecology*, Hall, M.A. & Balogh, S. (Eds).

(Springer, 2019).

14. Lehtinen, R. M. *et al*. Dispatches form the neighborhood watch: Using citizen science and

field survey data to document color morph frequency in space and time. *Ecol. Evol.* **10,**

1526-1538.

15. Perlut, N. G. Long-distance dispersal by eastern gray squirrels in suburban habitats.

Northeast. Nat. **27,** 195-200 (2020).

16. Goheen, J. R., Swihart, R. K, Gehring, T. M. & Miller, M. S. Forces structuring tree squirrel

communities in landscapes fragmented by agriculture: species differences in perceptions of

forest connectivity and carrying capacity. *Oikos* **102,** 95-103 (2003).

17. Ducharme, M. B., Larochelle, J. & Richard, D. Thermogenic capacity in gray and black

morphs of the gray squirrel, *Sciurus carolinensis*. *Physiol. Zool.* **62,** 1273-1292 (1989).

18. Linnen, C. R. & Hoekstra, H. E. Measuring natural selection on genotypes and phenotypes in

the wild. *Cold Spring Harb. Symp. Quant. Biol.* **74,** 155-168 (2010).

19. Campbell-Staton, S. C. *et al*. Parallel selection on thermal physiology facilitates repeated

adaptation of city lizards to urban heat islands. *Nat. Ecol. Evol.* **4,** 652-658 (2020).

20. Reid, N. M. *et al*. The genomic landscape of rapid repeated evolutionary adaptation to toxic

pollution in wild fish. *Science* **354,** 1305-1308 (2016).

21. Bowers, M. A. & Breland, B. Foraging of gray squirrels on an urban-rural gradient: use of

the GUD to assess anthropogenic impact. *Ecol. Appl.* **6,** 1135-1142 (1996).

22. McCleery, R. A., Lopez, R. R., Silvy, N.J. & Gallant, D. L. Fox squirrel survival in urban

and rural environments. *J. Wildl. Manage.* **72,** 133-137 (2008).

23. Benson, E. The urbanization of the eastern gray squirrel in the United States. *J. Am. Hist.*

100, 691-710 (2013).

24. Allen, D. L. *Michigan Fox Squirrel Management.* (Michigan Department of Conservation,

1943).

25. Schorger, A.W. Squirrels in early Wisconsin. *Trans. Wis. Acad. Sci. Arts Lett.* **39,** 195-247

(1949).

26. Robertson, G. I. *Distribution of Color Morphs of* Sciurus carolinensis *in Eastern North*

America. (University of Western Ontario, 1973).

27. MacCleery, D. W. *American Forests: A History of Resiliency and Recovery.* (Forest History

Society, 2011).

28. Foster, D. R. *et al. Wildlands and Woodlands: A Vision for the New England Landscape.*

(Harvard University Press, 2010).

29. Thompson, R. T., Carpenter, D. N., Cogbill, C. V., & Foster, D. R. Four centuries of change

in northeastern United States forests. *PLoS ONE* **8(9),** e72540 (2013).

30. Leveau, L. United colours of the city: A review about urbanization impact on animal colours.

Austral Ecol. **46,** 670-679 (2021).

31. Lambert, M. R. *et al.* Adaptive evolution in cities: progress and misconceptions. *Trends Ecol.*

Evol. **36,** 239-257 (2021).

32. Farquhar, D. N. *Some Aspects of Thermoregulation as Related to the Geographic*

Distribution of the Northern Melanic Phase of the Grey Squirrel. (York University, 1974).

33. Innes, S. & D. M. Lavigne. Comparative energetics of coat colour polymorphs in the eastern

gray squirrel *Sciurus carolinensis*. *Can. J. Zool.* **57,** 585-592 (1979).

34. Santangelo, J. S. *et al*. Predicting the strength of urban-rural clines in a Mendelian

polymorphism along a latitudinal gradient. *Evol. Lett.* **4,** 212-225 (2020).

35. Fidino, M. *et al*. Landscape-scale differences among cities alter common species' responses

to urbanization. *Ecol. Appl.* **31,** e02253 (2021).

36. Alberti, M. Global urban signatures of phenotypic change in animal and plant populations.

Proc. Natl. Acad. Sci. U.S.A. **114,** 8951-8956 (2017).

37. Aiello-Lammens, M. E. *et al*. spThin: an R package for spatial thinning of species occurrence

records for use in ecological niche models. *Ecography* **38,** 541-545 (2015).

38. R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. (2020).

39. Brown de Colstoun, E. C. *et al*. *Documentation for the Global Man-made Impervious*

Surface (GMIS) Dataset from Landsat. (NASA Socioeconomic Data and Applications

Center, 2017).

40. Steele, M. A. & Koprowski, J. L. *North American Tree Squirrels.* (Smithsonian Books,

2001).

41. Hansen, M. C. *et al*. High-resolution global maps of 21st-century forest cover change. *Science*

342, 850-853 (2013).

42. Fick, S. E. & Hijmans R. J. WorldClim 2: new 1km spatial resolution climate surfaces for

global land areas. *Int. J. Climatol.* **37,** 4302-4315 (2017).

43. Hijmans, R. L. raster: Geographic data analysis and modeling. R package version 3.3-13.

(2020). https://CRAN.R-project.org/package=raster.

44. Baston, D. exactextractr: Fast extraction from raster datasets using polygons. R package

version 0.5.1. (2020). https://CRAN.R-project.org/package=exactextractr.

45. Harrison, X. A. *et al*. A brief introduction to mixed effects modelling and multi-model

inference in ecology. *PeerJ* **6,** e4794 (2018).

46. Bates, D. , Maechler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using

lme4. *J. Stat. Softw.* **67,** 1-48 (2015).

47. Gelman, A. & Su, Y. arm: Data analysis using regression and multilevel/hierarchical models.

R package version 1.11-2. (2020). https://CRAN.R-project.org/package=arm.

48. Gelman, A. & Hill, J. *Data Analysis Using Regression and Multilevel/Hierarchical Models.*

(Cambridge University Press, 2007).

49. Crase, B., Liedloff, A. C., & Wintle, B. A. A new method for dealing with residual spatial

autocorrelation in species distribution models. *Ecography* **35,** 879-888 (2012).

50. Bivand, R. S. & Wong, D. W. S. Comparing implementations of global and local indicators

of spatial association. *TEST* **27,** 716-748 (2018).

51. Bardos, D. C., Guillera-Arroita, G. & Wintle, B. A. Valid auto-models for spatially

autocorrelated occupancy and abundance data. *Methods Ecol. Evol.* **6,** 1137-1149 (2015).

Acknowledgements

This research was supported by the National Science Foundation (DEB 2018249). We thank B.

Fischman for comments on the manuscript.

Author Contributions

B.J.C. and J.P.G designed the study. B.J.C. led the data analysis and wrote the first draft of the

manuscript. J.P.G. contributed ideas for data analysis and edited the manuscript.

Data Availability

The observations of eastern gray squirrels (*Sciurus carolinensis*) generated and analyzed for this

project can be found at https://github.com/bcosentino/squirrelMapper.

Competing Interests

The authors declare no competing interests.

Fig. 1. a, Distribution of coat color morphs of 26,924 eastern gray squirrels (*Sciurus*

carolinensis) among 43 cities in North America. Pie charts show the proportion of melanic and

gray color morphs in each city. Gray morph included rare color morphs that accounted for 2% of

observations (Supplementary Table 1). **b,** Clines in melanism along urbanization gradients in

each city. Urbanization was measured as standardized impervious cover. Regression lines

represent predicted effect of urbanization on melanism in each city based on a linear mixed

model including effects of city size, forest cover, and winter temperature (Fig. 2, Supplementary

Table 2). Photo 'grey squirrel, black squirrel' (tinyurl.com/pauha835) by Eyesplash Photography

is licensed under CC BY-NC-ND 2.0 (https://creativecommons.org/licenses/by-nc-nd/2.0/).

Fig. 2. a, Effect sizes for fixed effects of urbanization, winter temperature, forest cover, city size,

interaction terms with urbanization, and a spatial autocovariate from a linear mixed model of

melanism in 26,924 eastern gray squirrels *(Sciurus carolinensis)* among 43 cities in North

America. Effect sizes represent standardized regression coefficients and include 95% confidence

intervals. Urbanization was measured as standardized impervious cover. **b**, Relationship of the

probability of melanism to urbanization at low (-5°C) and high (5°C) winter temperature while

holding city size and forest cover constant. **c**, Relationship of the probability of melanism to

urbanization at varying levels of city size and forest cover. Effects of urbanization on melanism

are shown for combinations of small cities (25,000 ha), large cities (350,000 ha), low forest

cover (15%), and high forest cover (45%) while holding winter temperature constant at -5°C.