Charged sequence motifs increase affinity towards liquid-liquid phase separation

András L. Szabó, Anna Sánta, Zoltán Gáspári

Pázmány Péter Catholic University, Faculty of Information Technology and Bionics, 1083 Budapest, Práter u. 50/A

Abstract

Protein phase separation has been shown to be a major governing factor in multiple cellular processes, especially ones concerning RNA and RNA-binding proteins. Despite many key observations, the exact structural characteristics of proteins involved in the process are still not fully deciphered. In this work we show that proteins harbouring sequences with specific regions of charged residues are significantly associated with phase separation phenomena. In particular, regions with repetitive arrays of alternating charges (termed charged residue repeats, CRRs) show the strongest association, whereas segments with generally high charge density (charge-dense regions, CDRs) and single alpha-helices (SAHs) show also detectable but weaker connections.

It is known to contribute to the formation of membrane-less organelles (MLOs) and to an extent the aggregation of proteins. The causes and consequences of phase separation has been a rigorously researched topic in the last few years, as the condensation of specific phase-separating proteins is known to promote several diseases.

In this work we carried out a computational analysis to examine the presence of repetitive segments with high charge density in proteins prone to phase separation. Free resources such as the Charged Single α-Helix (CSAH) web server and the PhaSepDB online database were used to examine possible links between the charged side-chain content of protein sequences and their partition into membrane-less condensates. Furthermore, we carried out the development of a novel algorithm aimed to detect a larger variety of charged protein segments, in order to examine their relationship to the phenomenon. Fisher's exact test of independence was implemented on several generated data sets to confirm correlation between charged residue repeats (CRRs) and charge-dense regions (CDRs) within human protein sequences and their affinity for phase separation.

Introduction

The general phenomenon of phase separation

Cells are continuously conducting various complex biochemical processes, one of which is a set of intermolecular interactions resulting in the de-mixing of macromolecules such as multi-domain proteins and RNA molecules, and the formation of so-called membrane-less granules. As a relatively recent discovery, it has come to the focus of intensive research since the early 2010s. The general name for the phenomenon is phase separation, and it has been shown to have a role in critical processes of the living cell, such as chromatin regulation, RNA transcription, organization of the postsynaptic density etc... [39, 41, 40]

The molecular assemblies formed by these processes are typically micron-sized objects, containing multivalent proteins with multiple modular domains and / or charged elements. There are two main types of molecules participating in phase separation, as distinguished by Banani et al. [2], the "scaffolds" and the "clients". The former type consists of molecules essential to the integrity of the MLOs. The latter type contains the majority of the components; however, they only participate in functionalities under certain circumstances. P bodies are a good example for this duality, scaffolded by RNA-binding proteins and storing RNA as clients. Scaffold-scaffold interactions are more persistent than scaffold-client interactions, and the composition changes according to a set of factors such as stress and the cell cycle. [38, 39]

Liquid-liquid phase separation in particular

Liquid-liquid phase separation (LLPS) is responsible for the formation of membrane-less organelles (MLOs), intracellular granules whose main function is the concentration of proteins and nucleic acids – mainly RNAs – in response to external signals such as stress. The characteristic feature of LLPS that distinguishes it from other processes related to phase separation is that the solution transitions into distinct phases where the certain solutes are present in highly elevated concentrations, and these phases exhibit liquid-like properties. In this specific type of phenomenon, the term "scaffolds" is often replaced by "drivers", as in sets of proteins that are able to drive LLPS on their own, disregarding small molecules and ions even if they are required for the initiation of the condensation process. Clients in this context are molecules that may partition into MLOs but they hold no influence over their formation [3]. A category of proteins whose members are especially prone to such processes are intrinsically disordered proteins (IDPs). These biochemical components typically participate in multivalent interactions to self-assemble into MLOs through their modular domains and linear motifs which govern the transition from liquid phase to gel structure, and then solid state. Liquid-solid phase transitions – in specific cases called aggregation – are often associated with severe diseases such as amyotrophic lateral sclerosis (ALS), making the examination of IDPs and LLPS rigorously researched fields.

The role IDRs play in mediating LLPS was also stressed by Lin et al. in 2017 [4], when they fused IDRs derived from FUS with poly-Src homology 3 (polySH3) proteins. The fusion of the wild-type (WT) IDR lowered the concentration threshold for LLPS of polySH3-polyPRM system, where polyPRM stands for poly-proline-rich motif, a polypeptide consisting of regions known to bind SH3 domains. While WT IDR-fused SH3 domains exhibited increased LLPS-activity (indicated by higher turbidity in the figure below), SH3 domains either mixed or fused to mutant IDRs displayed a decrease in droplet formation. Such mutations were introduced to the tyrosine residues in the Gly/Ser-Tyr-Gly/Ser motifs of the IDR.

Earlier, Pak et al. [50] described the phase separation of the intrinsically disordered intracellular domain of Nephrin (NICD) via complex coacervation. When expressed as a soluble protein, NICD formed droplets in HeLa cells, and LLPS was also observed in vitro when mixed with positively charged supercharged GFP. While no specific sequence was identified in NICD responsible for LLPS, a pattern of blocks of negatively charged residues was observed. This means phase separation proved to be robust in withstanding mutations, as shuffle and deletion mutants were still able to drive phase separation as long as charged blocks were retained.

Many proteins involved in phase separation phenomena have been identified in the last 20 years, and thus it has become a relevant issue to catalogue them. PhaSepDB [5] is a manually curated database that has been created with the specific purpose of providing researchers with a detailed, reliable collection of information about the proteins connected to the phenomenon. There are other, more specific databases of phase-separating protein such as the stress granule protein database constructed by Catarina et al. [6], or PhaSePro that specifically collects LLPS drivers [42], but as of today one of the most comprehensive databases is PhaSepDB. Data collection and processing for such a library of proteins consists of several levels, and new entries are added into the database after their respective publications have been curated twice to confirm the relation to LLPS.

There are three categories of protein sets available in PhaSepDB: The first one is the "Reviewed" category that only contains articles published after January 1st of 2000, constituting the most reliable subset. The second category is called "UniProt reviewed" that includes a wider range of articles, not restricted by publication date, although its results must be confirmed by more recent studies. The third and final category is the "High-throughput" research data, generated by high-throughput methods categorized as either organelle purification, proximity labelling, immunofluorescence image-based screen or affinity purification. A plethora of computational tools is also applied for the prediction of protein sequences deemed worthy of being added to the database. Such programs include PONDR and ESpritz for disorder prediction, PLAAC for identifying sequences with prion-like amino acid composition(s), Pi-Pi for electrostatic interaction prediction, and CIDER for charged/hydrophobic residue distribution.

ESpritz is based on a bidirectional recurrent neural network that functions as an ensemble of three neural networks, one devoted to the N-terminal sequence context, one devoted to the C-terminal sequence context and one for the (middle)

sequence. The output contains two units producing the probability of order and disorder. [34] PLAAC uses a windowing function to scan individual sequences for domains with Prion-Like Amino Acid Composition. The minimum length of the prion domains can be specified, as well as a parameter for interpolation between organism-specific background frequencies. Per-residue scores are compiled into sliding averages and summary tables for the input sequence(s). Multiple sequences can be ranked by the maximum sum of per-residue log-likelihood ratios for any subsequence of the specified length that are identified as prion-like domains. [35] The ESpritz score as well as the prion-like domain probability from PLAAC are included in PhaSepDB as parts of per-residue plots describing the characteristics of a given sequence, justifying its inclusion into the high-throughput dataset.

Single α-Helices and other charged residue repeats

Single α-helices (SAHs) are protein segments that form stable and rigid helical structures even in isolation. [36, 37] These regions are rich in arginine, lysine and glutamate and exhibit a characteristic repetitive pattern of oppositely charged residues such as glutamic acid and lysine. Their length may vary from a few dozens up to about 200 residues, and they are stabilized by intrahelical salt bridges [7]. These regions show similar characteristics to coiled coils, as Peckham et al. [8] showed that ~4% of human proteins that were previously predicted to contain coiled coils actually have SAHs instead. SAH domains have been found to be prevalent in RNA-binding proteins, although the SAH region itself is unlikely to directly contribute to the interaction with RNA.

Inspired by the role of RNA molecules in phase separation, we intended to investigate whether proteins involved in LLPS are enriched in SAH segments. As SAH segments constitute a special case of sequences with high charge density, we have extended our analysis to segments with repeating patterns of charged residues, termed charged residue repeats, CRRs, as well as with segments enriched in charged residues but without any further specifications, denoted charge-dense regions, CDRs, below. We differentiate between "unsigned" and "signed" CRRs on the basis of their net charge. In this classification, most CRRs are special cases of CDRs, and SAHs constitute a sub-class of CRRs (Figure 1). We found that proteins with CRRs exhibit especially strong association with LLPS formation.

Figure 1. Venn diagram illustrating the relations between single α-helices, charged residue repeats, as well as signed and unsigned charge-dense regions within the human proteome.

It is important to note that while the majority of CRRs satisfy at least one of the two definitions for charge-dense regions, not all of them fall under those categories. The cleavage stimulation factor subunit 2 (CSTF2, UniProtKB ID: P33240) for example contains 12 X 5 AA tandem repeats on a 60 residue-long segment that was detected as a CRR by the FT_CHARGE algorithm - a 105 residue-long sub-sequence encompassing the annotated region was highlighted, to be exact:

```
magltvrdpavdrslrsvfvgnipyeateeqlkdifsevgpvvsfrlvydretgkpkgyg
fceyqdqetalsamrnlngrefsgralrvdnaaseknkeelkslgtgapviespygetis
pedapesiskavaslppeqmfelmkqmklcvqnspqearnmllqnpqlayallqaqvvmr
ivdpeialkilhrqtniptliagnpqpvhgagpgsgsnvsmnqqnpqapqaqslggmhvn
gapplmqasmqggvpapgqmpaavtgpgpgslapgggmqaqvgmpgsgpvsmergqvpmq
dpraamqrgslpanvptprgllgdapndprggtllsvtgeveprgylgpphqgppmhhvp
ghesrgppphelrggplpeprplmAEPRGPMLDQRGPPLDGRGGRDPRGIDARGMEARAMEA
RGLDARGLEARAMEARAMEARAMEARAMEARAMEVRGMEARGMDTRGPVPGPRGPIPSGMQG
PSPINmgavvpqgsrqvpvmqgtgmqgasiqggsqpggfspgqnqvtpqdhekaal
imqvlqltadqiamlppeqrqsililkeqiqkstgap
```

The region does not qualify as a SAH due to its slightly higher Fourier frequency, although there are multiple helical structures along the entire sequence of this protein. While up to this point we could not find a region that qualifies as a SAH while remaining outside our definitions for CDRs, we would reserve the possibility for their existence as the used representation of the human proteome only included a single isoform per protein.

Results

CDRs, CRRs and SAHs in the human proteome

We have restricted our studies to the human proteome as specified at the beginning of the Methods section. For the prediction of SAHs, we used our previously established methodology, involving the FT_CHARGE algorithm. CRR detection was carried out with FT_CHARGE by relaxing the requirement for the charge frequency characteristic for SAHs. CDRs were detected by an algorithm enumerating Asp, Glu, Arg and Lys residues in sequence windows of different length (see Methods for details). Since the vast majority of human proteins was found to contain a CDR according to our original detection threshold, we have applied a stricter criterion and investigated only the top 1% of CDRs.

Number of proteins				
		Redundancy-filtered proteome		
Sequence motifs	Full proteome	90%	70%	50%
All proteins	20 659	19 638	18 294	15 672
CDRs (unsigned)	9 731	9 314	8 792	7 669
CDRs (signed)	14 065	13 471	12 757	11 097
CRRs (all)	1 054	1 025	985	910
CRRs (>=90% overlap with uCDRs)	782	763	738	688
CRRs (>=90% overlap with sCDRs)	227	221	214	208
SAHs (all)	134	131	126	118
Percentage of residues				
All proteins	100%	96.74%	92.04%	80.73%
CDRs (unsigned)	10.02%	9.63%	9.16%	8.22%
CDRs (signed)	14.53%	13.98%	13.30%	11.58%
CRRs (all)	0.82%	0.80%	0.77%	0.72%
CRRs (>=90% overlap with uCDRs)	0.54%	0.52%	0.50%	0.47%
CRRs (>=90% overlap with sCDRs)	0.14%	0.13%	0.13%	0.12%
SAHs (all)	0.11%	0.11%	0.10%	0.10%

Table 1. Summary table exhibiting the amount of sequences associated with different sequence motifs, as well as the ratio of residues encompassed by those motifs. The table includes data for the full proteome and its variants with reduced redundancy.

Figure 2. Decimal logarithm of the P-values that belong to Fisher's tests where the functional attribute is the protein's relation to LLPS, and the structural attribute is the presence of different sequence motifs.

CRR-containing proteins show significant enrichment in proteins involved in LLPS, while proteins with SAHs show a weaker but still detectable association

Fisher's exact test of independence showed that there is an exceptionally high probability that charged residue repeats (CRRs) contribute to the LLPS of human proteins.

Figure 3. Decimal logarithm of the P-values that belong to Fisher's tests where the functional attribute is the protein's relation to LLPS, and the structural attribute is the presence of charged residue repeats with specific Fourier frequencies.

SAHs can be regarded as special subsets of CRRs detected with FT_CHARGE with the frequency region between 1/9 and 1/6. SAH regions still exhibit a statistically significant enrichment in LLPS-associated proteins, although this is considerably weaker than for CRRs in general. Another way to illustrate the level of association between LLPS and the presence of sequence motifs is to compare their enrichment in LLPS-related proteins. For example, the ratio of sequences with CRRs to other proteins is 1010:19418 in the full proteome in case of entries that are unrelated to LLPS. The same ratio in case of related sequences is 44:187. Which means that the ratio increases 4.52-fold between the two functional categories.

Figure 4. CRR-enrichment (proportional increase of the ratio of CRR-containing sequences) in case of the full proteome as well as its variants with reduced redundancy (with clusters and with representative sequences only).

Regions enriched in charged residues in general are prevalent in proteins associated with LLPS

We have investigated whether regions enriched in charged residues but not necessarily exhibiting regular repeating patterns are also associated with LLPS. To this end, besides the charged residue repeats detected by the FT_CHARGE algorithm, a more general type of protein sub-sequences called "Charge-Dense Regions / CDRs" was similarly probed for relations to LLPS. We have formulated two different approaches for their detection. One of them defines CDRs as protein sub-sequences where the density of charged residues is considerably higher than average. The other one defines them as protein sub-sequences with an overall charge the absolute value of which is significantly higher than average. We refer to the regions gained through the former approach as "unsigned CDRs", while regions resulting from using the other definition were named "signed CDR" - as their charges may have a negative sign, too. The protocols for detecting such components are described in detail in the Method section.

Unsigned and signed CDRs

If we consider the top ~1% of all hits to be charge-dense regions, then approximately 47.10% of human proteins contain some kind of charge-dense region. When these CDRs are compared to CRRs, it turns out that 90.99% of charged residue repeats display an above 90% overlap with CDRs, with 87.80% of all CRRs being entirely encompassed by charge-dense regions, while 2.53% of them are free from any overlap. Sequence-wise, 15.92% of the human proteome is categorized as CDRs with the above-mentioned parameters.

Considering the top 1% of all hits as signed CDRs results in 22.76% of charged residue repeats having at least 90% overlap with the top 1% of CDRs, and 20.25% having total coverage, while 37.14% having no coverage at all.

Additional investigations

Figure 5. Receiver operating characteristic (ROC) tests on CRRs, unsigned CDRs and signed CDRs, evaluating the presence of said motifs as an indicator of affinity towards LLPS. For CRRs the test shows the highest value at 0.5 FPR (TPR = 1) when the threshold is set to 0.38, meaning that sequences where at least 0.38 of all residues are assigned to charged residue repeats exhibit a high probability of participating in LLPS. These numbers are TPR = 0.667 and threshold = 0.80 for uCDRs, and TPR = 0.714 and threshold = 0.83 for sCDRs. The ROC curves suggest that the presence of CDRs/CRRs is not a strong predictor for LLPS but the absence of such motifs is an indicator that the protein is not associated with LLPS.

Figure 6. Number of FT_CHARGE hits organized by their Fourier frequency (1/64 - 32/64). Three separate groups can be distinguished by their prominent number of hits: 1/64-2/64, where charged residues are quite scarce, 30/64-32/64, where about every other residue is charged, and 8/64-10/64, which is within the range of SAHs (1/9-1/6). Hits with frequencies of 7/64 and 11/64 are just outside this range, and are therefore omitted from the group.

The Overrepresentation Test (OT) of Panther's Gene List Analysis tools was also utilized to compare the various special motifs in sequences that are related to LLPS versus the regions of proteins unrelated to the phenomenon. A total of eight tests were carried out, each comparing either LLPS-related or LLPS-unrelated SAHs, CRRs and signed as well as unsigned CDRs. From the OT-given Gene Ontology (GO) terms we manually selected the ones that are at the lowest level within their respective branch of the GO term hierarchy, meaning that only the most specific - and thus least redundant - terms were analyzed further. These terms were compiled into a table where each term's enrichment - or lack thereof - was noted in case of the four different kinds of LLPS-related and LLPS-unrelated motifs. From there it is possible to categorize each term, based on the number of LLPS-related categories they were enriched in (0-4) and the number of LLPS-unrelated categories they were enriched in (0-4). This method of categorization created a gradient table that shows the distribution of GO terms from highly related to LLPS to highly unrelated to LLPS. The table then could be reduced to contain GO terms with specific keywords in them, such as 'regulation', 'RNA' or 'signal'.

Total number of entries = 307					
Enrichment	0/4 Unrelated	1/4 Unrelated	2/4 Unrelated	3/4 Unrelated	4/4 Unrelated
4/4 Related	1	0	0	0	0
3/4 Related	10	0	0	0	0
2/4 Related	53	7	1	0	0
1/4 Related	90	5	3	0	0
0/4 Related	9	112	16	0	0

Total number of entries related to 'RNA' = 55					
Enrichment	0/4 Unrelated	1/4 Unrelated	2/4 Unrelated	3/4 Unrelated	4/4 Unrelated
4/4 Related	0	0	0	0	0
3/4 Related	5	0	0	0	0
2/4 Related	13	1	0	0	0
1/4 Related	26	0	3	0	0
0/4 Related	0	7	0	0	0

Total number of entries related to 'response' = 28					
Enrichment	0/4 Unrelated	1/4 Unrelated	2/4 Unrelated	3/4 Unrelated	4/4 Unrelated
4/4 Related	0	0	0	0	0
3/4 Related	0	0	0	0	0
2/4 Related	10	0	0	0	0
1/4 Related	9	0	0	0	0
0/4 Related	1	6	2	0	0

Total number of entries related to 'metabol' = 22					
Enrichment	0/4 Unrelated	1/4 Unrelated	2/4 Unrelated	3/4 Unrelated	4/4 Unrelated
4/4 Related	0	0	0	0	0
3/4 Related	0	0	0	0	0
2/4 Related	1	0	0	0	0
1/4 Related	2	1	0	0	0
0/4 Related	4	14	0	0	0

Figure 7. Gradient tables depicting the distribution of all GO terms (a), terms related to RNAs (b), terms related to responses (c), terms related to metabolism (d). In the latter case the actual keyword was 'metabol' to include GO terms containing 'metabolism' and 'metabolic' as well, much like the keyword 'RNA' encompasses 'mRNA', 'rRNA', 'snoRNA', 'RNA-binding', etc.

Figure 7. Distribution of human proteins according to their special charged motif (a-b) or CRR (c-d) content, LLPS association and their functional role in the PSD (a, c) or RNA binding (b, d).

It is apparent in the upper cases that there are barely any proteins related to either LLPS, the postsynaptic density or the molecular function of RNA-binding that are devoid of any motifs with either charged patterns or high charge density. Furthermore, the distributions implicate that RNA-binding proteins are more likely to participate in phase separation than PSD components as the ratio of entries in the middle (yellow) section doubles between the two figures. This enrichment can also be observed between PSD proteins and RNA-binding sequences when we only consider CRRs as a structural factor. Although the number of cases is small, it also seems that for RNA-binding proteins, the presence of charged motifs / CRRs is more prevalent when considering proteins participating in LLPS.

Discussion

Case studies

Our results clearly indicate that proteins with regions containing a high fraction of charged residues are preferentially associated with LLPS. Our approach allowed us to investigate multiple aspects of this association, consideration of the presence or absence of specific repetitive patterns. Below we discuss several proteins, the possible relationship between LLPS formation and the presence of SAHs/CRRs/CDRs. Examples, wherever it is possible, are based on the information available in the richly annotated PhaSePro database, where information about the region responsible for LLPS is provided if available.

Table 1. contained 1054 protein sequences that encompass CRRs but out of those only 44 sequences are considered to be LLPS-related. These sequences were investigated further to explore their connection to phase separation. The investigation revealed that a large variety of proteins may participate in the phenomenon, including ribonucleases [17], splicing factors [10, 11], transcriptional repressors [12, 13], translational initiation factors [18], transport- [14, 19] and Zinc finger proteins [15, 16], etc. However, there happens to be a common feature in most of the investigated proteins, as 31 of them carry out the molecular function of RNA-binding [17-28]. The fact that 70.45% of these proteins can bind RNA molecules is significant and might imply that RNA-binding is a process that is promoted by CRRs. Another plausible explanation may be that many membrane-less organelles – that are formed by LLPS – contain RNAs. According to the frequencies of the CRRs predicted by FT_CHARGE, 5/44 sequences possibly contain single α-helices, all of which constitute at least 10% of the respective sequence.

It is important to note that while RNA-binding connects most of these proteins, their sequences as well as their biological function vary greatly. There were only two cases where multiple sequences got clustered together. In the first case the two proteins were the probable global transcription activator SNF2L2 and the transcription activator BRG1. Both are involved in transcriptional activation and repression of select genes by chromatin remodelling [29]. They are also components of SWI/SNF chromatin remodelling complexes that carry out key enzymatic activities and change chromatin structure. While these two proteins proved to be participating in the same processes, they still only showed 59.75% sequence identity.

In the second case there were three proteins in the same cluster, proline- and glutamine-rich splicing factor SFPQ being the representative sequence, non-POU domain-containing octamer-binding protein NONO showing 58.39% identity, and paraspeckle component 1 PSPC1 exhibiting 54.49% sequence identity. The three proteins cooperatively regulate androgen receptor-mediated gene transcription activity in the Sertoli cell line [30, 31] and, besides the noncoding RNA NEAT1, are major components of the paraspeckle. These proteins contain a conserved SAH located at the C-terminal end of a right-handed coiled coil segment. [47] The region responsible for LLPS is the so-called NOPS region on the N-terminal side of the

coiled coil. The same goes for the trinucleotide repeat-containing gene 6B protein (TNRC6B) participates in RNA-mediated gene silencing, and acts as a scaffolding protein that can simultaneously interact with argonaute proteins - regulating miRNA and siRNA maturation - and recruit deadenylase complexes - carrying out poly(A) tail-shortening processes [32]. Finally, the E3 ubiquitin-protein ligase RNF168 accumulates repair proteins to sites of DNA-damage via binding to ubiquitinated histone H2A and H2AX [33]. Apart from SNF2L2 and BRG1, all of the above proteins do encompass at least one alpha helical structure, and all of them seem to exhibit some kind of RNA-binding trait, except RNF168.

SynGAP1 is part of the NMDAR complexes in excitatory synapses. The mouse ortholog has been explicitly shown to participate in LLPS, with the responsible region being its C-terminal segment forming a trimeric coiled coil and having a PDZ-domain binding motif that interacts with PSD-95. The role of SynGAP in LLPS formation in the postsynaptic density has been demonstrated in multiple experiments. [45] The key feature for LLPS in SynGAP is its multivalent nature provided by the trimerization via its coiled coil motif. SynGAP does not contain any SAHs or CRRs detected by our methods, but it bears a CDR and so does its partner protein PSD-95 - a.k.a. Disks large homolog 4. The guanylate kinase-associated protein GKAP - a.k.a. Disks large-associated protein 1 of the DLGAP1 gene - belongs to this protein network, and iss a scaffold protein that organizes the formation of the PSD together with PSD-95, Shank3 and Homer. [51] DLGAP1 contains CDRs and is involved in phase separation.

Synapsin-1 participates in neurotransmitter release in the presynaptic region. The region responsible for phase separation is its Pro/Gln-rich C-terminal half and the protein can form condensates on its own or with binding partners containing SH3 domains. The condensates can also contain synaptic vesicles (SVs), providing the basis of self-organization of SV clusters. Phosphorylation of Synapsin-1 by CamKII causes dissociation of the condensates. [46] Synapsin-1 does not seem to contain any SAHs/CRRs/CDRs.

Figure 8. Sequence motifs as well as PhaSePro-based sub-sequences that drive LLPS. The colour coding of individual motifs as well as their overlaps are in accordance with Figure 1. While currently there isn't any experimental evidence that the human protein SynGAP participates in the phenomenon, its *Mus musculus* homolog has, thus a PhaSePro region has been annotated according to that.

Role of charged regions in LLPS

While our results clearly demonstrate an association between charged segments and LLPS, the rationale behind this is not straightforward. Importantly, the detailed investigation of this association reveals that the absence of such a motif largely precludes the participation of a protein in LLPS and not that the presence of a CDR, CRR or SAH would be a strong indicator of LLPS formation. Here it should be noted that the accuracy of listings of proteins associated with LLPS is expected to increase with novel proteins recognized and curation efforts getting more intensive, probably resulting in an overall increase of proteins reliably proven to form condensates. That said, it is highly unlikely that charged regions identified here in this study in general might be directly responsible for LLPS. It is not the case for all proteins detailed above, let alone phase separating proteins without a CRR or CDR. Thus, it is more likely that the enrichment of CRRs and CDRs in LLPS-prone proteins has a more complex explanation. It should be noted that little is known about the structural features of CRRs and CDRs. Not surprisingly, many regions rich in charged residues, including SAHs, are predicted to be intrinsically disordered by several prediction methods. However, SAHs are known to possess a stable

well-characterized structure and can be recognized by specialized algorithms. Thus, the presence of specific structural preferences for other types of CRRs cannot be completely excluded, and the formation of more or less well-defined monomeric or oligomeric structures might occur for specific sequences. Regularly alternating blocks of positively and negatively charged residues might offer a platform for intermolecular associations in multiple mutual orientations.

Methods

Surveying protein sequences for charged residue repeats

We used the UP000005640 human reference proteome from UniProtKB (https://www.uniprot.org/proteomes/UP000005640), downloaded in fasta format on the 3rd of September 2020. The version of UP000005640 used for this study encompasses 20 659 human genes - with one isoform per gene - and represents the 2020 September 3 update.

To identify segments with regular charge patterns (CRRs), the method FT_CHARGE was used. FT_CHARGE was developed to identify SAH regions in proteins that contain a repeating charge pattern with a specific frequency between 1/9 and 1/6 but it is also capable of detecting other motifs with regularly alternating positively and negatively charged residues. [9] In this work the largest window size for FT_CHARGE was 64 residues, corresponding to a minimum frequency of 1/64. The frequency of 1/64 practically corresponds to a long, repeated segment of identically charged amino acids (e.g. a polylysine run), whereas the frequency of 1/2 corresponds to a region of residues with alternating charges (e.g. the sequence "KEKEKEKEKE"). Such detected sequences are hereafter denoted as "charged residue repeats" - or CRRs for short, to make a distinction between these and "genuine" SAHs. In this view, SAHs represent a subset of CRRs. For more details on the implementation of FT_CHARGE refer to [48, 49].

Clustering human proteins based on structural predictions

To remove redundancy from the sequence set, the webserver CD-HIT was employed to cluster the sequences on their similarity. The cut-off values for this were 0.9, 0.7 and 0.5, meaning 90%, 70% and 50% sequence identity, respectively. Clustering was carried out with global sequence identity, a 20-residue bandwidth of alignment, a minimal sequence length of 10 residues, and default alignment coverage parameters. Sequences were assigned to the best cluster that met the threshold. The output of CD-hit was further processed using MATLAB to ensure compatibility with our analysis pipeline. The three redundancy-filtered arrays obtained have contained clusters of UniProt IDs with a pre-set cut-off value (90%, 70% and 50% sequence identity).

Surveying protein sequences for Charge-Dense Regions (CDRs)

While SAHs - and CRRs in general - may contain a higher-than-average amount of charged residues, their features mainly result from the specific patterns these residues repeat within the sequence. Thus, a separate set of investigations must be carried out if we are to expand the analysis of the correlation between charged protein regions and phase separation. We applied two alternative approaches for identification of such regions, the first one considers a region highly charged if it contains a higher-than-average ratio of charged residues, while the other identifies sub-sequences whose overall charge significantly differs from zero. Based on this difference the recognized "Charge-Dense Regions" (CDRs) will be called "signed" and "unsigned" CDRs, or sCDR and uCDR motifs, respectively.

The scoring scheme for sCDR assigns a score of +1 to arginines, histidines and lysines, and -1 to aspartic and glutamic acids, while the one for uCDRs assigns a score of 1 to all five of the aforementioned residues. In both cases, the investigated sequence window is considered to contain a CDR if the absolute value of its overall score divided by its length reaches a pre-set cut-off value.

To determine adequate cut-off values for all the used window sizes, we run the algorithm on a randomized version of the human proteome with a cut-off value of zero. The randomized sequences had the same amino acid composition and length distribution as the natural ones. Based on the obtained scores, two cut-off values were determined, one that yields roughly 5% of all hits and one that yields only about 1% - note that the discrete nature of scoring (i.e. steps of 1/16 between 0 and 1 for a window length of 16 residues), selection of a threshold is somewhat approximated. Using these two cut-offs, the full proteome was scanned to identify signed and unsigned CDRs in the wild-type sequences.

The segments identified using different sequence windows were merged for each sequence. During the process, care was taken to ensure that the merged segments are still above the predefined threshold, which needs special treatment for signed CDRS where the overall charge should also be maintained. In the final step, all data were unified for all window sizes used, 16, 32, 64 and 128.

Conducting Fisher's exact test of independence on members - and clusters - of the human proteome

To assess the association of specific sequence sets with LLPS, 2x2 contingency tables were generated and then Fisher's exact test as implemented in R [R statistics package reference] was applied to determine the P-value.

In case of clusters, two different methods were applied for categorization. In the first one, evaluation is applied cluster-wise, meaning that association with LLPS for a given cluster was considered positive if any member (sequence) in the cluster was annotated as such in PhaSepDB. In the second, a cluster was only considered to be participating in LLPS if the representative sequence kept by CD-HIT was annotated in PhaSepDB. The presence or absence of CRRs was established in an analogous manner.

Preparing receiver operating characteristic (ROC) tests

The purpose of ROC tests is to evaluate an attribute as a viable indicator for classification. In the case of this study, they were used to investigate the presence of different sequence motifs as indicators for a protein's affinity towards liquid-liquid phase separation. We have sorted the proteins in the human proteome according to their score obtained in the CDR/CRR detection and calculated the true positive, true negative, false positive and false negative rates.

References

1. Nishikawa: Natively unfolded proteins: An overview, *Biophysics*, **2009**, *5*, 53-58
2. Banani et al.: Compositional Control of Phase-Separated Cellular Bodies, *Cell*, **2016**, *166*, 651-663
3. Pancsa et al.: Computational resources for identifying and describing proteins driving liquid-liquid phase separation, *Briefings in Bioinformatics*, **2021**, *Online ahead of print 00(00)*, 1-20
4. Lin et al.: Intrinsically disordered sequences enable modulation of protein phase separation through distributed tyrosine motifs, *JBC*, **2017**, *292(46)*, 19110-19120
5. You et al.: PhaSepDB: a database of liquid-liquid phase separation related proteins, *Nucleic Acids Research*, **2020**, *48*, D354-D359
6. Catarina et al.: MSGP: the first database of the protein components of the mammalian stress granules, *Database*, **2019**, *2019*, baz031
7. Wolny et al.: Characterization of long and stable *de novo* single alpha-helix domains provides novel insight into their stability, *Scientific Reports*, **2017**, 7, 44341
8. Peckham et al.: When a predicted coiled coil is really a single α-helix, in myosins and other proteins, *Soft Matter*, **2009**, *5*, 2493-2503
9. Gáspári et al.: Charged single alpha-helices in proteomes revealed by a consensus prediction approach, *BBA*, **2012**, *1824(4)*, 637-646
10. Kim et al.: De Novo Mutations in SON Disrupt RNA Splicing of Genes Essential for Brain Development and Metabolism, Causing an Intellectual-Disability Syndrome, *American Journal of Human Genetics*, **2016**, *99(3)*, 711-719
11. Heyd et al.: Phosphorylation-dependent regulation of PSF by GSK3 controls CD45 alternative splicing, *Mol Cell.*, **2010**, *40(1)*, 126-137
12. Klar et al.: Enhanceosome formation over the beta interferon promoter underlies a remote-control mechanism mediated by YY1 and YY2, *Mol Cell Biol.*, **2005**, *25(22)*, 10159-10170
13. Wang et al.: Prohibitin requires Brg-1 and Brm for the repression of E2F and cell growth, *EMBO J*, **2002**, *21(12)*, 3019-3028

14. Gaudet et al.: Phylogenetic-based propagation of functional annotations within the Gene Ontology consortium, *Briefings in Bioinformatics*, **2011**, *12(5)*, 449-462

15. Liang et al.: A Novel CCCH-Zinc Finger Protein Family Regulates Proinflammatory Activation of Macrophages, *Journal of Biological Chemistry*, **2008**, *283(10)*, 6337-6346

16. Yamada et al.: Functional analysis and the molecular dissection of zinc-fingers and homeoboxes 1 (ZHX1), *Biochem Biophys Res Commun.*, **2002**, *297(2)*, 368-374

17. Martínez et al.: A 54-kDa Fragment of the Poly(A)-specific Ribonuclease Is an Oligomeric, Processive, and Cap-interacting Poly(A)-specific 3' Exonuclease, *RNA: Structure Metabolism and Catalysis*, **2000**, *275(31)*, P24222-P24230

18. Lee et al.: eIF3 targets cell-proliferation messenger RNAs for translational activation or repression, *Nature*, **2015**, *522*, 111-114

19. Pardo et al.: pARIS-htt: an optimised expression platform to study huntingtin reveals functional domains required for vesicular trafficking, *Mol Brain.*, **2010**, *3*, 17

20. Egan et al.: An Enhanced H/ACA RNP Assembly Mechanism for Human Telomerase RNA, *American Society for Microbiology Journals*, **2012**, *32(13)*, 2428-2439

21. Okuwaki et al.: The RNA Binding Activity of a Ribosome Biogenesis Factor, Nucleophosmin/B23, is Modulated by Phosphorylation with a Cell Cycle-dependent Kinase and by Association with Its Subtype, *Molecular Biology of the Cell*, **2002**, *13(6)*, 2016-2030

22. Castello et al.: Insight into RNA Biology from an Atlas of Mammalian mRNA-binding Proteins, *Cell*, **2012**, *149(6)*, 1393-1406

23. Baltz et al.: The mRNA-bound proteome and its global occupancy profile on protein-coding transcripts, *Mol Cell.*, **2012**, *46(5)*, 674-690

24. Ahn et al.: SON Controls Cell-Cycle Progression by Coordinated Regulation of RNA Splicing, *Mol Cell.*, **2011**, *42(2)*, 185-198

25. Jeon et al.: YY1 Tethers Xist RNA to the Inactive X Nucleation Center, *Cell*, **2011**, *146(1)*, 119-133

26. Pereira et al.: CDX2 regulation by the RNA-binding protein MEX3A: impact on intestinal differentiation and stemness, *Nucleic Acids Research*, **2013**, *41(7)*, 3986-3999

27. Fritz et al.: RNA-Regulated Interaction of Transportin-1 and Exportin-5 with the Double-Stranded RNA-Binding Domain Regulates Nucleocytoplasmic Shuttling of ADAR1, *Mol Cell Biol.*, **2009**, *29(6)*, 1487-1497

28. Zhang et al.: The YTH Domain Is a Novel RNA Binding Domain, *Journal of Biological Chemistry*, **2010**, *285(19)*, 14701-14710

29. Bochar et al.: BRCA1 Is Associated with a Human SWI/SNF-Related Complex, *Cell,* **2000**, *102(2)*, 257-265

30. Sewer et al.: Transcriptional Activation of Human CYP17 in H295R Adrenocortical Cells Depends on Complex Formation among p54[nrb]/NonO,

Protein-Associated Splicing Factor, and SF-1, a Complex That Also Participates in Repression of Transcription, *Endocrinology*, **2002**, *143(4)*, 1280-1290

31. Passon et al.: Structure of heterodimer of human NONO and paraspeckle protein component 1 and analysis of its role of subnuclear body formation, *PNAS*, **2012**, *109(13)*, 4846-4850

32. Zipprich et al.: Importance of the C-terminal domain of the human GW182 protein TNRC6C for translational repression, *RNA*, **2009**, *15(5)*, 781-793

33. Mattiroli et al.: RNF168 Ubiquitinates K13-15 on H2A/H2AX to Drive DNA Damage Signaling, *Cell*, **2012**, *150(6)*, 1182-1195

34. Walsh et al.: ESpritz: accurate and fast prediction of protein disorder, *Bioinformatics*, **2012**, *28(4)*, 503-509

35. Lancaster et al.: PLAAC: a web and command-line application to identify proteins with prion-like amino acid composition, *Bioinformatics*, **2014**, *30(17)*, 2501-2502

36. Süveges et al.: Charged single alpha-helix: a versatile protein structural motif, *Proteins*, **2009**, *74(4)*, 905-916

37. Barnes et al.: Remarkable Rigidity of the Single α-Helical Domain of Myosin-VI As Revealed by NMR Spectroscopy, *Journal of the American Chemical Society*, **2019**, *141(22)*, 9004-9017

38. Molliex et al.: Phase Separation by Low Complexity Domains Promotes Stress Granule Assembly and Drives Pathological Fibrillization, *Cell*, **2015**, *163*, 123-133

39. Liu et al.: Phase separation drives decision making in cell division, *Journal of Biological Chemistry*, **2020**, *295(39)*, 13419-13431

40. Feng et al.: Phase separation as a mechanism for assembling dynamic postsynaptic density signalling complexes, *Current Opinion in Neurobiology*, **2019**, *57*, 1-8

41. Mitrea et al: Phase separation in biology; functional organization of a higher order, *Cell Communication and Signalling*, **2016**, *14*, article number: 1

42. Mészáros et al.: PhaSePro: the database of proteins driving liquid-liquid phase separation, *Nucleic Acids Research*, **2020**, *48(D1)*, D360-D367

43. Dannenhoffer-Lafage et al.: A Data-Driven Hydrophobicity Scale for Predicting Liquid-Liquid Phase Separation of Proteins, *The Journal of Physical Chemistry B*, **2021**, *125(16)*, 4046-4056

44. Gerstberger et al.: A census of human RNA-binding proteins, *Nature Reviews Genetics*, **2014**, *15(12)*, 829-845

45. Zeng et al.: Reconstituted Postsynaptic Density as a Molecular Platform for Understanding Synapse Formation and Plasticity, *Cell*, **2018**, *174(5)*, 1172-1187

46. Milovanovic et al.: A liquid phase of synapsin and lipid vesicles, *Science*, **2018**, *361(6402)*, 604-607

47. Dobson et al.: A conserved charged single α-helix with a putative steric role in paraspeckle formation, *RNA*, **2015**, *21*, 2023-2029

48. Kovács et al.: Detection of single alpha-helices in large protein sequence sets using hardware acceleration, *Journal of Structural Biology*, **2018**, *204*, 109-116

49. Dudola et al.: Consensus prediction of charged single alpha-helices with CSAHserver, *Methods in Molecular Biology*, **2017**, *1484*, 25-34

50. Pak et al.: Sequence Determinants of Intracellular Phase Separation by Complex Coacervation of a Disordered Protein, *Molecular Cell*, **2016**, *63(1)*, 72-85

51. Chen et al.: Phase separation at the synapse, *Nature Neuroscience*, **2020**, *23*, 301-310