Spatio-temporal coordination at the maternal-fetal interface promotes trophoblast invasion and vascular remodeling in the first half of human pregnancy

Shirley Greenbaum[1,2,*,+], Inna Averbukh[1,+], Erin Soon[1,3,+], Gabrielle Rizzuto[4], Alex Baranski[1], Noah Greenwald[1,5], Marc Bosse[1], Eleni G. Jaswa[6], Zumana Khair[1], Shirley Kwok[1], Shiri Warshawsky[1], Geneva Miller[7], Morgan Schwartz[7], Will Graf[7], David Van Valen[7], Leeat Keren[8], Travis Hollmann[9], Matt van de Rijn[1], Michael Angelo[1, *]

[+]Equal contribution

[*]Correspondence: greenbaumsh@gmail.com (S.G.), mangelo0@stanford.edu (M.A.)

[1]Department of Pathology, Stanford University

[2]Department of Obstetrics & Gynecology, Hadassah-Hebrew University Medical Center

[3]Immunology Program, Stanford University

[4]Department of Pathology, UCSF

[5]Cancer Biology Program, Stanford University

[6]Department of Obstetrics, Gynecology & Reproductive Sciences, UCSF

[7]Division of Biology and Bioengineering, California Institute of Technology

[8]Department of Molecular Cell Biology, Weizmann Institute of Science

[9]Department of Pathology, Memorial Sloan Kettering Cancer Center

Abstract

Beginning in the first trimester, fetally derived extravillous trophoblasts (EVTs) invade the uterus and remodel its spiral arteries, transforming them into large, dilated blood vessels that lack smooth muscle and are partially lined with EVTs instead of vascular endothelium. Several mechanisms have been proposed to explain how EVTs coordinate with decidual cells to promote a tissue microenvironment conducive to spiral artery remodeling (SAR). However, it remains a matter of debate which immune and stromal cell types participate in these interactions, how this process evolves with respect to gestational age, and which anatomic routes are the predominate path of EVT invasion in humans. To elucidate this complex interplay, we used multiplexed ion beam imaging by time of flight with a 37-plex antibody panel to build the first spatio-temporal atlas of the human maternal-fetal interface in the first half of pregnancy at single-cell resolution. We analyzed ~500,000 cells and 588 spiral arteries within intact decidua from 66 patients between 6-20 weeks of gestation. Using custom machine learning algorithms for cell segmentation and classification, we evaluated the spatial distributions and phenotype of 20 maternal and five EVT populations with respect to gestational age and SAR. Gestational age substantially influenced the frequency of most maternal immune and stromal cells, with tolerogenic subsets expressing CD206, CD163, TIM-3, Galectin-9, and IDO-1 preferentially enriched at later time points. In contrast, SAR progression, and not gestational age, preferentially correlated with local invasion of EVTs. Lastly, by comparing spatial co-occurrence and phenotype of decidual interstitial, perivascular and intravascular EVTs with respect to SAR progression, we developed a statistical model suggesting an intravasation mechanism as the predominant route of EVT invasion in superficial decidua. Taken together, these results support a coordinated model of decidualization in which increasing gestational age drives a transition in maternal decidua towards a tolerogenic niche conducive to locally regulated, EVT-dependent SAR.

Introduction

The maternal-fetal interface is established when the trophoectoderm cells of the blastocyst invade the decidualizing endometrial stroma, ultimately forming the placenta. From that point forward, normal development depends on a complex interplay between maternal cells and placental trophoblasts that ultimately transforms the nascent womb into a specialized niche capable of meeting the metabolic demands of a growing hemi-allogeneic fetus while maintaining maternal tolerance[1–5]. Rather than being a single monotonic trend, this process is multifaceted and dynamic with respect to both tissue structure and gestational age (GA). Subsequent to implantation, decidual cellular composition shifts to one that is enriched for invasive extravillous trophoblasts (EVTs)[6]. During this transition, maternal and fetal cells remodel uterine spiral arteries into highly dilated vessels with minimal smooth muscle where EVTs have partially replaced the

maternal endothelium within the arterial lumen[7–9]. Spiral artery remodeling (SAR) in

healthy pregnancies results in low-resistance vessels that can deliver blood to the

intervillous space at low flow velocities that prevent damage to the placental

architecture[10,11]. Conversely, impaired SAR, fewer tolerogenic maternal cells, and

abnormal decidual invasion of EVTs have each been implicated in placenta-related

obstetric complications, including preeclampsia, intrauterine growth restriction, and

preterm birth[12,13]. Therefore, detailed investigation of the cell population dynamics at the

maternal-fetal interface is key to understanding the biology of normal pregnancy and the

pathophysiology of placenta-related obstetric complications.

Due to the poor feasibility of controlled studies in pregnant humans, much of what

is known about maternal-fetal tolerance and SAR is based on pregnancy in small

mammals[14]. Although some similarities exist, key facets of hemochorial placentation in

humans are primate-specific, and in some cases are restricted even further to great

apes[15–17]. For example, EVTs in mice only invade the superficial decidua, do not replace

the vascular endothelium, and are thought to play a minor role in SAR compared to

maternal uterine natural killer (NK) cells[18]. In contrast, EVTs in humans invade completely

through the decidua into the inner third of the myometrium and are considered to be vital

for adequate SAR[3,19,20]. Since the most extensive EVT invasion has been observed in

humans, it may be a key adaptation that permitted upright, bipedal locomotion while

maintaining adequate blood flow in the third trimester when development of the large fetal

brain accounts for 60% of metabolic needs [21,22].

The study of human decidual remodeling is further complicated by additional

inherent challenges. First, cell composition and structure are temporally dynamic;

aggregating data across different GAs or observing a single time point may be misleading.

As endometrial stromal cells shift towards a decidualized phenotype[23,24], the functions of

maternal NK cells, T cells, and macrophages change dynamically in the first and second

trimesters to promote a permissive niche conducive to villus attachment and invasion.

This process necessarily establishes a gradient of EVT invasion that advances inward

from the superficial decidua. Consequently, decidual structure and composition in focal

regions can differ significantly from its bulk attributes. A second major challenge arises in

understanding how these global dynamics are coupled to processes requiring spatial

coordination, such as those between maternal and placental cells in the local tissue
microenvironment. For example, periarterial decidual NK cells are thought to contribute
to SAR both by initiating smooth muscle breakdown and by secreting chemokines that
attract invading EVTs, while phagocytic macrophages are thought to facilitate clearance
of the resultant apoptotic debris[25–27]. Overall, formation of the human maternal-fetal
interface involves sophisticated spatiotemporal coordination such that tissue composition,
structure, and function are inextricably coupled. Unraveling this interdependence requires
an approach that can ascertain how these facets change over time in intact human tissue.

With this in mind, we constructed the first high dimensional spatio-temporal atlas
of the human maternal-fetal interface. We leveraged archival tissue banks to assemble a
cohort of maternal decidua from 66 women who underwent elective terminations of
otherwise healthy pregnancies at 6-20 weeks gestation. We then performed high
dimensional, subcellular imaging with multiplexed ion beam imaging by time of flight
(MIBI-TOF) [28] using a 37-plex antibody panel designed to comprehensively identify the
location, lineage, and function of all major maternal and placental cells. To understand
how SAR relates to local decidual composition, we developed new algorithms for
quantifying vascular morphology that enabled us to assign a remodeling score to each
individual artery. Comparison of these scores for 588 arteries revealed that the extent of
SAR to varies significantly with respect to GA. We then leveraged these discordances to
discern which changes in decidual composition and structure were preferentially driven
by GA, SAR, or both. Overall, the frequencies and relative proportions of maternal
immune cells exhibited a robust temporal dependence that permitted us to predict GA
based on these features alone. In contrast, we found that EVT invasion and perivascular
localization were the dominant drivers of SAR in the tissue microenvironment. Given
these findings, we then used our atlas to compare two models for the path of EVT
migration from the cytotrophoblast cell columns to maternal spiral arteries that have been
proposed previously: (1) intravasation, where EVTs first invade the decidua and then
enter arteries by traversing the arterial wall, and (2) extravasation, where EVTs enter
arteries directly at the basal plate[9]. Using statistical analyses correlating EVT phenotype
and location with the extent of arterial smooth muscle and endothelial loss, we found that
our spatiotemporal atlas was most consistent with an intravasation model. Taken

together, these investigations support a cooperative interplay in the first half of pregnancy
in which temporally dependent changes in decidual function permit placental EVTs to
extensively alter the maternal uterine vasculature.

Results

Multiplexed imaging of human decidua reveals the tolerogenic composition of the maternal-fetal interface

As part of the Human BioMolecular Atlas Program (HuBMAP) initiative, we created
the first spatio-temporal tissue atlas of the human maternal-fetal interface in the first 20
weeks of pregnancy (Fig. 1a). The goal of this study was to comprehensively define the
structure and composition of decidua and to understand how it changes during the first
two trimesters with respect to two axes: GA and maternal SAR. To examine these issues,
we assembled a retrospective cohort of archival formalin-fixed, paraffin-embedded
placenta and decidua tissue from 74 patients who underwent elective termination of
pregnancies with no known fetal abnormalities. Archival tissue blocks were manually
screened by a perinatal pathologist in hematoxylin and eosin (H&E) stained tissue
sections to determine which samples contained decidua. Then, regions of decidua that
contained spiral arteries were demarcated, cored, and assembled into two tissue
microarrays (TMAs) of 1mm and 1.5 mm cores. The final dataset included samples for 6-
20 weeks of gestation (13.72±4.8 weeks) from 66 patients of varying parity (1.45±1.72),
age (28.17±5.9 years), body mass index (28.19±7.3 kg/m^2), and ethnicity (Fig. 1b-f,
Supplementary Table 1). Due to inherent limitations in how the tissue was procured,
precise anatomic locations could not be determined. However, 61 out 66 tissue blocks
contained placental villi, suggesting that the vast majority of this cohort was sampled from
decidua basalis (Supplementary Table 2, See methods).

Previous studies of intact tissue examining only one or a few cell populations at a
time reported shifts in maternal immune cells towards tolerogenic states that are
permissive to invasion by fetally derived EVTs[19]. To gain a more complete picture of the
complex cell-cell interactions that establish maternal tolerance in the first half of
pregnancy, we designed and validated a 37-plex antibody panel for simultaneously

Figure 1 | Study design and workflow. a. Diagram of a human embryo in utero at 6 weeks gestation. First inset: the maternal-fetal interface consisting of decidua basalis (purple) with maternal spiral arteries (light pink) and fetal chorionic villi in the intervillous space (bottom right corner). Second inset: early stage (6 weeks) unremodeled spiral artery and progression to late stage (20 weeks) remodeled artery and anchoring fetal villi. **b.** Cohort parity distribution. **c.** Cohort distribution of body mass index (BMI). **d.** Cohort age distribution. **e.** Cohort ethnicity distribution. **f.** Cohort distribution of gestational age (GA). **g.** TMA construction, antibody panel, MIBI workflow, and single-cell segmentation.

mapping the functional state and location of all major maternal and fetal cell populations

(Fig. 1g, See methods, Extended Data Fig. S1). In addition to canonical lineage defining

markers for fetal cells, maternal immune cells, fibroblasts, smooth muscle, endothelium,

and epithelium, we also quantified 10 functional markers previously implicated in maternal

immune tolerance, including TIM-3, Galectin-9, PD-1, PDL-1, and IDO-1 (Fig. 1g,

Extended Data Fig. S1)[29–34] . TMA sections were stained simultaneously with this

antibody panel and subsequently imaged at 500 nm resolution using MIBI-TOF (Fig. 1g).

Multiplexed images were denoised with a low-level image analysis pipeline as

described previously (Fig. 1g)[35]. To accurately capture the unique diversity of

morphologically distinct maternal and fetal cells, we used our previously validated custom

whole cell convolutional neural network, Mesmer[36] (See methods, Extended Data Fig.

S2a). We optimized this neural network for decidua-specific segmentation by training with

93,000 manually annotated single cell events from 25 decidual images, 15 of them from

our cohort. Applying this segmentation algorithm to our cohort images yielded 495,349

segmented cells in total, identified across 211 images (800µmx800µm, 2347±783 cells

per image). FlowSOM clustering[37] was used to assign 92% of whole cell segmented

events to 25 cell populations (Fig. 2a, b, See methods, Extended Data Fig. S2b, c,

Supplementary Table 3). These data (Fig. 2c-g) were then combined with whole-cell

segmentation masks to generate cell phenotype maps (CPM) in which each cell is colored

categorically by its respective population (Fig. 2h, Extended Data Fig. S2d). We then

determined whether cells expressed the functional markers by applying statistically

derived per-marker binary expression thresholds (see Methods, Extended Data Fig. S2e,

Supplementary Table 4). Noteworthy histological features—such as arteries, vessels,

glands, the cell columns, and decidual tissue boundaries were manually annotated in

collaboration with a perinatal pathologist.

Non-immune maternal (structural) cells accounted for the majority (56.3%) of all

segmented events in the decidua and were predominantly composed of decidual

fibroblasts (60.5%) and myofibroblasts (24.8%) with smaller contributions from vascular

endothelium (7.6%) and glandular epithelial cells (7.1%, Fig. 2b, c). Maternal immune

cells (31% of all cells) were dominated by macrophages (47.6% of immune) and NK cells

(42.6% of immune) with minor contributions from T (8% of immune), dendritic (1.3% of

immune), and mast cells (0.5% of immune). We identified a total of five decidual

Figure 2 | Multiplexed imaging of human decidua reveals the immune tolerance-conducive composition of the maternal fetal interface. a. Cell lineage assignments showing mean normalized expression of lineage markers (left) and functional marker positive cell frequency (right, Z-score). Columns (markers) are hierarchically clustered. **b.** Cell lineage abundances across our cohort. Placental mac: Placental macrophage; MyoFib: Myofibroblast. **c.** Representative MIBI field of view color overlay of a 20 week sample. Red = VIM, vimentin, purple = HLA-G, yellow = CD56, cyan = CD31, grey = H3, green = Ki67. **d.** Inset of **c**, interstitial fetal EVTs. Blue = Galectin-9, purple = HLA-G, yellow = CD14, grey = H3, red = VIM **e.** Inset of **c**, showing anchoring villous cell column to decidua interface. Blue = CK7, cytokeratin7, purple = HLA-G, green = Ki67, grey = H3. **f.** Inset of **c**, showing intravascular EVTs. Blue = PD-L1, purple = HLA-G, yellow = CD56, grey = H3. **g.** Inset of **c**, showing decidual stromal cells (fibroblasts) and macrophages. Blue = CD163, cyan = CD206, yellow = CD11c, grey = H3, red = vimentin. **h.** Cell lineage assignments overlaid onto the cell-segmentation output to produce a cell phenotype map.

macrophage (CD14$^+$) populations, ubiquitously co-expressing CD163 and CD206,

consistent with an M2-polarized, tolerogenic phenotype[38] (Fig. 2g). In line with previous

work showing pregnancy-specific recruitment, 77% of macrophages expressed DC-

SIGN[39] (Fig. 2a). We further classified DC-SIGN$^+$ macrophages into three subsets

(Mac2a, 2b, 2c) based on expression of CD11c (Mac2b, 2.7% of macrophages) or

absence of HLA-DR (Mac2c, 10.3% of macrophages). The majority (64%) of

macrophages were CD11c$^-$HLA-DR$^+$ (Mac2a). Macrophages lacking DC-SIGN (23% of

macrophages) were further categorized based on CD68 expression (CD68$^-$ Mac1b and

CD68$^+$ Mac1a) (Fig. 2a, b).

Four subsets of NK cells (CD3$^-$CD56$^+$) were identified based on combinatorial co-

expression of CD57, CD11c, and CD8. NK1 (CD57$^-$CD16^{low}) were the largest NK cell

population present, making up 59.7% of NK cells (Fig. 2a, b). The remaining three subsets

could be distinguished based on expression of CD57 (NK2, 25.8% of NK cells), CD11c

(NK3, 11.3% of NK cells), or CD8 (NK4, 3.2% of NK cells) (Fig. 2a, b). T cells consisted

of CD8$^+$ (53.2% of T cells), CD56$^+$ NKTs (28.8% of T cells), CD4$^+$ (17.1% of T cells), and

sparse numbers of regulatory T (Treg) cells (CD4$^+$FOXP3$^+$, 0.7% of T cells); while no B

cells were observed (Fig. 2a, b). Fetal cells (12.6% of all cells) primarily comprised four

subsets of EVTs that were delineated based on combinatorial expression of HLA-G, CK7,

CD57, and CD56 (Fig. 2a). HLA-G$^+$ interstitial EVT populations were CK7$^+$ (EVT1a,

44.6% of fetal cells), CK7$^-$ (EVT1b, 35.3% of fetal cells), or CD56$^+$ (EVT1c, 6.9% of fetal

cells) (Fig. 2c-f). EVT2 lacked HLA-G and were CD57$^-$CK7^{low} (EVT2, 9.4% of fetal cells).

Notably, placental macrophages (Hofbauer cells) located in chorionic villi constituted the

remainder (4.1%) of fetal cells and exhibited a cellular phenotype similar to that of Mac2c

(DC-SIGN$^+$HLA-DR$^-$) decidual macrophages (Fig. 2a).

As previously reported, we detected IDO-1 expression in glandular cells[29], but also

in vascular endothelium, where it has been previously reported to be scarce (12.3% of

endothelial cells were IDO-1$^+$, Fig. 2a)[34]. Our analysis revealed numerous functional

subsets of maternal cells, including TIM-3$^+$Galectin-9$^+$, iNOS$^+$, and HO-1$^+$ subsets of DC-

SIGN$^+$ macrophages, Galectin-9$^+$ fibroblasts (36.7% Galectin-9$^+$, Fig. 2d), and an

intriguing TIM-3$^+$Lck$^+$ subset of Tregs that accounted for >50% of this population (Fig. 2a,

b). Interestingly, both Tregs and NKT cells were highly proliferative (13.7% Ki67$^+$ Tregs,

17% Ki67$^+$ NKT cells), and with the notable exception of CD8$^+$ NK cells (22.9% GrB$^+$),
had higher frequencies of GrB (Granzyme B) expressing cells than any NK cell subset
(33.7% GrB$^+$ NKT, 19.5% GrB$^+$ Treg). Our highly multiplexed imaging platform confirmed
prior findings[32,39–41] and enabled us to enumerate an ensemble of functional states across
multiple lineages that were collectively consistent with maintaining a tolerogenic niche.

SAR progression is tightly correlated with the local tissue microenvironment

Perfusion of the intervillous space by uterine spiral arteries is the sole source of
oxygen and nutrients to the growing fetus after the establishment of arterial flow. During
the first half of pregnancy, these vessels undergo an extensive remodeling process that
culminates in dilated, non-contractile vessels depleted of smooth muscle and where the
maternal endothelium has been replaced by EVTs. While abnormal SAR is associated
with obstetric complications, such as intrauterine growth restriction and preeclampsia[12,13],
it is still not fully understood which cell populations directly participate in SAR, how this
process is locally regulated, and to what extent these changes are synchronized with GA.

We therefore used our spatiotemporal atlas of decidua to construct a SAR
trajectory to reveal how this process relates to temporal changes in decidua cell
composition and structure. Using artery size, smooth muscle layer disruption and loss,
endothelial continuity, EVT infiltration, and EVT endothelization to determine the extent
of SAR, we manually assigned each artery to one of five sequential remodeling stages
based on previously published criteria[42] (Fig. 3a). To ensure scoring was not biased by
patient demographics, the score of neighboring arteries, or the composition of nearby
stroma, scoring was performed on cropped images of each artery independently by
blinded experts. Out of 588 arteries, 186 were unremodeled and assigned to Stage 1 (Fig.
3b, c). Stage 2 arteries (300 arteries) were characterized by moderate smooth muscle
disruption and endothelial swelling (Fig. 3d, e). Stage 3 arteries (43 arteries) exhibited
more dilation, smooth muscle loss, and early endothelial disruption (Fig. 3f, g).
Progression to Stage 4 (34 arteries) was marked by the presence of EVTs within the
arterial lumen (Fig. 3h,i), while fully remodeled Stage 5 arteries (25 arteries) were
identified based on their very large size, near-complete smooth muscle loss, and EVT
endothelization (Fig. 3j, k, see Methods, Extended Data Fig. S3a, Supplementary Table

5).

Although SAR correlated with GA to some extent (Spearman's ρ=0.28, p-value =

$1.5*10^{-12}$), in many cases artery staging and GA were discordant. For example, at least

one late-stage artery (Stage 4-5) was present in 40% of week 8 samples, while minimally

remodeled arteries were present throughout (Fig. 3l). Moreover, SAR staging of arteries

from the same patient often varied significantly between tissue cores (32% of patients had

arteries that differed by at least two stages), suggesting that this discordance could be

highlighting aspects of SAR that are locally regulated by the tissue microenvironment (Fig.

3l, Extended Data Fig. S3b).

This decoupling of SAR and GA permitted us to identify changes in decidual

composition that were predominantly driven by one or the other. We first developed a

quantitative staging scheme for assigning a continuous, quantitative, and accurate

remodeling score in an automated fashion. For each artery, we extracted 35 parameters

describing the same aspects of arterial morphology that were used for manual scoring

(Fig. 3m, see Methods, Extended Data Fig. S3d). Together with manual staging, we used

this quantitative morphologic profile to construct a highly resolved pseudotime trajectory

of SAR using linear discriminant analysis (LDA)[43] (Fig. 3m, n). We generated this

trajectory by combining the 35 morphological features with our manually defined stage

labels and applying LDA to project each artery with respect to a two-dimensional LDA

space in which separation of arteries by their manually assigned stages is optimal (see

Methods). This separation was mainly driven by artery shape and size, properties of the

smooth muscle layer and EVT presence (see Methods, Supplementary Table 6).

We then defined a remodeling trajectory as the polynomial fit to artery points in this

space and subsequently mapped each artery (a_i) to the nearest point along this curve (b_i,

Fig. 3n inset, see Methods). Finally, a remodeling score (δ) was determined by calculating

the distance along this curve from the point of origin (x_o) to b_i for each artery (See integral

in Fig. 3n, Extended Data Fig. S3e, f, Supplementary Table 5). With our continuous

remodeling score δ, we next defined a simple scheme to differentiate GA- and SAR-driven

trends by performing linear regressions of cell frequency per image both as a function of

GA and as a function of δ. Regression R^2 and p-values were used as proxies for trend

a

Stage 1 | Stage 2 | Stage 3 | Stage 4 | Stage 5

Immune cell
Endothelial cell
Smooth muscle cell
Fibroblast

Swollen endothelium & disrupted smooth muscle layer

Substantial loss of smooth muscle

Intravascular extravillous trophoblasts (EVTs)

Endothelialized EVTs line lumen of spiral artery

b 50 µm **d** 50 µm **f** 50 µm **h** 50 µm **j** 50 µm

SMA VIM HLA-G CD31 CD45 | SMA VIM HLA-G CD31 CD45 | SMA VIM HLA-G CD31 CD45 | SMA VIM HLA-G CD31 CD45 | SMA VIM HLA-G CD31 CD45

c **e** **g** **i** **k**

l

Frequency vs Gestational Age (Weeks)

Stage

m

SAR Manual Staging

Digitizing Morphological Features

Thickness of endothelial & smooth muscle layers, presence/absence of EVTs

Continuity of endothelial & smooth muscle layers, organization of EVTs present

LDA → Continuous SAR Score δ

δ = 3.15

n

SAR Manual Stage
1 2 3 4 5

LD2 vs LD1

x_0 $f(x)$ a_i b_i

$\delta_i = \int_{x_0}^{b_i} \sqrt{1 + f'(x)^2}\, dx$

o

$-\log_{10}$(p-value) vs $\log_2(\delta/GA)$

Trend Size
-2 -1.5 -1 -0.5 0 0.5 1 2 3

Key
Macrophage
NK cell
Structural
T cell
Mast cell
Other immune
Fetal

Cell Frequency / Out of Category

GA-driven
SAR (δ)-driven

Macrophages / Immune cells
NK cells / Immune cells
Fibroblasts / Structural
Myofibroblasts / Structural
Mac2a / Macrophages
NK3 / NK cells
CD8+ T / T cells
T cells / Immune cells
EVT1a / Fetal cells
EVT / All cells
Immune cells / All cells
NK2 / NK cells
EVT2 / Fetal cells
NK1 / NK cells
Structural / All cells
EVT1c / Fetal cells

Figure 3 | Spiral Artery Remodeling (SAR) progression significantly influences maternal-fetal interface composition. a. Diagram showing key characteristics of SAR Stages 1-5, assessed manually. **b.** Representative MIBI color overlay of SAR manual Stage 1 arteries. VIM, vimentin; SMA, smooth muscle actin. **c.** Inset of **b**, showing SAR manual Stage 1 arteries. **d.** Representative MIBI color overlay of SAR manual Stage 2 arteries. **e.** Inset of **d**, showing one SAR manual Stage 2 artery. Arrowhead; swollen endothelial cells. **f.** Representative MIBI color overlay of SAR manual Stage 3 arteries. **g.** Inset of **f**, showing one SAR manual Stage 3 artery. Arrowhead; substantial loss of smooth muscle **h.** Representative MIBI color overlay of one SAR manual Stage 4 artery. **i.** Inset of **h**, showing one SAR manual Stage 4 artery. Arrowheads; intravascular EVTs. **j.** Representative MIBI color overlay of one SAR manual Stage 5 artery. **k.** Inset of **j**, showing one SAR manual Stage 5 artery. Arrowhead; endothelialized intravascular EVTs lining the spiral artery lumen. **l.** Distribution of SAR manual stages by gestational age (GA). **m.** Schematic of calculating the continuous SAR remodeling score (δ). Manual stages along with quantified digitized morphological features were used to construct a trajectory of SAR using LDA from which the continuous SAR score δ was calculated. **n.** Scatter of arteries in LDA space color coded by manually assigned stage. The polynomial fit depicts the remodeling trajectory. Inset: matching each artery point a_i to the SAR trajectory by finding the nearest point along trajectory b_i. The continuous SAR score δ was then defined as the distance from origin x_0 to b_i along the trajectory curve. **o.** Volcano plot distinguishing GA-driven from SAR (δ)-driven cell-type frequencies. X axis: $\log_2$ ratio of R^2 derived from linear regression against SAR (δ) and GA. Y axis: -$\log_{10}$ of the p-value for the better-fitting regression model. Points are color coded by the trend size observed in the better-fitting regression model.

quality and to assess significance, respectively. Trends where R^2 for GA and SAR differed

by at least two-fold were classified as being driven predominately by a single process,

while ones falling below this cutoff were classified as synchronized (Fig. 3o, see Methods,

Extended Data Fig. S3g, h, Supplementary Table 7).

Consistent with previous studies reporting fewer interstitial EVTs in pregnancy

complications that were related to impaired SAR, EVT decidual presence was better

correlated with SAR than with GA ($\log_2 R^2$ ratio(δ:GA) = 2.5, p-value for δ =1e^{-13}). This

increase in decidual EVTs with SAR was accompanied by a decrease in immune and

structural cell frequencies (Fig. 3o). Within the EVT compartment, the frequency of EVT1c

out of fetal cells exhibited the greatest bias for SAR ($\log_2 R^2$ ratio(δ:GA) = 10.3, p-value

for δ = 9e^{-4}, increase of 2.8 times the mean value, Fig. 3o), further highlighting this

dependence. Changes in the relative proportion of cell types within the immune

compartment were mostly driven by GA (Fig. 3o). Notably, the DC-SIGN$^+$ Mac2a subset

which was previously reported to be pregnancy-specific[39] was heavily GA-biased ($\log_2$

R^2 ratio(δ:GA) = -5.8, p-value for GA = 3e^{-8}, Fig. 3o), suggesting that this population is

recruited in a manner that is agnostic to SAR. In contrast, the trade-offs within the NK cell

compartment where the relative proportion of NK2 (CD57$^+$) diminished as NK1 (CD57$^-$)

increased were SAR-dependent ($\log_2 R^2$ ratio(δ:GA) $\geq$ 1.5, p-value for δ $\leq$ 0.003, Fig.

3o). Interestingly, CD57 expression in human NKs results in a highly cytotoxic phenotype
and has been shown to play a pivotal role in cancer immunosurveillance[44]. These trends
suggest that during pregnancy, the reduction in cytotoxic mature NKs promotes a
permissive niche where EVTs can invade and initiate SAR.

A lymphoid to myeloid shift in immune compartment composition is tightly
correlated with GA

299   Our analyses indicated a robust, GA dependent shift from a lymphoid to myeloid
300   dominated immune landscape, characterized by fewer NK and T cells and a concomitant

Figure 4 | A lymphoid-to-myeloid shift in immune-compartment composition is tightly correlated with gestational age (GA). a. Frequency of EVT, immune, and structural cell populations per patient, with patients ordered by GA. **b.** Frequency of immune cell populations per patient, by GA. MAC, macrophage; NKT, NK T cells; NK, total NK cells. **c.** Representative cell phenotype map of immune composition in decidual tissue in an early (6 weeks GA) sample. Green = NK, blue = macrophage, pink = T cell, purple = EVT, grey = other. **d.** Inset of **c**, showing a MIBI color overlay of NK cells with GrB expression. Grey = H3, yellow = CD56, blue = CD11c, red = GrB. **e.** Inset of **c**, showing a MIBI color overlay of T cells with PD-1 and Lck expression. Grey = H3, pink = CD3, orange = Lck, blue = PD-1. **f.** Representative cell phenotype map of immune composition in decidual tissue in a late (16 weeks GA) sample. Green = NK, blue = macrophage, pink = T cell, purple = EVT, grey = other. **g.** Inset of **f**, showing a MIBI color overlay of EVTs (1a, 1b) with PD-L1 expression. Grey = H3, purple = HLA-G, yellow = CD56, green = PD-L1. **h.** Inset of **f**, showing a MIBI color overlay of macrophages with TIM-3 expression. Grey = H3, green = DC-SIGN, blue = CD163, red = TIM-3. **i.** Predicted versus actual GA in days for a ridge regression model trained on GA-associated immune features, for a withheld test set (30%). Shaded region; 1 standard deviation. **j.** Ridge regression model coefficient loadings for GA-associated immune features.

increase in macrophage frequency (Log$_2$ R^2 ratio(δ:GA) ≤ -1.2, p-value for δ ≤ 1.2e^{-8}, Figs.

3o, 4a, b). In MIBI images at weeks 6-8 (Fig. 4c, e), show NK cells and T cells, including

those exhibiting cytotoxic (Fig. 4d) and immunosuppressive (Fig. 4e) phenotypes, greatly

outnumbered macrophages of all subsets (Fig. 4b, c). Contrastingly, images from weeks

16-20 were dominated by interstitial EVTs (Figs. 4a, 4f-g) and an accompanying increase

in tolerogenic macrophage populations (Fig. 4h) in relation to NK and T cells. To further

evaluate this relationship, we asked whether immune cell composition in the decidua

alone could be used to predict GA. Selecting immune features that were found to be

preferentially associated with GA rather than SAR (Fig. 3o), we trained and validated a

ridge regression model on a per-image basis using a random 70/30 test-train split

(Extended Data Fig. S4a). Remarkably, the trained model predicted GA in the withheld

test set within 19 days of the ground-truth value (R^2=0.7, Fig. 4i). On inspecting the model

weights, we found that the relative contribution of decidual immune cells was consistent

with the observed shift in the proportion of myeloid and lymphoid cells. In particular, the

relative frequencies of T and NK cells were negatively correlated with GA, while total

macrophage frequency was positively correlated with GA (Fig. 4j). Notably, a modified

regression model for predicting SAR (δ) based on the same immune cell population

parameters performed poorly (R^2=0.05, RMSE=0.85, Extended Data Fig. S4b),

reinforcing our hypothesis that these immune correlates are driven by GA and not SAR.

Coordinated up-regulation of tolerogenic functional markers with GA

Having examined the influence of GA and SAR in driving changes in the frequency

Figure 5 | Coordinated up-regulation of tolerogenic functional markers with gestational age (GA). a. Volcano plot distinguishing GA-driven from SAR (δ)-driven cell type-specific functional marker positivity fraction. X axis: $\log_2$ ratio of trend size is a relative measurement of R^2 derived from linear regression against GA or SAR (δ) and GA. Y axis: $-\log_{10}$ of the p-value for the better-fitting regression model. Points are color coded by the trend size observed in the better-fitting regression model **b.** Heatmap of changes in a subset of GA-driven functional markers as a function of GA in weeks. **c.** MIBI color overlay of IDO-1 expression in glandular cells (top inset) and endothelial cells (bottom inset) in an early (6 weeks GA) sample. Grey = H3, blue = VIM (vimentin), peach = SMA (smooth muscle actin), cyan = CD31, red = CK7, yellow = IDO-1. **d.** MIBI color overlay of IDO-1 expression in endothelial cells (inset) in spiral artery (SAR manual Stage 4) of a late (16 weeks GA) sample. Grey = H3, blue = vimentin, peach = SMA, cyan = CD31, magenta = HLA-G, yellow = IDO-1. **e.** Per-image Mac2a and Mac2b TIM-3$^+$ cell frequency versus Mac2a and Mac2b galectin-9$^+$ frequency, colored by GA. **f.** Cell phenotype map of macrophages and decidual fibroblasts. **g.** Inset of **f**; MIBI color overlay of TIM-3$^+$ and galectin-9$^+$ Mac2b and fibroblast cells. **h.** Inset of **f**; MIBI color overlay of TIM-3$^+$ and galectin-9$^+$ Mac1a, Mac1b, Mac2a, and fibroblast cells. Grey = H3, red = vimentin, pink = DC-SIGN, green = CD68, yellow = TIM-3, turquoise = Galectin-9.

of cell populations in the decidua, we next employed a similar approach to understand

how these two time axes correlate with shifts in decidual function. Using the same method

as our analysis of cell frequencies, we classified the temporal dynamics of functional

markers expression as GA-driven, SAR-driven, or synchronized (comparably correlated

with both GA and SAR) (Fig. 3o). Out of the 48 cell population-functional marker

combinations that were evaluated, 16 exhibited functional marker expression that

significantly correlated with one or both of these axes (Fig. 5a, see Methods,

Supplementary Table 8). These data revealed three overarching trends. First, both SAR

and GA are associated with dynamic changes in IDO-1 expression. For example, we
identified a GA-driven decline in IDO-1$^+$ glandular cells (Log$_2$ R^2 ratio(δ:GA) = -1.8, p-
value for GA = 2.3e^{-5}), a SAR-driven decline in IDO-1$^+$ dendritic cells (Log$_2$ R^2 ratio(δ:GA)
= 4.4, p-value for δ = 3e^{-3}), and an increase in IDO-1$^+$ vascular endothelium (p-value =
4e^{-4} ,Fig. 5c, d) that was comparably correlated with both GA and SAR (Fig. 5b, d).
Second, consistent with the cell frequency analysis (Fig. 3o) in which NK1 exhibited a
frequency increase preferentially associated with SAR, NK1 also exhibited a SAR
dependent increase in Ki67$^+$ frequency (Log$_2$ R^2 ratio(δ:GA) = 4.5, p-value for δ = 2e^{-4})
becoming more proliferative as arterial remodeling progresses (Fig. 5a). Third, functional
shifts in innate immunity were preferentially correlated with GA. All five macrophage
populations upregulated either TIM-3 and/or its cognate ligand Galectin-9 with GA (Fig.
5a, b). This trend was most prominent in the Mac2a and Mac2b populations, where a
tightly correlated up-regulation of both TIM-3 and Galectin-9 occurred (Fig. 5e-h,
Extended Data Fig. S5a). Interestingly, Galectin-9 upregulation was also detected in
fibroblasts (Fig. 5b, f-h). Notably, TIM-3 and Galectin-9 have been implicated in
suppressing anti-tumor surveillance by impairing the activity of cytotoxic NK and T cells
in various human cancers[30,45–47]. Taken with the SAR-driven decline in the proportion of
the cytotoxic NK2 observed here (Fig. 3o), these findings may suggest that TIM-3$^+$
Galectin-9$^+$ macrophages are serving a similar tolerogenic role in decidua.

Spatio-temporal EVT distribution suggests that intravasation is the predominant
route of EVT invasion in superficial decidua

Although intravascular EVTs are known to originate from the cytotrophoblast cell
columns, their path of migration remains a subject of debate primarily revolving around
two models: intravasation and extravasation (Fig. 6a). In the intravasation model, EVTs
detach from the cell columns and migrate through the decidua to first localize around the
spiral arteries. These perivascular EVTs then enter spiral arteries by migrating through
the arterial wall. In contrast, in the extravasation model, EVTs do not traverse the arterial
wall from within the decidua. Instead, detaching EVTs migrate retrograde against arterial
blood flow after entering at the basal plate where spiral arteries empty and merge into the
intervillous space[9].

a
Anchoring villous
Intervillous space
Intravasation
Extravasation

b
Interstitial
50 px
Intravascular
Perivascular
Cell column

c Cell column
100 μm
HLA-G CK7 CD57 H3 VIM PD-L1 Ki67

d Perivascular
50 μm
HLA-G CK7 CD56 H3 VIM CD31 SMA

e Intravascular
50 μm
HLA-G CK7 CD56 VIM CD31 SMA

f
% arteries ≤ δ, with EVT
Perivascular EVT
Intravascular EVT
SAR (δ)

g
% arteries ≤ s, with EVT
Perivascular EVT
Intravascular EVT
SMA loss (s)

h
% arteries ≤ e, with EVT
Perivascular EVT
Intravascular EVT
Endothelial loss (e)

i
HLA-G⁺
EVT1a EVT1b EVT1c EVT2
Frequency
Cell Column Interstitial Perivascular Intravascular

j
Kruskal Wallis, p = 1.1492e-19
Distance from artery (pixels)
EVT1c EVT1a EVT1b EVT2
HLA-G⁺

k
Z-score
-1 0 1
Ki67⁺ cell frequency
CD57 expression
HLA-G expression
PD-L1⁺ cell frequency
CD56 expression
Cell Column Interstitial Perivascular Intravascular

Figure 6 | Spatiotemporal EVT distributions suggest that intravasation is the predominant route of EVT invasion in superficial decidua. a. Two hypotheses for intravascular EVT invasion. (Left) Intravasation: orange arrows indicate movements of EVTs from the cell column of the anchoring villi into the decidua and through the wall of the artery and into the lumen. (Right) Extravasation: red arrows indicate movement of EVTs from the fetal villi through the intervillous space into the artery. **b.** Anatomical locations of interstitial, intravascular, perivascular, and cell column EVT populations in the decidua. **c.** MIBI overlay of anchoring villous and associated cell column EVT populations. Inset: cell column EVTs. Purple = HLA-G, yellow = CK7, green = CD57, grey = H3, red = VIM (vimentin), cyan = PD-L1, orange = Ki67 **d.** MIBI overlay of spiral arteries and associated perivascular EVT populations. Inset: perivascular EVT breaching artery wall. Purple = HLA-G, blue = CK7, yellow = CD56, grey = H3, red = VIM, green = CD31, orange = SMA (smooth muscle actin). **e.** MIBI overlay of remodeled spiral arteries and associated intravascular EVT populations. Inset; intravascular EVTs in a clump. Purple = HLA-G, blue = CK7, yellow = CD56, grey = H3, red = VIM, green = CD31, orange = SMA. **f.** Percentage of arteries with scores less than or equal to a given SAR (δ) threshold, by perivascular or intravascular EVTs present. **g.** Percentage of arteries with scores less than or equal to a given SMA loss (s) threshold, by perivascular or intravascular EVTs present. **h.** Percentage of arteries with scores less than or equal to a given endothelial loss (e) threshold, by perivascular or intravascular EVTs present. **i.** Frequency of EVT populations by anatomical location. **j.** Violin plot of distance from artery (in pixels) of EVTs grouped by EVT type. **k.** Heatmap of lineage and functional marker trends of EVT populations by anatomical location. Lineage marker (CD57, HLA-G, CD56) trends are mean expression values of EVT populations. Functional marker (Ki67, PD-L1) trends are the mean positive cell frequencies of EVT populations. Rows are z-scored and hierarchically clustered.

To determine which model best explains arterial invasion, we used our spatio-

temporal atlas to quantify how the phenotype and spatial distribution of EVTs evolve with

respect to SAR.   First, we manually defined feature masks demarcating cell column

anchoring villi and three decidual compartments—interstitial, perivascular, and

intravascular in our images (Fig. 6b) —to quantify EVT frequency in each (Fig. 6c-e).

Together with our SAR temporal trajectory, we used these data to ask a question that has

been qualitatively explored in previous work[20]: Where do EVTs accumulate first—in the

perivascular compartment (directly proximal to arteries) or within the intravascular

compartment? We quantified peri- and intravascular EVTs on a per-artery basis with

respect to their remodeling score δ and found that perivascular EVTs began accumulating

around less remodeled arteries in the decidua (Fig. 6f) and were consistently present at

earlier remodeling stages than intravascular EVTs (median δ = 2.2 vs. 3.2, Kruskal-Wallis

p-value = $5e^{-8}$, Extended Data Fig. S6a).   For arteries in which both were present, the

Log$_2$ ratio of EVTs present in these two compartments followed a continuous and smooth

trend as remodeling progressed, with intravascular EVTs increasing at the expense of

perivascular EVTs (R^2 = 0.5, p-value = $9e^{-12}$, Extended Data Fig. S6b). For a small

number of arteries, we observed perivascular EVTs breaching the artery wall, suggesting

they are in the process of invading the arterial lumen (Fig. 6d). These data are more

consistent with the intravasation model in which perivascular EVT are necessary before
intravascular EVT could appear.

Loss of smooth muscle and endothelium have defining roles in determining the
extent of SAR. Using morphometrics to quantify the extent of these concentric layers of
the arterial wall (see Methods), we examined how their integrity relates to EVT enrichment
in the perivascular and intravascular compartments. Similar to the trend seen with respect
to remodeling score δ, accumulation of perivascular EVTs was consistently present
around arteries at an earlier stage, with intravascular EVTs only appearing after 80%
smooth muscle loss (median smooth muscle loss for arteries with at least five
intravascular EVT present: 98%, Fig. 6g). Perivascular EVTs were present around
arteries irrespective of the degree of endothelium loss, while intravascular EVTs
increased with endothelium loss (Fig. 6h, linear regression on Log transformed
intravascular EVT as a function of endothelium loss: R^2=0.13, p-value =3e^{-4}) indicating
that endothelial disruption is a precursor to EVT entry into the arterial lumen. This
conclusion further supports the intravasation model, in which EVTs must transverse the
endothelial barrier to enter the arterial lumen.

Taken together, these data are consistent with a sequential process in which EVTs
detach from the cytotrophoblast cell columns and migrate through the decidua as
interstitial EVTs in order to accumulate in the perivascular compartment prior to
intravasation, as suggested previously[20]. To further evaluate this model, we posed the
following questions: Does EVT phenotype shift in a progressive manner that is consistent
with this stepwise intravasation model? If so, do intravascular EVTs more closely
resemble the cell column or perivascular compartment?

To answer these questions, we first compared the frequencies of the four EVT cell
populations within cell column, interstitial, perivascular, and intravascular masks. The
composition of each of these compartments shifted in a systematic manner along the
proposed path of migration: the cell columns consisted primarily of EVT1a, EVT1b and
EVT2 subsets with few CD56$^+$ EVT1c cells (99% vs. 1%). In the interstitial compartment,
the frequency of EVT2 dropped by 11-fold while the frequency of EVT1c cells increased
modestly to 2.4%. EVT1c cells were further enriched within the perivascular compartment

(7.3%) but were most prevalent in the intravascular compartment (54%, Fig. 6i). In
addition, EVT1c cells were found significantly closer to arteries than EVT2 cells (Kruskal
Wallis p-value = $1.15e^{-19}$, Fig. 6j).

Comparison of functional marker expression across all subsets within each
masked compartment, again revealed a progressive shift in EVT phenotype that best
aligned with a route of invasion consistent with an intravasation model.   Cell column
compartments were uniquely enriched for proliferative (Ki67$^+$), CD57$^+$ EVTs (Fig. 6k).
With decidual EVT invasion, a precipitous drop in CD57 and Ki67 expression was
accompanied by a progressive increase in PD-L1 that peaked in the intravascular
compartment (Fig. 6k). Notably, the intravascular compartment most closely resembled
the perivascular compartment in terms of functional marker expression (Fig. 6k,
perivascular 9.4% closer than interstitial to intravascular, Kruskal-Wallis p-value = $8e^{-7}$,
see Methods, Extended Data Fig. S6c). While the perivascular compartment is the most
similar to the intravascular compartment, a noticeable difference between the two
compartments was driven by PD-L1 and CD56 expression levels (Fig. 6k). This difference
stems from the highest prevalence of the CD56$^+$ PDL-1$^+$ EVT1c cells in the intravascular
compartment, which further increases with SAR (Fig. 6i, Extended Data Fig. S6d-f).

One potential explanation for the steep increase in EVT1c prevalence between the
perivascular and intravascular compartment is that arterial intravasation of perivascular
EVTs is accompanied by upregulation of CD56, such that EVT1a-b subsets would
effectively become the EVT1c subset. We therefore hypothesized that such a process
would involve an intermediate state between the two in which the EVT1a-b subsets
moderately express of CD56 en-route to the high expression observed in the EVT1c
subset. To test this hypothesis, we compared the average CD56 intensity of perivascular
and intravascular EVT1a-b EVTs on a per-artery basis (for arteries that initiated
remodeling: δ≥2). This analysis detected a statistically significant increase in CD56
expression between the perivascular and intravascular compartment by EVT1a-b subsets
(sided Wilcoxon signed rank test p-value = $5e^{-3}$, Extended Data Fig. S6g). An alternative
explanation for the disproportionate enrichment of EVT1c within vessels is that they are
more proliferative. However, only 0.5% of intravascular EVT1c were Ki67$^+$ compared to
9.6% and 1.8% of intravascular EVT1a and EVT1b cells, respectively (Fig. 6k, Extended

Data Fig. S6h).

Note that given the observational nature of our spatiotemporal atlas, neither model
can be definitively ruled in or out. However, taken together these analyses best align with
an intravasation model in which decidual invasion of cell column EVTs is accompanied
by pronounced downregulation of CD57 and Ki67 and upregulation of HLA-G.
Perivascular accumulation of EVTs occurs early in SAR, preceding the appearance of
intravascular EVTs and any loss in endothelium. In this model, as the endothelial barrier
is lost, perivascular EVTs invade the artery lumen and upregulate CD56.

Discussion

Decidualization is a fascinating process with no other normative precedent in
human biology, where the structure and function of the maternal endometrium transforms
to promote invasion of actively dividing, genetically dissimilar placental cells. Many
aspects of this process are primate-specific and some, such as deep arterial invasion of
EVTs into the myometrium, are thought to be largely restricted to humans with some
evidence in great apes[14–17,48]. Given this lack of tractable and relevant animal models
and the inability to study decidualization prospectively, our understanding of this process
is immature relative to other areas of human physiology. With this in mind, we used MIBI-
TOF and archival human tissue to generate the first spatiotemporal atlas of the maternal-
fetal interface during 6-20 weeks gestation. The central focus of our study was to
understand how global, temporally dependent changes in decidual composition are
coupled to local regulation of vascular remodeling in pregnancy. While initial invasion of
placental EVTs is prompted by a shift towards a permissive milieu, progression of SAR is
dependent on subsequent migration and perivascular accumulation of EVTs, where they
are thought to participate in cooperative cell-cell interactions with maternal fibroblasts, NK
cells, and macrophages[4,5]. Thus, formation of the maternal-fetal interface is mediated by
global, temporally dependent queues that serve as a gating function for remodeling
processes that are regulated in the local tissue microenvironment.

With this paradigm in mind, we set out to delineate which aspects of the first half
of pregnancy are driven globally by GA and how this relates to SAR. To achieve this, we

mapped the spatial distribution, composition, and functional state of ~500,000 maternal
cells and fetal EVTs with respect to glands, anchoring cell column villi, and spiral arteries
in >200 images from 66 patients.   Using LDA, image morphometrics, and expert
annotations, we assigned quantitative remodeling scores to every spiral artery in these
images. We then examined how cell frequency and function changed with respect to GA
and SAR. Our analysis of these changes determined GA to be the predominant driver of
maternal immune cell composition (Figs. 30, 4i, j). Progressive decreases in NK and T
cells drive a transition at 12-14 weeks GA from a lymphoid to myeloid predominant
decidua enriched for iNOS[+] NK cells, IDO-1[+] vascular endothelium, and DC-SIGN[+]
macrophages that co-express TIM-3 and Galectin-9 (Figs. 4b, 5a, b).   Notably, this
relationship between immune composition and GA was robust enough to allow us to
predict GA within 19 days based exclusively on immune population frequencies (Fig. 4i).

In contrast, all EVT subsets and only two maternal cell populations (NK1 and NK2)
preferentially correlated with progression of SAR. Higher remodeling scores were
correlated with more EVTs, more NK1s, and fewer NK2s. NK1 and NK2 primarily differ in
that the latter express CD57—a marker associated with a cytotoxic phenotype. Higher
proportions of presumptively more reactive NK2s early in SAR aligns well with previous
studies that have suggested that decidual NKs initiate early disruption of arterial smooth
muscle through secretion of GrB, MMP2, and MMP9[26,49]. Likewise, the proportional gains
seen here as SAR progresses of less reactive NK1s and invasive EVTs are consistent
with the tolerizing effects of HLA-G, which has been shown previously to decrease NK
cell cytotoxicity and induce production of IL-6 and IL-8 via binding of HLA-G to KIR2DL4,
LILRB1, and LILRB2[50,51]. Taken together, these data suggest that maternal and fetal cells
play cooperative, interdependent roles with SAR transitioning through NK- and EVT-
dependent phases.

We also examined a long-standing open question in the field: What is the path of
migration taken by EVTs that invade spiral arteries? On comparing cellular compositions
within cytotrophoblast cell columns of anchoring villi, decidua, and arteries, we found that
the local EVT frequency and phenotype within these regions shifted in a sequential,
coordinated manner consistent with an intravasation model in which EVTs within the
decidua enter spiral arteries through the arterial wall. Given the observational nature of

this study, we note that we cannot definitively rule out an extravasation model in which
EVTs migrate retrograde after entering spiral arteries directly at the basal plate. With this
limitation in mind, in our model EVTs detaching from proliferative cytotrophoblast cell
columns first invade the decidua and transition to a CD57- CK7$^+$ HLA-G$^+$ phenotype in
our proposed model. In line with previous work demonstrating EVT expression of MMP2
and MMP9[52], these cells migrate through the decidua and accumulate around spiral
arteries where they participate in removal of arterial smooth muscle. As this layer is
depleted, perivascular EVTs disrupt the underlying vascular endothelium and invade the
arterial lumen where they form multicellular clumps. Intravascular invasion is
accompanied by EVT upregulation of CD56, a homophilic binding molecule that has been
suggested to be necessary for heterotypic cell adhesion to endothelial cells[53]. Finally,
these multicellular clumps in fully remodeled arteries recede and are partially replaced by
trans-differentiated, endothelialized EVTs that have displaced the maternal endothelium.

Formation of the maternal-fetal interface is an organized and controlled invasive
process that is sometimes viewed as a template for understanding invasive and
immunosuppressive properties of tumors[54]. Both processes involve a genetically
dissimilar invasive cell type (haploidentical EVTs vs. clonal, mutated cancer cells),
extracellular matrix remodeling, and recruitment of a wide variety of tolerogenic immune
cells, including M2 polarized macrophages and proliferating Tregs. The intersection of
anchoring placental villi and maternal decidua morphologically resembles the invasive
margin of carcinomas and contains trophoblast cells expressing high levels of
immunomodulatory proteins and growth factors implicated in tumor severity including PD-
L1, IDO-1, TIM3, Her2, and EGFR [30,45,55,56]. In addition to these phenotypic and structural
similarities, recent work revealing mosaicism and clonal mutations in normal term
placentas demonstrate that this phenotypic overlap is even manifest at a genomic level[57].

Overall, we anticipate that this spatio-temporal atlas of the early human maternal-
fetal interface will provide a normative framework for elucidating etiological perturbations
in maternal-fetal tolerance and SAR in pregnancy complications. Likewise, this work may
also serve as a template for understanding how immune tolerance, tissue remodeling,
and angiogenesis are aberrantly recruited and synergized during tumor progression. With
this in mind, we plan in future studies to extend this comparative approach to archival

tissue from patients with preeclampsia, placenta accreta, and choriocarcinoma to further
elucidate cellular interactions involved in regulating SAR and EVT invasion.

Methods

Retrospective cohort design

The study cohort comprised decidua tissue from archival formalin-fixed, paraffin
embedded (FFPE) blocks, sampled after elective pregnancy terminations at the Women
Options Center at Zuckerberg San Francisco General Hospital, an outpatient clinic
located within a large public hospital affiliated with an academic medical center. Patients
at this clinic reflect a diverse population. The clinic serves women in the Bay Area as well
as referrals from California and out of state. While the patient population is predominantly
low-income mainly Medi-Cal patients, women of all economic backgrounds are cared for
at the clinic.

In the clinic, an ultrasound examination is performed to estimate GA, and medical
history is taken and logged as Electronic Medical Records ('eCW' - electronic clinical
works) or handwritten forms. A board-certified gynecologist reviewed medical records and
specifically extracted the following details: age, ethnicity, body mass index, gravidity,
parity, prior terminations, smoking, medications, HIV status, history of preeclampsia,
chronic hypertension, diabetes mellitus, renal disease, autoimmune disease, multifetal
pregnancy, and congenital anomalies (Supplementary Table 1). For procedures
occurring at less <14 weeks GA, suction aspiration is routinely used. For procedures at
>14 weeks GA, a combination of suction aspiration and grasping forceps is used. After
the procedure, tissue samples are routinely sent to pathology.

TMA construction

Regions of decidua with maternal spiral arteries, were cored and combined in two
TMA blocks by an experienced technician. Where possible, blocks containing the decidua
basalis were selected. Information on the histological characteristics of the blocks
retrieved, including the presence of cell column anchoring villi, is located in
Supplementary Table 2. The first TMA consisted of 205 cores (including three tonsil
cores, one endometrium core and one myometrium core) of 1 mm in diameter; the second

contained 86 cores of 1.5 mm in diameter. High resolution scans of each core were
uploaded to the Stanford Tissue Microarray Database (URL: http://tma.im/cgi-
bin/home.pl), a collaborative internal platform for designing, viewing, scoring, and
analyzing TMAs. Sequential recuts of the main experiment were stained with H&E, to aid
in choosing the imaging regions of interest (ROIs) and analyzing data.

Antibody preparation

Antibody staining was validated as described previously[28,58–60] . Briefly, each
reagent was first tested using single plex chromogenic IHC using multiple positive and
negative FFPE tissue controls prior to metal conjugation. Antibodies were then
conjugated to isotopic metal reporters as described previously[28,58–61] with the exception
of biotin-conjugated anti-PD-L1, for which a metal-conjugated secondary antibody was
used. Performance of metal conjugated antibody reagents were then tested within the
complete MIBI-TOF staining panel, under conditions identical to those in the main study
and compared with representative single plex chromogenic IHC to confirm equivalent
performance. Representative stains and information for each marker can be found in
Extended Data Fig. 1a and Supplementary Table 9 respectively. After conjugation,
antibodies were diluted in Candor PBS Antibody Stabilization solution (Candor
Bioscience). Antibodies were either stored at 4°C or lyophilized in 100 mM D-(+)-
Trehalose dehydrate (Sigma Aldrich) with ultrapure distilled H_2O for storage at −20°C.
Before staining, lyophilized antibodies were reconstituted in a buffer of Tris (Thermo
Fisher Scientific), sodium azide (Sigma Aldrich), ultrapure water (Thermo Fisher
Scientific), and antibody stabilizer (Candor Bioscience) to a concentration of 0.05 mg/mL.
Information on the antibodies, metal reporters, and staining concentrations is located in
Supplementary Table 9.

Tissue staining

Tissues were sectioned (4 μm in thickness) from tissue blocks on gold and
tantalum-sputtered microscope slides. Slides were baked at 70°C for 20 minutes followed
by deparaffinization and rehydration with washes in xylene (3x), 100% ethanol (2x), 95%
ethanol (2x), 80% ethanol (1x), 70% ethanol (1x), and ddH$_2$O with a Leica ST4020 Linear
Stainer (Leica Biosystems). Tissues next underwent antigen retrieval was carried out by

submerging sides in 3-in-1 Target Retrieval Solution (pH 9, DAKO Agilent) and incubating

them at 97°C for 40 minutes in a Lab Vision PT Module (Thermo Fisher Scientific). After

cooling to room temperature slides were washed in 1x PBS IHC Washer Buffer with

Tween 20 (Cell Marque) with 0.1% (w/v) bovine serum albumin (Thermo Fisher). Next,

all tissues underwent two rounds of blocking, the first to block endogenous biotin and

avidin with an Avidin/Biotin Blocking Kit (Biolegend). Tissues were then washed with wash

buffer and blocked for 1 hour at room temperature with 1x TBS IHC Wash Buffer with

Tween 20 with 3% (v/v) normal donkey serum (Sigma-Aldrich), 0.1% (v/v) cold fish skin

gelatin (Sigma Aldrich), 0.1% (v/v) Triton X-100, and 0.05% (v/v) Sodium Azide. The first

antibody cocktail was prepared in 1x TBS IHC Wash Buffer with Tween 20 with 3% (v/v)

normal donkey serum (Sigma-Aldrich) and filtered through a 0.1 µm centrifugal filter

(Millipore) prior to incubation with tissue overnight at 4°C in a humidity chamber. After the

overnight incubation slides were washed for 2 minutes in wash buffer. The second day,

the antibody cocktail was prepared as described (Supplementary Table 9) and incubated

with the tissues for 1 hour at 4°C in a humidity chamber. After staining, slides were

washed twice for 5 minutes in wash buffer and fixed in a solution of 2% glutaraldehyde

(Electron Microscopy Sciences) solution in low-barium PBS for 5 minutes. Slides were

washed in low-barium PBS for 20 seconds then, using a linear stainer, through 0.1 M Tris

at pH 8.5 (3x), ddH2O (2x), and then dehydrated by washing in 70% ethanol (1x), 80%

ethanol (1x), 95% ethanol (2x), and 100% ethanol (2x). Slides were dried under vacuum

prior to imaging.

MIBI-TOF imaging

Imaging was performed using a custom MIBI-TOF instrument with a Xe+ primary

ion source, as described previously[28,61]. 222 808 x 808um Fields of View (FOVs) were

acquired at approximately 600 nm resolution using an ion dose of 7nA*hr/mm^2. After

excluding 11 FOVs that contained necrotic or non-decidual tissue, or consisted of

duplicate tissue regions, the final dataset consisted of 211 FOVs from 66 patients.

Low-level image processing

Multiplexed image sets were extracted, slide background-subtracted, denoised,

and aggregate filtered as previously described[35,59–61]. For several markers, a

"background" channel consisting of signal from the mass 128 channel was used. All
parameters used as inputs for low-level processing are listed in Supplementary Table 9.

Feature annotation

Large tissue features were manually annotated in collaboration with a perinatal
pathologist. Pseudo-colored MIBI images with H3 to identify cell nuclei, vimentin for
decidual stromal cells, smooth muscle actin and CD31 for vessels, cytokeratin 7 (CK7)
for glands and the fetal cell columns, and HLA-G for EVTs were used to guide annotation.
Serial H&E sections, and an H&E recut of the entire block, if necessary, were additionally
used to supplement annotation. Labelling was performed in ImageJ and the annotated
features were exported as binary TIF masks.

Single cell segmentation

The Mesmer segmentation algorithm[36] was adapted specifically to segment the
cells in our dataset. First, training data were generated using a subset of 15 images out
of 211 in our cohort, in addition to 10 decidua MIBI-TOF images from titration data. 1024
x 1024 pixel crops were selected to encompass the range of different cell morphologies
present. The markers H3, vimentin, HLA-G, CD3, CD14 and CD56 were used to capture
the major cell lineages present. Subsequently, a team of annotators parsed these images
to identify the location of each unique cell using DeepCell Label, custom annotation
software   specifically   developed   for   this   task[36]   (code   URL:
https://github.com/vanvalenlab/deepcell-label). The manually annotated images were
used to generate partially overlapping crops of 256 x 256 pixels from each image. In total,
training data included 1600 distinct crops with 93,000 cells. This dataset was used to
retrain the Mesmer segmentation model, modifying the architecture to accept six distinct
channels of input. The output from the network was then post-processed using the default
model settings (Extended Data Fig. S2a).

Segmentation post-processing

Examination of the images revealed that glandular cells and chorionic villus
trophoblasts did not express any markers included in the training data; namely these cells
were predominantly CK7$^+$. This resulted in effectively nuclear-only segmentation being

predicted by the CNN within these features. To account for this, segmented cells that
overlapped with the gland mask were expanded radially by 5 pixels, and those in the cell
column mask by 2 pixels. This approach accounted for glandular cells and cell column
anchoring trophoblasts that were not expressing any markers but were included in the
training data, resulting in effectively nuclear-only segmentation being predicted by the
convolutional neural network. The number of pixels used for expansion was optimized to
approximate the observed cell size, based on systematic inspection of three images per
GA.

Single-cell phenotyping and composition

Single cell expression data were extracted for all cell objects and area-normalized.
Objects <100 pixels in area were deemed noncellular and excluded from subsequent
analyses. Single-cell data were linearly scaled with a scaling factor of 100 and ArcSinh-
transformed with a co-factor of 5. All mass channels were normalized to the 99th
percentile. To assign decidual cell populations (≥ 70% cell area in decidua) to a lineage,
the clustering algorithm FlowSOM (Bioconductor "FlowSOM" package in R)[37] was used,
which separated cells into 100 clusters based on the expression of 19 canonical lineage
defining markers (Extended Data Fig. S2b). Clusters were further classified into 21 cell
populations, with proper lineage assignments ensured by manual examination of
overlayed FlowSOM cluster identity with lineage-specific markers. Clusters containing
non biologically meaningful or distinct signals were assigned the label 'other'. Tregs were
identified by thresholding T cells (FlowSOM clusters 43, 53, 63) with CD3 signal ≥ the
mean CD3 expression of CD4$^+$ T cells and > 0.5 normalized expression of FOXP3. Mast
cells were identified as cells for which normalized expression of tryptase was >0.9. Mac2b
(CD11c$^+$) cells were identified as macrophages with >0.5 normalized expression of
CD11c. Placental macrophages (Hofbauer cells) were defined as CD14$^+$ >0.5 cells
located within the cell column. Cells from FlowSOM clusters 4, 5, and 15 ubiquitously and
predominantly expressed CK7 and were reassigned to the EVT2 subset if located within
the cell column feature mask, or as glandular cells otherwise (Extended Data Fig. S2b).
These thresholds were selected based on the distribution of lineage marker expression
(Extended Data Fig. S2c) as well as on systematic examination of the images by eye

since expression patterns varied significantly between markers.

Definition of thresholds for functional marker positivity

Cells were considered positive for a functional marker if their scaled expression
level was ≥ a set threshold, as described previously[61]. Thresholds for individual functional
markers were determined based on the distribution of functional marker expression and
by examining the images by eye, as expression patterns varied significantly between
markers (Extended Data Fig. S2e, Supplementary Table 4). To set the per marker
thresholds, 5 images for each functional marker were reviewed and increasing threshold
values were examined using custom software. Subsequently, cells defined as negative
for a marker based on the determined threshold value were re-examined to ensure the
thresholds were representative. For Ki67 positivity, only cells that had a nucleus in the
image were considered. Ki67 values were not cell size normalized because the Ki67
signal is exclusive to nuclei.

Blinded manual artery staging

Arteries were categorized into 5 remodeling stages based on criteria adapted from
the 4-stage model proposed by Smith et al[42]. These criteria were previously used to
describe spiral arteries observed in H&E and single channel IHC images and were
adapted to suit multiplexed MIBI data (Fig. 3a, details in Extended data Fig. S3a). 600
arteries were categorized according to these criteria by a single reviewer using
exclusively crops of MIBI pseudocolor overlays (SMA, Vimentin, CD31, H3, and HLA-G)
including only the artery (as defined by feature mask) and any EVTs in the lumen. The
reviewer was blinded to the rest of the image, serial H&E sections, gestational age, and
any clinical data. 12 partially captured arteries were excluded from the final dataset of
588 arteries.

Automated digitization of artery morphological features

The same format of cropped artery MIBI images that were manually scored by the
reviewer were used to calculate a set of geometric parameters for several selected
features. These features described the organization and structure of the vessel wall, the
continuity of the endothelium and its thickness, and the presence and structure of

intravascular EVTs. In order to capture these features, a structure of concentric circles

we termed the "onion" structure is defined, with the outer circle of this structure enclosing

the artery and the inner circles dividing it into layers. This structure is described below

using the two-dimensional cylindrical coordinate system with the radial axis r, azimuthal

(angular) axis ø, and origin of the axis at point (x,y). Point (x,y) is the user defined artery

center. For an artery in the binary mask M, the following algorithm was used to create the

"onion" structure (Extended Data Fig. S3c):

• Define a circle enclosing the artery, centered at point (x,y) with radius a as follows:

o (x,y) was taken as the user-defined artery center point

o a, the radius is defined as the maximum distance between (x,y) and the

717   edge of M – rounded up to the nearest integer multiple of n, such that $a=l*n$

718   for an integer l. n is a user defined thickness parameter for the "onion"

719   layers.

720   • Define the inner circles comprising the "onion" layers:

721   o Divide the radius a of the outer circle into l equal sections of length n,

722   creating layers along the radial r axis.

723   o The radii of the inner circles are then defined as: $0,1*n,2*n,…(l-1)*n$.

724   • Divide the "onion" into k equal sectors along the ø axis, k is a user defined integer.

725   • Subdivide each sector into segments:

726   o The sectors are internally divided by the circles, creating parts with 4

727   corners and 4 sides, with the 2 sides being straight (sector dividers), and

728   the 2 sides being arcs (parts of ellipse circumferences).

729   o The arcs are replaced with secants (straight line connecting the ends of the

730   arc), turning the segment into a trapezoid.

731   o The parameters n=10 pixels and k=100 were used to allow for segments

732   large enough to contain a sufficient number of pixels to average expression

733   over.

734   The following features are then extracted for each artery "onion":

735   1. Geometrical features:

736   a. radius - the maximum distance between any pixel within the mask and the

737   closest pixel on the edge of the mask.

738   b. perimeter – the Euclidean distance between all adjacent pixels on the edge

739   of the artery mask

740   c. area – the total number of pixels within the artery mask

741   2. Protein morphology features, for each of the following markers: CD31, CK7, H3,

742   HLA-G, SMA, VIM

743   a. Average signal – weighted-average over segments of marker expression,

744   where the weight of a segment corresponds to the number of pixels it

745   contains. Weighted average was used to avoid smaller inner segments

746   having disproportionate effect on the average.

747   b. Thickness –

748   i. For each sector we calculate the distance d between the inner-most

749   segment positive for the marker and the outer-most positive

750   segment. Positivity is measured by comparing the mean signal over

751   pixels the segment to a user defined threshold.

752   ii. The mean and standard deviation of thickness are calculated as the

753   mean and standard deviation of d over all sectors.

754   c. Radial coverage - the percentage of sectors positive for marker signal. A

755   sector is considered positive if the mean signal over sector pixels accedes

756   a user defined threshold.

757   d. Jaggedness – This feature measures the extent jaggedness of an artery

758   outline. To do so, first a skeletonization function written by Nicholas R.

759   Howe [62] is applied to the artery mask, this function returns a "skeleton" of

760   the artery outline. This "skeleton" also assigns values to the outline pixels

761   based on their distance from the core shape. Then, two different

762   binarization thresholds are chosen: a "non- branch" threshold (a high value

763   = 60 pixels, indicating greater topological distance and a "branch" threshold

764   (a low value = 5 pixels, indicating smaller topological distance). The ratio

765   between the total number of "non branch" and "branch" pixels is the

766   jaggedness.

Calculation of continuous SAR remodeling score δ

A supervised dimensionality reduction technique based on linear discriminant
analysis (LDA)[43] (code URL: https://github.com/davidrglass) was employed using the per
artery digitized morphological features and manually assigned remodeling stage labels
as inputs. All artery morphology feature values were standardized (mean subtracted and
divided by the standard deviation) and all arteries were used as training data. The LDA
output was:

a. The optimal linear combination of a subset of features, that maximized the
separation by manual stage between arteries in LDA space (Supplementary Table
6)
b. The coordinates of each artery in LDA space (Supplementary Table 5)

In order to define the SAR trajectory, a fourth-degree polynomial was fitted to the
artery coordinates in LDA space. To determine the optimal degree of the polynomial,
polynomials with degrees 1-6 were fitted and the degree that minimized the p-value for
separating δ distributions between arteries grouped by manual remodeling stage
(Extended Data Fig. S3f) was selected. The polynomial fit was implemented using the
MATLAB function fit and resulted in the following polynomial: $f(x) = 0.0005*x^4 - 0.01227*x^3$
$+ 0.1363*x^2 - 0.4354*x - 0.7425$. The polynomial was then numerically interpolated on a
dense 10^4 point grid and the distance from each artery point in LDA space to the
polynomial was calculated using this grid and the MATLAB exchange function
distance2curve [63]. δ per artery was then calculated as the line integral from the curve
origin to closest point to the artery on the curve (Fig. 3n, inset). This integral was
numerically calculated using a custom MATLAB script. δ values were linearly rescaled to
the range 1-5 using the MATLAB function rescale.

Cell type frequency as function of GA and SAR

For examining cell type frequencies within the decidua as function of GA and SAR
(Fig. 3, Fig. 4), per image cell frequency tables were constructed in which cell type
frequencies were calculated as the proportion of cells in the decidua feature mask of that
image. Cells located in other feature masks (artery, gland, vessel, or cell column masks)
were not counted, nor were cells of an unassigned type ('other'). In order to focus these

analyses on cell populations strictly found in the decidua, muscle and glandular cells were

also excluded; these cell types occasionally extended outside of their artery and gland

feature masks, respectively. Cell frequency as a function of GA for a cell type was defined

as the per image proportion values for that cell type, as function of the GAs associated

with the images. Similarly, cell frequency as a function of SAR for a cell type was defined

as the per image proportions of that cell type, as function of the mean δ values per image.

For the volcano plot in Fig. 3o, we fitted a linear regression model to the two above-

described functions. All linear regression models were implemented using the MATLAB

function fitlm and the volcano plot only shows points for which regression $R^2 \geq 0.05$. R^2

and p-values for all δ and GA based regressions can be found in Supplementary Table

7. The ration between R^2 in the two regression models was used to classify trends as GA-

driven, SAR-driven or synchronized. For example, the increase in EVTs out of all cells,

R_EVT, was classified as GA-driven because R^2 for R_EVT as a function of δ was 0.3,

but only 0.1 for R_EVT as a function of GA (Extended Data Figure S3g, Supplementary

Table 7). Another example is the increase in macrophages out of immune cells,

I_sumMac: it was classified as GA-driven since R^2 for I_sumMac as a function of GA was

0.6 but only 0.1 for I_sumMac as a function of δ (Extended Data Figure S3f,

Supplementary Table 7). For determining trend sizes depicted in Fig. 3o, the following

calculation was used: denote the per image frequencies of a cell type as V, and the

corresponding per image temporal stamps (either GA or mean image δ) as X. Trend size

is then calculated as the difference between the first and last time point in units of the

mean: $\frac{V(\max(X)) - V(\min(X))}{mean(V)}$.

Functional markers positivity rate per cell type as function of GA and SAR

For examining cell type specific temporal trends in the expression of functional

markers (Fig. 5a), 48 combinations of cell type- functional marker were selected. The

selected combinations were those for which the positivity frequency Z-score exceeded

0.5 (Fig. 2a, right panel). For each of these combinations, the frequency of cells positive

for the functional marker was calculated as the number of cells positive for the marker

(see "Definition of thresholds for functional marker positivity"), out of the total number of

cells of the same cell type in the image. All cells except those located within the cell

column mask were included to focus the analysis on functional marker trends of maternal
cells and EVTs that had infiltrated the decidua. For glandular cells, the location was further
restricted to the glands mask. The frequency of cells positive for a functional marker as a
function of GA, for a cell type, was defined as the per image positivity proportion values
as function of the GAs associated with the images. Similarly, marker positivity frequency
as a function of SAR for a cell type was defined as the per image proportions of that cell
type positive for the marker, as function of the mean δ values per image. For the volcano
plot in Fig. 5a, we fitted a linear regression model to the two above-described functions.
All linear regression models were implemented using the MATLAB function fitlm and the
volcano plot only shows points for which regression $R^2 \geq 0.05$. R^2 and p-values for all δ
and GA based regressions can be found in Supplementary Table 8. For determining trend
sizes depicted in Fig. 5a, the following calculation was used: denote the linear fit to the
per-image marker positivity proportion of a cell type as V, and the corresponding per
image temporal stamps (either GA or mean image δ) as X. Trend size is then calculated
as the difference between the first and last time point in units of the mean:
$\frac{V(\max(X)) - V(\min(X))}{mean(V)}$.

Ridge regression for predicting GA from immune composition

Ridge regression was implemented using the sklearn Python package
(sklearn.linear_model.Ridge, RidgeCV). Per-image immune frequencies were rescaled to
the range 0-1 prior to model fitting, using the sklearn scaling function. Images with fewer
than 10 immune cells were excluded (n=8). A randomly derived test-train split of 30/70
was used and GA distribution was verified to be equally represented in the test and train
sets (Extended Data Fig. S4a). Ridge regression adds a regularization penalty to the loss
function in order to prevent over or under representation of correlated variables, such as
immune cell populations. The penalty used for the test set (0.81) was selected using
Leave-One-Out Cross-Validation on the training set.

Definition of anatomical EVT location and associated arteries

Cell column EVTs were defined as EVTs located within cell column masks,
intravascular EVTs were located within artery masks, and interstitial EVTs were located

in the decidua. Perivascular EVTs were defined as interstitial EVTs located within 50
pixels of the edge of an artery, as defined by radial expansion of the artery masks (Fig.
6b). Arteries were said to have perivascular or intravascular EVT (Figs. 6f-h) if the number
of EVT in the appropriate artery compartment was $\geq$5. For Fig. 6j, only images that
contained all four EVT types were considered and cell to artery distance was measured
from the cell centroid as detected by segmentation to the border of the artery mask. For
Fig. 6i, one image was excluded (16_31762_20_8) due to abnormal tissue morphology.

SMA and endothelium loss scores

The loss scores presented in Fig. 6g, h were based on digitized morphological
features. For SMA, the average feature was used and for endothelium, the radial
coverage of CD31 (see "Automated digitization of artery morphological features"). The
values for each of the two features were then divided by their maximum across arteries
and subtracted from 1 to obtain a loss score. The resulting values were then linearly
rescaled to the range 0-1 using the MATLAB function rescale.

LDA of EVTs by compartment

A method similar to our calculation of the continuous SAR remodeling score δ was
used for compartment-wise analysis of EVT types. The input table consisted of marker
expression values per EVT. Lineage and functional markers expressed by EVTs were
included: CD56, CD57, HLA-G, CK7, PD-L1 and Ki67 (Fig. 2a). EVTs were labeled by
spatial compartment: cell column, interstitial, perivascular or intravascular (see "Definition
of anatomical EVT location"). Marker expression values were standardized (mean
subtracted and divided by the standard deviation) and cell column, interstitial, and
intravascular location labels per EVT were used for training the LDA model. Perivascular
EVTs were withheld as a test set. Due to the small number or features (markers) a one-
dimensional LDA was calculated yielding a single coordinate LD1. LD1 was the optimal
linear combination of a subset of markers, to maximize the separation by compartment
between EVTs (Supplementary Table 10). LD1 values were subsequently calculated for
the withheld test set of perivascular EVTs (Supplementary Table 11). To calculate the
difference in similarity to intravascular EVT between interstitial and perivascular, the
following calculation was used: intravascular-perivascular similarity was defined as

$sim_{intra-peri} = mean(Ld1_{intravascular}) - mean(Ld1_{perivascular})$. Similarly, intravascular-

interstitial similarity was defined as $sim_{intra-inter} = mean(Ld1_{intravascular}) -$

$mean(Ld1_{interstitial})$. The difference in these similarities was then calculated as:

$\frac{sim_{intra-inter} - sim_{intra-peri}}{im_{intra-inter}}$ in %.

Statistical analyses

Throughout the paper, the Kruskal-Wallis test was implemented using the
MATLAB function KruskalWallis. All linear regression models were implemented using
the MATLAB function fitlm unless stated otherwise. The sided Wilcoxon signed rank test
for paired analysis in Extended Data Fig. S6 was implemented using the MATLAB
function signrank. MATLAB version used throughout the paper for statistical analysis is
MATLAB 2020b.

Data availability

898   Imaging data, segmentation masks, and extracted features will be made publicly
899 available prior to publication. Code is currently available upon request and will be made
900 public prior to publication.

References

1. Moffett-King, A. Natural killer cells and pregnancy. *Nature reviews. Immunology* **2**,
656–663 (2002).

2. Red-Horse, K. *et al.* Trophoblast differentiation during embryo implantation and
formation of the maternal-fetal interface. *The Journal of clinical investigation* **114**,
744–754 (2004).

3. Menkhorst, E., Winship, A., van Sinderen, M. & Dimitriadis, E. Human extravillous
trophoblast invasion: intrinsic and extrinsic regulation. *Reproduction, fertility, and*
development **28**, 406–415 (2016).

4. Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal–fetal
interface in humans. *Nature* **563**, 347–353 (2018).

5. Suryawanshi, H. *et al.* A single-cell survey of the human first-trimester placenta and

decidua. *Science Advances* **4**, eaau4788 (2018).

6.   Mor, G., Aldo, P. & Alvero, A. The unique immunological and microbial aspects of
pregnancy. *Nature reviews. Immunology* **17**, 469–482 (2017).

7.   Kam, E. P. Y., Gardner, L., Loke, Y. W. & King, A. The role of trophoblast in the
physiological change in decidual spiral arteries. *Human Reproduction* **14**, (1999).

8.   Whitley, G. S. J. & Cartwright, J. E. Cellular and Molecular Regulation of Spiral
Artery Remodelling: Lessons from the Cardiovascular Field. *Placenta* **31**, 465–474
(2010).

9.   Pijnenborg, R., Vercruysse, L. & Hanssens, M. The Uterine Spiral Arteries In
Human Pregnancy: Facts and Controversies. *Placenta* vol. 27 939–958 (2006).

10.   Kliman, H. J. Uteroplacental Blood Flow: The Story of Decidualization,
Menstruation, and Trophoblast Invasion. *The American Journal of Pathology* **157**,
1759–1768 (2000).

11.   Burton, G. J., Woods, A. W., Jauniaux, E. & Kingdom, J. C. P. Rheological and
Physiological Consequences of Conversion of the Maternal Spiral Arteries for
Uteroplacental Blood Flow during Human Pregnancy. *Placenta* **30**, 473–482
(2009).

12.   Khong, T., de Wolf, F., Robertson, W. & Brosens, I. Inadequate maternal vascular
response to placentation in pregnancies complicated by pre-eclampsia and by
small-for-gestational age infants. *British journal of obstetrics and gynaecology* **93**,
1049–1059 (1986).

13.   Brosens, I., Pijnenborg, R., Vercruysse, L. & Romero, R. THE "GREAT
OBSTETRICAL SYNDROMES" ARE ASSOCIATED WITH DISORDERS OF DEEP
PLACENTATION. *American Journal of Obstetrics and Gynecology* **204**, 193
(2011).

14.   Carter, A. M. Animal Models of Human Placentation – A Review. *Placenta* **28**, S41–
S47 (2007).

15.   Schmidt, A., Morales-Prieto, D. M., Pastuschek, J., Fröhlich, K. & Markert, U. R.

Only humans have human placentas: molecular differences between mice and humans. *Journal of Reproductive Immunology* **108**, 65–71 (2015).

16. Pijnenborg, R., Vercruysse, L. & Carter, A. M. Deep trophoblast invasion and spiral artery remodelling in the placental bed of the lowland gorilla. *Placenta* **32**, 586–591 (2011).

17. Pijnenborg, R., Vercruysse, L. & Carter, A. M. Deep trophoblast invasion and spiral artery remodelling in the placental bed of the chimpanzee. *Placenta* **32**, 400–408 (2011).

18. Gaynor, L. M. & Colucci, F. Uterine Natural Killer Cells: Functional Distinctions and Influence on Pregnancy in Humans and Mice. *Frontiers in Immunology* **0**, 467 (2017).

19. Yang, F., Zheng, Q. & Jin, L. Dynamic Function and Composition Changes of Immune Cells During Normal and Pathological Pregnancy at the Maternal-Fetal Interface. *Frontiers in Immunology* **10**, 2317 (2019).

20. Pijnenborg, R., Bland, J. M., Robertson, W. B. & Brosens, I. Uteroplacental arterial changes related to interstitial trophoblast migration in early human pregnancy. *Placenta* **4**, 397–413 (1983).

21. Chaline, J. Increased cranial capacity in hominid evolution and preeclampsia. *Journal of reproductive immunology* **59**, 137–152 (2003).

22. Rockwell, C., Vargas, E. & Moore, L. G. Human physiological adaptation to pregnancy: Inter- and intraspecific perspectives. *American Journal of Human Biology* **15**, 330–341 (2003).

23. Zhu, H., Hou, C. C., Luo, L. F., Hu, Y. J. & Yang, W. X. Endometrial stromal cells and decidualized stromal cells: Origins, transformation and functions. *Gene* **551**, 1–14 (2014).

24. TANG, B., GULLER, S. & GURPIDE, E. Mechanism of Human Endometrial Stromal Cells Decidualization. *Annals of the New York Academy of Sciences* **734**, 19–25 (1994).

25.   Moffett, A. & Colucci, F. Uterine NK cells: active regulators at the maternal-fetal
interface. *The Journal of clinical investigation* **124**, 1872–1879 (2014).

26.   Robson, A. *et al.* Uterine natural killer cells initiate spiral artery remodeling in human
pregnancy. *The FASEB Journal* **26**, 4876–4885 (2012).

27.   Craven, C. M., Morgan, T. & Ward, K. Decidual spiral artery remodelling begins
before cellular interaction with cytotrophoblasts. *Placenta* **19**, 241–252 (1998).

28.   Keren, L. *et al.* MIBI-TOF: A multiplexed imaging platform relates cellular
phenotypes and tissue structure. *Science Advances* **5**, eaax5851 (2019).

29.   Chang, R.-Q., Li, D.-J. & Li, M.-Q. The role of indoleamine-2,3-dioxygenase in
normal and pathological pregnancies. *American Journal of Reproductive*
Immunology **79**, e12786 (2018).

30.   Yasinska, I. M. *et al.* The Tim-3-Galectin-9 Pathway and Its Regulatory
Mechanisms in Human Breast Cancer. *Frontiers in Immunology* **0**, 1594 (2019).

31.   Li, M. *et al.* Tim-3/CTLA-4 pathways regulate decidual immune cells-extravillous
trophoblasts interaction by IL-4 and IL-10. *FASEB journal : official publication of the*
Federation of American Societies for Experimental Biology **35**, (2021).

32.   Hu, X. *et al.* Newly characterized decidual Tim-3+ Treg cells are abundant during
early pregnancy and driven by IL-27 coordinately with Gal-9 from trophoblasts.
Human reproduction (Oxford, England) **35**, 2454–2466 (2020).

33.   Wang, S. *et al.* Tim-3 protects decidual stromal cells from toll-like receptor-
mediated apoptosis and inflammatory reactions and promotes Th2 bias at the
maternal-fetal interface. *Scientific Reports 2015 5:1* **5**, 1–9 (2015).

34.   Sedlmayr, P. *et al.* Localization of indoleamine 2,3-dioxygenase in human female
reproductive organs and the placenta. *Molecular Human Reproduction* **8**, 385–391
(2002).

35.   Baranski, A. *et al.* MAUI (MBI Analysis User Interface)-An image processing
pipeline for Multiplexed Mass Based Imaging. *PLoS computational biology* **17**,
(2021).

36.   Greenwald, N. F. *et al.* Whole-cell segmentation of tissue images with human-level
performance using large-scale data annotation and deep learning. *bioRxiv*
2021.03.01.431313 (2021) doi:10.1101/2021.03.01.431313.

37.   van Gassen, S. *et al.* FlowSOM: Using self-organizing maps for visualization and
interpretation of cytometry data. *Cytometry Part A* **87**, 636–645 (2015).

38.   Faas, M. M. & de Vos, P. Uterine NK cells and macrophages in pregnancy. *Placenta*
56, 44–52 (2017).

39.   Kämmerer, U. *et al.* Unique Appearance of Proliferating Antigen-Presenting Cells
Expressing DC-SIGN (CD209) in the Decidua of Early Human Pregnancy. *The*
American Journal of Pathology **162**, 887–896 (2003).

40.   Huhn, O. *et al.* Distinctive phenotypes and functions of innate lymphoid cells in
human decidua during early pregnancy. *Nature Communications* **11**, (2020).

41.   Salvany-Celades, M. *et al.* Three Types of Functional Regulatory T Cells Control T
Cell Responses at the Human Maternal-Fetal Interface. *Cell Reports* **27**, 2537-
2547.e5 (2019).

42.   Smith, S. D., Dunk, C. E., Aplin, J. D., Harris, L. K. & Jones, R. L. Evidence for
immune cell involvement in decidual spiral arteriole remodeling in early human
pregnancy. *American Journal of Pathology* **174**, 1959–1971 (2009).

43.   Tsai, A. G. *et al.* Multiplexed single-cell morphometry for hematopathology
diagnostics. *Nature Medicine* **26**, (2020).

44.   Nielsen, C. M., White, M. J., Goodier, M. R. & Riley, E. M. Functional significance
of CD57 expression on human NK cells and relevance to disease. *Frontiers in*
Immunology vol. 4 (2013).

45.   Gonçalves Silva, I. *et al.* The Tim-3-galectin-9 Secretory Pathway is Involved in the
Immune Escape of Human Acute Myeloid Leukemia Cells. *EBioMedicine* **22**, 44–
57 (2017).

46.   Blois, S. M. *et al.* Role of galectin-glycan circuits in reproduction: from healthy
pregnancy to preterm birth (PTB). *Seminars in Immunopathology 2020 42:4* **42**,

469–486 (2020).

47. Blois, S. M. *et al.* Pregnancy Galectinology: Insights Into a Complex Network of
Glycan Binding Proteins. *Frontiers in Immunology* **0**, 1166 (2019).

48. Robillard, P. Y., Dekker, G. A. & Hulsey, T. C. Evolutionary adaptations to pre-
eclampsia/eclampsia in humans: Low fecundability rate, loss of oestrus,
prohibitions of incest and systematic polyandry. *American Journal of Reproductive*
Immunology vol. 47 104–111 (2002).

49. Naruse, K. *et al.* Localization of matrix metalloproteinase (MMP)-2, MMP-9 and
tissue inhibitors for MMPs (TIMPs) in uterine natural killer cells in early human
pregnancy. *Human Reproduction* **24**, 553–561 (2009).

50. Hunt, J., Petroff, M., McIntire, R. & Ober, C. HLA-G and immune tolerance in
pregnancy. *FASEB journal : official publication of the Federation of American*
Societies for Experimental Biology **19**, 681–693 (2005).

51. Tilburgs, T., Evans JH, Crespo, Â. & Strominger, J. The HLA-G cycle provides for
both NK tolerance and immunity at the maternal-fetal interface. *Proceedings of the*
National Academy of Sciences of the United States of America **112**, 13312–13317
(2015).

52. Isaka, K. *et al.* Expression and Activity of Matrix Metalloproteinase 2 and 9 in
Human Trophoblasts. *Placenta* **24**, 53–64 (2003).

53. Harris, L. K., Jones, C. J. P. & Aplin, J. D. Adhesion Molecules in Human
Trophoblast – A Review. II. Extravillous Trophoblast. *Placenta* **30**, 299–304 (2009).

54. None, K. *et al.* Evolution of placental invasion and cancer metastasis are causally
linked. *Nature ecology & evolution* **3**, 1743–1753 (2019).

55. Bulmer, J., Thrower, S. & Wells, M. Expression of epidermal growth factor receptor
and transferrin receptor by human trophoblast populations. *American journal of*
reproductive immunology (New York, N.Y. : 1989) **21**, 87–93 (1989).

56. Wright JK *et al.* HER1 signaling mediates extravillous trophoblast differentiation in
humans. *Biology of reproduction* **83**, 1036–1045 (2010).

57. Coorens, T. H. H. *et al.* Inherent mosaicism and extensive mutation of human placentas. *Nature 2021 592:7852* **592**, 80–85 (2021).

58. Angelo, M. *et al.* Multiplexed ion beam imaging of human breast tumors. *Nature Medicine* **20**, 436–442 (2014).

59. McCaffrey, E. F. *et al.* Multiplexed imaging of human tuberculosis granulomas uncovers immunoregulatory features conserved across tissue and blood. *bioRxiv* 2020.06.08.140426 (2020) doi:10.1101/2020.06.08.140426.

60. Risom, T. *et al.* Transition to invasive breast cancer is associated with progressive changes in the structure and composition of tumor stroma. *bioRxiv* 2021.01.05.425362 (2021) doi:10.1101/2021.01.05.425362.

61. Keren, L. *et al.* A Structured Tumor-Immune Microenvironment in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging. *Cell* **174**, 1373-1387.e19 (2018).

62. Howe, N. Better Skeletonization - File Exchange - MATLAB Central. https://www.mathworks.com/matlabcentral/fileexchange/11123-better-skeletonization.

63. D'Errico, J. distance2curve - File Exchange - MATLAB Central. https://www.mathworks.com/matlabcentral/fileexchange/34869-distance2curve.

Acknowledgements

We thank V. Winn, N. Vivanco, A. Moore, E. McCaffrey, D. Glass, T. Risom, J. P. Oliveria, K. O'Neill and C. Coutifaris for comments. M.A. is supported by 5U54CA20997105, 5DP5OD01982205, 1R01CA24063801A1, 5R01AG06827902, 5UH3CA24663303, 5R01CA22952904, 1U24CA22430901, 5R01AG05791504, and 5R01AG05628705 from NIH, W81XWH2110143 from DOD, and other funding from the Bill and Malinda Gates Foundation, Cancer Research Institute, the Parker Center for Cancer Immunotherapy, and the Breast Cancer Research Foundation. S.G is supported by the Bill and Melinda Gates Foundation OPP1113682. I.A. is an awardee of the

1081 Weizmann Institute of Science – Israel National Postdoctoral Award Program for
1082 Advancing Women in Science. E.S is supported by National Science Scholarship, Agency
1083 for Science, Technology, and Research (A*STAR), Singapore.

Author contributions

S.G. assembled the tissue cohort, performed and designed experiments,
annotated images, analyzed and interpreted data and wrote the manuscript. I.A. analyzed
and interpreted data, wrote the manuscript. E.S. performed and designed experiments,
annotated images, analyzed and interpreted data, wrote the manuscript. G.R. advised on
cohort design, assembled tissue cohort, annotated images. A.B, N.G., G.M., M.S., W.G.
and D.V.V. wrote software for image analysis. M.B. advised on experimental design and
reagent validation. E.J. assembled cohort patient metadata. L.K. advised on
computational analysis. Z.K. Prepared and validated reagents. S.K. Constructed the
tissue microarray. S.W. annotated images. T.H. validated reagents and advised on
experimental design. M.R. oversaw tissue microarray construction. M.A. conceived the
study, advised on experimental design and data analysis, wrote the manuscript.

Supplementary table legends

Supplementary Table 1 - Patients table. This table provides patient meta-data
such as age, ethnicity, body mass index, parity and relevant medical conditions such as
HIV.

**Supplementary Table 2 - Information on the histological characteristics of
the blocks retrieved, including the presence of cell column anchoring villi.** This
table shows, for each patient's block, whether cell column anchoring villi were present (1)
or absent (0), and the number of regions containing spiral arteries annotated as
appropriate for TMA construction by the pathologist (Methods). In blocks containing ≥ 2
distinct, separate pieces of tissue, cell column villi were considered present if they were
present on any piece containing pathologist annotations.

Supplementary Table 3 - Cell table. This table enumerates all single cells in this
study and provides their location, morphological characteristics such as size and shape,

marker expression, FlowSOM cluster assignment and cell type assignment.

Supplementary Table 4 - Positivity binary threshold for functional markers.
This table provides binary expression thresholds per functional marker- used to determine
whether cells are positive for that marker (See methods).

Supplementary Table 5 - Artery properties and staging. This table provides
arteries meta-data, including their measured digitized morphological features (see
Methods), manual stage and remodeling score δ.

Supplementary Table 6 - LDA coefficients for artery morphological features.
This table contains the coefficients per artery feature that define the LDA space used for
digitized artery staging (Fig. 3n). For the features selected by the algorithm, their ld1 and
ld2 coefficients are listed. Additional columns show the Z scored absolute values of these
coefficients.

Supplementary Table 7 - Regression results for cell type proportions as a
function of GA and δ. This table provides the values plotted in Fig. 3o. Each row
represents a cell proportion with those starting with R_ indication proportion out of all cells
in the image, I_ - proportion out of immune cells in image, N_- proportion out of NK cells
in image, M_ proportion out of macrophages in image, T_- proportion out of T cells in
image, F_- proportion out of EVT in image, S_ proportion out of structural cells in image
(Fig. 2b). The columns show values for the linear regression on per image proportions as
function of GA and remodeling score δ: the log transformed ratio of R^2, the maximal
obtained regression R^2 (maximal between GA and δ), the minimal obtained regression p-
value and trend size (see Methods).

Supplementary Table 8 - Regression results for functional markers
expression as a function of GA and δ. This table provides the values plotted in Fig. 5a.
Each row represents a combination of a cell type and a functional marker. The columns
show values for the linear regression on per image marker positivity rates for the marker-
cell type as function of GA and remodeling score δ: the log transformed ratio of R^2, the
maximal obtained regression R^2 (maximal between GA and δ), the minimal obtained
regression p-value and trend size (see Methods).

Supplementary Table 9 - Information on antibodies, metal reporters, staining
concentrations, and parameters used for low-level processing of MIBI data. This
table contains, for each marker, relevant antibody information including clone, vendor,
vendor ID, channel and elemental reporter, and final staining titers used. The parameters
used for marker-specific low-level processing of MIBI data (background removal,
denoising, and aggregate removal steps as previously described) are also shown.

Supplementary Table 10 - LD1 coefficients for markers expressed by EVT.
This table contains the coefficients per EVT expressed marker that define the LDA space
used for measuring similarity between anatomical tissue compartments (see Methods,
Extended Data Fig. 6c). For the markers selected by the algorithm, their ld1 coefficients
are listed. An additional column shows the Z scored absolute values of these coefficients.

Supplementary Table 11 - LD1 values per EVT. This table shows ld1 values per
single EVT with LDA input- standardized marker expression values (see Methods,
Extended Data Fig. 6c). Additional metadata such as EVT type, anatomical location, the
image the cell was taken from is also provided.