Characterization of apicomplexan amino acid transporters (ApiATs) in the malaria parasite

Plasmodium falciparum

Jan Stephan Wichers[1,2,3], Carolina van Gelder[1,2,3†*], Gwendolin Fuchs[1,2,3†§], Julia Mareike

Ruge[1,2,3], Emma Pietsch[1,2,3], Josie L. Ferreira[1,4], Soraya Safavi[1,2,3], Heidrun von Thien[1,2,3], Paul-

Christian Burda[1,2,3], Paolo Mesén-Ramirez[2], Tobias Spielmann[2], Jan Strauss[1,2,3,5], Tim-Wolf

Gilberger[1,2,3,#], Anna Bachmann[1,2,3,6,#]

[1] Centre for Structural Systems Biology, 22607 Hamburg, Germany

[2] Bernhard Nocht Institute for Tropical Medicine, 20359 Hamburg, Germany

[3] Biology Department, University of Hamburg, 22609 Hamburg, Germany

[4] Heinrich-Pette-Institute, Leibniz Institute for Experimental Virology, 20251 Hamburg, Germany

[5] GEOMAR Helmholtz Centre for Ocean Research Kiel, 24105 Kiel, Germany

[6] German Center for Infection Research (DZIF), partner site Hamburg-Borstel-Lübeck-Riems, Germany

† contributed equally

* Present address: University of Duisburg-Essen, 45141 Essen, Germany

§ Present address: Karolinska Institutet, 171 77 Stockholm, Sweden

Corresponding author: gilberger@bnitm.de; bachmann@bnitm.de

ORCID:

Jan Stephan Wichers: 0000-0002-0599-1742

Carolina van Gelder: 0000-0002-8109-3098

Gwendolin Fuchs: 0000-0001-9294-6984

Julia Mareike Ruge: 0000-0002-6295-0293

Emma Pietsch: 0000-0003-4744-1666

Josie L. Ferreira: 0000-0002-4411-6131

Soraya Safavi: 0000-0001-9989-0952

Paul-Christian Burda: 0000-0003-0461-4352

Paolo Mesén-Ramírez: 0000-0001-7842-5867

Tobias Spielmann: 0000-0002-3968-4601

Jan Strauss: 0000-0002-6208-791X

Tim-Wolf Gilberger: 0000-0002-7965-8272

Anna Bachmann: 0000-0001-8397-7308

ABSTRACT

During the symptomatic human blood phase, malaria parasites replicate within red blood

cells. Parasite proliferation relies on the uptake of nutrients, such as amino acids, from the

host cell and the blood plasma, requiring transport across multiple membranes. Amino acids

are delivered to the parasite through the parasite surrounding vacuolar compartment by

specialized nutrient-permeable channels of the erythrocyte membrane and the

parasitophorous vacuole membrane (PVM). However, further transport of amino acid across

the parasite plasma membrane (PPM) is currently not well characterized. In this study, we

focused on a family of Apicomplexan amino acid transporters (ApiATs) that comprises five

members in *Plasmodium falciparum*. First, we localized four of the *Pf*ApiATs at the PPM using

endogenous GFP-tagging. Next, we applied reverse genetic approaches to probe into their

essentiality during asexual replication and gametocytogenesis. Upon inducible knockdown

and targeted gene disruption a reduced asexual parasite proliferation was detected for

*Pf*ApiAT2 and *Pf*ApiAT4. Functional inactivation of individual *Pf*ApiATs targeted in this study

had no effect on gametocyte development. Our data suggest that individual *Pf*ApiATs are

partially redundant during asexual *in vitro* proliferation and fully redundant during

gametocytogenesis of *P. falciparum* parasites.

IMPORTANCE

Malaria parasites live and multiply inside cells. To facilitate their extremely fast intracellular

proliferation they hijack and transform their host cells. This also requires the active uptake of

nutrients, such as amino acids, from the host cell and the surrounding environment through

various membranes that are the consequence of the parasite's intracellular lifestyle. In this

manuscript we focus on a family of putative amino acid transporters termed ApiAT. We show

expression and localization of four transporters in the parasite plasma membrane of

Plasmodium falciparum-infected erythrocytes that represent one interface of the pathogen

to its host cell. We probed into the impact of functional inactivation of individual transporters

on parasite growth in asexual and sexual blood stages of *P. falciparum* and reveal that only

two of them show a modest but significant reduction in parasite proliferation but no impact

on gametocytogenesis pointing towards redundancy within this transporter family.

INTRODUCTION

Malaria parasites replicate within human erythrocytes during the asexual blood phase, which

is responsible for the symptoms of the disease. Although *Plasmodium falciparum* is able to

synthesize some amino acids *de novo*[1–6] during the intraerythrocytic development, the

parasite rely heavily on amino acid acquisition from its host. Amino acids are derived either

from the digestion of hemoglobin endocytosed from the infected erythrocyte[1,7] or from the

uptake of free amino acids from the blood plasma[1]. Both processes are important for

efficient parasite growth[8]. The parasite is able to import all 20 naturally occurring α-amino

acids from the external medium and uses them for its own protein synthesis[9–12]. Especially,

the import of isoleucine[13–15] – and for some *P. falciparum* strains also methionine[13] – is

crucial for the survival of the parasite as adult human hemoglobin lacks isoleucine.

Accordingly, an increase in the permeability to a range of amino acids has been reported for

erythrocytes upon *Plasmodium* infection[10,11,16,17]. This is mainly mediated via the New

Permeability Pathways (NPPs) established by the parasite within the membrane of the

infected erythrocyte[16,17]. While the subsequent transport across the parasitophorous vacuole

membrane (PVM) is linked to nutrient-permeable channel activity[18–21] the molecular

machinery responsible for the further transport across the parasite plasma membrane (PPM)

is not well defined. Neutral amino acids like isoleucine and methionine traverse the PPM

more rapidly than anionic and cationic amino acids, which may be coupled to the transport of

other substrates like H^+ or Na^+ [16,22]. However, neither the membrane transporters nor the

exact mechanism(s) by which amino acids cross the PPM have been characterized so far[8,22].

The *P. falciparum* transportome is predicted to be encoded by 144 genes[23], of which at least

eleven are classified as putative amino acid transporters[23,24]. Five of these putative amino

acid transporters belong to the Apicomplexan amino acid transporter (ApiAT)[25,26] family. This

Apicomplexan-specific family of transmembrane transporters can be subdivided into the

eleven subfamilies ApiAT1–11. Some of the subfamilies are lineage-specific. For instance,

ApiAT4, ApiAT8, ApiAT9 and ApiAT10 are only present in the genomes of *Plasmodium spp.*[1,25].

Others, such as the most ancient variant ApiAT2, can be found in many different

Apicomplexan classes[25]. The main feature of the ApiAT family is the possession of multiple,

typically twelve, transmembrane domains characteristic of solute transporters with a

signature sequence between transmembrane domains 2 and 3[25]. This classifies them as

members of the major facilitator superfamily (MFS)[24,27,28]. However, overall they have limited

sequence similarity to other known eukaryotic or prokaryotic transporters[25].

P. falciparum possesses five of the eleven ApiAT subfamilies: *Pf*ApiAT2 (*Pf*ApiAT2/MFR4:

PF3D7_0914700), *Pf*ApiAT4 (MFR5: PF3D7_1129900), *Pf*ApiAT8 (NPT1: PF3D7_0104800),

*Pf*ApiAT9 (MFR2: PF3D7_0104700) and *Pf*ApiAT10 (MFR3: PF3D7_0312500). Previous work in

the rodent malaria species *Plasmodium berghei* (*Pb*) and the related Apicomplexan parasite

Toxoplasma gondii (*Tg*) showed that *Pb*ApiAT8 (or *Pb*NPT1) and several other *Tg*ApiATs are

localized at the PPM[29–32] and possess amino acid transport activity[25,26,29–34]. To date, only

*Pb*ApiAT4 has been shown to play an important role in parasite proliferation within

erythrocytes[35]. Apart from this, *Pb*ApiAT8 is essential for gametocyte development[29,30,35] and

all variants except *Pb*ApiAT2 appear to be necessary for exflagellation of male gametocytes[35].

Additionally, sporozoite development likely requires *Pb*ApiAT2, *Pb*ApiAT4, *Pb*ApiAT9 and

*Pb*ApiAT10[35] and a lack of *Pb*ApiAT2 further impairs oocyst development. Furthermore, an

association of episomally overexpressed *Pf*ApiAT10-GFP (termed MFR3-GFP) with the

mitochondrion has been reported[36].

In this study, we characterize the localization and essentiality of the ApiAT family members of

P. falciparum during intraerythrocytic development in order to further dissect the function of

these unique transporters.

RESULTS

P. falciparum ApiATs localize at the parasite plasma membrane

The five members of the ApiAT family in *P. falciparum* show different gene expression

patterns and mRNA levels during the intraerythrocytic developmental cycle (IDC)[37]. While

*Pf*ApiAT2 has its maximum transcript level in early ring stage parasites (8 hpi) and *Pf*ApiAT4

and *Pf*ApiAT10 mRNA levels peak in late ring stage parasites (16 hpi), *Pf*ApiAT8 shows a

maximum of transcripts in late stage schizonts (48 hpi) and mRNA of *Pf*ApiAT9 is almost

absent during the IDC (Figure 1A). Overall, *Pf*ApiAT2 and *Pf*ApiAT4 are most abundantly

expressed on mRNA level.

To determine their protein expression and localization, we tagged each of the five members

of the *Pf*ApiAT family endogenously with GFP using the selection-linked integration (SLI)

system[38]. Correct integration of the corresponding targeting plasmids into the respective

genomic loci was verified by PCR (Figure S1A). Except for 3D7-ApiAT9-GFP, all generated

transgenic cell lines expressed the full-length fusion protein (Figure S1B) to a sufficient level

that allowed its subcellular localization. All of them are localized at the PPM (Figure 1B). This

localization was confirmed by co-localization with the episomally expressed PPM marker

Lyn[39]-mCherry[38] and becomes particularly evident in schizont stage parasites and free

merozoites, when PPM and PVM can clearly be separated (Figure 1C).

In contrast to published data[36], we also found endogenously GFP-tagged *Pf*ApiAT10 localizing

at the PPM. To re-probe into the apparent PPM localization of endogenously GFP-

tagged   ApiAT10 we also over-expressed this gene as a GFP and mCherry fusion protein

using two different promotors (*crt*[40] *or ama1*[41]). This allowed us to assess the influence of the

tags as well as differential expression profiles on   ApiAT10 protein localization. All cell lines

showed PPM localization of the   ApiAT10 fusion proteins (Figure 1B-E, S1). Of note,

*Pf*ApiAT10-TGD parasites showed also no conferral of drug resistance to Atovaquone (Figure

S1B).

Individual *Pf*ApiATs are not essential for asexual blood stage development

To probe into the essentiality of the ApiAT family for asexual parasite proliferation, we first

targeted the most abundantly expressed *Pf*ApiATs, *Pf*ApiAT2 and *Pf*ApiAT4 by conditional

knockdown. Down-regulation was achieved by introducing a glmS ribozyme sequence[40]

before the 3'UTR of either the *apiat2* or *apiat4* genomic locus (Figure S1). The ribozyme was

activated by addition of 2.5 mM D-(+)-glucosamine hydrochloride (GLCN), which resulted in

degradation of mRNA and therefore decreased protein levels. This was assessed and

quantified by live cell fluorescence microscopy after two cycles. GLCN treatment resulted in a

decreased GFP-fluorescence of 85.9 % ± SD 0.9 for 3D7-*Pf*ApiAT2-GFP or 74.8 % ± SD 14.0 for

3D7-*Pf*ApiAT4-GFP (Figure 2A–D) and led to a moderate, but significant reduction of parasite

growth of 30–35 % compared to the control (Figure 2E). This data indicates that *Pf*ApiAT2

and *Pf*ApiAT4 play a role in efficient blood cell proliferation but imply that they might be non-

essential. Therefore, we targeted these genes with deletion constructs using the SLI system[38]

that lead to the expression of severely truncated versions of the ApiATs. In this targeted gene

disruption (TGD) approach, we also included *Pf*ApiAT8 and *Pf*ApiAT10.

Imaging of the TGD cell lines revealed a more diffuse, but still membrane-associated GFP

signal (Figure 3A–D). This might be due to the remaining transmembrane domains of the

truncated *Pf*ApiAT mutants; however, our approach deleted at least ¾ of their predicted TM

domains and thus most likely abolished transporter activity. In concordance with the

inducible knockdown data, functional inactivation by truncation of *Pf*ApiAT2 and *Pf*ApiAT4 in

the corresponding transgenic cell lines (3D7-ApiAT2-TGD, 3D7-ApiAT4-TGD) led to a

moderate decrease of parasite proliferation of 20.2 % ± SD 3.2 and 19.8 % ± SD 8.6 after two

parasite replication cycles (Figure 3E). No significant reduction of growth was observed upon

disruption of *Pf*ApiAT8 and *Pf*ApiAT10 (Figure 3E). Interestingly, cultivation in amino acid

depleted medium (approximately 90 % reduced concentration) did not indicate a higher

susceptibility of any of the TGD cell lines to low amino acid concentrations compared to wild

type parasites (Figure 3F).

To probe into potential transcriptional perturbations within this gene family due to functional

inactivation of a single member, quantitative real-time PCR (qPCR) analysis was performed

using RNA from four different time points during asexual blood stage replication. However,

no consistent upregulation of RNA levels of other *Pf*ApiAT family members was observed in

individual *Pf*ApiAT TGDs (Figure S3).

_Pf_ApiATs are dispensable during gametocyte development

Previous data[30,35] indicated a role of ApiAT8 during gametocyte development of the rodent

malaria parasite *P. berghei*. Therefore, we re-engineered the GFP-tagged gene knockdown

(Figure S4) and deletion cell lines (Figure S5) for *Pf*ApiAT2, *Pf*ApiAT4, *Pf*ApiAT8 and *Pf*ApiAT10

in an inducible gametocyte producing parasite line (3D7-iGP-GDV1GFP-DD[41]) using the same

SLI approach. The resulting parasite lines allowed a robust, efficient and synchronized

induction of gametocytogenesis and therefore a solid basis for phenotypic analysis. First,

using the C-terminal GFP tag, we confirmed expression of all four *Pf*ApiATs in gametocytes.

As expected, most *Pf*ApiATs remain PPM localized during gametocytogenesis, which was

additionally confirmed by the colocalization with the episomally expressed PPM marker

Lyn[39]-mCherry[38] (Figure 4A–F). The exception was *Pf*ApiAT9, which showed only a faint

background staining in all gametocyte stages in the 3D7-ApiAT9-GFP line (Figure 4D). The

prominent GFP signal at the food vacuole in gametocytes is most likely an unspecific

hemozoin signal, as it is also observed in 3D7-iGP control parasites (Figure S4C). Of note,

*Pf*ApiAT2 was observed to be strongest expressed in early-stage gametocytes and weaker in

late-stages (Figure 5B, D; Figure S4D), while *Pf*ApiAT4 showed strongest expression in late-

stage gametocytes (Figure 5C, E). Next, we investigated the consequence of glmS-based

conditional knockdown for *Pf*ApiAT2 and *Pf*ApiAT4 (Figure 5A). Although 75–80 % knockdown

of *Pf*ApiAT2 or *Pf*ApiAT4 expression was achieved (Figure 5B–E), no significant reduction in

gametocytemia or aberrant gametocyte development could be detected (Figure 5F). This was

re-investigated by targeted gene disruption. Likewise, deletion of these two genes as well as

of *Pf*ApiAT8 and *Pf*ApiAT10 did not result in any measurable impairment of gametocyte

development or morphology, indicating the dispensability of these individual *Pf*ApiATs for the

sexual stage development of the parasite (Figure 6, Figure S5).

DISCUSSION

We localized four putative amino acid transporters (*Pf*ApiAT2, *Pf*ApiAT4 *Pf*ApiAT8, P*f*ApiAT10)

of the ApiAT-family to the PPM in asexual blood stage parasites and gametocytes. Due to the

low expression of *Pf*ApiAT9-GFP – in agreement with the transcript levels of *apiat9* in these

stages[37,42,43] – no conclusive localization could be delivered. The observed PPM localization of

the investigated ApiATs is in concordance with published data, since i) the *P. berghei* ApiAT8

homologue was shown to be a general cationic amino acid transporter of the PPM[29,30], ii)

*Pf*ApiAT8 has recently been localized to the PPM using an overexpression approach[31], and iii)

in *T. gondii* several ApiATs (*Tg*ApiAT1, *Tg*ApiAT2, *Tg*ApiAT3-1, *Tg*ApiAT3-2, *Tg*ApiAT3-3,

*Tg*ApiAT5-3, *Tg*ApiAT6-1 and *Tg*ApiAT6-3) have been located at the PPM as well[25,29,30,32,33,44].

Recent work using episomally overexpressed *Pf*ApiAT10-GFP implied an association of this

transporter with the mitochondrial membrane[36] in *Pf*Dd2 parasites. This localization differs

from the observed PPM localization of endogenously GFP-tagged *Pf*ApiAT10 in both, 3D7 as

well as 3D7-iGP parasites (Figure 1B, 4C, S2A) reported in this study. We confirmed the PPM

localization of *Pf*ApiAT10 by its overexpression either as a GFP or mCherry fusion using two

different promoters (*crt*[45], *ama1*[46]). In line with that, the reported reduced sensitivity to the

mitochondrial electron transport chain inhibitor Atovaquone[47] upon knockout of

*Pf*ApiAT10[36], was not observed upon targeted gene disruption in our study (Figure S2B). It is

possible but appears unlikely that the reported mitochondrial association as well as reduced

sensitivity to atovaquone upon overexpression is due to the different parasite strains

(*Pf*Dd2[48] versus 3D7 this study), given that *PfDd2_apiat10* (PFDd2_030017500) has only one

silent mutation at position G1080A compared to *Pf3D7_apiat10*[49].

During the intraerythrocytic development of the parasite, the amino acid requirements are

largely covered by degradation of the globin polypetide[14,50], although – for instance – the

import of isoleucine is crucial for the survival of the parasite as *P. falciparum* lacks the

canonical pathways for its biosynthesis[51] and adult human hemoglobin lacks this amino acid.

Dedicated amino acid transporter could fill this gap. Therefore, we tested the impact of

functional inactivation of individual *Pf*ApiATs on parasite growth in asexual and sexual blood

stages of *P. falciparum*. We only observed a minor but significant reduction in parasite

growth upon knockdown or gene disruption of *Pf*ApiAT2 and *Pf*ApiAT4 in asexual blood

stages without compensatory up-regulation of other *Pf*ApiATs on the transcriptional level, as

indicated by qPCR analysis. The phenotypes are in agreement with a previously reported

reduced parasite multiplication rate of 36 % in *Pb*ApiAT4 knockout parasites[35]. Moreover our

data is also in line with the finding that *Pb*ApiAT8 is not essential for asexual

replication[29,30,35,52]. This might be explained by functional redundancy either within the ApiAT

family, as recently observed for *T. gondii*[34], or by the presence of yet unassigned transporters

capable of transporting essential amino acids such as isoleucine across the PPM. Like in *T.*

gondii, overlapping substrate specificities and lower transport levels might be sufficient for

parasite growth *in vitro*[34]. Of note, in *T. gondii* the arginine transporter *Tg*ApiAT1 has been

shown to be differently regulated on the translational level in dependence of arginine

mediated by an upstream open reading frame (uORF) present on the 5' leader sequence of

the transcript[26]. A similar layer of regulation might also be present in *Pf*ApiATs. However,

since *Toxoplasma* and *Plasmodium* only share one ApiAT, the likely most ancestral ApiAT2[25],

the regulatory elements as well as the general characteristics and substrates of the different

*Pf*ApiATs remain largely unknown.

Interestingly, a knockout of the ApiAT8 homologue of *P. berghei* (PBANKA_0208300) resulted

in strongly reduced number of mature gametocytes with an aberrant morphology of the

remaining parasites[30] and strongly reduced exflagellation[35]. In our study, functional

inactivation of *Pf*ApiAT8 via targeted gene disruption had no impact on gametocyte

development and morphology, which might reflect the pronounced differences in
gametocyte development between the rodent infecting *P. berghei* and the human infecting
P. falciparum parasites[53]. For future work, it will be interesting to target multiple ApiATs by
gene disruption in parallel to assess their putative synergy and to probe into *Pf*ApiAT function
in other fast-replicating stages of *P. falciparum* such as liver stages, for which the essentiality
of several metabolic processes has recently been shown and that primarily rely on the amino
acid uptake from their host[54].

METHODS

P. falciparum culture

Blood stages of *P. falciparum* 3D7[55] were cultured in human red blood cells (O+ or B+).
Cultures were maintained at 37°C in an atmosphere of 1 % O_2, 5 % CO_2 and 94 % N_2 using
RPMI complete medium containing 0.5 % Albumax according to standard protocols[56]. In
order to obtain highly synchronous parasite cultures late schizonts were isolated by percoll
gradient[57] and cultured with fresh erythrocytes for 4 hours. Afterwards sorbitol
synchronization[58] was applied in order to remove remaining schizonts resulting in a highly
synchronous ring stage parasite culture with a four-hour age window.
Induction of gametocytogenesis was done as previously described[41,59]. Briefly, GDV1-GFP-DD
expression was achieved by addition of 2 or 4 µM Shield-1 to the culture medium and
gametocyte cultures were treated with 50 mM N-acetyl-D-glucosamine (GlcNAc) for five days
starting 72 h post Shield-1 addition to eliminate asexual parasites[60]. Alternatively, asexual
ring stage cultures with >10 % parasitemia, cultured in the presence of choline, were
synchronized with Sorbitol[58] and washed twice in choline-free RPMI medium. Cells were kept
in choline-free medium for the entirety of the assay. After one reinvasion cycle cultures at
trophozoite stage were treated with 50 mM N-acetyl-D-glucosamine (GlcNAc)[60] and kept on
this for five days. Gametocytes were maintained in RPMI complete medium containing 0.25
% Albumax and 0.25 % sterile filtered human serum (Interstate Blood Bank, Inc. Memphis,
TN, USA).
Growth assays in low amino acid medium were performed using amino acid restricted RPMI
medium prepared as previously described[20]. Briefly, complete medium was added in a 1/20

dilution to amino acid-free RPMI medium 1640 (US Biological, R9010-01). This resulted in a

1:20 of the concentration of every amino acid compared to the standard RPMI complete

based medium.

Cloning of plasmid constructs for parasite transfection

For endogenous tagging using the SLI system[38] and glmS based conditional knockdown[40] a

855 (*Pf*ApiAT4/MFR5/PF3D7_1129900) and 1001 bp (*Pf*ApiAT2/MFR4/PF3D7_0914700)

homology region was amplified using 3D7 gDNA and cloned into pSLI-PIC1-GFP-glmS[61] using

the NotI/MluI restriction site.

For endogenous GFP-tagging a 886 bp (*Pf*ApiAT8/NPT1/PF3D7_0104800), 953 bp

(*Pf*ApiAT9/MFR2/PF3D7_0104700), 881 bp (*Pf*ApiAT10/MFR3/PF3D7_0312500) homology

region was amplified using 3D7 gDNA and cloned into pSLI-GFP[38] using the NotI/MluI

restriction site.

For SLI-based targeted gene disruption (SLI-TGD) a 396 bp (*Pf*ApiAT4, *Pf*ApiAT8), 402 bp

(*Pf*ApiAT9), 393 bp (*Pf*ApiAT10) and 435 bp (*Pf*ApiAT2) homology region was amplified using

3D7 gDNA and cloned into the pSLI-TGD plasmid[38] using NotI and MluI restriction sites.

For overexpression constructs the full length *Pf*ApiAT10 sequence was amplified from

parasite gDNA and cloned into pARL-ama1[46]-AIP-mCherry-yDHODH[61] using the XhoI/KpnI

restriction site or into the pARL-crt[45]-PF3D7_0324600-GFP-hDHFR[37] plasmid using the

KpnI/AvrII restriction site. Oligonucleotides used to generate the DNA fragments are

summarized in Table S1.

For co-localization experiments the plasmid pLyn-FRB-mCherry[38] was used.

Western blot analysis

Immunoblots were performed using saponin-lysed, infected erythrocytes. Parasite proteins

were separated on a 12 % SDS-PAGE gel as described previously[62,63] and transferred to a

nitrocellulose membrane (Amersham Protran; 0.45-µm pore nitrocellulose membrane; GE

Healthcare) using a Trans-Blot device (Bio-Rad) according to the manufacturer's instructions.

The membranes were blocked with 3 % skim milk in TBS for 30 minutes and then probed with

mouse anti-GFP (clone 7.1 and 13.1; 1:1,000, Roche) or rabbit anti-aldolase[64] (1:2,000). The

chemiluminescent signal of the horseradish peroxidase-coupled secondary antibodies

(Dianova) was visualized using a Chemi Doc XRS imaging system (Bio-Rad) and processed with

Image Lab 5.2 software (Bio-Rad).

To perform loading controls and ensure equal loading of parasite material, rabbit anti-

aldolase[64] antibodies were used. The corresponding immunoblots were incubated twice in

stripping buffer (0.2 M glycine, 50 mM dithiothreitol, 0.05 % Tween 20) at 55°C for 1 h and

washed 3 times with Tris-buffered saline for 10 min before re-probing.

Transfection of *P. falciparum*

For transfection, Percoll-purified[57] parasites at late schizont stage were transfected with 50

µg plasmid DNA using Amaxa Nucleofector 2b (Lonza, Switzerland) as previously described[65].

Transfectants were selected using either 4 nM WR99210 (Jacobus Pharmaceuticals), 0.9 µM

DSM1[66] (BEI Resources), or 2 µg/mL blasticidin S (Life Technologies, USA). In order to select

for parasites carrying the genomic modification via the SLI system[38], G418 (ThermoFisher,

USA) at a final concentration of 400 µg/mL was added to a culture with about 5 %

parasitemia. The selection process and integration test were performed as previously

described[38].

Imaging

All fluorescence images were captured using a Zeiss Axioskop 2plus microscope with a

Hamamatsu Digital camera (Model C4742-95) or a Leica D6B fluorescence microscope

equipped with a Leica DFC9000 GT camera and a Leica Plan Apochromat 100x/1.4 oil

objective.

Microscopy of live parasite-infected erythrocytes was performed as previously described[67].

Briefly, parasites were incubated in standard culture medium with 1 µg/mL Hoechst-33342

(Invitrogen) for 15 minutes at 37°C prior to imaging. 5.4 µL of infected erythrocytes were

added on a glass slide and covered with a cover slip. Nuclei were stained with 1 µg/mL

Hoechst-33342 (Invitrogen). Mitochondria were visualized by incubation of parasites with 20

nM MitoTracker Red 665 CMXRos (Invitrogen) for 15 min at 37°C prior to imaging. Images

were processed using Fiji[68] and Adobe Photoshop CC 2021 was used for display purposes

only.

Growth assay

For growth assays of TGD cell lines a flow cytometry assay, adapted from previously

published assays[69,70], was performed to measure proliferation over five days. For growth

under low amino acid conditions TGD and wild type cell lines were cultured in parallel in

standard and amino acid-depleted medium for five days. Each day parasite cultures were

resuspended and 20 µL samples were transferred to an Eppendorf tube. 80 µL RPMI

containing Hoechst-33342 and dihydroethidium (DHE) was added to obtain final

concentrations of 5 µg/mL and 4.5 µg/mL, respectively. Samples were incubated for 20 min

(protected from UV light) at room temperature, and parasitemia was determined using an

LSRII flow cytometer by counting 100,000 events using the FACSDiva software (BD

Biosciences) or using an ACEA NovoCyte flow cytometer.

Gametocyte quantification assay

Giemsa-stained blood smears at day 10 post induction of GDV1 expression were obtained

and at least 10 fields of view were recorded using a 63x objective per treatment and time

point. Erythrocyte numbers were then determined using the automated Parasitemia

software (http://www.gburri.org/parasitemia/) while the number of gametocytes was

determined manually in >700 erythrocytes per sample.

GlmS-based knockdown

GlmS-based knockdown assay was adapted from previously published assays[40,71]. To induce

knockdown, highly synchronous early ring stage parasites were split in two dishes, 2.5 mM

glucosamine was added to one of them and parasite growth was measured by FACS after two

and four parasite replication cycles. Parasite cultures were inspected daily by Giemsa smears

and, if necessary, diluted to avoid growth bias caused by high parasitemia. As an additional

control, the same amount of glucosamine was also added to 3D7 wildtype parasites. For all

analyses, medium was changed daily, and fresh glucosamine was added every day. GlmS-
based knockdown assay was adapted from previously published assays[40,71]. To induce
knockdown, highly synchronous early ring stage parasites were split into two dishes, 2.5 mM
glucosamine was added to one of them and parasite growth was measured by FACS after two
and four parasite replication cycles. Parasite cultures were inspected daily by Giemsa smears
and, if necessary, diluted to avoid growth bias caused by high parasitemia. As an additional
control, the same amount of glucosamine was also added to 3D7 wildtype parasites. For all
analyses, medium was changed daily, and fresh glucosamine was added every day.
Knockdown was quantified by fluorescence live cell microscopy using schizonts about 40 h
post glucosamine treatment. Parasites with similar size were imaged, and fluorescence was
captured with the same acquisition settings to obtain comparable measurements of the
fluorescence intensity. Fluorescence intensity (integrated density) was measured with Fiji[68],
and background was subtracted in each image. The data was visualized with GraphPad Prism
version 8 (GraphPad Software, USA).
For knockdown experiments in gametocytes synchronized ring stage cultures were induced
by the addition of Shield-1, as described above. At day 3 post induction the culture was spilt
into two dishes and one dish was cultured in the presence of 2.5 mM glucosamine for the
remaining ten days. Knockdown was quantified by fluorescence live cell microscopy at day 7
and 10 post induction, as described above.
404
405 **Drug assays**
406
Drug assays were adapted from previously described assays[72–74]. Briefly, 3D7-iGP and 3D7-
iGP-ApiAT10-TGD parasites were synchronized to a 4 h time window resulting in 0–4 h ring
stage parasites. At 24 hpi, parasitemia was determined by flow cytometry and the drug
susceptibility assays were set up in black 96-well microtiter plates (Thermo Scientific) with 0.1
% starting parasitemia and 2 % hematocrit in a final volume of 200 µl. In each plate, infected
erythrocytes in the absence of drugs treated with DMSO only served as positive controls,
while uninfected RBCs served as negative controls (for background subtraction). Parasites
were incubated with varying concentrations of dihydroartemisinin (DHA) (Adipogen,
Switzerland, AG-CN2-0468) (0–50 nM) and Atovaquone (Cayman, Item No.23802) (0–16 nM).

After 96 h of incubation, parasite growth was determined by measuring the fluorescence of

SYBR Gold (Invitrogen). Therefore, 100 µl/well supernatant was discarded without disturbing

the RBC layer and 100 µl of lysis buffer (20 mM Tris, 0.008 % saponin, 0.08 % Triton X-100, 1x

SYBR Gold) was added to each well. Plates were incubated in the dark for 2 h at room

temperature before measuring fluorescence using the EnVision Multimode Plate Reader

(PerkinElmer), as described previously[74]. In order to calculate IC_{50} values, the measured

values were normalized to the uninfected erythrocytes and plotted in GraphPad Prism

version 8 (GraphPad Software, USA) as % of DMSO control. Dose-response curves were

generated using nonlinear regression (curve fit > dose-response inhibition > (log) inhibitor vs.

normalized response—variable slope).

Quantitative real-time PCR (qPCR)

Parasites at different time-points (8, 16, 32 and 44 hpi) were harvested for 3D7-ApiAT2-TGD,

3D7-ApiAT4-TGD, 3D7-ApiAT8-TGD, 3D7-ApiAT10-TGD and 3D7-WT to obtain RNA samples

for quantitative real-time PCR. Highly synchronous ring stage parasite cultures were grown

for another 40 h and TRIzol samples were harvested in the following cycle. Volumes of

prewarmed TRIzol used for infected erythrocyte lysis and storage of RNA samples depended

on the parasite stage: ring stages were lysed in 5x volumes, trophozoites in 10x volumes and

schizonts in 20x volumes of the settled cell pellet. RNA was purified and checked for absence

of genomic DNA. cDNA synthesis with random hexamers and quantitative real-time PCR was

performed exactly as previously described[75]. Primers for each of the *apiat* genes, for genes to

control for parasite stages (*sbp1*[76]*, tom22, ama1*) and for housekeeping genes (*arginyl-tRNA*

synthetase[75]*, fructose bisphosphate aldolase*[77]) are listed in Table S1. Amplification

efficiencies of the primer pairs were determined over six log10 dilutions of gDNA (10 ng –

0.0001 ng) and were shown to have similar values between 1.915 and 2.001 (Table S1).

Expression of *apiat* genes and controls were analyzed in relation to expression of the *arginyl-*

tRNA synthetase gene (normalizer).

Software

Schematic protein representations were designed using IBS[78], predicted protein domains

were obtained from plasmodDB[79] inferred from TMHMM[80]. Parasite icons were generated

using BioRender (biorender.com). Statistical analyses were performed with GraphPad Prism

version 8 (GraphPad Software, USA).

Acknowledgements

We thank Till Voss for 3D7-iGP parasites, Jacobus Pharmaceuticals for WR99210 and Greg

Burri for the parasitemia software. DSM1 (MRA-1161) was obtained from MR4/BEI

Resources, NIAID, NIH.

Author contribution

Conceptualization: JSW, JS, TWG, AB; Methodology: JSW, EP, PCB, PMR, TS, JS, TWG, AB;

Investigation: JSW, CVG, GF, JMR, EP, JLF, HVT, SS, PCB; Formal Analysis: JS, AB; Writing

original manuscript: JSW, AB; Review & Editing: JSW, TWG, AB; Funding Acquisition: TWG, AB,

JS; Resources: PMR, TS; Project Administration: TWG, AB; Supervision: TWG, AB. All authors

read and approved the manuscript.

Funding

AB and JSW were funded by the German Research Foundation (DFG) grant BA 5213/3-1. This

project was also supported by Partnership of Universität Hamburg and DESY (PIER) project ID

PIF-2018-87 and by a Joachim Herz Foundation Project grant. JLF is supported by a HFSPO

long-term postdoctoral fellowship (LT000024/2020-L). PCB was funded by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) – project number 414222880.

Figures

Figure 1: *P. falciparum* ApiATs localize to the parasite plasma membrane (PPM) during asexual

blood stage development. (A) Heatmap of RNA-seq gene expression profiles[37] of *Pf*ApiAT2,

*Pf*ApiAT4, *Pf*ApiAT8, *Pf*ApiAT9 and *Pf*ApiAT10 during the asexual blood stage development.

Timepoints indicated as hours post infection (hpi) plus merozoites (m). **(B)** Localization of

*Pf*ApiAT2-GFP, *Pf*ApiAT4-GFP, *Pf*ApiAT8-GFP, *Pf*ApiAT9-GFP and *Pf*ApiAT10-GFP by live-cell

microscopy across the IDC of 3D7 parasites. **(C)** Co-localization of the GFP-tagged ApiAT

fusion proteins with the PPM marker protein lyn-mCherry in schizonts and merozoites. **(D)**

Live-cell microscopy of 3D7-crt-ApiAT10-GFP parasites across the IDC. **(E)** 3D7-iGP-ApiAT10-

GFP/ama1-ApiAT10-mCherry parasites at trophozoite and schizont stage. Nuclei were stained

with Hoechst-33342. Scale bar, 2 µm. Parasite stages as indicated; ring stage (R), early

trophozoite (ET), late trophozoite (LT) and schizont (S).

Figure 2: Conditional knockdown of *Pf*ApiAT2 and *Pf*ApiAT4 reveals a minor growth defect

during asexual blood stage development. (A, B) Live-cell microscopy of **(A)** 3D7-ApiAT2-GFP

and **(B)** 3D7-ApiAT4-GFP parasites, which were treated for 40 hours with 2.5 mM

glucosamine (GLCN) or that were left untreated (control). Scale bar, 2 µm. **(C, D)**

Quantification of knockdown efficiency by measuring mean fluorescence intensity (MFI)

density and parasite size (area) of **(C)** 3D7-ApiAT2-GFP and **(D)** 3D7-ApiAT4-GFP parasites 40

hours after treatment with or without 2.5 mM glucosamine. Data is displayed as mean +/- SD

of three independent experiments and individual data points are displayed on a scatterplot

color-coded by experiments according to Superplot guidelines[81]. P-values displayed were

determined using a two-tailed unpaired t-test with Welch's correction. **(E)** Growth of

parasites treated with or without 2.5 mM GLCN after two and four parasite replication cycles

as determined by flow cytometry. Shown are relative parasitemia values, which were

obtained by dividing the parasitemia of glucosamine-treated cultures by the parasitemia of

the corresponding untreated ones. Displayed are means +/- SD of independent growth

experiments with the number of experiments (n) indicated. Adjusted p-values displayed were

determined with a two-tailed unpaired t-test with Welch's correction and using the

Benjamini-Hochberg correction afterwards accounting for multiple testing by comparing

ApiAT2-GFP-glmS or ApiAT4-GFP-glmS cultured with 2.5mM GLCN to 3D7 parasites cultured

with 2.5mM GLCN.

Figure 3: Targeted gene disruption (TGD) of *Pf*ApiAT2, *Pf*ApiAT4, *Pf*ApiAT8 and *Pf*ApiAT10

reveals the dispensability of *Pf*ApiATs for *in vitro* parasite proliferation. (A–D) Schematic

representation of the full-length and truncated protein versions (upper panel). Protein length

(number of amino acids) and putative transmembrane protein domains (blue) are indicated.

Localization of **(A)** *Pf*ApiAT8-TGD-GFP, **(B)** *Pf*ApiAT4-TGD-GFP, **(C)** *Pf*ApiAT2-TGD-GFP and **(D)**

*Pf*ApiAT10-TGD in ring, trophozoite and schizont stage parasites is shown in lower panels.

Nuclei were stained with Hoechst-33342. Scale bar, 2 µm. **(E)** Growth of 3D7-ApiAT-TGD cell

lines as percentage of 3D7 parasites growth, monitored over two intracellular development

cycles by flow cytometry. The number of independent growth experiments (n) per 3D7-

ApiAT-TGD cell line is indicated. 3D7 wildtype parasites were measured in parallel. Statistical

differences were analyzed using a one-sample t-test with Benjamini-Hochberg correction

accounting for multiple comparisons. **(F)** Growth of TGD and 3D7 cell lines cultivated in low

amino acid medium relative to their growth in standard medium is shown as percentage of

growth after two parasite replication cycles. The number of individual growth experiments

(n) performed is indicated for each 3D7-ApiAT-TGD line, additionally 3D7 wildtype parasites

were analyzed with n=9. No statistical differences were observed by comparing relative

growth of TGD cell lines to 3D7 using a two-tailed unpaired t-test with Bonferroni correction.

Figure 4: *P. falciparum* ApiATs localize to the parasite plasma membrane (PPM) during

gametocyte development. (A–D) Localization of **(A)** *Pf*ApiAT2-GFP, **(B)** *Pf*ApiAT4-GFP, **(C)**

*Pf*ApiAT8-GFP and **(D)** *Pf*ApiAT10-GFP in individual 3D7-iGP-ApiAT-GFP cell lines during

gametocyte development (stage I - V) as determined by live-cell microscopy. White arrow

heads indicate remaining GDV1-GFP signal observed in close proximity to the Hoechst signal

in the 3D7-iGP-ApiAT8-GFP cell line **(C)**, as previously described[59,82,83]. **(E)** Localization of

*Pf*ApiAT9-GFP during gametocytogenesis was assessed with the 3D7-ApiAT9-GFP cell line (see

Figure 1) upon induction with choline depletion. **(F)** Co-localization of the GFP-tagged ApiAT

fusion proteins with the PPM marker protein lyn-mCherry. Nuclei were stained with Hoechst-

33342. Scale bar, 2 µm.

Figure 5: Conditional knockdown of *Pf*ApiAT2 and *Pf*ApiAT4 reveals dispensability for

gametocyte development. (A) Schematic representation of the experimental setup. **(B, C)**

Live-cell microscopy of parasites with identical settings of **(B)** 3D7-iGP-ApiAT2-GFP and **(C)**

3D7-iGP-ApiAT4-GFP stage I – V gametocytes. Scale bar, 2 µm. **(D, E)** Quantification of

knockdown by measuring mean fluorescence intensity (MFI) density and size (area) of **(D)**

3D7-iGP-ApiAT2-GFP **(E)** 3D7-iGP-ApiAT4-GFP parasites at day 7 and day 12 post induction of

gametocytogenesis cultured either with or without (control) 2.5 mM GLCN. Scale bar, 2 μm.

Data are displayed as mean +/- SD of three (3D7-iGP-ApiAT2-GFP) or four (3D7-iGP-ApiAT4-

GFP) independent experiments and individual data points are displayed on a scatterplot

color-coded by experiments according to Superplots guidelines[81]. P-values displayed were

determined with a two-tailed unpaired t-test with Welch's correction. **(F)** For each condition

gametocytemia at day 10 post gametocyte induction was determined by counting between

702-7693 (mean 2210) cells per condition in Giemsa-stained thin blood smears. The relative

gametocytemia values (%) displayed were obtained by dividing the gametocytemia of

glucosamine-treated cultures by the gametocytemia of the corresponding untreated

cultures. Displayed are means +/- SD of independent growth experiments with the number of

experiments (n) indicated. A two-tailed unpaired t-test with Welch's and Benjamini-Hochberg

correction was used to calculate multiplicity adjusted p-values for ApiAT2-GFP-glmS or

ApiAT4-GFP-glmS versus 3D7-iGP parasites all cultured with 2.5mM GLCN.

Figure 6: Targeted gene disruption (TGD) of *Pf*ApiAT2, *Pf*ApiAT4, *Pf*ApiAT8 and *Pf*ApiAT10

reveals dispensability of ApiATs for gametocyte development. Representative images from two

(*Pf*ApiAT2, *Pf*ApiAT4) or three (*Pf*ApiAT8, *Pf*ApiAT10) independent experiments derived from

Giemsa-stained thin blood smears of gametocyte stages I-V of 3D7-iGP, 3D7-iGP-ApiAT2-TGD,

3D7-iGP-ApiAT4-TGD, 3D7-iGP-ApiAT8-TGD, 3D7-iGP-ApiAT10-TGD parasites. Scale bar, 5 μm.

Figure S1: PCR analysis and western blots of 3D7-ApiAT-GFP and 3D7-ApiAT-TGD cell lines. (A)

PCR-based analysis of unmodified wildtype (WT) and transgenic knock-in (KI) cell lines (3D7-

ApiAT2-GFP-glmS, 3D7-ApiAT4-GFP-glmS, 3D7-ApiAT8-GFP, 3D7-ApiAT9-GFP, 3D7-ApiAT10-

GFP). Specific genomic modifications resulting from correct integration of the respective SLI-

based vectors were tested targeting the 3'- and 5'-end of the locus. **(B)** Western blot analysis

of wildtype (3D7) and knock-in (KI) cell lines (3D7-ApiAT2-GFP-glmS, 3D7-ApiAT4-GFP-glmS,

3D7-ApiAT8-GFP, 3D7-ApiAT10-GFP) using mouse anti-GFP to detect the tagged full-length

protein (upper panel) and rabbit anti-aldolase to control for equal loading (lower panel).

Protein sizes are indicated in kDa. **(C)** PCR-based analysis of unmodified wildtype (WT) and

transgenic parasites modified by targeted gene disruption (TGD).

574 **Figure S2: *Pf*ApiAT10 is PPM localized and its functional inactivation has no influence on**

575 Atovaquone sensitivity. (A) Live-cell microscopy of 3D7-iGP-ApiAT10-GFP parasites stained

576 with MitoTracker Red CMXRos across the IDC and in gametocytes. Stages as indicated: R =

577 ring stage, ET = early trophozoite, LT = late trophozoite, ES = early schizont, LS = late schizont,

578 G = gametocyte; Nuclei were stained with Hoechst-33342. Scale bar, 2 μm. **(B)** Drug

579 susceptibility assays of 3D7-iGP and 3D7-iGP-ApiAT10-TGD parasites were performed with

580 Atovaquone (left) and dihydroartemisinin (DHA, right). Parasite growth was determined by

581 measuring the DNA content using SYBR gold when exposed to varying concentrations of

582 drugs for 96 h. The growth of DMSO-treated control parasites was set to 100%. Shown are

583 means +/- SD of 4 or 5 independent biological replicates performed in technical duplicates.

584 Calculated IC_{50} values with 95% confidence intervals are shown above each graph.

586 **Figure S3: Quantitative real-time PCR of individual *Pf*ApiAT-encoding genes and controls in**

587 **3D7-ApiAT-TGD and wildtype parasites.** Expression of *apiat* genes and control genes (*sbp1*,

588 *tom22*, *ama1*, *fructose-bisphosphate aldolase*) normalized to the housekeeping control

589 *arginyl-tRNA synthetase* over four different time points during intraerythrocytic development

590 cycle of *P. falciparum* parasites. Bars with individual measurements from n=2 (3D7-ApiAT2-

591 TGD, 3D7-ApiAT4-TGD) or n=3 (3D7-ApiAT8, 3D7-ApiAT10, 3D7 wildtype) biological

592 replicates. Gene accession numbers, primer sequences and amplification efficiencies are

593 listed in Table S1.

595 **Figure S4: PCR analysis of 3D7-iGP-ApiAT-GFP cell lines and Giemsa smears from *Pf*ApiAT2 and**

596 ***Pf*ApiAT4 conditional knockdown experiments. (A)** PCR-based analysis of unmodified wildtype

597 (WT) and transgenic knock-in (KI) cell lines (3D7-iGP-ApiAT2-GFP-glmS, 3D7-iGP-ApiAT4-GFP-

598 glmS, 3D7-iGP-ApiAT8-GFP, 3D7-iGP-ApiAT10-GFP). **(B)** Giemsa smears of stage I – V 3D7-iGP-

599 ApiAT2-GFP-glmS and 3D7-iGP-ApiAT4-GFP-glmS gametocytes cultured either without

600 (control) or with 2.5 mM glucosamine (GLCN). Scale bar, 5 μm. **(C)** Live cell imaging of 3D7-

601 iGP stage IV and V gametocytes. Nuclei were stained with Hoechst-33342. Scale bar, 2 μm.

602 **(D)** Live cell microscopy of 3D7-iGP-ApiAT2 stage IV and V gametocytes from Figure 5B, with

603 adjusted brightness. Scale bar, 2 μm.

Figure S5: PCR analysis of 3D7-iGP-ApiAT-TGD cell lines and Giemsa smears from 3D7-iGP-
ApiAT8-TGD stage IV and V gametocytes showing normal morphology. (A) PCR-based analysis
of unmodified wildtype (WT) and transgenic TGD cell lines (3D7-iGP-ApiAT2-TGD, 3D7-iGP-
ApiAT4-TGD, 3D7-iGP-ApiAT8-TGD, 3D7-iGP-ApiAT10-TGD). (**B**) Giemsa smears of 3D7-iGP-
ApiAT8-TGD and 3D7-iGP stage IV and V gametocytes from three independent experiments.
Scale bar, 5 µm.

Table S1: Oligonucleotides used for cloning and quantitative real-time PCR (qPCR).

References

1. Krishnan, A. & Soldati-Favre, D. Amino acid metabolism in apicomplexan parasites.
Metabolites **11**, 1–17 (2021).

2. Wrenger, C. *et al.* Specific inhibition of the aspartate aminotransferase of plasmodium
falciparum. *J. Mol. Biol.* **405**, 956–971 (2011).

3. Nagaraj, V. A. *et al.* Asparagine requirement in Plasmodium berghei as a target to
prevent malaria transmission and liver infections. *Nat. Commun.* **6**, 1–13 (2015).

4. Meireles, P. *et al.* Uptake and metabolism of arginine impact Plasmodium
development in the liver. *Sci. Rep.* **7**, (2017).

5. MacRae, J. I. *et al.* Mitochondrial metabolism of glucose and glutamine is required for
intracellular growth of toxoplasma gondii. *Cell Host Microbe* **12**, 682–692 (2012).

6. Sopitthummakhun, K. *et al.* Plasmodium serine hydroxymethyltransferase as a
potential anti-malarial target: Inhibition studies using improved methods for enzyme
production and assay. *Malar. J.* **11**, 194 (2012).

7. Spielmann, T., Gras, S., Sabitzki, R. & Meissner, M. Endocytosis in Plasmodium and
Toxoplasma Parasites. *Trends in Parasitology* **36**, 520–532 (2020).

8. Kirk, K. & Lehane, A. M. Membrane transport in the malaria parasite and its host
erythrocyte. *Biochem. J.* **457**, 1–18 (2014).

9. Divo, A. A., Geary, T. G., Davis, N. L. & Jensen, J. B. Nutritional Requirements of
Plasmodium falciparum in Culture. I. Exogenously Supplied Dialyzable Components
Necessary for Continuous Growth. *J. Protozool.* **32**, 59–64 (1985).

10. Ginsburg, H., Kutner, S., Krugliak, M. & Ioav Cabantchik, Z. Characterization of
permeation pathways appearing in the host membrane of Plasmodium falciparum

infected red blood cells. *Mol. Biochem. Parasitol.* **14**, 313–322 (1985).

11.   Kirk, K., Horner, H. A., Elford, B. C., Ellory, J. C. & Newbold, C. I. Transport of diverse

substrates into malaria-infected erythrocytes via a pathway showing functional

characteristics of a chloride channel. *J. Biol. Chem.* **269**, 3339–3347 (1994).

12.   Olszewski, K. L. *et al.* Host-Parasite Interactions Revealed by Plasmodium falciparum

Metabolomics. *Cell Host Microbe* **5**, 191–199 (2009).

13.   Liu, J., Istvan, E. S., Gluzman, I. Y., Gross, J. & Goldberg, D. E. Plasmodium falciparum

ensures its amino acid supply with multiple acquisition pathways and redundant

proteolytic enzyme systems. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 8840–8845 (2006).

14.   Babbitt, S. E. *et al.* Plasmodium falciparum responds to amino acid starvation by

entering into a hibernatory state. *Proc. Natl. Acad. Sci. U. S. A.* **109**, E3278–E3287

(2012).

15.   McLean, K. J. & Jacobs-Lorena, M. The response of Plasmodium falciparum to

isoleucine withdrawal is dependent on the stage of progression through the

intraerythrocytic cell cycle. *Malar. J.* **19**, 147 (2020).

16.   Martin, R. E. & Kirk, K. Transport of the essential nutrient isoleucine in human

erythrocytes infected with the malaria parasite Plasmodium falciparum. *Blood* **109**,

2217–2224 (2007).

17.   Cobbold, S. A., Martin, R. E. & Kirk, K. Methionine transport in the malaria parasite

Plasmodium falciparum. *Int. J. Parasitol.* **41**, 125–135 (2011).

18.   Desai, S. A., Krogstad, D. J. & McCleskey, E. W. A nutrient-permeable channel on the

intraerythrocytic malaria parasite. *Nature* **362**, 643–646 (1993).

19.   Desai, S. A. & Rosenberg, R. L. Pore size of the malaria parasite's nutrient channel.

Proc. Natl. Acad. Sci. U. S. A. **94**, 2045–2049 (1997).

20.   Mesén-Ramírez, P. *et al.* EXP1 is critical for nutrient uptake across the parasitophorous

vacuole membrane of malaria parasites. *PLoS Biol.* **17**, e3000473 (2019).

21.   Garten, M. *et al.* EXP2 is a nutrient-permeable channel in the vacuolar membrane of

Plasmodium and is essential for protein export via PTEX. *Nat. Microbiol.* 1 (2018).

doi:10.1038/s41564-018-0222-7

22.   Counihan, N. A., Modak, J. K. & de Koning-Ward, T. F. How Malaria Parasites Acquire

Nutrients From Their Host. *Front. Cell Dev. Biol.* **9**, 582 (2021).

23.   Martin, R. E. The transportome of the malaria parasite. *Biol. Rev. Camb. Philos. Soc.* **95**,

305–332 (2020).

24.   Martin, R. E., Henry, R. I., Abbey, J. L., Clements, J. D. & Kirk, K. The 'permeome' of the

malaria parasite: an overview of the membrane transport proteins of Plasmodium

falciparum. *Genome Biol.* **6**, R26 (2005).

25.   Parker, K. E. R. *et al.* The tyrosine transporter of Toxoplasma gondii is a member of the

newly defined apicomplexan amino acid transporter (ApiAT) family. *PLoS Pathog.* **15**,

e1007577 (2019).

26.   Rajendran, E. *et al.* Substrate-mediated regulation of the arginine transporter of

Toxoplasma gondii. *PLoS Pathog.* **17**, e1009816 (2021).

27.   Reddy, V. S., Shlykov, M. A., Castillo, R., Sun, E. I. & Saier, M. H. The major facilitator

superfamily (MFS) revisited. *FEBS J.* **279**, 2022–2035 (2012).

28.   Pao, S. S., Paulsen, I. T. & Saier, M. H. Major Facilitator Superfamily. *Microbiol. Mol.*

Biol. Rev. **62**, 1–34 (1998).

29.   Rajendran, E. *et al.* Cationic amino acid transporters play key roles in the survival and

transmission of apicomplexan parasites. *Nat. Commun.* **8**, 1–13 (2017).

30.   Boisson, B. *et al.* The novel putative transporter NPT1 plays a critical role in early

stages of Plasmodium berghei sexual development. *Mol. Microbiol.* **81**, 1343–1357

(2011).

31.   Blomqvist, K. *et al.* Influence of Plasmodium falciparum Calcium-Dependent Protein

Kinase 5 (PfCDPK5) on the Late Schizont Stage Phosphoproteome. *mSphere* **5**, (2020).

32.   Kloehn, J., Lunghi, M., Varesio, E., Dubois, D. & Soldati-Favre, D. Untargeted

Metabolomics Uncovers the Essential Lysine Transporter in Toxoplasma gondii.

Metabolites **11**, 476 (2021).

33.   Wallbank, B. A. *et al.* Characterisation of the *Toxoplasma gondii* tyrosine transporter

and its phosphorylation by the calcium-dependent protein kinase 3. *Mol. Microbiol.*

111, 1167–1181 (2019).

34.   Fairweather, S. J. *et al.* Coordinated Action of Multiple Transporters in the Acquisition

of Essential Cationic Amino Acids by the Intracellular Parasite Toxoplasma gondii 2 3

Running Title: Essential Cationic Amino Acid Transporters in Toxoplasma parasites.

bioRxiv 2021.06.25.450001 (2021). doi:10.1101/2021.06.25.450001

35.   Kenthirapalan, S., Waters, A. P., Matuschewski, K. & Kooij, T. W. A. Functional profiles

of orphan membrane transporters in the life cycle of the malaria parasite. *Nat.*

Commun. **7**, 10519 (2016).

36.   Rocamora, F. *et al.* PfMFR3: A Multidrug-Resistant Modulator in Plasmodium

falciparum. *ACS Infect. Dis.* acsinfecdis.0c00676 (2021).

doi:10.1021/acsinfecdis.0c00676

37.   Wichers, J. S. *et al.* Dissecting the gene expression, localization, membrane topology,

and function of the plasmodium falciparum STEVOR protein family. *MBio* **10**, e01500-

19 (2019).

38.   Birnbaum, J. *et al.* A genetic system to study Plasmodium falciparum protein function.

Nat. Methods **14**, 450–456 (2017).

39.   Inoue, T., Heo, W. Do, Grimley, J. S., Wandless, T. J. & Meyer, T. An inducible

translocation strategy to rapidly activate and inhibit small GTPase signaling pathways.

Nat. Methods **2**, 415–418 (2005).

40.   Prommana, P. *et al.* Inducible Knockdown of Plasmodium Gene Expression Using the

glmS Ribozyme. *PLoS One* **8**, e73783 (2013).

41.   Boltryk, S. D. *et al.* CRISPR/Cas9-engineered inducible gametocyte producer lines as a

valuable tool for Plasmodium falciparum malaria transmission research. *Nat. Commun.*

12, 4806 (2021).

42.   Otto, T. D. *et al.* New insights into the blood-stage transcriptome of Plasmodium

falciparum using RNA-Seq. *Mol. Microbiol.* **76**, 12–24 (2010).

43.   López-Barragán, M. J. *et al.* Directional gene expression and antisense transcripts in

sexual and asexual stages of Plasmodium falciparum. *BMC Genomics* **12**, 587 (2011).

44.   Sangaré, L. O. *et al.* Unconventional endosome-like compartment and retromer

complex in Toxoplasma gondii govern parasite integrity and host infection. *Nat.*

Commun. **7**, 1–14 (2016).

45.   Struck, N. S. *et al.* Re-defining the Golgi complex in Plasmodium falciparum using the

novel Golgi marker PfGRASP. *J. Cell Sci.* **118**, 5603–5613 (2005).

46.   Hu, G. *et al.* Transcriptional profiling of growth perturbations of the human malaria

parasite Plasmodium falciparum. *Nat. Biotechnol.* **28**, 91–98 (2010).

47.   Fry, M. *et al.* Novel anti-malarial hydroxynaphthoquinones with potent broad

spectrum anti-protozoal activity. *Parasitology* **90**, 45–55 (1985).

48.   Wellems, T., Oduola, A. M., Fenton, B., Desjardins, R. & Panton, L. Chromosome size

variation occurs in cloned Plasmodium falciparum on in vitro cultivation. *Rev. bras.*

genét **11**, 813–25 (1988).

49. Otto, T. D. *et al.* Long read assemblies of geographically dispersed Plasmodium

falciparum isolates reveal highly structured subtelomeres. *Wellcome open Res.* **3**, 52

(2018).

50. Goldberg, D. E. Complex nature of malaria parasite hemoglobin degradation.

Proceedings of the National Academy of Sciences of the United States of America **110**,

5283–5284 (2013).

51. Payne, S. H. & Loomis, W. F. Retention and loss of amino acid biosynthetic pathways

based on analysis of whole-genome sequences. *Eukaryot. Cell* **5**, 272–276 (2006).

52. Sayers, C. P., Mollard, V., Buchanan, H. D., McFadden, G. I. & Goodman, C. D. A genetic

screen in rodent malaria parasites identifies five new apicoplast putative membrane

transporters, one of which is essential in human malaria parasites. *Cell. Microbiol.* **20**,

e12789 (2018).

53. Ngotho, P. *et al.* Revisiting gametocyte biology in malaria parasites. *FEMS Microbiol.*

Rev. **43**, 401–414 (2019).

54. Stanway, R. R. *et al.* Genome-Scale Identification of Essential Metabolic Processes for

Targeting the Plasmodium Liver Stage. *Cell* **179**, 1112-1128.e26 (2019).

55. Walliker, D. *et al.* Genetic analysis of the human malaria parasite Plasmodium

falciparum. *Science (80-.).* **236**, 1661–6 (1987).

56. Trager, W. & Jensen, J. B. Continuous culture of Plasmodium falciparum: its impact on

malaria research. *Int. J. Parasitol.* **27**, 989–1006 (1997).

57. Rivadeneira, E., Wasserman, M. & Espinal, C. Separation and Concentration of

Schizonts of Plasmodium falciparum by Percoll Gradients. *J. Protozool.* **30**, 367–370

(1983).

58. Lambros, C. & Vanderberg, J. P. Synchronization of Plasmodium falciparum

Erythrocytic Stages in Culture. *J. Parasitol.* **65**, 418 (1979).

59. Filarsky, M. *et al.* GDV1 induces sexual commitment of malaria parasites by

antagonizing HP1-dependent gene silencing. *Science (80-.).* **359**, 1259–1263 (2018).

60. Ponnudurai, T., Lensen, A. H. W., Meis, J. F. G. M. & Meuwissen, J. H. E.

Synchronization of Plasmodium falciparum gametocytes using an automated

suspension culture system. *Parasitology* **93**, 263–274 (1986).

61. Wichers, J. S. *et al.* Identification of novel inner membrane complex and apical annuli

765   proteins of the malaria parasite Plasmodium falciparum. *Cell. Microbiol.* **23**, e13341

766   (2021).

767   62.   Bachmann, A. *et al.* A comparative study of the localization and membrane topology of

768   members of the RIFIN, STEVOR and PfMC-2TM protein families in Plasmodium

769   falciparum-infected erythrocytes. *Malar. J.* **14**, 274 (2015).

770   63.   Heiber, A. & Spielmann, T. Preparation of Parasite Protein Extracts and Western Blot

771   Analysis. *BIO-PROTOCOL* **4**, (2014).

772   64.   Mesén-Ramírez, P. *et al.* Stable Translocation Intermediates Jam Global Protein Export

773   in Plasmodium falciparum Parasites and Link the PTEX Component EXP2 with

774   Translocation Activity. *PLoS Pathog.* **12**, e1005618 (2016).

775   65.   Moon, R. W. *et al.* Adaptation of the genetically tractable malaria pathogen

776   Plasmodium knowlesi to continuous culture in human erythrocytes. *Proc. Natl. Acad.*

777   *Sci.* **110**, 531–536 (2013).

778   66.   Ganesan, S. M. *et al.* Yeast dihydroorotate dehydrogenase as a new selectable marker

779   for Plasmodium falciparum transfection. *Mol. Biochem. Parasitol.* **177**, 29–34 (2011).

780   67.   Grüring, C. & Spielmann, T. Imaging of live malaria blood stage parasites. *Methods*

781   *Enzymol.* **506**, 81–92 (2012).

782   68.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

783   *Methods* **9**, 676–682 (2012).

784   69.   Malleret, B. *et al.* A rapid and robust tri-color flow cytometry assay for monitoring

785   malaria parasite development. *Sci. Rep.* **1**, 118 (2011).

786   70.   Jonscher, E. *et al.* PfVPS45 Is Required for Host Cell Cytosol Uptake by Malaria Blood

787   Stage Parasites. *Cell Host Microbe* **25**, 166-173.e5 (2019).

788   71.   Burda, P.-C. *et al.* Structure-Based Identification and Functional Characterization of a

789   Lipocalin in the Malaria Parasite Plasmodium falciparum. *Cell Rep.* **31**, 107817 (2020).

790   72.   Burda, P.-C. *et al.* Global analysis of putative phospholipases in the malaria parasite

791   Plasmodium 1 falciparum reveals critical factors for parasite proliferation 2 3. *bioRxiv*

792   2021.06.28.450158 (2021). doi:10.1101/2021.06.28.450158

793   73.   Smilkstein, M., Sriwilaijaroen, N., Kelly, J. X., Wilairat, P. & Riscoe, M. Simple and

794   Inexpensive Fluorescence-Based Technique for High-Throughput Antimalarial Drug

795   Screening. *Antimicrob. Agents Chemother.* **48**, 1803–1806 (2004).

74.   Schwed, S. I., Alder, A., Gilberger, T. & Kunick, C. 4-Arylthieno[2,3-b]pyridine-2-

carboxamides Are a New Class of Antiplasmodial Agents. *Molecules* **25**, (2020).

75.   Bachmann, A. *et al.* Controlled human malaria infection with Plasmodium falciparum

demonstrates impact of naturally acquired immunity on virulence gene expression.

PLOS Pathog. **15**, e1007906 (2019).

76.   Petter, M. *et al.* Expression of P. falciparum var genes involves exchange of the histone

variant H2A.Z at the promoter. *PLoS Pathog.* **7**, (2011).

77.   Salanti, A. *et al.* Selective upregulation of a single distinctly structured var gene in

chondroitin sulphate A-adhering Plasmodium falciparum involved in pregnancy-

associated malaria. *Mol. Microbiol.* **49**, 179–191 (2003).

78.   Liu, W. *et al.* IBS: An illustrator for the presentation and visualization of biological

sequences. *Bioinformatics* **31**, 3359–3361 (2015).

79.   Aurrecoechea, C. *et al.* PlasmoDB: a functional genomic database for malaria parasites.

Nucleic Acids Res. **37**, D539–D543 (2009).

80.   Sonnhammer, E. L., von Heijne, G. & Krogh, A. A hidden Markov model for predicting

transmembrane helices in protein sequences. *Proceedings. Int. Conf. Intell. Syst. Mol.*

Biol. **6**, 175–82 (1998).

81.   Lord, S. J., Velle, K. B., Mullins, R. D. & Fritz-Laylin, L. K. SuperPlots: Communicating

reproducibility and variability in cell biology. *J. Cell Biol.* **219**, (2020).

82.   Tibúrcio, M. *et al.* A 39-Amino-Acid C-Terminal Truncation of GDV1 Disrupts Sexual

Commitment in Plasmodium falciparum. *mSphere* **6**, (2021).

83.   Eksi, S. *et al.* Plasmodium falciparum Gametocyte Development 1 (Pfgdv1) and

Gametocytogenesis Early Gene Identification and Commitment to Sexual

Development. *PLoS Pathog.* **8**, (2012).

Figure 1