Trypanosomal variant surface glycoprotein expression in human African trypanosomiasis patients

Jaime So[1]**, Sarah Sudlow[1]**, Abeer Sayeed[1], Tanner Grudda[1], Stijn Deborggraeve[2]#, Dieudonné Mumba Ngoyi[3], Didier Kashiama Desamber[4], Bill Wickstead[5], Veerle Lejon[6], and Monica R. Mugnier[1]*

[1]Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, United States of America

[2]Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

[3]Department of Parasitology, Institut National de Recherche Biomédicale, Kinshasa, Democratic Republic of the Congo

[4]Programme Nationale de Lutte contre la Trypanosomiase Humaine Africaine, (PNLTHA), Ministry of Health, Kinshasa, Democratic Republic of the Congo

[5] School of Life Sciences, Queen's Medical Centre, University of Nottingham, Nottingham, NG7 2UH, United Kingdom

[6] UMR-177 Intertryp, Institut de Recherche pour le Développement, Centre de Coopération Internationale en Recherche Agronomique pour le Développement, University of Montpellier, Montpellier, France

**These authors contributed equally to the work
Current address: Médecins Sans Frontières - Access Campaign, Geneva, Switzerland

* Corresponding author
E-mail: mmugnie1@jhu.edu (MM)

Abstract

Trypanosoma brucei gambiense, an extracellular protozoan parasite, is the primary causative agent of human African trypanosomiasis. *T. b. gambiense* is endemic to West and Central Africa, where it is transmitted by the bite of infected tsetse flies. In the bloodstream of an infected host, the parasite evades antibody recognition by altering the Variant Surface Glycoprotein (VSG) that forms a dense coat on its cell surface through a process known as antigenic variation. Each VSG has a variable N-terminal domain exposed to the host and a less variable C-terminal domain that is partially hidden from host antibodies. Our lab developed VSG-seq, a targeted RNA-seq method, to study *VSG* gene expression in *T. brucei*. Studies using VSG-seq to characterize antigenic variation in a mouse model have revealed marked diversity in *VSG* expression within parasite populations, but this finding has not yet been validated in a natural human infection. Here, we used VSG-seq to analyze *VSGs* expressed in the blood of 12 patients infected with *T. b. gambiense*. The number of *VSGs* identified per patient ranged from one to 14; notably, two *VSGs* were shared by more than one patient. Analysis of expressed *VSG* N-terminal domain types revealed that 82% of expressed *VSGs* encoded a type B N-terminus, a bias not seen in datasets from other *T. brucei* subspecies. C-terminal types in *T. b. gambiense* infection were also restricted. These results demonstrate a bias either in the underlying *VSG* repertoire of *T. b. gambiense* or in this subspecies' expression of *VSGs* during infection. This work demonstrates the feasibility of using VSG-seq to study antigenic variation in human infections and highlights the importance of understanding *VSG* repertoires in the field.

Author Summary

Human African Trypanosomiasis is a neglected tropical disease primarily caused by the extracellular parasite *Trypanosoma brucei gambiense*. To avoid elimination by the host, these parasites repeatedly replace their dense surface coat of Variant Surface Glycoprotein (VSG). Despite the important role of VSGs in prolonging infection, *VSG* expression during natural human infections is poorly understood. A better understanding of natural *VSG* gene expression dynamics can clarify the mechanisms that *T. brucei* uses to alter its VSG coat and improve how trypanosomiasis is diagnosed in humans. We analyzed the expressed *VSGs* detected in the blood of patients with trypanosomiasis. Our findings indicate that parasites in both natural and experimental infections express diverse sets of *VSGs.*

Introduction

Human African Trypanosomiasis (HAT) is caused by the protozoan parasite *Trypanosoma brucei*. *T. brucei* and its vector, the tsetse fly, are endemic to sub-Saharan Africa [1]. There are two human-infective *T. brucei* subspecies: *T. b. gambiense,* which causes chronic infection in West and Central Africa (~98% of cases), and *T. b. rhodesiense,* which causes acute infection in East and Southern Africa (~2% of cases) [2,3]. In humans, infections progress from an early stage, usually marked by a fever and body aches, to a late stage associated with severe neurological symptoms that begins when the parasite crosses the blood-brain barrier [4]. HAT is considered fatal without timely diagnosis and treatment. While around 50 million people are at risk of infection [5], the number of annual human infections has declined significantly in recent years, with only 864 cases reported in 2019 [6]. The World Health Organization is working towards zero human transmissions of HAT caused by *T. b. gambiense* (gHAT) by 2030 [7]. Current public health initiatives to control the disease depend on case detection and treatment, complemented with vector control.

Prospects for developing a vaccine are severely confounded by the ability of African trypanosomes to alter their surface antigens [8]. As *T. brucei* persists extracellularly in blood, lymph, and tissue fluids, it is constantly exposed to host antibodies [9–12]. The parasite periodically changes its dense Variant Surface Glycoprotein (VSG) coat to evade immune recognition. This process, called antigenic variation, relies on a vast collection of thousands of VSG-encoding genes[13–16]. *T. brucei* also continually expands the number of usable antigens by constructing mosaic *VSGs* through one or more recombination events between individual *VSG* genes [17,18].

The VSG contains two domains: a variable N-terminal domain of ~350-400 amino acids, and a less variable C-terminal domain of ~40-80 amino acids, characterized by one or two conserved groups of four disulfide-bonded cysteines [13,19]. On the surface of trypanosomes, the VSG N-terminal domain is readily exposed to the host. In contrast, the C-terminal domain is proximal to the plasma membrane and largely hidden from host antibodies [20–22]. The N-terminal domain is classified into two types, A and B, each further distinguished into subtypes (A1-3 and B1-2), while the C-terminal domain has been classified into six types (1-6) [13,19]. These classifications are based on protein sequence patterns anchored by the conservation of cysteine residues, but the biological implications of VSG domain types have not been investigated.

Little is known about how the large genomic repertoire of *VSGs* is used in natural infections; the number and diversity of *VSGs* expressed by wild parasite populations remain unknown. One VSG in particular, LiTat 1.3, has been identified as an antigen against which many gHAT patients have antibodies [23]. As a result, LiTat 1.3 is the main target antigen in the primary serological screening tool for gHAT: the card agglutination test for trypanosomiasis (CATT/*T.b. gambiense*) [24]. Despite the widespread use of the CATT to screen for gHAT, there are shortcomings. Most notably, it relies on immunity against a variable antigen, and some *T. b. gambiense* strains lack the LiTat 1.3 gene entirely [25,26]. More recently developed rapid diagnostic tests use a

119 combination of native LiTat1.3 and another VSG, LiTat1.5 [27,28], or the combination of
120 a VSG with the invariant surface glycoprotein ISG 65 [29]. Currently, there is no
121 serological test for diagnosis of infection with *T. b. rhodesiense.*
122
Given the role of VSGs during infection and their importance in gHAT screening tests, a
better understanding of *VSG* expression dynamics could inform the development of
improved screening tests while providing insight into the molecular mechanisms of
antigenic variation. We developed VSG-seq, a targeted RNA-sequencing method that
identifies the *VSGs* expressed in any population of *T. brucei* and measures the
prevalence of each VSG in the population [30]. In a proof-of-principle study, we used
VSG-seq to gain insight into the number and diversity of *VSGs* expressed during
experimental mouse infections [30]. This proof-of-principle study revealed significant
VSG diversity within parasite populations in each animal, with many more variants
expressed at a time than the few thought to be sufficient for immune evasion. This
diversity suggested that the parasite's genomic VSG repertoire might be insufficient to
sustain a chronic infection, highlighting the potential importance of the recombination
mechanisms that form new VSGs[13,17]. The study also showed that many of the same
VSGs were expressed in different infections, supporting previous reports of a "semi-
predictable" order to *VSG* switching in *T. brucei* culture and animal models of infection
[17,31,32].
139
Recently, similar high-throughput sequencing methods have been used to characterize
antigenic variation in experimental infections of natural hosts for two related African
trypanosome species, *T. vivax* and *T. congolense* [33–35]. These studies suggest that
the mechanism for antigen production in some animal parasites may be different from
the *T. brucei* model. It remains unknown whether findings from studies of antigenic
variation using experimental models translate to natural *T. brucei* infections. To our
knowledge, only one study has investigated *VSG* expression in wild *T. brucei* isolates
[36]. For technical reasons, this study relied on RNA isolated from parasites passaged
through small animals after collection from the natural host. As *VSG* expression may
change during passage, the data obtained from these samples is somewhat difficult to
interpret. To better understand the characteristics of antigenic variation in natural *T.*
brucei infections, we sought to analyze *VSG* expression in *T. brucei* field isolates from
which RNA was directly extracted.
153
In the present study, we used VSG-seq to determine the number and diversity of *VSGs*
expressed by *T. b. gambiense* in the blood of 12 patients with a *T. b. gambiense*
infection. To complement these data, we also analyzed published datasets from both
experimental mouse infections and *T. b. rhodesiense* patients. Our analysis revealed
diverse expression of *VSGs* in natural *T. brucei* infections and distinct biases in *VSG*
expression that may be unique to the *T. b. gambiense* subspecies.

Methods

Ethics statement

The blood specimens from *T.b. gambiense* infected patients were collected within the projects, "Longitudinal follow-up of CATT seropositive, trypanosome negative individuals (SeroSui)" and "An integrated approach for identification of genetic determinants for susceptibility for trypanosomiasis (TrypanoGEN)" [37]. In France, the SeroSui study received approval from the Comité Consultatif de Déontologie et d'Ethique (CCDE) of the French National Institute for Sustainable Development Research (IRD), May 2013 session. In Belgium, the study received approval from the Institutional Review Board of the Institute of Tropical Medicine (reference 886/13) and the Ethics Committee of the University of Antwerp (B300201318039). In the Democratic Republic of the Congo, the projects SeroSui and TrypanoGEN were approved by the Ministry of Health through the Ngaliema Clinic of Kinshasa (references 422/2013 and 424/2013). Participants gave their written informed consent to participate in the projects. For minors, additional written consent was obtained from their legal representative.

Patient enrollment and origin map

Patients originated from the Democratic Republic of the Congo (DR Congo) and were identified over six months in the second half of 2013. This identification occurred either during passive screening at the center for HAT diagnosis and treatment at the hospital of Masi Manimba, or during active screening by the mobile team of the national sleeping sickness control program (PNLTHA) in Masi Manimba and Mosango health zones (Kwilu province, DR Congo).

Individuals were screened for the presence of specific antibodies in whole blood with the CATT test. For those reacting blood positive in CATT, we also tested twofold serial plasma dilutions of 1/2-1/32 were also tested and determined the CATTCATT end titer was determined. CATTCATT positives underwent parasitological confirmation by direct microscopic examination of lymph (if enlarged lymph nodes were present), and examination of blood by the mini-anion exchange centrifugation technique on buffy coat [38]. Individuals in whom trypanosomes were observed underwent lumbar puncture. The cerebrospinal fluid was examined for white blood cell count and the presence of trypanosomes to determine the disease stage and select the appropriate treatment. Patients were questioned about their place of residence. The geographic coordinates of their corresponding villages were obtained from the Atlas of HAT [39] and plotted on a map of the DR Congo using ArcGIS® software by Esri. Distances were determined and a distance matrix generated (see Supplemental Table 2).

Patient blood sample collection and total RNA isolation

A 2.5 mL volume of blood was collected from each patient in a PAXgene Blood RNA Tube. The blood was mixed with the buffer in the tube, aliquoted in 2 mL volumes and frozen in liquid nitrogen for a maximum of two weeks. After arrival in Kinshasha, tubes

205 were stored at -70°C. Total RNA was extracted and isolated from each blood sample as
206 previously described [40].
207
208
Estimation of parasitemia
Two approaches were used to estimate parasitemia. First, a 9 mL volume of blood on
heparin was centrifuged, 500 microliters of the buffy coat were taken up and
trypanosomes were isolated using the mini-anion exchange centrifugation technique.
After centrifugation of the column eluate, the number of parasites visible in the tip of the
collection tube were estimated. Second, Spliced Leader (SL) RNA expression levels
were measured by real-time PCR as previously described [40]. A Ct value was
determined for each patient blood sample. Real-time PCR was performed on RNA
samples before reverse transcription to verify the absence of DNA contamination.
218
219
RNA sequencing
DNase I-treated RNA samples were cleaned up with 1.8x Mag-Bind TotalPure NGS
Beads (Omega Bio-Tek, # M1378-01). cDNA was generated using the SuperScript III
First-strand synthesis system (Invitrogen, 18080051) according to manufacturer's
instructions. 8 microliters of each sample (between 36 and 944 ng) were used for cDNA
synthesis, which was performed using the oligo-dT primer provided with the kit. This
material was cleaned up with 1.8x Mag-Bind beads and used to generate three replicate
library preparations for each sample. These technical replicates were generated to
ensure that any *VSGs* detected were not the result of PCR artifacts[41,42].
229
Because we expected a low number of parasites in each sample, we used a nested
PCR approach to prepare the VSG-seq libraries. First, we amplified *T. brucei* cDNA
from the parasite/host cDNA pool by PCR using a spliced leader primer paired with an
anchored oligo-dT primer (SL-1-nested and anchored oligo-dT; Supplemental Table 1).
20 cycles of PCR were completed (55ºC annealing, 45s extension) using Phusion
polymerase (Thermo Scientific, #F530L). PCR reactions were cleaned up with 1.8x
Mag-Bind beads. After amplifying *T. brucei* cDNA, a *VSG*-specific PCR reaction was
carried out using M13RSL and 14-mer-SP6 primers (see primers; Supplemental Table
1). 30 cycles of PCR (42ºC annealing, 45s extension) were performed using Phusion
polymerase. Amplified *VSG* cDNA was then cleaned up with 1X Mag-Bind beads and
quantified using a Qubit dsDNA HS Assay (Invitrogen Q32854).
241
Sequencing libraries were prepared from 1 ng of each VSG PCR product using the
Nextera XT DNA Library Preparation Kit (Illumina, FC-131-1096) following the
manufacturer's protocol except for the final cleanup step, which was performed using
1X Mag-Bind beads. Single-end 100bp sequencing was performed on an Illumina HiSeq
2500. Raw data are available in the National Center for Biotechnology Information
(NCBI) Sequence Read Archive under accession number PRJNA751607.
248
249

VSG-seq analysis of *T. b. gambiense* and *T. b. rhodesiense* sequencing libraries

250
To analyze both *T. b. gambiense* (VSG-seq preparations) and *T. b. rhodesiense*
(traditional mRNA sequencing library preparations; sequences were obtained from ENA,
accession numbers PRJEB27207 and PRJEB18523), we processed raw reads using
the VSG-seq pipeline available at https://github.com/mugnierlab/VSGSeqPipeline.
Briefly, *VSG* transcripts were assembled *de novo* from quality- and adapter-trimmed
reads for each sample (patient or patient replicate) from raw reads using Trinity (version
5.26.2) [43]. Contigs containing open reading frames (ORFs) were identified as
previously described [30]. ORF-containing contigs were compared to Lister 427 and
EATRO1125 *VSGs* as well as a collection of known contaminating non-*VSG*
sequences. Alignments to *VSGs* with an E-value below 1×10^{-10} that did not match any
known non-*VSG* contaminants were identified as *VSG* transcripts. For *T. b. gambiense*
replicate libraries, *VSG* ORFs identified in any patient replicate were consolidated into a
sole reference genome for each patient using CD-HIT (version 4.8.1)[44] with the
following arguments: -d 0 -c 0.98 -n 8 -G 1 -g 1 -s 0.0 -aL 0.0. Final *VSG* ORF files were
manually inspected.
266
Two *T. b. gambiense* patient *VSG*s (Patients 11 and 13) showed likely assembly errors.
In one case, a *VSG* was duplicated and concatenated, and in another, two *VSGs* were
concatenated. These reference files were manually corrected (removing the duplicate or
editing annotation to reflect two *VSGs* in the concatenated ORF) so that each *VSG*
could be properly quantified. *VSG* reference databases for each patient are available at
https://github.com/mugnierlab/Tbgambiense2021/. For *T. b. gambiense,* we then
aligned reads from each patient replicate to that patient's consolidated reference
genome using Bowtie with the parameters *-v 2 -m 1 -S* (version 1.2.3) [45].
275
For *T. b. rhodesiense,* we aligned each patient's data to its own *VSG* ORF assembly.
RPKM values for each *VSG* in each sample were generated using MULTo (version 1.0)
[46], and the percentage of parasites in each population expressing a *VSG* was
calculated as described previously [30]. For *T. b. gambiense* samples, we included only
VSGs with an expression measurement above 1% in two or more patient replicates in
our analysis. For *T. b. rhodesiense* samples, we included only *VSGs* with expression
>0.01%. To compare *VSG* expression between patients, despite the different reference
genomes used for each patient, we used CD-HIT to cluster *VSG* sequences with
greater than 98% similarity among patients, using the same parameters used to
consolidate reference *VSG* databases before alignment. We gave each unique *VSG*
cluster a numerical ID (e.g., Gambiense #) and chose the longest sequence within each
group to represent the cluster. Before analysis, we manually removed clusters
representing TgsGP and SRA from the expressed *VSG* sets. UpSet plots were made
with the UpSetR package[47]. The R code used to analyze resulting data and generate
figures is available at https://github.com/mugnierlab/Tbgambiense2021/.
291
292

Analysis of VSG N-terminal domains

Genomic *VSG* sequences
The *VSG* repertoires of *T. b. brucei* Lister 427 ("Lister427_2018" assembly), *T. b. brucei*
TREU927/4 and *T. b. gambiense* DAL972 were taken from TriTrypDB (v50), while the
T. b. brucei EATRO 1125 VSGnome was used for analysis of the EATRO1125 *VSG*
repertoire (vsgs_tb1125_nodups_atleast250aas_pro.txt, available at
https://tryps.rockefeller.edu/Sequences.html or GenBank accession KX698609.1 -
KX701858.1). VSG sequences from other strains (except those generated by VSG-seq)
were taken from the analysis in Cross, et al. [15]. Likely VSG N-termini were identified
as predicted proteins with significant similarity (e-value ≤ 10^{-5}) to hidden Markov models
(HMMs) of aligned type A and B VSG N-termini taken from [15].

N-terminal domain phylogenies
Phylogenies of VSG N-termini based on unaligned sequence similarities were
constructed using the method described in [48] and used previously to classify VSG
sequence [15]. We extracted predicted N-terminal domain protein sequences by using
the largest bounding envelope coordinates of a match to either type A or type B HMM. A
matrix of similarities between all sequences was constructed from normalized
transformed BLASTp scores as in Wickstead, et al. [48] and used to infer a neighbor-
joining tree using QuickTree v1.1 [49]. Trees were annotated and visualized in R with
the package APE v5.2 [50].

HMM
For N-terminal typing by HMM, we used a python analysis pipeline available at
(https://github.com/mugnierlab/find_VSG_Ndomains). The pipeline first identifies the
boundaries of the VSG N-terminal domain using the type A and type B HMM profiles
generated by Cross *et al.* which includes 735 previously-typed VSG N-terminal domain
sequences [15]. N-terminal domains are defined by the largest envelope domain
coordinate that meets e-value threshold ($1x10^{-5}$, --domE 0.00001). In cases where no
N-terminal domain is identified using these profiles, the pipeline executes a "rescue"
domain search in which the VSG is searched against a 'pan-VSG' N-terminus profile we
generated using 763 previously-typed VSG N-terminal domain sequences. This set of
VSGs includes several *T. brucei* strains and/or subspecies: Tb427 (559), TREU927
(138), *T. b. gambiense* DAL972 (28), EATRO795 (8), EATRO110 (5), *T. equiperdum*
(4), and *T. evansi* (21). The N-terminal domain type of these VSGs were previously
determined by Cross et. al (2014) by building neighbor-joining trees using local
alignment scores from all-versus-all BLASTp similarity searches [15]. Domain
boundaries are called using the same parameters as with the type A and B profiles.

After identifying boundaries, the pipeline extracts the sequence of the N-terminal
domain, and this is searched against five subtype HMM profiles. To generate N-terminal
domain subtype HMM profiles, five multiple sequence alignments were performed using
Clustal Omega [51] with the 763 previously-typed VSG N-terminal domain sequences
described above; each alignment included the VSG N-terminal domains of the same
subtype (A1, A2, A3, B1, and B2). Alignment output files in STOCKHOLM format were

used to generate distinct HMM profiles for type A1, A2, A3, B1, and B2 VSGs using the
pre-determined subtype classifications of the 763 VSGs using HMMer version 3.1b2
[52]. The number of sequences used to create each subtype profile ranged from 75 to
211. The most probable subtype is determined by the pipeline based on the highest
scoring sequence alignment against the subtype HMM profile database when HMMscan
is run under default alignment parameters. The pipeline generates a FASTA file
containing the amino acid sequence of each VSG N-terminus and a CSV with
descriptions of the N-terminal domain including its type and subtype.
347
<u>Network graph</u>
N-terminal network graphs were made using VSG N-terminal domains from the
TriTrypDB Lister427_2018 and *T. b. gambiense* DAL972 (v50) VSG sets described
above, and the *T. b. gambiense* and *T. b. rhodesiense* patient VSG N-termini which met
our expression thresholds. Identified N-terminal domains were then subjected to an all-
versus-all BLASTp. A pairwise table was created that includes each query-subject pair,
the corresponding alignment E-value, and N-terminal domain type of the query
sequence if previously typed in Cross, et al. [15]. Pseudogenes and fragments were
excluded from the Lister427_2018 reference prior to plotting by filtering for *VSG* genes
annotated as pseudogenes and any less than 400 amino acids in length, as the
remaining sequences are most likely to be full length VSG. Network graphs were
generated with the igraph R package[53] using undirected and unweighted clustering of
nodes after applying link cutoffs based on E-value $< 10^{-2}$. The leading eigenvector
clustering method [54] was used to detect and assign nodes to communities based on
clustering (cluster_leading_eigen() method in igraph).
363
364
Analysis of VSG C-terminal domains
VSG C-termini were extracted from expressed *T. b. gambiense VSGs*, *T.b. gambiense*
DAL972 (v50), and 545 previously-typed VSG C-termini from the Lister 427 strain using
the C-terminal HMM profile generated by Cross et al.[15] and the same HMMscan
parameters as for N-termini (E-value $< 1x10^{-5}$; largest domain based on envelope
coordinates). An all-vs-all BLASTp was performed on these sequences, and network
graphs were generated in the same manner as the N-terminal network graphs. Links
were drawn between C-termini with a BLASTp E-value $1x10^{-3}$. The leading eigenvector
method for clustering [54] was used to detect and assign nodes to communities based
on clustering (cluster_leading_eigen() method in igraph).
375
376
Comparison of gHAT patient VSGs to the DAL972 reference
The nucleotide sequences of expressed *T. b. gambiense* patient VSGs were compared
to the complete *T. b. gambiense* DAL972 reference genome sequence (v54) from
TriTrypDB using BLASTn. To find evidence of VSG diversification by mosaicism or
segmental gene conversion, permissive parameters were used so short regions of
homology would be reported (blastn -word_size 11 -evalue 0.1). Both query and subject
alignment coordinates were recorded in the output for downstream analysis, along with

E-value and percent identity. Alignment ranges were plotted with the IRanges R
package[55].

Results

Parasites in gHAT patients express diverse sets of VSGs

To investigate *VSG* expression in natural human infections, we performed VSG-seq on RNA extracted from whole blood collected from 12 human African trypanosomiasis patients from five locations in the Kwilu province of DR Congo (Figure 1A). We estimated the relative parasitemia of each patient by SL-QPCR [56], and we estimated the number of parasites after mAECT on buffy coat for all patients except patient 29 (Table 1). Using RNA extracted from 2.5 mL of whole blood from each patient, we prepared libraries for VSG-seq in three separate batches for each technical replicate. We amplified *T. brucei* RNA from host/parasite total RNA using a primer against the *T. brucei* spliced leader sequence and an anchored oligo-dT primer. The resulting trypanosome-enriched cDNA was used as a template to amplify *VSG* cDNA, and *VSG* amplicons were then submitted to VSG-seq sequencing and analysis. To determine whether a *VSG* was expressed within a patient, we applied the following stringent cutoffs:

1) We conservatively estimate that each 2.5 mL patient blood sample contained a minimum of 100 parasites. At this minimum parasitemia, a single parasite would represent 1% of the population (and consequently ~1% of the parasite RNA in a sample). As a result, we excluded all *VSGs* expressed by <1% of the population as estimated by VSG-seq.
2) We classified *VSGs* as expressed if they met the expression cutoff in at least two of three technical library replicates.

1112 unique *VSG* open reading frames were assembled *de novo* in the patient set and 44 met our expression criteria. Only these 44 *VSGs*, which we will refer to as "expressed *VSGs*," were considered in downstream analysis, except when otherwise noted. TgsGP, the VSG-like protein which partially enables resistance to human serum in *T. b. gambiense* [57]*,* assembled in samples from patients 2, 11, 13, and 17, and met the expression threshold in patients 2, 11, and 17. The absence of this transcript in most samples is likely due to the low amount of input material used to prepare samples. Notably, none of the expressed *VSGs* shared significant nucleotide sequence similarity outside the conserved C-terminal domain with any *VSGs* in the *T. b. gambiense* DAL972 genome (Supplemental Figures 6 and 7).

At least one *VSG* met our expression criteria in each patient, and in most cases, multiple *VSGs* were detected. Patient 2 showed the highest diversity, with 14 *VSGs* expressed (Figure 1B, Supplemental Figure 1). We observed a correlation between parasitemia, as estimated by qPCR, and the number of *VSGs* (Supplemental Figure 2), suggesting that these samples do not reflect the full diversity of each population. Nevertheless, two *VSGs* were shared between patients: *VSG* 'Gambiense 195' was expressed in both patient 12 and patient 17 from Village C; *VSG* 'Gambiense 38' was expressed in patient 12 from Village C and patient 23 from Village D (Figure 1C). Because our sampling did not reach saturation, resulting in some variability between

432 technical replicates, we chose to focus only on the presence/absence of individual
433 VSGs rather than relative expression levels within each population.
434
435

Patient	Location	est. parasites in 500µL buffy coat	mean SL-RNA Ct	WBC	Parasites in CSF	Stage
1	Village A	>50	22.155	1	-	First
2	Village A	>50	19.020	6	-	Early 2nd
3	Village A	2-5	28.780	6	-	Early 2nd
11	Village C	>50	22.030	9	-	Early 2nd
12	Village C	6-20	25.430	6	-	Early 2nd
13	Village C	6-20	26.635	12	-	Early 2nd
17	Village C	21-50	24.495	13	-	Early 2nd
19	Village C	1	28.245	7	-	Early 2nd
23	Village D	6-20	27.085	2	-	First
29	Village B	-	28.320	3	-	First
30	Village B	>50	22.960	694	+	Severe 2nd
33	Village E	1	32.385	2	-	First

436
437
438 **Table 1. Patient stage and parasitemia data.** We used the following staging definitions: First:
439 0-5 WBC/µl, no trypanosomes in cerebrospinal fluid (CSF). Second: >5 WBC/µl or
440 trypanosomes in CSF (with early 2nd: 6-20 WBC/µl and no trypanosomes in CSF; severe 2nd:
441 >100 WBC/µl). WBC: white blood cells.
442

Figure 1. Parasites isolated from gHAT patients express multiple *VSGs*. (A) Map showing
the location of each patient's home village. Maps were generated with ArcGIS® software by
Esri, using world imagery and National Geographic style basemaps. (B) Graph depicting the
total number of *VSGs* expressed in each patient. (C) The intersection of expressed *VSG* sets in
each patient. Bars on the left represent the size of the total set of VSGs expressed in each
patient. Dots represent an intersection of sets with bars above the dots representing the size of
the intersection. Color indicates patient origin.

Natural *T. b. gambiense* infections show a strong bias towards the expression of type B VSG

453
454

To further characterize the set of expressed *VSGs* in these samples, we sought a fast
and unbiased method for determining the type and subtype of each VSG's N-terminal
domain. We evaluated two approaches. The first approach was to create a bioinformatic
pipeline to determine each N-terminal domain subtype, using HMM profiles we created
for each subtype from sets of N-terminal domains previously typed by Cross et al. [15].
The second approach was to create a BLASTp network graph based on a published
method [58] where the N-terminal subtype of a VSG is determined by the set of VSGs it
clusters with, and clusters are identified using the leading eigenvector method [54]. We
used each approach to determine the N-terminal subtype of each expressed *VSG* in our
patient sample dataset, along with 863 VSG N-termini from the Lister 427 genome. We
compared these results to either existing N-terminal classification (for Lister 427 VSGs)
or classification based on position in a newly-generated BLASTp-tree [15] (for *T. b.*
gambiense VSGs; Figure 2A).

Both the new HMM profile and BLASTp network graph approaches generally
recapitulated previous VSG classification based on BLASTp-tree, with all three methods
agreeing 93.7% of the time (Figure 2B). The HMM pipeline method agreed with
BLASTp-tree typing for all patient VSGs, while the network graph approach agreed for
43/44 VSGs (Figure 2B, Supplemental Figure 3, Supplemental Table 1) [15]. It is not
surprising that the HMM pipeline would better reflect the results of the BLASTp-tree
method, as the N-terminal subtype HMM profiles were generated using VSGs classified
by this method. Based on these data, we determined that the HMM method is a fast and
accurate approach for determining the N-terminal domain types of unknown VSGs.

Our N-terminal domain typing pipeline identified the domain sequence and subtype for
all 44 patient VSGs (Figure 2C). Of the expressed *T. b. gambiense* VSGs, 82% had
type B N-terminal domains, and 50% or more of expressed *VSGs* within each patient
were type B. This bias was not restricted to highly expressed *VSGs*, as 74.5% of all
assembled *VSG* (813 of 1091 classifiable to an N-terminal subtype) were also type B.
This observation motivated further investigation into the expressed N-terminal domains
in infections by other *T. brucei* subspecies.

488
Figure 2. *T. b. gambiense* samples show a bias towards the expression of type B VSG. (A)
Visualization of relatedness between N-terminal domain peptide sequences inferred by
Neighbor-Joining based on normalized BLASTp scores. Legend indicates classification by HMM
pipeline (for Lister 427 VSGs, to highlight agreement between the two methods) or by
subspecies for *VSGs* expressed in patients. (B) Agreement between three VSG typing methods
for Lister 427 *VSG* set and the expressed *T. b. gambiense* patient *VSG* set. (C) N-terminal
domain subtype composition of expressed *T. b. gambiense VSGs* as determined by HMM
analysis pipeline.
497
498
499

500 **Type B VSG bias is not observed in natural *T. b. rhodesiense* infection**
501
To determine whether the bias towards type B VSGs was unique to *T. b. gambiense*
infections, we analyzed RNA-seq data from a published study measuring gene
expression in the blood and cerebrospinal fluid (CSF) of *T. b. rhodesiense* patients in
Northern Uganda [59]. These libraries were prepared conventionally after either rRNA-
depletion for blood or poly-A selection for CSF samples. We analyzed only those
samples for which at least 10% of reads mapped to the *T. brucei* genome. Raw reads
from these samples were subjected to the VSG-seq analysis pipeline. Because the
estimated parasitemia of these patients was much higher than our *T. b. gambiense*
study, we adjusted our expression criteria accordingly to 0.01%, the published limit of
detection of VSG-seq [30]. Using this approach, we identified 77 unique *VSG*
sequences across all blood and CSF samples (Figure 3A, Supplemental Figure 4).
SRA, the VSG-like protein that confers human serum resistance in *T. b. rhodesiense*
[60]*,* was expressed in all patient samples.
515
The HMM pipeline determined types for 74 of these VSG sequences; the remaining
sequences appeared to be incompletely assembled, presumably due to insufficient read
depth from their low level of expression. Multiple *VSGs* assembled in each patient
(Figure 3A), and a large proportion of *VSGs* were expressed in multiple patients (Figure
3C), in line with our observations in experimental mouse infections. Although most
VSGs detected in these patients were type B (57%, Figure 3B), this VSG type was
much less predominant than in *T. b. gambiense* infection. Interestingly, *T. b.*
rhodesiense patient CSF revealed another possible layer of diversity in *VSG*
expression, with 5 *VSGs* expressed exclusively in this space.
525
526
527
528

Figure 3. *T. b. rhodesiense* samples reveal diverse *VSG* expression but little N-terminal type bias. (A) The total number of expressed *T. b. rhodesiense VSGs* in each patient and sample type. Bar color represents the sample type from which RNA was extracted. (B) N-terminal domain subtype composition of all expressed *VSGs*. (C) Intersections of *VSGs* expressed in multiple infections. Bars on the left represent the size of the total set of VSGs expressed in each patient. Dots represent an intersection of sets, with bars above the dots representing the size of the intersection. Color indicates patient origin.

537
538

The predominant VSG N-terminal type fluctuates over time during experimental *T. b. brucei* infection

One explanation for the bias towards type B VSG in *T. b. gambiense* could be that VSG type stabilizes over time and type B predominates at later infection timepoints. This explanation is plausible because patient samples only represent a single moment during infection, and *T. b. gambiense* samples are more likely to be obtained at much later timepoints post-infection. To investigate whether N-terminal type predominance is stable or fluctuates over the course of infection, we took advantage of our published VSG-seq analysis of parasites isolated from mice infected with the *T.b. brucei* EATRO1125 strain. Blood was collected over time during this study, providing data from days 6/7, 12, 14, 21, 24, 26, and 30 post-infection in all four mice, and from days 96, 99, 102, and 105 in one of the four mice (Mouse 3). Of 192 unique VSGs identified between days 0-30, the python HMM pipeline typed 190; of 97 unique VSGs identified between days 96-105, the pipeline typed 93 VSGs. The remaining VSGs were incompletely assembled by Trinity. Our analysis of VSG types over time revealed that the predominantly expressed N-terminal domain type fluctuates between type A and type B throughout the early stages of infection and in extended chronic infections (Figure 4). Parasitemia did not correlate with either the diversity of *VSG* expression or N-terminal domain type predominance (Supplemental Figure 3). Because all patient samples were collected from a single timepoint in varying stages of disease, it remains unclear whether the predominant N-terminal type continues to fluctuate in human infections over time as it does in mice that have reached chronic stages of infection.

Figure 4. VSG N-terminal type composition fluctuates over the course of infection in mice. Proportions of N-terminal domain types expressed in *T. b. brucei* infected mice over time. The black dotted line represents the total number of identified VSGs. A) N-terminal type composition days 0-30. B) Type composition days 96-105.

The composition of the genomic VSG repertoire is reflected in expressed VSG N-terminal domain types

Another source for bias in expressed VSG type is the composition of the genomic *VSG* repertoire. We could directly compare the genomic and expressed *VSG* repertoire for EATRO1125 mouse infections, as the 'VSGnome' for this strain has been sequenced. Although the predominant N-terminal VSG type fluctuates throughout these infections, the expressed *VSG* repertoire across all time points generally reflects the composition of the genomic repertoire (chi-squared p = 0.0515, Figure 5A).

Unfortunately, the entire repertoire of *VSGs* encoded by most trypanosome strains is unknown, so such a direct comparison is impossible for *T. b. gambiense* and *T. b. rhodesiense* patient samples. Although the content of the 'core' *T. brucei* genome (containing the diploid, housekeeping genes) is similar enough among subspecies for resequencing projects to be scaffolded using the TREU927 or Lister 427 reference genomes [61–63], it is not clear whether the *VSG* repertoires of subspecies (or even individual parasite strains [36]) share this degree of similarity. Because no near-complete VSGnome for any *T. b. rhodesiense* strain was available, we compared the makeup of *T. b. rhodesiense* expressed *VSGs* with the closely related and near-complete *T. b. brucei* Lister 427 repertoire [62]. We observed no difference in the proportions of N-terminal types (chi-squared p-value = 0.2422) (Figure 5B). Similarly, the proportion of N-terminal domains identified in the *T. b. gambiense* patient samples is not statistically different from the incomplete *T. b. gambiense* DAL972 genomic repertoire (chi-squared p-value = 0.0575) (Figure 5B). Both *T. b. gambiense* patient VSGs (chi-squared p-value = 2.413e-4) and the 54 VSGs identified in *T. b. gambiense* DAL972 (chi-square p-value = 0.0301) have A and B type frequencies that differ significantly from Lister427. These results should be interpreted with caution given the limitations of the reference genomes used. Taken together, however, they support a model in which VSG types are drawn from the repertoire at a roughly equal frequency to their representation in the genome, with *T. b. gambiense* exhibiting an N-terminal type composition that differs from other subspecies.

Figure 5. VSG expression reflects the genomic VSG repertoire of the infecting parasites.
(A) Columns show the proportion of VSG types identified in each mouse infection over all time
points and the proportion of VSG types in the infecting *T. b. brucei* strain, EATRO 1125. The
total number of unique VSG sequences is displayed above each column. (B) A comparison of
the frequencies of type A and B VSGs expressed in patients and those present in Lister 427 and
DAL972 reference genomes. Relevant statistical comparisons are shown.

608
T. b. gambiense expressed VSG C-terminal domains show a bias for type 2
domains
611
In addition to examining N-terminal types in our *T. b. gambiense* dataset, we also
examined expressed VSG C-termini. Previous studies defined six C-terminal types,
although resolving these types in larger VSG sets has been difficult due to the extensive
similarity among VSG C-termini [15,58]. In line with previous observations, a BLASTp-
tree analysis of assembled *T. b. gambiense* C-terminal domains revealed frequent
sequence similarity between expressed C-terminal types but did not provide sufficient
resolution to confidently assign types (Fig 6A). To supplement this analysis, we also
performed a network graph analysis. Although this method had previously performed
poorly in resolving VSG C-termini [58], using the leading eigenvector clustering method
[54] to define community membership within the graph allowed a faithful reconstitution
of the C-terminal types previously determined by BLASTp-tree analysis.
623
Using the network graph approach, we tentatively assigned C-terminal domain types to
the *T. b. gambiense* VSGs (Figure 6B). Most patient C-terminal domain types were type
2, while the remaining types were predominantly type 1. Only one type 3 C-terminus
was identified in the patient set. Although there are very few VSG C-termini available in
the *T. b. gambiense* DAL972 genome, these sequences show a predominance of types
3 and 5. In contrast, the genomic repertoire of C-termini in Lister 427 shows a roughly
equivalent representation of types 1, 2, and 3. Unlike *T. b. gambiense* N-termini, the
expressed *T. b. gambiense* C-termini were more restricted than the sets of C-termini
apparently available in the *T. b. gambiense* DAL972 or *T. b. brucei* Lister 427 genomes
(Fisher's exact test, p-value $< 1 \times 10^{-5}$).
634
In line with previous observations, we saw no evidence of domain exclusion: a C-
terminal domain of one type could be paired with any type of N-terminal domain (Figure
6E)[20]. As observed in our analysis of expressed N-termini, C-terminal domain types
were not correlated to geographical origin (Figure 6F). Overall, these data suggest that,
like N-termini, expressed VSG C-termini are also biased towards certain C-terminal
types. Unlike N-termini, however, C-terminal types expressed in *T. b. gambiense*
infection may not reflect the parasite's genomic repertoire composition.

Figure 6. Expressed VSG C-termini are primarily type 1 and type 2. A) BLASTp-tree of C-
terminal domains. Points are colored based on previously determined C-terminal type from
Cross et al. or by the source of the sequence (genomic or expressed) for *T. b. gambiense*
VSGs. B) Network plot showing peptide sequence relatedness between C-terminal domains in
T. b. gambiense expressed VSGs. Each point represents a VSG C-terminus. A link was drawn
between points if the BLASTp e-value was less than $1x10^{-3}$. Points are colored by the cluster
determined by the clustering algorithm. Shaded circles also indicate clusters. C) Same network
plot as in B but colored by previously determined C-terminal type from Cross et al., or by source
for unclassified genomic or expressed *VSGs*. D) VSG C-terminal types, based on cluster
assignment visualized in panel B, in genomic and expressed *VSG* sets. E) Pairing of C- and N-
termini in *T. b. gambiense* patients. F) C-termini detected in each patient village.

Discussion

African trypanosomes evade the host adaptive immune response through a process of antigenic variation where parasites switch their expressed *VSG* [64]. The genome of *T. brucei* encodes a large repertoire of *VSG* genes, pseudogenes, and gene fragments that can be expanded continuously through recombination to form entirely novel "mosaic" *VSGs* [17]. While antigenic variation has been studied extensively in culture and animal infection models, our understanding of the process in natural infections, particularly human infection, is limited. Most experimental mouse infections are sustained for weeks to months, while humans and large mammals may be infected for several months or even years. Additionally, laboratory studies of antigenic variation almost exclusively use *T. b. brucei*, a subspecies of *T. brucei* that, by definition, does not infect humans.

The primary hurdle to exploring antigenic variation in nature has been technical: it is difficult to obtain sufficient parasite material for analysis. This is especially true for infection with *T. b. gambiense,* which often exhibits extremely low parasitemia. VSG-seq, which relies on PCR and requires very small amounts of material for analysis, provides a new tool for exploring *VSG* expression in natural human infections. Here we have demonstrated the feasibility of VSG-seq to analyze *VSG* expression in RNA samples isolated directly from HAT patients. Our analyses show that the diversity seen in mouse infection models is mostly recapitulated in natural infection; however, there may be unique aspects of antigenic variation in *T. b. gambiense* that can only be explored by studying natural infections.

The most notable result in our previous analysis of mouse infections was the diversity of *VSGs* expressed within each infection. Rather than a few *VSGs* expressed at a time, we saw many *VSGs* expressed simultaneously in each parasite population. This finding confirms previous estimates of antigenic diversity in experimental mouse infection [17] and suggests that the genomic *VSG* repertoire might be exploited very rapidly. The study presented here detected several expressed *VSGs* in most *T. b. gambiense* samples. Although diversity in *T. b. gambiense* infection appeared lower overall, the correlation we observed between parasitemia and diversity in *T. b. gambiense* isolates suggests that our sampling was incomplete. Indeed, in our analysis of *T. b. rhodesiense* infection (a more reasonable comparison to mouse infection given similar expression cutoffs and parasitemia), we observed diversity similar to or higher than what we have detected in *T. b. brucei* mouse infections. Moreover, *T. b. rhodesiense* patient CSF revealed another layer of diversity in *VSG* expression, with 5 *VSGs* expressed exclusively in this space. Although this observation is difficult to interpret without information about the precise timing of sample collection, it is exciting to speculate that different organs or body compartments could harbor different sets of VSGs. Overall, our analysis of *VSG* expression in *T. b. gambiense* and *T. b. rhodesiense* patients confirmed the long-held assumption that *VSG* diversity is a feature of natural infection.

While analyzing the sets of expressed *VSGs* in *T. b. gambiense* and *T. b. rhodesiense* infections, we found evidence for another feature of experimental infection that holds

true in a natural host: hierarchical *VSG* expression. Both *in vitro* and *in vivo* studies
have shown that *VSG* switching is not entirely stochastic but rather hierarchical. Specific
variants dominate expression in the parasite population in a reproducible order that
appears to be independent of the starting *VSG* [17,31,32,65,66]. In the *T. b. gambiense*
samples, we found two *VSGs* that met our detection threshold in multiple patients. In *T.
b. rhodesiense* samples*,* multiple patients shared a large proportion of expressed *VSGs*.
Given the large size of the genomic *VSG* repertoire, any overlap in expressed *VSG*
repertoire is likely indicative of a semi-predictable hierarchy of switching and preference
for the expression of certain *VSGs*.
709
Of the two shared *VSGs* we identified in the *T. b. gambiense* patients, one was
identified in two patients from the same village, while the other was found in two
patients from villages 40km apart. At this short distance, it is possible that the infecting
parasites were genetically similar. However, it would be interesting to investigate
whether preference for the expression of certain *VSGs* occurs even between parasites
isolated at greater distances. Indeed, past research has shown that the sensitivity of
serological tests for gHAT, which detect antibodies against the LiTat 1.3 VSG, vary
regionally, potentially due to differences in the underlying genomic or expressed *VSG*
repertoire in circulating strains [67,68].
719
Our data are consistent with the notion that there could be geographic variations in *T.
brucei VSG* repertoires. None of our assembled *T. b. gambiense VSGs* from patients in
the DR Congo shared significant N-terminal sequence similarity with those in the
genome of DAL972, a parasite isolate from Côte d'Ivoire. In contrast, the C-termini of
patient *VSGs* were well represented by similar sequences within the DAL972 reference.
Geographic variation has been observed in *var* gene repertoires of *Plasmodium
falciparum* [69] and the *VSG* repertoire of *Trypanosoma vivax*, another African
trypanosome [35]. A better understanding of such differences in *T. brucei*, if they exist,
could lead to development of more sensitive HAT diagnostics.
729
To better understand the VSG proteins expressed in natural infections, we developed
an HMM VSG typing pipeline that revealed an intriguing bias in *T. b. gambiense*
infection towards type B VSGs, which appears to be specific to *T. b. gambiense* patient
samples. Small sample sizes and important differences between each *T. brucei*
subspecies' dataset limit the conclusions that can be drawn. However, comparisons
between these sets suggest that the genomic *VSG* repertoire determines the
distribution of VSG N-terminal types expressed during *T. brucei* infection and may
account for the bias we have observed in *T. b. gambiense* patients. Thus, the *T. b.
gambiense VSG* repertoire may contain a larger proportion of type B VSGs than its
more virulent counterparts.
740
Another possibility we cannot rule out, however, is that the gHAT samples are biased
due to selection by the serological test used for diagnosis. Patients were screened for *T.
b. gambiense* infection using the CATT, a serological test that uses parasites
expressing VSG LiTat 1.3 as an antigen. LiTat 1.3 contains a type B2 N-terminal
domain [67,68]. Patients infected with parasites predominantly expressing type B VSGs

may be more likely to generate antibodies that cross-react with LiTat1.3, resulting in
preferential detection of these cases. In contrast, *T.b. rhodesiense* can only be
diagnosed microscopically, removing the potential to introduce bias through screening.
It remains to be investigated whether samples from patients diagnosed using newer
screening tests, which include the invariant surface glycoprotein ISG65 and the type A
VSG LiTat 1.5 [29], would show similar bias towards the expression of type B VSGs.
752
Analysis of expressed VSG C-terminal domains in *T. b. gambiense* patients showed a
bias towards C-terminal types 1 and 2. The diagnostic VSG LiTat1.3 contains a type 3
C-terminus, a C-terminal type underrepresented in the patient set. Therefore, it is
unlikely that a bias in expressed C-terminal types is related to the screening method.
Notably, the bias toward C-terminal types 1 and 2 was not reflected in the limited *VSG*
repertoire of the DAL972 reference genome or the repertoire of the Lister 427 *T. b.*
brucei reference genome. This finding could be related to the limited set of *VSG*s
present in the DAL972 reference genome or reflect a true expression bias.
761
Could a bias towards certain VSG types, whether due to a difference in repertoire
composition or expression preference, account for unique features of *T. b. gambiense*
infection*,* including its chronicity and primarily anthroponotic nature [70]? While the
genomic *VSG* repertoire has been analyzed extensively in laboratory strains, little is
known about how differences in VSG proteins relate to parasite biology or whether there
could be biological consequences to the expression of specific VSG N- or C-terminal
types. Type A *var* genes in *Plasmodium falciparum* infection are associated with severe
malaria [71–75], and similar mechanisms have been hypothesized to exist in *T. vivax*
and *T. congolense* infections [33,35,76,77].
771
In *T. brucei,* several VSGs have evolved specific functions besides antigenic variation
[77]. The first type B VSG structure was recently solved [78], revealing a unique *O-*
linked carbohydrate in the VSG's N-terminal domain. This modification interfered with
the generation of protective immunity in a mouse infection model; perhaps structural
differences between each VSG type, including glycosylation patterns, could influence
infection outcomes. Further research will be needed to determine whether the observed
predominance of type B VSGs could influence the biology of *T. b. gambiense* infection.
779
Currently, it is unclear why this collection of gHAT patient isolates demonstrates a bias
towards the expression of certain VSG types. More research will be needed to
determine whether the *T. b. gambiense VSG* repertoire contains a unique distribution of
VSG types, whether these parasites preferentially express certain VSG types, and
whether this bias could have functional consequences. This study has shown, however,
is that it is feasible to explore antigenic variation in natural infection. Moreover, although
mouse models do reflect the general dynamics of antigenic variation in natural *T. brucei*
infection, unique biology remains to be uncovered by studying this process in its natural
context.
789
790
791

Acknowledgments

We are very grateful to the patients without whom this work would not have been
possible. We thank George Cross and Danae Schulz for comments on the manuscript,
and Mary Gebhardt for help with GIS. The Atlas of HAT is an initiative of the World
Health Organization (WHO), jointly implemented with the Food and Agriculture
Organization of the United Nations (FAO) in the framework of the Programme Against
African Trypanosomiasis (PAAT). Field work and specimen collection in DR Congo
were funded through the Wellcome Trust (study number 099310/Z/12/Z) awarded to the
TrypanoGEN Consortium (www.trypanogen.net), members of H3Africa (h3africa.org).
Sample work-up was supported by the Research Foundation Flanders (FWO grant
1501413N). Work by BW was supported by University of Nottingham/Wellcome Trust
Institutional Strategic Support Fund award 204843/Z/16/Z. MRM and SS were
supported by Office of the Director, NIH (DP5OD023065). JS is supported by NIH
T32AI007417.

Supplement

Supplemental Table 1. Primer sequences.

Supplemental Table 2. gHAT patient distance matrix.

Supplemental Table 3. gHAT VSG expression data.

Supplemental Table 4. Tables comparing BLAST-tree, HMMscan, and network plot typing methods.

Supplemental Table 5. rHAT VSG expression data.

Supplemental Figure 1. Heatmap of all assembled *T.b. gambiense* patient VSGs.
Greyscale shows $\log_{10}$ of the estimated percentage of the parasite population expressing each
VSG. Variants expressed by less than 1% of parasites considered not detected (n.d.).

828
Supplemental Figure 2. Correlation between parasitemia and diversity and N-terminal type distribution. (A) Correlation plots for VSG diversity and percent of N-terminal domain type B for *T.b. brucei* infected mice from Mugnier et al. 2015. (B) Correlation plots for *T.b. rhodesiense* infected patients from Mulindwa et al. 2018. (C) Correlation plots for *T.b. gambiense* infected patients.

Supplemental Figure 3. (A) Network plot showing peptide sequence relatedness between N-terminal domains. Each point represents a VSG N-terminus. A link was drawn between points if the BLASTp e-value was less than 10^{-2}. Colors and shaded circles represent community assignments determined by the clustering algorithm. (B) The same graph as in (A), but points are manually colored by known N-terminal subtype from Cross et al. or by subspecies for VSGs identified in patients.

847   **Supplemental Figure 4. BLASTp-tree of all *T. b. gambiense VSGs.*** File attached.

Supplemental Figure 5. Heatmap of all assembled *T. b. rhodesiense* patient VSGs.
Greyscale shows log$_{10}$ of the estimated percentage of the parasite population expressing each
VSG. Variants expressed by less than 0.01% of parasites considered not detected (n.d.).

Supplemental Figure 6. Ranges of all BLASTn hits with e-value < 0.1 for each assembled
gHAT patient VSG against the TbgDAL972 whole genome reference. The full length of each
patient VSG in base pairs is plotted as the X-axis of each plot facet. Bars color-coded by
percent identity show each hit's alignment length and position within the query VSG. Alignments
within the same query coordinates that hit multiple locations in the reference genome are
stacked in bins. The maximum number of bins is plotted on the y-axis.
862

863
864   **Supplemental Figure 7. Patient VSG representation within TbgDAL972.** A histogram of the
865   percentage of each patient VSG that was represented anywhere within the Tbg DAL972
866   genome by BLASTn with a threshold E-value < 0.1.
867

References

1. Romero-Meza G, Mugnier MR. Trypanosoma brucei. Trends in Parasitology. 2019. doi:10.1016/j.pt.2019.10.007
2. Büscher P, Cecchi G, Jamonneau V, Priotto G. Human African trypanosomiasis. www.thelancet.com. 2017;390. doi:10.1016/S0140-6736(17)31510-6
3. Franco JR, Cecchi G, Priotto G, Paone M, Diarra A, Grout L, et al. Monitoring the elimination of human African trypanosomiasis at continental and country level: Update to 2018. PLoS Negl Trop Dis. 2020;14: e0008261. doi:10.1371/JOURNAL.PNTD.0008261
4. Kennedy PGE, Rodgers J. Clinical and neuropathogenetic aspects of human African trypanosomiasis. Frontiers in Immunology. Frontiers Media S.A.; 2019. doi:10.3389/fimmu.2019.00039
5. Franco JR, Cecchi G, Priotto G, Paone M, Diarra A, Grout L, et al. Monitoring the elimination of human African trypanosomiasis at continental and country level: Update to 2018. PLoS Negl Trop Dis. 2020;14: e0008261. doi:10.1371/JOURNAL.PNTD.0008261
6. WHO. The Global Health Observatory. In: WHO [Internet]. World Health Organization; [cited 17 Aug 2021]. Available: https://www.who.int/data/gho/data/themes/topics/indicator-groups/indicator-group-details/GHO/human-african-trypanosomiasis
7. WHO. Ending the neglect to attain the sustainable development goals: a road map for neglected tropical diseases 2021–2030. Geneva; 2020.
8. Magez S, Caljon G, Tran T, Stijlemans B, Radwanska M. Current status of vaccination against African trypanosomiasis. Parasitology. 2010;137: 2017–2027.
9. Trindade S, Rijo-Ferreira F, Carvalho T, Pinto-Neves D, Guegan F, Aresta-Branco F, et al. Trypanosoma brucei Parasites Occupy and Functionally Adapt to the Adipose Tissue in Mice. Cell Host Microbe. 2016;19: 837–848. doi:10.1016/j.chom.2016.05.002
10. Pereira SS, Trindade S, De Niz M, Figueiredo LM. Tissue tropism in parasitic diseases. Open Biology. Royal Society Publishing; 2019. doi:10.1098/rsob.190036
11. Camara M, Soumah AM, Ilboudo H, Travaillé C, Clucas C, Cooper A, et al. Extravascular Dermal Trypanosomes in Suspected and Confirmed Cases of gambiense Human African Trypanosomiasis. Clin Infect Dis. 2021;73: 12–20. doi:10.1093/cid/ciaa897
12. Alfituri OA, Bradford BM, Paxton E, Morrison LJ, Mabbott NA. Influence of the Draining Lymph Nodes and Organized Lymphoid Tissue Microarchitecture on Susceptibility to Intradermal Trypanosoma brucei Infection. Front Immunol. 2020;11. doi:10.3389/fimmu.2020.01118
13. Marcello L, Barry JD. Analysis of the VSG gene silent archive in Trypanosoma brucei reveals that mosaic gene expression is prominent in antigenic variation and is favored by archive substructure. Genome Res. 2007;17: 1344–1352.
14. Berriman M, Ghedin E, Hertz-Fowler C, Blandin G, Renauld H, Bartholomeu DC, et al. The genome of the African trypanosome Trypanosoma brucei. Sci (New York, NY). 2005;309: 416–422.
15. Cross GAM, Kim H-S, Wickstead B. Capturing the variant surface glycoprotein

repertoire (the VSGnome) of Trypanosoma brucei Lister 427. Mol Biochem
Parasitol. 2014;195: 59–73. doi:10.1016/j.molbiopara.2014.06.004

16. Müller LSM, Cosentino RO, Förstner KU, Guizetti J, Wedel C, Kaplan N, et al.
Genome organization and DNA accessibility control antigenic variation in
trypanosomes. Nature. 2018;563: 121–125.

17. Hall JPJ, Wang H, Barry JD. Mosaic VSGs and the scale of Trypanosoma brucei
antigenic variation. Horn D, editor. PLoS Pathog. 2013;9: e1003502.

18. Jayaraman S, Harris C, Paxton E, Donachie A-M, Vaikkinen H, McCulloch R, et
al. Application of long read sequencing to determine expressed antigen diversity
in Trypanosoma brucei infections. Acosta-Serrano A, editor. PLoS Negl Trop Dis.
2019;13: e0007262. doi:10.1371/journal.pntd.0007262

19. Carrington M, Miller N, Blum M, Roditi I, Wiley D, Turner M. Variant specific
glycoprotein of Trypanosoma brucei consists of two domains each having an
independently conserved pattern of cysteine residues. J Mol Biol. 1991;221: 823–
835.

20. Hutchinson OC, Smith W, Jones NG, Chattopadhyay A, Welburn SC, Carrington
M. VSG structure: similar N-terminal domains can form functional VSGs with
different types of C-terminal domain. Mol Biochem Parasitol. 2003;130: 127–131.

21. Jones NG, Nietlispach D, Sharma R, Burke DF, Eyres I, Mues M, et al. Structure
of a glycosylphosphatidylinositol-anchored domain from a trypanosome variant
surface glycoprotein. J Biol Chem. 2008;283: 3584–3593.

22. Schwede A, Jones N, Engstler M, Carrington M. The VSG C-terminal domain is
inaccessible to antibodies on live trypanosomes. Mol Biochem Parasitol.
2011;175: 201–204.

23. Van Meirvenne N, Magnus E, Büscher P. Evaluation of variant specific
trypanolysis tests for serodiagnosis of human infections with Trypanosoma brucei
gambiense. Acta Trop. 1995;60: 189–199. doi:10.1016/0001-706X(95)00127-Z

24. Magnus E, Vervoort T, Van Meirvenne N. A card-agglutination test with stained
trypanosomes (C.A.T.T.) for the serological diagnosis of T. b. gambiense
trypanosomiasis. Ann Soc Belg Med Trop (1920). 1978;58: 169–176.

25. Dukes P, Gibson WC, Gashumba JK, Hudson KM, Bromidge TJ, Kaukus A, et al.
Absence of the LiTat 1.3 (CATT antigen) gene in Trypanosoma brucei gambiense
stocks from Cameroon. Acta Trop. 1992.

26. Enyaru JCK, Allingham R, Bromidge T, Kanmogne GD, Carasco JF. The isolation
and genetic heterogeneity of Trypanosoma brucei gambiense from north-west
Uganda. Acta Trop. 1993.

27. Bisser S, Lumbala C, Nguertoum E, Kande V, Flevaud L, Vatunga G, et al.
Sensitivity and Specificity of a Prototype Rapid Diagnostic Test for the Detection
of Trypanosoma brucei gambiense Infection: A Multi-centric Prospective Study.
Acosta-Serrano A, editor. PLoS Negl Trop Dis. 2016;10: e0004608.
doi:10.1371/journal.pntd.0004608

28. Büscher P, Mertens P, Leclipteux T, Gilleman Q, Jacquet D, Mumba-Ngoyi D, et
al. Sensitivity and specificity of HAT Sero-K-SeT, a rapid diagnostic test for
serodiagnosis of sleeping sickness caused by Trypanosoma brucei gambiense: A
case-control study. Lancet Glob Heal. 2014;2: e359–e363. doi:10.1016/S2214-
109X(14)70203-7

29.   Lumbala C, Biéler S, Kayembe S, Makabuza J, Ongarello S, Ndung'u JM.
Prospective evaluation of a rapid diagnostic test for Trypanosoma brucei
gambiense infection developed using recombinant antigens. PLoS Negl Trop Dis.
2018;12: e0006386. doi:10.1371/JOURNAL.PNTD.0006386

30.   Mugnier MR, Cross GAM, Papavasiliou FN. The in vivo dynamics of antigenic
variation in Trypanosoma brucei. Science (80-). 2015;347: 1470–1473.
doi:10.1126/science.aaa4502

31.   Morrison LJ, Majiwa P, Read AF, Barry JD. Probabilistic order in antigenic
variation of Trypanosoma brucei. Int J Parasitol. 2005;35: 961–972.

32.   Lythgoe KA, Morrison LJ, Read AF, Barry JD. Parasite-intrinsic factors can
explain ordered progression of trypanosome antigenic variation. Proc Natl Acad
Sci U S A. 2007;104: 8095–8100.

33.   Silva Pereira S, Heap J, Jones AR, Jackson AP. VAPPER: High-throughput
variant antigen profiling in African trypanosomes of livestock. Gigascience.
2019;8. doi:10.1093/gigascience/giz091

34.   Pereira SS, Casas-Sánchez A, Haines LR, Ogugo M, Absolomon K, Sanders M,
et al. Variant antigen repertoires in Trypanosoma congolense populations and
experimental infections can be profiled from deep sequence data using universal
protein motifs. Genome Res. 2018;28: 1383–1394. doi:10.1101/gr.234146.118

35.   Silva Pereira S, de Almeida Castilho Neto KJG, Duffy CW, Richards P, Noyes H,
Ogugo M, et al. Variant antigen diversity in Trypanosoma vivax is not driven by
recombination. Nat Commun. 2020;11: 844. doi:10.1038/s41467-020-14575-8

36.   Hutchinson OC, Picozzi K, Jones NG, Mott H, Sharma R, Welburn SC, et al.
Variant Surface Glycoprotein gene repertoires in Trypanosoma brucei have
diverged to become strain-specific. BMC Genomics. 2007;8: 234.
doi:10.1186/1471-2164-8-234

37.   Ilboudo H, Noyes H, Mulindwa J, Kimuda MP, Koffi M, Kaboré JW, et al.
Introducing the TrypanoGEN biobank: A valuable resource for the elimination of
human African trypanosomiasis. PLoS Negl Trop Dis. 2017;11: e0005438.
doi:10.1371/JOURNAL.PNTD.0005438

38.   Camara M, Camara O, Ilboudo H, Sakande H, Kaboré J, N'Dri L, et al. Sleeping
sickness diagnosis: use of buffy coats improves the sensitivity of the mini anion
exchange centrifugation test. Trop Med Int Heal. 2010;15: 796–799.
doi:10.1111/J.1365-3156.2010.02546.X

39.   Simarro PP, Cecchi G, Paone M, Franco JR, Diarra A, Ruiz JA, et al. The Atlas of
human African trypanosomiasis: A contribution to global mapping of neglected
tropical diseases. Int J Health Geogr. 2010;9: 57. doi:10.1186/1476-072X-9-57

40.   González-Andrade P, Camara M, Ilboudo H, Bucheton B, Jamonneau V,
Deborggraeve S. Diagnosis of trypanosomatid infections: Targeting the spliced
leader RNA. J Mol Diagnostics. 2014. doi:10.1016/j.jmoldx.2014.02.006

41.   Brakenhoff RH, Schoenmakers JG, Lubsen NH. Chimeric cDNA clones: a novel
PCR artifact. Nucleic Acids Res. 1991;19: 1949. doi:10.1093/NAR/19.8.1949

42.   Meyerhans A, Vartanian J-P, Wain-Hobson S. DNA recombination during PCR.
Nucleic Acids Res. 1990;18: 1687–1691. doi:10.1093/NAR/18.7.1687

43.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-
length transcriptome assembly from RNA-Seq data without a reference genome.

Nat Biotechnol. 2011;29: 644–652.

44.   Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. Bioinformatics. 2006;22: 1658–1659.

45.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biol. 2009;10:
R25.

46.   Storvall H, Ramsköld D, Sandberg R. Efficient and comprehensive representation
of uniqueness for next-generation sequencing by minimum unique length
analyses. PLoS One. 2013;8: e53822.

47.   Conway JR, Lex A, Gehlenborg N. UpSetR: An R Package for the Visualization of
Intersecting Sets and their Properties. doi:10.1101/120600

48.   Wickstead B, Gull K. Dyneins across eukaryotes: a comparative genomic
analysis. Traffic. 2007;8: 1708–1721.

49.   Howe K, Bateman A, Durbin R. QuickTree: Building huge neighbour-joining trees
of protein sequences. Bioinformatics. 2002;18: 1546–1547.
doi:10.1093/bioinformatics/18.11.1546

50.   Paradis E, Schliep K. Ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics. 2019;35: 526–528.
doi:10.1093/bioinformatics/bty633

51.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal
Omega. Mol Syst Biol. 2011;7: 539–539. doi:10.1038/msb.2011.75

52.   Eddy SR. Accelerated Profile HMM Searches. PLOS Comput Biol. 2011;7:
e1002195. doi:10.1371/JOURNAL.PCBI.1002195

53.   Csárdi G, Nepusz T. The igraph software package for complex network research.

54.   Newman MEJ. Finding community structure in networks using the eigenvectors of
matrices. Phys Rev E. 2006;74: 036104. doi:10.1103/PhysRevE.74.036104

55.   Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al.
Software for Computing and Annotating Genomic Ranges. PLOS Comput Biol.
2013;9: e1003118. doi:10.1371/JOURNAL.PCBI.1003118

56.   González-Andrade P, Camara M, Ilboudo H, Bucheton B, Jamonneau V,
Deborggraeve S. Diagnosis of trypanosomatid infections: targeting the spliced
leader RNA. J Mol Diagn. 2014;16: 400–404.

57.   Berberof M, Pérez-Morga D, Pays E. A receptor-like flagellar pocket glycoprotein
specific to Trypanosoma brucei gambiense. Mol Biochem Parasitol. 2001;113:
127–138.

58.   Weirather JL, Wilson ME, Donelson JE. Mapping of VSG similarities in
Trypanosoma brucei. Mol Biochem Parasitol. 2012;181: 141–152.

59.   Mulindwa J, Leiss K, Ibberson D, Kamanyi Marucha K, Helbig C, Melo do
Nascimento L, et al. Transcriptomes of Trypanosoma brucei rhodesiense from
sleeping sickness patients, rodents and culture: Effects of strain, growth
conditions and RNA preparation methods. Salavati R, editor. PLoS Negl Trop Dis.
2018;12: e0006280.

60.   De Greef C, Hamers R. The serum resistance-associated (SRA) gene of
Trypanosoma brucei rhodesiense encodes a variant surface glycoprotein-like
protein. Mol Biochem Parasitol. 1994;68: 277–284.

61. Jackson AP, Sanders M, Berry A, McQuillan J, Aslett MA, Quail MA, et al. The
genome sequence of Trypanosoma brucei gambiense, causative agent of chronic
human African Trypanosomiasis. PLoS Negl Trop Dis. 2010;4.
doi:10.1371/journal.pntd.0000658
62. Sistrom M, Evans B, Benoit J, Balmer O, Aksoy S, Caccone A. De Novo Genome
Assembly Shows Genome Wide Similarity between Trypanosoma brucei brucei
and Trypanosoma brucei rhodesiense. Louis EJ, editor. PLoS One. 2016;11:
e0147660. doi:10.1371/journal.pone.0147660
63. Sistrom M, Evans B, Bjornson R, Gibson W, Balmer O, Maser P, et al.
Comparative genomics reveals multiple genetic backgrounds of human
pathogenicity in the trypanosoma brucei complex. Genome Biol Evol. 2014;6:
2811–2819. doi:10.1093/gbe/evu222
64. Mugnier MR, Stebbins CE, Papavasiliou FN. Masters of Disguise: Antigenic
Variation and the VSG Coat in Trypanosoma brucei. PLOS Pathog. 2016;12:
e1005784. doi:10.1371/journal.ppat.1005784
65. Aitcheson N, Talbot S, Shapiro J, Hughes K, Adkin C, Butt T, et al. VSG switching
in Trypanosoma brucei: antigenic variation analysed using RNAi in the absence of
immune selection. Mol Microbiol. 2006;57: 1608–1622. doi:10.1111/j.1365-
2958.2005.04795.x.VSG
66. Liu D, Albergante L, Newman TJ, Horn D. Faster growth with shorter antigens can
explain a VSG hierarchy during African trypanosome infections: a feint attack by
parasites. Sci Rep. 2018;8: 10922. doi:10.1038/s41598-018-29296-8
67. Chappuis F, Loutan L, Simarro P, Lejon V, Büscher P. Options for field diagnosis
of human african trypanosomiasis. Clin Microbiol Rev. 2005;18: 133–146.
68. Truc P, Lejon V, Magnus E, Jamonneau V, Nangouma A, Verloo D, et al.
Evaluation of the micro-CATT, CATT/Trypanosoma brucei gambiense, and
LATEX/T b gambiense methods for serodiagnosis and surveillance of human
African trypanosomiasis in West and Central Africa. Bull World Health Organ.
2002;80: 882–886.
69. Tonkin-Hill G, Ruybal-Pesántez S, Tiedje KE, Rougeron V, Duffy MF, Zakeri S, et
al. Evolutionary analyses of the major variant surface antigen-encoding genes
reveal population structure of Plasmodium falciparum within and between
continents. PLOS Genet. 2021;17: e1009269.
doi:10.1371/JOURNAL.PGEN.1009269
70. Büscher P, Bart JM, Boelaert M, Bucheton B, Cecchi G, Chitnis N, et al. Do
Cryptic Reservoirs Threaten Gambiense-Sleeping Sickness Elimination? Trends
Parasitol. 2018;34: 197–207. doi:10.1016/J.PT.2017.11.008
71. Kirchgatter K, del Portillo HA. Association of Severe Noncerebral Plasmodium
falciparum Malaria in Brazil With Expressed PfEMP1 DBL1α Sequences Lacking
Cysteine Residues. Mol Med 2002 81. 2002;8: 16–23. doi:10.1007/BF03401999
72. Bull PC, Berriman M, Kyes S, Quail MA, Hall N, Kortok MM, et al. Plasmodium
falciparum Variant Surface Antigen Expression Patterns during Malaria. PLOS
Pathog. 2005;1: e26. doi:10.1371/JOURNAL.PPAT.0010026
73. Abdi AI, Hodgson SH, Muthui MK, Kivisi CA, Kamuyu G, Kimani D, et al.
Plasmodium falciparum malaria parasite var gene expression is modified by host
antibodies: longitudinal evidence from controlled infections of Kenyan adults with

varying natural exposure. BMC Infect Dis 2017 171. 2017;17: 1–11.
doi:10.1186/S12879-017-2686-0

74.   Kyriacou HM, Stone GN, Challis RJ, Raza A, Lyke KE, Thera MA, et al.
Differential var gene transcription in Plasmodium falciparum isolates from patients
with cerebral malaria compared to hyperparasitaemia. Mol Biochem Parasitol.
2006;150: 211–218. doi:10.1016/J.MOLBIOPARA.2006.08.005

75.   Duffy F, Bernabeu M, Babar PH, Kessler A, Wang CW, Vaz M, et al. Meta-
analysis of plasmodium falciparum var signatures contributing to severe Malaria in
African children and Indian adults. MBio. 2019;10. doi:10.1128/MBIO.00217-19

76.   Pereira SS, Casas-Sánchez A, Haines LR, Ogugo M, Absolomon K, Sanders M,
et al. Variant antigen repertoires in Trypanosoma congolense populations and
experimental infections can be profiled from deep sequence data using universal
protein motifs. Genome Res. 2018;28: 1383. doi:10.1101/GR.234146.118

77.   Silva Pereira S, Jackson AP, Figueiredo LM. Evolution of the variant surface
glycoprotein family in African trypanosomes. Trends Parasitol. 2021.
doi:10.1016/J.PT.2021.07.012

78.   Pinger J, Nešić D, Ali L, Aresta-Branco F, Lilic M, Chowdhury S, et al. African
trypanosomes evade immune clearance by O-glycosylation of the VSG surface
coat. Nat Microbiol. 2018;19: 53.

1117
1118