1
2
Sperm Histone H3 Lysine 4 tri-methylation serves as a metabolic sensor of paternal obesity and is associated with the inheritance of metabolic dysfunction
5

Authors
Anne-Sophie Pepin[1], Christine Lafleur[2], Romain Lambrot[2], Vanessa Dumeaux[3], Sarah Kimmins[1,2,4]*

Affiliations
[1]Department of Pharmacology and Therapeutics, Faculty of Medicine, McGill University, Montreal, QC H3G 1Y6, Canada
[2]Department of Animal Science, Faculty of Agricultural and Environmental Sciences, McGill University, Montreal, QC H9X 3V9, Canada
[3]Department of Biology, PERFORM Center, Concordia University, Montreal, QC H4B 1R6, Canada
[4]Lead Contact
*Correspondence: sarah.kimmins@mcgill.ca

Running title: Paternal obesity alters sperm H3K4me3 and metabolic outcomes in descendants

Summary sentence: Paternal obesity impacts sperm H3K4me3 and is associated with placenta, embryonic and metabolic outcomes in descendants.

Keywords: obesity, metabolism, chromatin, sperm, epigenetic inheritance

Abstract

Parental environmental exposures can strongly influence descendant risks for adult disease. Metabolic
disorders arise from the intersection of environmental and genetic risk factors, with epigenetic inheritance
being at the center of the familial cycle of transgenerational disease. How paternal high-fat diet changes
the sperm chromatin leading to the acquisition of metabolic disease in offspring remains controversial and
ill-defined. Using a genetic model of epigenetic inheritance, we investigated the role of histone H3 lysine 4
methylation (H3K4me3) in the paternal transmission of metabolic dysfunction. We show that obesity-
induced alterations in sperm H3K4me3 associated with offspring phenotypes and corresponded to
embryonic and placental chromatin profiles and gene expression. Transgenerational susceptibility to
metabolic disease was only observed when grandsires had a pre-existing modified sperm epigenome due
to overexpression of the histone demethylase KDM1 in the male germline. This non-DNA based knowledge
of inheritance has the potential to improve our understanding of how environment shapes heritability and
may lead to novel routes for the prevention of disease.

Introduction
The prevalence of obesity and type II diabetes is growing globally at rates indicating that environment rather
than genes is the principal driver. Exposures to high-fat diet, toxicants or micronutrient deficiency can
impact our health and that of future generations (Gernand *et al.*, 2016; Braun, Messerlian and Hauser,
2017; Donkin and Barrès, 2018; Eberle *et al.*, 2020). Only now are we beginning to identify mechanisms
linking these exposures to parental and offspring health. One connection between environment and health
is the epigenome. The epigenome refers to the biochemical content associated with DNA that impacts gene
expression, chromatin organization, and is transmitted via the gametes to alter phenotypes across
generations. Uncovering how genomic information is organized and regulated through epigenetic
processes to control gene expression and cell functions in the next generation is still in a nascent stage.
We and others have shown that errors in epigenomic profiles in sperm can be induced by environmental
exposure to toxicants such as those in insecticides and plastics, obesity, and poor diet (Lambrot *et al.*,
2013; Radford *et al.*, 2014; Donkin *et al.*, 2016; Wu *et al.*, 2017; Lismer, Dumeaux, *et al.*, 2021; Pilsner *et
al.*, 2021). We recently demonstrated that these epigenome changes at the level of chromatin can be
transmitted via sperm to alter embryonic gene expression, development, and offspring health (Lismer,
Dumeaux, *et al.*, 2021). Historically, parental health and fertility have focused predominantly on the mother,
although it is clear a father's health and lifestyle can also impact his children's health. How epimutations in
sperm functionally impact the embryo urgently require elucidation to prevent transmission of disease from
father to offspring.
57
58   Metabolic disease including obesity and type II diabetes can in part be attributed to genetic factors with a
59   5-10% increased risk (Voight *et al.*, 2010). The remaining risk is attributable to environmental-epigenetic
60   interactions including potentially those of our ancestors. This possibility is supported by epidemiological

and animal studies. Transgenerational effects are suggested by studies in humans that linked the food supply of grandfathers to obesity and cardiovascular disease in their grandchildren (Kaati, Bygren and Edvinsson, 2002; Pembrey *et al.*, 2006; Lumey and Poppel, 2013). However, the ability for diet to induce transgenerational effects in animal models remains controversial and requires more in depth studies addressing the underlying molecular mechanisms (Siklenka *et al.*, 2015; Dalgaard *et al.*, 2016; Miska and Ferguson-smith, 2016). To date, studies using mice to assess the impact of diet and obesity in relation to the sperm epigenome, have focused on the DNA methylome and ncRNA as the potential sperm-borne mediators of metabolic disease (Carone *et al.*, 2010; Radford *et al.*, 2014; Grandjean *et al.*, 2015; Chen *et al.*, 2016; Cropley *et al.*, 2016; de Castro Barbosa *et al.*, 2016; Sharma *et al.*, 2016). The role of sperm chromatin in the non-genetic inheritance of metabolic disorders is unknown. In human and mouse sperm, histone H3 lysine 4 trimethylation (H3K4me3) localizes to genes involved in metabolism and development (Lambrot *et al.*, 2019; Lambrot, 2021; Lismer, Dumeaux, *et al.*, 2021). Moreover, sperm H3K4me3 can be altered by folate deficiency and influences embryonic development and gene expression (Siklenka *et al.*, 2015; Lismer *et al.*, 2020). This association of histone modifications in sperm with offspring phenotypes has since been confirmed in other mouse models (Stringer *et al.*, 2018; Lesch *et al.*, 2019). Based on these observations, we hypothesized that sperm H3K4me3 may serve as a metabolic sensor that is implicated in the paternal transmission of obesity-associated disease in offspring.

A focus of this study was to identify whether paternal obesity impacts the F_1-F_2, and if so, to identify potential mechanisms of this non-genetic inheritance. In our transgenic (TG) mouse model of epigenetic inheritance, male mice over-express the histone demethylase KDM1A (TG) specifically in spermatogenesis, resulting in sperm with alterations in H3K4me2 and me3. Of note, only H3K4me3 has been implicated in transgenerational inheritance in this mouse model (Lismer *et al.*, 2020). Therefore, as sperm H3K4me3 is responsive to paternal folate deficiency (Lismer, Dumeaux, *et al.*, 2021), and has been implicated in transgenerational inheritance (Lismer *et al.*, 2020), we targeted this mark in sperm to probe in response to paternal obesity and as a potential mediator of inheritance. In this study, we aimed to: 1) assess the impact of high-fat diet (HFD) induced paternal obesity on sperm H3K4me3 and its association with metabolic dysfunction across generations, and 2) determine if descendants of obese TG sires with a previously altered sperm epigenome would show more severe metabolic dysfunction? To address these aims, we used wildtype (WT), or the germline specific KDM1A-overexpressing TG mice, in combination with a diet-induced obesity. These TG sires have descended from males that have an altered sperm epigenome and whose ancestors had compromised health (see Materials and Methods for details). This TG model is used to represent an at-risk population that may be more susceptible to poor health when challenged with obesity. Here, we demonstrate that a paternal high-fat diet induces F_0 obesity and metabolic dysfunction in the F_1. Remarkably, transgenerational phenotypes were only observed in descendants of obese KDM1A TG males, and this was associated with enhanced alterations in H3K4me3 enrichment in obese TG sperm. This suggests that the risk of transgenerational disease transmission may be greater if an ancestor has had

prior exposures that cause pre-existing damage to the sperm epigenome. Concordant with the metabolic

phenotypes observed in offspring, obesity-induced alterations in sperm H3K4me3 occurred at genes

involved in development, placenta formation, inflammatory processes, glucose and lipid metabolic

pathways. These sperm altered H3K4me3 regions persist in the embryo and placenta, supporting a role for

sperm H3K4me3 in paternal origins of adult-onset metabolic disorders.

Results

Paternal obesity induces metabolic phenotypes in a sex-specific manner that are enhanced in

KDM1A F_1 and F_2 transgenic descendants

Impact of paternal obesity on offspring bodyweight and fat accruement

Beginning at weaning until 20 weeks, inbred C57BL/6NCrl control mice (WT), or KDM1A heterozygous

transgenics (TG) were fed either a calorie-dense high-fat diet (HFD; 60% kcal fat), or a sucrose- and

vitamin-matched low fat control diet (CON; 10% kcal fat) (Fig. 1A-C and Table S1). Table S2 provides the

animal numbers by sex, generation, and genotype for each metabolic analysis. In the 2 weeks post-

weaning, F_0 males on the HFD consumed more calories and gained significantly more weight than CON

males irrespective of genotype (Fig. S1A-B). These effects persisted throughout the diet intervention (Fig.

S1A-C), with TG HFD males weighing the most at 4 months (Fig. S1C_i). This trend continued in the TG

male F_1 and F_2 descendants (fed regular chow), with weights being significantly more than the F_1 and F_2 of

TG CON and WT HFD (Fig. S1C_{ii-iii}). Indicating sex-specific responses to paternal obesity, in female

descendants the changes in body weight and fat deposition differed from males (Fig. S1C-E). For example,

female F_1 HFD descendants showed no changes in body weight, while F_2 TG HFD females weighed more

in comparison to F_2 WT HFD females (Fig. S1C_{iv-v}, respectively). Unexpectedly, F_2 WT HFD females

weighed less than F_2 WT CON. To assess fat accruement, we measured visceral mesenteric (mWAT) and

gonadal white adipose tissue (gWAT). All male (F_0) on the HFD accumulated more mWAT compared to

CON males, with no genotype effect (Fig. S1D_i). Male and female F_1 offspring sired by WT HFD or TG HFD

had increased mWAT fat mass compared to WT CON and TG CON (Fig. S1D_{ii} and S1D_{iv}, respectively).

Strikingly, mWAT stores were greater in TG HFD F_1 and F_2 males and females compared to WT HFD

descendants (Fig. S1D_{ii-v}). Gonadal fat depots in F_0 males were not impacted by the HFD (gWAT; Fig. S1E_i),

while male WT HFD F_1 showed increased gWAT, and TG HFD F_1 did not (gWAT; Fig. S1E_{ii}). In the male

F_2, TG HFD descendants had increased gWAT, but not WT HFD (Fig. S1E_{iii}). Consistent with the unusual

decrease in body weight in female F_2 WT HFD vs F_2 WT CON, mWAT and gWAT were also reduced in

female F_2 WT HFD (Fig. S1D_v and S1E_v, respectively). Perhaps indicative of a sex-specific over-correction

in the female F_2 to grand-sire obesity. Like for body weight and mWAT, female F_2 TG HFD had increased

gWAT in comparison to WT HFD (Fig. S1E_v). Overall analysis of body weight and fat accruement revealed

sex-specific responses in descendants with transgenerational effects of paternal obesity being detected

only in the TG HFD descendants of both males and females.

Impact of paternal obesity on glucose homeostasis

Next, we assessed glucose metabolism and insulin sensitivity by glucose tolerance (GTT), and insulin tolerance tests (ITT). These were conducted following the standard operating procedures of the NIH mouse Metabolic Phenotyping Center (Ayala *et al.*, 2010). First, we assessed the effects of the HFD on baseline fasting blood glucose. Consumption of a HFD resulted in elevated baseline glucose in male (F_0) WT HFD and TG HFD in comparison to WT CON and TG CON, respectively (Fig. S2Ai). Male TG HFD descendants (F_1 and F_2), but not WT HFD descendants had significantly elevated fasting blood glucose (Fig. S2A$_{ii-iii}$). In contrast, the glycemic status of all descendant females (F_1 and F_2) did not differ (Fig. S2A$_{iv-v}$). The same animals used to assess baseline glucose were then given an intraperitoneal glucose challenge and the rate of glucose disposal measured. Analysis of GTT data showed that F_0 WT HFD and TG HFD were glucose intolerant following glucose injection in comparison to F_0 CON males as shown by their elevated blood glucose curves that persisted across the time course of 15-120 min (Fig. 1D$_i$). Note that at T0, before the glucose challenge, F_0 WT HFD males had significantly elevated blood glucose in comparison to F_0 WT CON which is in line with the baseline glucose measures (Fig. 1D$_i$ and Fig. S2Ai). In contrast, the T0 blood glucose levels did not differ between male F_0 TG CON and F_0 TG HFD even though baseline measures after fasting were significantly different (Fig. 1D$_i$ and Fig. S2Ai). These non-congruent findings are attributable to multiple testing correction to adjust for the high number of simultaneous tests conducted in the analysis of GTT curves. Intergenerational effects of the paternal HFD were apparent as male (F_1) sired by WT HFD and TG HFD were also glucose intolerant (Fig. 1D$_{ii}$). Indicating that there was enhanced metabolic disruption in the male F_1 TG HFD, elevated glucose levels persisted across the GTT time-course for the F_1 TG HFD but not the F_1 WT HFD. Interestingly, glycemic response impairments persisted in the F_2 generation of male descendants of TG HFD only (Fig. 1D$_{iii}$). Although fat measures were impacted in female F_1 and F_2 HFD, they did not exhibit glucose impairment (Fig. 1D$_{iv-v}$). Analysis of the area under the curve (AUC) for the GTT was consistent with the male and female glycemic responses shown in the glucose curves (Fig. S2B$_{i-v}$). In line with the observed glycemic responses, the insulin tolerance test and the corresponding AUC demonstrated that male F_0 WT HFD and TG HFD were insulin insensitive (Fig. 1E$_i$ and S2C$_i$). Analysis of the AUC indicated that F_1 WT HFD and F_1 TG HFD were insulin insensitive (Fig. S2C$_{ii}$). Like the glucose tolerance test, there were more pronounced impairments revealed by the ITT for the F_1 TG HFD in comparison to the F_1 WT HFD and only the F_2 TG HFD showed impaired whole body insulin action (Fig. 1E$_{iii}$ and Fig. S2C$_{ii-iii}$). Like the GTT, there was no indication of insulin impairment in female HFD F_1 nor F_2 (Fig. 1E$_{iv-v}$, S2C$_{iv-v}$).

To summarize, the effects of paternal high-fat diet on glucose homeostasis were sex-specific; male descendants had impaired glucose homeostasis, whereas females did not. Many factors can influence the outcomes of GTT and ITT measures such as the length of fasting, age and strain of mice (Ayala *et al.*, 2010; Bowe *et al.*, 2012). It is worth considering that even though female descendants of HFD (F_{1-2}) showed alterations in fat accretion but not in glucose homeostasis, there remains the possibility that findings may have been different had measurements been taken after a reduced fasting period, or beyond 4 months of

age. Taken together, the assessments of weight and metabolic testing indicate that the TG descendants

had enhanced responses to paternal obesity in comparison to WT descendants.

Paternal obesity was associated with altered liver gene expression in the F_0-F_1 with unique genes

being differentially expressed in KDM1A descendants (F_1-F_2)

Obesity contributes to pathophysiological changes in gene expression in liver gene expression (Kim *et al.*,

2004). To determine whether the altered metabolic status of HFD sires and their descendants (F_1-F_2) was

associated with differential gene expression in the liver, we performed RNA-sequencing on the left lateral

lobe (*lobus hepatis sinister lateralis*) of adult mice (F_0-F_2). Sequencing quality was high with most RNA

profiles having a Pearson correlation coefficient > 0.8 (Fig. S3A). Interestingly, principal component analysis

of sequencing data revealed distinct hepatic transcriptomic profiles between males and females that was

independent of experimental group and genotype (Fig. S3C). We compared hepatic transcriptome profiles

by diet, sex, genotype and generation using a gene-level analysis at single-transcript resolution (Yi *et al.*,

2018). As expected, obesity was associated with differential liver gene expression. Liver from obese F_0 WT

males showed differential expression of 2,136 genes in comparison to non-obese F_0 WT males (Fig. 2A,

Lancaster p<0.05). Similarly, when comparing obese F_0 TG to non-obese F_0 TG, 1,476 genes were

differentially expressed (Fig. 2B, Lancaster p<0.05). Of these differentially expressed genes (DEGs), 448

were commonly altered by obesity in both the F_0 WT and F_0 TG (p<0.0001; Fig. 2i). To identify which genes

were altered due to genotype, we compared WT obese to TG obese and identified 524 DEGs, suggesting

that obesity had a unique effect in TG mice due to an interaction between diet and genotype (Fig. 2C,

Lancaster p<0.05).

192   To determine if the effects of paternal obesity on liver function were intergenerational, we compared

the liver transcriptome of male and female F_1. In comparison to F_1 WT CON and TG CON males, livers of

F_1 WT HFD and TG HFD, showed differential expression of 1,015 and 794 genes (Fig. 2D and 2E,

respectively, Lancaster p<0.05). A total of 165 DEGs overlapped between F_1 WT and TG (p<0.0001; Fig.

2ii). Of the DEGs between the WT CON and HFD in the F_1, 139 were the same deregulated genes as

identified in the F_0 WT CON vs HFD males (p=0.76; Fig 2iv). Similarly, there were 103 shared transcripts

identified as differentially expressed between the F_1 TG CON vs HFD, that were also altered in the F_0 TG

CON vs HFD (p=0.003; Fig 2v). This suggests that a common set of genes maintain dysfunction as a

consequence of direct exposures to obesity and these changes are maintained in the non-obese F_1. When

comparing genes altered by genotype in the F_1 (WT HFD vs TG HFD), 961 were significantly altered (Fig

2F, Lancaster p<0.05), with 78 overlapping DEGs between the F_0 and the F_1 (p<0.0001; Fig 2vi). The

overlap in deregulated genes between the F_0 and F_1 indicates that the metabolic phenotypes generated by

the paternal HFD persist intergenerationally despite the F_1 being fed a regular chow diet. There were also

novel genes in the F_1 that were deregulated, suggesting secondary intergenerational effects of the paternal

HFD. As expected, there was a reduction in the number of deregulated genes in the F_1 WT and TG males

compared to the F_0 which is consistent with the milder metabolic phenotypes observed in the F_1 in

comparison to the directly HFD-exposed F_0. Likely reflecting the enhanced genotype influence of the F_0 TG

HFD sires that resulted in more severe weight changes and higher baseline glucose in their descendants

compared to the WT HFD descendants, there were more DEG in the male F_0 compared to the male F_1 TG

HFD (524 in F_0 versus 961 in F_1; Fig. 2_{vi}).

The last comparisons in liver transcriptomes were between the F_1 male and female. Despite the

female F_1 having no metabolic phenotype detected by our measures, there was significantly altered gene

expression in the livers of F_1 female offspring of WT HFD vs WT CON sires (830; Fig 2G, Lancaster $p<0.05$)

Of these, 153 were in common with the F_1 male WT HFD sired offspring ($p<0.0001$; Fig 2_{vii}). Likewise, the

F_1 female sired by TG HFD had 1,125 DEGs in comparison to females sired by TG CON (Fig. 2H, Lancaster

$p<0.05$) with 148 in common with F_1 male TG HFD sired offspring ($p<0.0001$; Fig. 2_{viii}). Of these altered

transcripts, 160 were in common between F_1 female descendants of WT HFD and TG HFD ($p<0.0001$; Fig.

2_{iii}). Like the F_1 male TG HFD offspring, there were unique transcripts altered in F_1 female TG HFD offspring

(1,370; Fig. 2I, Lancaster $p<0.05$), with 181 differentially expressed in both F_1 males and females

($p=<0.0001$; Fig. 2_{ix}). These may reflect genes impacted by genotype regardless of sex. An interesting

finding from the F_2 phenotyping was those transgenerational metabolic effects of the HFD were only

detected in the male descendants of TG. Therefore, we only profiled F_2 male livers by RNA-seq. This

analysis revealed differential expression of 2,141 genes between the F_2 WT HFD and TG HFD (Fig. 2J,

Lancaster $p<0.05$) with 129 overlapping with the F_1 WT HFD vs TG HFD males ($p=0.06$; Fig 2_x). The number

of differentially expressed genes increased every generation in comparisons between the WT HFD and the

TG HFD ($F_0=524$, $F_1=961$, $F_2 = 2,141$). This sustained deregulated gene expression in the livers of TG

HFD F_2, matches the enhanced metabolic phenotypes observed in only F_2 TG HFD males but not in the F_1

WT HFD. It also indicates a dilution in phenotype by reprogramming in the WT HFD descendants.

Paternal diet-induced obesity disrupts gene expression in functional processes that differ

between genotypes, sex and generations

To gain insight into the physiological implications of obesity-induced altered hepatic transcriptomes, we

used a gene ontology (GO) approach combined with functional similarity clustering to compare processes

in the liver impacted by diet across genotype and sex, and those impacted by genotype across generation

(Fig. 3A-C, Supplemental files 1-3 and Table S5-7) (Brionne, Juanchich and Hennequet-Antier, 2019).

Interactive heatmaps that facilitate in-depth probing of the gene frequency and the $-log_{10}$ p-value of enriched

GO terms within each cluster are found in Supplemental files 1-3. The non-interactive heatmaps are shown

in Fig. 3. Overall, there were similar processes altered by obesity in F_0 WT and TG livers, including lipid,

amino acid, and small molecule metabolism (Fig. 3A, Supplemental file 1 and Table S5; clusters 1-5),

homeostasis and environmental responses (clusters 8-10), and cellular differentiation and signalling

(clusters 11-13). However, the gene frequency (# of genes annotated to that process) within processes

differed by genotype. For example, genes involved in lipid metabolic processes (cluster 2), specifically the

carboxylic acid metabolic process was more enriched at 19.03% gene frequency in the F_0 WT HFD ($-log_{10}$

p-value 6.18), than in the F_0 TG HFD at 14.7% gene frequency (-$\log_{10}$ p-value 4.37). In contrast, genes

involved in environmental responses (cluster 8) such as to alcohol, were similarly enriched in both the F_0

WT HFD with a gene frequency of 19.19% (-$\log_{10}$ p-value 1.3) and of 19.19% in the F_0 TG HFD (-$\log_{10}$ p-

value 2.29). Interestingly, deregulated genes involved in insulin, protein and metal ion transport were only

enriched in F_0 TG HFD livers (clusters 6-7).

When the altered functional pathways were compared between F_1 WT CON vs WT HFD males and

females, there were clear impacts of obesity on the liver functional pathways of offspring, and these differed

by sex (Fig. 3B, Supplemental file 2 and Table S6). Reflecting sex differences, a greater number of GO

terms related to inflammation (cluster 4), and cell cycle, differentiation and signalling regulation (clusters

10-11) were significantly enriched in males compared to females. Of note, genes involved in the regulation

of proinflammatory cytokines were particularly enriched in males but not females (clusters 4 and 6). This

concurs with the more severe phenotypes observed in the males. Conversely, genes involved in DNA/RNA

biosynthesis, transcription factors and telomere activity (clusters 1-3), and macromolecule and nitrogen

metabolism (cluster 5) were more enriched in females. Interestingly, pathways associated with chromatin

and cellular organization and protein metabolism (clusters 8-9) were differentially enriched by paternal

obesity in both sexes.

Next, we compared the intergenerational and transgenerational effect of the interaction between

the KDM1A transgene with obesity in terms of differences in process enrichment across generations when

comparing F_{0-2} WT HFD with F_{0-2} TG HFD (Fig. 3C, Supplemental file 3 and Table S7). Reflecting the

increasing generational changes in liver gene expression in the TG HFD descendants, there was an

increase in the number of significantly enriched GO terms when comparing WT HFD vs TG HFD across

generations (F_0 male=79; F_1 male=118; F_1 female=159; F_2 male=206; Supplemental file 4). This finding is

concordant with the metabolic phenotypes detected in the TG HFD male F_2 descendants, but not in the F_2

male WT HFD (Fig. 1, Fig. S1, Fig. S2; Fig. 3C and Supplemental file 3). An interesting finding that supports

a unique genotype response in the male F_1 TG HFD in comparison to the male F_1 WT HFD, is the over-

representation of genes implicated in chromatin remodelling and transcription (clusters 17-19). This may

reflect the activation of genetic response to reprogram the KDM1A-diet-induced sperm epimutations. In

addition, there was an enrichment in the differential expression of genes with functions related to

inflammation and environmental response (clusters 3-5), and metabolic processes (clusters 11-14). Genes

involved in lipid catabolic processes (cluster 13) were increasingly enriched across generations, with 3.54%

(F_0), 6.69% (F_1 male), 8.66% (F_1 female) and 20.87% (F_2) gene frequency. Like the males, female F_1 WT

HFD vs F_1 TG HFD showed differences in enrichment pathways that reflect the interaction between obesity

and the KDM1A transgene in the F_0 and the sex differences observed in the phenotypes (Fig. 3C)

Obesity in combination with expression of the KDM1A transgene increases differential enrichment

of sperm H3K4me3 at genes involved in metabolism and development

Previously we associated paternal folate deficient diets with altered H3K4me3 in sperm and showed that these alterations are transmitted to the embryo and in line with offspring phenotypes (Lismer et al., 2021). We hypothesized that the sperm epigenome at the level of H3K4me3 would be altered by obesity and that this effect would be enhanced in KDM1A TG males with pre-existing alterations in sperm H3K4me3. To test these hypotheses, we performed native chromatin immunoprecipitation followed by deep sequencing (ChIP-seq) targeting histone H3K4me3, using sperm from individual WT or TG males fed either a CON or HFD (N=5 per experimental group, on average 33.3 million reads per sample with an alignment rate of 97%, Table S8). H3K4me3 localized to 30,745 genomic regions, with a Spearman correlation coefficient of 0.98 between samples (Fig. 4A and S4). Principal component analysis of H3K4me3 profiles revealed a clear separation of samples according to dietary treatment within genotype groups (Fig. 4 B-C). WT samples separated along Principal Component 1 (PC1) with 37.41% of variance attributed to diet (Fig. 4B; PERMANOVA, permutation-based p=0.01). TG samples separated on PC1 with 32.68% of the variability, with diet as the second source of variance (PC2), at 25.56% (Fig. 4C; PERMANOVA, permutation-based p = 0.009).

To focus our analysis on the regions most impacted by diet we selected the top 5% differentially enriched H3K4me3 regions (deH3K4me3, n=1,538) in each genotype (PC1 in WT, PC2 in TG) (Fig. 4Di-iv). The genome distribution analysis for specific annotations showed that obesity-sensitive H3K4me3 regions were predominantly located in CpG islands, promoters, exons, and intergenic regions (Fig. S5). To a lesser extent, deH3K4me3 also occurred at transposable elements (LINE, SINE and LTR), where epigenetic de-repression is associated with the use of alternative promoters and long- and short-range enhancers that are implicated in embryo development and pluripotency (Gerdes *et al.*, 2016) (Fig. S5). Representative genome browser tracks (Fig. S5) showing enrichment gains and losses for H3K4me3 at gene promoters are shown for *Pde1c* (phosphodiesterase 1C; affects the olfactory system), *Bcdin3d* (RNA methyltransferase; highly expressed in embryonic development), *Sh2d4a* (Sh2 domain containing protein 4A; expressed during development and associated with endocrine and liver function), and *Col15a1* (collagen alpha-1; cell differentiation and development, endocrine, cardiovascular system and more) (*Mouse Genome Informatics*, no date).

Next, we compared the regions of H3K4me3 that were altered by obesity, their genomic location, directionality change and functions between diets and genotype (Fig. 4). As a response to obesity, H3K4me3 enrichment gains were more predominant than losses for both F_0 WT HFD and TG HFD (Fig. 4D). In the WT HFD 1,323 regions gained and 215 lost H3K4m3 in comparison to the WT CON (Fig. 4Di-ii). Similarly, in the F_0 TG HFD sperm, 1,067 regions gained and 471 lost H3K4me3 in comparison to the F_0 TG CON (Fig 4Diii-iv). Regions with deH3K4me3 in WT HFD had an 15.6% overlap (240/1,538 regions) with those of TG HFD (Fig. 4E). Of those common 240 regions, 162 had the same directionality change in both WT and TG HFD, with 159 regions with a gain and 3 regions with a reduction in H3K4me3 enrichment (Fig. 4 F and G, respectively). The non-overlapping regions of deH3K4me3 in WT HFD and TG HFD sperm could be a consequence of genetic-epigenetic interactions where the TG mice respond uniquely to obesity

as was observed in the phenotypic characterization. The proximity to the TSS of the deH3K4me3 regions
in sperm altered by obesity in the F_0 WT HFD and TG HFD were similar (Fig. 4H).

Next, we performed a gene ontology (GO) enrichment analysis on promoters to gain functional
insight into the genes disturbed by obesity and how they may relate to offspring phenotypes. This analysis
revealed that changes in H3K4me3 occurred at promoters of genes that function in reproduction and
development which is expected as these processes are known to be enriched for H3K4me3 in sperm (Fig.
4I and Tables S7-10). Supporting the idea that deH3K4me is specifically impacted by obesity and may be
implicated in the transmission of metabolic disease to offspring is the finding that enriched genes were
identified in processes related to metabolism and the impact of obesity. These included inflammatory
processes, glucose and lipid metabolic pathways, and one-carbon cycle metabolism (Fig. $4I_{i-iv}$; Tables S7-
10). Notably, some of the significantly enriched pathways are concordant with disturbed metabolic
phenotypes of the F_0-F_2 including, for example, carbohydrate metabolic processes, glycolysis, growth
hormone signaling and insulin signaling (Fig 4I, Tables S7-10).

The metabolic phenotypes of WT HFD and TG HFD descendants were similar, although the F_{1-2}
TG HFD showed enhanced abnormalities. We hypothesized that these differences in offspring metabolic
disturbances may relate to the degree of H3K4me3 alteration in F_0 sperm, the directionality of the change
(gain versus loss), and the functionality of genes bearing alterations. Interestingly, when comparing WT
CON with TG HFD sperm, samples separated along PC2, with 26.69% of variance associated with
genotype and diet (Fig. 5A; PERMANOVA, permutation-based p=0.006). Of the top 5% impacted regions
selected (n=1,538), a greater proportion showed a gain of enrichment for H3K4me3 in TG HFD sperm in
comparison to WT CON (Fig. 5B, n=1,071 regions with gains; Fig. 5C. n=467 regions with losses). We
analyzed the detected regions impacted by genotype and diet (n=1,538) for differential enrichment to
determine whether obesity in combination with KDM1A overexpression led to greater changes in H3K4me3
enrichment. This analysis identified 264 regions with a significant linear trend, where TG HFD sperm
showed a greater degree of change in enrichment, and TG CON and WT HFD showed intermediate
changes in comparison to WT CON (Fig. 5D-E, adjusted p<0.2). There were only 9 significant regions with
further increase in H3K4me3 in the TG HFD (Fig. 5 D), whereas 255 regions showed a greater loss of
H3K4me3 enrichment in the TG HFD (Fig. 5E). Consistent with the stronger metabolic phenotypes
observed in the TG HFD F_{1-2}, the functional analysis of the promoters showing significant linear trends
(n=104) for H3K4me3 across experimental groups occurred at genes implicated in metabolic and
cardiovascular disease progression (Fig. 5F, Table S13).

Paternal obesity impacts sperm H3K4me3 at regions that coincide with open chromatin and gene
expression in pre-implantation embryos
We recently demonstrated that sperm H3K4me3 is transmitted to the embryo and associated with gene
expression (Lismer, Dumeaux, *et al.*, 2021). We hypothesized that obesity-altered sperm H3K4me3 is
transmitted and associated with chromatin accessibility in the early embryo, which in turn could influence

gene expression and offspring phenotypes. To assess this possibility, we investigated the relationship between deH3K4me3 in sperm in relation to H3K4me3 in the embryo, the oocyte and open chromatin (Zhang *et al.*, 2016; Jung *et al.*, 2017; Liu *et al.*, 2019). We used existing datasets from mouse 2-cell embryos (Zhang *et al.*, 2016), generated by crossing males and females of different strains, permitting the assignment of reads to the paternal-specific allele. In line with a preferential paternal contribution of H3K4me3 to the 2-cell embryo, regions enriched for H3K4me3 in sperm, including those altered by obesity are not enriched in the oocyte (Fig 6A).

Next, we examined the relationship between sperm H3K4me3, chromatin accessibility and embryonic gene expression at the 4-cell and morula stages (Fig. 6 and Fig. S6A). We focused on the 4-cell and morula stages as this is when large-scale embryonic transcription directs development (Jukam, Shariati and Skotheim, 2017). There is an association between sperm H3K4me3 and embryonic gene expression (Lismer *et al.*, 2020; Lismer, Dumeaux, *et al.*, 2021) (Fig. S6 A_i). Strikingly, sperm H3K4me3 including obesity-sensitive regions are associated with ATAC-seq signal in pre-implantation embryos (Fig. 6A and 6B). To determine the functional relationship between the H3K4me3 obesity-altered regions and embryonic gene expression, we compared these with 4-cell and morula expressed genes and performed a gene ontology analysis. Of the sperm deH3K4me3 regions overlapping promoters (n=738), 51.8% (n=382) are expressed in the 4-cell embryos, 44.3% (n=327) are expressed in the morula embryos, and 39.7% (292) overlap in both (Fig. S6A_{ii}). To gain insight into what obesity-altered H3K4me3 associated genes in sperm relate to embryonic gene expression, we performed a GO analysis on the regions that are deH3K4me3 in sperm and the corresponding genes expressed in 4-cell and morula embryos (Fig. 6C_{i-ii}). Again, supporting a role for sperm H3K4me3 in paternal transmission of metabolic disease, with both the 4-cell and the morula gene processes significantly enriched specific to metabolism (Fig. 6C_{i-ii} and Tables S12-13).

Unlike our previous studies on the function of H3K4me3 sperm where we found strong associations between deH3K4me3 and birth defects (Lismer *et al.*, 2020; Lismer, Dumeaux, *et al.*, 2021), in this study the defects in descendants appear to be limited to metabolic dysfunction and not development. To further probe the bias of obesity altered H3K4me3 for genes that function in metabolism and not post-implantation development, we examined the relationship between deH3K4me3 and H3K4me3/H3K27me3 bivalency. Genes that are bivalent for H3K4me3/H3K27me3 in sperm and embryos are termed as poised genes and are expressed later in embryo development (Hammoud *et al.*, 2009; Brykczynska *et al.*, 2010). Some promoters with high H3K4me3 enrichment in sperm are associated with bivalency in the 2-cell embryos (Fig. S6B_i). The obesity-altered H3K4me3 regions are predominantly not associated with H3K4me3/H3K27me3 bivalent genes in the embryo (Fig. S6C_{ii}). Taken together these findings suggest a preferential contribution of H3K4me3 on the paternal chromatin in the early embryo that includes obesity-sensitive regions that may be instructive of metabolic-associated gene expression and a direct route for epigenetic inheritance.

HFD alters the sperm epigenome at regions instructive for placenta development

The placenta is a key extra-embryonic organ that represents the uterine-fetal interface and plays a
central role in energy allocation, nutrient exchange, and developmental progression. Placental
abnormalities have been linked to late onset cardiometabolic diseases, highlighting the importance of the
in utero environment for adulthood metabolic health (Perez-Garcia *et al.*, 2018). Our gene ontology analysis
on diet-induced deH3K4me3 regions in sperm revealed significant enrichment of genes involved in placenta
development (Fig. 4I and Tables S7-10). Given the sperm epigenome influences placental gene expression
(Wang *et al.*, 2013), we were interested in the prospect that diet-induced epimutations in sperm affect
placenta gene expression that could influence metabolic phenotypes across generations. To investigate
this possibility, we used existing H3K4me3 and transcriptomic datasets from mouse trophectoderm – the
embryonic precursor of placenta lineage – and placenta (Shen *et al.*, 2012; Wu *et al.*, 2016; Chu *et al.*,
2019). We compared the enrichment profiles of H3K4me3 in sperm, trophectoderm and placenta, at all
H3K4me3-enriched regions in sperm (n=30,745) and at those sensitive to diet (n=2,836). Most regions
showed presence of H3K4me3 in both the trophectoderm and the placenta (Fig. 7A). Of the 738
deH3K4me3 regions localizing to promoters in sperm, 56.8% (n=418) were expressed in the
trophectoderm, 76.8% (n=567) were expressed in the placenta, and 54.6% (n=403) were expressed in both
(Fig. S6C$_{ii}$). Gene ontology analysis of the shared H3K4me3 in sperm with TE and placenta revealed that
there was an association with placenta function including at deH3K4me3 regions (Fig. 7Bi and iii, Tables
S14 and S16). The GO analysis of the H3K4me3 regions that were not common with TE and placenta were
involved in spermatogenesis, fertilization and sperm function (Fig. 7Bii and iv, Tables S15 and S17).

Next, we compared gene enrichment of sperm H3K4me3 with low- and high-expressed genes in
the TE and placenta. Suggesting an influential role of sperm H3K4me3, the highly expressed genes and to
a lesser extent the lowly expressed genes in placenta were positively correlated with sperm H3K4me3 (Fig.
7Ci-iv). Notably when the same comparisons were made with the deH3K4me3 there was a significant
relationship with both lowly- and highly-expressed placenta genes (p=1.2e-11 and p=0.008, respectively;
Fig. 7Cv-viii). In addition, the GO analysis of TE- and placenta-expressed genes that overlap with
deH3K4me3 promoters are in line with the metabolic phenotypes in offspring (Fig. 7D i-ii, Tables S18-19).
Taken together this analysis raises the possibility that obesity-induced alterations in sperm may influence
embryonic and placenta gene expression to alter metabolic function of offspring.
420

Obesity-induced sperm epigenomic and hepatic transcriptomic alterations are unrelated
In a recent study, paternal low-protein diet was associated with reduced H3K9me2 at genes in sperm and
were suggested to modulate gene expression profiles in the liver (Yoshida *et al.*, 2020). We aimed to assess
whether a similar association between obesity-induced deH3K4me3 in sperm would relate to differential
expression in the livers of the next generation. We focused on the obesity-associated sperm deH3K4me3
at promoters in F_0 sires and their relationship to differentially expressed genes in the liver (DEGs) of F_1
males. This analysis revealed that genes with differential expression in livers (n=1,644) were by in large
unrelated to genes bearing deH3K4me3 in sperm. Only 9.1% (n=67) of promoters with deH3K4me3 in

sperm were differentially expressed in the liver of F_1 males sired by HFD-fed sires (Fig. S7 A-B). We then

asked if deH3K4me3 promoters in sperm and liver DEGs had related biological functions. Strikingly, sperm-

and liver-altered genes showed few functional similarities (Fig. S7C, Supplemental file 5 and Table S22).

Functional pathways specifically enriched in deH3K4me3 promoters involved development and

differentiation processes (clusters 12-15). As expected in a paternal obesity model, gene processes altered

in offspring livers included: regulation of transcription and RNA splicing (clusters 1-3), protein and histone

post-translational modifications (clusters 4-5), and metabolism of lipid, nitrogen and glucose (clusters 6-8).

Pathways enriched in both the deH3K4me3 promoters in sperm and the DEGs in liver were involved in cell

cycle, transport and signaling (clusters 16-19), and response to stress and inflammation (clusters 20-22).

These commonly enriched pathways might reflect obesity-associated systemic inflammation which could

affect multiple organs in a similar manner. These findings indicate that paternal obesity alters the sperm

epigenome at distinct genes and functional pathways than those differentially expressed in offspring livers

and fits with a developmental origin of adult metabolic dysfunction that could be related to alterations in

gene expression in the embryo and placenta.

Discussion

445
Epidemiological studies and animal models point to the world-wide increases in childhood obesity and

diabetes being in part attributed to environment-epigenetic interactions in the gametes (Huypens *et al.*,

2016; Isganaitis, Suehiro and Cardona, 2017; Sales, Ferguson-Smith and Patti, 2017). With the prevalence

of childhood obesity predicted to reach 70 million by 2025 (Brown *et al.*, 2015), it is urgent to gain a better

understanding of the underlying heritable mechanisms. Studies of obesity in mice and men have focussed

on the role of sperm DNA methylation and non-coding RNA as mediators of paternal non-genetic

inheritance of metabolic dysfunction (Carone *et al.*, 2010; Fullston *et al.*, 2013; Lambrot *et al.*, 2013;

Martínez *et al.*, 2014; Chen *et al.*, 2016; Soubry *et al.*, 2016; Youngson *et al.*, 2016; Zhang *et al.*, 2018;

Salas-Huetos *et al.*, 2021). At the mechanistic level, how sperm altered DNA methylation and non-coding

RNA lead to metabolic phenotypes in offspring remains unresolved. In contrast to the voluminous studies

on the response of sperm DNA methylation to paternal obesity/diets, there have been only a handful of

studies on the role of chromatin. The few chromatin studies have been limited by either the state of the

available technology, methodology used, or gene targeting approaches (Carone *et al.*, 2011; Palmer *et al.*,

2011; Martínez *et al.*, 2014; Donkin *et al.*, 2016).

Diets high in fat alter epigenetic programming (Ng *et al.*, 2010), likely through the alteration of

cellular metabolism, which influences the availability of methyl donors and/or the activation or inactivation

of chromatin modifying enzymes. In overweight and obese individuals, homocysteine is consistently

elevated, and associated with reduced B12 and folate (Sánchez-Margalet *et al.*, 2002; Karatela and

Sainani, 2009). In spermatogenesis there are dynamic changes to the sperm epigenome including histone

methylation which is susceptible to alterations induced by changes in methyl donor availability (Lambrot *et*

al., 2013; Lismer, Dumeaux, *et al.*, 2021). It follows that the obesity-induced alterations in H3K4me3 we

report here could be a consequence of an altered methyl donor pool. Intriguingly, the effects of obesity on the paternal epigenome were linked with the metabolic dysfunction in the descendants; deH3K4me3 occurred at the promoters of genes involved in fertility, metabolism, and placenta processes. The association with metabolic processes is apparently specific to the HFD and obesity. In our analysis of sperm from folate deficient males, deH3K4me3 was associated more with developmental processes such as skeletal formation which aligned with the developmental defects observed in the offspring (Lismer, Dumeaux, *et al.*, 2021). Indicative of paternal transmission of sperm altered H3K4me3 as a mechanism of metabolic dysfunction was the strong relationship between deH3K4me3, an open chromatin state and gene expression in embryos and placenta.

The enhanced metabolic abnormalities observed in the descendants of obese F_0 TG revealed an increased susceptibility to metabolic disease in the TG line. An explanation for this response is that the F_0 TG were descendants from a lineage with pre-existing alterations in the sperm epigenome due to the genetic modification causing KDM1A overexpression. This genetic stress in combination with the environmental challenge of the HFD resulted in a more severely altered sperm epigenome in comparison to the WT, with consequent enhanced offspring phenotypes. Admittedly speculative, these findings suggest that the higher incidence of poor health in at-risk populations may be attributed to generational exposures to poor diet that leads to an accumulation of sperm epigenome errors that escape reprogramming.

Notably, paternal obesity-induced transgenerational metabolic disturbances in offspring were only observed in descendants of obese TG males. The phenomena of transgenerational inheritance has been most documented in genetic mouse models of epigenetic inheritance and studied in relation to DNA methylation patterns. These include the Avy agouti model (Morgan *et al.*, 1999; Dolinoy *et al.*, 2006; Cropley *et al.*, 2016), the kinky tail model (Axin[Fu] allele) (Rakyan *et al.*, 2003), and in mice bearing a mutation in the *Mtrr* gene, a folate metabolism enzyme (Padmanabhan *et al.*, 2013). In the context of environmental challenges, paternal transgenerational inheritance has been associated with altered sperm DNA methylation when there has been gestational exposure to toxicants and undernutrition (Anway *et al.*, 2005; Martínez *et al.*, 2014), and in a non-genetic pharmacologically-induced prediabetes model begun at weaning (Wei *et al.*, 2014). Taken together, this growing body of evidence indicates that transgenerational inheritance occurs under genetic influence, or when exposures coincide with developmental programming. The male F_0 mice in this study were exposed to the paternal HFD from weaning and not *in utero*, which may account for why transgenerational effects were not observed in WT HFD descendants. Another possibility is that transgenerational responses in the WT may have become detectable in older mice.

Our analysis indicates that the inherited metabolic disturbances observed in adult descendants originated early in development. In rodent models, paternal obesity and *in utero* undernutrition has been linked to altered gene expression in offspring livers and pancreatic islets with some minor links to concordant DNA methylation changes (Carone *et al.*, 2011; Martínez *et al.*, 2014; Wei *et al.*, 2014). It has been suggested that diet-associated alterations in DNA methylation in sperm are retained through embryogenesis and maintained in adult tissues mediating paternally-induced phenotypes (Martínez *et al.*,

2014; Wei *et al.*, 2014). Consistent with these studies, altered hepatic gene expression occurred in F_{1-2}

offspring of obese sires. In contrast, we observed minimal overlap of genes and functional pathways

between altered H3K4me3 enrichment in sperm, with those differentially expressed in F_1 livers. Instead, we

demonstrate a significant overlap of obese sperm H3K4me3 profiles with the expression of metabolic-

related genes in the embryo and placenta. Based on these findings, we suggest that the metabolic

phenotypes we observe originate in early embryogenesis and through changes in placental gene

expression.

There is a bounty of research linking maternal obesity to adverse metabolic consequences for the

offspring that coincide with altered placental gene expression and function (Kerr *et al.*, 2018; Franzago *et*

al., 2019). On the other hand, it is an emerging concept that the paternal environment including factors

such as diet and age can influence placental development and function. It is known that paternally

expressed genes contribute to placental growth, trophoblast invasion and insulin resistance and adiposity

(Moore, 2001; Binder, Hannan and Gardner, 2012; Wang *et al.*, 2013; Rosenfeld, 2015; Naruse *et al.*, 2019;

Michelle M Denomme *et al.*, 2020; Michelle M. Denomme *et al.*, 2020). In humans, errors in epigenomic

programming have been associated with gestational trophoblast disease and pre-eclampsia, but the role

of the obese father in these conditions has been entirely unexplored (Gabory, Attig and Junien, 2011;

Nelissen *et al.*, 2011). Previous studies support a connection between paternal diets, obesity, and placental

dysfunction as a developmental route to metabolic disease in children. For example, we have shown that

a folate deficient paternal diet and altered sperm DNA methylation coincided with deregulated placenta

gene expression of *Cav1* and *Txndc16* (Lambrot *et al.*, 2013). Moreover, paternal obesity in mice has been

attributed to defective placental development (Binder, Hannan and Gardner, 2012; Binder *et al.*, 2015). In

women, altered DNA methylation in the regulation of some genes in preeclampsia has been established.

However, many genes with deregulated expression were not associated with DNA methylation raising the

possibility of altered chromatin signatures leading to abnormal gene expression in this placental disorder

(Leavey *et al.*, 2018). Indeed, upregulated expression of LncRNA by increased H3K4me3 has been

observed in preeclampsia placentas (Sun *et al.*, 2020), and the levels of H3K4me3 as detected by

immunocytochemistry are decreased (Meister *et al.*, 2021). Until now the connection between sperm

chromatin and placenta function has been unexplored. Our analyses revealed that most of the obesity-

altered H3K4me3 at promoters occurred at loci involved in placental development and inflammatory

processes (56.6% and 76.8% of deH3K4me3 occurred at promoters expressed in the trophectoderm and

placenta, respectively). Remarkably, deregulated expression of genes implicated in inflammation have

been implicated in hypertensive disorders in pregnancy including pre-eclampsia. This raises the possibility

that the paternal sperm epigenome may influence maternal health during pregnancy in addition to that of

the developing fetus. Hypertensive disorders in pregnancy have been associated with increased risk for

developing cardiovascular disease (Naruse *et al.*, 2019).

As in previous studies we found that paternal obesity resulted in sex-specific differences in

metabolism and fat accruement with males being more impacted. The underlying mechanisms that lead to

the greater susceptibility of males may be related to sexually dimorphic placental gene expression (Eriksson
et al., 2010). In support of this possibility, paternal environment (diet) influenced placental function in a sex-
specific manner (Binder *et al.*, 2015). Alternatively, different metabolic responses in male and female
offspring may be due to hormonal responses where estrogen has been shown to protect against altered
glucose homeostasis (Gupte, Pownall and Hamilton, 2015; Lainez *et al.*, 2018).

In summary, we provide evidence that paternal obesity induced altered H3K4me3 signatures in
sperm contribute to the inheritance of metabolic disease. Our findings indicate further studies on the
relationship between sperm chromatin, the placental chromatin and gene expression are warranted and
could lead to a better understanding of paternal sperm epigenome transmission of metabolic disease. The
translational validation of these findings will be important in developing intervention strategies focused on
paternal health that could impact the health of future generations (Barratt *et al.*, 2021).

Materials and Methods

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by
the Lead Contact, Sarah Kimmins (sarah.kimmins@mcgill.ca).
Materials availability
This study did not generate new unique reagents.
Data and code availability
The sperm H3K4me3 ChIP-Seq and liver RNA-Seq data generated during this study are available at the
following GEO accession number: GSE178096.
EXPERIMENTAL MODEL AND SUBJECT DETAILS
Animals
All animal procedures were carried out in accordance with the guidelines of the Faculty Animal Care
Committee of McGill University, Montreal. For the wild-type line (WT), C57BL/6NCrl 8-week old males and
6-week old females were purchased from Charles Rivers Laboratory and were allowed one week of
acclimation before breeding. For the KDM1A transgenic line (TG), mice were generated as previously
described (Siklenka et al., 2015), with the same genetic background as the wild-type line. The F_0 transgenic
mice used in this study were from the 11th generation. The earlier generations of mice in this KDM1A TG
had severe developmental abnormalities, pre-implantation loss and early post-natal death (Siklenka *et al.*,
2015) (Siklenka et al., 2015). Over time we have selected against the severe phenotype by breeding the
mice that survive and are normal. Single males were housed with two females to generate the F_0
generation. All animals were given access to water and food *ad libitum* and were maintained on a controlled
light/dark cycle.
METHODS DETAILS

Diet experiments and animal breeding

The low-fat control diet (CON; D12450J) and high-fat diet (HFD; D12492) were obtained from Research Diets, and selected based on the matched amounts of sucrose, vitamin mix and folate. Diets' macronutrients composition are listed in Table S1. Males of the F_0 generation were generated from at least 7 different sires per group. F_0 males were weaned at 3 weeks of age and randomly assigned to either a CON or HFD. The number of animals per group, per sex and per generation, used for all metabolic characterization tests can be found in Table S2. Total body weights were monitored weekly. Cumulative caloric intake was recorded weekly by weighting pellets from the food hopper and calculated as kilocalorie per animal. The diet intervention spanned 10-12 weeks followed by 2 weeks of metabolic testing (at 4 months of age), 1 week of rest and 1-2 weeks of breeding with 7-week old C57BL/6NCrl females. In order to minimize maternal exposure to the intervention diets, females were housed with males overnight and separated during the day, for a maximum of 3 nights per week for 2 weeks, or until a vaginal plug was detected. Litter sizes (number of pups per litter) were recorded, and sex ratios (ratio of male pups over total number of pups) were calculated for all litters generated and can be found in Tables S3 and S4, respectively. The same timeline was used to generate the F_1 and F_2 animals. All females used for breeding and all F_1 and F_2 were fed a regular chow diet (2020X Teklad global soy protein-free extruded rodent diet, Envigo). All animals were sacrificed at 22 weeks (±2 weeks) by carbon dioxide asphyxiation under isoflurane anesthesia.

Metabolic testing

Assessment of metabolic parameters was conducted at 4 months of age within 2 consecutive weeks according to the standard operating procedures of the NIH mouse Metabolic Phenotyping Center (Ayala *et al.*, 2010). For the glucose tolerance test, animals were fasted overnight for 15 hours (± 1 hour) starting at 6:00PM with free access to water. Blood glucose was measured before and 15, 30, 60 and 120 minutes following an intraperitoneal injection of 2 g/kg of a 20% glucose solution (D-glucose, G7021, Sigma Aldrich) with one drop of blood from the tail-tip using a glucometer (Accu-Chek Aviva Nano). For the insulin tolerance test, animals were fasted for 6 hours (1 hour), starting at 9:00AM with free access to water. Blood glucose was measured before and 15, 30, 60 and 120 minutes following an intraperitoneal injection of 1 IU/kg insulin (Insulin solution, I9278, Sigma Aldrich), with one drop of blood from the tail-tip using a glucometer (Accu-Chek Aviva Nano). The area under the curves (AUCs) for the tolerance tests were calculated using the trapezoidal rule (GraphPad Prism version 8). For the baseline blood glucose levels, blood glucose levels were measured after an overnight fasting of 15 hours (± 1 hour) with one drop of blood from the tail-tip using a glucometer (Accu-Chek Aviva Nano).

Tissue collection

At necropsy, mice were dissected to collect adipose tissue (gonadal and mesenteric white adipose depots; gWAT and mWAT, respectively) and a liver lobe (left lateral lobe or *lobus hepatis sinister lateralis* for RNA-sequencing). All tissues were weighed, transferred to a clean tube, snap frozen in liquid nitrogen and stored

612 at -80°C until subsequent downstream experiments. Cauda epididymides were weighed and immediately
613 used for sperm isolation.

Sperm isolation

Spermatozoa were isolated from paired caudal epididymides (Hisano *et al.*, 2013; Lismer, Lambrot, *et al.*,
2021). Cauda epididymides were cut into 5 mL of freshly-prepared Donners medium (25 mM NaHCO$_3$, 20
mg ml^{-1} BSA, 1 mM sodium pyruvate, 0.53% vol/vol sodium DL-lactate in Donners stock) and gently agitated
to allow to swim out for 1 hour at 37°C. The solution was passed through a 40-µm cell strainer (Fisher
Scientific, #22363547) and washed three times with phosphate-buffered saline (PBS). The sperm pellet
was cryopreserved in freezing medium (Irvine Scientific, cat. #90128) and kept in a -80°C freezer until the
chromatin immunoprecipitation experiment.

RNA-Sequencing and library preparation

RNA extraction was performed using the RNeasy Mini Kit (Qiagen, cat. #74104) following the
manufacturer's protocol with slight modifications. In brief, 15-20 mg of liver lobes were cut on dry ice using
a sterile scalpel and Petri dish. Samples were lysed in 350 µL of a denaturing buffer (*Buffer RLT* with beta-
mercaptoethanol) and homogenized with homogenizer pestles. Lysates were centrifuged at maximum
speed for 3 minutes and the supernatants transferred to a clear tube. Ethanol (50%) was added to lysates
to promote selective binding of RNA molecules to the silica-based membrane when applied to the spin
columns. To avoid genomic DNA contamination, an additional DNase digestion was performed. Finally,
membranes of the spin columns were washed twice with 500 µL of *Buffer RPE* and total RNA was eluted
using 30 µL of RNase-free water. Libraries were prepared and sequenced at the *Génome Québec*
Innovation Centre with single-end 50 base-pair reads on the illumina HiSeq 4000 and paired-end 100 base-
pair reads on the illumina NovaSeq 6000 S2 sequencing platforms.

ChIP-Sequencing and library preparation

Chromatin immunoprecipitation was performed as we have previously described (Hisano *et al.*, 2013;
Lismer, Lambrot, *et al.*, 2021). In brief, spermatozoa samples in freezing media were thawed on ice and
washed with 1 mL phosphate-buffered saline. For each sample, two aliquots of 10 µL were used to count
spermatozoa in a hemocytometer under microscope, and 10 million spermatozoa were used per
experiment. Sperm chromatin was decondensed in 1 M dithiothreitol (DTT; Bio Shop, #3483-12-3) and the
reaction quenched with N-ethylmaleimide (NEM). Samples were lysed in lysis buffer (0.3M sucrose, 60mM
KCl, 15mM Tris-HCl pH 7.5, 0.5mM DTT, 5mM McGl2, 0.1mM EGTA, 1% deoxycholate and 0.5% NP40).
An MNase enzyme (15 units; Roche, #10107921001) was added to aliquots containing 2 million
spermatozoa in an MNase buffer (0.3 M sucrose, 85 mM Tris-HCl pH 7.5, 3mM MgCl$_2$ and 2 mM CaCl$_2$),
for exactly 5 minutes at 37°C. The digestion was stopped with 5 mM EDTA. Samples were centrifuged at
maximum speed for 10 minutes, and the supernatants of aliquots from each sample were pooled back
together. Each tube was supplemented with a protease inhibitor to obtain an 1X solution (complete Tablets
EASYpack, Roche, #04693116001). Magnetic beads (DynaBeads, Protein A, Thermo Fisher Scientific,
#10002D) were pre-blocked in a 0.5% Bovine Serum Albumin (BSA, Sigma Aldrich, #BP1600-100) solution

for 4 hours at 4°C and then used to pre-clear the chromatin for 1 hour at 4°C. Pulling down of the pre-
cleared chromatin was performed with the use of magnetic beads that were previously incubated with 5 µg
of antibody (Histone H3 Lysine 4 trimethylation; H3K4me3; Cell Signaling Technology, cat. #9751) for 8
hours at 4°C. Immunoprecipitation of the chromatin with the beads-antibody suspension was performed
overnight at 4°C. Beads bound to the chromatin were subjected to a 3-step wash, one wash with Washing
Buffer A (50 mM Tris-HCl pH 7.5, 10 mM EDTA, 75 mM NaCl) and two washes with Washing Buffer B (50
mM Tris-HCl pH 7.5, 10 mM EDTA, 125 mM NaC). The chromatin was eluted in 250 µL of Elution Buffer
(0.1 M HaHCO3, 0.2% SDS, 5 mM DTT) by incubating the beads twice (2 x 125 µL) shaking at 400 rpm for
10 minutes at 65°C, vortexing vigorously and transferring the chromatin elute in a clean tube. The eluted
chromatin was finally treated with 5 µL of RNase A (Sigma Aldrich, #10109169001) by shaking in a
thermomixer at 400 rpm for 1 hour at 37°C, and then with 5 µL of Proteinase K (Sigma Aldrich, #P2308)
overnight at 55°C. DNA was extracted and purified using the *ChIP DNA Clean and Concentrator kit* (Zymo
Research, #D5201) using the manufacturer's protocol, eluted with 25 µL of the provided elution buffer. Size
selection of the mononucleosomes (147 bp) was performed with the use of Agencourt AMPure XP beads
(Beckman Coulter, #A63880). Libraries were prepared in-house using the Ultra-low Input Library kit
(Qiagen; #180495). Libraries were sequenced with single-end 50 base-pair reads on the illumina HiSeq
4000 sequencing platform (n=5 samples per experimental group).

Pre-processing

Liver RNA-Sequencing data

All samples were processed with the same parameters with the exception of those sequenced on the
NovaSeq platform to adapt for paired-end sequencing and sequencing read length. Reads were trimmed
using *Trim Galore* (version 0.5.0, parameters for HiSeq: --phred33 --length 36 -q 5 --stringency 1 -e 0.1;
parameters for NovaSeq: --paired --retain_unpaired --phred33 --length 36 -q 5 --stringency 1 -e 0.1)
(Krueger, 2015). Trimmed reads were aligned to the *Ensembl* Genome Reference Consortium mouse
reference 38 (GRCm38) primary assembly using *hisat2* (version 2.1.0, parameters: -p 8 --dta) (Kim,
Langmead and Salzberg, 2015). Aligned files with SAM format were converted to binary SAM format (BAM)
and sorted by genomic position using *SAMtools* (version 1.9) (Li *et al.*, 2009). Transcripts were assembled
and gene abundances calculated using *Stringtie* (version 2.1.2, parameters: -p 8 -e -B -A) (Pertea *et al.*,
2015).

Sperm ChIP-Sequencing data

Sequencing reads were trimmed using *Trimmomatic* on single-end mode to remove adapters and filter out
low-quality reads (version 0.36, parameters: 2:30:15 LEADING:30 TRAILING:30) (Bolger, Lohse and
Usadel, 2014). Trimmed reads were aligned to the *Mus Musculus* mm10 genome assembly using *Bowtie2*
(version 2.3.4) (Salzberg, 2013). Unmapped reads were removed using *SAMtools* (version 1.9) (Li *et al.*,
2009), and those with 3 mismatches or more were filtered out using *Perlcode*. BAM coverage files (BigWig)
were generated using *deeptools2 bamCoverage* function (version 3.2.1, parameters: -of bigwig -bs 25 -p
20 --normalizeUsing RPKM -e 160 --ignoreForNormalization chrX) (Ramírez *et al.*, 2016).

Other publicly available ATAC-Sequencing or ChIP-Sequencing datasets

Raw files for 2-cell H3K4me3 and H3K27me3 ChIP-Seq (Liu *et al.*, 2016) (GEO: GSE73952), MII oocyte H3K4me3 ChIP-Seq (Zhang *et al.*, 2016) (GEO: GSE71434), sperm ATAC-Seq (Jung *et al.*, 2017) (GEO: GSE79230), 4-cell and morula ATAC-Seq (Liu *et al.*, 2019) (NCBI SRA: SRP163205),TE H3K4me3 ChIP-Seq (Liu *et al.*, 2016) (GEO: GSE73952), and placenta H3K4me3 ChIP-Seq (Shen *et al.*, 2012) (GEO: GSE29184) were downloaded from the National Centre for Biotechnology Information (NCBI) using the Sequencing Read Archive (SRA) Toolkit. Files were pre-processed as described above for the sperm H3K4me3 ChIP-Sequencing with slight modifications to adapt for datasets with paired-end reads and for different sequencing read lengths.

Other publicly available RNA-Sequencing data

Raw files for 4-cell and morula (Liu *et al.*, 2019) (NCBI SRA: SRP163205), TE (Liu *et al.*, 2016) (GEO: GSE73952), and placenta (Chu *et al.*, 2019) (NCBI SRA: SRP137723) RNA-Seq were downloaded from the National Centre for Biotechnology Information (NCBI) using the Sequencing Read Archive (SRA) Toolkit. Files were pre-processed as described above for the sperm H3K4me3 ChIP-Sequencing with slight modifications to adapt for datasets with paired-end reads and for different sequencing read lengths.

Paternal allele 2-cell embryo ChIP-Sequencing data

Raw files for 2-cell H3K4me3 ChIP-Seq (Zhang *et al.*, 2016) (GEO: GSE71434) were downloaded from the National Centre for Biotechnology Information (NCBI) using the Sequencing Read Archive (SRA) Toolkit. *SNPsplit* (version 0.3.2) was used to build a reference genome with PWK_PhJ SNPs masked (Krueger and Andrews, 2016). Reads were aligned to the generated PWK_PhJ SNPs N-masked reference genome using *Bowtie2* (parameters: -p 10 -t -q -N 1 -L 25 -X 2000 --no-mixed --no-discordant). Aligned files with SAM format were converted to binary SAM format (BAM) and sorted by genomic position using *SAMtools* (version 1.9) (Li *et al.*, 2009). *SNPsplit* (version 0.3.2) was used to assign reads to either the paternal (PWK_PhJ) or the maternal (C57BL/6) genome based on SNPs origin. BAM coverage files (BigWig) were generated using *deeptools2 bamCoverage* function (parameters: -of bigwig -bs 25 -p 20 --normalizeUsing RPKM -e 160 --ignoreForNormalization chrX).

QUANTIFICATION AND STATISTICAL ANALYSIS

Visualization and statistical analyses for metabolic characterization

Visualization of the metabolic characterization data was performed using Jupyter Notebook (version 6.0.1) with Python (version 3.7.4), with the use of the following packages: *seaborn* (version 0.9.0) (Waskom, 2021), *numpy* (version 1.17.2) (Harris *et al.*, 2020), and *panda* (version 0.25.2) (Mckinney, 2010). The *pyplot* and *patches* modules were loaded from the *matplotlib* library (version 3.4.2) (Hunter, 2007). Statistical analyses were conducted using GraphPad Prism 8. For all tests, a p-value less than 0.05 was considered significant. Significance for individual time points on the blood glucose curves for the glucose and insulin tolerance tests, for cumulative energy intake and for growth trajectories during the diet intervention, was tested using multiple t-test with a Holm-Sidak correction. Significance for total body

weight, mesenteric and gonadal white adipose tissue weight, baseline blood glucose and the area under
the curve for the glucose and insulin tolerance tests, was tested using two-way ANOVA followed by Fisher's
LSD. Litter sizes and sex ratios were analyzed by 2-way ANOVA to assess genotype and diet effects and
interaction, followed by Tukey's multiple comparisons test.

Bioinformatics analysis

All bioinformatics analyses were conducted using R version 4.0.2 (R Core Team, 2018).

Liver RNA-Sequencing data

Transcripts with a mean count below 10 were filtered out, conferring a total of 27,907 and 45,992 detected
expressed transcripts in samples sequenced on the illumina HiSeq and NovaSeq platforms, respectively.
The samples tended to cluster by RNA Integrity Number (RIN), which was corrected for in the differential
analysis (Fig. S3B). Differential expression analysis was conducted using *DESeq2* (version 1.28.1) (Love,
Huber and Anders, 2014), by including sample's RIN value and group in the design formula. Independent
hypothesis weighting (IHW, version 1.16.0) was used to correct for multiple testing and prioritization of
hypothesis testing based on covariate (i.e. the means of normalized counts) (Ignatiadis *et al.*, 2016). IHW
calculates weight for each individual p-value and then applies the Benjamini-Hochberg (BH) procedure to
adjust weighted p-values (Benjamini and Hochberg, 1995). Finally, we used the Lancaster method to
perform a gene-level analysis at single transcript resolution (*aggregation* package, version 1.0.1) (Yi *et al.*,
2018). Lancaster applies aggregation of individual transcripts p-values to obtain differentially expressed
genes while capturing changes at the transcript level. Genes with a Lancaster p-value below 0.05 were
considered significant.

For data visualization, transcript counts were normalized using variance stabilizing transformation without
the use of blind dispersion estimation (i.e. with parameter *blind=FALSE*) (Love, Huber and Anders, 2014).
This transformation approach translates data on a $\log_2$ scale, allows correction for library size and removes
the dependence of the variance on the mean (heteroscedasticity). Variance-stabilized transcript counts
were corrected for RIN values using *limma's removeBatchEffect* function (version 3.44.3) (Ritchie *et al.*,
2015). Pearson correlation heatmaps were generated using the *corrplot* package (version 0.88) (Taiyun,
Wei, Simko, 2021), with samples ordered by hierarchical clustering. Principal component analysis was
performed using *DEseq's plotPCA* function, with RIN values and sexes labeled. Heatmaps of differentially
expressed genes were generated with the *Pheatmap* package (version 1.0.12) (Kolde, 2019), with
transcripts ordered by k-means clustering (n kmeans=2) and samples ordered by hierarchical clustering
using complete-linkage clustering based on Euclidean distance. Alluvial plots were generated with *ggplot2*
(version 3.3.3) (Wickham, 2016), and overlap of differentially expressed genes across genotypes,
generations and sexes were determined by the *GeneOverlap* package (version 1.24.0) (Shen, 2014), which
uses a Fisher's exact test to compute p-values.

Visualization, Semantic similarity, and Enrichment Analysis of Gene Ontology (ViSEAGO)

Gene ontology (GO) analysis was performed using the *ViSEAGO* package (version 1.2.0) (Brionne,
Juanchich and Hennequet-Antier, 2019). Gene symbols and *EntrezGene* IDs from the *org.Mm.eg.db*
database were retrieved using the *AnnotationDbi* package. GO annotations were retrieved from
EntrezGene for the *Mus Musculus* species (ID="10090") using the *ViSEAGO EntrezGene2GO* followed by
annotate functions. ViSEAGO uses topGO to perform GO terms enrichment tests on the sets of genes of
interest (differentially expressed genes). We used the Biological Process (BP) ontology category with
Fisher's exact test (classic algorithm), and a p-value below 0.01 was considered significant. Results of
enrichment tests for each set of genes of interest were then merged and hierarchical clustering was
performed based on Wang's semantic similarity distance and *ward.D2* aggregation criterion. Results are
visualized on a heatmap where GO terms are ordered by hierarchical clustering based on their functional
similarity and GO terms enrichment significance is shown as a color gradient (-$\log_{10}$ p-value) in each set of
differentially expressed genes of interest.

Sperm ChIP-Sequencing data

To detect genomic regions enriched with H3K4me3 in sperm, we used *csaw* (version 1.22.1) (Lun and
Smyth, 2016) to scan the genome into windows of 150-bp. Windows with a fold-change enrichment of 4
over bins of 2,000 bp (background) were considered enriched. Enriched regions less than 100 bp apart
were merged for a maximum width of 5,000 bp, conferring a total of 30,745 merged enriched regions.
Counts in enriched regions were normalized using TMM normalization followed by *ComBat's* correction for
batch effects (*sva* package, version 3.36.0) (Leek *et al.*, 2012; Zhang, Parmigiani and Johnson, 2020).
Spearman correlation heatmaps and MA-plots were generated using raw and normalized counts at
enriched regions using *corrplot* (version 0.88) (Taiyun, Wei, Simko, 2021), and *graphics* packages,
respectively.

Principal component analysis was conducted on normalized counts in enriched regions, by comparing WT
HFD vs WT CON (effect of diet in WT), TG HFD vs TG CON (effect of diet in TG), and WT CON vs TG
HFD (effects of genotype and HFD). Based on visual assessment of the separation of samples according
to dietary or genotype groups along Principal Component 1 (PC1; x axis) or 2 (PC2; y axis), the top 5%
regions contributing the PC of interest were selected. Permutational multivariate analysis of variance
(PERMANOVA) was conducted to determine whether variation is attributed to dietary/genotype group,
using the *adonis* function (*vegan* package, version 2.5-7) (Oksanen *et al.*, 2007). Euclidean distances were
used as a metric, 999 permutations were performed, and a $p<0.05$ was considered significant. The
directionality change in enrichment was identified based on the positive (up-regulated regions) and negative
(down-regulated regions) $\log_2$ fold change values of the median of normalized counts using *gtools'*
foldchange2logratio function. Regions with increased and decreased enrichment for each comparison of
interest were visualized using *Pheatmap* (version 1.0.12) (Kolde, 2019). Regions distance relative to

transcription start site (TSS) were annotated and visualized using the package *chipenrich* (version 2.12.0)

(Welch *et al.*, 2014). Gene ontology analysis was performed using *topGO* (version 2.40.0) for genes with

increased or decreased H3K4me3 enrichment at the promoter region for each comparison of interest. We

used the Biological Process (BP) ontology category with Fisher's exact test *weight01Fisher* algorithm

(Alexa, Rahnenführer and Lengauer, 2006), and a p-value less than 0.05 was considered significant.

Genomic regions with deH3K4me3 were annotated using *annotatr* (version 1.14.0) (Cavalcante and Sartor,

2017) including CpG annotations and basic genes genomic features. Upset plots were generated using

UpsetR (version 1.4.0) (Conway, Lex and Gehlenborg, 2017)*,* by ordering each set by frequency and

displaying 12 sets. Z-scores were calculated using *regioneR's overlapPermTest* (version 1.20.1) which

performs a permutation test (n=1,000 permutations) to assess whether a set of regions is significantly

enriched to a specific genomic feature compared to genomic regions from the whole genome (Gel *et al.*,

2016). Genome browser snapshots were generated using *trackplot* (Pohl and Beato, 2014).

To assess linear trends associated with the cumulative exposure of KDM1A overexpression and high-fat

feeding in sperm, we ran *DESeq2* (version 1.28.1) on the top 5% regions contributing to Principal

Component 2 (PC2; n=1,538 regions) associated with sample separation when comparing WT CON and

TG HFD normalized counts. In the design formula, we included sample's batch information, and assigned

a numerical value for each sample based on their group category (WT CON=1, WT HFD=2, TG CON=2,

TG HFD=3). Independent hypothesis weighting (IHW) was used to correct for multiple testing and

prioritization of hypothesis testing based on covariate (i.e. the means of normalized counts) (Ignatiadis *et*

al., 2016). Median of normalized counts were used to depict the increased and decreased trend of

significant regions (adjusted p-value less than 0.2) across groups recoded on a numerical scale as defined

above.

Competing Interest Statement

The authors declare no competing interests.

Acknowledgements

We thank the team at the McGill University and Génome Québec Innovation Centre for performing the

sequencing and Dr. Deborah Sloboda (McMaster University) for advice on metabolic phenotyping methods.

This research was funded by the Canadian Institute of Health Research (CIHR) grants to SK (358654 and

350129).

Author Contributions

SK conceived, designed, funded and guided the study. ASP developed the model and conducted the animal

studies with assistance from CL and RL with associated laboratory analysis. ASP conducted the

bioinformatic analysis with assistance and oversight from VD. ASP and SK assembled and wrote the

manuscript which was edited by VD.

References

Alexa, A., Rahnenführer, J. and Lengauer, T. (2006) 'Improved scoring of functional groups from gene

expression data by decorrelating GO graph structure', *Bioinformatics*, 22(13), pp. 1600–1607. doi:

10.1093/bioinformatics/btl140.

Anway, M. D. *et al.* (2005) 'Epigenetic Transgenerational Actions of Endocrine Disruptors and Male

Fertility', *Science*, 308(5727), pp. 1466 LP – 1469. doi: 10.1126/science.1108190.

Ayala, J. E. *et al.* (2010) 'Standard operating procedures for describing and performing metabolic tests of

glucose homeostasis in mice', *DMM Disease Models and Mechanisms*, 3(9–10), pp. 525–534. doi:

10.1242/dmm.006239.

Barratt, C. L. R. *et al.* (2021) 'A global approach to addressing the policy, research and social challenges

of male reproductive health.', *Human reproduction open*, 2021(1), p. hoab009. doi:

10.1093/hropen/hoab009.

Benjamini, Y. and Hochberg, Y. (1995) 'Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing', *Journal of the Royal Statistical Society. Series B (Methodological)*. [Royal

Statistical Society, Wiley], 57(1), pp. 289–300. Available at: http://www.jstor.org/stable/2346101.

Binder, N. K. *et al.* (2015) 'Paternal obesity in a rodent model affects placental gene expression in a sex-

specific manner', *Reproduction*, 149(5), pp. 435–444. doi: 10.1530/REP-14-0676.

Binder, N. K., Hannan, N. J. and Gardner, D. K. (2012) 'Paternal Diet-Induced Obesity Retards Early

Mouse Embryo Development, Mitochondrial Activity and Pregnancy Health', *PLoS ONE*, 7(12). doi:

10.1371/journal.pone.0052304.

Bolger, A. M., Lohse, M. and Usadel, B. (2014) 'Trimmomatic : a flexible trimmer for Illumina sequence

data', *Bioinformatics*, 30(15), pp. 2114–2120. doi: 10.1093/bioinformatics/btu170.

Braun, J. M., Messerlian, C. and Hauser, R. (2017) 'Fathers Matter: Why It's Time to Consider the Impact

of Paternal Environmental Exposures on Children's Health.', *Current epidemiology reports*, 4(1), pp. 46–

55. doi: 10.1007/s40471-017-0098-8.

Brionne, A., Juanchich, A. and Hennequet-Antier, C. (2019) 'ViSEAGO: A Bioconductor package for

clustering biological functions using Gene Ontology and semantic similarity', *BioData Mining*. BioData

Mining, 12(1), pp. 1–13. doi: 10.1186/s13040-019-0204-1.

Brown, C. L. *et al.* (2015) 'Addressing Childhood Obesity: Opportunities for Prevention.', *Pediatric clinics

of North America*, 62(5), pp. 1241–1261. doi: 10.1016/j.pcl.2015.05.013.

Brykczynska, U. *et al.* (2010) 'Repressive and active histone methylation mark distinct promoters in

human and mouse spermatozoa', *Nature Structural and Molecular Biology*. Nature Publishing Group,

17(6), pp. 679–687. doi: 10.1038/nsmb.1821.

Carone, B. R. *et al.* (2010) 'Paternally induced transgenerational environmental reprogramming of

metabolic gene expression in mammals', *Cell*. Elsevier Inc., 143(7), pp. 1084–1096. doi:

10.1016/j.cell.2010.12.008.

Carone, B. R. *et al.* (2011) 'Paternally-induced transgenerational environmental reprogramming of

metabolic gene expression in mammals Benjamin', 143(7), pp. 1084–1096. doi:

10.1016/j.cell.2010.12.008.Paternally-induced.

de Castro Barbosa, T. *et al.* (2016) 'High-fat diet reprograms the epigenome of rat spermatozoa and

transgenerationally affects metabolism of the offspring', *Molecular Metabolism*. Elsevier GmbH, 5(3), pp.

184–197. doi: 10.1016/j.molmet.2015.12.002.

Cavalcante, R. G. and Sartor, M. A. (2017) 'annotatr: genomic regions in context.', *Bioinformatics (Oxford,*

England), 33(15), pp. 2381–2383. doi: 10.1093/bioinformatics/btx183.

Chen, Q. *et al.* (2016) 'Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic

disorder', *Science*, 351(6271), pp. 397–400. doi: 10.1126/science.aad7977.

Chu, A. *et al.* (2019) 'The Placental Transcriptome in Late Gestational Hypoxia Resulting in Murine

Intrauterine Growth Restriction Parallels Increased Risk of Adult Cardiometabolic Disease.', *Scientific*

reports, 9(1), p. 1243. doi: 10.1038/s41598-018-37627-y.

Conway, J. R., Lex, A. and Gehlenborg, N. (2017) 'UpSetR: an R package for the visualization of

intersecting sets and their properties', *Bioinformatics*, 33(18), pp. 2938–2940. doi:

10.1093/bioinformatics/btx364.

Cropley, J. E. *et al.* (2016) 'Male-lineage transmission of an acquired metabolic phenotype induced by

grand-paternal obesity', *Molecular Metabolism*. Elsevier GmbH, 5(8), pp. 699–708. doi:

10.1016/j.molmet.2016.06.008.

Dalgaard, K. *et al.* (2016) 'Trim28 Haploinsufficiency Triggers Bi-stable Epigenetic Obesity', *Cell*. Elsevier,

164(3), pp. 353–364. doi: 10.1016/j.cell.2015.12.025.

Denomme, Michelle M. *et al.* (2020) 'Advanced paternal age directly impacts mouse embryonic placental

imprinting', *PLoS ONE*, 15(3), pp. 1–13. doi: 10.1371/journal.pone.0229904.

Denomme, Michelle M *et al.* (2020) 'The inherited methylome landscape is directly altered with paternal

aging and associated with offspring neurodevelopmental disorders', *Aging Cell*. John Wiley & Sons, Ltd,

19(8), p. e13178. doi: https://doi.org/10.1111/acel.13178.

Dolinoy, D. C. *et al.* (2006) 'Maternal genistein alters coat color and protects Avy mouse offspring from

obesity by modifying the fetal epigenome', *Environmental Health Perspectives*, 114(4), pp. 567–572. doi:

10.1289/ehp.8700.

Donkin, I. *et al.* (2016) 'Obesity and bariatric surgery drive epigenetic variation of spermatozoa in

humans', *Cell Metabolism*, 23(2), pp. 369–378. doi: 10.1016/j.cmet.2015.11.004.

Donkin, I. and Barrès, R. (2018) 'Sperm epigenetics and influence of environmental factors', *Molecular*

Metabolism, 14, pp. 1–11. doi: https://doi.org/10.1016/j.molmet.2018.02.006.

Eberle, C. *et al.* (2020) 'Paternal metabolic and cardiovascular programming of their offspring: A

systematic scoping review.', *PloS one*, 15(12), p. e0244826. doi: 10.1371/journal.pone.0244826.

Eriksson, J. G. *et al.* (2010) 'Boys live dangerously in the womb.', *American journal of human biology : the*
official journal of the Human Biology Council, 22(3), pp. 330–335. doi: 10.1002/ajhb.20995.

Franzago, M. *et al.* (2019) 'Nutrigenetics , epigenetics and gestational diabetes : consequences in mother
and child', *Epigenetics*. Taylor & Francis, 14(3), pp. 215–235. doi: 10.1080/15592294.2019.1582277.

Fullston, T. *et al.* (2013) 'Paternal obesity initiates metabolic disturbances in two generations of mice with
incomplete penetrance to the F2 generation and alters the transcriptional profile of testis and sperm
microRNA content', *FASEB Journal*, 27(10), pp. 4226–4243. doi: 10.1096/fj.12-224048.

Gabory, A., Attig, L. and Junien, C. (2011) 'Developmental programming and epigenetics', *The American*
Journal of Clinical Nutrition, 94(suppl_6), pp. 1943S-1952S. doi: 10.3945/ajcn.110.000927.

Gel, B. *et al.* (2016) 'regioneR: an R/Bioconductor package for the association analysis of genomic
regions based on permutation tests', *Bioinformatics*, 32(2), pp. 289–291. doi:
10.1093/bioinformatics/btv562.

Gerdes, P. *et al.* (2016) 'Transposable elements in the mammalian embryo: pioneers surviving through
stealth and service.', *Genome biology*, 17, p. 100. doi: 10.1186/s13059-016-0965-5.

Gernand, A. D. *et al.* (2016) 'Micronutrient deficiencies in pregnancy worldwide: health effects and
prevention.', *Nature reviews. Endocrinology*, 12(5), pp. 274–289. doi: 10.1038/nrendo.2016.37.

Grandjean, V. *et al.* (2015) 'RNA-mediated paternal heredity of diet-induced obesity and metabolic
disorders', *Scientific Reports*. Nature Publishing Group, 5(June), pp. 1–9. doi: 10.1038/srep18193.

Gupte, A. A., Pownall, H. J. and Hamilton, D. J. (2015) 'Estrogen : An Emerging Regulator of Insulin
Action and Mitochondrial Function', 2015.

Hammoud, S. S. *et al.* (2009) 'Distinctive chromatin in human sperm packages genes for embryo
development', *Nature*. Nature Publishing Group, 460(7254), pp. 473–478. doi: 10.1038/nature08162.

Harris, C. R. *et al.* (2020) 'Array programming with NumPy', *Nature*, 585(7825), pp. 357–362. doi:
10.1038/s41586-020-2649-2.

Hisano, M. *et al.* (2013) 'Genome-wide chromatin analysis in mature mouse and human spermatozoa',
Nature Protocols, 8(12), pp. 2449–2470. doi: 10.1038/nprot.2013.145.

Hunter, J. D. (2007) 'Matplotlib: A 2D Graphics Environment', *Computing in Science & Engineering*, 9(3),
pp. 90–95. doi: 10.1109/MCSE.2007.55.

Huypens, P. *et al.* (2016) 'Epigenetic germline inheritance of diet-induced obesity and insulin resistance',
Nature Genetics, 48(5), pp. 497–499. doi: 10.1038/ng.3527.

Ignatiadis, N. *et al.* (2016) 'Data-driven hypothesis weighting increases detection power in genome-scale
multiple testing', *Nature Methods*, 13(7), pp. 577–580. doi: 10.1038/nmeth.3885.

Isganaitis, E., Suehiro, H. and Cardona, C. (2017) 'Who's your daddy?: paternal inheritance of metabolic
disease risk.', *Current opinion in endocrinology, diabetes, and obesity*. England, 24(1), pp. 47–55. doi:
10.1097/MED.0000000000000307.

Jukam, D., Shariati, S. A. M. and Skotheim, J. M. (2017) 'Zygotic Genome Activation in Vertebrates.',
Developmental cell, 42(4), pp. 316–332. doi: 10.1016/j.devcel.2017.07.026.

Jung, Y. H. *et al.* (2017) 'Chromatin States in Mouse Sperm Correlate with Embryonic and Adult

Regulatory Landscapes', *Cell Reports*. ElsevierCompany., 18(6), pp. 1366–1382. doi:

10.1016/j.celrep.2017.01.034.

Kaati, G., Bygren, L. O. and Edvinsson, S. (2002) 'Cardiovascular and diabetes mortality determined by

nutrition during parents' and grandparents' slow growth period', *European Journal of Human Genetics*,

10(11), pp. 682–688. doi: 10.1038/sj.ejhg.5200859.

Karatela, R. A. and Sainani, G. S. (2009) 'Plasma homocysteine in obese, overweight and normal weight

hypertensives and normotensives.', *Indian heart journal*. India, 61(2), pp. 156–159.

Kerr, B. *et al.* (2018) 'Foetoplacental epigenetic changes associated with maternal metabolic

dysfunction.', *Placenta*. Netherlands, 69, pp. 146–152. doi: 10.1016/j.placenta.2018.04.006.

Kim, D., Langmead, B. and Salzberg, S. L. (2015) 'HISAT: a fast spliced aligner with low memory

requirements', *Nature Methods*, 12(4), pp. 357–360. doi: 10.1038/nmeth.3317.

Kim, S. *et al.* (2004) 'Hepatic gene expression profiles in a long-term high-fat diet-induced obesity mouse

model.', *Gene*. Netherlands, 340(1), pp. 99–109. doi: 10.1016/j.gene.2004.06.015.

Kolde, R. (2019) 'pheatmap: Pretty Heatmaps. R package'. Available at: https://cran.r-

project.org/package=pheatmap.

Krueger, F. (2015) 'Trim galore', *A wrapper tool around Cutadapt and FastQC to consistently apply

quality and adapter trimming to FastQ files*, 516, p. 517.

Krueger, F. and Andrews, S. R. (2016) 'SNPsplit: Allele-specific splitting of alignments between genomes

with known SNP genotypes [version 1; referees: 3 approved]', *F1000Research*, 5, pp. 1–15. doi:

10.12688/F1000RESEARCH.9037.1.

Lainez, N. M. *et al.* (2018) 'Diet-induced obesity elicits macrophage infiltration and reduction in spine

density in the hypothalami of male but not female mice', *Frontiers in Immunology*, 9(SEP), pp. 1–16. doi:

10.3389/fimmu.2018.01992.

Lambrot, R. *et al.* (2013) 'Low paternal dietary folate alters the mouse sperm epigenome and is

associated with negative pregnancy outcomes', *Nature Communications*, 4. doi: 10.1038/ncomms3889.

Lambrot, R. *et al.* (2019) 'The genomic distribution of histone H3K4me2 in spermatogonia is highly

conserved in sperm†.', *Biology of reproduction*. United States, 100(6), pp. 1661–1672. doi:

10.1093/biolre/ioz055.

Lambrot, R. (2021) 'Whole genome sequencing of H3 lysine 4 tri-methylation and DNA methylation in

human sperm reveals regions of overlap and exclusion linked to fertility, development and epigenetic

inheritance', *Cell Reports*, CELL-REPOR.

Leavey, K. *et al.* (2018) 'Epigenetic regulation of placental gene expression in transcriptional subtypes of

preeclampsia', *Clinical Epigenetics*, 10(1), p. 28. doi: 10.1186/s13148-018-0463-6.

Leek, J. T. *et al.* (2012) 'The sva package for removing batch effects and other unwanted variation in

high-throughput experiments.', *Bioinformatics (Oxford, England)*, 28(6), pp. 882–883. doi:

10.1093/bioinformatics/bts034.

Lesch, B. J. *et al.* (2019) 'Intergenerational epigenetic inheritance of cancer susceptibility in mammals',

eLife, 8, pp. 1–29. doi: 10.7554/elife.39380.

Li, H. *et al.* (2009) 'The Sequence Alignment/Map format and SAMtools.', *Bioinformatics (Oxford,*

England), 25(16), pp. 2078–2079. doi: 10.1093/bioinformatics/btp352.

Lismer, A. *et al.* (2020) 'Sperm histone H3 lysine 4 trimethylation is altered in a genetic mouse model of

transgenerational epigenetic inheritance', *Nucleic Acids Research*. doi: 10.1093/nar/gkaa712.

Lismer, A., Lambrot, R., *et al.* (2021) 'ChIP-seq protocol for sperm cells and embryos to assess

environmental impacts and epigenetic inheritance', *STAR Protocols*, 2(2), p. 100602. doi:

https://doi.org/10.1016/j.xpro.2021.100602.

Lismer, A., Dumeaux, V., *et al.* (2021) 'Histone H3 lysine 4 trimethylation in sperm is transmitted to the

embryo and associated with diet- induced phenotypes in the offspring Article Histone H3 lysine 4

trimethylation in sperm is transmitted to the embryo and associated with diet-induce', *Developmental Cell*.

Elsevier Inc., pp. 1–16. doi: 10.1016/j.devcel.2021.01.014.

Liu, L. *et al.* (2019) 'An integrated chromatin accessibility and transcriptome landscape of human pre-

implantation embryos', *Nature communications*. Nature Publishing Group UK, 10(1), p. 364. doi:

10.1038/s41467-018-08244-0.

Liu, X. *et al.* (2016) 'Distinct features of H3K4me3 and H3K27me3 chromatin domains in pre-implantation

embryos', *Nature*. Nature Publishing Group, 537(7621), pp. 558–562. doi: 10.1038/nature19362.

Love, M. I., Huber, W. and Anders, S. (2014) 'Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2', *Genome Biology*, 15(12), pp. 1–21. doi: 10.1186/s13059-014-0550-8.

Lumey, L. H. and Poppel, F. W. A. Van (2013) 'The Dutch Famine of 1944-45 as a Human Laboratory :

Changes in the Early Life Environment and Adult Health', in Lumey, L. H. and Vaiserman, A. (ed.) *Early*

Life Nutrition and Adult Health and Development. Nova Science Publishers, Inc., pp. 59–76.

Lun, A. T. L. and Smyth, G. K. (2016) 'csaw: a Bioconductor package for differential binding analysis of

ChIP-seq data using sliding windows.', *Nucleic acids research*, 44(5), p. e45. doi: 10.1093/nar/gkv1191.

Martínez, D. *et al.* (2014) 'In utero undernutrition in male mice programs liver lipid metabolism in the

second-generation offspring involving altered Lxra DNA methylation.', *Cell metabolism*. United States,

19(6), pp. 941–951. doi: 10.1016/j.cmet.2014.03.026.

Mckinney, W. (2010) 'Data Structures for Statistical Computing in Python', in *Proceedings of the 9th*

Python in Science Conference, pp. 56–61.

Meister, S. *et al.* (2021) 'Epigenetic modification via H3K4me3 and H3K9ac in human placenta is reduced

in preeclampsia', *Journal of Reproductive Immunology*, 145, p. 103287. doi:

https://doi.org/10.1016/j.jri.2021.103287.

Miska, E. A. and Ferguson-smith, A. C. (2016) 'Transgenerational inheritance: Models and mechanisms

of non – DNA sequence – based inheritance', 354(6308), pp. 778–782.

Moore, T. (2001) 'Genetic conflict, genomic imprinting and establishment of the epigenotype in relation to

growth.', *Reproduction (Cambridge, England)*. England, 122(2), pp. 185–193. doi:

10.1530/rep.0.1220185.

Morgan, H. D. *et al.* (1999) 'Epigenetic inheritance at the agouti locus in the mouse', *Nature Genetics*, 23(3), pp. 314–318. doi: 10.1038/15490.

Mouse Genome Informatics (no date).

Naruse, K. *et al.* (2019) 'Preliminary evidence of a paternal-maternal genetic conflict on the placenta: Link between imprinting disorder and multi-generational hypertensive disorders', *Placenta*, 84, pp. 69–73. doi: https://doi.org/10.1016/j.placenta.2019.02.009.

Nelissen, E. C. M. *et al.* (2011) 'Epigenetics and the placenta', *Human Reproduction Update*, 17(3), pp. 397–417. doi: 10.1093/humupd/dmq052.

Ng, S.-F. *et al.* (2010) 'Chronic high-fat diet in fathers programs β-cell dysfunction in female rat offspring', *Nature*, 467(7318), pp. 963–966. doi: 10.1038/nature09491.

Oksanen, J. *et al.* (2007) 'The vegan package', *Community ecology package*, 10(631–637), p. 719.

Padmanabhan, N. *et al.* (2013) 'Mutation in Folate Metabolism Causes Epigenetic Instability and Transgenerational Effects on Development', *Cell*. Elsevier Inc., 155(1), pp. 81–93. doi: 10.1016/j.cell.2013.09.002.

Palmer, N. O. *et al.* (2011) 'SIRT6 in mouse spermatogenesis is modulated by diet-induced obesity.', *Reproduction, fertility, and development*. Australia, 23(7), pp. 929–939. doi: 10.1071/RD10326.

Pembrey, M. E. *et al.* (2006) 'Sex-specific, male-line transgenerational responses in humans', *European Journal of Human Genetics*, 14(2), pp. 159–166. doi: 10.1038/sj.ejhg.5201538.

Perez-Garcia, V. *et al.* (2018) 'Placentation defects are highly prevalent in embryonic lethal mouse mutants', *Nature*, 555(7697), pp. 463–468. doi: 10.1038/nature26002.

Pertea, M. *et al.* (2015) 'StringTie enables improved reconstruction of a transcriptome from RNA-seq reads', *Nature Biotechnology*, 33(3), pp. 290–295. doi: 10.1038/nbt.3122.

Pilsner, J. R. *et al.* (2021) 'Aging-induced changes in sperm DNA methylation are modified by low dose of perinatal flame retardants', *Epigenomics*. Future Medicine, 13(4), pp. 285–297. doi: 10.2217/epi-2020-0404.

Pohl, A. and Beato, M. (2014) 'bwtool: a tool for bigWig files', *Bioinformatics*, 30(11), pp. 1618–1619. doi: 10.1093/bioinformatics/btu056.

Radford, E. J. *et al.* (2014) 'In utero undernourishment perturbs the adult sperm methylome and intergenerational metabolism', *Science*, 345(6198), p. 1255903. doi: 10.1126/science.1255903.

Rakyan, V. K. *et al.* (2003) 'Transgenerational inheritance of epigenetic states at the murine AxinFu allele occurs after maternal and paternal transmission', *Proceedings of the National Academy of Sciences of the United States of America*, 100(5), pp. 2538–2543. doi: 10.1073/pnas.0436776100.

Ramírez, F. *et al.* (2016) 'deepTools2: a next generation web server for deep-sequencing data analysis.', *Nucleic acids research*, 44(W1), pp. W160-5. doi: 10.1093/nar/gkw257.

Ritchie, M. E. *et al.* (2015) 'limma powers differential expression analyses for RNA-sequencing and microarray studies.', *Nucleic acids research*, 43(7), p. e47. doi: 10.1093/nar/gkv007.

Rosenfeld, C. S. (2015) 'Sex-Specific Placental Responses in Fetal Development', *Endocrinology*,

156(10), pp. 3422–3434. doi: 10.1210/en.2015-1227.

Salas-Huetos, A. *et al.* (2021) 'Sperm DNA methylation changes after short-term nut supplementation in

healthy men consuming a Western-style diet.', *Andrology*. England, 9(1), pp. 260–268. doi:

10.1111/andr.12911.

Sales, V. M., Ferguson-Smith, A. C. and Patti, M. E. (2017) 'Epigenetic Mechanisms of Transmission of

Metabolic Disease across Generations', *Cell Metabolism*. Elsevier Inc., 25(3), pp. 559–571. doi:

10.1016/j.cmet.2017.02.016.

Salzberg, B. L. and S. L. (2013) 'Fast gapped-read alignment with Bowtie 2', *Nature Methods*, 9(4), pp.

357–359. doi: 10.1038/nmeth.1923.Fast.

Sánchez-Margalet, V. *et al.* (2002) 'Elevated plasma total homocysteine levels in hyperinsulinemic obese

subjects', *The Journal of Nutritional Biochemistry*, 13(2), pp. 75–79. doi: https://doi.org/10.1016/S0955-

2863(01)00197-8.

Sharma, U. *et al.* (2016) 'Biogenesis and function of tRNA fragments during sperm maturation and

fertilization in mammals', 351(6271), pp. 391–396. doi: 10.1126/science.aad6780.Biogenesis.

Shen, L. (2014) 'GeneOverlap : An R package to test and visualize gene overlaps', pp. 1–10.

Shen, Y. *et al.* (2012) 'A map of the cis-regulatory sequences in the mouse genome.', *Nature*, 488(7409),

pp. 116–120. doi: 10.1038/nature11243.

Siklenka, K. *et al.* (2015) 'Disruption of histone methylation in developing sperm impairs offspring health

transgenerationally', *Science*, 350(6261). doi: 10.1126/science.aab2006.

Soubry, A. *et al.* (2016) 'Obesity-related DNA methylation at imprinted genes in human sperm: Results

from the TIEGER study', *Clinical Epigenetics*, 8(1), p. 51. doi: 10.1186/s13148-016-0217-2.

Stringer, J. M. *et al.* (2018) 'Reduced PRC2 function alters male germline epigenetic programming and

paternal inheritance', *BMC Biology*. BMC Biology, 16(1), pp. 1–20. doi: 10.1186/s12915-018-0569-5.

Sun, N. *et al.* (2020) 'H3K4me3-Mediated Upregulation of LncRNA-HEIPP in Preeclampsia Placenta

Affects Invasion of Trophoblast Cells', 11(December), pp. 1–11. doi: 10.3389/fgene.2020.559478.

Taiyun, Wei, Simko, V. (2021) 'R package "corrplot": Visualization of a Correlation Matrix.' Available at:

https://github.com/taiyun/corrplot.

Team, R. C. (2018) 'R: A language and environment for statistical computing.' Vienna, Austria: R

Foundation for Statistical Computing.

Voight, B. F. *et al.* (2010) 'Twelve type 2 diabetes susceptibility loci identified through large-scale

association analysis', *Nature Genetics*, 42(7), pp. 579–589. doi: 10.1038/ng.609.

Wang, X. *et al.* (2013) 'Paternally expressed genes predominate in the placenta', *Proceedings of the

National Academy of Sciences*, 110(26), pp. 10705–10710. doi: 10.1073/pnas.1308998110.

Waskom, M. L. (2021) 'seaborn : statistical data visualization Statement of need', *The Journal of Open

Source Software*, 6, pp. 1–4. doi: 10.21105/joss.03021.

Wei, Y. *et al.* (2014) 'Paternally induced transgenerational inheritance of susceptibility to diabetes in

mammals', *Proceedings of the National Academy of Sciences*, 111(5), pp. 1873–1878. doi:

10.1073/pnas.1321195111.

Welch, R. P. *et al.* (2014) 'ChIP-Enrich: gene set enrichment testing for ChIP-seq data', *Nucleic Acids*

Research, 42(13), pp. e105–e105. doi: 10.1093/nar/gku463.

Wickham, H. (2016) *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-Verlag. Available

at: https://ggplot2.tidyverse.org.

Wu, H. *et al.* (2017) 'Preconception urinary phthalate concentrations and sperm DNA methylation profiles

among men undergoing IVF treatment: a cross-sectional study.', *Human reproduction (Oxford, England)*,

32(11), pp. 2159–2169. doi: 10.1093/humrep/dex283.

Wu, J. *et al.* (2016) 'The landscape of accessible chromatin in mammalian preimplantation embryos',

Nature. Nature Publishing Group, 534(7609), pp. 652–657. doi: 10.1038/nature18606.

Yi, L. *et al.* (2018) 'Gene-level differential analysis at transcript-level resolution', *Genome Biology*.

Genome Biology, 19(1), pp. 1–11. doi: 10.1186/s13059-018-1419-z.

Yoshida, K. *et al.* (2020) 'ATF7-Dependent Epigenetic Changes Are Required for the Intergenerational

Effect of a Paternal Low-Protein Diet', *Molecular Cell*. Elsevier Inc., 78(3), pp. 445-458.e6. doi:

10.1016/j.molcel.2020.02.028.

Youngson, N. *et al.* (2016) 'Obesity-induced sperm DNA methylation changes at satellite repeats are

reprogrammed in rat offspring YR - 2016/12/1', *Asian Journal of Andrology*, (6 UL-

https://www.ajandrology.com/article.asp?issn=1008-

682X;year=2016;volume=18;issue=6;spage=930;epage=936;aulast=Youngson;t=5), pp. 930 OP-936

VO–18. doi: 10.4103/1008-682X.163190.

Zhang, B. *et al.* (2016) 'Allelic reprogramming of the histone modification H3K4me3 in early mammalian

development', *Nature*. Nature Publishing Group, 537(7621), pp. 553–557. doi: 10.1038/nature19361.

Zhang, Y., Parmigiani, G. and Johnson, W. E. (2020) 'ComBat-seq: batch effect adjustment for RNA-seq

count data', *NAR Genomics and Bioinformatics*, 2(3). doi: 10.1093/nargab/lqaa078.

Zhang, Yunfang *et al.* (2018) 'Dnmt2 mediates intergenerational transmission of paternally acquired

metabolic disorders through sperm small non-coding RNAs', *Nature Cell Biology*. Springer US, 20(May).

doi: 10.1038/s41556-018-0087-2.

1114

A

C57BL/6NCrl WT

C57BL/6NCrl
KDM1A⁺/⁻ TG

B

3 weeks — 15 weeks — 17 weeks — 20 weeks

Weaning — Metabolic testing — Breeding — Necropsy

Adiposity (mWAT, gWAT)

RNA-seq

ChIP-seq

C

— WT CON   ···· WT HFD   — TG CON   ···· TG HFD

F₀♂   F₁♂   F₂♂   F₁♀   F₂♀

D Glucose tolerance test

E Insulin tolerance test

Figure 1: Paternal obesity induces transgenerational metabolic phenotypes in a sex-specific manner that are enhanced in KDM1A descendants A) Experimental mouse model depicting breeding scheme and generations studied. Male C57BL6NCrl (WT) and KDM1A$^{+/-}$ transgenics (TG, C57BL6NCrl) were fed either a control diet (CON) or high-fat diet (HFD) from weaning for 10-12 weeks, then mated to 8-week-old C57BL6NCrl females fed a regular chow diet (CD). Animals studied per experimental group: F_0 (n=15-25 males), F_1 (n=28-49 per sex) and F_2 (n=8-21 per sex) B) Experimental timeline for metabolic testing and downstream experiments performed for each generation (F_{0-2}). Metabolic profiles were measured after the diet intervention at 15 weeks of age and included: baseline blood glucose, and intraperitoneal glucose and insulin tolerance tests (ipGTT and ipITT, respectively). Visceral adipose depots were weighed (mWAT: mesenteric white adipose tissue and gWAT: gonadal white adipose tissue) and the left lateral lobe of the liver used for RNA-sequencing (RNA-seq). Sperm from cauda epididymides were used for chromatin immunoprecipitation followed by sequencing (ChIP-seq), targeting histone H3 lysine 4 tri-methylation (H3K4me3). C) Age-matched male mice fed either a control diet (left) or a high-fat diet (right) for 12 weeks. D) Glucose tolerance test. Blood glucose levels before and after (shaded in grey) an intraperitoneal glucose injection, after overnight fasting (15 ±1 hour) at 4 months of age in F_0 males (i), F_1 males (ii), F_2 males (iii), F_1 females (iv) and F_2 females (v). E) Insulin tolerance test. Blood glucose levels before and after (shaded in grey) an intraperitoneal insulin injection, after a 6-hour (±1 hour) fasting at 4 months of age in F_0 males (i), F_1 males (ii), F_2 males (iii), F_1 females (iv) and F_2 females (v). Results are shown as mean ± SEM. Statistical analyses were performed using multiple t-test with Holm-Sidak correction. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001 (in blue; WT CON vs WT HFD, in green; TG CON vs TG HFD) and [#]P<0.05, [##]P<0.01 (WT HFD vs TG HFD).

Figure 2: Paternal obesity is associated with altered gene expression in the livers of the F_0-F_2
A-J) Heatmaps of normalized expression values scaled by row (z-score) for transcripts that code for differentially expressed hepatic genes (Lancaster p-value<0.05) for each comparison assessed across sex and generation. Individual transcripts (rows) are ordered by k-means clustering and samples (columns) are arranged by hierarchical clustering, using complete-linkage clustering based on Euclidean distance. F_0 WT CON vs WT HFD males (A), F_0 TG CON vs TG HFD males (B), F_0 WT HFD vs TG HFD males (C), F_1 WT CON vs WT HFD males (D), F_1 TG CON vs TG HFD males (E), F_1 WT HFD vs TG HFD males (F), F_1 WT CON vs WT HFD females (G), F_1 TG CON vs TG HFD females (H), F_1 WT HFD vs TG HFD females (I), and F_2 WT HFD vs TG HFD males (J). i-x) Alluvial plots depicting frequency distributions of significant (colored boxes) and non-significant (grey boxes) genes for each comparison and their overlap across genotype (i-iii), across F_0 and F_1 males (iv-vi), across F_1 males and females (vii-ix) and across F_1 and F_2 males (x). Significance of overlap between differentially expressed genes lists was calculated by Fisher's exact test. P-values are included for each comparison above the respective alluvial plot.

Figure 3: Obesity-induced hepatic transcriptome disturbances show functional similarities across genotype, sex and generation A-C) Heatmaps of significant gene ontology (GO) terms clustered by functional similarity, comparing enriched biological functions for each comparison of interest across genotype (A), sex (B) and generation (C). Columns represent enriched GO terms which are ordered by hierarchical clustering based on Wang's semantic similarity distance and *ward.D2* aggregation criterion. Each row represents a comparison of interest for which enriched GO terms were annotated based on the list of significant genes. The color gradient depicts the GO term enrichment significance (-$\log_{10}$ p-value). Interactive versions of these figures can be found in Supplemental files 1-3 and the complete lists of significantly enriched GO terms can be found in Tables S5-7.

Figure 4: Genomic location, directionality change and functions of regions with altered H3K4me3 enrichment by obesity A) Histogram showing frequency distributions of read abundances in 150 bp windows throughout the genome. Windows with an abundance below $\log_2(4)$ fold over background bins of 2,000 bp were filtered out as indicated by the vertical red line. Enriched regions less than 100 bp apart were merged for a maximum width of 5,000 bp, conferring a total of 30,745 merged enriched regions. Reads were counted in merged enriched regions and normalized counts were used for downstream analyses. (see Material and Methods) B-C) Principal component analysis on normalized counts at merged enriched regions comparing WT CON vs WT HFD (B) and TG CON vs TG HFD (C). The top 5% regions contributing to separation of samples along Principal Component 1 (in B; PC1; x axis) or PC2 (in C; y axis) were selected. The PERMANOVA p-values indicating significance associated with dietary treatment are included under each PCA plot. D) Heatmaps of log2 normalized counts of deH3K4me3 regions in sperm with increased enrichment in WT HFD (i; n=1,323), decreased enrichment in WT HFD (ii; n=215), increased enrichment in TG HFD (iii; n=1,067) and decreased enrichment in TG HFD (iv; n=471) in each group. Samples (columns) and regions (rows) are arranged by hierarchical clustering using complete-linkage clustering based on Euclidean distance. Colored boxes indicate sample groups (light blue=WT CON, dark blue=WT HFD, light green=TG CON, dark green=TG HFD). E-G). Venn diagrams showing the overlap of deH3K4me3 in sperm of WT HFD (blue) and in TG HFD (green), for all detected regions (E), those gaining H3K4me3 (F) and those losing H3K4me3 (G). H) Barplots showing the distribution of altered regions based on the distance from the TSS of the nearest gene, for regions with increased enrichment in WT HFD (i; n=1,323), decreased enrichment in WT HFD (ii; n=215), increased enrichment in TG HFD (iii; n=1,067), and decreased enrichment in TG HFD (iv; n=471). The color gradient represents the distance of the regions to TSS in kilobase. I) Gene ontology analysis of diet-induced deH3K4me3 regions at promoters with increased enrichment in WT HFD (i; n=381), decreased enrichment in WT HFD (ii; n=34), increased enrichment in TG HFD (iii; n=230) and decreased enrichment in TG HFD (iv; n=150). Barplots show 8 selected significant GO terms with their respective $-\log_2$(p-value). Tables S9-12 include the complete lists of significantly enriched GO terms.

Pepin_Fig5

Figure 5: Additive effects of KDM1A overexpression and diet-induced obesity in the sperm epigenome at the level of H3K4me3

A) Principal component analysis on normalized counts at merged enriched regions comparing WT CON vs TG HFD. The top 5% regions contributing to separation of samples along Principal Component 2 (PC2; y axis) were selected. The PERMANOVA p-value under the plot indicates significance. B-C) Profile plots of RPKM H3K4me3 counts +/- 1 kilobase around the center of regions with increased H3K4me3 (B) and +/- 2.5 kilobase around the center of regions with decreased H3K4me3 enrichment in TG HFD (C). D-E) Line plots showing the median of normalized sperm H3K4me3 counts for each experimental group at regions showing a significant trend (n=264, adjusted p-value<0.2) with a linear increase in H3K4me3 enrichment (D; n=9) or a linear decrease in H3K4me3 enrichment (E; n=255) from WT CON, WT HFD, TG CON to TG HFD groups. F) Gene ontology analysis on the regions associated with a significant linear trend at promoters (n=104). Barplots show 8 selected significant GO terms with their respective $-\log_2$(p-value). Table S13 includes the complete list of significantly enriched GO terms.

Figure 6: Sperm H3K4me3 regions sensitive to obesity occur at genes with an open chromatin state and expressed in the pre-implantation embryo A) Heatmaps of RPKM counts signal +/- 10 kilobase around the center of regions enriched with H3K4me3 in sperm (i; n=30,745) and regions with obesity-induced deH3K4me3 in sperm (ii; n=2,836) for H3K4me3 enrichment levels in sperm (this study), 2-cell embryo (Liu *et al.*, 2016), 2-cell embryo on the paternal allele and MII oocyte (Zhang *et al.*, 2016), and for chromatin accessibility signal in sperm (Jung *et al.*, 2017), 4-cell embryo and morula embryo (Liu *et al.*, 2019). B) Scatterplots showing H3K4me3 enrichment in sperm (x axis; log2 counts + 10), chromatin accessibility signal (y axis; log2 counts + 10; (Jung *et al.*, 2017)) and gene expression levels (color gradient; log2 FPKM + 10; (Liu *et al.*, 2019)) in 4-cell (i,ii,v,vi) or in morula (iii,iv,vii,viii) embryos, at either all genes with promoters enriched with H3K4me3 in sperm (i-iv) or at diet-sensitive genes (v-viii). The top row of scatterplots includes lowly-expressed genes (bottom 50%) in 4-cell (i and v) or morula (iii or vii) embryos. The bottom row of scatterplots includes highly-expressed genes (top 50%) in 4-cell (ii and iv) or morula (vi and viii) embryos. Pearson's correlation coefficients and their associated p-values are indicated above each scatterplot, comparing H3K4me3 enrichment in sperm versus H3K4me3 enrichment in 4-cell or morula embryos. C) Gene ontology analysis of genes expressed in the 4-cell (i) or the morula (ii) embryos, overlapping with diet-sensitive promoters in sperm. Barplots show 8 selected significant GO terms with their respective -$\log_2$(p-value). Tables S14-15 include the complete lists of significantly enriched GO terms.

Figure 7: Obesity-induced deH3K4me3 regions overlap with genes marked by H3K4me3 and expressed in the trophectoderm and placenta A) Heatmaps of RPKM counts signal +/- 5 kilobase around the center of regions enriched with H3K4me3 in sperm (i; n=30,745) and at regions with diet-induced deH3K4me3 in sperm (n=2,836) for H3K4me3 enrichment levels in sperm (this study), trophectoderm (TE) (Liu *et al.*, 2016) and placenta (Shen *et al.*, 2012). B) Gene ontology analysis of regions enriched with H3K4me3 in sperm, TE and placenta (top 75% from A i) (i), regions enriched with H3K4me3 in sperm only (bottom 25% from A i) (ii), diet-sensitive regions enriched with H3K4me3 in sperm, TE and placenta (top 75% from A ii) (iii), and diet-sensitive regions enriched with H3K4me3 in sperm only (bottom 25% from A ii) (iv). Barplots show 8 selected significant GO terms with their respective -$\log_2$(p-value). Tables S16-19 include the complete lists of significantly enriched GO terms. C) Scatterplots showing H3K4me3 enrichment at promoters in sperm (x axis; log2 counts + 10), H3K4me3 enrichment (y axis; log2 counts + 10) and gene expression levels (color gradient; log2 FPKM + 10) in the trophectoderm (i,ii,v,vi; (Liu *et al.*, 2016)) or in the placenta (iii,iv,vii,viii; (Shen *et al.*, 2012; Chu *et al.*, 2019)), at either all genes with promoters enriched with H3K4me3 in sperm (i-iv) or at diet-sensitive genes (v-viii). The top row of scatterplots includes lowly-expressed genes (bottom 50%) in trophectoderm (i and v) or placenta (iii or vii). The bottom row includes highly-expressed genes (top 50%) in trophectoderm (ii and iv) or placenta (vi and viii). Pearson's correlation coefficients and associated p-values are indicated above each scatterplot, comparing H3K4me3 enrichment in sperm versus H3K4me3 enrichment in the trophectoderm or placenta. D) Gene ontology analysis of genes expressed in the trophectoderm (i) or the placenta (ii), overlapping with diet-sensitive promoters in sperm. Barplots show 8 selected significant GO terms with their respective -$\log_2$(p-value). Tables S20-21 include the complete lists of significantly enriched GO terms.