SHORT TITLE

Multiplex knockout using a single gRNA

ARTICLE TITLE

Multiplex knockout of trichome-regulating MYB duplicates in hybrid poplar using a single gRNA

ONE SENTENCE SUMMARY

Targeting conserved sequences with a single gRNA allowed efficient mutagenesis of a multigene family

and the recovery of trichomeless and triterpene-free poplar mutants.

AUTHORS

W. Patrick Bewg[1,2,3], Scott A. Harding[1,2,3], Nancy L. Engle[4], Brajesh N. Vaidya[5,a], Ran Zhou[1,2,3], Jacob

Reeves[6,b], Thomas W. Horn[6], Nirmal Joshee[5], Jerry W. Jenkins[7,8], Shengqiang Shu[8], Kerrie W. Barry[8], Yuko

Yoshinaga[8], Jane Grimwood[7,8], Robert J. Schmitz[2], Jeremy Schmutz[7,8], Timothy J. Tschaplinski[4], and

Chung-Jui Tsai[1,2,3*]

[1] *School of Forestry and Natural Resources, University of Georgia, Athens, Georgia, 30602, USA*

[2] *Department of Genetics, University of Georgia, Athens, Georgia, 30602, USA*

[3] *Department of Plant Biology, University of Georgia, Athens, Georgia, 30602, USA*

[4] *Oak Ridge National Laboratory, Oak Ridge, Tennessee, 37830, USA*

[5] *Fort Valley State University, Fort Valley, Georgia, 31030, USA*

[6] *Department of Computer Science, University of Georgia, Athens, Georgia, 30602, USA*

[7] *HudsonAlpha Institute for Biotechnology, Huntsville, Alabama, 35806, USA*

[8] *U.S. Department of Energy Joint Genome Institute, Berkeley, California, 94720, USA*

*Senior author and author for correspondence: cjtsai@uga.edu (C.-J.T.)

[a] *Present address: GreenVenus, LLC, Wimauma, Florida 33598, USA*

[b] *Present address: SoundHound Inc., Boulder, Colorado 80302, USA*

AUTHOR CONTRIBUTIONS

C.-J.T. and W.P.B. conceived the study and designed the experiments; W.P.B. performed all experiments and analyzed data; S.A.H. provided guidance on physiological and metabolic analyses; J.R., T.W.H, and R.Z. provided bioinformatic support; B.N.V. and N.J. performed SEM analysis; J.J., K.B., R.J.S., Y.Y., S.S., J.G. and J.S. contributed the *Populus tremula* x *P. alba* INRA 717-1B4 draft genome; N.L.E. and T.J.T. performed wax compositional analysis; W.P.B. and C.-J.T. wrote the manuscript, with input from S.A.H.

FUNDING

The work was funded in part by The Center for Bioenergy Innovation, a U.S. Department of Energy Research Center supported by the Office of Biological and Environmental Research in the DOE Office of Science, the Division of Integrative Organismal Systems (grant nos. IOS-1546867) of the National Science Foundation, the Community Science Program of the Joint Genome Institute, a DOE Office of Science User Facility. The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under Contract No DE-AC02-05CH11231.

Keywords: multiplex editing, gene duplication, triterpenes, glabrous, genomic dropout

ABSTRACT

As the focus for CRISPR edited plants moves from proof-of-concept to real world applications, precise gene manipulation will increasingly require concurrent multiplex editing for polygenic traits. A common approach for editing across multiple sites is to design one gRNA per target; however, this complicates construct assembly and increases the possibility of off-target mutations. In this study, we utilized one gRNA to target *MYB186*, a known positive trichome regulator, as well as its paralogs *MYB138* and *MYB38* at a consensus site for mutagenesis in *Populus tremula* × *P. alba* INRA 717-1B4. Unexpected duplications of *MYB186* and *MYB138* resulted in a total of eight alleles for the three targeted genes in the hybrid poplar. Deep sequencing and PCR analyses confirmed editing across all eight targets in nearly all of the resultant glabrous mutants, ranging from small indels to large genomic dropouts, with no off-target activity detected at four potential sites. This highlights the effectiveness of a single gRNA targeting conserved exonic regions for multiplex editing. Additionally, cuticular wax and whole leaf analyses showed a complete absence of triterpenes in the trichomeless mutants, hinting at a previously undescribed role for the non-glandular trichomes of poplar.

INTRODUCTION

CRISPR (clustered regularly interspaced short palindromic repeats) technology has been adopted for plant genome editing in an increasing number of species for both basic and applied research (Bewg et al., 2018; Chen et al., 2019; Nasti and Voytas, 2021). The power of CRISPR is due in part to its simplicity with just two core components (Jinek et al., 2012): a nuclear-localized endonuclease, such as CRISPR-associated Cas9 that works universally across all domains of life and a synthetic guide RNA (gRNA) that is customizable and scalable for sequence-specific targeting. With its proven precision and efficiency (Li et al., 2013; Nekrasov et al., 2013; Shan et al., 2013; Endo et al., 2016) and given the polygenic nature of many agronomic traits, there is growing interest in targeting multiple loci for simultaneous CRISPR editing to aid gene function investigation and/or trait engineering (Armario Najera et al., 2019).

Multiplex editing usually involves coexpression of multiple guide RNAs (gRNAs). For the classic CRISPR/Cas9 system, this has been demonstrated using individual gRNA cassettes each driven by a separate RNA polymerase III (Pol III) promoter (Xing et al., 2014; Lowder et al., 2015; Ma et al., 2015). Alternatively, multiple gRNAs can be expressed in tandem with tRNAs as a single polycistronic transcript and processed into individual gRNAs using endogenous tRNA processing machinery (Xie et al., 2015). Polycistronic gRNA transcripts have also been engineered with built-in RNA cleavage sites for processing by ribozymes or the CRISPR-associated endoribonuclease Csy4 (Qi et al., 2012; Gao and Zhao, 2014; Tang et al., 2016; Čermák et al., 2017; Tang et al., 2019). In several cases, functional gRNAs were generated from a single transcriptional unit of *Cas9* fused with an artificial gRNA array without specific flanking sequences (Mikami et al., 2017; Wang et al., 2018). It has been reported that up to eight gRNAs have been successfully deployed for multiplex editing (Ma et al., 2015; Xie et al., 2015; Čermák et al., 2017).

An understudied approach is the use of single gRNAs to target homologous sequences at discrete loci. The capability was showcased by effective inactivation of all 62 copies of porcine endogenous retroviruses in an immortalized pig cell line using two gRNAs to target a highly conserved region of the polymerase (*pol*) gene (Yang et al., 2015). Besides parasitic elements, a single consensus gRNA has also been used to edit paralogs derived from various gene duplication events in soybean and sorghum (Jacobs et al., 2015; Li et al., 2018) or homoeologs in polyploid wheat and oilseed rape (Braatz et al., 2017; Zhang et al., 2017). Multiplex targeting of duplicated genes is especially important for

investigation of functional redundancy in plant genomes that are shaped by whole-genome, segmental,

tandem, and/or transposon-mediated duplications (Flagel and Wendel, 2009; Panchy et al., 2016).

Depending on the duplication age and subsequent selection constraints, sequence similarity can be very

high among duplicates, enabling identification of consensus target sites for multiplex editing by a single

gRNA. This approach greatly simplifies construct design and assembly, reduces off-target potential that

increases with the number of gRNAs (McCarty et al., 2020), and can be bundled with other multi-gRNA

editing strategies discussed above for higher-order multiplex targeting of distinct gene families.

The present study explored the utility of a single gRNA for multiplex editing in an outcrossing woody

perennial, *Populus tremula* × *P. alba* INRA 717-1B4 (hereon referred to as 717). As an interspecific

hybrid, the 717 genome is highly heterozygous which presents additional challenges to gRNA design and

edit outcome determination (Xue et al., 2015). Using trichomes as visual reporter, we targeted a known

positive regulator of trichome development, *PtaMYB186* (Plett et al., 2010), and its close paralogs

PtaMYB138 and *PtaMYB38* for knockout (KO). We show that a single gRNA with SNP-aware design is

effective for multiplex KO of paralogous genes and robust against copy number variations in a hybrid

genome with an unexpected tandem duplication in one of its sub-genomes. We employed multiple

approaches to address the analytical challenge of discriminating among highly similar target sites to

discern mutations that ranged from small indels to large genomic dropouts. Finally, analysis of the

resultant trichomeless mutants revealed a complete absence of triterpenes, and implicated a role for

poplar trichomes in triterpene accrual.

RESULTS

Multiplex CRISPR/Cas9 editing of trichome-regulating MYBs

The known positive regulator of trichome initiation *PtaMYB186* (Plett et al., 2010) corresponds to gene

model Potri.008G089200 in the *P. trichocarpa* v3.1 genome. It belongs to clade 15 of the R2R3-MYB

protein family tree (Wilkins et al., 2009), which is expanded in poplar and contains three additional

members, *MYB138*, *MYB38* and *MYB83*, with as yet unclear functions. The four clade 15 members are

derived from multiple duplication events, based on whole paranome K_S (synonymous distance)

distribution and gene collinearity analyses using the wgd program (Zwaenepoel and Van de Peer, 2018).

These include an ancient (gamma) whole genome duplication (*MYB186* and *MYB83*, K_S = 3.76), a Salicoid

duplication (*MYB186/MYB138* and *MYB38*, K_S = 0.21-0.22), and a tandem duplication (*MYB186* and

MYB138, K_S = 0.0001) (Figure 1). MYB186, MYB138 and MYB38 share higher levels (88-96%) of amino

acid sequence similarity than with MYB83 (55-57%). To ascertain these MYB involvement in trichome

development, we mined RNA-seq data from different stages of 717 leaf development. Transcript levels

of *MYB186*, *MYB138* and *MYB38* were highest in newly emerged leaves (Leaf Plastochron Index LPI-1)

when trichome initiation occurs (Plett et al., 2010), but quickly declined thereafter in expanding (LPI-5)

and mature (LPI-15) leaves (Figure 1). In contrast, *MYB83* transcripts were detected throughout leaf

maturation (Figure 1), weakening support for its potential involvement in trichome development.

We designed a single gRNA to target a conserved region in exon two of *MYB186*, *MYB138* and *MYB38*

(Figure 2A) based on the *P. trichocarpa* v3.1 reference genome and cross-checked using the 717 variant

database (Xue et al., 2015; Zhou et al., 2015) to assure the gRNA target sites were SNP-free in 717. Two

CRISPR/Cas9 constructs were generated (see Methods); the first erroneously omitted a guanine

between the gRNA and the scaffold sequences (referred to as ΔG, Figure 2B), which was corrected in the

second construct (Figure 2A). Both constructs were used for 717 transformation in order to learn

whether ΔG would affect CRISPR/Cas9 editing. In total, 28 independent events generated from the ΔG

construct were all phenotypically indistinguishable from the wild type (WT) and Cas9-only controls

(Figure 2C-J). In contrast, 37 independent events generated from the correct knock out (KO) construct

were glabrous (Figure 2N-R), and one glabrous-like event (KO-27) had a greatly reduced trichome

density across all shoot tissues (leaf, petiole and stem) independent of age (Figure 2K-M). SEM imaging

revealed no trichome initiation or development on the abaxial leaf surface of the glabrous mutants

(Figure 2Q). Epidermal cell morphology of young leaves from tissue cultured plants did not differ

between control and mutant genotypes on either their abaxial (Figure 2F, N) or adaxial surfaces (Figure

2J, R). These results are consistent with roles for MYB186 (Plett et al., 2010) and its paralogs MYB138

and MYB38 in trichome initiation and development.

Mutation spectrum of duplicated 717 *MYB* alleles

A random selection of 30 glabrous events, 28 ΔG events, two Cas9-only events and four WT plants were

subject to amplicon deep-sequencing using consensus primers for *MYB186*, *MYB138* and *MYB38*. Initial

analysis by AGEseq (Xue and Tsai, 2015) showed numerous chimeric edits (mix of edited and unedited

sequences at a given site) not observed in other CRISPR/Cas9-edited 717 transgenics in our experience

(Zhou et al., 2015; Bewg et al., 2018; Tsai et al., 2020). *De novo* assembly of amplicon reads from control

samples revealed seven distinct sequences, more than the expected six alleles of the three target genes.

Blast search against the preliminary 717 genome assemblies by the Joint Genome Institute uncovered an

unexpected copy number variation in 717 relative to the *P. trichocarpa* reference genome. The region

containing paralogous *MYB186* and *MYB138* on Chromosome (Chr) 8 is found as a tandem duplicate in

one of the 717 subgenomes (Figure 3A). This results in three alleles each for *MYB186* and *MYB138* (two

on the main subgenome [Chr8m] and one on the alternative subgenome [Chr8a]) and two alleles for

MYB38 on Chr10 (Chr10m and Chr10a, Figure 3A). Two of the eight alleles were identical in the (original)

amplicon region, explaining the seven distinct sequences we recovered from *de novo* assembly. Based

on the 717 assemblies, we redesigned primers to ensure the amplicons span allele-specific SNP(s) to aid

mutation pattern determination of the eight alleles.

Amplicon-sequencing showed no editing in the 28 ΔG events, except one (ΔG-24) with a 9 bp deletion at

one of the eight target sites (Supplemental Dataset S1). This translates into a mutation rate of 0.45%

(one out of 224 potential target sites), which suggests a negative effect of the ΔG on CRISPR/Cas9

function (hereafter, the ΔG plants were treated as transformation controls). In contrast, we confirmed

successful editing across the eight alleles in all glabrous mutants except KO-27 (Figure 3B-C,

Supplemental Dataset S1). This event showed six edited and two WT (unedited) alleles, consistent with

trichome detection in this line (Figure 2K-M). In aggregate, small insertions and deletions (indels) were

the predominant edits at all sites (Figure 3B-D), with frameshift deletions of 1 bp (-1), 2 bp (-2) and 4 bp

(-4) accounting for over three quarters of the indel mutations (Figure 3D). In-frame deletions (-3 or -6)

accounted for 10% of indels and were detected in 14 events, including KO-27 (Figure 3B-DThese in-

frame mutations are unlikely functional because the gRNA target site is located within the third α-helix

of the R2 domain critical for MYB-DNA interaction (Wang et al., 2020), and because 13 of the events

with in-frame mutations are glabrous. We therefore conclude that all small indels we detected are null

mutations.

Large genomic dropouts between tandem genes

The vast majority (80%) of the sequenced mutants also harbored potentially large deletions as

evidenced by the dearth of mapped amplicon reads at the target sites, referred to as no-amplification

(NA) alleles (Figure 3B-D, Supplemental Dataset S1). The NA frequencies differed by chromosome

position and were positively correlated with copy number, being highest at the Chr8m sites (four

tandem copies), followed by the Chr8a sites (two tandem copies) and least at the single-copy Chr10 sites

(Figure 3A-B). The NA alleles on Chr8 often spanned consecutive copies, suggesting large dropouts

between two gRNA cleavage sites. To support this idea, we examined a subset of mutant lines using

allele-specific primers for PCR amplification of the target genes. As expected, NA alleles yielded no PCR

products, whereas alleles previously detected by amplicon sequencing produced observable PCR

products (Supplemental Figure S1). We next used consensus primers for PCR amplification of all six Chr8

(*MYB186* and *MYB138*) alleles, approximately 850 – 950 bp, from three control plants and four KO lines

each with 4-5 NA alleles on Chr8. These KO lines had reduced PCR band intensity when compared with

controls (Figure 4A-B). Sanger sequencing of the PCR products resulted in clean chromatograms with

clear nucleotide peaks throughout the sequenced length for KO-5 and KO-69 (Figure 4C), two mutant

lines with only one detectable Chr8 allele (Figure 4B). In contrast, the chromatograms for KO-63, KO-70

(both containing two detectable Chr8 alleles) and WT samples were noisy as would be expected for

mixed template (Figure 4B-C). The Sanger sequencing data of KO-5 and KO-69 not only confirmed the

indel pattern (-2 in both cases) detected by amplicon sequencing, but also supported the occurrence of

gene fusion between two gRNA cleavage sites, based on SNP patterns upstream and downstream of the

gRNA target (Figure 4B-C). KO-5 harbors a fusion junction between *MYB186m1* and *MYB138m1* with a

~29 Kb genomic dropout, whereas KO-69 contains a fusion of *MYB138m1* and *MYB138m2* with a ~62 Kb

genomic dropout (Figure 4B-C, Supplemental Figure S2). Both events likely contain additional large

deletions or genomic fusions, as allele(s) downstream (KO-5) or upstream (KO-69) of the respective

fusion point could not be PCR amplified (Figure 4B). Regardless, our findings show that a single gRNA is

highly effective for multiplex KO of tandem duplicates via either small indels or large deletions.

Assessment of off-target activity in mutants

A combination of computational prediction and experimental verification was used to assess off-target

effects. Potential off-target sites of the gRNA were predicted by CCTop (Stemmer et al., 2015) using the

P. trichocarpa v3.1 reference genome as well as the two SNP-substituted Pta717 v2 (*P. alba* and *P.*

tremula) subgenomes (Xue et al., 2015). The same four exonic locations were ranked among the top

potential off-target sites (excluding intergenic or unassembled scaffold sequences) across the three

genomes, each having three mismatches with the gRNA sequence. We designed three sets of primers to

examine potential editing at the four off-target sites; OT1 (Potri.004G115600 and Potri.004G118000),

OT2 (Potri.004G138000), and OT3 (Potri.014G024400). Amplicon sequencing of 20 trichomeless mutants

found no off-target activity across these four sites (Supplemental Dataset S2).

Absence of triterpenes in trichomeless leaves

Trichomes as epidermal outgrowths are covered with waxy cuticles like other epidermis cells (Hegebarth

et al., 2016). The striking glabrous phenotype of the mutants prompted us to compare leaf wax

composition between control and trichomeless plants. Total wax load of mature leaves (extractable wax

from leaf surface) did not change significantly between genotypes (Figure 5A). Alkanes were the most

abundant class of leaf cuticular waxes detected in 717 and differed little between control and

trichomeless plants (Figure 5B). In contrast, levels of triterpenes, fatty alcohols and β-sitosterol were

significantly reduced in the mutants (Figure 5B-D). Specifically, the wax of mutant leaves was devoid of

any triterpenes, including α-amyrin, β-amyrin, β-amyrone and lupenone (Figure 5E). Two primary

alcohols, 1-octacosanol (C28) and 1-hexacosanol (C26), were depleted in the mutants by >50% (Figure

5C), and β-sitosterol, by 42% (Figure 5D). To further investigate the absence of triterpenes in the mutant

wax, whole leaf tissues were also profiled for compounds that were significantly reduced in cuticular

wax. Again, triterpenes were not detected in the leaves of trichomeless mutants (Figure 5E), whereas 1-

octacosanol, 1-hexacosanol and β-sitosterol were detected at levels comparable with controls (Figure

5C,D). The data support a previously unsuspected link between triterpene accrual and non-glandular

trichomes in poplar.

DISCUSSION

The present study demonstrates that a single gRNA targeting conserved genomic sites is highly effective

for multiplex editing in poplar. The 30 independent KO lines experienced an average of 5.4 CRISPR/Cas9-

mediated cleavages per line based on indel alleles, which is likely an underestimate because many NA

alleles may also result from CRISPR/Cas9 cleavages as shown for KO-5 and KO-69 (Figures 3 and 4). The

unexpected genomic complexity in the hybrid 717 highlights the importance of ensuring SNP-free

targets for gRNA design (Zhou et al., 2015), as well as the challenge of decoding multiplexed edits among

highly homologous gene duplicates.

The negligible editing by the ΔG construct (Supplementary Dataset S1) provides insight into scaffold

structure and stability. The ΔG configuration can lead to two hypothetical outcomes: either the guanine

is omitted from the scaffold and the gRNA remains intact and capable of base pairing to the target sites

for Cas9 cleavage, or the guanine is sequestered for secondary structure folding of the scaffold, resulting

in a 3▯-truncated gRNA no longer PAM-adjacent at the target sites (Figure 2B). The lack of mutations in

ΔG transformants supports the latter scenario and is consistent with transcription and folding of gRNA

molecules preceding their base-pairing with genomic targets. Our finding suggests that 3'-truncated

gRNA could serve as an alternative approach for generating transgenic controls.

A number of methods are commonly used for decoding CRISPR-mediated mutation patterns, including

restriction digestion, endonuclease-based mismatch detection, gene-/allele-specific PCR sometimes in

conjunction with cloning and/or Sanger sequencing (Figure 4), or amplicon deep sequencing (Figure 3).

The pros and cons of these methods have been discussed elsewhere (e.g., Germini et al., 2018). Analysis

of genome editing across multiple target sites poses additional challenges over mono-targeted

experiments, especially when highly homologous target and flanking sequences are encountered. These

multiplex scenarios generally exceed the resolution of most methods or may require additional assays

(*e.g.*, allele-specific PCR) to determine editing outcomes. In the case exemplified here, deep sequencing

of a pooled library of amplicons obtained with consensus primers for all eight target alleles was highly

effective for decoding multiplexed edits. The use of consensus primers provides built-in controls for

each PCR, allowing for high confidence calling of NA alleles (see Supplemental Dataset S1) which are

otherwise difficult to distinguish from failed PCR in individual reactions. As another advantage, the

amplicon deep sequencing data can be used for *de novo* assembly which in our case led to the discovery

of unexpected copy number variations of MYB186 and MYB138 in the experimental poplar 717

genotype. Although technical limitations remain in short-read mapping to highly homologous sites,

inclusion of allele-specific SNPs within the amplicon region and adoption of bioinformatic programs with

parameter tuning capabilities (*e.g.*, AGEseq) are key to multiplexed mutation pattern determination.

The glabrous mutants (Figure 2) provide strong support for an essential role of PtaMYB186/138/38 in

the initiation of trichome development in 717. Additionally, the low trichome density of KO-27 suggests

that MYB38 plays a redundant but minor role in leaf/stem trichome initiation (Figures 2 and 3). Follow-

up research, including allele-specific KOs, is needed to dissect the functional redundancy and allele-dose

response of clade 15 MYB members more fully. The unedited (WT) *MYB38* alleles in KO-27 appear stable

during vegetative propagation as this event has maintained a low trichome density for over two years in

both tissue culture and greenhouse environments. This adds to previously reported stability of CRISPR

editing outcomes in clonally propagated poplar (Bewg et al., 2018)

The loss of trichomes did not significantly affect the total epidermal wax load but led to a complete

absence of triterpenes both in cuticular wax and whole leaves of the mutants. It is unlikely that

MYB186/138/38 have an additional role in triterpene biosynthesis (*i.e.*, lack of triterpenes as a direct KO

effect) because of their recent duplication history (Figure 1) and because a recent report implicated

phylogenetically distinct MYBs in triterpene regulation (Falginella et al., 2021). We interpret the absence

of triterpenes in trichomeless leaves as suggesting a role for non-glandular trichomes in triterpene

accrual in poplar. While glandular trichomes are well known for their roles in biosynthesis and storage of

terpenes (Lange and Turner, 2013), the presence of terpenes in non-glandular trichomes has only been

reported recently (Santos Tozin et al., 2016; Dmitruk et al., 2019). The genetic evidence presented

herein provides strong support for a functional link between triterpenes and non-glandular trichomes

that warrants further investigation.

The glabrous phenotype of the null mutants we obtained highlights the potential utility of trichomes as

a visual reporter. Assessments of CRISPR/Cas functionality often target the chlorophyll biosynthetic

enzyme phytoene desaturase (PDS) (Norris et al., 1995), as mutations result in an albino phenotype

(Shan et al., 2013; Ma et al., 2015; Xie et al., 2015). Whilst phenotypically obvious, *PDS* mutations are

lethal for the regenerated plant, thus limiting follow-up investigations. Alternatively, the glabrous

phenotype achieved by KO of trichome-regulating *MYBs* is non-lethal and no inhibition to plant growth

was detected. This allows stacked mutagenesis of these mutants, including reparative transformations

to restore trichome initiation. The use of trichomes as a visual reporter for CRISPR/Cas9 mutation or

repair of a defective allele has been established in Arabidopsis (Hahn et al., 2017; Hahn et al., 2018)

which provides support for further developing this system in poplar.

MATERIALS AND METHODS

Generation of KO mutants

The ΔG and KO constructs in p201N-Cas9 (Jacobs et al., 2015) were prepared by Gibson assembly. PCR

was used to amplify the p201N-Cas9 binary vector following *Swa*I (New England BioLabs) digestion, and

the *Medicago truncatula Mt*U6.6 promoter and scaffold fragments from HindIII and EcoRI (New England

BioLabs) digested pUC-gRNA shuttle vector (Jacobs et al., 2015), with Q5 High-Fidelity DNA Polymerase

(New England BioLabs) and primers (Sigma) listed in Supplemental Table S1. The p201N-Cas9 (Addgene

59175) and pUC-gRNA (Addgene 47024) plasmids were both gifts from Wayne Parrott. Two pairs of

oligos (Sigma) corresponding to the consensus gRNA target site in exon two of *MYB186*

(Potri.008G089200), *MYB138* (Potri.008G089700) and *MYB38* (Potri.010G165700) were assembled with

p201N-Cas9. The NEBuilder HiFi DNA Assembly Cloning Kit (New England Biolobs) was used to assemble

p201N-Cas9, *Mt*U6.6 promoter and scaffold fragments with a pair of oligos containing the gRNA target

sequence (Supplemental Table S1). Following transformation into DH5α *E. coli* (Zymo Research Mix &

Go! Competent Cells), PCR-positive colonies were used for plasmid purification before Sanger

sequencing (Eurofins) confirmation. Plasmids were then heat-shocked into *Agrobacterium tumefaciens*

strain C58/GV3101 (pMP90) (Koncz and Schell, 1986) and confirmed by colony PCR.

Populus tremula x *alba* (IRNA 717-1B4) transformation and regeneration was performed as outlined in

Meilan and Ma (2006), except 0.05 mg/L 6-benzylaminopurine was used in shoot elongation media, and

200 mg/L L-glutamine was added to all media, with 3 g/L gellan gum (PhytoTechnology Lab) as a gelling

agent. Following a 2-day agrobacterial cocultivation, leaf discs were washed in sterile water followed by

washing in 200 mg/L cefotaxime and 300 mg/L timentin with shaking for 1.5 hr. Transformants were

selected on media supplemented with 100 mg/L kanamycin, 200 mg/L cefotaxime and 300 mg/L

timentin for callus induction and shoot regeneration and with kanamycin and timentin for shoot

elongation and rooting. All cultures were grown and maintained at 22°C under a 16-hr light/8-hr dark

photoperiod with Growlite® FPV24 LED (Barron Lighting Group) at ~150 μmol/m^2/s.

RNA-seq analysis

For developmental profiling, LPI-1, LPI-5 and LPI-15 were collected from three greenhouse-grown WT

plants (~5 ft in height) for RNA extraction using Direct-zol RNA MiniPrep kit (Zymo Research) with Plant

RNA Purification Reagent (Invitrogen). RNA-seq library preparation and Illumina NextSeq 500 sequencing

was performed at the Georgia Genomics and Bioinformatics Core. We obtained 10.8-13.3 PE75 reads

per sample. After pre-processing to remove adapter and rRNA sequences, reads were mapped to the

717 SNP-substituted genome sPta717 v2 (Xue et al., 2015) using STAR v2.5.3a (Dobin and Gingeras,

2015). Transcript abundance in FPKM (fragments per kilobase of transcript per million mapped reads)

was estimated by featureCounts v1.5.2 (Liao et al., 2014).

Amplicon sequencing determination of mutation spectrums

Newly emerged leaves were excised from individual events in tissue culture for genomic DNA extraction

(Dellaporta et al., 1983). The DNA pellet was resuspended in water with RNase A (10 µg/mL) for

amplicon library preparation using GoTaq G2 Green Master Mix (Promega) and primers (Supplemental

Table S1) spanning the gRNA target site (between 264 bp to 280 bp). Samples were then barcoded with

Illumina amplicon indexing primers and pooled for Illumina MiSeq nano PE150 sequencing performed at

the University of Georgia's Georgia Genomics and Bioinformatics Core. Demultiplexed sequence reads

were analyzed by the AGEseq (Analysis of Genome Editing by Sequencing) program (Xue and Tsai, 2015),

with mismatch allowance set at 1%, followed by manual curation.

Because initial amplicon data analysis revealed lower editing efficiencies (<90%) than we typically

observed in 717 (Zhou et al., 2015; Bewg et al., 2018) at several target sites, we performed *de novo*

assembly of WT amplicon reads using Geneious, and recovered seven distinct alleles. We then searched

the JGI draft 717 genome assembly v1.0 with the *P. trichocarpa* Nisqually-1 v3.1 (Phytozome v12)

MYB186, *MYB138* and *MYB38* gene models and extracted the surrounding 50-150 Kb regions from Chr8

and Chr10 for manual annotation against the *P. trichocarpa* Nisqually-1 reference (Figure 3A). The

matching *MYB* gene sequences were extracted for error correction using 717 resequencing data (Xue et

al., 2015). Curated sequences were used for new (amplicon and allele-/gene-specific) primer design and

as references in amplicon data analysis. In the case of WT and transgenic controls with no editing,

erroneous read assignments—and hence indel calls—still remained because the amplicon region

between some alleles differs only in the number of intronic dinucleotide (GT) repeats (Supplemental

Dataset S1). Misassigned reads led to erroneous indel calls of -2, +2 or their multiples outside of the

gRNA target site. For this reason, WT and control samples were processed by ustacks from Stacks 2.3

(Catchen et al., 2011). Parameters were adjusted to avoid collapsing reads with SNPs and/or Indels from

paralogous alleles into the same tag group and gapped alignments were disabled. Tags from the output

were then used for allele assignment.

Determination of leaf and cuticle wax compositions

Leaf punches (25 mm diameter) were taken from mature leaves of similar size (between LPI-10 and LPI-

15) of soil-grown plants in a growth chamber and washed in 4 mL of methylene chloride for 30 sec. The

washes were dried under a continuous N_2 stream before resuspension in 400 µL chloroform. A 200 µL

aliquot was subsequently dried under vacuum and the residues shipped to the Oak Ridge National

Laboratory for analysis. Sorbitol (1 mg/mL) was added to the residues as an internal standard and re-

dried under N_2. For whole leaf analysis, liquid nitrogen-ground and freeze-dried powders from LPI-5 (25

mg) of control and KO plants were extracted by 80% ethanol to which sorbitol (1 mg/mL) was added and

dried under N_2. The samples were derivatized prior to analysis on an Agilent Technologies 7890A GC

coupled to a 5975C inert XL MS fitted with an Rtx-5MS capillary column with a 5m Integra-Guard column

(Restek) as described in Holwerda et al. (2014). Compound identification was based on mass spectral

fragmentation patterns against the NIST08 database and an in-house library built with authentic

standards.

Tissue Imaging and SEM analysis

Images of poplar were taken with either a Google Pixel 3a running Android v11, or a Leica M165 FC

dissection microscope attached to a Leica DFC500 camera running Leica Application Suite software

v3.8.0. Scanning electron microscopic (SEM) observations were obtained using Hitachi 3400 NII (Hitachi

High Technologies America) microscope following optimized protocols at the Center for Ultrastructural

Research at the Fort Valley State University. LPI-1 from growth chamber plants or young leaves of tissue

culture plants were processed for primary fixation at 25°C in 2 % glutaraldehyde (Electron Microscopy

Sciences, EMS) prepared with Sorensen's Phosphate buffer, pH 7.2 (EMS) for one hour and then washed

three times for 15 min each with the same buffer before secondary fixation in 1% osmium tetroxide

(EMS) prepared in Sorensen's Phosphate buffer, pH 7.2 for 1 hour at 25°C. After three washes with dH_2O

for 15 min each, fixed tissues were dehydrated with ethanol series passing through 25%, 50%, 75%, and

95% for 15 min each, followed by three changes of 100% ethanol for 15 min each. Critical point drying of

fixed samples was conducted using a critical point dryer (Leica) and then samples were placed on Hitachi

M4 aluminum specimen mounts (Ted Pella) using double sided carbon adhesive tabs (EMS) for coating.

Gold coating of 50 Å thickness was done for 60 sec using sputter coater (Denton Desk V) under a vacuum

pressure of 0.05 torr. Image acquisition in various magnification was done at accelerating voltage of 5

KV.

ACCESSION NUMBERS

The RNA-seq data has been deposited to the National Center for Biotechnology Information's Sequence

Read Archive under accession No. PRJNA753499.

ACKNOWLEDGEMENTS

The authors thank Gilles Pilate of the Institut National de la Recherche Agronomique, France for

providing poplar clone INRA 717-1B4, Hongduyen Pham and Margot Chen for tissue culture assistance,

Yingying Zhu for RNA from developmentally staged leaves and Liang-Jiao Xue for guidance on RNA-seq

data processing. We additionally thank the Department of Energy Joint Genome Institute and

collaborators for prepublication access to the *Populus tremula* x *P. alba* (IRNA 717-1B4) genome

sequence and annotation.

SUPPORTING INFORMATION

Table S1. Primers used in this study.

Figure S1. PCR confirmation of NA alleles using allele-specific primers.

Figure S2. Sequence alignment of wild type and fusion *MYB* alleles from KO-5 and KO-69.

Dataset S1. CRISPR/Cas9 mutation patterns of the eight target *MYB* alleles in ΔG and KO lines.

Dataset S2. Assessment of off-target activity in trichomeless mutants.

FIGURE LEGENDS

Figure 1. Expression of clade 15 *MYB* transcription factors during *Populus* leaf maturation.

A simplified phylogenetic tree is shown with duplication history noted on the left. Data are mean±SD of

n=3. LPI, leaf plastochron index; FPKM, fragments per kilobase of transcript per million mapped reads;

MYB186, Potri.008G089200; MYB138, Potri.008G089700; MYB38, Potri.010G165700; and MYB83,

Potri.017G086300.

Figure 2. CRISPR/Cas9 KO of trichome-regulating *MYBs*.

A, Schematic illustrations of the *MYB* gene structure, gRNA target site, and base pairing between the

genomic target (black) and the gRNA spacer (red)-scaffold (blue) molecule. Black line denotes the

protospacer adjacent motif (PAM). **B**, Zoomed-in view of the ΔG vector configuration at the gRNA

spacer-scaffold junction with a guanine omission. **C-R**, Representative shoot tip (C, G, K, O) and LPI-1

abaxial (D, H, L, P) phenotypes and SEM images (E, F, I, J, M, N, Q, R) of soil-grown WT (D, E), Cas9 vector

control (C), ΔG control (G-I), KO-27 (K-M), and null mutant (O-Q) plants, and leaf abaxial (F, N) or adaxial

(J, R) images of tissue cultured ΔG (F, J) and null mutant (N, R) plants. Scale bar = 3 mm (D, H, L, P), 500

μm (E, I, M), 1 mm (Q), or 25 μm (F, J, N, R).

Figure 3. Mutation analysis of trichomeless mutants.

A, Schematic illustration of *MYB186* and *MYB138* on Chr8 subgenomes (main and alternative, or Chr8m

and Chr8a, respectively) and *MYB38* on Chr10m and Chr10a of the 717 genome. Neighboring genes are

color coded for synteny and the putative duplication block containing *MYB186* and *MYB138* on Chr8 is

marked by red brackets. Black triangles denote the eight gRNA target sites. **B**, Mutation spectrum

determined by amplicon sequencing. The eight alleles are arranged by genomic position for each plant

line and color-coded for the editing outcomes: green, unedited; orange, indel; and grey, no amplification

(NA). **C**, Representative amplicon sequencing output of three mutant events. All eight alleles, their

detection frequencies and indel patterns (mapped read count and percentage with the indicated

pattern) are shown and colored as in B. The gRNA target sequence is shown on top and protospacer

adjacent motif underlined. Allele-discriminating SNPs are shown in pink (see Supplemental Dataset S1

for the full data). **D**, Pie chart summary of the overall (left) and indel (right) editing patterns.

Figure 4. PCR analysis of selected mutant lines.

A, PCR amplification of the six *MYB* alleles on Chr8 from two WT, one ΔG and four KO lines. The four KO

lines were selected to represent one (KO-5 and KO-69) or two (KO-63 and KO-70) remaining Chr08

alleles. *UBC* (ubiquitin-conjugating enzyme) was included as loading control. M, molecular weight

marker; ntc, no-template control. **B**, Mutation patterns of Chr08 alleles in the selected mutant lines as

shown in Figure 3B. Orange triangles connected by dashes represent fusion junctions shown in C. **C**,

Sanger sequencing of PCR products from A. Sequence alignment of the six alleles flanking the gRNA

target site (red) is shown on top and chromatograms of the same region are shown below. Grey shaded

alignments are introns, with allele-discriminating SNPs shown in pink and homologous intron 2

sequences in blue (shifted upstream by 21 bp in *138m1* and *186m2* due to gaps). PAM is underlined and

boxed in blue for correspondence with the sequence traces below. Black triangles denote the Cas9

cleavage site and black dashed box corresponds to the 2-bp deletion (-2) detected in KO-5 and KO-69.

The two fusion alleles as determined by SNPs are marked below the KO-5 and KO-69 traces (see

Supplemental Figure S2 for the full sequence alignment).

Figure 5. Cuticular wax composition of trichomeless and control leaves.

A, Total wax load. **B**, Major classes of cuticular wax. **C**, Fatty alcohols (C26, 1-hexacosanol; C28, 1-

octacosanol) in wax (left) or whole leaves (right). **D**, β-sitosterol detected in wax (left) or whole leaves

(right). **E**, Triterpenes detected in wax (top) or whole leaves (bottom). Ergosterone, 14,24-dimethyl-

ergosta-8,25-dien-3-one; cycloartanone, 24-methylene cycloartan-3-one; lanosterone, lanosta-8,24-

dien-3-one. Data are mean±SD of n=5. All concentration estimates were based on sorbitol equivalent.

Statistical significance was determined by 2-tailed *t*-test (* *P*<0.05, ** *P*<0.01, *** *P*<0.001). nd, not

detected.

REFERENCES

Armario Najera V, Twyman RM, Christou P, Zhu C (2019) Applications of multiplex genome editing in
higher plants. Current Opinion in Biotechnology **59**: 93-102
Bewg WP, Ci D, Tsai C-J (2018) Genome editing in trees: From multiple repair pathways to long-term
stability. Frontiers in Plant Science **9**: 1732
Braatz J, Harloff H-J, Mascher M, Stein N, Himmelbach A, Jung C (2017) CRISPR-Cas9 targeted
mutagenesis leads to simultaneous modification of different homoeologous gene copies in
polyploid oilseed rape (*Brassica napus*). Plant Physiology **174**: 935-942
Catchen JM, Amores A, Hohenlohe P, Cresko W, Postlethwait JH (2011) Stacks: Building and genotyping
loci *de novo* from short-read sequences. Genes Genomes Genetics **1**: 171-182
Čermák T, Curtin SJ, Gil-Humanes J, Čegan R, Kono TJY, Konečná E, Belanto JJ, Starker CG, Mathre JW,
Greenstein RL, Voytas DF (2017) A multipurpose toolkit to enable advanced genome
engineering in plants. Plant Cell **29**: 1196-1217
Chen K, Wang Y, Zhang R, Zhang H, Gao C (2019) CRISPR/Cas Genome Editing and Precision Plant
Breeding in Agriculture. Annual Review of Plant Biology **70**: 667-697
Dellaporta SL, Wood J, Hicks JB (1983) A plant DNA minipreparation: Version II. Plant Molecular Biology
Reporter **1**: 19-21
Dmitruk M, Sulborska A, Żuraw B, Stawiarz E, Weryszko-Chmielewska E (2019) Sites of secretion of
bioactive compounds in leaves of *Dracocephalum moldavica* L.: anatomical, histochemical, and
essential oil study. Brazilian Journal of Botany **42**: 701-715
Dobin A, Gingeras TR (2015) Mapping RNA-seq reads with STAR. Curr Protoc Bioinformatics **51**:
11.14.11-11.14.19
Endo A, Masafumi M, Kaya H, Toki S (2016) Efficient targeted mutagenesis of rice and tobacco genomes
using Cpf1 from Francisella novicida. Scientific Reports **6**: 38169
Falginella L, Andre CM, Legay S, Lin-Wang K, Dare AP, Deng C, Rebstock R, Plunkett BJ, Guo L, Cipriani
G, Espley RV (2021) Differential regulation of triterpene biosynthesis induced by an early failure
in cuticle formation in apple. Horticulture Research **8**: 75
Flagel LE, Wendel JF (2009) Gene duplication and evolutionary novelty in plants. New Phytol **183**: 557-
564
Gao Y, Zhao Y (2014) Self-processing of ribozyme-flanked RNAs into guide RNAs in vitro and in vivo for
CRISPR-mediated genome editing. Journal of Integrative Plant Biology **56**: 343-349
Germini D, Tsfasman T, Zakharova VV, Sjakste N, Lipinski M, Vassetzky Y (2018) A comparison of
techniques to evaluate the effectiveness of genome editing. Trends in Biotechnology **36**: 147-
159
Hahn F, Eisenhut M, Mantegazza O, Weber APM (2018) Homology-directed repair of a defective
Glabrous gene in *Arabidopsis* with Cas9-based gene targeting. Frontiers in Plant Science **9**: 424
Hahn F, Mantegazza O, Greiner A, Hegemann P, Eisenhut M, Weber AP (2017) An efficient visual screen
for CRISPR/Cas9 activity in *Arabidopsis thaliana*. Frontiers in Plant Science **8**: 39
Hegebarth D, Buschhaus C, Wu M, Bird D, Jetter R (2016) The composition of surface wax on trichomes
of Arabidopsis thaliana differs from wax on other epidermal cells. Plant Journal **88**: 762-774

Holwerda EK, Thorne PG, Olson DG, Amador-Noguez D, Engle NL, Tschaplinski TJ, van Dijken JP, Lynd
LR (2014) The exometabolome of *Clostridium thermocellum* reveals overflow metabolism at high
cellulose loading. Biotechnology for Biofuels **7**: 155
Jacobs TB, LaFayette PR, Schmitz RJ, Parrott WA (2015) Targeted genome modifications in soybean with
CRISPR/Cas9. BMC Biotechnology **15**: 16
Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E (2012) A Programmable dual-RNA-
guided DNA endonuclease in adaptive bacterial immunity. Science **337**: 816-821
Koncz C, Schell J (1986) The promoter of TL-DNA gene *5* controls the tissue-specific expression of
chimaeric genes carried by a novel type of *Agrobacterium* binary vector. Molecular & General
Genetics **204**: 383-396
Lange BM, Turner GW (2013) Terpenoid biosynthesis in trichomes--current status and future
opportunities. Plant Biotechnol J **11**: 2-22
Li A, Jia S, Yobi A, Ge Z, Sato SJ, Zhang C, Angelovici R, Clemente TE, Holding DR (2018) Editing of an
alpha-kafirin gene family increases, digestibility and protein quality in sorghum. Plant Physiology
177: 1425-1438
Li J-F, Norville JE, Aach J, McCormack M, Zhang D, Bush J, Church GM, Sheen J (2013) Multiplex and
homologous recombination–mediated genome editing in *Arabidopsis* and *Nicotiana*
benthamiana using guide RNA and Cas9. Nature Biotechnology **31**: 688-691
Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. Bioinformatics **30**: 923-930
Lowder LG, Zhang D, Baltes NJ, Paul JW, Tang X, Zheng X, Voytas DF, Hsieh T-F, Zhang Y, Qi Y (2015) A
CRISPR/Cas9 toolbox for multiplexed plant genome editing and transcriptional regulation. Plant
Physiology **169**: 971-985
Ma X, Zhang Q, Zhu Q, Liu W, Chen Y, Qiu R, Wang B, Yang Z, Li H, Lin Y, Xie Y, Shen R, Chen S, Wang Z,
Chen Y, Guo J, Chen L, Zhao X, Dong Z, Liu Y-G (2015) A robust CRISPR/Cas9 system for
convenient, high-efficiency multiplex genome editing in monocot and dicot plants. Molecular
Plant **8**: 1274-1284
McCarty NS, Graham AE, Studená L, Ledesma-Amaro R (2020) Multiplexed CRISPR technologies for
gene editing and transcriptional regulation. Nature Communications **11**: 1281
Meilan R, Ma C (2006) Poplar (*Populus* spp.). *In* K Wang, ed, Agrobacterium Protocols Volume 2.
Methods in Molecular Biology, Vol 344. Humana Press, Totowa, NJ
Mikami M, Toki S, Endo M (2017) In planta processing of the SpCas9–gRNA complex. Plant and Cell
Physiology **58**: 1857-1867
Nasti RA, Voytas DF (2021) Attaining the promise of plant gene editing at scale. Proceedings of the
National Academy of Sciences **118**: e2004846117
Nekrasov V, Staskawicz B, Weigel D, Jones JDG, Kamoun S (2013) Targeted mutagenesis in the model
plant Nicotiana benthamiana using Cas9 RNA-guided endonuclease. Nature Biotechnology **31**:
691-693
Norris SR, Barrette TR, DellaPenna D (1995) Genetic dissection of carotenoid synthesis in Arabidopsis
defines plastoquinone as an essential component of phytoene desaturation. Plant Cell **7**: 2139-
2149
Panchy N, Lehti-Shiu M, Shiu S-H (2016) Evolution of gene duplication in plants. Plant Physiology **171**:
2294-2316
Plett JM, Wilkins O, Campbell MM, Ralph SG, Regan S (2010) Endogenous overexpression of *Populus*
MYB186 increases trichome density, improves insect pest resistance, and impacts plant growth.
Plant Journal **64**: 419-432
Qi L, Haurwitz RE, Shao W, Doudna JA, Arkin AP (2012) RNA processing enables predictable
programming of gene expression. Nature Biotechnology **30**: 1002-1006

Santos Tozin LRd, de Melo Silva SC, Rodrigues TM (2016) Non-glandular trichomes in Lamiaceae and
Verbenaceae species: morphological and histochemical features indicate more than physical
protection. New Zealand Journal of Botany **54**: 446-457
Shan Q, Wang Y, Li J, Zhang Y, Chen K, Liang Z, Zhang K, Liu J, Xi JJ, Qiu J-L, Gao C (2013) Targeted
genome modification of crop plants using a CRISPR-Cas system. Nat Biotech **31**: 686-688
Stemmer M, Thumberger T, del Sol Keyer M, Wittbrodt J, Mateo JL (2015) CCTop: An intuitive, flexible
and reliable CRISPR/Cas9 target prediction tool. PLoS ONE **10**: e0124633
Tang X, Ren Q, Yang L, Bao Y, Zhong Z, He Y, Liu S, Qi C, Liu B, Wang Y, Sretenovic S, Zhang Y, Zheng X,
Zhang T, Qi Y, Zhang Y (2019) Single transcript unit CRISPR 2.0 systems for robust Cas9 and
Cas12a mediated plant genome editing. Plant Biotechnology Journal **17**: 1431-1445
Tang X, Zheng X, Qi Y, Zhang D, Cheng Y, Tang A, Voytas Daniel F, Zhang Y (2016) A single transcript
CRISPR-Cas9 system for efficient genome editing in plants. Molecular Plant **9**: 1088-1091
Tsai C-J, Xu P, Xue L-J, Hu H, Nyamdari B, Naran R, Zhou X, Goeminne G, Gao R, Gjersing E, Dahlen J,
Pattathil S, Hahn MG, Davis M, Ralph J, Boerjan WA, Harding SA (2020) Compensatory guaiacyl
lignin biosynthesis at the expense of syringyl lignin in *4CL1*-knockout poplar. Plant Physiology
183: 123-136
Wang B, Luo Q, Li Y, Yin L, Zhou N, Li X, Gan J, Dong A (2020) Structural insights into target DNA
recognition by R2R3-MYB transcription factors. Nucleic Acids Research **48**: 460-471
Wang M, Mao Y, Lu Y, Wang Z, Tao X, Zhu J-K (2018) Multiplex gene editing in rice with simplified
CRISPR-Cpf1 and CRISPR-Cas9 systems. Journal of Integrative Plant Biology **60**: 626-631
Wilkins O, Nahal H, Foong J, Provart NJ, Campbell MM (2009) Expansion and diversification of the
Populus R2R3-MYB family of transcription factors. Plant Physiology **149**: 981-993
Xie K, Minkenberg B, Yang Y (2015) Boosting CRISPR/Cas9 multiplex editing capability with the
endogenous tRNA-processing system. Proceedings of the National Academy of Sciences **112**:
3570-3575
Xing H-L, Dong L, Wang Z-P, Zhang H-Y, Han C-Y, Liu B, Wang X-C, Chen Q-J (2014) A CRISPR/Cas9
toolkit for multiplex genome editing in plants. BMC Plant Biology **14**: 327
Xue L-J, Alabady MS, Mohebbi M, Tsai C-J (2015) Exploiting genome variation to improve next-
generation sequencing data analysis and genome editing efficiency in Populus tremula x alba
717-1B4. Tree Genetics & Genomes **11**: 82
Xue LJ, Tsai CJ (2015) AGEseq: Analysis of genome editing by sequencing. Molecular Plant **8**: 1428-1430
Yang L, Güell M, Niu D, George H, Lesha E, Grishin D, Aach J, Shrock E, Xu W, Poci J, Cortazio R,
Wilkinson RA, Fishman JA, Church G (2015) Genome-wide inactivation of porcine endogenous
retroviruses (PERVs). Science **350**: 1101-1104
Zhang Y, Bai Y, Wu G, Zou S, Chen Y, Gao C, Tang D (2017) Simultaneous modification of three
homoeologs of TaEDR1 by genome editing enhances powdery mildew resistance in wheat. Plant
Journal **91**: 714-724
Zhou X, Jacobs TB, Xue L-J, Harding SA, Tsai C-J (2015) Exploiting SNPs for biallelic CRISPR mutations in
the outcrossing woody perennial *Populus* reveals 4-coumarate:CoA ligase specificity and
redundancy. New Phytologist **208**: 298-301
Zwaenepoel A, Van de Peer Y (2018) wgd—simple command line tools for the analysis of ancient whole-
genome duplications. Bioinformatics **35**: 2153-2155

Figure 1. Expression of clade 15 *MYB* transcription factors during *Populus* leaf maturation. A simplified phylogenetic tree is shown with duplication history noted on the left. Data are mean ± SD of n=3. LPI, leaf plastochron index; FPKM, fragments per kilobase of transcript per million mapped reads; MYB186, Potri.008G089200; MYB138, Potri.008G089700; MYB38, Potri.010G165700; and MYB83, Potri.017G086300.

Figure 2. CRISPR/Cas9 KO of trichome-regulating *MYBs*. **A**, Schematic illustrations of the *MYB* gene structure, gRNA target site, and base pairing between the genomic target (black) and the gRNAspacer (red)-scaffold (blue) molecule. Black line denotes the protospacer adjacent motif (PAM). **B**, Zoomed-in view of the ΔG vector configuration at the gRNA spacer-scaffold junction with a guanine omission. **C-R**, Representative shoot tip (C, G, K, O) and LPI-1 abaxial (D, H, L, P) phenotypes and SEM images (E, F, I, J, M, N, Q, R) of soil-grown WT (D, E), Cas9 vector control (C), ΔG control (G-I), KO-27 (K-M), and mutant (O-Q) plants, and leaf abaxial (F, N) or adaxial (J, R) images of tissue cultured ΔG (F, J) and mutant (N, R) plants. Scale bar = 3 mm (D, H, L, P), 500 μm (E, I, M), 1 mm (Q), or 25 μm (F, J, N, R).

Figure 3. Mutation analysis of trichomeless mutants. **A**, Schematic illustration of *MYB186* and *MYB138* on Chr8 subgenomes (main and alternative, or Chr8m and Chr8a, respectively) and *MYB38* on Chr10m and Chr10a of the 717 genome. Neighboring genes are color coded for synteny and the putative duplication block containing *MYB186* and *MYB138* on Chr8 is marked by red brackets. Black triangles denote the eight gRNA target sites. **B**, Mutation spectrum determined by amplicon sequencing. The eight alleles are arranged by genomic position for each plant line and color-coded for the editing outcomes: green, unedited; orange, indel; and grey, no amplification (NA). **C**, Representative amplicon sequencing output of three mutant events. All eight alleles, their detection frequencies and the indel patterns (mapped read count and percentage with the indicated pattern) are shown and colored as in B. The gRNA target sequence is shown on top and protospacer adjacent motif underlined. Allele-discriminating SNPs are shown in pink (see Supplemental Dataset S1 for the full data). **D**, Pie chart summary of the overall (left) and indel (right) editing patterns.

Figure 4. PCR analysis of selected mutant lines. **A**, PCR amplification of the six *MYB* alleles on Chr8 from two WT, one ΔG and four KO lines. The four KO lines were selected to represent one (KO-5 and KO-69) or two (KO-63 and KO-70) remaining Chr08 alleles. *UBC* (ubiquitin-conjugating enzyme) was included as loading control. M, molecular weight marker; ntc, no-template control. **B**, Mutation patterns of Chr08 alleles in the selected mutant lines as shown in Figure 3B. Orange triangles connected by dashes represent fusion junctions shown in C. **C**, Sanger sequencing of PCR products from A. Sequence alignment of the six alleles flanking the gRNA target site (red) is shown on top and chromatograms of the same region are shown below. Grey shaded alignments are introns, with allele-discriminating SNPs shown in pink and homologous intron 2 sequences in blue (shifted upstream by 21 bp in *138m1* and *186m2* due to gaps). PAM is underlined and boxed in blue for correspondence with the sequence traces below. Black triangles denote the Cas9 cleavage site and black dashed box corresponds to the 2-bp deletion (-2) detected in KO-5 and KO-69. The two fusion alleles as determined by SNPs are marked below the KO-5 and KO-69 traces (see Supplemental Figure S2 for the full sequence alignment).

Figure 5. Cuticular wax composition of trichomeless and control leaves. **A**, Total wax load. **B**, Major classes of cuticle wax. **C**, Fatty alcohols (C26, 1-hexacosanol; C28, 1-octacosanol) detected in wax (left) or whole leaves (right). **D**, β-sitosterol detected in wax (left) or whole leaves (right). **E**, Triterpenes detected in wax (top) or whole leaves (bottom). Ergosterone, 14,24-dimethyl-ergosta-8,25-dien-3-one; cycloartanone, 24-methylene cycloartan-3-one; lanosterone, lanosta-8,24-dien-3-one. Data are mean±SD of n=5. All concentration estimates were based on sorbitol equivalent. Statistical significance was determined by 2-tailed *t*-test (* *P*<0.05, ** *P*<0.01, *** *P*<0.001). nd, not detected.