Osteoblast-osteoclast co-cultures: a systematic review and map of available literature

Remmers, S. J. A[1]., de Wildt, B. W. M.[1], Vis, M. A. M[1]., Spaander, E. S. R. [1], de Vries, R.B.M.[2], Ito, K. [1], Hofmann, S. [1]*

[1] Orthopaedic Biomechanics, Department of Biomedical Engineering and the Institute of Complex Molecular Systems, Eindhoven University of Technology, Eindhoven, the Netherlands.

[2] SYRCLE, Department for Health Evidence, Radboud Institute for Health Sciences, Radboudumc, Nijmegen, the Netherlands.

*corresponding author

E-mail: s.hofmann@tue.nl

Abstract

Drug research with animal models is expensive, time-consuming and translation to clinical trials is often poor, resulting in a desire to replace, reduce, and refine the use of animal models. One approach to replace and reduce the use of animal models in research is using *in vitro* cell-culture models.

To study bone physiology, bone diseases and drugs, many studies have been published using osteoblast-osteoclast co-cultures. The use of osteoblast-osteoclast co-cultures is usually not clearly mentioned in the title and abstract, making it difficult to identify these studies without a systematic search and thorough review. As a result, researchers are all developing their own methods from the ground up, leading to conceptually similar studies with many methodological differences and, as a direct consequence, incomparable results.

The aim of this study was to systematically review existing osteoblast-osteoclast co-culture studies published up to 6 January 2020, and to give an overview of their methods, predetermined outcome measures (formation and resorption, and ALP and TRAP quantification as surrogate markers for formation and resorption, respectively), and other useful parameters for analysis. Information regarding these outcome measures was extracted and collected in a database, and each study was further evaluated on whether both the osteoblasts and osteoclasts were analyzed using relevant outcome measures. From these studies, additional details on methods, cells and culture conditions were extracted into a second database to allow searching on more characteristics.

The two databases presented in this publication provide an unprecedented amount of information on cells, culture conditions and analytical techniques for using and studying osteoblast-osteoclast co-cultures. They allow researchers to identify publications relevant to their specific needs and allow easy validation and comparison with existing literature. Finally, we provide the information and tools necessary for others to use, manipulate and expand the databases for their needs.

Introduction

Bone is a highly dynamic tissue with mechanical and metabolic functions that are maintained by the process of bone remodeling by the bone forming osteoblasts (OBs), bone resorbing osteoclasts (OCs), and regulating osteocytes. In healthy tissue, bone resorption and formation are in equilibrium, maintaining the necessary bone strength and structure to meet the needs of the body. In diseases such as osteoporosis and osteopetrosis this equilibrium is disturbed, leading to pathological changes in bone mass that adversely affect the bone's mechanical functionality (1).

Studies on bone physiology, bone disease and drug development are routinely performed in animal models, which are considered a fundamental part of preclinical research. The use of animals raises ethical concerns and is generally more time consuming and more expensive than *in vitro* research. Laboratory animals are also physiologically different from humans and their use in pre-clinical studies leads to poor translation of results to human clinical trials (2,3), and the subsequent failure of promising discoveries to enter routine clinical use (4,5). These limitations and the desire to reduce, refine and replace animal experiments gave rise to the development of *in vitro* models (6,7). Over the last four decades, significant incremental progress has been made towards developing OB-OC co-culture models.

The development of *in vitro* OB-OC co-cultures started with a publication of T.J. Chambers in 1982 (8), where the author induced quiescence of isolated tartrate resistant acid phosphatase (TRAP)-positive rat OCs with calcitonin and reversed their quiescence by co-culturing them with isolated rat OBs in direct contact. At that time, studies involving OCs resorted to the isolation of mature OCs by disaggregation from fragmented animal bones. The first account of *in vitro* osteoclastogenesis in co-culture was realized in 1988 when Takahashi and co-authors (9) cultured mouse spleen cells and isolated mouse OBs in the presence of 1α,25-dihydroxyvitamin D3 and found TRAP-positive dentine-resorbing cells. The herein described methods were used and adapted to generate OCs for the

following decade. Most of the studies published until this point in time used co-cultures as a tool for

achieving osteoclastogenesis, as opposed to a model for bone remodeling. At that time, a co-culture

of OBs with spleen cells or monocytes was the only way of generating functional OCs *in vitro*. It

wasn't until 1999 that Suda (10) discovered Receptor Activator of Nuclear Factor Kappa Ligand

(RANKL) and Macrophage Colony Stimulating Factor (M-CSF) as the necessary and sufficient proteins

required for differentiating cells from the monocyte/macrophage lineage into functioning OCs (11–

13). This discovery marked the start of co-culture models developed for studying bone remodeling.

In recent years, many research groups have ventured into the realm of OB-OC co-cultures with the

intent of studying both formation and resorption, but each group seems to be individually

developing the tools to suit their needs resulting in many functionally related experiments that are

methodologically completely different. In addition, the use of such methods is often not clearly

stated within title and abstracts. Simple title/abstract searches such as 'OB + OC +co-culture' tend to

scratch only the surface of the base of evidence available using OB-OC co-cultures. Finding and

comparing different co-culture approaches and results is thus virtually impossible and forces each

group to develop and use their own methods instead of building upon those of others.

The aim of this study was to construct a systematic review of all OB-OC co-cultures published up to

January 6, 2020. With this systematic review, we aimed at identifying all existing OB-OC co-culture

studies and analyze these within two comprehensive databases, allowing researchers to quickly

search, sort and select studies relevant for their own research. Database 1 contains all OB-OC co-

culture studies in which at least one relevant primary outcome measure was investigated (formation

and/or resorption) or secondary outcome measure (alkaline phosphatase (ALP) and/or tartrate

resistant acid phosphatase (TRAP) quantification as surrogate markers for formation and resorption,

respectively) (S1_File_Database_1). A sub-selection of studies that investigated these relevant

outcome measures on both OBs and OCs in the co-culture was included in Database 2, accompanied

by additional details on methods, culture conditions and cells (S2_File_Database_2). The collection

of the two databases will further be referred to as a systematic map.

Methods

For this systematic map a structured search protocol was developed using the SYRCLE protocol

format (14). This protocol format is tailored to the preparation, registration, and publication of

systematic reviews of preclinical studies, and helps authors predefine the methodological approach

of their review from research question to data synthesis. The protocol and search strings were made

publicly available before completion of the study selection via Zenodo (15) to ensure transparency of

the publication. In short, three online bibliographic literature sources were consulted with a

comprehensive search string and the resulting publications were combined and screened using a

four-step procedure (Fig. 1): 1) identification of OB-OC co-cultures, 2) identification of relevant

outcome measures, 3) categorization in Databases 1 and 2 (Fig 2), 4) search for additional articles in

the reference lists of studies included in Database 2 and relevant reviews.

Fig 1. Flow diagram of systematic literature search and screening. Screening step 1: Hits from 3

online bibliographic literature sources were combined, primary studies were selected, and duplicates

were removed. Title and abstracts were screened for the presence of OB-OC co-cultures. Screening

step 2: OB-OC co-cultures were screened in full text for relevant outcome measures. All studies in

which at least one relevant outcome measure was studied were included into Database 1. Screening

step 3: Papers in which both cell types were studied with relevant outcome measures were included

into Database 2. Screening step 4: Papers included into Database 2 and relevant reviews were

screened for potentially missing relevant studies and identified studies were screened in the same

manner as described here. Each screening step is marked with a separate background color. Each

selection step within the screening steps is marked with a colored header. Blue header: used as input

for the review. Grey header: selection step. Red header: excluded studies. Yellow header: Database as

presented in this systematic map. Abbreviations: outcome measures (OM), Database 2 (DB2),

osteoblast (OB), osteoclast (OC).

Fig 2. Schematic overview of Databases 1 and 2. *All identified studies were searched for OB-OC co-*

cultures, where co-culture was defined as OB and OC being present simultaneously and able to

exchange biochemical signals. In addition to direct-contact cultures, cultures such as transwell

cultures, 3D or scaffold cultures and bioreactor cultures were allowed as well. OB-OC co-culture

studies which used relevant outcome measures were included into Database 1. Of these, only the

relevant outcome measures were analyzed. All studies where relevant outcome measures were used

for both OB and OC were included into Database 2 as well. Of these, cells and culture conditions were

analyzed. The figure was modified from Servier Medical Art, licensed under a Creative Common

Attribution 3.0 Generic License (http://smart.servier.com, accessed on 2 July 2021).

Database Search

The online bibliographic literature sources Pubmed, Embase (via OvidSP) and Web of Science were

searched on January 6, 2020 with a predefined search query developed to identify as many studies

as possible employing OB-OC co-cultures. The search strings used a combination of thesaurus and

free text terms where possible and consisted of the following components: ([OBs] OR ([OB

precursors] AND [bone-related terms])) AND ([OCs] OR ([OC precursors] AND [bone-related terms]))

AND [co-culture], where each component in square brackets represents a list of related thesaurus

and free-text search terms, and where parentheses indicate the order of operations within the

search query. The full search strings can be found via Zenodo (15). The results of all three searches

were combined. Conference abstracts and duplicates were removed using the duplicate removal

tools of Endnote X7 and Rayyan web-based systematic review software (16).

The remaining entries were screened using a four-step procedure that resulted in the generation of

2 databases: Database 1 containing all co-culture studies that measured at least one relevant

outcome measure (formation, resorption, ALP or TRAP), and Database 2 containing all studies that

measured at least one relevant outcome measure on both OB and OC: either formation or ALP for

OB, and either resorption or TRAP for OC. The entire data collection and screening process was

performed independently by two researchers starting from the moment that conference abstracts

and duplicates had been removed and ending the moment that the relevant information of the

publications was extracted into the two databases. Disagreements between researchers were

discussed and publications were re-screened if necessary. A third independent researcher was

consulted for disagreements that could not be solved by the first two independent researchers.

Screening step 1: Identification of OB-OC co-cultures

This step was performed to identify and extract OB-OC co-cultures from the complete list of studies

identified from the three online bibliographic literature sources after automatic removal of

conference abstracts and duplicates. All further steps were done on these studies or a sub-selection

thereof. Using Rayyan web-based systematic review software (16), the titles and abstracts of all

entries were screened for the presence of primary studies using OB-OC co-cultures. Reviews, theses,

chapters, and conference abstracts that were not automatically detected were excluded at this

point. Potentially relevant reviews were saved separately to serve as an additional source of studies

that could have been missed by the systematic search.

In the selection process, co-culture was defined as the simultaneous presence (verified) or assumed

presence (expected) of OBs and OCs (or OB-like and/or OC-like cells) within the same culture system

at a moment during the described experiment such that the cells were able to communicate either

via soluble factors in the medium and/or direct cell-cell contact. Both primary cells and cell lines of

any origin were admitted including heterogeneous cell populations, if these were clearly defined and

expected to result in a biologically relevant number of the desired cell type, precursor type, or

terminally differentiated cell type. The presence of progenitor cells (such as monocytes or

mesenchymal stem/stromal cells) was allowed only if these were either verified or expected to

differentiate into OBs and/or OCs. Studies using a single animal or human donor for both cell types

were allowed, but only if the two (progenitor) cell types were at one point separated, counted, and

reintroduced in a controlled manner. In addition, trans-well systems (no physical contact but shared

medium compartment with or without membrane), scaffolds (3-dimensional porous structure of any

material including decellularized matrix), and bioreactor culture systems (culture exposed to physical

stimuli such as rotation, mechanical loading or fluid flow) were included. Conditioned media

experiments were excluded because these do not allow real-time two-way exchange of cell signals.

Explant cultures or organ cultures were excluded because these studies contain a living *ex vivo*

culture element, whereas the focus of this systematic map is limited to *in vitro* studies.

When the study used any type of OB-OC co-culture as defined above, the study was included. When

there was no indication that there was an OB-OC co-culture, the study was excluded. When, based

on the title and abstract, it was likely that there was a co-culture, but this was not described as such,

the full-text publication was screened.

181 Screening step 2: Identification of relevant outcome measures in the co-

182 culture experiments

This step was used to identify co-cultures that specifically investigated outcome measures related to

bone remodeling: measuring formation or resorption (primary outcome measures), or quantitative

measurements of activity markers ALP or TRAP in a dedicated assay (secondary outcome measures).

The primary outcome measures of measuring resorption and formation were chosen because these

are the processes that are directly affected in bone diseases. Measuring these outcome measures

usually requires a specific methodological setup such as a specific surface analysis for measuring

formation, or a resorbable substrate for measuring resorption. The secondary outcome measures of

ALP and TRAP were included because these are regarded as viable alternatives for the direct

measurement of formation and resorption. The full texts of the studies identified in screening step 1

were screened for experimental techniques and outcome measures. Studies in which for at least one

of the cell types a relevant outcome measure was used, were selected to be used in Database 1

(S1_File_Database_1). The measurement of formation was defined as any method that directly

measures the area or volume of (tissue) mineralization by OBs, any method that measures by-

products of formation, and any method that measures biochemical markers that directly and

exclusively correlate to formation. The measurement of resorption was defined as any method that

directly measures the surface area or volume that has been resorbed by OCs, that biochemically

measures products or by-products of resorption, or that measures biochemical markers that directly

and exclusively correlate to resorption. The measurement of ALP and TRAP was defined as the

detection of either the direct measurement of the enzymatic activity of these proteins, or the direct

quantification of the amount of those proteins present in a dedicated assay. Studies that determined

ALP or TRAP gene expression using PCR were excluded because PCR was not considered a dedicated

assay for this map and did not directly measure the amount of protein present. However, the use of

PCR was recorded in the generated databases in a separate column. Immuno-histological stainings of

ALP or TRAP were not considered relevant outcome measures, even when followed by image

analysis because at best these quantify stained surface area and not actual protein content.

All co-cultures that did not contain at least a single outcome measure that met these criteria were

excluded from further use. Because this was the first step at which the availability of the full text

publication was required, publications written in languages other than English with no translation

available, and publications of which the full text could not be found were excluded at this point.

213   Screening step 3: Categorization within Database 1

In this step, a distinction was made between studies in which only one of either OB or OC was

studied, or both were studied. This distinction was made because ideally, a model for bone

remodeling should show effects on both OBs and OCs. Each study selected from screening step 2

was assessed on the methods to study OBs and OCs. Each study was categorized into one of five

categories within Database 1: 1) The relevant outcome measures were measured in both OBs and

OCs in the co-culture. These studies were also included in the in-depth screening for Database 2

(S2_File_Database_2). 2) Both cell types were studied, but relevant outcome measures were only

measured in OCs or 3) Both cell types were studied, but relevant outcome measures were only

measured in OBs. 4) Only OCs were studied in co-culture, the other cell type was neglected or 5) OBs

were studied in co-culture, the other cell type was neglected. Thus, category 1 contained the studies

in which both formation and resorption were investigated, either directly or by ALP or TRAP

quantification. Category 2 and 3 contained studies in which both OBs and OCs were studied, but only

one of the two was studied with the relevant outcome measures. The other cell type was studied

using other methods instead such as stainings or PCR. Categories 4 and 5 contain studies in which

only one of the two cell types was analyzed with one of the relevant outcome measures while the

other cell type was present but not analyzed in co-culture at all. Note that for this categorization, it

was necessary that the cells that were used in co-culture were studied, and not for example a mono-

culture conducted in parallel.

233   Screening step 4: Review and reference list screening

To find additional studies that may have been missed during bibliographic searches, relevant review

articles identified during the selection process and studies labeled as category 1 in step 3 were

screened for additional publications that could be relevant to the current systematic map. Of these

studies, the relevant passages within the text were screened, followed by a thorough screening of

the complete reference lists of these studies. All potentially relevant studies were first cross-checked

with the original search results of the bibliographic literature search, and if these were not identified

there, were screened in the same manner as all other studies used in this systematic map. Unique

relevant studies were then added to the corresponding databases and analyzed as described earlier.

243 Database 1 generation and analysis – All co-cultures with relevant

244 outcome measures

Every study included in Database 1 was screened for the relevant outcome measures resorption,

formation, ALP and TRAP during screening steps 2 and 3. To provide useful and specific information

of each of the studies included in this database, all potentially relevant information related to the

relevant outcome measures was collected and organized. For resorption, additional information on

the resorbed substrate, the methodological procedure and quantification of results was collected.

For formation, additional information on the type of analysis, the methodological procedure and

quantification of results was collected. For both ALP and TRAP, additional information on the

mechanism of the biochemical assay, whether it was conducted on lysed cells or supernatant, and

information regarding the quantification was collected. In addition, the following information was

extracted, whether: the authors described their setup as a model specifically for remodeling, the

experiment was conducted in 3D, the experiment applied bioreactors, more than 2 cell types were

cultured simultaneously, the culture used a trans-well setup, the culture used PCR and components

in the supernatant of the culture were analyzed by ELISA or a similar quantification method. If the

answer to these questions was yes, then the applicable details were collected as well. Finally, a

column for additional remarks was introduced for details that did not fit in another column. Studies

where the authors are color coded in pink were those not found through the initial database search

but by the screening of the review articles or reference lists. Studies categorized as category 1 in

screening step 3 were selected for use in Database 2 and had their title color coded in orange.

Quality assessment and scripting

In Database 1 only the methods used for analyzing relevant outcome measured are reported, and

not the data obtained from them or the results described in the publication. Quality assessment in

Database 1 is thus limited to assessing the completeness of the necessary elements of the collected

methodological details, to the extent that the methods are properly represented in Database 1 and

related tables. Please note that the methods themselves were not investigated on a complete

description for a perfect replication of the study, but only on the description necessary to accurately

classify the method within this systematic map. For example, a study claiming to investigate

resorption on dentine discs using Toluidine Blue was deemed sufficiently described to accurately

classify, regardless of whether the information presented was sufficient to duplicate that specific

method precisely. Publications in which information was missing are here represented as 'not

reported' if no information was provided, 'reference only' if no information was provided but

another study was referenced, and 'undefined kit', when a commercial kit was used but the content

or methodology was not further described. Instances of missing information can easily be identified

in figures, tables and databases, but were not further used in this systematic map. Studies where an

instance of information was missing were still used for other analyses for which the corresponding

provided information was present.

A script was written in Excel Visual Basics programming language to analyze Database 1 and extract

relevant statistical information on the collected information. On sheet 2 "Data" of the Database 1

excel file, the descriptive statistical data and collected information are presented in the form of lists

and tables and together with a button to re-run the analysis based on the reader's requirements.

The script is integrated within the excel file and can be used only when the file is saved as a 'macro-

enabled' file (.xlsm).

Database 2 generation and analysis – All co-cultures in which both cell types had relevant outcome measures.

In addition to the information already collected for Database 1, additional information was extracted

from the studies in which relevant outcome measures were studied of both OB and OC: the species

(17) and type (cell line or primary) and actual used cell type (6) of both the OB and OC were

collected. Seeding numbers and densities (18) for both OB and OC were collected or calculated

where possible, separated by 2D (cell density per area) and 3D (cell density per volume), and the

seeding ratio (19) between OB and OC was noted or calculated. The culture surface (bio-)material

(20), sample size (samples per group), culture duration and medium refreshing rate in units as

reported in the study, environmental conditions or variations such as CO_2 or O_2 alterations or

mechanical loading (21), and pre-culture duration (22) were collected, where pre-culture is defined

as a different co-culture condition (such as a different supplement cocktail) lasting for a short

duration (such as 2 days) prior to the 'main' co-culture. The medium composition (23) was collected

and organized by base medium type such as Dulbecco's Modified Eagle Medium (DMEM) and alpha-

Modified Eagle Medium (αMEM), glucose content (if provided separately in the text), Fetal Bovine

Serum (FBS) / Fetal Calf Serum (FCS) in percentages, antibiotics (types and concentrations or

percentages as provided), OB supplement concentrations (ascorbic acid, β-glycerophosphate and

dexamethasone, OC supplement concentrations (M-CSF and RANKL) and other supplements, as well

as medium content of any monoculture prior to the co-culture. Finally, the tested genes of all studies

applying PCR and any proteins studied with ELISA or other supernatant analyses executed on the co-

culture were noted.

Quality assessment and scripting

In Database 2 the culture conditions, cells and materials used are reported, and not the data obtained from them or the results described in the publication. Quality assessment in Database 2 is thus limited to assessing the completeness of the necessary elements of the collected methodological details, to the extent that the methods are properly represented in Database 2 and related figures and tables. Please note that the methods themselves were not investigated on a complete description for a perfect replication of the study, but only on the description necessary to accurately classify the method within this systematic map. For example, a study claiming to use human primary monocytes and human primary osteoblasts for the OB-OC co-culture was deemed sufficiently described to accurately classify respectively the OB and OC origin, regardless of whether the information presented was sufficient to perfectly replicate that part of the study. Publications in which information was missing are here represented as 'not reported' (NR) if no information was provided, or 'reference only' if no information was provided but another study was referenced. If studies were missing information critical to reproduce the outcome measures (for example seeding ratio's, culture surface material, medium or supplement information, critical steps in analyses), the cells in the database missing this information were labeled in red. If the missing information was not critical for the outcome measures but necessary for a replication of the study (for example sample size, medium refresh rate, control conditions), the cells were labeled in orange. The sum of both orange and red cells for each color in each study is shown as well to indicate how many instances of missing information were identified in each study. The color coding was determined by the authors of this map but can be adjusted within Database 2 if other criteria for critical information and completeness are desired. Instances of missing information can easily be identified in the corresponding figures, tables and databases, but were not further used in this systematic map. Studies where an instance of information was missing were still used for other analyses for which the corresponding provided information was present.

Using Excel Visual Basics programming language, three scripts were written to analyze and process

Database 2. One script was created to count all instances of cells labeled as 'missing info' and

present this number in two dedicated columns (missing critical or non-critical info). One script was

created to count the frequency of occurrence of all (co-)authors and years of publication. Finally,

one script was created to analyze this database and extract relevant descriptive statistical data on

the collected information. On sheet 2 "Data" of the Database 2 excel file, the statistical data and

collected information are presented in the form of lists and tables and together with the buttons to

re-run the analyses based on the reader's requirements. The scripts are integrated within the excel

file and can be used only when the file is saved as a 'macro-enabled' file (.xlsm).

Results

Search results

From three online bibliographic literature sources, 7687 studies were identified (Pubmed: 1964,

Embase via OvidSP: 2709, Web of Science: 3014). After removal of conference abstracts, 6874

studies remained. After duplicate removal, 3925 unique studies were identified to be screened.

Studies included into Database 1

After title-abstract screening and when in doubt full text screening (screening step 1), 694 studies

were identified as OB-OC co-cultures. A list of these studies is available as a supplementary file

(S4_File_List of all OB-OC co-cultures). Of these, one study was excluded from further analysis

because the full text could not be obtained, 35 were excluded because they were in a language

other than English and 406 were excluded because no relevant outcome measure was used in the

study (screening step 2). The qualifying 252 studies with at least one relevant outcome measure

were included in Database 1.

Studies included into Database 2

358 In 77 of these studies, both the OB and OC were studied, and in 39 of these, both OB and OC were

359 studied using relevant outcome measures (screening step 3). These 39 studies were included in

360 Database 2.

Additional screening of review articles and reference lists for missing studies

The 39 studies of Database 2 and 10 additional review publications were screened for other relevant

studies that the initial search may have missed (screening step 4). An additional 25 unique studies

were identified in the 10 reviews, and 34 unique studies were identified from the reference lists of

the included studies. These additional 59 studies were reviewed as described previously and

resulted in an additional 3 OB-OC co-cultures with only relevant outcome measures measured on

one cell type, resulting in a total of 255 studies with relevant outcome measures on at least one cell

type for Database 1, and 39 studies in which relevant outcome measures were studied in both cell

types for Database 2. A detailed overview of the search and selection process is shown in Fig 1.

Publications per year

The publications included in Database 1 were published between 1983 and 2019, with a peak in

publications around the year 2000, followed by a dip and then a more or less slight but steady

increase until now (Fig 3a). The peak roughly coincides with the discovery that M-CSF and RANKL

were both necessary and sufficient to induce osteoclastic differentiation in monocytes in 1999 (10).

The included publications in Database 2 span the time between 1997 and 2019, with only 8

publications before 2010 (Fig 3b). This coincides with the progress in development of *in vitro* co-

cultures of OB and OC, moving beyond co-cultures with OB to generate OC, and moving towards co-

cultures of OB and OC to study for example cell-cell interactions (6).

Fig 3. Relevant publications per year. *A) All 255 publications that contain relevant outcome*

measures counted by year ranging from 1983 to 2019 (Database 1). B) The 39 selected publications

of Database 2 counted by year ranging from 1998 to 2019 (Database 2).

Database 1 results

Database 1 provides an overview of all OB-OC co-culture studies published until January 6, 2020 in

which at least one relevant outcome measure was studied. Of the 255 studies included, resorption

was analyzed in 181 studies, formation was analyzed in 37 studies and both were analyzed in 16

studies. ALP was analyzed in 42 studies, TRAP was analyzed in 61 studies and both were analyzed in

22 studies (Table 1).

Table 1. Combinations and frequencies of primary and secondary outcome measures.

Combinations of primary and secondary outcome measures in each study		Primary outcome measures				
		No resorption or formation	Resorption only	Formation only	Resorption and formation	Total
Secondary	No ALP or TRAP	0	151	14	9	174
	ALP only	16	0	2	2	20

		23	9	3	4	39
	TRAP only	23	9	3	4	39
	ALP + TRAP	14	5	2	1	22
	Total	53	165	21	16	255

Table 1: *This table can be referenced to identify the number of studies using any combination of*

primary and secondary outcome measures. All 255 studies that investigate at least one of the

primary or secondary outcome measures are represented once in this table. Each study is

represented by a combination of primary outcome measures (horizontal) and secondary outcome

measures (vertical). Marginal totals of each row and column are counted under 'total' with the grand

total in the bottom-right cell. These marginal totals sum the total number of studies that studied only

that combination of either primary or secondary outcome measures, with no regard of the outcome

measures on the opposite axis. The total numbers of each individual outcome measure can be

calculated from this table but are presented in the following paragraphs and tables.

Resorption

Resorption is the process by which osteoclasts remove old and damaged bone tissue through

enzymatic degradation or acidic dissolution. Out of all 255 OB-OC co-culture publications included in

Database 1, resorption was studied directly on 188 occasions in 181 studies and quantified 142 times

(Table 2). In some publications, more than one material or method of analysis for resorption was

used. In cases where multiple materials were used, each material was counted as an individual study

of resorption and for each material, the corresponding analyses were counted, even if these were

identical per material. In those cases where multiple methods of analysis are used on the same

material, all methods are counted, and the material is counted only once. This resulted in a counted

number of studies that is higher than the actual number of publications. When numbers of studies

are referenced, these are the 'counted' number of studies defined above, and not the actual number

of publications.

Table 2a. Occurrences of resorption on different types of substrates and subsequent analyses.

		Materials used as a resorbable substrate for measuring resorption										
Shapes, structures and types of materials used as resorbable substrate for analysis of resorption. Per-material total number of studies		Dentine	Bone	HA	Silk	Collagen	CaP	PLLA	Chitosan	Osteologic	Mineralized	Not reported	Per-row total
		76	66	6	5	2	4	1	1	19	6	2	188
Per material quantified studies		55	52	3	4	1	4	0	0	17	4	2	142
Shape or structure of material	Discs	76	63	2			2			13			156
	Films			2	4			1	1				8
	Coatings			2			1						3
	Scaffolds				1	1					3		5
	Hydrogels					1							1
	ECM										2		2
	Nodule										1		1
	Fragments		3										3
	Substrates						1						1
	Plates									6			6
	Not reported											2	2
Analysis techniques for analzying resorption on resorbable substrates.		Dentine	Bone	HA	Silk	Collagen	CaP	PLLA	Chitosan	Osteologic	Mineralized	Not reported	Per-row total
Staining	Toluidine Blue	36	19										55
	Haematoxylin	16	2										18
	Eosin		1										1
	H&E		1										1
	Alum / Coomassie Blue		1										1
	TRAP	1											1
	Von Kossa						2			4	1		7
Microscopy	Phase contrast						1			4			5
	SEM	12	37	5	3		1			1			59
	TEM					1					1		2
	2-Photon										1		1
	Atomic force				1			1	1				3
	Reflected light	8	2										10
	Dark field										1		1

	Light microscopy									6			6
Other	Assay					1							1
	Immuno-assay		3								3	1	7
	MicroCT				1					1			2
	Reference only	2											2
	Not reported	1		1						2		1	5
	Total per material	76	66	6	5	2	4	1	1	19	6	2	188

Table 2a: Each column signifies a different material used as a substrate for measuring resorption. If other cells, prior to the introduction of OC, were used to deposit mineralized matrix on another material, then the material was listed in the column 'Mineralized'. If the material was not reported, the study was listed in the column 'Not reported'. The first rows show how many instances of each material were included into this systematic map in total, and how many times the results were quantified. The final column shows incremental totals per material type or analysis type. This table consists of two sections. The top section shows in what form or shape the corresponding materials were used as a substrate for resorption. The bottom section shows the techniques that were used to study the resorption described on the materials described in the top section. Each individual study is represented exactly once in the top section of the table to signify the type and form of the substrate used, and exactly once in the bottom section of the table to signify the method used to analyze the resorption that occurred on that substrate. This required the selection of the most 'important' part of the methods used. In the cases where first a staining was used followed by microscopy, only the staining is listed. Only in those cases where resorption was investigated directly with a microscope without prior staining, the type of microscopy is listed.

Table 2b Supernatant resorption techniques

Supernatant Analysis techniques per material used for analysis of resorption.		dentine	bone	HA	silk	collagen	CaP	PLLA	chitosan	osteologic	mineralized	not reported	Per-row total
Supernatant analysis	NTx	1									2		3
	CTx										1	1	2
	ICTP		1										1
	Phosphate release					1					2		3
	Radioactive proline release		2										2

433 **Table 2b:** *This table presents five types of resorption analysis where measurements can be performed in the culture supernatant and not on the material itself. In the corresponding studies, these were done in addition to 'regular' analysis presented in Table 2a, and for that reason are presented separately. These have the advantage that they can be used to monitor changes over time in a non-destructive way.*

438 Most of these studies used discs or fragments of either bone or dentine. Due to the flat nature of these discs, the surface can be considered 2D, and resorption pits can be visualized directly using conventional microscopy techniques, such as for example Scanning Electron Microscopy (SEM) or Reflected Light Microscopy (RLM). To enhance the contrast of the resorption pits, stains such as Toluidine Blue (TB) and Hematoxylin (H) were used. Resorption on bone fragments was quantified using radioimmunological assays measuring the release of *in vivo* pre-labeled [3]H-proline or type I collagen telopeptide.

445 Synthetic resorbable discs or coatings on culture plates are designed specifically for studying resorption, and usually the exact composition has not been revealed. These will further be referred to as 'osteologic' plates or discs. The discs were analyzed in roughly the same way as bone or dentine discs. Thin resorbable coatings on translucent culture plates offer another interesting approach. Resorbed areas reveal the translucent culture plate, while unresorbed areas are less translucent and can be stained with for example von Kossa's method to provide even more contrast, making quantification with image analysis easy.

Hydroxyapatite (HA) and other calcium phosphates were used in the form of discs, films, coatings, or

scaffolds and were analyzed using various types of microscopy, both with and without prior staining.

These can be used in a similar way as biological and synthetic materials mentioned earlier, with the

main advantage being their known composition.

Resorption of ECM or nodules produced by OBs and scaffolds mineralized by OBs were investigated

with transmission electron microscopy, light microscopy after staining, using 2-photon Second

Harmonic Generation microscopy (24), supernatant phosphate levels, or with an ELISA for C-terminal

telopeptide (CTx) or N-terminal telopeptide (NTx), which are bone turnover marker more commonly

used for testing urine and serum samples.

Formation

Formation is the process by which osteoblasts create new bone tissue through the mineralization of

deposited collagenous extracellular matrix. Out of all OB-OC co-cultures included in Database 1,

formation was studied directly 39 times in 37 studies and quantified 29 times. (Table 3) In some

studies, more than one method of measuring, analyzing and quantifying formation was used. In

those cases, all methods are counted as individual studies. The methods of formation analysis were

divided into 5 types: nodule analysis, volume analysis, surface analysis, supernatant analysis and 3D

scans.

Table 3: Formation statistics and analyses.

		Type of analysis used to measure formation					
	Technique	Scan	Nodule analysis	Supernatant analysis	Surface analysis	Volume analysis	Per-row Total

Measured shape or structure / Staining / Assays / Other	Analysis	Scan	Nodule analysis	Supernatant	Surface analysis	Volume analysis	Per-row Total
	Total	3	20	6	5	5	39
	Quantified	3	12	6	3	5	29
Measured shape or structure	Scaffold	2	1	1	3	2	9
	Film	1		1	1	2	5
	Hydrogel					1	1
	Pellet		1			1	2
	Dye release		5				5
Staining	H&E				1		1
	Von Kossa		2		1		3
	Alizarin Red		16				16
	Lentiviral fluorescence		1				1
Assays	Calcium					3	3
	Calcium + Phosphate					2	2
	CICP			6			6
Other	SEM		1		3		4
	microCT	3					3
Per-analysis Total		3	20	6	5	5	39

Table 3: *Each column signifies a different type of analysis used for measuring formation. These include any type of non-destructive scan, a form of analysis of mineralized nodules, supernatant analysis, surface analysis, and destructive analysis of a mineralized volume. The first rows show how many instances of each type of analysis were included into this systematic map in total, and how*

many times the results were quantified. The final column shows marginal totals per row of each row.

This table consists of two distinct sections, each starting with a row showing all analysis types for

convenience. The first section lists defining characteristics of studies such as using films, scaffolds,

hydrogels or pellets, or using a technique to first stain tissue, and then releasing and measuring the

released dye. Not each study had such defining characteristics, and the total of section one does not

add up to 39 studies. Section two shows either which materials was measured, or which technique

was used for measuring formation. Each instance of formation is represented in section two of this

table exactly once. Stainings were followed by microscopy or an assay in those cases where dye was

released to be measured.

The most common method to quantify formation was to investigate mineralized nodule formation.

This was done by using staining techniques such as Alizarin Red (calcium) (25) or von Kossa

(phosphate) (26) followed by imaging, or directly imaging the nodules. While any staining specific for

mineralized matrix or even plain light microscopy images could be quantified using appropriate

imaging techniques and software, Alizarin Red offers an additional way of quantification: the amount

of dye binding to the mineral correlates to the amount of mineralization, after imaging the dye can

be released from the minerals using acetic acid and can then be quantified using colorimetric

spectrophotometry (27). Surface analysis was used in a similar way to study formation on scaffolds,

films, or particles. Scaffolds were stained and/or imaged, and the area of matrix deposition was

visualized or quantified. Volume analysis was used to describe the measurement of mineralized

tissue components calcium and phosphate, which were released after destruction of the matrix.

These three types of formation measurement are destructive methods, meaning that the samples

must be sacrificed for each time point.

The remaining two types of formation methods are non-destructive. Supernatant analysis was used

to describe the measurement of Collagen type I C-terminal propeptide (CICP), a byproduct of

collagen deposition, in cell culture supernatant. 3D scan was used to describe the use of (in this

case) µCT quantify the three-dimensional structure of mineralized matrix.

TRAP measurements as a surrogate marker of osteoclastic resorption

TRAP is a protein that has long been used as the predominant OC marker (28). Out of all OB-OC

studies in Database 1, TRAP was studied 63 times in 61 publications by a dedicated assay (Table 4).

TRAP can be measured intracellularly or excreted into the medium in two ways. Its enzymatic

phosphatase activity can be measured directly, or the amount of TRAP molecules present can be

quantified. TRAP release was studied both on cell lysate and on supernatant, and in some cases on

both. The most frequently used method to study TRAP activity was using 4-nithophenylphosphate

(pNPP), a substrate that is cleaved by phosphatases into phosphate and detectable yellow 4-

nithophenol. Others used the fluorophore Naphthol ASBI-phosphate, which is converted into the

fluorescent Naphthol-ASBI (29) and shows specificity for TRAP isoform 5b, making this method more

specific for the detection of OC when compared to the measurement of TRAP enzyme in general

(30). Naphtol ASMX phosphate (31) and an otherwise undisclosed diazonium salt function in a

similar manner. Enzyme linked Immunosorbent Assay (ELISA) can be used to detect TRAP in a slightly

different manner; by binding a detectable substrate directly to the TRAP enzyme instead of using the

enzyme to produce detectable substrate. Relying on conjugated enzymes or fluorescence, these

techniques should be more effective at low concentrations of TRAP because multiple conjugates

could bind to a single TRAP molecule. Others used a kit to detect TRAP, but no description of the

assay other than the manufacturer were given.

Table 4: TRAP measurement techniques and analyses.

Type	pNPP	N-ASBI-P	N-ASMX-P	ELISA	Diazonium salt	Undefined kit	Reference	Not reported	Total
Total	33	5	1	9	1	9	4	1	63
Lysed cells	29	5	1	1		3	2		41
Supernatant	6			7	1	6	2		22
Reference only				1					1
Not reported						1		1	2
Analysis	pNPP	N-ASBI-P	N-ASMX-P	ELISA	Diazonium salt	Kit	Reference	Not reported	Total
absorbance	33		1	8		6	2	1	51
Fluorescence		5							5
Radiography									0
Reference only	2			1	1		2		6
Not reported						4			4

Table 4: *Each column in Table 4 signifies a different technique to measure TRAP. This table consists of*

two distinct sections. The first section shows the number of studies that used each technique, and

whether these were used on (lysed) cells or on culture supernatant. If only a reference to other

literature was provided, that instance was listed in the row 'Reference only', and when these details

were not reported, that instance was listed in the row 'Not reported'. Note that in a single study

TRAP can be measured with the same technique on both cell lysate and culture supernatant,

resulting in a higher count of occurrences than number of studies that analyzed TRAP. The second

section shows with which method of analysis the TRAP content was measured. If only a reference to

other literature was provided, that instance was listed in the row 'Reference only', and when these

details were not reported, that instance was listed in the row 'Not reported'. If one study measured

TRAP on both cells and supernatant, then that study is represented twice in the second section. In all

other cases, each study is represented once in each section.

ALP measurements as a surrogate marker of osteoblastic tissue formation

Alkaline phosphatase (ALP), a bone turnover marker that is commonly used to investigate OBs, was

studied in 42 publications (Table 5). ALP is a phosphatase, that like TRAP in OCs can be found both

within and on the OBs surface and can be excreted into culture medium soluble or via extracellular

vesicles (32,33). The most frequently used method to measure ALP was to use the substrate pNPP,

which is cleaved by ALP into phosphate and detectable yellow 4-nitrophenol. Enzyme Immuno

Assays (EIA) and ELISAs are similar immunoenzymatic assays (34) that rely on an labelling ALP

molecules with a detectable substrate or other enzymes. This is in contrast with the pNPP-based

methods, where the ALP enzyme itself through its inherent enzymatic activity is responsible for

generating the colored substance. An advantage of the EIA and ELISA methods is that these are

generally more sensitive; multiple detectable molecules or enzymes can be bound to each ALP

molecule. Others used a kit to measure ALP, but no description of the assay other than the

manufacturer were given.

Table 5: ALP measurement techniques and analysis.

		ALP measurement techniques				
	Type	pNPP	EIA	ELISA	Undefined kit	Total
Substrate	Total	26	8	1	7	42
	Lysed cells	19	1		6	26
	supernatant	8	7	1	2	18
Detection	absorbance	25	8	1	3	37
	Reference only	2				2
	Not reported				5	5

Table 5: Each column signifies a different technique to measure ALP. The first rows show the

occurrence of each technique and whether these were used on (lysed) cells, or on culture

supernatant. Note that in a single study ALP can be measured with the same technique on both cell

lysate and culture supernatant, resulting in a higher count of occurrences than number of studies

that analyzed ALP. The final three rows show with which method of analysis the ALP content was

measured.

560   Database 2 results

While Database 1 was created to provide an overview of all reported methods to study the relevant

outcome measures (resorption, formation, TRAP and ALP) without other experimental details,

Database 2 was created to provide more insight into what culture conditions were used for co-

cultures. From Database 1, studies that investigated relevant outcome measures on both OB and OC

were regarded as co-cultures capable of showing OB-OC interaction, versus using one cell type only

to stimulate an effect or differentiation in the other. Of these qualifying studies, more information

on the used cells and culture conditions was extracted and analyzed in Database 2.

569   Osteoblasts

Osteoblasts are the bone forming cells responsible for depositing mineralized matrix. From all 39

studies included in Database 2, the cell types that were present at the start of the co-culture were

recorded and are shown in Table 6. More than half used human primary cells, whereas the others

used animal primary cells or any type of cell line. Whether OBs or their progenitor cells were applied

differed greatly between studies: almost half of the studies started the co-culture with OBs, the

others started the co-culture with a type of progenitor cell. It needs to be noted that some of cell

descriptions in Table 6 might refer to identical cell populations. This is a result of ambiguous

isolation methods and nomenclature which is subjective and can evolve over time (35). This

systematic map reflects the nomenclature used by the authors or an unambiguous translation of the

provided nomenclature to nomenclature used in this map and does not interpret the provided

information if it was ambiguous.

One interesting observation regarding the cells used as OBs is that there is little variation in the

different types of cells introduced into the co-cultures. Except for the oldest 6 studies that used

chicken and rat cells, all studies used human or mouse cells, most of which were primary cells. While

the studies using rat and mouse cells mostly directly introduced OBs (either isolated as such or

differentiated before seeding), those that used human cells predominantly resorted to using

progenitor cells (35). Such OB precursors can be obtained from blood and bone marrow donations

and can be expanded to the required number of cells *in vitro*. The main difference between OB

versus progenitors is the presence or absence of the osteogenic differentiation phase.

Differentiation within the experiment could be desired for the research question or must be

considered in case it is not. Those that used primary OBs purchased expandable human OBs (36) or

used OBs (37), undefined expanded bone cells (38), or differentiated MSCs (39) from bone material

obtained during a surgical procedure.

Table 6: Osteoblast origins and occurrences.

Cell Origin	Osteoblasts	Mesenchymal stem cells	Mesenchymal stromal cells	Stromal cells	Stromal vascular Fraction	Osteoprogenitor cells	Per-row Total
Human primary	4	9	2	6	1		22
Human cell line	1						1
Mouse primary	3	2					5
Mouse cell line	4						4
Rat primary	3					1	4
Chicken primary				2			2

Reference only	1						1
Total	16	11	2	8	1	1	39

Table 6: *From Database 2, the origin of the cells that were used as OB was extracted. Each column*

represents a different cell type of OB-like cells or their precursors. Each row represents a different

source of cells, differentiating between both the origin species and whether the cells are primary cells

or cell lines. Incremental totals are presented in the last row and column.

Seeding densities plays a major role in proliferation and cell function of OBs (18,83). Seeding density

of OBs in 2D could be extracted or calculated for 26 studies and ranged from approximately 900

cells/cm^2 to approximately 60.000 cells/cm^2 with a median of approximately 6500 cells/cm^2 and a

mean of approximately 11000 cells/cm^2 (Fig 4a). Seeding density of OBs in 3D could be extracted or

calculated for 6 studies and ranged from approximately 300 cells/cm^3 to approximately 7*10^7

cells/cm^3 with a median of approximately 4*10^6 cells/cm^3 and a mean of 15*10^6 cells/cm^3 (Fig 4d). It

is important to note that these numbers are taken from the entire base of studies in Database 2, and

as such are not representative for any type of OB or precursor used. These numbers can be further

sorted and selected based on the individual researchers' needs.

Fig 4. Seeding densities and seeding ratios. *Violin plots of 2D and 3D seeding ratios of OB (A+D), OC*

(B+E) and respective seeding ratios (C+F). Values are calculated based on reported seeding numbers

of the cells or precursors thereof by authors per surface are or volume. No distinction was made

between different types of cells or precursors in these figures and this introduces a considerable

spread in data due to possible cell proliferation (OB) and cell fusion (OC) that might have occurred

after seeding. This distinction can be made in the database itself. Please take note that the ranges

along the Y-axis are not the same for each figure. Each seeding density of each study is represented

by a blue dot.

Osteoclasts

Osteoclasts are the bone resorbing cells that remove old and damaged bone tissue to make place for

the deposition of new mineralized matrix. Out of all 39 studies included in Database 2, 20 used

human primary cells, the others used animal primary cells or any type of cell line (Table 7). In most

cases cultures were initiated with OC progenitors: 16 studies introduced monocytes, 11 introduced

mononuclear cells, the rest used other precursors. Again, it needs to be noted that some of these

descriptions are ambiguous. What is reported here is the definition used by the authors of the

respective studies.

The origin of the cells used as OCs is remarkably like those of the OBs. The 6 oldest included studies

used chicken and rat cells, and all others used mouse or human cells. With only one exception

combining a mouse ST-2 cell line with human monocytes (40), all studies used cells of exclusively a

single species for the OB and OC source. Such a similarity was not found regarding the use of cell

lines versus primary cells. While many studies introduced OBs directly into the co-culture, only a

single study claimed to introduce OCs directly into co-culture but failed to provide any information

regarding either cell source and was therefore ignored from further use.

Seeding density of OC in 2D could be extracted or calculated for 25 studies and ranged from $5*10^3$

cells/cm^2 to $15*10^6$ cells/cm^2 with a median of $42*10^3$ cells/cm^2 and a mean of $190*10^3$ cells/cm^2

(Fig 4b). Seeding density of OC in 3D could be extracted or calculated for 6 studies and ranged from

$2*10^4$ cells/cm^3 to $7*10^7$ cells/cm^3 with a median of $4*10^6$ cells/cm^3 and a mean of $17*10^6$ cells/cm^3

(Fig 4e). Seeding ratios of OB:OC in 2D varied highly and ranged from 1:1500 to 1:1 (Fig 4c) and

seeding ratios of OB:OC in 3D ranged from 100:1 to 1:25 (Fig 4f). In human bone tissue, the ratio of

OB:OC is estimated to be approximately 7:1 (41). It must be noted that in these numbers, no

distinction has been made between the use of precursors versus OB or OC or any type of expansion

phase within experiments. These distinctions can be made within Database 2 for each individual

need.

Table 7: Osteoclast origins and occurrences.

Cell Origin	Monocytes	Mononuclear cells	Macrophages	Osteoclast precursors	Osteoclasts	Spleen cells	Total
Human primary	10	6	1	3			20
Human cell line	4						4
Mouse primary	2		2	2			6
Mouse cell line			2				2
Rat primary		3				1	4
Chicken primary		2					2
Reference only					1		1
Total	16	11	5	5	1	1	39

647 ***Table 7:*** *From Database 2, the origin of the cells that were used as OC was extracted. Each column*

648 *represents a different cell type of OC-like cells or their precursors. Each row represents a different*

649 *source of cells, differentiating between both the origin species and whether the cells are primary cells*

650 *or cell lines. If the cell source was indicated using only a reference, that instance was listed in the row*

651 *'reference only'. Incremental totals are presented in the last row and column.*

Co-culture medium composition and culture conditions

The behavior of cells is highly dependent on their environment, of which the biochemical part is predominantly determined by the culture medium composition. The main components of typical culture media are a base medium, fetal bovine serum (FBS) and specific supplements such as growth factors, especially when progenitor cells need to be differentiated first. Within the scope of this study, the base medium, FBS content and concentration of typical OB and OC supplements were analyzed. It became obvious that culture conditions are manifold and differ much between studies: A total of 8 different base (or complete) media were reported (Fig 5a), with αMEM and DMEM accounting for approximately 80% of all studies. FBS content ranged from 0% to 20%, with most studies using 10% (Fig 5b). Those without supplemented FBS used forms of complete media of which the composition was not described, but possibly including a type of serum or equivalent serum-free supplements.

Fig 5. Medium components used by studies in Database 2. A) The occurrence of all identified base and complete media used during the co-culture phase of each study. B) Serum concentrations during the co-culture phase of each study. Numbers report exclusively the use of separately introduced FBS or FCS. Serum as part of a complete medium kit that was not described in the methods section is not reported here. C) OC supplements administered during the co-culture phase of each study. OC supplements were exclusively reported in ng/ml and are reported as such in the violin plot with all individual concentrations as blue dots. Please note that the x-axis has a linear distribution. D) Osteogenic supplements during the co-culture phase of each study. Osteogenic supplements were recalculated to molarity where necessary for comparability. Individual molarities are shown as blue dots. Please note that the x-axis has a logarithmic scale.

OC supplements were reported exclusively in ng/ml (Fig 5c). M-CSF concentration was reported in 11

studies and ranged from 10 ng/ml to 100 ng/ml with a mean of 39,82 ng/ml. RANKL concentration

was reported in 14 studies and ranged from 10 ng/ml to 100 ng/ml with a mean of 49 ng/ml. All OB

supplements were recalculated to molarity if they were reported in concentrations (Fig 5d). Ascorbic

Acid (AA), which was also referred to as ascorbic acid-2-phosphate, L-ascorbic acid or L-ascorbate-2-

phosphate, was used in 19 studies. AA concentration ranged from 0.05 mM to 0.57 mM, with mean

of 0.18 mM and one outlier at 200 mM that was disregarded for this calculation. Dexamethasone

was used and reported in 13 studies and was used in 2 different molarities: 6 times at 10^{-7} M and 7

times at 10^{-8} M. β-Glycerophosphate (βGP) use was reported in 17 studies, and ranged from 1 mM to

46 mM, with a mean of 13 mM.

Other culture conditions and techniques

In addition to the cell and medium characteristics, there are other factors that define an experiment.

Out of the 39 studies of Database 2, 9 studies used a type of transwell or well insert culture (where

cell populations are separated, but factor exchange is possible), 16 studies used a form of 3D culture,

3 studies reported the use of bioreactors, 2 studies used more than the required 2 cell types to form

a tri- or tetra-culture (39,42), 7 studies reported using non-standard environmental conditions such

as gas concentrations or mechanical loading. Polymerase chain reaction (PCR) was used in 13

studies, and supernatant analyses such as ELISA were used in 14 studies. The target genes, proteins

or compounds were extracted from the publications and the occurrence of each target was recorded

in the analysis of Database 2.

Discussion

In recent years, many research groups have ventured into the realm of OB-OC co-cultures with the intent of studying both formation and resorption. Due to a lack of standardization within the field and the difficulty of finding publications based on methods instead of results, each group seems to be individually developing the tools to suit their needs resulting in many functionally related experiments that are methodologically completely different. The use of OB-OC co-cultures is usually not clearly mentioned in the title and abstract, making it difficult to find these studies without a systematic search and thorough review. The aim of this study was to generate a systematic map to give an overview of existing osteoblast-osteoclast co-culture studies published up to 6 January 2020, and present their methods, predetermined outcome measures and other useful parameters for analysis in 2 databases which can be filtered, sorted, searched and expanded.

The Database 1 contains all OB-OC co-culture studies in which at least one relevant primary outcome measure (formation and/or resorption) or secondary outcome measure (ALP and/or TRAP quantification as surrogate markers for formation and resorption, respectively) was investigated (S1_File_Database_1). A sub-selection of studies that have relevant outcome measures investigated on both OBs and OCs in the co-culture are shown in Database 2, accompanied by additional details on methods, culture conditions and cells (S2_File_Database_2).

Resorption

Most studies in Database 1 investigating resorption did so in 2D cultures using a resorbable substrate such as bone, dentine, or synthetic osteological discs. This is not unexpected, as these three options are either the actual *in vivo* material (bone), a similar material with excellent properties for studying resorption (dentine) (43), or a material designed specifically for the purpose

of studying resorption (osteologic discs or coated wells). One crucial advantage of using dentine

discs over bone is related to the native structure of dentine itself: it does not contain canaliculi and

has fewer other irregularities because it is not actively remodeled, providing more contrast between

the native structure and resorption pits to accurately visualize them (43,44). Because of that reason,

dentine is often favored over bone. The advantages of bone over dentine are that bone is the actual

tissue of interest as opposed to a bone-like material, it can be obtained from many different species

in relevant quantities and sizes, can be more easily be prelabeled *in vivo* with for example

radioactive markers such as 3H-proline (45), it is cheaper and more readily available, and could be

used in conjunction with cells from the same species or even same animal, although the latter was

not observed in this map. Dentine is a component of ivory, usually obtained from elephants (46),

hippo's (47) or sperm whales (48). Regulations regarding ivory are strict and the material is rare,

making it difficult and expensive to import and obtain. Synthetic osteologic discs have the advantage

of being produced in a uniform manner and should show little sample-to-sample variation compared

to discs made from animal tissue, or hand-made discs. Using well plates with thin osteologic coatings

has the advantage that once the coating is resorbed, the translucent well below is revealed, which

facilitates imaging with light microscopes. Combined with certain stainings, it makes quantifying

resorbed area using conventional light microscopy easier.

Choosing the surface that will be resorbed by the osteoclasts will result in a compromise. For

example, HA and other calcium phosphates are a likely choice for studying resorption since they are

a major constituent of bone. While not optimized to facilitate resorption per se, they are simple to

create, have a known composition and should offer good between-lab reproducibility. This contrasts

with resorbable discs and plates with undisclosed ingredients and likely between-manufacturer

variation. They are however synthetic, and do not contain any organic ECM components, which

means that techniques such as measuring bone turnover markers NTx (49) and CTx (50) do not work.

It is believed that the deposition of collagen type I by osteoblasts is a vital step in the formation of

mineralized tissue (51), and similarly could play a role in the resorption thereof. It is also possible to

generate the to-be-resorbed material *in vitro* by the OBs (50), even within the same experiment. This

essentially simulates a bone remodeling environment that is a step closer to the physiological

process of bone remodeling versus only resorption, although *in vivo* the order in which this typically

happens is reversed: first, damaged ECM is resorbed by OC, then new ECM is deposited by OB (52).

However, the process of creating a mineralized matrix may introduce a variation in substrate size

even prior to initiating the co-culture (53). Also, many *in vitro* formation experiments, while being

able to produce the ECM constituents collagen and mineral, are not (yet) producing real bone ECM

(51). An advantage specific to using a collagen-based material in favor of a pure ceramic material is

that techniques such as NTx (49) and CTx (50) can be used. These bone turnover markers are used in

the clinic and can quantify resorption by directly analyzing the liberated collagen fragments that

were present in the resorbed mineralized matrix (54).

Because most studies were conducted in 2D, most resorted to using various types of 2D microscopy

to analyze resorption, usually after staining to increase contrast. This can facilitate the quantification

of resorbed area using image analysis software but is usually limited to a quantification of surface

area, whereas resorption is a three-dimensional process. While methods exist to reconstruct a set of

stereoscopic 2D images into 3D height maps (55), these were not identified within the studies in

either database of this systematic map. It would be better to consider imaging techniques that can

directly quantify the resorbed volume. Examples are 2-photon microscopy for thin samples and

micro computed tomography (μCT) (56). Due to the non-destructive nature of μCT, it is well suited

to monitor mineralized volume over time within the same samples (57) and images can be compared

for changes over time (53,56). The usefulness of such a monitoring tool is however dependent on

the envisaged resolution versus the corresponding potential cell-damage caused by radiation

exposure (58,59). Registering consecutive images can even show both formation and resorption of

mineralized tissue within the same set of images of the same sample if both mineralizing OBs and

resorbing OCs are present (53). While μCT in this map is predominantly used on 3D samples, one

study used it to quantify the thickness of mineralized films and combined that data with surface

metrological data (60).

Overall, the golden standard (bone and dentine discs) remains the most-used method to study 2D

resorption, although alternatives such as osteological coatings offer new and easy ways of

quantification. Compared to 2D cultures however, 3D cultures are under-represented in this

systematic map. While the systematic search covers all publications until January 6 2020 available,

only 24 studies were labeled as 3D co-cultures in Database 1, the first being published only in 2006

(61). From these we learn that studying 3D resorption remains a challenge, with the only identified

viable options for quantification being μCT imaging and supernatant analysis techniques such as NTx

and CTx.

Formation

The result of bone formation is the deposition of mineralized matrix. This is however a multi-step

process of the presence of properly stimulated OBs that lay down a framework of type I collagen,

which in turn is mineralized by the addition of calcium phosphates (51). No single method of

measuring formation confirms the occurrence of each step in this process, instead relying on the

assumption that the confirmed presence of one step indicates the presence of the entire process.

With most studies being two-dimensional co-cultures, it is no surprise that most formation analyses

extracted from Database 1 were stainings. Of these, Alizarin Red is particularly interesting due to the

option of quantifying the amount of bound dye, which correlates to the amount of calcium (27). A

risk when using this method on larger samples is that it is not certain how far both dye application

and dye extraction penetrate the material. This should not affect relative comparisons between

different sample groups but could lead to underestimations of calcium deposition. By completely

lysing the samples and directly measuring the exact amount of calcium or phosphate (62,63) this risk

could be avoided, at the cost of not gaining information on the location and distribution of calcium

or phosphate through the sample.

The two types of non-destructive formation measurements, CICP and μCT, are coincidently well-

suited for the analysis of three-dimensional co-cultures as well. A major advantage of these is that

because of their non-destructive nature, they can be used to measure the same samples over time,

and they can be used prior to other destructive techniques. CICP measurements (64) have no

negative effects on the co-culture, requiring only that culture supernatant samples can be taken at

the desired timepoints, usually at medium exchange. The use of μCT leads to both quantification and

visualization of mineralization within the same sample over time, but it has some aspects to

consider. Most importantly, to use it as a non-destructive technique the samples must be cultured in

sterile vessels capable of being scanned. This means that experiments are limited by severe practical

constraints. Additionally, there is a direct correlation between the resolution of the images (and thus

the minimal detectable size of mineral deposits) and exposure to radiation and subsequent cell

damage (58,59). Radiation damage directly affects the usefulness as a monitoring tool, and a careful

balance between minimal acceptable resolution and maximal radiation exposure must be found.

Overall, 2D nodule stainings were the most frequently used method to measure formation.

Combined with Alizarin Red dye release these provide an easy way to quantify mineralization,

though CICP supernatant analysis and μCT techniques provide a non-destructive alternative that can

also be used for 3D co-cultures.

ALP and TRAP

ALP and TRAP are the two major markers used for indirectly quantifying respectively OB and OC

activity that were included into Database 1. Their presence is no conclusive proof that formation and

resorption are occurring because ALP is expressed in differentiating MSCs already (65) and TRAP is

expressed on monocytes as well (53), but there is a correlation between their presence and that of

OB and OC, respectively. ALP is an enzyme that makes phosphates available to be incorporated into

the matrix (66), while TRAP has been associated with migration and activation of OC (67). These

enzymes can be measured both after lysis of the cells or within the culture supernatant. The former

allows the quantification of enzyme per DNA content when combined with a DNA assay, whereas

the latter allows the monitoring of relative enzyme release over time. The precise methodological

details and experimental setup are of lesser importance for measuring ALP and TRAP than they are

for measuring formation and resorption. All that is required is the possibility to use the supernatant

or cell lysate, which is possible in most common experimental setups. The most frequently used

methods are the pNPP-based methods where ALP and TRAP directly convert a substrate into a

measurable compound. Napthtol-based methods (29) rely on a similar principle, and show an

increased specificity for TRAP isoform 5B in particular (30). The main advantage of these methods is

that they use the inherent enzymatic activity of ALP and TRAP, reducing the complexity and cost of

the assay. However, the reliance on the inherent enzymatic activity of the enzymes is also a practical

limitation as inherent activity can be affected by freezing and long-term storage. Especially when

monitoring ALP or TRAP release over time, samples are commonly frozen and stored for different

periods of time, and enzyme activity could be affected by this. A workaround would be to directly

analyze the samples after taking them, or to use methods that rely on the presence and not the

activity of these enzymes.

One of those methods is the immunoenzymatic assay, of which ELISA is the most well-known. With a

traditional ELISA the antigen is first bound to the assay plate, and then labeled with one or a series

of antibodies that are conjugated with an enzyme to convert a substrate to a chromogenic product

(68). These methods have the capacity to detect lower concentrations of protein because it is

possible to label each individual protein with an excess of new enzymes each capable of converting

substrate. In the case of TRAP, ELISA kits exist that are specific for TRAP isoform 5b which is

expressed almost exclusively in OCs (69), whereas isoform 5a is also expressed by macrophages and

dendritic cells (70). While in a co-culture with pure populations of OB and OC this distinction would

not be relevant, macrophages or macrophage-like cells can be used as a precursor for OCs (24), and

thus express isoform 5a which could be detected in a pNPP based assay. Similarly, most co-cultures

use a precursor or heterogeneous population that either contains macrophages or contains cells

capable of differentiating into macrophages such as mononuclear cells (71), which means that the

presence of other isoforms or even other phosphatases is likely. Whether this negatively affects the

results is another matter that can only be determined by comparison between the two types of

assay. Another factor to consider in co-cultures is the fact that both ALP and TRAP are phosphatases.

Assays that rely on their inherent phosphatase activity may show cross-reactivity of other

phosphatases, although this should be mitigated by controlling the pH during the test.

To conclude, pNPP based methods are the most frequently used methods for detecting ALP and

TRAP due to their affordability and simplicity. However, immunoenzymatic detection methods are

more sensitive and specific, and do not rely on the intrinsic enzymatic activity of ALP and TRAP which

can be affected by freeze-thaw cycles, long-term storage, and could show cross-reactivity with other

phosphatases.

Osteoclasts

OCs are the bone resorbing cells, and together with bone forming OBs they keep the bone mass and

bone strength in equilibrium with the required loads placed upon it. OCs are created when OC

precursors such as monocytes exit the bloodstream because of chemotactic cues followed by the

correct biochemical signals that result in cell-fusion into OCs. Cells are currently considered to be

OCs when expressing TRAP, having an actin ring, and having at least 3 nuclei (6). Osteoclastic

resorption *in vivo* is an integral part of bone maintenance. Old and damaged bone tissue is resorbed

and quickly replaced by OBs with new bone tissue.

There is a clear preference in the studies identified for Database 2 for using human cells to generate

OCs, most notably monocytes and mononuclear cells. These have in the past two decades proven to

be a reliable and relatively straight-forward precursor population for OCs (6), they can be obtained

from human blood donations, and are thought to be better representatives for studying human

physiology than cells of animal origin (2,3).

The choice of using precursors versus differentiated OCs is forced sharply into one direction because

of both biological and experimental limitations. The extraction of OCs from bone is possible but

cumbersome, requires access to fresh bone material and generally does not yield relevant numbers

of OCs. Generating OCs from circulating precursors has proven to be an easier way of obtaining OCs.

However, OCs have an average life span of approximately 2 weeks (72,73), some of which would

already be lost if OCs would be created prior to the actual experiments. In contrast to most cells,

differentiation happens by fusion of several precursors into a single OC. Fused multi-nucleated OCs

can become large and hard to handle without damaging them. For those reasons they are usually

generated within the experiment itself instead of in a prior culture. In fact, the first OB-OC co-

cultures were designed specifically to generate OCs by using osteoblastic cell signals (9), as opposed

to generating a model to study both OBs and OCs simultaneously as this systematic map has indexed

(74).

OCs can currently be obtained *in vitro* without the need for OBs thanks to the discovery in 1999 that

M-CSF and RANKL are the necessary and sufficient proteins to induce osteoclastic differentiation

from precursors (10). The cells are predominantly introduced into the co-culture as precursors to

differentiate within the co-culture, regardless of whether these two proteins are used or not. Where

in the past researchers used spleen cells for this, the studies included in this systematic map

predominantly use (blood-derived) mononuclear cells, monocytes, or macrophages as precursor

cells. These four sources are closely related, and the main differences between them are the purity

of the population and how far along the path to differentiated and active OCs they are. In short:

Spleen cells contain many cells, among others mononuclear cells. A part of the mononuclear cell

population consists of monocytes which are currently regarded as 'the' OC precursors (75,76).

Monocytes can differentiate into macrophages or fuse together into OCs, depending on the

biochemical cues received. Macrophage-like cell-lines are being used to generate OCs as well.

There are risks associated with each method of generating OCs. Animal cells introduce a between-

species variation and can respond differently than human cells (17), human donor cells tend to

exhibit large between-donor variation compared to cell lines (77) and the number of cells acquired is

limited and variable (78). The large variation between donors again highlights the need for patient-

specific disease models instead of generic bone models. By using cells of a single diseased donor, the

reaction of that patient's cells on potential treatment options can be studied. Immortalized cell-lines

result in immortal subsequently generated OC-like cells. This is however not the case *in vivo* and

while it can greatly reduce between-experiment and between-lab variation, it is also physiologically

less relevant. While these risks and characteristics do not discredit any source as a viable source of

OCs for any experiment, the results of the corresponding studies should be interpreted with these

characteristics in mind.

913 Osteoblasts

OBs are the bone forming cells, and together with bone resorbing OCs they keep the bone mass and

bone strength in equilibrium with the required loads placed upon it. In addition to their role in bone

formation, they excrete the exact biochemical cues necessary to generate OCs out of their

circulating precursors. Before the identification and commercial synthesis of these factors, a co-

culture with OB was the only way to generate OCs *in vitro*.

The preference for the use of human primary cells identified in the studies included in Database 2

can be explained by the good availability of donor material, expandability of OB precursors, and

because human cells have the potential to better reflect human physiology than cells from other

species (2,3). The choice of OB progenitors versus OBs is not as crucial here as it is with OCs. MSCs,

the most commonly used precursors, have a tri-lineage potential (79) and should be able to

differentiate into OBs on a 1-1 ratio. The advantage of osteoprogenitors such as MSCs is that these

are capable of extensive proliferation before differentiation and could be used to migrate into and

populate hard-to-reach areas within 3D scaffolds. Additionally, using progenitors opens possibilities

to study osteoblastogenesis in addition to bone formation. When the effect of an intervention on

mineralization but not osteogenesis is under investigation, care must be taken that the intervention

is not applied before differentiation is has been achieved.

The advantage of directly introducing OBs instead of precursors, whether obtained directly from

primary material or pre-differentiated *in vitro*, is that these do not need to be differentiated within

the experiment anymore, and all seeded cells are already OBs, and by extension, any experimental

conditions affect only mature OBs and not osteoblastogenesis in parallel. Actual OBs or to-be-

differentiated MSCs isolated from orthopedic surgery are the most common source of primary

human OBs. However, healthy human donor OBs are scarce because they are mostly isolated after

surgery of mainly diseased patients. Whether the use of OBs from unhealthy donors affects

experimental results needs to be elucidated. On the other hand, using patient cells to create a

personalized *in vitro* disease model is the first step towards personalized medicine, especially if all

cells are of that same patient. Finally, the use of any type of animal cell instead of human cells

carries the risk of finding inter-species differences that can affect the results and conclusions, and

everything based on that, because animal cells can behave differently than human cells (17). While

none of these risks directly discredit any of the methods obtaining OBs, the results must be

interpreted with these risks and characteristics in mind.

Culture conditions

The success of a cell-culture experiment is dependent on many factors related to culture conditions.
For most cell-types, standard culture conditions have been established. During co-culture
experiments however, the needs of two or more cell types need to be met. Medium components
and factors may be needed in different concentrations, as they can be beneficial to one cell type but
inhibitory to the other (80).

There is a clear preference for medium based on DMEM and αMEM, but the choice of base medium
for a culture is not an easy one. Base media are generally chosen based on the intended cell type,
recommendations by a manufacturer or supplier of either cells or medium, preferred effect on cells,
interaction with other supplements, and earlier experience. These factors make direct comparison of
experimental results by literature virtually impossible. Additionally, none of the studies mentioned
why they specifically chose the base media they used.

Another variable in medium composition is the use of FBS (or FCS). It is commonly known that there
can be batch-to-batch and between-brand differences in FBS (81) which can impact the results of an
experiment tremendously. While different concentrations are being used, the most common FBS
concentration is 10%. However, no study explains why each type and concentration of FBS was used.

Although there was no clear predictor for using or not using any of the osteoblastic or osteoclastic
supplements, when they were used, the concentrations were usually within the same order of
magnitude in all studies, except for ascorbic acid. However, only 2 studies used all 5 of the
supplements indexed in this study (AA, βGP, Dexamethasone, M-CSF and RANKL) and many
combinations of supplements have been registered in this map. Looking at OC supplements, it is
generally accepted that RANKL and M-CSF are both necessary and sufficient for osteoclastogenesis
(10). However, OBs can produce RANKL and M-CSF themselves to trigger differentiation (9) and

therefore the supplements are not necessarily required in co-culture. The need for all osteoblastic

supplements is not as great considering osteoblasts can be introduced in various stages of

development. Still, each supplement contributes to a specific function. Dexamethasone upregulates

osteogenic differentiation, βGP acts as a phosphate source, and AA is a co-factor involved in collagen

synthesis (82). Depending on the type of cells introduced, the aim of the experiment and other

methodological details, their inclusion could be beneficial.

Finally, many studies used or omitted specific supplements related to their research question

regarding the activity of OBs or OCs or used less common supplements for differentiation such as

vitamin D3, human serum or Phorbol 12-myristate 13-acetate. What is seldom addressed however,

is the compromise that must be made in choosing the right supplements and concentrations. Adding

too high doses of supplements could cause an excess of these signals in the culture medium,

effectively overshadowing any other ongoing cell-signaling over the same pathway by other cells.

This is of critical importance when the goal is not to achieve only OBs and/or OCs activity, but to

obtain a homeostasis in which the two cell types regulate each other, with experimental conditions

or interventions that are expected to affect this balance. Here, it may be beneficial to experiment

with lower concentrations of factors, supplemented only during critical phases of the cells'

development or differentiation.

The choice of medium in a co-culture is most likely going to be a compromise and must be based on

the exact research question to be addressed, where the advantages and disadvantages of base

media and supplements for both cell types are carefully weighed. Most likely, the ultimate goal for

the envisaged co-culture would be to reach tissue homeostasis, in which the environment is as

similar to the *in vivo* environment in tissue homeostasis where cell interactions with each other can

be monitored.

Seeding densities and seeding ratios

Using the correct seeding densities plays a major role in proliferation and cell function of OBs (18,83) and osteoclastic differentiation (84). The seeding densities reported in this map show an enormous spread. Many factors could have influenced these numbers. For example, some studies report the numbers prior to expansion, others expand the cells in (co-)culture. Similarly, the percentages of relevant precursor cells in heterogenous cell populations can vary widely. The cell numbers present and OB:OC ratio most likely even change during a co-culture due to ongoing cell-division, differentiation, fusion and different expected life spans and the corresponding cell death. Regrettably, the available documentation of exact cell numbers introduced is often lacking, and open to some interpretation. While the figures show this large spread in data points, the included databases can be manipulated to filter and select studies that match criteria according to the readers' specific needs.

Animal type, cell type, cell line versus primary cells and even passage number may also directly influence the choice of seeding densities in addition to various experimental choices. At the same time, the purpose of the experiment and more specifically the purpose of the cells and type of interaction required should determine the necessary seeding density. Are the cells required to actively deposit or resorb measurable amounts of minerals, or are they just supposed to be there to facilitate OB-OC communication? The combination of all these factors suggests that there in fact is no one ideal seeding density, that the best density for a certain experiment can only be determined by taking all the above factors into account, learning from others that did a similar experiment, and most importantly verifying assumptions and predictions in the lab.

Looking at the cell seeding ratio, here reported as number of seeded OB/OB-precursors per seeded OC/OC-precursor, outliers can be normalized against their seeded counterparts. In 2D studies, there are never more OBs/OB-precursors than OCs/OC-precursors. At most, they are seeded at a 1:1 OB:OC ratio. Even though in human bone tissue the ratio of OB:OC is estimated to be approximately

1017 7:1 (41), higher OC numbers than OB numbers are not unexpected. OB precursors can still

1018 proliferate, whereas OC precursors usually still need to fuse together to form mature OC or OC-like

1019 cells. In 3D we do not see the same trend, with ratio's ranging from 1:20 to 100:1. These differences

1020 are again affected by the same factors that influence individual OB and OC seeding densities, further

1021 enhanced by the extra layer of complexity that are inherent to 3D cultures.

1022

1023 Limitations of the systematic search

While the authors took great care to construct a series of search queries fine-tuned for each of the

three online bibliographic literature sources, the authors cannot be certain that all relevant OB-OC

co-cultures have been included into the two databases. The search was limited by the necessary

addition of a 'co-culture' search element. Co-culture studies without any indication thereof in the

title or abstract simply cannot be identified through the initial search. To compensate for this,

screening step 4, searching through identified reviews and publications included into Database 2,

was executed. The publications included into Database 1 or the complete list of identified OB-OC co-

cultures could have been screened for references as well, but the authors decided against this.

Database 2 was specifically chosen for this because the likelihood of a publication that matches all

relevant inclusion criteria citing other such publications was deemed high, whereas less relevant

papers (included into Database 1, or not included at all) were considered much less likely to cite

publications relevant to this systematic map that had not already been identified by the search itself

or the screening of reviews and Database 2. Publications in languages other than English, Dutch or

German were excluded because none of the researchers involved in data curation and analysis were

fluent in those languages. No budget was available to hire a professional translator for the remaining

languages. The consequence of that is that there is a likelihood that relevant publications were

missed.

1041

Limitations of the databases

1042

The use of co-culture models is a field that is still developing, and we are now aware that it is not

only about adding an additional cell type, but that the complexity of such a culture is more than just

doubled. The applied choice of methods, cells, and culture conditions should be tailored to the

research question to be investigated, and ideally would be comparable to other studies within the

field. This systematic map shows that the currently applied methods are far from standardized and

that many research groups have developed their own approach attempting to overcome each

challenge, making comparison between research groups virtually impossible. There is no consensus

on cell types, seeding densities, seeding ratios or medium composition, and many of these are

predominantly determined by the research question and whatever has been done before in each

respective laboratory. For each study, 86 columns worth of data has been extracted including in

some cases extrapolation and recalculation of numbers, which are now available for sorting and

filtering for individual needs. Still these databases only scratch the surface of each study, and to fully

understand the collected information and the context on which it was gathered, one must still read

the full publication.

It must be noted that the quality of reporting in many cases is lacking. Both missing information

critical and non-critical for reproducing the methods of the studies was identified, and only 13 out of

39 studies included in Database 2 did not miss at least a basic description of all indexed

characteristics. However, more relevant details of these characteristics may have been omitted that

describe exactly how each method or culture conditions was executed that were not required for

this systematic map. Instead, this systematic map focuses on a high-level indexing and evaluation of

defining characteristics of methods and culture conditions.

This systematic map is not intended to provide a definitive answer to the question of how to set up

the perfect OB-OC co-culture. Instead, it allows searching through all relevant co-culture studies

looking for specific matching experimental characteristics or culture details that may be applicable

to one's own research. For this, it contains the possibility to search, sort and filter through many

relevant characteristics. This allows one to find relevant studies that may have already (partly)

studied one's research question, or that can be used as a guide to design comparable experiments.

1070

Conclusion

1071

With this systematic map, we have generated an overview of existing OB-OC co-culture studies

published until January 6, 2020, their methods, predetermined outcome measures (formation and

resorption, and ALP and TRAP quantification as surrogate markers for formation and resorption,

respectively), and other useful parameters for analysis. The two constructed databases are intended

to allow researchers to quickly identify publications relevant to their specific needs, which otherwise

would have not been easily available or findable. The presented high-level evaluation and discussion

of the major extracted methodological details provides important background information and

context, suggestions and considerations covering most of the used cell sources, culture conditions

and methods of analysis. Finally, this map includes the instructions for others to expand and

manipulate the databases to answer their own more specific research questions.

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

References

1.   Feng X, McDonald JM. Disorders of Bone Remodeling. Annu Rev Pathol. 2011;6:121–45.

2.   Burkhardt AM, Zlotnik A. Translating translational research: mouse models of human disease.
Cell Mol Immunol. 2013 Sep;10(5):373–4.

3.   Contopoulos-Ioannidis DG, Ntzani E, Ioannidis JPA. Translation of highly promising basic
science research into clinical applications. Am J Med. 2003 Apr 15;114(6):477–84.

4.   Montagutelli X. Animal models are essential to biological research: issues and perspectives.
Futur Sci OA. 2015;1(4).

5.   Thomas DW, Burns J, Audette J, Adam Carroll, Dow-Hygelund C, Hay M. Clinical Development

Success Rates 2006-2015. 2016.

6.   Owen R, Reilly GC. In vitro Models of Bone Remodelling and Associated Disorders. Front

Bioeng Biotechnol. 2018;6:134.

7.   Holmes A, Brown R, Shakesheff K. Engineering tissue alternatives to animals: applying tissue

engineering to basic research and safety testing. Regen Med. 2009;4(4):579–92.

8.   Chambers TJ. Osteoblasts release osteoclasts from calcitonin-induced quiescence. J Cell Sci.

1982;57:247–60.

9.   Takahashi N, Akatsu T, Udagawa N, Sasaki T, Yamaguchi A, Moseley JM, et al. Osteoblastic

cells are involved in osteoclast formation. Endocrinology. 1988 Nov;123(5):2600–2.

10.   Suda T, Takahashi N, Udagawa N, Jimi E, Gillespie MT, Martin TJ. Modulation of osteoclast

differentiation and function by the new members of the tumor necrosis factor receptor and

ligand families. Vol. 20, Endocrine Reviews. Endocrine Society; 1999. p. 345–57.

11.   Simonet WS, Lacey DL, Dunstan CR, Kelley M, Chang MS, Lüthy R, et al. Osteoprotegerin: A

novel secreted protein involved in the regulation of bone density. Cell. 1997 Apr

18;89(2):309–19.

12.   Udagawa N, Takahashi N, Akatsu T, Tanaka H, Sasaki T, Nishihara T, et al. Origin of

osteoclasts: mature monocytes and macrophages are capable of differentiating into

osteoclasts under a suitable microenvironment prepared by bone marrow-derived stromal

cells. Proc Natl Acad Sci U S A. 1990 Sep 1;87(18):7260–4.

13.   Teitelbaum SL. Bone resorption by osteoclasts. Science. 2000 Sep 1;289(5484):1504–8.

14.   de Vries RBM, Hooijmans CR, Langendam MW, van Luijk J, Leenaars M, Ritskes-Hoitinga M, et

al. A protocol format for the preparation, registration and publication of systematic reviews

of animal intervention studies. Evidence-based Preclin Med. 2015 Aug 1;2(1):e00007.

15.   Remmers SJA, Wildt BWM, Vis MVA, Spaander ESR, de Vries R, Hofmann S, et al. Osteoblast-

osteoclast co-culture models of bone-remodelling: A systematic review protocol [Internet].

2020 [cited 2021 Jan 16]. Available from:

https://doi.org/10.5281/zenodo.3969441#.YAK3zqvkF6c.mendeley

16.   Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for

systematic reviews. Syst Rev. 2016 Dec 5;5(1):210.

17.   Jemnitz K, Veres Z, Monostory K, Kóbori L, Vereczkey L. Interspecies differences in

acetaminophen sensitivity of human, rat, and mouse primary hepatocytes. Toxicol In Vitro.

2008;22(4):961–7.

18.   Bitar M, Brown RA, Salih V, Kidane AG, Knowles JC, Nazhat SN. Effect of cell density on

osteoblastic differentiation and matrix degradation of biomimetic dense collagen scaffolds.

Biomacromolecules. 2008;9(1):129–35.

19.   Jolly JJ, Chin K-Y, Farhana MFN, Alias E, Chua KH, Hasan WNW, et al. Optimization of the

Static Human Osteoblast/Osteoclast Co-culture System. Iran J Med Sci. 2018 Mar;43(2):208–

13.

20.   Jones GL, Motta A, Marshall MJ, El Haj AJ, Cartmell SH. Osteoblast: osteoclast co-cultures on

silk fibroin, chitosan and PLLA films. Biomaterials. 2009;30(29):5376–84.

21.   Kadow-Romacker A, Duda GN, Bormann N, Schmidmaier G, Wildemann B. Slight changes in

the mechanical stimulation affects osteoblast- and osteoclast-like cells in co-culture. Transfus

Med Hemother. 2013 Dec;40(6):441–7.

22.   De Vries TJ, Schoenmaker T, Aerts D, Grevers LC, Souza PPC, Nazmi K, et al. M-CSF priming of

osteoclast precursors can cause osteoclastogenesis-insensitivity, which can be prevented and

overcome on bone. J Cell Physiol. 2015 Jan;230(1):210–25.

23.   Kyllönen L, Haimi S, Mannerström B, Huhtala H, Rajala KM, Skottman H, et al. Effects of

different serum conditions on osteogenic differentiation of human adipose stem cells in vitro.

Stem Cell Res Ther. 2013 Feb 15;4(1):17.

24.   Hikita A, Iimura T, Oshima Y, Saitou T, Yamamoto S, Imamura T. Analyses of bone modeling

and remodeling using in vitro reconstitution system with two-photon microscopy. Bone.

2015;76:5–17.

25.   Puchtler H, Meloan SN, Terry MS. On the history and mechanism of alizarin and alizarin red S

stains for calcium. J Histochem Cytochem Off J Histochem Soc. 1969;17(2):110–24.

26.   Meloan SN, Puchtler H. Chemical Mechanisms of Staining Methods: Von Kossa's Technique:

What von Kossa Really Wrote and a Modified Reaction for Selective Demonstration of

Inorganic Phosphates. J Histotechnol. 1985;8(1):11–3.

27.   Schroder HC, Wang XH, Wiens M, Diehl-Seifert B, Kropf K, Schlossmacher U, et al. Silicate

modulates the cross-talk between osteoblasts (SaOS-2) and osteoclasts (RAW 264.7 cells):

inhibition of osteoclast growth and differentiation. J Cell Biochem. 2012;113(10):3197–206.

28.   Hayman AR. Tartrate-resistant acid phosphatase (TRAP) and the osteoclast/immune cell

dichotomy. Autoimmunity. 2008 Jan 7;41(3):218–23.

29.   Vaughan A, Guilbault GG, Hackney D. Fluorometric methods for analysis of acid and alkaline

phosphatase. Anal Chem. 1971;43(6):721–4.

30.   Janckila AJ, Takahashi K, Sun SZ, Yam LT. Naphthol-ASBI phosphate as a preferred substrate

for tartrate-resistant acid phosphatase isoform 5b. J Bone Miner Res. 2001;16(4):788–93.

31.   Davidov M. Histochemical detection of alkaline phosphatase by means of naphthol

phosphate AS-MX in association with fast blue salt BB in the large acinous glands - PubMed.

Nauchni Tr Viss Med Inst Sofiia. 1967;46(2):13–8.

32.   Stagni N, Vittur F, de Bernard B. Solubility properties of alkaline phosphatase from matrix

1171   vesicles. BBA - Gen Subj. 1983 Dec 27;761(3):246–51.

1172   33.   Millán JL. The role of phosphatases in the initiation of skeletal mineralization. Calcif Tissue Int.

1173   2013 Oct;93(4):299–306.

1174   34.   Lequin RM. Enzyme immunoassay (EIA)/enzyme-linked immunosorbent assay (ELISA). Clin

1175   Chem. 2005 Dec;51(12):2415–8.

1176   35.   Lindner U, Kramer J, Rohwedel J, Schlenke P. Mesenchymal stem or stromal cells: Toward a

1177   better understanding of their biology? Vol. 37, Transfusion Medicine and Hemotherapy.

1178   Karger Publishers; 2010. p. 75–83.

1179   36.   Clarke MSF, Sundaresan A, Vanderburg CR, Banigan MG, Pellis NR. A three-dimensional tissue

1180   culture model of bone formation utilizing rotational co-culture of human adult osteoblasts

1181   and osteoclasts. Acta Biomater. 2013 Aug;9(8):7908–16.

1182   37.   Kadow-Romacker A, Duda GN, Bormann N, Schmidmaier G, Wildemann B. Slight changes in

1183   the mechanical stimulation affects osteoblast- and osteoclast-like cells in co-culture. Transfus

1184   Med Hemother. 2013;40(6):441–7.

1185   38.   Hammerl A, Diaz Cano CE, De-Juan-Pardo EM, van Griensven M, Poh PSP. A Growth Factor-

1186   Free Co-Culture System of Osteoblasts and Peripheral Blood Mononuclear Cells for the

1187   Evaluation of the Osteogenesis Potential of Melt-Electrowritten Polycaprolactone Scaffolds.

1188   Int J Mol Sci. 2019;20(5).

1189   39.   Bongio M, Lopa S, Gilardi M, Bersini S, Moretti M. A 3D vascularized bone remodeling model

1190   combining osteoblasts and osteoclasts in a CaP nanoparticle-enriched matrix. Nanomedicine

1191   (London, England). 2016 May;11(9):1073–91.

40.   Domaschke H, Gelinsky M, Burmeister B, Fleig R, Hanke T, Reinstorf A, et al. In vitro

ossification and remodeling of mineralized collagen I scaffolds. Tissue Eng. 2006

Apr;12(4):949–58.

41.   Gruber HE, Ivey JL, Thompson ER, Chesnut CH, Baylink DJ. Osteoblast and osteoclast cell

number and cell activity in postmenopausal osteoporosis. Miner Electrolyte Metab. 1986 Jan

1;12(4):246–54.

42.   Gremare A, Aussel A, Bareille R, Paiva Dos Santos B, Amedee J, Thebaud NB, et al. A Unique

Triculture Model to Study Osteoblasts, Osteoclasts, and Endothelial Cells. Tissue Eng Part C

Methods. 2019;25(7):421–32.

43.   Boskey AL. Mineralization of Bones and Teeth. Elements. 2007 Dec 1;3(6):385–91.

44.   Rumpler M, Würger T, Roschger P, Zwettler E, Sturmlechner I, Altmann P, et al. Osteoclasts

on bone and dentin in vitro: mechanism of trail formation and comparison of resorption

behavior. Calcif Tissue Int. 2013 Dec;93(6):526–39.

45.   Teti A, Grano M, Colucci S, Cantatore FP, Loperfido MC, Zallone AZ. Osteoblast-osteoclast

relationships in bone resorption: osteoblasts enhance osteoclast activity in a serum-free co-

culture system. Biochem Biophys Res Commun. 1991;179(1):634–40.

46.   Armour KJ, van 't Hof RJ, Armour KE, Torbergsen AC, Del Soldato P, Ralston SH. Inhibition of

bone resorption in vitro and prevention of ovariectomy-induced bone loss in vivo by

flurbiprofen nitroxybutylester (HCT1026). Arthritis Rheum. 2001;44(9):2185–92.

47.   Buckley KA, Hipskind RA, Gartland A, Bowler WB, Gallagher JA. Adenosine triphosphate

stimulates human osteoclast activity via upregulation of osteoblast-expressed receptor

activator of nuclear factor-kappa B ligand. Bone. 2002;31(5):582–90.

48.   Lalande A, Roux S, Denne MA, Stanley ER, Schiavi P, Guez D, et al. Indapamide, a thiazide-like

diuretic, decreases bone resorption in vitro. J Bone Min Res. 2001;16(2):361–70.

49.   Rossi E, Mracsko E, Papadimitropoulos A, Allafi N, Reinhardt D, Mehrkens A, et al. An In Vitro

1217   Bone Model to Investigate the Role of Triggering Receptor Expressed on Myeloid Cells-2 in

1218   Bone Homeostasis. Tissue Eng Part C Methods. 2018;24(7):391–8.

1219   50.   Krishnan V, Vogler EA, Sosnoski DM, Mastro AM. In vitro mimics of bone remodeling and the

1220   vicious cycle of cancer in bone. J Cell Physiol. 2014;229(4):453–62.

1221   51.   de Wildt BWM, Ansari S, Sommerdijk NAJM, Ito K, Akiva A, Hofmann S. From bone

1222   regeneration to three-dimensional in vitro models: tissue engineering of organized bone

1223   extracellular matrix. Vol. 10, Current Opinion in Biomedical Engineering. Elsevier B.V.; 2019.

1224   p. 107–15.

1225   52.   Delaisse J-M. The reversal phase of the bone-remodeling cycle: cellular prerequisites for

1226   coupling resorption and formation. BoneKEy reports. 2014 Jan;3:561.

1227   53.   Remmers S, Mayer D, Melke J, Ito K, Hofmann S. Measuring mineralised tissue formation and

1228   resorption in a human 3d osteoblast-osteoclast co-culture model. Eur Cell Mater.

1229   2020;40:189–202.

1230   54.   Shetty S, Kapoor N, Bondu J, Thomas N, Paul T. Bone turnover markers: Emerging tool in the

1231   management of osteoporosis. Vol. 20, Indian Journal of Endocrinology and Metabolism.

1232   Medknow Publications; 2016. p. 846–52.

1233   55.   Chan EKM, Darendeliler MA, Petocz P, Jones AS. A new method for volumetric measurement

1234   of orthodontically induced root resorption craters. Eur J Oral Sci. 2004 Apr;112(2):134–9.

1235   56.   Hagenmüller H, Hofmann S, Kohler T, Merkle HP, Kaplan DL, Vunjak-Novakovic G, et al. Non-

1236   invasive time-lapsed monitoring and quantification of engineered bone-like tissue. Ann

1237   Biomed Eng. 2007 Oct;35(10):1657–67.

57.   Melke J, Zhao F, van Rietbergen B, Ito K, Hofmann S. Localisation of mineralised tissue in a

complex spinner flask environment correlates with predicted wall shear stress level

localisation. Eur Cell Mater. 2018 Jul 31;36:57–68.

58. Kraehenbuehl TP, Stauber M, Ehrbar M, Weber F, Hall H, Muller R. Effects of muCT radiation

on tissue engineered bone-like constructs. Biomed Tech. 2010;55(4):245–50.

59. Yang B, Zhou H, Zhang X-D, Liu Z, Fan F-Y, Sun Y-M. Effect of radiation on the expression of

osteoclast marker genes in RAW264.7 cells. Mol Med reports. 2012 Apr;5(4):955–8.

60. Hayden RS, Quinn KP, Alonzo CA, Georgakoudi I, Kaplan DL. Quantitative characterization of

mineralized silk film remodeling during long-term osteoblast-osteoclast co-culture.

Biomaterials. 2014;35(12):3794–802.

61. Domaschke H, Gelinsky M, Burmeister B, Fleig R, Hanke T, Reinstorf A, et al. In vitro

ossification and remodeling of mineralized collagen I scaffolds. Tissue Eng. 2006;12(4):949–

58.

62. Hayden RS, Fortin JP, Harwood B, Subramanian B, Quinn KP, Georgakoudi I, et al. Cell-

tethered ligands modulate bone remodeling by osteoblasts and osteoclasts. Adv Funct Mater.

2014;24(4):472–9.

63. Loomer PM, Ellen RP, Tenenbaum HC. Effects of Porphyromonas gingivalis 2561 extracts on

osteogenic and osteoclastic cell function in co-culture. J Periodontol. 1998;69(11):1263–70.

64. Boanini E, Torricelli P, Sima F, Axente E, Fini M, Mihailescu IN, et al. Strontium and

zoledronate hydroxyapatites graded composite coatings for bone prostheses. J Colloid

Interface Sci. 2015;448:1–7.

65. Kim YH, Yoon DS, Kim HO, Lee JW. Characterization of different subpopulations from bone

marrow-derived mesenchymal stromal cells by alkaline phosphatase expression. Stem Cells

Dev. 2012 Nov 1;21(16):2958–68.

66.   Golub EE, Boesze-Battaglia K. The role of alkaline phosphatase in mineralization. Vol. 18,

Current Opinion in Orthopaedics. 2007. p. 444–8.

67.   Sheu TJ, Schwarz EM, Martinez DA, O'Keefe RJ, Rosier RN, Zuscik MJ, et al. A phage display

technique identifies a novel regulator of cell differentiation. J Biol Chem. 2003 Jan

3;278(1):438–43.

68.   Engvall E, Perlmann P. Enzyme-linked immunosorbent assay (ELISA) quantitative assay of

immunoglobulin G. Immunochemistry. 1971;8(9):871–4.

69.   Halleen JM, Tiitinen SL, Ylipahkala H, Fagerlund KM, Väänänen HK. Tartrate-resistant acid

phosphatase 5b (TRACP 5b) as a marker of bone resorption. Clin Lab. 2006;52(9–10):499–

509.

70.   Halleen JM, Ylipahkala H, Alatalo SL, Janckila AJ, Heikkinen JE, Suominen H, et al. Serum

tartrate-resistant acid phosphatase 5b, but not 5a, correlates with other markers of bone

turnover and bone mineral density. Calcif Tissue Int. 2002;71(1):20–5.

71.   Young PS, Tsimbouri PM, Gadegaard N, Meek R, Dalby MJ.

Osteoclastogenesis/osteoblastogenesis using human bone marrow-derived cocultures on

nanotopographical polymer surfaces. Nanomedicine (London, England). 2015 Mar;10(6):949–

57.

72.   Parfitt AM. Osteonal and hemi-osteonal remodeling: The spatial and temporal framework for

signal traffic in adult human bone. J Cell Biochem. 1994 Jul;55(3):273–86.

73.   Manolagas SC. Birth and death of bone cells: basic regulatory mechanisms and implications

for the pathogenesis and treatment of osteoporosis. Endocr Rev. 2000 Apr;21(2):115–37.

74.   Bongio M, Lopa S, Gilardi M, Bersini S, Moretti M. A 3D vascularized bone remodeling model

combining osteoblasts and osteoclasts in a CaP nanoparticle-enriched matrix. Nanomed.

2016 May;11(9):1073–91.

75.   Udagawa N, Takahashi N, Akatsu T, Tanaka H, Sasaki T, Nishihara T, et al. Origin of

osteoclasts: Mature monocytes and macrophages are capable of differentiating into

osteoclasts under a suitable microenvironment prepared by bone marrow-derived stromal

cells. Proc Natl Acad Sci U S A. 1990;87(18):7260–4.

76.   Kylmäoja E, Nakamura M, Turunen S, Patlaka C, Andersson G, Lehenkari P, et al. Peripheral

blood monocytes show increased osteoclast differentiation potential compared to bone

marrow monocytes. Heliyon. 2018 Sep 1;4(9):e00780.

77.   Susa M, Luong-Nguyen N-H, Cappellen D, Zamurovic N, Gamse R. Human primary osteoclasts:

in vitro generation and applications as pharmacological and clinical assay. J Transl Med. 2004

Mar 16;2(1):6.

78.   Yang J, Qiao M, Li Y, Hu G, Song C, Xue L, et al. Expansion of a population of large monocytes

(atypical monocytes) in peripheral blood of patients with acute exacerbations of chronic

obstructive pulmonary diseases. Mediators Inflamm. 2018 May 17;2018:1–13.

79.   Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, et al. Multilineage

potential of adult human mesenchymal stem cells. Science. 1999 Apr 2;284(5411):143–7.

80.   Vis MAM, Ito K, Hofmann S. Impact of Culture Medium on Cellular Interactions in in vitro Co-

culture Systems. Front Bioeng Biotechnol. 2020 Aug 4;8:911.

81.   van der Valk J, Bieback K, Buta C, Cochrane B, Dirks WG, Fu J, et al. Fetal Bovine Serum (FBS):

Past - Present - Future. ALTEX. 2018;35(1):99–118.

82.   Langenbach F, Handschel J. Effects of dexamethasone, ascorbic acid and β-glycerophosphate

on the osteogenic differentiation of stem cells in vitro. Vol. 4, Stem Cell Research and

Therapy. BioMed Central; 2013. p. 117.

83.   Wiedmann-Al-Ahmad M, Gutwald R, Lauer G, Hübner U, Schmelzeisen R. How to optimize

seeding and culturing of human osteoblast-like cells on various biomaterials. Biomaterials.

2002;23(16):3319–28.

84.   Motiur Rahman M, Takeshita S, Matsuoka K, Kaneko K, Naoe Y, Sakaue-Sawano A, et al.

Proliferation-coupled osteoclast differentiation by RANKL: Cell density as a determinant of

osteoclast formation. Bone. 2015 Dec 1;81:392–9.

1314

1315

Supporting information

S1 File. Database 1. This database contains all studies in which at least one relevant outcome measure was investigated. Characteristics of outcome measures and descriptive statistics are listed in this database.

S2 File. Database 2. This database contains all studies in which at least one relevant outcome measure was investigated for both OB and OC. Characteristics of cells, methods and culture conditions, and descriptive statistics are listed in this database.

S3 File. Using the databases. This document provides instructions on how to operate the databases, how to add publications and expand the analyses with more elements.

S4 File. List of all OB-OC co-cultures. This list contains the initial list of 694 OB-OC cocultures obtained after screening, before full-text investigation and exclusion based on outcome measures.

S5 File PRISMA checklist. The PRISMA checklist describing all elements of the systematic review, and on what page or which section of the submitted manuscript to find them.

1329 **S6 File Systematic Review Protocol and Search Queries.** The protocol and search queries as they

1330 were published prior to execution of the fulltext screening phase.

Systematic literature search

Identification of cocultures

Trans well · Direct · 3D/scaffold · Bioreactor

Database 1

Database 2

Primary outcome

Formation · Resorption

Secondary outcome

ALP · TRAP

Or

And

Data extraction

Resorption or formation related outcome measures

Resorption and formation related outcome measures and culture conditions

A: Database 1

B: Database 2