Lateral surface pressure generated by nascent ribosomal RNA suppresses growth of fibrillar centers in the nucleolus

Tetsuya Yamamoto[a,b], Tomohiro Yamazaki[c], Kensuke Ninomiya[c], Tetsuro Hirose[c,d]

[a]Institute for Chemical Reaction Design and Discovery, Hokkaido University, Kita 21, Nishi 10, Kita-ku, Sapporo, 001-0021, Japan.
[b]PRESTO, Japan Science and Technology Agency (JST), 4-1-8, Honcho, Kawaguchi, Saitama 332-0012, Japan.
[c]Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita 565-0871, Japan.
[d]Institute for Open and Transdisciplinary Research Initiatives, Osaka University, 1-3 Yamadaoka, Suita 565-0871, Japan

The authors declare no conflict of interest.
[1]T. Yamamoto and T. Yamazaki contributed equally.
[2]To whom correspondence should be addressed. E-mail: (Theory) tyamamoto@icredd.hokudai.ac.jp and (Experiment) tyamazaki@fbs.osaka-u.ac.jp

All the authors designed and discussed the research and wrote and revised the paper. T. Yamamoto developed the theoretical model and performed calculations. T. Yamazaki performed the experiments.

ABSTRACT

Liquid-liquid phase separation (LLPS) has been thought to be the biophysical principle governing the assembly of the multiphase structures of nucleoli, the site of ribosomal biogenesis. Condensates assembled through LLPS increase their sizes to minimize the surface energy as far as their components are available. However, multiple microphases, fibrillar centers (FCs), dispersed in a nucleolus are stable and their sizes do not grow unless the transcription of pre-ribosomal RNA (pre-rRNA) is inhibited. To understand the mechanism of the suppression of the FC growth, we here construct a minimal theoretical model by taking into account the nascent pre-rRNAs tethered to the FC surfaces by RNA polymerase I. Our theory predicts that nascent pre-rRNAs generate the lateral osmotic pressure that counteracts the surface tension of the FCs and this suppresses the growth of the FCs over the stable size. The stable FC size decreases with increasing the transcription rate and decreasing the RNA processing rate. This prediction is supported by our experiments showing that RNA polymerase inhibitors increase the FC size in a

dose-dependent manner. This theory may provide insight into the general mechanism of
the size control of nuclear bodies.

Nucleolus | Transcription | Phase separation

Significance statement

The nucleolus, a site of pre-ribosomal RNA (pre-rRNA) production, has a characteristic
multiphase structure, which has been thought to be assembled through liquid-liquid
phase separation (LLPS). Although condensates assembled through LLPS grow by
coarsening or coalescence as far as the components are available, the multiple inner
phases, fibrillar centers (FCs), are dispersed in a nucleolus. To investigate the underlying
mechanism, we constructed a minimal theoretical model by considering nascent pre-
rRNAs tethered to RNA polymerase I at the FC surface. This model is supported by our
experiments and explains previous experimental observations. This work shed light on
the role of nascent RNAs to control the size of nuclear bodies.

INTRODUCTION

In the interchromatin spaces, there are a variety of nuclear bodies, such as nucleoli (1-
4), nuclear speckles (5), and paraspeckles (6). Many of the nuclear bodies are scaffolded
by ribonucleoprotein (RNP) complexes, which are composed of RNA and RNA-binding
proteins (RBPs). The class of RNAs, which are essential to the assembly of specific
subcellular bodies, is called architectural RNA (arcRNA) (6-8). A growing number of
researches suggest that some nuclear bodies are assembled via liquid-liquid phase
separation (LLPS), which is driven by the multivalent interaction between RBPs bound
to arcRNAs (6-8). Condensates produced by LLPS are spherical and increase their size
by coarsening and/or coalescence due to the surface tension (macroscopic phase
separation) (9).

Nucleoli are nuclear bodies, where ribosome assembly takes place (1,2). In the nucleoli,
pre-ribosomal RNAs (pre-rRNAs) are transcribed by RNA polymerase I (Pol I) and
maturated to construct ribosomes with ribosomal proteins. A nucleolus is not a uniform
disordered liquid, but multiple microphases, called fibrillar centers (FCs), are dispersed
in the sea of a granular component (GC) (Fig. 1A). Fibrillarins (FBLs), which are RBPs
interacting with pre-rRNAs, are condensed to form another phase, called the dense
fibrillar component (DFC), at the interfaces between FCs and the GC. The multiphase
structures of nucleoli have been thought to be assembled via simple LLPS (10). However,

this picture may not be complete because the FCs do not show coalescence or coarsening to increase their sizes. Indeed, when the Pol I transcription of pre-rRNA is inhibited, FCs show coalescence, as in the case of LLPS, and are excluded to the surface of the nucleolus (the excluded FCs are called nucleolar caps) (11). This implies that transcription somehow suppresses the growth of FCs. Ribosomal DNA (rDNA), from which pre-rRNAs are transcribed, is a repeat sequence of coding units (10 kb) intervened by the intergenic regions (30 kb). The transcriptionally active rDNA units and Pol I are localized at the surfaces of FCs where the transcription of pre-rRNAs takes place (3, 11-13). The DFC layer is probably assembled by the RNP complexes of nascent pre-rRNAs and the associated RBPs, such as FBLs (18). FBLs show phase separation with the physiological concentration and bind to nascent pre-rRNAs (10,13). These experimental results imply that nascent pre-rRNAs may act as `surfactants' that suppress the growth of FCs.

Multiple paraspeckles scaffolded by NEAT1_2 arcRNAs are dispersed in a nucleoplasm (14). This situation is somewhat analogous to FCs in a nucleolus. Paraspeckles form the core-shell structure, where the two terminal regions of NEAT1_2 form the shell and the middle region of NEAT1_2 forms the core (15,16). We have recently shown that the terminal regions of NEAT1_2 in the shell suppresses the growth of paraspeckles, analogous to micelles of triblock copolymers (14,17). The size of paraspeckles increases with the NEAT1_2 transcription upregulation (14,18), which is opposite to the response of FCs in a nucleolus to transcription upregulation. This implies that the size control mechanism of FCs in a nucleolus is different from that of paraspeckles.

We here construct a simple theoretical model that predicts the contributions of nascent pre-rRNAs to the assembly of the multiphase structure of nucleolus. This model takes into account the phase separation, the transcription kinetics, and the interfacial effects, which are probably the minimum to account for the multiphase structure. Our theory predicts that the nascent pre-rRNAs are stretched to accommodate FBLs in the DFC layer and generate the lateral pressure that counteracts with the interfacial tensions. The size of FCs is determined by the balance of the interfacial tensions and the lateral pressure. Our theory predicts that the radius of FCs is proportional to the inverse of the square root of the transcription rate. The suppression of the FC growth by the pre-rRNA transcription results from the fact that these complexes are end-grafted to the FC surfaces via Pol I, in contrast to other condensates in which arcRNAs diffuse freely and increase their sizes by the transcription upregulation. To test our prediction, we experimentally measured the FC sizes and the pre-rRNA levels while the transcription

rate is changed by the dose of BMH-21 or CX-5461, which specifically inhibits the Pol I

transcription. The scaling exponent predicted by our theory is consistent with our

experimental results, implying that the lateral pressure generated by nascent pre-rRNAs

are possible mechanism of the size control of FCs. We anticipate that this theory can be

a base theory to further look into the contribution of the multiphase structure in the

function of nucleoli and can be extended to study the mechanism of the size control and

the functions of other nuclear bodies, such as nuclear speckles and transcriptional

condensates.

RESULTS

Model of nucleolus

We here construct a minimal model of a nucleolus to predict the size of FCs in the steady

state. The nucleolus is composed of a GC in which multiple spherical FCs are dispersed

(Fig. 1A). Pol I molecules (shown by white particles in Fig. 1B) are entrapped in the FCs

(shown by a light blue droplet in Fig. 1B). The transcriptionally active repeat units of rDNA

(shown by the black line in Fig. 1B) are localized at the surfaces of the FCs. We assume

that the number of Pol I molecules N_{pol} and the number of transcriptionally active rDNA

units N_{act} in the nucleolus are constant. Nascent pre-rRNAs (shown by chains of green

particles in Fig. 1B) are localized at the surfaces of FCs and form a DFC layer (Fig. 1B).

Recent experiment revealed that only the 5' terminal external transcribed spacer (ETS)

regions of nascent pre-rRNAs are localized at the DFC layer and that, among the

proteins localized in the DFC layer, FBLs contributed most significantly to the localization

of the terminal regions of nascent pre-rRNAs in the DFC layer (13). Motivated by this

result, we take into account only the FBL-binding terminal regions of nascent pre-rRNAs

as arcRNAs that scaffolds the DFC layers and only FBLs as RBPs that bind to these

regions (see also the Discussion section). Recent experiments also revealed that the

upstream half of ETSs are localized at the DFC layer and the downstream half is

localized at the FC (13). This motivated us to assume that the terminal regions are

spanning from the top to the bottom of the DFC layer. Some RBPs bind to the nascent

pre-rRNAs and the others diffuse freely in the DFC layer (shown by magenta particles in

Fig. 1B). The interface between FC and DFC is located at a distance r_{in} from the center

and the interface between DFC and GC is located at a distance r_{ex} from the center (see

Fig. 1B). For simplicity, we assume that all the FCs have equal volume and the sum of

the volumes of microphases is fixed to V_{m}. This assumption does not seem so far off

although it has not been quantified (19).

The number of nascent pre-rRNAs at the surfaces of FCs is determined by the kinetics
of transcription and RNA processing. We treat the transcription dynamics of pre-rRNA by
using an extension of the model used by Stasevich and coworkers (20) (Fig. 2). Leaving
the kinetic equations to Materials and Methods, we here discuss the processes of
transcription and RNA processing that are taken into account in our model, see eqs. 6-
10. Pol I in an FC binds to the transcription start site (TSS) of a transcriptionally active
rDNA unit. The Pol I bound to the TSS starts transcription (Fig. 2A) or returns to the FC
without starting transcription (Fig. 2A). The rate with which the Pol I bound to the TSS
starts transcription is smaller than the rate with which this Pol I returns to the FC without
starting transcription. The Pol I polymerizes a pre-rRNA while it moves uni-directionally
towards the transcription termination site (TTS). The nascent pre-rRNAs are subject to
co-transcriptional RNA processing (Fig. 2B). The FBL-binding terminal region is located
at the ETS of pre-rRNA and is cleaved co-transcriptionally by endoribonuclease. The
average time between the transcription initiation and the cleavage of the FBL-binding
region is τ_{pr}, where this time includes the elongation time to the cleavage site and the
time necessary for the enzymatic reaction, see Fig. 2B and the Discussion section. Pol I
reaches TTS at the average elongation time τ_{e} and is then released to the FC ($\tau_{\mathrm{pr}} <$
τ_{e}), see Fig. 2C. This model is designed to focus on the transcription of the FBLs of
nascent pre-rRNAs and to treat the rest of the process as simple as possible. The kinetic
equations of the Pol I transcription and co-transcriptional processing are derived by
taking into account the above-mentioned processes, see Fig. 2 and eqs. 6-8. By solving
these kinetic equations for the steady state, we found that the surface density σ_{in} is
proportional to the radius r_{in} of FCs, see eq. 11 in Materials and Methods. We thus
introduce a dimensionless parameter

$$\zeta = N_{\mathrm{r}} \frac{\sigma_{\mathrm{in}} b^3}{r_{\mathrm{in}}}, \qquad [1]$$

which is proportional to the transcription rate and processing time τ_{pr}, while it is
independent of the radius r_{in}.

The stability of the system is quantified by the free energy. The free energy F_{d} of a DFC
layer has 4 contributions: 1) the elastic free energy f_{ela} of nascent pre-rRNA, 2) the
mixing free energy f_{mix} of RBPs and solvent, 3) the interaction free energy f_{int}
between RBPs (both freely diffusing in the DFC layer and bound to nascent pre-rRNA),
and 4) the binding free energy f_{bnd} of RBPs to the pre-rRNA terminal regions. The
expressions of these free energy contributions are shown in eqs. 12–17 in Materials and

Methods. The elastic free energy f_{ela} of flexible polymer chains, such as nascent pre-
rRNAs, increases as the chains stretch because the number of possible conformations
decreases (9). The thermal fluctuation mixes the components (nascent pre-rRNA units,
RBPs, and solvent) and this contribution is quantified as the mixing free energy f_{mix}.
The interaction free energy f_{int} is the free energy contribution of the multivalent
interactions between RBPs. The binding free energy f_{bnd} represents the free energy
gain due to the binding of RBPs to nascent pre-rRNA units and the thermal fluctuations
that dissociate RBPs from nascent pre-rRNA units. The free energy F_{d} takes into
account the binding of RBPs to the terminal regions of nascent pre-rRNAs and the
spherical geometry of the system in an extension of the Alexander model of polymer
brush (21).

The free energy F of the system has the form

179
$$\frac{F}{V_{\text{m}}} = \frac{3}{4\pi r_{\text{in}}^3}\left[F_{\text{d}} + 4\pi r_{\text{in}}^2 \gamma_{\text{in}} + 4\pi r_{\text{ex}}^2 \gamma_{\text{ex}}\right], \qquad [2]$$

where γ_{in} is the interfacial energy per unit area (interfacial tension) at the interface
between FC and DFC ($r = r_{\text{in}}$) and γ_{ex} is the surface energy per unit area (interfacial
tension) at the interface between DFC and GC ($r = r_{\text{ex}}$). The external radius r_{ex} is
determined by the conservation of the number of RNA units in the DFC layer in the steady
state, see eq. 11 in the Materials and Methods. The interfacial tension results from the
interactions between the components of interfacing phases. We therefore assume that
the surface tensions, γ_{in} and γ_{ex}, are proportional the local volume fraction of RBPs at
the interfaces, with the proportional coefficient γ_{p} (see eqs. 18 and 19 in the Materials
and Methods). The free energy F_{d} is a functional of the occupancy α_{p} of the pre-rRNA
terminal regions by RBPs, the volume fraction ϕ_{p} of RBPs freely diffusing in the DFC
layer, and the volume fraction ϕ_{r} of the FBL units, where these are functions of the
distance r from the center of the FC.

In our theory, we take into account the non-equilibrium nature of the system via nascent
pre-rRNAs produced at the surfaces of FCs by Pol I transcription. One may also think
that the continuous addition of RNA units to nascent pre-rRNAs makes the conformations
of their terminal regions different from those at the equilibrium. In a simple estimate, the
local equilibrium approximation that uses the equilibrium chain conformations is effective
in the length scales smaller than the diffusion length λ_{d} ($= \sqrt{D_{\text{r}}\tau_{\text{s}}}$), where D_{r} is the
diffusion constant of a pre-rRNA unit and τ_{s} is the elongation time to produce a pre-
rRNA unit. The diffusion length λ_{d} is estimated as $\sim 3\ \mu\text{m}$ (see Table 1) and is larger

than the typical thickness, $\sim 200\,\mathrm{nm}$, of the DFC layer: the local equilibrium

approximation is effective to treat the conformation of the pre-rRNAs in the DFC layer.

The occupancy α_p, the volume fractions ϕ_p and ϕ_r of RBPs and nascent pre-rRNA,

and the radius r_in thus turn over towards the minimum of the free energy, eq. 2.

Parameter	Meaning	Value	Ref.
b	Length of a RNA unit	$4\,\mathrm{nm} \approx 12\,\mathrm{b}$	22
η_w	Water viscosity	$1\,\mathrm{mPa \cdot s}$	
ν_e	Elongation rate	$6\,\mathrm{kb/min}$	23,24
T	Absolute Temperature	$300\,\mathrm{K}$	

204   Table 1 The values of the parameters used to estimate the diffusion length λ_d. The

205   diffusion constant of a RNA unit is estimated as $D_r \sim \frac{k_\mathrm{B}T}{4\pi\eta_\mathrm{w}b} \sim 8 \times 10^{-11}\,\mathrm{m^2/s}$ and the

206   time to synthesize a RNA unit is estimated as $\tau_\mathrm{s} = b/\nu_\mathrm{e} \sim 0.12\,s$ by using the values

207   listed in this table. We estimated b from the persistence length of single-stranded DNA.

Parameter estimate suggests that the terminal regions of nascent pre-rRNAs are
highly stretched

The independent parameters involved in our theory and their estimates are summarized

in Table 2. The kinetic constants involved in the Pol I transcription are included in ζ, see

eq. 1. The energy gain due to the binding of RBPs to the pre-rRNA terminal regions is

represented by $\epsilon k_\mathrm{B}T$. We represent the magnitudes of the multivalent interaction

between RBPs by using the Flory interaction parameter χ. We neglect the dependence

of the magnitudes of the interaction between RBPs on their binding state and also

assume that solvent molecules and naked pre-rRNA units are equivalent in terms of the

magnitudes of the interaction. This highlights the fact that the multivalent interaction

between FBLs drives phase separation *in vitro*, not to lose the essence by introducing

many interaction parameters. The volume fraction ϕ_p0 of RBPs in the nucleoplasm is

taken into account via the chemical potential $\mu_\mathrm{p} = k_\mathrm{B}T \log \phi_\mathrm{p0}$.

The interaction parameter χ and the binding energy ϵ are important quantities of

polymer physics, but are not experimentally well characterized. In the limit of vanishing

volume fraction of nascent RNAs, $\sigma_{\text{in}} \to 0$, our model returns to the theory of the binary
mixture of RBPs and solvent molecules (25), see eqs. 3 and 4. The latter theory predicts
that the phase separation happens at the threshold interaction parameter χ_{tr0} (=
$-\mu_{\text{p}}/(k_{\text{B}}T)$). FBL molecules show phase separation *in vitro* with the physiological
concentration (10), implying that $\chi > \chi_{\text{tr0}}$. If FBLs bind to the pre-rRNA terminal regions
even in a dilute solution, it implies that $\epsilon > -\mu_{\text{p}}/(k_{\text{B}}T)$. The size of the FBL of the
terminal region of a nascent pre-rRNA is $\approx 30\,\text{nm}$ in the relaxed state ($\approx 130\,\text{nm}$ in the
maximally stretched state), while the thickness of a DFC layer is $\approx 100\,\text{nm}$, implying that
FBLs are highly stretched.

Parameter	Meaning	Estimate	Value	Ref
ζ	Rescaled pre-rRNA surface density	< 0.12	Varied	13
$\dfrac{\gamma_{\text{p}} b^2}{N_r k_{\text{B}} T}$	Rescaled surface tension per RBP	~ 0.03	0.03	
χ	Interaction parameter	> 10	12	
ϵ	RBP-RNA binding energy	> 10	-12	
$\mu_{\text{p}}/(k_{\text{B}}T)$	RBP chemical potential	-10	-10	10

Table 2 List of independent (dimensionless) parameters involved in our theory. Estimate
of the unit length b of pre-rRNA and the absolute temperature are shown in Table 1. The
surface density σ_{in} was estimated by the number of Pol I (50 Pol I per rDNA, 300 copies
of active rDNA repeat per cell, ~ 10 FCs per nucleolus, ~ 2 nucleoli per cell) and the
typical radius of FCs $r_{\text{in}} \sim 100\,\text{nm}$ (3,13). The surface tension γ_{p} can be estimated as
$\sim k_{\text{B}}T/b_{\text{p}}^2$ (25,26), where the size b_{p} of FBL is $\approx 4\,\text{nm}$ (27). The number of units N_r in
the FBL-binding region of pre-rRNA is estimated by using its length $\sim 400\,b$ (13). The
chemical potential of FBL was estimated by using its concentration $\sim 1\,\mu\text{M}$ (10).

The multivalent interaction between FBLs localizes these RBPs to the DFC layer
and stretches the terminal regions of nascent pre-rRNAs

The occupancy α_{p}, the volume fractions ϕ_{p} and ϕ_{r} of RBPs and nascent pre-rRNA at
the free energy minimum is derived by using the variational principle, see sec. S1B in
the SI Appendix. For cases in which the curvature of the surfaces of FCs is not negligible,

α_p, ϕ_p, and ϕ_r are not uniform in the DFC layer. To avoid complexity arising from the
geometry of the system, we first analyze the occupancy α_p and the volume fractions,
ϕ_p and ϕ_r, in a DFC layer on a planer surface. This is indeed the limit of large FCs and
the essential feature of the assembly of DFC layers is included in this analysis. In the
planer geometry, the chemical potential of RBPs and the osmotic pressure is uniform in
the DFC layer.

The magnitude of the multivalent interactions between RBPs (either bound to the pre-
rRNA terminal regions or freely diffusing) increases with increasing the interaction
parameter χ. Although it is not straightforward to experimentally control the interaction
parameter χ of FBLs, it is instructive to analyze the composition of the layer occupied
by the FBLs of nascent pre-rRNAs as a function of the interaction parameter χ, see Fig.
3. For cases in which the interaction parameter χ is small (which corresponds to self-
association defective FBL ΔIDR mutant (13)), the major component of the layer is solvent
(nucleoplasm), see the cyan line in Fig. 3. For $\chi < \chi_\mathrm{s}$, the excluded volume interaction
between nascent pre-rRNA units dominates the attractive interaction between RBPs
bound to these nascent pre-rRNAs (good solvent regime). The interaction parameter χ_s
at the crossover is 2 in the limit of $\epsilon \ll \mu_\mathrm{p}/(k_\mathrm{B}T)$, such as the case of Fig. 3, see also eq.
S31 in the SI Appendix for the general expression. The inverse of the volume fraction ϕ_r
of nascent pre-rRNA is proportional to the extent of the stretching of the terminal regions
of nascent pre-rRNAs. The terminal regions of these RNAs are somewhat stretched for
$\chi \approx 0$ due to the excluded volume interaction between the pre-rRNA units (good solvent
regime). The terminal regions shrink as the magnitude χ of the attractive interaction
between RBPs bound to the terminal regions of nascent pre-rRNAs, see the green line
in Fig. 3. For $\chi_\mathrm{s} < \chi < -\mu_\mathrm{p}/(k_\mathrm{B}T)$, the attractive interaction dominates the excluded
volume interaction (poor solvent regime) and the terminal regions are collapsed with
RBPs bound to these regions (melt regime), see the green line in Fig. 3.

At a threshold interaction parameter, $\chi_\mathrm{th} \approx -\mu_\mathrm{p}/(k_\mathrm{B}T)$, the volume fraction ϕ_p of freely
diffusing RBPs jumps to a large value, see the magenta line in Fig. 3. In contrast, the
volume fraction of the solvent (nucleosol) jumps to almost zero, see the cyan line in Fig.
3. The feature of the layer for $\chi > \chi_\mathrm{th}$ is analogous to the DFC layer observed
experimentally (which we thus call DFC regime). The volume fraction ϕ_r of nascent pre-
rRNA units jumps to a small value at $\chi \approx \chi_\mathrm{th}$, see the green line in Fig. 3, implying that
the terminal regions of the nascent pre-rRNAs are stretched at the transition. Because
the volume fraction ϕ_r is smaller than the case of $\chi = 0$, the stretching is not solely due
to the fact that RBPs in the DFC layer behave as an athermal solvent to the RNP

complexes of RBPs and the pre-rRNA terminal regions. The terminal regions stretch so
that the DFC layer can accommodate more RBPs to decrease the free energy due to the
multivalent interactions between them. The stretching thus may be a transport
mechanism of the ETS of pre-rRNAs towards the DFC layer.

The lateral pressure generated by nascent pre-rRNAs suppresses the growth of
FCs

Now we discuss the DFC layers at the surfaces of spherical FCs to derive the radius of
FCs in the steady state. The first (functional) derivatives of the free energy with respect
to α_p, ϕ_p, and ϕ_r are zero in the steady state. In the spherical geometry, these
conditions correspond to the fact that the chemical potential of RBPs and the chemical
potential of pre-rRNA units, instead of the osmotic pressure, are uniform, see sec. S1B
for the mathematics of the free energy minimization. We treat the DFC regime in which
the volume fraction of solvent molecules is negligibly small and the occupancy α_p of the
pre-rRNA terminal regions is approximately unity, see Table 2 and Fig. 3. We performed
numerical and analytical calculations within these conditions, see sec. S3A in the SI
Appendix.

In the spherical geometry, the volume fraction ϕ_r of pre-rRNA units is not uniform in the
DFC layer and decreases with the distance r from the center ($r_\mathrm{in} < r < r_\mathrm{ex}$). It is
because the elastic stress (proportional to the number of pre-rRNAs per unit area)
generated by pre-rRNA decreases with increasing r, see the solid green line in Fig. 4.
We performed the asymptotic analysis for $\phi_\mathrm{r} \ll 1$ by expanding the chemical potentials
of proteins and pre-rRNA units in a power series and neglected higher order term of ϕ_r.

This leads to an asymptotic expression, $\phi_\mathrm{r} \propto \frac{1}{r^2}$, which reasonably explains the

numerical calculation, see the solid dark green line and broken light green line in Fig. 4
and also eq. S55 in the SI Appendix for the explicit form. The rest of the space in the
DFC layer is filled by RBPs, see the inset of Fig. 4.

We used the profile of the pre-rRNA units to derive the free energy as a function of the
radius r_in of FCs. For cases in which the Pol I transcription is inhibited, the free energy
of the system is a monotonically decreasing function of the radius r_in, implying that, the
radius of FCs increases with time by coarsening or coalescence. For cases in which the
Pol I transcription is active, the free energy generated by the terminal regions of the
nascent pre-rRNA is an intriguing function of r_in, which has a minimum, see the solid
black line in Fig. 5A. Indeed, the surface free energy, the second and third terms of eq.

2, is not significant for the values of parameters estimated in Table 2, see the broken
cyan line in Fig. 5A. In such case, the free energy is simpler if it is shown as a function
of the ratio r_{ex}/r_{in}, see Fig. 5B and eq. S59 in the SI Appendix.

The first derivative of the free energy with respect to the radius r_{in} is zero at the
minimum of the free energy,

318
$$\Delta\Pi_{\parallel} \approx \gamma_{in} + \gamma_{ex}\frac{r_{ex}^2}{r_{in}^2}. \qquad [3]$$

319   Eq. 3 represents the surface tensions, γ_{in} and γ_{ex}, are balanced to the lateral osmotic
320   pressure $\Delta\Pi_{\parallel}$ generated in the DFC layer (the complete form is shown in eq. 21). The
321   radius r_{in} at the minimum of the free energy decreases with increasing the transcription
322   level, see Fig. 6. The radius r_{in} at the minimum of the free energy has an asymptotic
323   form

324
$$\frac{r_{in}}{bN_r} = \frac{2}{\sqrt{3}}\left(\chi + \frac{\mu_p}{k_B T}\right)^{\frac{1}{2}}\zeta^{-\frac{1}{2}}, \qquad [4]$$

see also the broken line in Fig. 6. Eq. 4 is derived by expanding the free energy in a
power series of ϕ_r and by using the asymptotic form of ϕ_r, see sec. S3A in the SI
Appendix for the details of the derivation. We also neglected the surface free energy, the
second and third terms of eq. 2, because it is not significant with our estimate in Table 2.
Our theory therefore predicts that the radius of FCs decreases with the inverse of the
square root of the transcription rate. At the minimum of the free energy, the ratio of the
radii is $\frac{r_{ex}}{r_{in}} \approx 2$, which roughly agrees with the fact that the radius of FCs and the
thickness of DFC layers are the same order of length scales, $\propto 100\,\text{nm}$, see Fig. 5B.

Mild inhibition of Pol I increases the size of FCs

The main prediction from our theoretical model is that the size of FCs increases by
reducing the expression level of nascent pre-rRNAs. To experimentally test this
prediction, we used BMH-21 and CX-5461, specific Pol I inhibitors, which reduce nascent
pre-rRNA levels in a dose-dependent manner (28,29) (Fig. S1A and B). In untreated cells,
small foci of UBF (a marker for FCs) and FBL (a marker for DFC) proteins were dispersed
within the nucleoli as previously reported (11) (Figs. 7A, S2A, S2B, and S3A). In contrast,
UBF and FBL proteins were relocalized to nucleolar caps in the presence of high doses

of BMH-21 (0.5 μM) or CX-5461 (2 μM), as reported (11,28) (Figs. 7A, S2A, S2B, and S3A). Strikingly, the FCs were larger in cells treated with the medium doses of BMH-21 (0.0625, 0.125, and 0.25 μM) or CX-5461 (0.25, 0.5, and 1 μM) than in untreated cells (Figs. 7A, S2A, S2B, and S3A). We then quantified the size of the FCs under these untreated and mildly treated conditions (Figs. 7B, 7C, S3B, S3C, and S4). Indeed, the longest axis (Lx) and area of the FCs increase with increasing the dose of BMH-21 and CX-5461. As theoretically predicted, the Lx and area of the FCs increase according to pre-rRNA level reduction (Figs. 7D, 7E, S3D, and S3E).

FC radius is a power function of the pre-rRNA transcription level with an exponent of approximately -0.5

To compare our theory and experiments more quantitatively, we curve-fit the square root of the area of FCs as a function of the transcription level by a power function (Fig. 8). The scaling exponent that describes the dependence of the square root of the area of FCs on the transcription level is a universal quantity that does not depend on specific values of the parameters. Checking such scaling exponents with experiments is the first thing to do to test theories in polymer physics (30). By fitting the data with a power function, we found that the scaling exponent is -0.49 for the case of the BMH-21 treatment, -0.46 for the case of the CX-5461, and -0.48 if both data are fitted. The scaling exponent predicted by our theory -0.5 reasonably agrees with our experiments, see eq. 4.

DISCUSSION

Our theory predicts that the terminal regions of nascent pre-rRNAs at the surfaces of FC microphases in a nucleolus generate the lateral osmotic pressure and suppress the growth of microphases. Our theory predicts that the radius of FCs is proportional to the inverse of the square root of the transcription rate. This prediction is consistent with our experiments. The parameter ζ is also proportional to the inverse of the rate τ_{pr}^{-1} of co-transcriptional processing that cleaves the terminal regions of pre-rRNA. Our theory thus predicts that the radius of FCs is proportional to the square root of the processing rate τ_{pr}^{-1}. The inhibition of processing induced by ribosomal protein depletion, such as uL18 (RPL5) and uL5 (RPL11), changes the multiphase structure of nucleoli (31). Indeed, recent experiments have demonstrated that long pre-rRNAs are accumulated in RPL5 knockdown cells and the size of FCs in such cells is smaller than that in the wild type (32). The consistency between our theory and experiments suggests that the growth of FCs is suppressed by the terminal regions of nascent pre-rRNAs localized at the

surfaces of FCs. This implies that the elastic energy of the terminal region is the driving
force of the transport of the 5' ETSs cleaved by RNA processing towards GC.

In many cases, the multivalent interaction between complexes of RNA and RBPs drives
the growth of condensates (33,34) (Fig. 8A). The size of such condensates increases
with increasing the transcription rate (14,17,35). In contrast, for the case of FCs in
nucleoli, the multivalent interaction between the complexes of nascent RNA and RBPs
rather suppresses the growth of FCs (11). The size of FCs indeed decreases with
increasing the transcription rate. Our theory predicts that the suppression of the growth
of FCs by transcription results from the fact that pre-rRNAs are end-grafted to the
surfaces of FCs via Pol I (Fig. 8B).

Our theory predicts that the function of the terminal regions of nascent pre-rRNAs is
analogous to surfactants, such as lipids. If the surfactants at an interface between two
immiscible liquids are dense enough to form a monomolecular liquid film, these
surfactants decrease the interfacial tension to almost zero to disperse multiple droplets
even in the thermodynamic equilibrium (microemulsion) (25,36). The surfactant
monolayer acts as a kinetic barrier to suppress the coalescence and decreases the
hydrostatic pressure of the droplet to suppress the coarsening (9,25). These features are
analogous to the terminal regions of pre-rRNAs at the surfaces of FCs. The fact that
some RNAs and proteins can act as surfactants was discussed in recent researches
(37,38). There are also differences between nascent pre-rRNA and surfactants. First and
foremost, surfactants are localized at interfaces at the thermodynamic equilibrium, while
the terminal regions of nascent pre-rRNAs are localized at the FC-DFC interfaces only
during the transcription and their surface activity is regulated by Pol I transcription
dynamics. Second, the hydrophobic chains of surfactants in a liquid monolayer tend to
form a melt, while the terminal regions of pre-rRNA are extended due to the RBPs freely
diffusing in the DFC layers. Last, but not least, surfactants can freely diffuse in a
monolayer, while the terminal regions of nascent pre-rRNAs are anchored to the rDNA
at the interface via Pol I and thus are not mobile.

In our model, we focused on FBLs because these RBPs contribute most significantly to
the localization of the terminal regions of pre-rRNAs to the DFC layer (13). FRAP
experiments on the condensates of FBLs reconstituted *in vitro* suggest that these
condensates are thixotropic (and probably also be viscoelastic) and become gels in a
long timescale (10). Indeed, recent experiments showed that long non-coding RNA
SLERT facilitates the transition of the RNA helicase DDX21 to the closed conformation

and ensures the fluidity of proteins in the DFC layer (39). This mechanism ensures the
validity of our assumption that unbound FBLs can diffuse freely in the DFC layer,
although we did not take into account these factors explicitly.

Although our theory probably could capture the essential features that are necessary to
understand the assembly mechanism of the multiphase structure of nucleolus, it is not
complete. First, the layers at the interface between FCs and GC are not uniform, but
DFC regions are flanked by the regions, in which FBLs are not observed by super-
resolution microscopy and are not visible by electron microscopy. Second, we did not
take into account the downstream pre-rRNA regions that are not bound by FBLs. These
downstream regions were observed at the surfaces of FCs (13). The direct contribution
of rDNAs localized at the surfaces of FCs are not significant, see S4 in the SI Appendix.
Third, we did not explicitly take into account the transport dynamics of pre-rRNAs in GCs.
Fourth, RNAs can be not only cleaved, but also can be folded and/or chemically modified
during the co-transcriptional processing. The processing of pre-rRNAs in yeast has been
well-documented (40,41), but much less in higher eukaryotes (42,43), in which the
multiphase structure of nucleolus is usually observed. It is of interest to understand the
mechanism of the processed pre-rRNAs, which have the affinity to FCs, penetrate
through the DFC layer towards the GC. One possible hypothesis is that the processed
pre-rRNAs pass the flanking regions of the DFC layer. These flanking regions increase
the density of the terminal regions of pre-rRNAs in the DFC compartments and enhance
the lateral osmotic pressure. The enhanced lateral pressure and/or the kinetic trapping
of the coalescence of FCs by nascent pre-rRNAs may explain the fact that the radius of
FCs observed in experiments is smaller than that predicted by our theory. Whether the
multiphase structure of the nucleolus contributes to its function is not yet clear. It is of
interest to extend our theory to understand the structure-function relationship of the
nucleolus (44). However, following the approach of the classical theoretical physics, our
present theory should be experimentally tested before adding more assumptions that
have not been experimentally explored.

The key assumption of our theory is that the terminal regions of nascent pre-rRNAs span
between the top to the bottom of the DFC layer. This is motivated by a recent experiment
that the upstream half of ETSs are localized at the DFC layer and the downstream half
is localized at the FC (13). The stretching of the terminal regions results from the
multivalent interactions between FBLs and thus our theory should be tested in the
condition of high FBL volume fraction. In the conventional picture, the DFC layer is
composed of 5' ETSs that are already cleaved and folded by RNA processing. Such

diffusive molecules tend to make the interaction between condensates attractive (the
depletion interaction) (25), however, the non-equilibrium nature of the transcription may
change the situation. We envisage a future theoretical study to predict the stable size of
FCs with the conventional picture to check if the latter agrees with existing experiments,
including ours.

Our theory may provide insight into the size control mechanism of other nuclear bodies.
The size of nuclear speckles (5) increases by the suppression of transcription. The
stability of transcriptional condensate depends on the frequency of transcription and the
length of transcripts (45,46). Transcriptional condensates are actively studied in recent
experiments (45-49). The simulations by Henninger and coworkers predict that RNAs
diffuse freely in the interior of condensates and destabilize the condensates due to the
electrostatic repulsion (45). In contrast, Hilbert and coworkers argued that DNA-Pol II-
RNA complexes are localized at the interface between transcriptional condensates (rich
in RNA and proteins) and the exterior (rich in DNA) and act as surfactants that suppress
the coalescence and coarsening of the condensates (46), analogous to microemulsion
(see above). Our theory is somewhat similar to the idea of Hilbert and coworkers, but
different in the fact that the complexes of nascent pre-rRNAs and RBPs form distinct
phase at the interface between FCs and GC due to the transcription and FCs and GC
are somewhat similar in terms of the relative affinity to the nucleoplasm, as characterized
by the interfacial tension (10). The Pol II transcription is regulated by the phosphorylation
of the C-terminal domains and is somewhat different from the Pol I transcription. It is of
interest to take into account such differences in an extension of our present theory to
understand the assembly mechanism of nuclear bodies and their functions.

The multiphase structure of nucleoli may be somewhat analogous to that observed in
nuclear stress bodies (nSBs) by electron microscopy (50). nSBs are assembled during
the thermal stress condition and are composed of HSATIII architectural RNA and specific
RBPs (51). HSATIII is transcribed by Pol II from the pericentromeric Satellite III regions,
which are enriched in tandem repeats (52,53). These features common with nucleoli
motivate us to think of a general mechanism involved in the assembly of the multiple
subcompartments. The nascent RNAs, which are produced by transcription and are
modulated by RNA processing, are important elements that regulate the multiphase
structures, and possibly their functions, of nuclear bodies.

MATERIALS AND METHODS

Transcription and RNA processing

The occupancy n_{ps} of the TSS of the active rDNA by Pol I follows the kinetic equation

$$\frac{d}{dt}n_{\text{ps}} = k_{\text{on}}\rho(1 - n_{\text{ps}}) - k_{\text{off}}n_{\text{ps}} - k_{\text{e}}n_{\text{ps}}. \qquad [6]$$

Eq. 6 suggests that the occupancy n_{ps} changes due to the binding of Pol I from the FC
(the first term), the unbinding of Pol I from the TSS to the FC without starting transcription
(the second term), and the initiation of the elongation of Pol I (the third term), see Fig.
2A. k_{on} is the rate constant for the binding of Pol I to TSS and ρ is the concentration
of Pol I in a FC (which will be determined later). k_{off} is the rate constant for the unbinding
of Pol I from TSS. k_{e} is the rate of the transition of Pol I from the bound state to the
elongation state.

The number n_{pe} of elongating Pol I protein complexes with pre-processed nascent pre-
rRNA follows the kinetic equation

$$\frac{d}{dt}n_{\text{pe}} = k_{\text{e}}n_{\text{ps}} - \frac{n_{\text{pe}}}{\tau_{\text{pr}}}. \qquad [7]$$

Eq. 7 suggests that the number n_{pe} changes due to the release of Pol I from TSS for
elongation (the first term) and the cleavage of the FBL-binding terminal region of nascent
pre-rRNA (the second term), see Fig. 2B. τ_{pr} is the average time between the
transcription initiation and the cleavage of the terminal regions of the pre-rRNA.

The number n_{p0} of elongating Pol I with processed nascent pre-rRNA follows the kinetic
equation

$$\frac{d}{dt}n_{\text{p0}} = \frac{n_{\text{pe}}}{\tau_{\text{pr}}} - \frac{n_{\text{p0}}}{\tau_0}. \qquad [8]$$

Eq. 8 suggest that the number n_{p0} changes due to the the cleavage of the terminal
regions of pre-rRNAs (the first term) and the transcription termination (the second term),
see Fig. 2C. τ_0 is the average time between the cleavage of FBL-binding region and the
transcription termination.

The numbers, n_{ps}, n_{pe}, and n_{p0}, of Pol I in each state are derived for the steady state,
$\frac{dn_{\text{ps}}}{dt} = \frac{dn_{\text{pe}}}{dt} = \frac{dn_{\text{p0}}}{dt} = 0$, by using eqs. 6-8. The inverse time k_{e} of the transcription
initiation is much smaller than the inverse k_{off} of the unbinding of Pol I from TSS, $k_{\text{e}} \ll$

k_{off}. The concentration ρ of Pol I in a FC is determined by the condition

$$(n_{\mathrm{ps}} + n_{\mathrm{pe}} + n_{\mathrm{p0}})N_{\mathrm{a}} + \rho V_{\mathrm{m}} = N_{\mathrm{pol}} \qquad [9]$$

that the sum of the number of Pol I during transcription (the first term) and the number of
Pol I diffusing in microphases (the second term) in the system is constant. All the
parameters involved in eq. 9 do not depend on the size of FCs, see eqs. 6-8. The
concentration ρ of Pol I in FCs thus does not depend on the size of FCs.

The surface density σ_{in} of the FBL-binding terminal regions of nascent pre-rRNAs at
each FC (the number of nascent pre-rRNAs per unit area of each FC surface) is
represented as

$$\sigma_{\mathrm{in}} = \frac{n_{\mathrm{pe}}}{4\pi r_{\mathrm{in}}^2}\left(\frac{4\pi r_{\mathrm{in}}^3}{3V_{\mathrm{m}}}N_{\mathrm{a}}\right) = \frac{1}{3}k_{\mathrm{e}}\tau_{\mathrm{pr}}\frac{\rho}{\rho + K_{\mathrm{pl}}}\frac{N_{\mathrm{a}}}{V_{\mathrm{m}}}r_{\mathrm{in}}. \qquad [10]$$

514   by assuming that the terminal regions of nascent pre-rRNA is distributed uniformly at the
515   surfaces of FCs. We note that $3V_{\mathrm{m}}/(4\pi r_{\mathrm{in}}^3)$ is the number of FCs in a nucleolus. The last
516   form of eq. 10 is derived by using the solution of eqs. 6-8.

Thickness of DFC layer

518   The exterior radius r_{ex} (see also Fig. 1B) is determined by the relationship

$$\int_{r_{\mathrm{in}}}^{r_{\mathrm{ex}}} dr\, 4\pi r^2 \phi_{\mathrm{r}} = 4\pi r_{\mathrm{in}}^2 \sigma_{\mathrm{in}} N_{\mathrm{r}}, \qquad [11]$$

where N_{r} is the average number of units in the terminal region of a nascent pre-rRNA.
It is a mean field treatment that assumes that nascent RNAs are composed of the same
number of units and is effective within the Alexander-de Gennes approximation, with
which the brush height is determined by the distance between neighboring grafting points
and the average number of units per chain (if one neglects the fact that the lateral
fluctuations of a chain composed of N_{r} units is limited to $\sim N_{\mathrm{r}}^{1/2}b$) (21, 54).

Free energy of DFC layer

527   The free energy of each DFC layer has the form

$$F_{\mathrm{d}} = \int_{r_{\mathrm{in}}}^{r_{\mathrm{ex}}} dr\, \frac{4\pi r^2}{b^3} f_{\mathrm{d}} \qquad [12]$$

529   with

$$f_{\mathrm{d}} = f_{\mathrm{ela}} + f_{\mathrm{mix}} + f_{\mathrm{int}} + f_{\mathrm{bnd}} - \mu_{\mathrm{p}}(\phi_{\mathrm{p}} + \alpha_{\mathrm{p}}\phi_{\mathrm{r}}) + \Pi_{\mathrm{ex}}b^{3}. \qquad [13]$$

Eq. 12 is the functional of the local volume fraction ϕ_{r} of nascent pre-rRNA units, the
local volume fraction ϕ_{p} of RBPs, and the occupancy α_{p} of nascent pre-rRNA units by
RBPs, which are functions of the distance r from the center of the FC ($r_{\mathrm{in}} < r < r_{\mathrm{ex}}$),
see Fig. 1B. b is the length of a pre-rRNA unit. This free energy is composed of 4
contributions: f_{ela} is the elastic free energy density of the terminal regions of nascent
RNAs, f_{mix} is the free energy density due to the mixing entropy of RBPs and solvent
molecules, f_{int} is the free energy density due to the interactions between RBPs, and
f_{bnd} is the free energy density due to the binding of RBPs to the terminal regions of
nascent RNAs. μ_{p} is the chemical potential of RBPs and Π_{ex} is the osmotic pressure
from the exterior.

The (entropic) elastic free energy density has the form

$$\frac{f_{\mathrm{ela}}}{k_{\mathrm{B}}T} = \frac{3}{2}\frac{b^{4}\sigma^{2}(r)}{\phi_{\mathrm{r}}}, \qquad [14]$$

where $\sigma(r)$ is the surface density of FBL-binding regions of nascent pre-rRNAs. $\sigma(r) =$
$\sigma_{\mathrm{in}}r_{\mathrm{in}}^{2}/r^{2}$ for cases in which the surface of FCs is sphere and $\sigma(r) = \sigma_{\mathrm{in}}$ for cases in
which the surface of FCs is planer (which corresponds to the limit of $\frac{r_{\mathrm{ex}}-r_{\mathrm{in}}}{r_{\mathrm{in}}} \ll 1$). The
volume fraction ϕ_{r} of pre-rRNA units decreases as the terminal regions of the nascent
pre-rRNAs are stretched. Eq. 14 represents the fact that the elastic free energy f_{ela}
increases as these terminal regions are stretched. Eq. 14 takes into account the
spherical geometry of the system in an extension of the Alexander model (that assumes
that the concentration of the terminal regions of nascent pre-rRNA is uniform in the DFC
layer in the limit of $\frac{r_{\mathrm{ex}}-r_{\mathrm{in}}}{r_{\mathrm{in}}} \ll 1$) in the spirit of the Daoud-Cotton theory (55), see sec.
S5 in the SI Appendix for the derivation. k_{B} is the Boltzmann constant and T is the
absolute temperature.

The free energy due to the mixing entropy of RBPs and solvent molecules has the form

$$\frac{f_{\mathrm{mix}}}{k_{\mathrm{B}}T} = \phi_{\mathrm{p}}\log\phi_{\mathrm{p}} + \left(1 - \phi_{\mathrm{p}} - (1 + \alpha_{\mathrm{p}})\phi_{\mathrm{r}}\right)\log\left(1 - \phi_{\mathrm{p}} - (1 + \alpha_{\mathrm{p}})\phi_{\mathrm{r}}\right). \qquad [15]$$

556   The mixing free energy, eq. 15, has the contributions of the mixing entropy of RBPs (the

557   first term) and the mixing entropy of solvent molecules (the second term).

558   The free energy due to the interactions between RBPs has the form

559
$$\frac{f_{\text{int}}}{k_{\text{B}}T} = -\chi\left(\phi_{\text{p}} + \alpha_{\text{p}}\phi_{\text{r}}\right)^2,$$
[16]

where χ is the interaction parameter that accounts for the attractive interactions. For
simplicity, we assumed that RBPs bound to the terminal regions of nascent pre-rRNAs
are equivalent to RBPs freely diffusing in the DFC layer and that solvent molecules are
equivalent to nascent RNA units in terms of the interactions, see also the discussion in
Parameter estimate.

The free energy due to the binding of RBPs to nascent RNAs has the form

566
$$\frac{f_{\text{bnd}}}{k_{\text{B}}T} = \phi_{\text{r}}\left[\alpha_{\text{p}} \log \alpha_{\text{p}} + \left(1 - \alpha_{\text{p}}\right) \log\left(1 - \alpha_{\text{p}}\right) + \epsilon\alpha_{\text{p}}\right],$$
[17]

where $\epsilon k_{\text{B}}T$ is the energy increase due to the binding of RBPs to nascent RNAs. For
simplicity, we assumed that each nascent RNA unit has one binding site of RBPs.

Free energy of system

The free energy F of the system, eq. 2 is derived by using the fact that the number of
microphases in the system is $3V_{\text{m}}/(4\pi r_{\text{in}}^3)$ to derive this form. The interfacial tension
between GC and nucleosol is relative small (3,10) and FC is excluded out from the
nucleolus by the transcription inhibition (3,11), implying that the interfacial tension
between FC and nucleosol is even smaller. In vitro experiments suggest that the
attractive interaction between FBLs is much larger than that between nucleophosmins,
a major component of GC (10). It implies that the interfacial tension between DFC and
GC and that between DFC and FC are larger than their difference. By neglecting the
difference between these interfacial tensions, the interfacial tensions γ_{in} and γ_{ex} at the
interior and the exterior interfaces are represented by the forms

580
$$\gamma_{\text{in}} = \gamma_{\text{p}}\left(\phi_{\text{p}}(r_{\text{in}}) + \alpha_{\text{p}}(r_{\text{in}})\phi_{\text{r}}(r_{\text{in}})\right)$$
[18]

581
$$\gamma_{\text{ex}} = \gamma_{\text{p}}\left(\phi_{\text{p}}(r_{\text{ex}}) + \alpha_{\text{p}}(r_{\text{ex}})\phi_{\text{r}}(r_{\text{ex}})\right)$$
[19]

582 where γ_p is the surface tension between a nucleosol and a liquid of the RBPs and is
583 proportional to χ.

Lateral osmotic pressure

585 The lateral osmotic pressure $\Pi_\parallel$ can be defined by

$$\Delta\Pi_\parallel = \sigma_{in}^2 \frac{\partial}{\partial\sigma_{in}}\left(\frac{F_d}{4\pi r_{in}^2 \sigma_{in}}\right), \qquad [20]$$

587 where the derivative with respect to σ_0 is taken with the condition that $4\pi\sigma_{in}r_{in}^2$ is
588 constant (this increases the area of the surface of a FC without changing the composition
589 of the DFC layer). $\Delta\Pi_\parallel$ is the lateral osmotic pressure, from which the contribution of the
590 isotropic osmotic pressure is subtracted.

Radius of FCs at free energy minimum

592 The free energy, eq. 1, is represented as a function only of the radius r_{in} of FCs if the
593 occupancy, α_p, and the volume fractions, ϕ_p and ϕ_r, derived by using eqs. 2-5 are
594 substituted into eq. 1. The radius r_{in} at the steady sate is derived by the condition $\frac{dF}{dr_{in}} =$
595 0 with the condition that $\sigma_0 \propto r_{in}$ (this increases the area of the surfaces of FCs with
596 the optimization of the arrangement of the active rDNA repeats). This leads to the form

$$\Delta\Pi_\parallel - \left(\gamma_{in} + \gamma_{ex}\frac{r_{ex}^2}{r_{in}^2}\right) + r_{in}\left(\frac{\partial\gamma_{in}}{\partial r_{in}} + \frac{\partial\gamma_{ex}}{\partial r_{in}}\frac{r_{ex}^2}{r_{in}^2}\right) = 0, \qquad [21]$$

598 which represent the fact that the surface tension is balanced to the surface pressure
599 generated by the FBRs of nascent pre-rRNAs.

Cell culture, drug treatment, reverse transcription-quantitative PCR (RT-qPCR), immunofluorescence, and quantification of the size of FCs

HeLa cells were maintained in DMEM containing high glucose (Nacalai Tesque, Cat#
08458-16) supplemented with 10% FBS (Sigma) and Penicillin-Streptomycin (Nacalai
Tesque). Cells were treated with BMH-21 (Selleckchem, Cat# S7718) or CX-5461
(AdooQ Bioscience) for 3 hours. Quantification of nascent pre-rRNAs (forward primer:
5'-CCTTCCCCAGGCGTCCCTCG-3', reverse primer: 5'-
GGCAGCGCTACCATAACGGA-3') (11) by RT-qPCR was performed using LightCycler
480 II (Roche). GAPDH mRNAs were used as a loading control (14). The antibodies to

DATA AVAILABILITY

Mathematica file used for the numerical calculations are available in figshare with the identier (https://doi.org/10.6084/m9.figshare.16599446).

ACKNOWLEDGMENT

This research was supported by KAKENHI grants from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan [to   T. Yamamoto (20H05934, 21K03479, 21H00241), T. Yamazaki (21H00253, 22H02545), KN (19K06478, 22K06083),and TH (20H00448, 20H05377,21H05276)], JST, PRESTO Grant Number JPMJPR18KA (to T. Yamamoto), the Mochida Memorial Foundation for Medical and Pharmaceutical Research (to T. Yamazaki), the Naito Foundation (to T. Yamazaki), the Takeda Science Foundation (to T. Yamazaki), and JST CREST Grant Number JPMJCR20E6 (to T.H.). T. Yamazaki thanks Toyofumi Kameoka (NIKON SOLUTIONS CO., LTD.) and Takayuki Funato (NIKON SOLUTIONS CO., LTD.) for support of microscopic image analyses. T. Yamamoto acknowledges the fruitful discussion with

Noriko Saito (Japanese Foundation for Cancer Research), Yuma Ito (Tokyo Institute of Technology), Satoru Ide (National Institute of Genetics), Tsutomu Suzuki (Univ. of Tokyo), Takahiko Kobayashi (Univ. of Tokyo), Yutetsu Kuruma (JAMSTEC), Shintaro Iwasaki (Univ. of Tokyo), Sumio Sugano (Chiba University), Haruhiko Siomi (Keio Univ.), Hideaki Matsubayashi (Johns Hopkins Univ.), and Naomichi Takemata (Kyoto Univ.).

REFERENCE

1. S Klinge, J Woolford Jr, Ribosome assembly coming into focus. *Nat. Rev. Mol. Cell Biol.* **20,** 116–131 (2019).

2. B McStay, I Grummt, The Epigenetics of rRNA genes: From Molecular to Chromosome Biology. *Annu. Rev. Cell Dev. Biol.* **24,** 131–157 (2008).

3. D Lafontaine, J Riback, R Bascetin, C Brangwynne, The nucleolus as a multiphase liquid condensate. *Nat. Rev. Mol. Cell Biol.* **22,** 165–182 (2021).

4. J Berry, S Weber, N Vaidya, M Haataja, C Brangwynne, RNA transcription modulates phase transition driven nuclear body assembly. *Proc. Natl. Acad. Sci. USA* **112,** E5237–E5245 (2015).

5. A Lamond, D Spector, Nuclear speckles: a model for nuclear organelles. *Nat. Rev. Mol. Cell Biol.* **4,** 605–612 (2003).

6. S Nakagawa, T Yamazaki, T Hirose, Molecular dissection of nuclear paraspeckles: towards understanding the emerging world of the RNP milieu. *Open Biol.* **8,** 180150 (2018).

7. T Chujo, T Yamazaki, T Hirose, Architectural RNA (arcRNAs): A class of long noncoding RNAs that function as the scaffold of nuclear bodies. *Biochim. Biophys. Acta* **1859,** 139–146 (2015).

8. T Yamazaki, S Nakagawa, T Hirose, Architectural RNAs for Membraneless Nuclear Body Formation. *Cold Spring Harb. Symp. Quant. Biol.* **84,** 227–237 (2019).

9. M Doi, *Soft Matter Physics.* (Oxford University Press, Oxford, UK.), (2013).

10. M Feric, et al., Coexisting Liquid Phases Underlie Nucleolar Subcompartments. *Cell* **165,** 1686–1697 (2016).

11. S Ide, R Imai, H Ochi, K Maeshima, Transcriptional suppression of ribosomal DNA
with phase separation. *Sci. Adv.* **6,** eabb5953 (2020).

12. A Maiser, et al., Super-resolution in situ analysis of active ribosomal DNA chromatin
organization in the nucleolus. *Sci. Rep.* **10,** 7462 (2020).

13. R Yao, et al., Nascent Pre-rRNA Sorting via Phase Separation Drives the Assembly
of Dense Fibrillar Components in the Human Nucleolus. *Mol. Cell* **76,** 767–783 (2019).

14. T Yamazaki, et al., Paraspeckles are constructed as block copolymer micelles.
EMBO J **40,** e107270 (2021).

15. S Souquere, G Beauclair, F Harper, A Fox, G Pierron, Highly Ordered Spatial
Organization of the Structural Long Noncoding Neat1 RNAs Within Paraspeckle Nuclear
Bodies. *Mol. Biol. Cell* **21,** 4020–4027 (2010).

16. J West, et al., Structural, super-resolution microscopy analysis of paraspeckle
nuclear body organization. *J. Cell Biol.* **214,** 817–830 (2016).

17. T Yamamoto, T Yamazaki, T Hirose, Triblock copolymer micelle model of spherical
paraspeckles. *Front. Mol. Biosci.* **9,** 925058 (2022).

18. T Hirose, et al., NEAT1 long noncoding RNA regulates transcription via protein
sequestration within subnuclear bodies. *Mol. Biol. Cell* **25,** 169–183 (2014).

19. P Tchelidze, A Benassarou, H Kaplan, MF O'Donohue, L Lucas, C Terryn, L Rusishvili,
G Mosidze, N Lalun D. Ploton, Nucleolar sub-compartments in motion during rRNA
synthesis inhibition: Contraction of nucleolar condensed chromatin and gathering of
fibrillar centers are concomitant. PLOS ONE **12,** e0187977 (2017).

20. T Stasevich, et al., Regulation of RNA polymerase II activation by histone acetylation
in single living cells. *Nature* **516,** 272–275 (2014).

21. S Alexander, Adsorption of chain molecules with a polar head a scaling description.
J. Phys. France **38,** 983–987 (1980).

22. E Roth, A G Azaria, O Girshevitz, A Bitler, and Y Garini, Measuring the conformation
and persistence length of single-strand DNA using a DNA origami structure. *Nano Lett.*
18, 6703-6709.

23. M Dundr, U Hoffmann-Rohrer, Q Hu, I Grummt, L I Rothblum, R D Phair, and T Misteli, Mammalian RNA Polymerase in Vivo. *Science* **298,** 1623-1626 (2002).

24. V Stefanovsky, F Langlois, T Gagnon-Kugler, L I Rothblum, and T Moss, Growth factor signaling regulates elongation of RNA polymerase I transcription in mammals via UBF phosphorylation and r-chromatin remodeling. *Mol Cell* **21,** 629-639.

25. S Safran, *Statistical Thermodynamics of Surfaces, Interfaces, and Membranes.* (Westview Press, CO, US.), (2003).

26. P G de Gennes, F Brochard-Wyart, and D Quéré, Wetting phenomena Drops, Bubbles, Pearls, Waves. (Springer, NY, US.) (2004).

27. H Wang, D Boisvert, K K Kim, R Kim, and S H Kim, Crystal structure of a fibrillarin homologue from Methanococcus jannaschii, a hyperthermophile, at 1.6 Å resolution. *EMBO J.* **19,** 317-323 (2000).

28. K Peltonen, et al., A Targeting Modality for Destruction of RNA Polymerase I that Possesses Anticancer Activity. *Cancer Cell* **25,** 77–90 (2014).

29. D Drygin, et al., Targeting RNA Polymerase I with an Oral Small Molecule CX-5461 Inhibits Ribosomal RNA Synthesis and Solid Tumor Growth. *Cancer Res.* **71,** 1418-1430 (2011).

30. P de Gennes, *Scaling Concepts in Polymer Physics*. (Cornell Univ. Press, NY, US.), (1979).

31. E Nicolas, et al., Involvement of human ribosomal proteins in nucleolar structure and p53-dependent nucleolar stress. *Nat. Comm.* **7,** 11390 (2016).

32. H Matsumori, et al., Ribosomal protein L5 facilitates rDNA-bundled condensate and nucleolar assembly. *Life Sci. Alliance* **5,** e202101045 (2022).

33. S Banani, H Lee, A Hyman, M Rosen, Biomolecular condensates: organizers of cellular biochemistry. *Nat. Rev. Mol. Cell Biol.* **18,** 285–298 (2017).

34. Y Shin, C Brangwynne, Liquid phase condensation in cell physiology and disease. *Science* **357,** eaaf4382 (2017).

35. T Yamamoto, T Yamazaki, T Hirose, Phase separation driven by production of
architectural RNA transcripts. *Soft Matter* **16,** 4692–4698 (2020).

36. D Langevin, Microemulsion, *Acc. Chem. Res.* **21,** 255-260 (1988).

37. AW Folkmann, A Putnam, CF Lee, and G Seydoux, Regulation of biomolecular
condensates by interfacial protein clusters. Science. 373, 1218-1224 (2021).

38. MGJ Navarro, S Kashida, R Chouaib, S Souquere, G Pierron, D Weil, and Z Gueroui,
RNA is a critical element for the sizing and the composition of phase-separated RNA-
protein condensates. Nat. Comm., 10, 3230 (2019).

39. M Wu, et al., lncRNA SLERT controls phase separation of FC/DFCs to facilitate Pol
I transcription. *Science* **373,** 547–555 (2021).

40. T W Turowski and D Tollervery, Cotranscriptional events in eukaryotic ribosome
synthesis. *WIREs RNA* **6,** 129-139.

41. J Barandun, M Hunziker, and S Klinge, Assembly and structure of the SSU
processome – a nucleolar precursor of the small ribosomal submit. *Curr. Opin. Struct.*
Biol. **49,** 85-93.

42. R. Tomecki, P J Sikorski, M Zakrzewska-Placzek. Comparison of preribosomal RNA
processing pathways in yeast, plant, and human cells – focus on coordinated action of
endo- and exoribonulease. *FEBS Lett.* **591,** 1801-1850.

43. C E Scull and D A Schneider. Coordinated control of rRNA processing by RNA
polymerase I. *Trends Genet.* **35,** 724-733.

44. A Tartakoff, P DiMario, E Hurt, B McStay, V G Panse, and D Tollervery, The dual
nature of the nucleolus. *Genes Dev.* **36,** 765-769 (2022).

45. J Henninger, et al., RNA-Mediated Feedback Control of Transcriptional Condensates.
Cell **184,** 207–225 (2021).

46. L Hilbert, et al., Transcription organizes euchromatin via microphase separation. *Nat.*
Comms. **12,** 1360 (2021).

47. T Nozaki, et al., Dynamic Organization of Chromatin Domains Revealed by Super-
Resolution Live-Cell Imaging. *Mol. Cell* **67,** 282–293 (2017).

48. B Sabari, et al., Coactivator condensation at super-enhancers links phase separation
and gene control. *Science* **361,** eaar3958 (2018).

49. M Wei, et al., Nucleated transcriptional condensates amplify gene expression. *Nat.*
Cell Biol. **22,** 1187–1196 (2020).

50. T Kawaguchi, et al., SWI/SNF chromatin-remodeling complexes function in
noncoding RNA-dependent assembly of nuclear bodies. *Proc. Natl. Acad. Sci. USA* **112,**
4304–4309 (2015).

51. G Biamonti, C Vourc'h, Nuclear Stress Bodies. *Cold Spring Harb. Perspect. Biol.* **2,**
a000695 (2010).

52. K Ninomiya, et al., LncRNA-dependent nuclear stress bodies promote intron
retention through SR protein phosphorylation. *EMBO J.* **39,** e102729 (2020).

53. K Ninomiya, T Hirose, Short Tandem Repeat-Enriched Architectural RNAs in Nuclear
Bodies: Functions and Associated Disease. *non-coding RNA* **6,** 6 (2020).

54. P de Gennes, Conformation of polymers attached to an interface. *Macromolecules*
13, 1069–1075 (1980).

55. M Daoud, J Cotton, Star shaped polymers: a model for the conformation and its
concentration dependence. *J. Phys. France* **43,** 531–538 (1982).

 FIGURE LEGENDS

Figure 1   Multiphase structure of a nucleolus. (A) A nucleolus is composed of
multiple fibrillar center (FC) microphases in the sea of the granular component (GC).
There is a layer of dense fibrillar component (DFC) between each FC and GC. (B) RNA
polymerase I (Pol I) molecules (white particles) are entrapped in FC microphases (light
blue) and the active rDNA units (black line) are localized at the surfaces of microphases.
Nascent pre-rRNAs (green particles) are thus at the surfaces of the microphase and form
a DFC layer with RNA-binding proteins (magenta particles). The interface between FC
and DFC is located at a distance r_in from the center and the interface between DFC
and GC is located at a distance r_ex.

Figure 2   Model of transcription dynamics. DNA (black solid line) is localized at the
surface of an FC (cyan). (A) RNA polymerase I (Pol I) in a microphase binds to the
transcription starting site (TSS) of an active rDNA repeat unit. The bound Pol I starts
transcription with the rate k_e or returns to the microphase without starting transcription.
(B) During the transcription, Pol I migrates uni-directionally towards the transcription
terminating site (TTS) while polymerizing a nascent pre-rRNA. The terminal region of the
nascent pre-ribosomal RNA (pre-rRNA), to which fibrillarin binds, are cleaved by the co-
transcriptional RNA processing with the rate τ_pr^{-1}. The terminal region is released to GC.
(C) After the cleavage of the terminal region, Pol I continues transcription until it reaches
the TTS. At the TTS, Pol I is releaed to the FC with the rate τ_0^{-1}.

Figure 3   Composition of planer DFC layer vs interaction parameter χ. (A) The
volume fraction ϕ_r of pre-rRNA units (green), the volume fraction ϕ_p of freely diffusing
RBPs (magenta), the volume fraction ϕ_s of solvent molecules (cyan) in a DFC layer are
shown as functions of the interaction parameter χ. (B) The occupancy α_p of pre-rRNA
terminal region by RBPs is shown as a function of the interaction parameter χ. We used
$\mu_p/(k_\mathrm{B}T) = -10.0$, $\epsilon = -12.0$, $\sigma_\mathrm{in}b^2 = 0.05$, and $\Pi_\mathrm{ex}b^3/(k_\mathrm{B}T) = 0.0$ for the
calculations, see also Table 2. The solid lines are derived by numerically solving eqs.
S15, S16, and S20 in the SI Appendix. The broken lines are derived by using eqs. S47
(shown for $\chi > 10.1$).

Figure 4   Profile of volume fraction ϕ_r of pre-rRNA units in DFC layer. The volume
fraction of nascent pre-rRNA units is shown as a function of the position r in the DFC
layer ($r_\mathrm{in} < r < r_\mathrm{ex}$). The solid dark green line is derived by numerically calculating eqs.
S13, S15, and S16 in the SI Appendix for $\zeta = 0.06$ with the condition that the volume
fraction of solvent is zero and the occupancy of the terminal regions of pre-rRNAs by

RBPs is unity. The values of other parameters are summarized in Table 2. The broken

light green line is derived by using eq. S59 in the SI Appendix with $\phi_{\text{ex}} = 0.0157$ and

$r_{\text{ex}} = 1.98$ (which were derived from the numerical calculation to obtain the light green

line). The volume fraction of freely diffusing RBPs is shown in the inset.

Figure 5   **Free energy F of nucleolus.** The free energy F of the system is shown

as a function of the radius r_{in} of FCs (A) and the ratio $r_{\text{ex}}/r_{\text{in}}$ of the external radius to

the internal radius (B). The black solid line is the total free energy, including the free

energy of DFC layers (shown by the magenta broken line, the first term of eq. 2) and the

surface free energy (shown by the cyan broken line, the second and third terms of eq. 2)

for $\zeta = 0.06$. The values of other parameters are summarized in Table 2.

Figure 6 Radius r_{in} of FCs vs rescaled transcription rate ζ. The radius r_{in} of FCs

at the free energy minimum is shown as a function of rescaled transcription rate ζ. The

solid dark green line is derived by numerically calculating eqs. S13, S15, and S16 in the

SI Appendix with the condition that the volume fraction of solvent is zero and the

occupancy of the terminal regions of pre-rRNAs by RBPs is unity. The orange broken

line is derived by using eq. 5. The parameters used for the calculations are summarized

in Table 2.

Figure 7   Mild Pol I inhibition by BMH-21 increases the size of FCs. (A)

Immunofluorescence of UBF (FC) and NPM1 (GC) in HeLa cells with or without BMH-21

treatments. Scale bar, 10 μm. (B and C) Quantification of longest axis (B) and area (C)

of the FCs in cells under indicated conditions. Each scatter dot plot shows the mean

(black line). Dots indicate all points of quantified data (n = 250). Mean longest axes of

the FCs is shown below: 0 μM: 0.4907 μm, 0.0625 μM: 0.7440 μm, 0.125 μM: 0.8710

μm, 0.25 μM: 1.131 μm. Mean areas of the FCs are shown below: 0 μM: 0.170 μm^2,

0.0625 μM: 0.3255 μm^2, 0.125 μM: 0.4347 μm^2, 0.25 μM: 0.7312 μm^2. Statistical

analyses using Kruskal-Wallis test with Dunn's multiple comparison test were performed

and the results are shown as follows. (B) 0 μM vs 0.0625 μM: P < 0.0001, 0 μM vs 0.125

μM: P < 0.0001, 0 μM vs 0.25 μM: P < 0.0001, 0.0625 μM vs 0.125 μM: P < 0.0001,

0.0625 μM vs 0.25 μM: P < 0.0001, 0.125 μM vs 0.25 μM: P = 0.0008. (C) 0 μM vs 0.0625

μM: P = 0.0003, 0 μM vs 0.125 μM: P < 0.0001, 0 μM vs 0.25 μM: P < 0.0001, 0.0625

μM vs 0.125 μM: P < 0.0001, 0.0625 μM vs 0.25 μM: P < 0.0001, 0.125 μM vs 0.25 μM:

P = 0.0010. (D and E) Graphs showing the mean longest axis (D) and area (E) of the

FCs with SEM vs pre-rRNA expression levels. The pre-rRNA expression level in

untreated cells is defined as 1.

Figure 8   Exponent that accounts for the dependence of the size of FCs on pre-
rRNA expression level. The data in Fig. 7E was shown in the double-logarithm plot and
fitted with a power function. The magenta and cyan dots are the results by suppressing
the Pol I transcription by using BMH-21 and CX-5461. The slope of the double-logarithm
plot is the exponent that accounts for the dependence of the radius of FCs on the
transcription level. The curve fitting shows that the exponent is -0.49 for the case of the
BMH-21 treatment, -0.46 for the case of the CX-5461, and -0.48 if both data are fitted.

Figure 9   Summary of results.   RNP complexes enhance or suppress the growth of
condensates depending on whether the RNP complexes are mobile in the interior or
tethered to the surfaces of the condensates. A. The multivalent interaction between the
RNP complexes enhance the growth of the condensates if these condensates are
assembled by the RNP complexes. B. The multivalent interaction between the RNP
complexes suppress the growth of the condensates if these complexes are tethered to
the surfaces of the condensates assembled by other RNAs and proteins.
854
855

LEGENDS OF SUPPLEMENTARY FIGURES

Figure S1   BMH-21 and CX-5461 treatment reduce pre-rRNA expression levels in
a dose-dependent manner.
(*A*) Quantification of pre-rRNA expression levels by RT-qPCR in BMH-21-untreated and
-treated conditions. Data are represented as mean ± SD (*n* =3). (*B*) Quantification of pre-
rRNA expression levels by RT-qPCR in CX-5461-untreated and -treated conditions. Data
are represented as mean ± SD (*n* =3).

Figure S2   FCs and DFCs in the BMH-21 and CX-5461-untreated and -treated cells.
(*A* and *B*) Immunofluorescence of UBF (FC) and FBL (DFC) in HeLa cells with or without
BMH-21 (*A*) or CX-5461 (*B*) treatments. Scale bar, 10 μm.

Figure S3   Mild Pol I inhibition by CX-5461 increases the size of FCs.
(*A*) Immunofluorescence of UBF (FC) and NPM1 (GC) in HeLa cells with or without CX-
5461 treatments. In the cells treated with 2 μM CX-5461, the cells with large granules
(upper) or nucleolar caps (lower) were observed. Scale bar, 10 μm. (*B* and *C*)
Quantification of longest axis (*B*) and area (*C*) of the FCs in cells under indicated
conditions. Each scatter dot plot shows the mean (black line). Dots indicate all points of
quantified data (*n* = 500). Mean longest axes of the FCs is shown below: 0 μM: 0.5196
μm, 0.25 μM: 0.6367 μm, 0.5 μM: 0.7192 μm, 1 μM: 1.059 μm. Mean areas of the FCs
are shown below: 0 μM: 0.1155 μm^2, 0. 25 μM: 0.2171 μm^2, 0.5 μM: 0.3058 μm^2, 1 μM:
0.6544 μm^2. Statistical analyses using Kruskal-Wallis test with Dunn's multiple
comparison test were performed and the results are shown as follows. (*B*) 0 μM vs 0.25
μM: *P* < 0.0001, 0 μM vs 0.5 μM: *P* < 0.0001, 0 μM vs 1 μM: *P* < 0.0001, 0.25 μM vs 0.5
μM: *P* = 0.0211, 0.25 μM vs 1 μM: *P* < 0.0001, 0.5 μM vs 1 μM: *P* < 0.0001. (*C*) 0 μM vs
0.25 μM: *P* < 0.0001, 0 μM vs 0.5 μM: *P* < 0.0001, 0 μM vs 1 μM: *P* < 0.0001, 0.25 μM
vs 0.5 μM: *P* = 0.0012, 0.25 μM vs 1 μM: *P* < 0.0001, 0.5 μM vs 1 μM: *P* < 0.0001. (*D*
and *E*) Graphs showing the mean longest axis (*D*) and area (*E*) of the FCs with SEM vs
pre-rRNA expression levels. The pre-rRNA expression level in untreated cells is defined
as 1.

Figure S4   Quantification of the FCs in the nucleoli
Screenshots of detections of FCs and GCs by NIS Elements Advanced Research
software (NIKON) are shown. Green circles indicate UBF foci, and red circles indicate
nucleoli labeled by NPM1 antibodies. UBF foci detected in the nucleoli (lower right) were
quantified.

Figure 1 Multi-phase structure of a nucleolus. (A) A nucleolus is composed of multiple fibrillar center (FC) microphases in the sea of the granular component (GC). There is a layer of dense fibrillar component (DFC) between each FC and GC. (B) RNA polymerase I (Pol I) molecules (white particles) are entrapped in FC microphases (light blue) and the active rDNA units (black line) are localized at the surfaces of microphases. Nascent pre-rRNA transcripts (green particles) are thus at the surfaces of the microphase and form a DFC layer with RNA-binding proteins (magenta particles). The interface between FC and DFC is located at a distance r_{in} from the center and the interface between DFC and GC is located at a distance r_{ex}.

A Promoter binding/transcription start

B RNA processing

C Termination

A Pol I occupancy at TSS

$$\frac{dn_{\mathrm{ps}}}{dt} = k_{\mathrm{on}}\rho(1 - n_{\mathrm{ps}}) - k_{\mathrm{off}}n_{\mathrm{ps}} - k_{\mathrm{e}}n_{\mathrm{ps}}$$

B Pol I with nascent pre-rRNA terminal region

$$\frac{dn_{\mathrm{pe}}}{dt} = k_{\mathrm{e}}n_{\mathrm{ps}} - \frac{n_{\mathrm{pe}}}{\tau_{\mathrm{pr}}}$$

C Pol I without nascent pre-rRNA terminal region

$$\frac{dn_{\mathrm{p0}}}{dt} = \frac{n_{\mathrm{pe}}}{\tau_{\mathrm{pr}}} - \frac{n_{\mathrm{p0}}}{\tau_{0}}$$

Figure 2 Model of transcription dynamics. DNA (black solid line) is localized at the surface of an FC (cyan). (A) RNA polymerase I (Pol I) in a microphase binds to the transcription starting site (TSS) of an active rDNA repeat unit. The bound Pol I starts transcription with the rate k_{e} or returns to the microphase without starting transcription. (B) During the transcription, Pol I migrates uni-directionally towards the transcription terminating site (TTS) while polymerizing a nascent pre-rRNA. The terminal region of the nascent pre-ribosomal RNA (pre-rRNA), to which fibrillarin binds, are cleaved by the co-transcriptional RNA processing with the rate τ_{pr}^{-1}. The terminal region is released to GC. (C) After the cleavage of the terminal region, Pol I continues transcription until it reaches the TTS. At the TTS, Pol I is releaed to the FC with the rate τ_{0}^{-1}.

Figure 3 Composition of planer DFC layer vs interaction parameter χ. (A) The volume fraction φ_r of pre-rRNA units (green), the volume fraction φ_p of freely diffusing RBPs (magenta), the volume fraction φ_s of solvent molecules (cyan) in a DFC layer are shown as functions of the interaction parameter χ. (B) The occupancy α_p of RNA terminal region by RBPs is shown as a function of the interaction parameter χ. We used $\mu_p/(k_B T)=-10.0$, $\varepsilon=-12.0$, $\sigma_{in}b^2=0.05$, and $\Pi_{ex}b^3/(k_B T)=0.0$ for the calculations, see also Table 1. The solid lines are derived by numerically solving eqs. S15, S16, and S20 in the SI Appendix. The broken lines are derived by using eqs. S47 (shown for $\chi>10.1$).

Figure 4 Profile of volume fraction φ_r of pre-rRNA units in DFC layer. The volume fraction of nascent pre-rRNA units is shown as a function of the position r in the DFC layer ($r_{in} < r < r_{ex}$). The solid dark green line is derived by numerically calculating eqs. S13, S15, and S16 in the SI Appendix for ζ=0.06 with the condition that the volume fraction of solvent is zero and the occupancy of the terminal regions of pre-rRNAs by RBPs is unity. The values of other parameters are summarized in Table 1. The broken light green line is derived by using eq. S59 in the SI Appendix with φ_{ex}=0.0157 and r_{ex}=1.98 (which were derived from the numerical calculation to obtain the light green line). The volume fraction of freely diffusing RBPs is shown in the inset.

Figure 5   **Free energy F of nucleolus.** The free energy F of the system is shown as a function of the radius r_{in} of FCs (a) and the ratio r_{ex}/r_{in} of the external radius to the internal radius (b). The black solid line is the total free energy, including the free energy of DFC layers (shown by the magenta broken line, the first term of eq. 2) and the surface free energy (shown by the cyan broken line, the second and third terms of eq. 2) for $\zeta=0.06$. The values of other parameters are summarized in Table 1.

Figure 6 Radius r_{in} of FCs vs rescaled transcription rate ζ. The radius r_{in} of FCs at the free energy minimum is shown as a function of rescaled transcription rate ζ. The solid dark green line is derived by numerically calculating eqs. S13, S15, and S16 in the SI Appendix with the condition that the volume fraction of solvent is zero and the occupancy of the terminal regions of pre-rRNAs by RBPs is unity. The orange broken line is derived by using eq. 5. The parameters used for the calculations are summarized in Table 1.

Figure 7   Mild Pol I inhibition by BMH-21 increases the size of FCs. (A) Immunofluorescence of UBF (FC) and NPM1 (GC) in HeLa cells with or without BMH-21 treatments. Scale bar, 10 μm. (B and C) Quantification of longest axis (B) and area (C) of the FCs in cells under indicated conditions. Each scatter dot plot shows the mean (black line). Dots indicate all points of quantified data (n = 250). Mean longest axes of the FCs is shown below: 0 μM: 0.4907 μm, 0.0625 μM: 0.7440 μm, 0.125 μM: 0.8710 μm, 0.25 μM: 1.131 μm. Mean areas of the FCs are shown below: 0 μM: 0.170 μm^2, 0.0625 μM: 0.3255 μm^2, 0.125 μM: 0.4347 μm^2, 0.25 μM: 0.7312 μm^2. Statistical analyses using Kruskal-Wallis test with Dunn's multiple comparison test were performed and the results are shown as follows. (B) 0 μM vs 0.0625 μM: P < 0.0001, 0 μM vs 0.125 μM: P < 0.0001, 0 μM vs 0.25 μM: P < 0.0001, 0.0625 μM vs 0.125 μM: P < 0.0001, 0.0625 μM vs 0.25 μM: P < 0.0001, 0.125 μM vs 0.25 μM: P = 0.0008. (C) 0 μM vs 0.0625 μM: P = 0.0003, 0 μM vs 0.125 μM: P < 0.0001, 0 μM vs 0.25 μM: P < 0.0001, 0.0625 μM vs 0.125 μM: P < 0.0001, 0.0625 μM vs 0.25 μM: P < 0.0001, 0.125 μM vs 0.25 μM: P = 0.0010. (D and E) Graphs showing the mean longest axis (D) and area (E) of the FCs with SEM vs pre-rRNA expression levels. The pre-rRNA expression level in untreated cells is defined as 1.

Figure 8 Exponent that account for the dependence of the size of FCs on pre-rRNA expression level.
The data in Fig. 7E was shown in the double-logarithm plot and fitted with a power function. The magenta and cyan dots are the results by suppressing the Pol I transcription by using BMH-21 and CX-5461. The slope of the double-logarithm plot is the exponent that accounts for the dependence of the radius of FCs on the transcription level. The curve fitting shows that the exponent is -0.49 for the case of the BMH-21 treatment, -0.46 for the case of the CX-5461 treatment, and -0.48 if both data are used.

A

B

Condensate growth enhanced Condensate growth suppressed

Figure 9 Summary of results. RNP complexes of RNAs and RBPs enhance or suppress the growth of condensates depending on whether the RNP complexes are mobile in the interior or tethered to the surfaces of the condensates. A. The multi-valent interaction between the RNP complexes enhance the growth of the condensates if these condensates are assembled by the RNP complexes. B. The multi-valent interaction between the RNP complexes suppress the growth of the condensates if these complexes are tethered to the surfaces of the condensates assembled by other RNAs and proteins.