**Social hosts evade predation but have deadlier parasites**

Jason Walsman[a,*], Mary J. Janecka[a], David R. Clark[a], Rachael D. Kramp[a], Faith Rovenolt[a],

Regina Patrick[a], Ryan S. Mohammed[b], Mateusz Konczal[c], Clayton E. Cressler[d], Jessica F.

Stephenson[a]

[a]*Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260*

[b]*Department of Life Sciences, The University of the West Indies, St. Augustine, Trinidad and*

*Tobago*

[c]*Evolutionary Biology Group, Faculty of Biology, Adam Mickiewicz University, 60-614 Poznań,*

*Poland*

[d]*School of Biological Sciences, University of Nebraska, Lincoln, NE 68588*

[*]*Author for correspondence: Jason Walsman; Department of Biological Sciences, University of*

*Pittsburgh, Pittsburgh, PA 15260; walsmanjason@gmail.com*

**Abstract.** Parasites exploit hosts to replicate and transmit, but overexploitation kills both host and

parasite[1]: parasite virulence evolves to balance these costs and benefits. Predators can in theory

shift this balance by consuming hosts[2–4]. However, the non-consumptive effects of predators may

be as important as their consumptive effects[5]. Here, we use an eco-coevolutionary model to show

that predators select for host grouping, a common anti-predator, defensive social behaviour[6]. Host

grouping simultaneously increases parasite transmission, thus within-host parasite competition,

and therefore favours more exploitative, virulent, parasites[7]. When parametrized with data from

the guppy-*Gyrodactylus* spp. system, including our experimentally demonstrated trade-off

between virulence and transmission, our model accurately predicted the common garden-assayed

virulence of 18 parasite lines collected from four Trinidadian guppy populations under different

predation regimes. The quantitative match between theory and data lends credence to the model's

insight that the non-consumptive, social behaviour pathway is entirely responsible for the observed

increase in virulence with predation pressure. Our results indicate that parasites play an important,

underappreciated role in guppy evolutionary ecology. Moreover, group living is a common anti-

predator defence[6] and our general model accommodates host-parasite interactions across taxa: its

insight into the interactions among predation, sociality, and virulence evolution may apply broadly.

Our results additionally suggest that social distancing, by reducing host-host contact, can select

for less virulent parasites and pathogens.

**Main.** Unprecedented infectious disease emergence among human and wildlife populations[8,9]

demands that we improve our understanding of the evolutionary ecology of parasites and

pathogens (hereafter 'parasites'). A key tenet of infectious disease evolutionary theory is that

parasites face a trade-off between virulence and transmission[10–12]. Parasites exploit their hosts to

replicate and transmit, but increasing this exploitation harms parasite fitness by killing hosts

('virulence'), which reduces the time window for successful transmission, or the 'infectious

period'[1]. With appropriate curvature, this trade-off should result in stabilising selection for

intermediate virulence[13–16]. While theory predicts various mechanisms can shift the costs and

benefits of exploitation, altering virulence evolution[2–4,7] and host disease outcomes[12,15,17,18],

empirical evaluation of their relative importance remains scant[12,19,20].

Predation, a ubiquitous ecological interaction, can theoretically select for higher or lower

virulence[2–4,12]. Predation directly increases host mortality, selecting for faster exploitation and

transmission before the host dies, and hence increased virulence[2,3,19]. Conversely, predation may

select for lower virulence through a more complex pathway. By reducing the density of infected

hosts, predation reduces the per-host rate at which new infections arise ('force of infection');

lower force of infection reduces infections with multiple parasite genotypes[4] ('multiple

infections') and thus the within-host competition between parasite genotypes that selects for

higher virulence[20]. In addition to these consumptive effects, predators can also affect host traits[5],

shifting how they interact with their parasites[21–24], and probably virulence evolution. One

example is host grouping, a common defence[6] which effectively decreases predator-induced

mortality[25,26], potentially selecting for lower virulence[7]. However, grouping also increases host-

host contact rates, increasing the force of infection, so host grouping should increase multiple

infections and therefore select for higher virulence[7,20,27]. Both consumptive and non-consumptive

effects of predation act simultaneously in natural communities[23], but their relative importance for

selection on parasite virulence lacks empirical and theoretical clarification.

Here, we elucidate how predators affect virulence evolution using theory and data from

Trinidadian guppies, *Poecilia reticulata* and their *Gyrodactylus* spp. parasites (Fig. 1). Persistent,

natural variation in predation risk drives population-level variation in guppy grouping for

defence ('shoaling')[26,28,29], which likely controls the contact rate for directly transmitted,

*Gyrodactylus* spp. ectoparasites[30,31]: high-predation populations shoal more and suffer higher

*Gyrodactylus* spp. infection prevalence[30,32,33]. First, we experimentally confirm that

*Gyrodactylus* spp. parasite lines fall along the commonly assumed virulence-transmission trade-

off (Fig. 2) and show that *Gyrodactylus* species (*G. turnbulli* and *G. bullatarudis*) and genotypes

co-occur on individual guppies in the wild (Extended Data [ED] Table 1). These factors make

this system ideal for testing how predation shifts virulence evolution. We then used laboratory

and field measurements to parametrize a model with selection on host and parasite phenotypes

interacting with host and parasite densities ('eco-coevolutionary model'), revealing one

dominant pathway (Fig. 3): predators shift coevolution toward higher shoaling rate, increasing

multiple infections, and thus also higher virulence. The model accurately predicted that wild-

collected, *Gyrodactylus* spp. parasites from high shoaling populations had higher laboratory-

assayed virulence (Fig. 4). Because we assayed parasite virulence in common garden conditions,

this difference was due to differences in parasite exploitation of hosts, not host defences. Overall,

the quantitative model-data match lends credence to the model's inference into parasite evolution

in wild communities.

**Figure 1. Natural guppy populations differ in predation, driving evolutionary divergence in shoaling rate.** *Waterfalls divide upper course and lower course guppy populations and prevent the upstream migration of large piscivores. Natural populations have therefore evolved under different predation regimes, replicated across rivers. Shoaling rate differences likely drive population-level differences in transmission rate of*[30,34,35]*, and thus selection on the virulence of, their highly prevalent, directly transmitted monogenean ectoparasites* Gyrodactylus *spp.. The map shows locations and data types that we use.*

**Gyrodactylus spp. transmission trades off with virulence, mediated by parasite intensity**

**Figure 2. Infection intensity links transmission rate and virulence with a stabilizing trade-off. a,** *Across four parasite lines, transmission rate increased with infection intensity. Points are transmission events jittered vertically.* **b,** *Across 22 ['domestic' (circles) or 'wild' (triangles)] parasite lines, higher intensity (a measure of parasite growth rate) induced higher host death rate ($d+v_i$, a measure of virulence; back-transformed partial residuals).* **c,** *(a) allows a link between transmission rate and virulence across lines. Points in (b, c) are mean line traits. In (a-c), bands are 95% C.I.* **d,** *The curvature in (c) maximizes $R_{0i}$ at intermediate transmission rate/virulence.*

We found a virulence-transmission trade-off among *Gyrodactylus* spp. lines. First, we

found that transmission rate, given arena area and time until infection, increased asymptotically

as intensity increased (Fig. 2a; $\beta = 6.08 \times 10^{-3}$ intensity$^{0.138}$; $\chi^2_{(1, N = 101)} = 17.1$, $P = 3.48 \times 10^{-5}$, $r =$

0.35). This pattern did not differ across two experiments using different, domestic parasite lines

(isolated from commercial guppies instead of wild-caught; line effect: $\chi^2_{(3, N = 101)} = 1.89$, $P =$

0.596).

Second, 22 laboratory-maintained lines (including three of four from Fig. 2a) differed in

the mean intensity they attained ($F_{21, 1149} = 8.49$, $P = 9.36 \times 10^{-25}$, $\eta^2 = 0.13$), and higher intensity

lines imposed a higher mean death rate on infected hosts (Fig. 2b; $\chi^2_{(1, N = 22)}, = 6.97$, $P = 0.008$, $r$

$= 0.49$). Across intensities, domestic lines imposed less death ($\chi^2_{(1, N = 22)} = 14.73$, $P = 1.24 \times 10^{-}$

$^4$, $r = 0.63$; domestic $= 0.016$ *vs*. wild $= 0.052$ at overall mean intensity). We used back-

transformed, partial residual death rate to control for non-focal predictors (line origin: Fig. 2b

shows the relationship as if all lines were wild) so that we could determine the relationship

relevant to wild parasites (all results are very similar if instead we set all lines as domestic, see

SI).

Third, we predicted the mean transmission rate for each line, given its intensity

measurements and the transmission-intensity relationship in Fig. 2a. We found that the mean

transmission rate for each line traded-off with the mean virulence ($d+v$ minus a reasonable

background death rate, $d = 1.30 \times 10^{-3}$ day$^{-1}$, see ED Table 2 for symbols, meanings, values, and

sources): parasite lines that attained higher intensity benefitted from higher transmission rate but

higher virulence shortened their infections, on average (Fig. 2c). Since *Gyrodactylus* spp. are

capable of sexual and asexual reproduction, we tested whether any of our lines were clones (i.e.,

identical multi-locus genotypes, MLGs). Genotyping a subset (13/22, based on sample

availability) found no identical MLGs (all pairs within populations differed at $\geq$ 16 loci,

representing at least half of all variable loci for each source population, see SI).

The curvature of this trade-off indicated stabilizing selection on virulence ($v_i = 1.38$ x $10^6$

$\beta_i^{3.61}$; $d$ $v_i$ and $\beta_i$ are average traits for a line). Bootstrapping parasite lines included in the

analysis showed this curvature was significant (exponent > 1 in all $10^4$ bootstrapped samples)

and AIC indicates a better fit to the data than for a linear model ($\Delta$AIC = 13.3). Theory predicts

that the trade-off curvature leads to optimal parasite fitness at intermediate transmissibility ($\beta_i =$

$cT_i$, $c$ is host shoaling rate, $T_i$ is a line's transmissibility) and virulence [Fig. 2d; $R_{0i} =$

$cT_iS/(d+v_i+y)$ for pure infections where we use reasonable values of $S = 10$ susceptible host m$^{-2}$

and $\gamma = 0.020$ day$^{-1}$ $d$, see ED Table 2]. Unsurprisingly, parasite lines fell to the right of this

optimum: higher background mortality in wild than laboratory guppies shifts the optimum

higher; the mixed-stock guppies in our common garden assay may differ substantially from local,

wild hosts; and multiple infections (common in the wild, ED table 1) select for higher virulence

than would optimize $R_{0i}$ of pure infections[4].

**Eco-coevolutionary model resolves complexity to show that social hosts have deadlier**

**parasites**

Our model provides two general insights. First, it demonstrates that the net, selective

effect of predation, shoaling rate, or other ecological factors on virulence depends only on

whether they increase or decrease the infectious period (see SI for proof). Second, the model

shows that shoaling rate evolves to balance predator-induced mortality against parasite-induced

mortality. When predation increases, hosts evolve higher shoaling rate, increasing transmission

rates. When parasites are more abundant and virulent, hosts evolve lower shoaling rate to prevent

infection. For some parameter values, higher virulence can select for higher shoaling rate if

parasites become so virulent that the density of infected hosts and force of infection decline

strongly (ED Fig. 1).

160           When parametrized with empirical data from the guppy-*Gyrodactylus* spp. system, the

model complexity resolves into one dominant pathway: predation increases coevolutionarily

stable contact rate and thus virulence (hereafter 'coevolutionary' traits; Fig. 3). When predators

only have consumptive effects on predation (i.e., hosts do not evolve), predation decreases

prevalence and virulence (grey curves in Fig. 4a-e; presence/absence of parasite evolution has

little effect on grey curves). However, as hosts evolve higher shoaling rate in response to

predation (Fig. 4a), predation also increases prevalence (black curve in Fig. 4b) in all tested

parameter sets (see sensitivity analysis methods). Increasing predation increases overall host

mortality and reduces host density, particularly when hosts evolve (Fig. 4c). Further, most of the

increase in mortality (76%) with predation is due to increased parasite-induced mortality

(prevalence × virulence), rather than increased predator-induced mortality (ED Fig. 2; robust to

parameters: ED Fig. 3). Higher shoaling rate increases the force of infection (and thus multiple

infections), selecting for higher parasite intensity and virulence (Figs. 4d, e; note intensity was

not modelled directly but back-calculated from transmissibility according to Fig. 2b); in all

parameter sets, higher predation increased virulence. Importantly, predators increase virulence

through their non-consumptive effects on shoaling rate (Fig. 4e; robust to parameters: ED Fig.

3). The consumptive effects of predation only slightly affect virulence (grey curves in Fig. 4e)

because decreasing multiple infections is as important as increasing mortality.

**Figure 3. Predation drives increased shoaling rate and virulence in the eco-coevolutionary**

**model.** *Curves give the continuously stable strategy (CSS) for parasite ($v_{CSS}$ black) or host ($c_{CSS}$*

*grey) evolution at predation levels corresponding to natural populations ($P_{Low}$; dashed or $P_{High}$;*

*solid). Parasites evolve higher virulence with higher shoaling rate. The consumptive effects of*

*predation have little effect on evolved virulence. Hosts evolve lower shoaling rate with increasing*

*virulence (grey curves move toward lower* c *as* v *increases). Predation substantially increases*

*evolved shoaling rate. Host and parasite curves intersect at coevolutionarily stable points (green*

*and blue points). As predation increases (from green point to blue point), coevolutionarily stable*

*shoaling rate and virulence increase.*

**190**  **Empirical results quantitatively match model predictions that social hosts have deadlier**

**191**  **parasites**

**192**      The model's coevolutionary outcomes successfully fit training data (Fig. 4a-c, f-h; ED

**193**  table 2). High predation fish spent more time shoaling in our assay ($\chi^2_{(1, N = 68)}$ = 7.68, $P$ = 0.006,

**194**  $r$ = 0.32) and across population estimates extracted from the literature ($\chi^2_{(1, N = 22)}$ = 11.17, $P$ =

**195**  8.31 x 10$^{-4}$, $r$ = 0.58; Fig. 4f). While percent time shoaling is unlikely precisely proportional to

**196**  host-host contact rate (as we use it, see ED Table 2), shoaling is relevant to transmission in this

**197**  system[34,35]. This helps explain why high predation populations suffered higher parasite

**198**  prevalence across multiple rivers, sites, and years ($df$ = 1, $\chi^2_{(1, N = 107)}$ = 5.02, $P$ = 0.025, $r$ = 0.21;

**199**  Fig. 4g). Higher levels of predation and parasitism may depress host density (though not

**200**  significantly so: $df$ = 1, $\chi^2_{(1, N = 23)}$ = 2.18, $P$ = 0.140, $r$ = -0.29; Fig. 4h). Overall, the model fit the

**201**  training data reasonably well, with an average relative error of 10.5% per fitted quantity (shown

**202**  in Fig. 4f-h and ED table 2).

**203**

**204**      The trained model predicted higher virulence in high predation populations. We validated

**205**  this prediction by quantifying the traits of 18 genotypically-distinct parasite lines isolated from

**206**  wild populations and maintained in the lab under common garden conditions for 65 days. Lines

**207**  from high predation populations attained higher intensity on ($df$ = 1, $\chi^2_{(1, N = 425)}$ = 4.80, $P$ =

**208**  0.028, $r$ = 0.11; Fig. 4i) and induced higher death rate for infected, mixed-stock hosts ($df$ = 1,

**209**  $\chi^2_{(1, N = 216)}$ = 4.75, $P$ = 0.029, $r$ = 0.15; Fig. 4j). The two parasite species, *G. turnbulli* (11 known

**210**  lines) and *G. bullatarudis* (3 known lines) did not differ significantly in intensity ($df$ = 1, $\chi^2_{(1, N = }$

**211**  $_{345)}$ = 2.73, $P$ = 0.098, $r$ = 0.09) or virulence ($df$ = 1, $\chi^2_{(1, N = 177)}$ = 0.891, $P$ = 0.345, $r$ = 0.07;

**212**  unlike[36]). Restricting the analysis for Fig. 4j to *G. turnbulli* found marginally higher intensity ($df$

$= 1, \chi^2_{(1, N = 208)} = 3.15, P = 0.076, r = 0.12$) and significantly higher virulence in high predation

populations ($df = 1, \chi^2_{(1, N = 102)} = 6.86, P = 0.009, r = 0.25$). The overall quantitative match

between the model-predicted mortality and our empirical results supports our model's inferences

($d+v$ in the model *vs.* mean of $d+v$ back-transformed partial residuals: low predation 0.020 *vs.*

0.021; high predation: 0.044 *vs.* 0.041). We compared theoretical predictions to partial residuals

to control for non-focal predictors in the statistical model, especially duration of parasite

maintenance in the laboratory, to obtain the most biologically relevant estimate of virulence. Our

theoretical model makes this prediction based on higher shoaling rate increasing multiple

infections, which may be more frequent in high predation populations (see ED table 1).

The model-data agreement on evolved virulence aids model-data agreement on how

mortality changes across predation regime. In our model, 76% of the increased mortality across

predation regime is increased parasite-induced mortality while increased predator-induced

mortality accounts for 24%. Prevalence in natural populations and virulence data on laboratory

fish ($p_{Hi}v_{Hi}$-$p_{Low}v_{Low}$) indicated that parasitism would explain 64% of the mortality difference

from Reznick *et al*[37] while predation may account for the remaining 36%.

**Figure 4. Predation increases shoaling rate and thus selects for higher virulence.** *a-e, Model results. Black curves: eco-coevolution (connecting Fig. 3 coloured points). Grey curves: no host evolution. f-h, empirical results. Horizontal segments: predation regime means. Grey points: one river/regime/year mean. Coloured points: one focal river/predation regime mean (Aripo left of Guanapo). Vertical bars: standard errors. Training data: violins. Non-training data: boxplots. (a, f) Shoaling increased with predation. (b, g) Prevalence increased with predation while (c, h) host density decreased non-significantly. (d, i) Parasite intensity and (e, j) virulence increased with predation (boxplots are back-transformed partial residuals; centre line, median; box limits, first and third quartiles; whiskers, 1.5x interquartile range; points, outliers).*

**Discussion.** Our theoretical-empirical approach clarifies that predation drives the evolution of parasite virulence by increasing host shoaling rate. As a result, shoaling in response to increasing predation pressure leads to parasite-induced mortality rising more than that induced by predators. In contrast to these strong non-consumptive effects, we found that the consumptive effects of predation are small, balance one another out, and barely alter virulence evolution. We discuss key emergent patterns.

    How shoaling rate drives virulence evolution depends on multiple infections. Without multiple infections, shoaling decreases predator-induced mortality and thus selects for decreased virulence[38]. With multiple infections, we and others[7] find that increased shoaling rate selects for higher virulence. While we do not know of previous empirical tests of the effect of host-host contact rate on virulence evolution, our result is analogous to more host dispersal[18,39,40] and

parasite founder diversity[17] selecting for higher host exploitation through high local parasite

diversity[41].

We assayed parasite traits using infections on mixed-stock, wildtype guppies: this

approach allowed us to draw robust conclusions about parasite evolution but leaves untested how

host defences besides shoaling rate may affect disease dynamics in natural communities. Thus

far, guppy defences against parasites have not been robustly characterised across predation

regime, although illustrative data suggest that guppies from low predation regimes are better

defended against, and better able to evolve in response to *Gyrodactylus* spp.[30,32,33,42,43]: they need

not balance this investment with defence against predators[24], and sexual selection, which acts on

gyrodactylid resistance[44,45], is more effective in the absence of predators[46]. If high predation

guppies are indeed less well defended against *Gyrodactylus* spp., our common garden assay

likely underestimates the difference in virulence that would be expressed in local hosts across

predation regimes. How host behavioural and immunological defences may coevolve with

parasite virulence in natural communities remains an outstanding and complex theoretical and

empirical challenge.

The importance and implications of predation and social parasite transmission, evident

from our model and data, may hold across directly transmitted parasites of group-living hosts[47].

Predators drive defensive group-living in animals across taxa[6], which can increase parasitism[48],

creating ecological and evolutionary feedbacks between host sociality and parasites[47]. Parasites

evolve along virulence-transmission trade-offs in systems ranging from viruses of humans[16],

bacterial pathogens of birds[14], protozoan pathogens of insects[15], and our monogenean fish

ectoparasite. Multiple infections are common for many parasites, allowing the simple mechanism

of competition for within-host resources to select for higher virulence[20]. Thus, diverse systems

likely meet the essential assumptions of our model: predators may frequently shift the

antagonistic interplay of host sociality and parasite virulence, driving hosts into the arms of more

virulent parasites. Conversely, these results also indicate that social distancing may select for

lower virulence when parasites exhibit multiple infections and a virulence-transmission trade-off

(e.g., influenza A virus[13,49]). Host behaviour that reduces contact may thus effectively control

both the spread and virulence evolution of pathogens and parasites.

**Methods.**

*1.  Molecular methods*

Identifying parasites to species. We used two methods of molecular species identification

for individual parasites: we sequenced the mitochondrial COII gene for a subset of domestic

lines established before March 2020, and we used a newly developed restriction enzyme assay

for wild and domestic lines established after March 2020 (see SI for further details). To prevent

disrupting data collection, sample collection was restricted to already-dead hosts, stored in 70 %

EtOH. This conservative precaution meant we were unable to collect useable samples from some

of our parasite lines.

Distinguishing parasite genotypes. We developed panels of single nucleotide

polymorphisms (SNPs) for *G. turnbulli* and *G. bullatarudis* to examine the incidence of multi-

genotype coinfections from wild collections, and to verify that the lines we established in the lab

were genetically distinct. Genotypes were called using Fluidigm SNP genotyping analysis

software. A subset of individuals was re-genotyped to identify and remove error-prone loci and

estimate error rates (the proportion of mismatches to matches for re-genotyped loci). Loci which

amplified consistently and with score calls greater than 90% were selected for use in the analysis.

This resulted in a total of 140 variable loci for *G. turnbulli* (error rate 1.5%) and 83 loci for *G.*

*bullatarudis* (error rate 2 %) across all populations. However, due to significant local variation in

informative loci, the total number of loci used for each source population varied (see SI). The

resulting multilocus genotypes (MLGs) were assessed using GENALEX 6.502 to estimate the

fraction of multi-genotype infections at six sites across Trinidad including our four focal

populations (low- and high-predation in the Aripo and Guanapo rivers), and to assess the

genotypes of the established lines.

As a conservative estimate, two individuals were considered to have different genotypes

if they varied in at least 50% of the total variable loci from a given source population (see SI for

further method details). This applied both to determining that our lines were different genotypes

and finding the frequency of multiple infections. In finding the frequency of multiple infections,

we genotyped worms from fish with at least two worms and accounted for the proportion of fish

with only one worm, estimated for each site from our 2020 survey and Stephenson *et al*.[30] (as our

2020 survey lacked power for some sites).

2. *Parasite trait data*

General maintenance. We established parasite lines by transferring a single worm to an

uninfected host from our mixed, laboratory-bred stocks descended from wild populations. Each

founding worm was obtained from a single guppy either from a commercial supplier ("domestic"

lines) or wild-caught in Trinidad ("wild" lines from Caura, Aripo, Guanapo, and Lopinot rivers

in Caroni drainage). Wild-caught adult guppies (~50/population) were shipped from Trinidad to

the University of Pittsburgh in March 2020 (see ethics for permit information). We established

43 parasite lines (18 domestic and 25 wild, 10 from low predation populations and 15 from high

predation). Different lines were included in different analyses, as explained below. Lines were

maintained under common garden conditions for 65 days (some domestic lines were maintained

in the lab much longer) on groups of 3-6 guppies. We added uninfected fish to each line as

required to replace those that either died or were found parasite-free during twice-weekly

screening of all fish in each line under anaesthetic (tricaine methanesulfonate; 4g/L) using a

dissecting microscope. During these screens, we recorded the number of parasites infecting each

fish. Each line was housed in a single 1.8 L tank on a recirculating system (Aquaneering Inc.;

12L:12D; 24ºC). Recirculated water passes through fine foam, sand, and ultraviolet filters before

re-entering other tanks: individual parasites cannot transmit between lines, as supported by our

genotyping.

<u>Trait measurements.</u> We estimated transmission rate by re-analysing data from

Stephenson *et al.*[50] and conducting a similar new experiment. Donor fish received an infection of

one parasite line or multiple infection and were individually housed in 1.8 L tanks (donor fish

received line A, B, C, or A and B; no lines in the new experiment were used by Stephenson *et*

*al.*[50]). We added a parasite-naïve recipient fish on day 8 of the donor's infection, and screened

both fish for infection every 24 hrs.

We also assayed intensity and death rate. For intensity (a proxy for within-host growth

rate), we counted the number of parasites on each host at each date (1 observation). We

estimated the per-capita host death rate from the number of fish found dead divided by the

number of infected fish in the tank at the previous observation time point and the days between

observations (1 observation; 79% of observations were 3±1 days after the previous one).

Trade-off statistics. We calculated transmission rate from the number of days until

successful infection of the recipient and intensity as the number of worms on the donor fish on

the day of transmission, following Stephenson et al.[50] (total $n = 101$ transmission events). Worm

numbers changed slowly on the scale of the number of days until successful infection. In these

assays, $dS/dt = -cT_iIS$ ($c$ is contact rate, $T_i$ is the transmissibility of parasite line i, $I$ is infected

host density, and $S$ is susceptible host density) so susceptible hosts follow an exponential decay

pattern; thus, mean time until infection is $1/(cT_iI)$. If days until transmission are $D$, then the

transmission rate estimated by a given transmission event is $cT_i = 1/(DI)$. Based on the area of

enclosures, $I = 88.5$ hosts m$^{-2}$ for Stephenson et al.[50] and $I = 72.0$ hosts m$^{-2}$ for the follow-up

experiment. We fit these estimates of transmission rate to $\log_{10}$ intensity with parasite line (line

A, B, C, or coinfection of lines A and B) as a fixed effect using a generalized linear model with a

Gamma error family and log link function. Parasite line was non-significant, so we refit the

model without parasite line (P-value for intensity reported from this model; all P-values

represent two-sided, Type II Wald chi-square or F tests for this and other methods). Effect sizes

are $\eta^2$ for a simple ANOVA or partial correlation coefficients, $r$. For all analyses, we provide the

code, output, and validation of the model fit in the SI.

When determining average traits for a line, we only included lines maintained in the

laboratory for more than 30 days ($n = 22$; 10 domestic, 12 wild lines, including 1-4 lines from

each of the four focal populations). We used an ANOVA to determine whether lines differed

significantly in log intensity (1171 observations of intensity for the 22 parasite lines, at least 25

for each line). We modelled mean death rate for each line (344 observations and at least 13 for

each line) as a function of mean intensity and line origin (wild *vs.* domestic) with a beta error

family and logit link function. We estimated virulence ($v$) by subtracting background death rate

(see ED Table 2) from partial residuals of infected death rate (to get $v$ from $d+v$). We used partial

residuals of infected death rate back-transformed onto the response scale to control for non-focal

predictors (here, line origin) and find the relationship for wild parasites. We estimated mean

transmission rate for a line by mapping the individual intensity measurements for each line onto

the relationship observed between intensity and transmission in the experiments described above

(Fig. 2a); we then calculated the mean transmission rate for each line. With estimates of

transmission rate and virulence for each line, we examined the relationship between them using a

generalized linear model of virulence with Gaussian error distribution and log link function as a

function of natural log transmission rate. The statistical model was free to fit a positive (exponent

$> 1$), neutral (exponent $= 1$), or negative curvature (exponent $< 1$) to these data. We bootstrapped

this fit by sampling lines to include in the analysis from the 22 parasite lines with replacement to

determine how often the exponent was more than one.

Statistics on predation regime differences in parasite traits. To examine the patterns in

Fig. 4i and 4j, we used our estimates of parasite intensity and host death, (calculated as above) as

response variables in GLMMs. These traits were assayed from parasites collected from low and

high predation populations of our focal rivers (low/high predation; river distances: low Guanapo-

high Guanapo = 11.1 km, high Guanapo-high Aripo = 23.9 km, high Aripo-low Aripo = 6.6 km;

3 lines from high-predation Aripo and 5 from each of the other focal populations; 18 lines total).

Each observation was of one line at one date but intensity on and death of multiple hosts for each

line. This analysis included some lines maintained in the laboratory for less than 30 days but

these lines necessarily had fewer observations and thus did not influence the analysis as much as

lines with more observations. For intensity ($n = 425$ observations), we used predation regime,

river, and number of fish in the common garden at time of measurement (since *Gyrodactylus*

spp. can move relatively freely among fish) as fixed effects. Parasite line and lineday (days the

line had been in lab) nested within parasite line were included as random effects. We used

log(intensity) as the response variable. For death rate ($n = 216$), we used predation regime and

river as the fixed effects with parasite line and lineday nested within line as random effects, a

Tweedie error family, and a log link function. The same analyses were used when determining

the effect of predation regime on intensity ($n = 208$) and host death rate ($n = 102$) for known *G.*

*turnbulli* lines. We compare the theoretical model to mean partial residuals of infected host death

rate, back-transformed onto the response scale, to control for non-focal predictors, such as

lineday, to find the most representative estimate of death rate in each predation regime.

Statistics on species differences in parasite traits. To determine trait differences by

species, we considered all wild lines that were identified to species (including 7 from rivers

considered non-focal due to lack of high predation *vs.* low predation comparisons). Three of

these lines were *G. bullatarudis* and 11 were *G. turnbulli*. Because all three *G. bullatarudis* lines

were from high predation populations, we only compared species differences within that regime.

Species identity was used as a predictor instead of predation regime but statistical models were

otherwise identical to those for the effect of predation regime for intensity ($n = 345$) and infected

host death rate ($n = 177$).

*3. Theoretical modelling*

Model ecology. We wrote a relatively simple, ordinary differential equation model that

retains the key biology of the predator-prey/host-parasite system. In the model, predators are not

limited by prey/host density (e.g., because predators are generalists), allowing us to represent

predation as a parameter ($P$) for simplicity instead of a state variable. Susceptible hosts ($S$)

become infected ($I$) and can recover to an immune state ($R$) with waning immunity (see ED

Table 2 for symbols, values, and units):

$$dS/dt \ = \ [a - q(S + I + R)](S + I + R) - (d + \frac{Pc_1}{c})S - cTIS + zR \quad (1a)$$

$$dI/dt \ = \ cTIS - (d + \frac{Pc_1}{c})I - vI - \gamma I \quad (1b)$$

$$dR/dt \ = \ \gamma I - (d + \frac{Pc_1}{c})R - zR \quad (1c)$$

Susceptible hosts are born from all hosts with density dependent, per-capita birth rate $a$-

$q(S+I+R)$ (i.e., hosts grow logistically, eq. 1a). Hosts die at background rate $d$ and from

predation at rate $Pc_1/c$ that decreases with host shoaling rate ($c$; that $P$ and $c_1$ together determine

the strength of predation). Along with parasite transmissibility given contact ($T$), the rate of

density-dependent transmission from $I$ to $S$ depends on the host-host contact rate, dictated by

shoaling rate. Recovered hosts also move back into the susceptible class as immunity wanes at

rate $z$ (average duration of immunity is $1/z$).

Infected hosts (eq. 1b) suffer background mortality and predation (predation is not

selective in this model) while suffering additional mortality due to virulence, $v$. If predation was

selective, especially to remove the sickest hosts, the consumptive effects of predation would be

even more likely to select against virulence[3], strengthening our conclusion that shoaling rate

drives virulence (Fig. 4). Infected hosts recover with rate $\gamma$ (eq. 1c). Recovered hosts suffer

background mortality and predation while losing immunity over time.

Model evolution: We modelled parasite and host evolution via Adaptive Dynamics[51].

Biologically, parasite evolution is rendered more complex by the presence of multiple lines as

well as two phenotypically similar species; host evolution is also complex because shoaling rate

has important genetic and plastic components[31]. For both host and parasite, our model is agnostic

regarding the basis of adaptation but simply examines competition between phenotypes based on

fitness of the rarer phenotype (i.e., invasion analysis). Each phenotype corresponds to an asexual

genotype in the model. Parasite fitness when invading depends on the traits of the mutant (m)

and resident genotypes (r):

441
$$\frac{1}{I_\mathrm{m}}\frac{dI_\mathrm{m}}{dt} = cT_\mathrm{m}S_\mathrm{r}^* - d - \frac{Pc_1}{c} - v_\mathrm{m} - \gamma + \sigma c T_\mathrm{m}I_\mathrm{r}^* - \sigma c T_\mathrm{r}I_\mathrm{r}^* \quad (2a)$$

442
$$v_\mathrm{i} = k_1 T_\mathrm{i}^{k_2} \quad (2b)$$

The resident sets host densities at $S_\mathrm{r}$ and $I_\mathrm{r}$ which depend on its traits, $T_\mathrm{r}$ and $v_\mathrm{r}$. Mutant fitness

depends on transmission to susceptible hosts, background death, predation, virulence, recovery,

and gains and losses due to superinfection (eq. 2a). In superinfection, parasites of one genotype

take over a host already infected by another genotype. We assume that each genotype has the

same probability of successful superinfection given transmission, $\sigma$. This assumption is likely

conservative as the probability of successful superinfection is often expected to increase with

virulence[4], which could amplify the impact of contact rate to select for higher virulence. From

parasite fitness when invading, we find the continuously stable strategy[51] (CSS) for parasite

virulence; a CSS represents a trait that can be reached by gradual trait changes but that cannot be

invaded by rare phenotypes with slightly different traits.

For host fitness, we used the Next Generation Matrix technique[52] that accounts for fitness

within each class and the rates of movement between classes (see SI). In the model, host

genotypes differ in their contact rates, $c_i$. An invading host genotype (contact rate $c_\mathrm{m}$) contacts

the resident genotype ($c_\mathrm{r}$) at rate sqrt($c_\mathrm{m}c_\mathrm{r}$) (for the purposes of transmission and predation),

following Bonds et al.[38]. An invading genotype suffers crowding (-$q$ term, eq. 1a) from all hosts.

Only intersections of a host CSS curve and parasite CSS curve (Fig. 3) can be coevolutionary

stable points (coCSSs). We confirmed that a potential coCSS is indeed a coCSS with the strong

convergence stability criterion[53].

Model parametrization. We have empirical estimates of some, but not all, of the parameters

of the eco-coevolutionary model. As such, we used a simple evolutionary algorithm (described

below) to estimate all model parameters, using empirical parameter estimates and field estimates

of model outputs (e.g., prevalence) to fit the model to the data. Importantly, we did not use

empirical data on evolved virulence in the fitting process, i.e., model virulence is free to evolve

along the virulence-transmissibility trade-off however best fits the other data; instead, we used

virulence data to validate the model.

Many empirical estimates came from the literature (see below and ED table 2). We estimated

maximum birth rate ($a$) from lifetime fecundity of guppies with high food[54] in the laboratory. We

estimated background mortality in the absence of parasites and predators ($d$) as the inverse of life

expectancy in the laboratory[54]. We estimated recovery rate ($\gamma$) as the inverse of the time required

for fish to clear infection in the laboratory[55]. We estimated the rate at which immunity wanes ($z$)

from the observation that fish were roughly half resistant (averaging across totally resistant,

partially resistant, and not resistant) 21 days after initial infection[56]. Our trait measurements

provided estimates for the trade-off parameters (Fig. 2c) given an estimate of $d$.

For parameters where we lacked estimates [importantly $Pc_1$ (which acts as one parameter) in

low- and high-predation populations and the superinfection parameter, $\sigma$], we also fit the model's

eco-coevolutionary outputs at low and high predation to estimates of such outcomes in the field.

We converted recapture rates from a mark-recapture experiment[37] into instantaneous mortality

rates to determine overall mortality rate in the field ($d+Pc_1/c+pv$) for both a high predation

population and a low predation population. We found a transmission rate reasonable for low

predation populations ($cT_{Low}$) by fitting transmission rate to peak prevalence and timing of peak

prevalence found for low predation Aripo guppies and parasites in a stream mesocosm[57]. We

also used estimates of shoaling rate, prevalence, and host density (see SI). We were able to fit

some outputs to estimates from low predation populations, high predation populations, and the

ratio of the two (representing the impact of changing predation regime). For others, we only have

estimates of one of these three. This scheme weights the model training toward quantities that are

well-characterized. See ED Table 2 for estimates used and sources.

We optimized the model's fit to both these empirical parameter estimates and field-estimated

model outputs with the evolutionary algorithm[58]. We began with parameters at their estimated

values (for $a$, $d$, $k_1$, $k_2$, $\gamma$, and $z$). Initial values of other parameters were chosen arbitrarily [$q = 1$ x

$10^{-2}$, $(Pc_1)_{Low} = 1$ x $10^{-2}$, $(Pc_1)_{Hi} = 3$ x $10^{-2}$, $\sigma = 0.5$]. Each parameter was mutated by a factor of

$10^{\wedge}x$ where $x$ is a single sample from the normal distribution $N(0,0.05)$ to create a parameter set;

each parameter set formed one strategy in the evolutionary algorithm. One un-mutated strategy

and 299 mutated strategies were evaluated in one "generation" of the algorithm by their summed

relative error from all inputs and outputs with known estimates: summed relative error of

strategy x $= |a_x\text{-}a_{data}|/a_{data} +\ldots+|cT_{Low,x}\text{-}cT_{Low,data}|/cT_{Low,data}$. There are four constrained

parameters, four unconstrained parameters, and 8 independent scored outputs (e.g., $p_{Low}$ is

independent from $p_{Hi}$ but not $p_{Hi}/p_{Low}$). The strategy with the lowest summed relative error was

passed to the next generation of the algorithm along with 299 mutated versions of itself. This

process continues for 20 generations of the algorithm, as the model had asymptotically

converged to a fit, yielding the parameter values used for the model (see ED Table 2).

Sensitivity analysis. We performed a sensitivity analysis to determine how key outcomes

responded to parameter values. We performed this analysis for coCSS parasite prevalence, host

density, parasite virulence, host contact rate, overall host mortality, transmission rates, and the

ratios of these quantities between high and low predation populations. We also examine two

other key 'proportion change metrics' representing how key factors change between the two,

focal predation levels. First was the proportion of total host mortality change with predation

regime that is caused by increased parasite-induced mortality: $(p_{Hi}v_{Hi}\text{-}p_{Low}v_{Low})/[(d+p_{Hi}v_{Hi}$

$+P_{High}c_1/c_{Hi})\text{ - }(d+p_{Low}v_{Low} +P_{Low} c_1/c_{Low})]$. Second was the proportion of change in

coevolutionary virulence across predation regime that is via predator's non-consumptive effects

(evolution of higher shoaling rate). We defined this proportion as 1- (virulence change when

predators only have consumptive effects/virulence change when predators have consumptive and

non-consumptive effects); the virulence change when predators only have consumptive effects

was calculated as mean($v_{CSS\ high\ P,\ low\ c}$ - $v_{CSS\ Plow,\ low\ c}$, $v_{CSS\ Phigh,\ high\ c}$ - $v_{CSS\ Plow,\ high\ c}$) and virulence

change when predators have both is $v_{coCSS\ Phigh}$ - $v_{coCSS\ Plow}$ (Fig. 3). To generate random

parameter sets, we used Latin Hypercube sampling with the 'lhs' R package[59] and then computed

the partial rank correlation coefficients of the parameters with respect to each model outcome of

interest (coevolutionary outcomes and key proportion change metrics) for $5 \times 10^3$ runs using the

R package 'sensitivity'[60]. We provide the code for the model and sensitivity analysis in the SI.

*4. Analysis of shoaling rate, prevalence, and host density*

We extracted estimates of shoaling rate, prevalence, and host density from the literature and

supplemented them with our own, new data on prevalence and shoaling rate (see SI). We used

mixed models to test if guppy shoaling behaviour, *Gyrodactylus* spp. prevalence, and guppy

density differed between predation regimes. In these models we used predation regime as a fixed

effect along with river and year as random effects. We only included rivers with paired estimates

from high and low predation populations because of variation across rivers. Each population

average for a given site+year was one data point in these analyses. For % time shoaling (22

estimates, 4 rivers)[61–64] and prevalence (107 estimates, 11 rivers)[30,32,33,43,57,65,66], we used a

generalized linear mixed model beta regression with a logit link. For host density (23 estimates,

4 rivers)[67–71], we used a linear mixed model.

*5.  Data availability statement*

The authors declare that all data used in this study are available within this publication and its

supplementary files.

*6.  Code availability statement*

Custom computer code used to conduct the theoretical modelling and statistical tests is

included in the supporting information.

*7.  Animal use ethics statement*

All collections and fish handling protocols were approved by IACUC protocol 18072155

and 21069471 from the corresponding author's institution.  The permit to collect and export

guppies was granted by the Director of Fisheries in the Ministry of Agriculture, Land and

Fisheries Division, Aquaculture Unit of the Republic of Trinidad and Tobago on 2/26/2020

(copy available on request).  The United States Import/Export License for live and preserved fish

was issued by the U.S. Fish and Wildlife Service, Office of Law Enforcement Permit number

A107080 on February 21, 2020 for the shipment of live guppies to the corresponding author's

institution. Samples were declared and cleared via USFWS form 3-177 on 03/11/2020.

**References**

1.    Anderson, R. M. & May, R. M. Coevolution of hosts and parasites. *Parasitology* **85**, 411–

426 (1982).

2.    Choo, K., Williams, P. D. & Day, T. Host mortality, predation and the evolution of parasite

virulence. *Ecol. Lett.* **6**, 310–315 (2003).

3.    Williams, P. D. & Day, T. Interactions between sources of mortality and the evolution of

parasite virulence. *Proceedings of the Royal Society of London. Series B: Biological*

*Sciences* **268**, 2331–2337 (2001).

4.    Gandon, S., Jansen, V. A. A. & Van Baalen, M. Host life history and the evolution of

parasite virulence. *Evolution* **55**, 1056–1062 (2001).

5.    Peacor, S. D. & Werner, E. E. The contribution of trait-mediated indirect effects to the net

effects of a predator. *Proceedings of the National Academy of Sciences* **98**, 3904–3908

(2001).

6.    Krause, J. & Ruxton, G. D. *Living in Groups*. (Oxford University Press, 2002).

7.    Prado, F., Sheih, A., West, J. D. & Kerr, B. Coevolutionary cycling of host sociality and

pathogen virulence in contact networks. *J. Theor. Biol.* **261**, 561–569 (2009).

8.    Everard, M., Johnston, P., Santillo, D. & Staddon, C. The role of ecosystems in mitigation

and management of Covid-19 and other zoonoses. *Environ. Sci. Policy* **111**, 7–17 (2020).

9.    Fisher, M. C. *et al.* Emerging fungal threats to animal, plant and ecosystem health. *Nature*

**484**, 186–194 (2012).

10.  Handel, A. & Rohani, P. Crossing the scale from within-host infection dynamics to

between-host transmission fitness: a discussion of current assumptions and knowledge.

*Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370**, (2015).

11.  Alizon, S., Hurford, A., Mideo, N. & Van Baalen, M. Virulence evolution and the trade-off

hypothesis: history, current state of affairs and the future. *J. Evol. Biol.* **22**, 245–259 (2009).

12.  Cressler, C. E., McLeod, D. V., Rozins, C., van den Hoogen, J. & Day, T. The adaptive

evolution of virulence: a review of theoretical predictions and empirical tests. *Parasitology*

**143**, 915–930 (2016).

13.  McKay, B., Ebell, M., Dale, A. P., Shen, Y. & Handel, A. Virulence-mediated

infectiousness and activity trade-offs and their impact on transmission potential of influenza

patients. *Proc. Biol. Sci.* **287**, 20200496 (2020).

14.  Bonneaud, C. *et al.* Experimental evidence for stabilizing selection on virulence in a

bacterial pathogen. *Evolution Letters* **73**, 636 (2020).

15.  De Roode, J. C., Yates, A. J. & Altizer, S. Virulence-transmission trade-offs and population

divergence in virulence in a naturally occurring butterfly parasite. *Proceedings of the*

*National Academy of Sciences* **105**, 7489–7494 (2008).

16.  Fraser, C., Hollingsworth, T. D., Chapman, R., de Wolf, F. & Hanage, W. P. Variation in

HIV-1 set-point viral load: epidemiological analysis and an evolutionary hypothesis.

*Proceedings of the National Academy of Sciences* **104**, 17441–17446 (2007).

17.  Herre, E. A. Population structure and the evolution of virulence in nematode parasites of fig

wasps. *Science* **259**, 1442–1445 (1993).

18.  Boots, M. & Mealor, M. Local interactions select for lower pathogen infectivity. *Science*

**315**, 1284–1286 (2007).

19. Bull, J. J. & Lauring, A. S. Theory and empiricism in virulence evolution. *PLoS Pathog.* **10**,

e1004387 (2014).

20. Alizon, S., de Roode, J. C. & Michalakis, Y. Multiple infections and the evolution of

virulence. *Ecol. Lett.* **16**, 556–567 (2013).

21. Rohr, J. R. *et al.* Predator diversity, intraguild predation, and indirect effects drive parasite

transmission. *Proceedings of the National Academy of Sciences* **112**, 3008–3013 (2015).

22. Seppälä, O., Karvonen, A. & Valtonen, E. T. Shoaling behaviour of fish under parasitism

and predation risk. *Anim. Behav.* **75**, 145–150 (2008).

23. Lopez, L. K. & Duffy, M. A. Mechanisms by which predators mediate host–parasite

interactions in aquatic systems. *Trends Parasitol.* (2021) doi:10.1016/j.pt.2021.06.006.

24. Rigby, M. C. & Jokela, J. Predator avoidance and immune defence: costs and trade–offs in

snails. *Proceedings of the Royal Society of London B: Biological Sciences* **267**, 171–176

(2000).

25. Godin, J.-G. J. Antipredator function of shoaling in teleost fishes: a selective review. *Nat.*

*Can.* **113**, 241–250 (1986).

26. Seghers, B. H. Schooling behavior in the guppy (*Poecilia reticulata*): An evolutionary

response to predation. *Evolution* **28**, 486–489 (1974).

27. Gandon, S., van Baalen, M. & Jansen, V. A. A. The evolution of parasite virulence,

superinfection, and host resistance. *Am. Nat.* **159**, 658–669 (2002).

28. Magurran, A. E. *Evolutionary ecology: the Trinidadian guppy*. 1–224 (Oxford Univeristy

Press, 2005).

29. Magurran, A. E. & Seghers, B. H. Variation in schooling and aggression amongst guppy

(*Poecilia reticulata*) populations in Trinidad. *Behaviour* **118**, 214–234 (1991).

30.  Stephenson, J. F., van Oosterhout, C., Mohammed, R. S. & Cable, J. Parasites of

Trinidadian guppies: evidence for sex- and age-specific trait-mediated indirect effects of

predators. *Ecology* **96**, 489–498 (2015).

31.  Huizinga, M., Ghalambor, C. K. & Reznick, D. N. The genetic and environmental basis of

adaptive differences in shoaling behaviour among populations of Trinidadian guppies,

*Poecilia reticulata . J. Evol. Biol.* **22**, 1860–1866 (2009).

32.  Gotanda, K. M. *et al.* Adding parasites to the guppy-predation story: insights from field

surveys. *Oecologia* **172**, 155–166 (2013).

33.  Fraser, B. A., Ramnarine, I. W. & Neff, B. D. Temporal variation at the MHC class IIb in

wild populations of the guppy (*Poecilia reticulata*). *Evolution* **64**, 2086–2096 (2010).

34.  Johnson, M. B., Lafferty, K. D., van Oosterhout, C. & Cable, J. Parasite transmission in

social interacting hosts: monogenean epidemics in guppies. *PLoS One* **6**, e22634 (2011).

35.  Richards, E. L., van Oosterhout, C. & Cable, J. Sex-specific differences in shoaling affect

parasite transmission in guppies. *PLoS One* **5**, e13285 (2010).

36.  Cable, J. & Van Oosterhout, C. The impact of parasites on the life history evolution of

guppies (*Poecilia reticulata*): The effects of host size on parasite virulence. *Int. J. Parasitol.*

**37**, 1449–1458 (2007).

37.  Reznick, D. N., Butler, M. J., IV, Rodd, F. H. & Ross, P. Life-history evolution in guppies

(*Poecilia reticulata*) 6. Differential mortality as a mechanism for natural selection.

*Evolution* **50**, 1651–1660 (1996).

38.  Bonds, M. H., Keenan, D. C., Leidner, A. J. & Rohani, P. Higher disease prevalence can

induce greater sociality: a game theoretic coevolutionary model. *Evolution* **59**, 1859–1866

(2005).

39.  Kerr, B., Neuhauser, C., Bohannan, B. J. M. & Dean, A. M. Local migration promotes

competitive restraint in a host–pathogen'tragedy of the commons'. *Nature* **442**, 75–78

(2006).

40.  Boots, M. & Sasaki, A. 'Small worlds' and the evolution of virulence: infection occurs

locally and at a distance. *Proceedings of the Royal Society of London. Series B: Biological*

*Sciences* **266**, 1933–1938 (1999).

41.  Wild, G., Gardner, A. & West, S. A. Adaptation and the evolution of parasite virulence in a

connected world. *Nature* **459**, 983–986 (2009).

42.  Van Oosterhout, C. *et al.* Balancing selection, random genetic drift, and genetic variation at

the major histocompatibility complex in two wild populations of guppies (Poecilia

reticulata). *Evolution* **60**, 2562–2574 (2006).

43.  van Oosterhout, C. *et al.* Selection by parasites in spate conditions in wild Trinidadian

guppies (*Poecilia reticulata*). *Int. J. Parasitol.* **37**, 805–812 (2007).

44.  Stephenson, J. F., Stevens, M., Troscianko, J. & Jokela, J. The size, symmetry, and color

saturation of a male guppy's ornaments forecast his resistance to parasites. *Am. Nat.* **196**,

000–000 (2020).

45.  Houde, A. E. & Torio, A. J. Effect of parasitic infection on male color pattern and female

choice in guppies. *Behav. Ecol.* **3**, 346–351 (1992).

46.  Godin, J.-G. J. & McDonough, H. E. Predator preference for brightly colored males in the

guppy: a viability cost for a sexually selected trait. *Behav. Ecol.* **14**, 194–200 (2003).

47.  Hawley, D. M., Gibson, A. K., Townsend, A. K., Craft, M. E. & Stephenson, J. F.

Bidirectional interactions between host social behaviour and parasites arise through

ecological and evolutionary processes. *Parasitology* 1–15 (2020).

48.  Briard, L. & Ezenwa, V. O. Parasitism and host social behaviour: a meta-analysis of

insights derived from social network analysis. *Anim. Behav.* **172**, 171–182 (2021).

49.  Tao, H., Li, L., White, M. C., Steel, J. & Lowen, A. C. Influenza A Virus Coinfection

through Transmission Can Support High Levels of Reassortment. *J. Virol.* **89**, 8453–8461

(2015).

**Methods-only references**

50.  Stephenson, J. F. *et al.* Host heterogeneity affects both parasite transmission to and fitness

on subsequent hosts. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **372**, 20160093–20160010

(2017).

51.  Eshel, I. Evolutionary and continuous stability. *J. Theor. Biol.* **103**, 99–111 (1983).

52.  Hurford, A., Cownden, D. & Day, T. Next-generation tools for evolutionary invasion

analyses. *J. R. Soc. Interface* **7**, 561–571 (2009).

53.  Leimar, O. Multidimensional convergence stability. *Evol. Ecol. Res.* **11**, 191–208 (2009).

54.  Reznick, D., Bryant, M. & Holmes, D. The evolution of senescence and post-reproductive

lifespan in guppies (*Poecilia reticulata*). *PLoS Biol.* **4**, e7 (2006).

55.  Stephenson, J. F. Parasite-induced plasticity in host social behaviour depends on sex and

susceptibility. *Biol. Lett.* **15**, (2019).

56.  Lopez, S. Acquired resistance affects male sexual display and female choice in guppies.

*Proceedings of the Royal Society of London. Series B: Biological Sciences* **265**, 717–723

(1998).

57.  Pérez-Jvostov, F., Hendry, A. P., Fussmann, G. F. & Scott, M. E. Are host–parasite

interactions influenced by adaptation to predators? A test with guppies and *Gyrodactylus* in

experimental stream channels. *Oecologia* **170**, 77–88 (2012).

58.   Eiben, A. E. & Smith, J. E. *Introduction to Evolutionary Computing*. (Springer, Berlin,

Heidelberg, 2015).

59.   Carnell, R. *Package "lhs."* (cran.pau.edu.tr, 2016).

60.   Iooss, B., Da Veiga, S., Janon, A. & Pujol, G. *Package 'sensitivity.'* (CRAN, 2015).

61.   Seghers, B. H. & Magurran, A. E. Predator inspection behaviour covaries with schooling

tendency amongst wild guppy, *Poecilia reticulata*, populations in Trinidad. *Behaviour* **128**,

121–134 (1994).

62.   Griffiths, S. W. & Magurran, A. E. Sex and schooling behaviour in the Trinidadian guppy.

*Anim. Behav.* **56**, 689–693 (1998).

63.   Magurran, A. E., Seghers, B. H., Carvalho, G. R. & Shaw, P. W. Behavioural consequences

of an artificial introduction of guppies (*Poecilia reticulata*) in N. Trinidad: evidence for the

evolution of anti-predator behaviour in the wild. *Proceedings of the Royal Society of*

*London. Series B: Biological Sciences* **248**, 117–122 (1992).

64.   Sievers, C. *et al.* Reasons for the invasive success of a guppy (*Poecilia reticulata*)

population in Trinidad. *PLoS One* **7**, e38404 (2012).

65.   Mohammed, R. S. *et al.* Parasite diversity and ecology in a model species, the guppy

(*Poecilia reticulata*) in Trinidad. *Royal Society Open Science* **7**, 191112 (2020).

66.   Lyles, A. M. Genetic variation and susceptibility to parasites: *Poeclia reticulata* infected

with *Gyrodactylus turnbulli*. (Princeton University, 1990).

67.   Reznick, D. & Endler, J. A. The impact of predation on life history evolution in Trinidadian

guppies (*Poecilia reticulata*). *Evolution* **36**, 160–177 (1982).

68.   El-Sabaawi, R. W. *et al.* Assessing the effects of guppy life history evolution on nutrient

recycling: from experiments to the field. *Freshw. Biol.* **60**, 590–601 (2015).

69.  Croft, D. P. *et al.* Mechanisms underlying shoal composition in the Trinidadian guppy,

*Poecilia reticulata*. *Oikos* **100**, 429–438 (2003).

70.  Liley, N. R. & Luyten, P. H. Geographic variation in the sexual behaviour of the guppy,

*Poecilia Reticulata* (Peters). *Behaviour* **95**, 164–179 (1985).

71.  Reznick, D. N. *et al.* Eco-evolutionary feedbacks predict the time course of rapid life-

history evolution. *Am. Nat.* **194**, 671–692 (2019).

**Extended Data Table 1. Coinfection rates in the wild.** *For each site (river+predation regime),*

*we genotyped multiple worms per a sample of fish hosting more than one worm. Coinfection*

*within species = (no. fish with >1 genotype of species x/no. fish with > 1 worm of species x) × %*

*of infections with >1 worm. Overall coinfection = (no. fish with at least two different worms/no.*

*fish with at least two worms) × % of infections with >1 worm. Infections were diverse overall.*

*Coinfection by identical multilocus genotypes was rare (31 out of the 73 worms possible).* G.t. =

G. turnbulli *and* G.b. = G. bullatarudis.

| Site | | Sample size | | Species presence on fish | | | Infections >1 worm | Coinfection within species | | Overall coinfection rate |
|---|---|---|---|---|---|---|---|---|---|---|
| River | Predation | Fish | Worms/fish (mean±SE) | %*G.t.* only | %*G.b.* only | %Both | % | *G.t.* | *G.b.* | % |
| Aripo | High | 7 | 2.43±0.20 | 14.3 | 85.7 | 0 | 70.8 | 0 | 35.4 | 30.3 |
| Aripo | Low | 4 | 3.25±0.75 | 100 | 0 | 0 | 58.2 | 14.6 | NA | 14.6 |
| Guanapo | High | 7 | 2.29±0.18 | 14.3 | 85.7 | 0 | 62.5 | 62.5 | 52.1 | 53.6 |
| Guanapo | Low | 5 | 2 | 80 | 20 | 0 | 61.9 | 15.5 | 0 | 12.4 |
| Santa Cruz | High | 7 | 4.86±1.30 | 0 | 71.4 | 28.6 | 92.6 | 92.6 | 92.6 | 92.6 |
| Santa Cruz | Low | 3 | 3±1 | 66.7 | 33.3 | 0 | 61.7 | 30.9 | 61.7 | 41.1 |

725
726
727

728 **Extended Data Table 2. Meaning, value, and units for state variables, parameters, or**

729 **outputs.** *Training values may be for a parameter, an output, or outputs related to a parameter*

730 *or output. Training values were used in the model fitting and their sources are provided. Low/hi.*

731 *pred. indicates predation regime. Values result from the model training. Note that the model was*

732 *not trained with evolved virulence data but instead validated against it.*

| Symbol | Meaning | Value and/or units | Training value | Source |
|---|---|---|---|---|
| $S$ | Susceptible host density | hosts m$^{-2}$ | | |
| $I$ | Infected host density | hosts m$^{-2}$ | | |
| $R$ | Recovered host density | hosts m$^{-2}$ | | |
| $(S+I+R)_{Low}$ | Total host density, low pred. | 5.61 hosts m$^{-2}$ | 8.19 | 67–71 |
| $(S+I+R)_{Hi}$ | Total host density, hi. pred. | 4.69 hosts m$^{-2}$ | 4.68 | 67–71 |
| $(S+I+R)_{Hi}/(S+I+R)_{Low}$ | Host density ratio | 0.837 hosts m$^{-2}$ | 0.571 | 67–71 |
| $p$ | Prevalence: $p = I/(S+I+R)$ | unitless | | |
| $p_{Low}$ | Prevalence, low pred. | 0.288 | 0.311 | 30,32,33,43,57,66 |
| $p_{Hi}$ | Prevalence, hi. pred. | 0.400 | 0.394 | 30,32,33,43,57,66 |
| $p_{Hi}/p_{Low}$ | Prevalence ratio | 1.39 | 1.27 | 30,32,33,43,57,66 |
| $a$ | Maximum host per-capita fecundity | 0.106 day$^{-1}$ | $a = 0.106$ | 54 |
| $c$ | Host social contact rate | 1.7-5.5 m$^2$ host$^{-1}$ day$^{-1}$ | | |
| $c_{Hi}/c_{Low}$ | Contact rate ratio | 2.23 | 1.93 | 61–64 |
| $c_1$ | Scaling parameter for contact rate | 1 m$^2$ host$^{-1}$ day$^{-1}$ | | |
| $d$ | Background host death rate | 1.30x10$^{-3}$ day$^{-1}$ | $d = 1.41 \times 10^{-3}$ | 54 |
| $k_1$ | Trade-off parameter: Virulence with transmissibility = 1 | 1.38x10$^6$ day$^{-1}$ | See text | Fig. 2c |
| $k_2$ | Trade-off parameter: Exponent | 3.61 | See text | Fig. 2c |
| $P$ | Predation regime | 7.4-33.0x10$^{-3}$ day$^{-1}$ | | |
| $(d+Pc_1/c+pv)_{Low}$ | Overall host mortality, low pred. | 0.011 day$^{-1}$ | 0.011 | 37 |
| $(d+Pc_1/c+pv)_{Hi}$ | Overall host mortality, hi. pred. | 0.026 day$^{-1}$ | 0.026 | 37 |
| $(d+Pc_1/c+pv)_{Hi}/(d+Pc_1/c+pv)_{Low}$ | Overall host mortality ratio | 2.46 | 2.47 | 37 |
| $q$ | Density dependence of host fecundity | 0.017 m$^2$ host$^{-1}$ day$^{-1}$ | | |
| $T$ | Transmissibility given contact | 6.2-8.81x10$^{-3}$ | | |
| $(cT)_{Low}$ | Transmission rate, low pred. | 0.015 m$^2$ host$^{-1}$ day$^{-1}$ | 0.014 | 57 |
| $cTI$ | Force of infection | day$^{-1}$ | NA | |
| $v$ | Mortality virulence | 0.015-0.053 day$^{-1}$ | Intentionally not trained | |
| $v_{Low}$ | Virulence, low pred. | 0.020 day$^{-1}$ | Validated: 0.021 | Fig. 4j |
| $v_{Hi}$ | Virulence, hi. pred. | 0.044 day$^{-1}$ | Validated: 0.041 | Fig. 4j |
| $z$ | Rate of waning of host immunity | 0.033 day$^{-1}$ | $z = 0.033$ | 56 |
| $\gamma$ | Infected host recovery rate | 0.020 day$^{-1}$ | $y = 0.033$ | 55 |
| $\sigma$ | Probability of successful superinfection given transmission | 1.21 | | |

733
734

**Extended data figure 1. Virulence can select for increased contact rate.** *Hosts can evolve*

*increasing contact rate in response to increased virulence, especially at very high virulence*

*(beyond range used in main text). **a**, Increasing virulence (and transmissibility along the trade-*

*off) can decrease prevalence. **b**, Overall parasite-induced mortality can decrease if prevalence*

*declines sharply enough. This decrease occurs because, while parasites are very virulent, very*

*few hosts are infected and suffering that virulence. **c**, At high virulence, increasing virulence can*

*select for higher host contact rates.* c = 2 *and* P = 0.074 *used. All other parameters at default.*

**Extended data Figure 2. Host evolution in response to increasing predation causes parasite-**

**induced mortality (red curves) to increase more than predator-induced mortality (black**

**curves). a,** *Without host evolution (contact rate,* c*, set to the green point in Figs. 3 while*

*parasites evolve to some CSS), parasite-induced mortality declines with predation while*

*predator-induced mortality increases. Death from background sources (*d*, grey line) does not*

*change.* ***b,*** *This trend is similar for a higher* c *(set to high, blue point in Fig. 3).* ***c,*** *When hosts*

*evolve increasing* c *with increasing predation (coCSS curve in Fig. 3), parasite-induced*

*mortality increases more than predator-induced mortality. This pattern is due to host evolution*

*and is qualitatively unchanged if hosts evolve but parasites do not.*

**Extended data Figure 3. Sensitivity analysis shows the impact of parameters on key outputs**

**and the range for those outputs.** *Key model outputs: the proportion of the mortality increase*

*from increasing predation that is via increased parasite-induced mortality (**a-b**) and the*

*proportion of the virulence increase from increasing predation that is via increased contact rate*

*(**c-d**). Vertical lines are 95% confidence intervals. (**a & c**). **a,** Notable among other parameters,*

*the trade-off exponent,* $k_2$, *strongly increases the proportion of mortality increase due to*

*parasites. **b,** The mortality increase via parasites is typically well over half (0.5). **c,** Both* $k_2$ *and*

σ, *the superinfection parameter, strongly increase the proportion of virulence increase via*

*contact rate. **d,** The proportion of virulence increase via contact rate is typically well over half.*

*Note here that* $P_H = P_{Hi}$ *and* $P_L = P_{Low}$.

**Extended data table 3. Parameter effects based on 95% CIs.** *For each model output (first*

*column), we report which parameters increased (second column) or decreased (third column)*

*that model output in the sensitivity analysis. A positive or negative effect is only reported if the*

*95% CI for that parameter's effect did not overlap 0. Trait outputs are the coCSS trait values.*

| Model output | Parameters w. + effect | Parameters w. − effect |
|---|---|---|
| $p_{Low}$ | $a, k_2, P_{Low}, z$ | $k_1, q, \gamma, \sigma$ |
| $p_{Hi}$ | $a, k_2, P_{Hi}, z$ | $k_1, q, \gamma, \sigma$ |
| $p_{Hi}/p_{Low}$ | $d, k_1, P_{Hi}, q, z, \gamma, \sigma$ | $a, k_2, P_{Low}$ |
| $(S+I+R)_{Low}$ | $a, k_1, \gamma$ | $k_2, P_{Low}, q$ |
| $(S+I+R)_{Hi}$ | $a, k_1, \gamma$ | $k_2, P_{Hi}, q$ |
| $(S+I+R)_{Hi}/(S+I+R)_{Low}$ | $a, k_1, P_{Low}, q, \gamma$ | $k_2, P_{Hi}$ |
| $v_{Low}$ | $a, k_2, P_{Low}, \gamma, \sigma$ | $k_1, q, z$ |
| $v_{Hi}$ | $a, k_2, P_{Hi}, \gamma, \sigma$ | $k_1, q, z$ |
| $v_{Hi}/v_{Low}$ | $a, k_2, P_{Hi}$ | $P_{Low}, \gamma, z$ |
| $c_{Low}$ | $k_1, P_{Low}, q, \gamma$ | $a, k_2, z, \sigma$ |
| $c_{Hi}$ | $k_1, P_{Hi}, q, \gamma$ | $a, k_2, z, \sigma$ |
| $c_{Hi}/c_{Low}$ | $k_2, P_{Hi}$ | $P_{Low}, q, z, \sigma$ |
| $(d+Pc_1/c+pv)_{Low}$ | $a, d, k_2, P_{Low}$ | $k_1, q, \gamma$ |
| $(d+Pc_1/c+pv)_{Hi}$ | $a, k_2, P_{Hi}$ | $k_1, q, \gamma$ |
| $(d+Pc_1/c+pv)_{Hi}/(d+Pc_1/c+pv)_{Low}$ | $a, k_1, P_{Hi}, \gamma$ | $d, k_2, P_{Low}$ |
| Proportion of mortality change via parasites | $a, k_2, P_{Hi}, P_{Low}, \gamma$ | $k_1, q, z, \sigma$ |
| $cT_{Low}$ | $k_2, P_{Low}, q, \gamma$ | $a, k_1, z, \sigma$ |
| $cT_{Hi}$ | $k_2, P_{Hi}, q, \gamma$ | $a, k_1, z, \sigma$ |
| $cT_{Hi}/cT_{Low}$ | $k_2, P_{Hi}$ | $P_{Low}, q, z, \sigma$ |
| Proportion of virulence change via contact rate | $k_2, P_{Hi}, P_{Low}, z, \sigma$ | $a, k_1, q, \gamma$ |