The Arabidopsis SAC9 Enzyme is enriched in a cortical population of early endosomes and restricts PI(4,5)P$_2$ at the Plasma Membrane

Alexis Lebecq†[1], Mehdi Doumane†[1], Aurélie Fangain[1], Vincent Bayle[1], Jia Xuan Leong[4], Frédérique Rozier[1], Maria del Mar Marquès-Bueno[1,3], Laia Armengot[1,7], Romain P. Boisseau[2], Mathilde Laetitia Audrey Simon[1], Mirita Franz-Wachtel[6], Boris Macek[6], Suayb Üstün[4,5], Yvon Jaillais[1*], Marie-Cécile Caillaud[1*]

[1] Laboratoire Reproduction et Développement des Plantes, Université de Lyon, ENS de Lyon, CNRS, INRA, F-69342 Lyon, France.

[2] Division of Biological Science, University of Montana, Missoula, MT 59812, USA.

[3] Current address: Department of Molecular Genetics, Center for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB, Bellaterra (Cerdanyola del Vallès), 08193, Barcelona, Spain.

[4] University of Tübingen, Center for Plant Molecular Biology (ZMBP), 72076 Tübingen, Germany

[5] Faculty of Biology & Biotechnology, Ruhr-University Bochum, 44780 Bochum, Germany

[6] Interfaculty Institute for Cell Biology, Department of Quantitative Proteomics, University of Tübingen, 72076 Tübingen, Germany

[7] Current address: Genetics Section, Universitat de Barcelona (UB), 08028 Barcelona, Catalonia, Spain and Centre for Research in Agricultural Genomics (CSIC-IRTA-UAB-UB). Edifici CRAG, Campus UAB, 08193, Bellaterra, Catalonia,Spain.

† The authors contributed equally to this work

*Corresponding authors: yvon.jaillais@ens-lyon.fr; marie-cecile.caillaud@ens-lyon.fr

Short title: SAC9 prevents the intracellular accumulation of PI(4,5)P$_2$ in plant cells.

One-sentence summary: SAC9 prevents the accumulation of PI(4,5)P$_2$ in intracellular compartments, thereby contributing to clathrin-mediated endocytosis at the plasma membrane and regulating the cortical localization of its protein partner SH3P2.

Keywords: Arabidopsis, endocytic trafficking, phosphoinositides, early endosome, plasma membrane

ABSTRACT

Membrane lipids, and especially phosphoinositides, are differentially enriched within the
eukaryotic endomembrane system. This generates a landmark code by modulating the
properties of each membrane. Phosphatidylinositol 4,5-bisphosphate [PI(4,5)P$_2$] specifically
accumulates at the plasma membrane in yeast, animal and plant cells, where it regulates a
wide range of cellular processes including endocytic trafficking. However, the functional
consequences of mispatterning PI(4,5)P$_2$ in plants are unknown. Here, we functionally
characterized the putative phosphoinositide phosphatase *SUPPRESSOR OF ACTIN9* (*SAC9*)
in *Arabidopsis thaliana* (Arabidopsis). We found that SAC9 depletion led to the ectopic
localization of PI(4,5)P$_2$ on cortical intracellular compartments, which depends on PI4P and
PI(4,5)P$_2$ production at the plasma membrane. SAC9 localizes to a subpopulation of *trans*-
Golgi Network/early endosomes that are enriched in a region close to the cell cortex and that
are coated with clathrin. Furthermore, it interacts and colocalizes with Src Homology 3
Domain Protein 2 (SH3P2), a protein involved in endocytic trafficking. In the absence of
SAC9, SH3P2 localization is altered and the clathrin-mediated endocytosis rate is reduced.
Together, our results highlight the importance of restricting PI(4,5)P$_2$ at the plasma
membrane and illustrate that one of the consequence of PI(4,5)P$_2$ misspatterning in plants is
to impact the endocytic trafficking.

INTRODUCTION

Phosphoinositides constitute a family of low abundance lipids differentially enriched in
the membranes of eukaryotic cells (Platre and Jaillais 2016; Balla 2013; Noack and Jaillais
2020). These versatile lipids can be interconverted into one another. For example,
Phosphatidylinositol 4,5-biphosphate [PI(4,5)P$_2$] is synthetized from Phosphatidylinositol 4-
phosphate (PI4P) by PI4P-5 kinases, and is dephosphorylated into PI4P by PI(4,5)P$_2$ 5-
phosphatase (Noack and Jaillais 2017). Furthermore, PI(4,5)P$_2$ and PI4P are hydrolyzed by
phospholipases C (PLC) into diacylglycerol and soluble phosphorylated inositol (Balla 2013).
PI(4,5)P$_2$ strictly localizes at the plasma membrane in plants and animal cells (van Leeuwen
et al. 2007; Carim et al. 2019; Del Signore et al. 2017; Simon et al. 2014; Ben El Kadhi et al.
2011) despite the plasma membrane being constantly turned-over by endocytosis and

exocytosis. Throughout this paper, we define endocytosis in its broader sense, including the

internalization step *sensus stricto* (i.e., recruitment of cargo and coat components of the

"clathrin- mediated endocytosis" machinery, formation of clathrin-coated pits, scission of the

clathrin-coated vesicles and uncoating) followed by the subsequent transport of lipid and

proteins through the endosomal system. As such, the endocytosis, or here after the endocytic

trafficking, is the process that allows 1) cells to transport particles and molecules across the

plasma membrane and 2) the termination of signaling through transport toward the vacuole

for degradation. At the plasma membrane, $PI(4,5)P_2$ interacts with a variety of extrinsic

membrane proteins such as endocytic protein adaptors (Zhang et al. 2015) and actin-

regulatory proteins (Paez Valencia et al. 2016), which are recruited and/or activated by the

binding to $PI(4,5)P_2$. Therefore, $PI(4,5)P_2$ subcellular patterning is likely critical to regulate

the recruitment of proteins that act at the plasma membrane, and the cellular processes they

mediate, including clathrin-mediated endocytosis.

Consistent with a critical role of $PI(4,5)P_2$ in the recruitment of early clathrin-mediated

endocytosis factors, a *pip5k1 pip5k2* double mutant in *Arabidopsis thaliana* (Arabidopsis)

lacking two ubiquitously expressed PI4P-5 kinases, has abnormal auxin distribution and

defective endocytic trafficking of the transmembrane auxin efflux carriers PIN-FORMED 1

(PIN1) and PIN2 (Tejos et al. 2014; Ischebeck et al. 2013; Mei et al. 2012). Furthermore, the

pip5k1 pip5k2 double mutant has an altered dynamic of CLATRHIN LIGHT CHAIN2

(CLC2), with the density of CLC2 foci at the plasma membrane being reduced in the mutant

(Ischebeck et al. 2013). Overexpression of Arabidopsis PI4P-5 kinase 6 in tip-growing

pollen tubes induced massive aggregation of the plasma membrane in pollen tube tips due to

excessive clathrin-dependent membrane invagination, supporting a role for $PI(4,5)P_2$ in

promoting early stages of clathrin-mediated endocytosis (Zhao et al. 2010). The inducible

overexpression of a highly active human PI4P-5 kinase leads to an increase $PI(4,5)P_2$

production, very strong developmental phenotypes and heightened endocytic trafficking

toward the vacuole (Gujas et al. 2017). In addition, we recently showed that inducible

depletion of the $PI(4,5)P_2$ from the plasma membrane using the iDePP system leads to a

decrease of the fraction of the clathrin adaptor protein AP2-μ and the Src Homology (SH3)-

domain containing protein 2 (SH3P2) at the plasma membrane (Doumane et al. 2021).

Furthermore, FM4-64 uptake experiments confirmed an impact of $PI(4,5)P_2$ depletion from

the plasma membrane on bulk endocytic flow.

In animal cells, several $PI(4,5)P_2$ phosphatases are required for the late stages of clathrin-
mediated endocytosis (He et al. 2017; Pirruccello et al. 2014). Many $PI(4,5)P_2$ phosphatases
belong to the 5-Phosphatase enzyme family, including OCRL and synaptojanins (Syn1/2) in
animals, and synaptojanin-like proteins (Inp51p/Snjl1p, Inp52p/Sjl2p and Inp53p/Sjl3p) in
Saccharomyces cerevisiae. The Arabidopsis genome contains 15 genes encoding 5-
phosphatases, but only a few are characterized. Mutation in the 5-phosphatase 9 leads to
osmotic stress tolerance, with reduced reactive oxygen species production and Ca^{2+} influx
(Golani et al. 2013). The 5-phosphatase 6/COTYLEDON VASCULAR PATTERN2 (CVP2)
and 5-phosphatase 7/CVP2 LIKE 1 (CVL1) are specifically required for vascular
differentiation (Rodriguez-Villalon et al. 2015; Carland and Nelson 2009; Carland and
Nelson 2004). Finally, the *5-phosphatase 15/FRAGILE FIBER 3 (FRA3)* is expressed in
developing fibers and vascular cells, which is consistent with the defective fiber and vessel
phenotypes seen in the loss-of-function *fra3* mutant (Zhong et al. 2004).

Proteins containing SUPPRESSOR OF ACTIN (or Sac1-like) domains constitute another
family of phosphoinositide phosphatases (Zhong and Ye 2003). In Arabidopsis, there are nine
SAC proteins, forming three clades (Zhong and Ye 2003). The first clade is composed of
SAC1, a $PI(3,5)P_2$ 5-phosphatase (Zhong et al. 2005), and its relatives SAC2 to 5, putative
$PI(3,5)P_2$ 5-phosphatases (Novakova et al. 2014). The second clade corresponds to
SAC7/RHD4 a PI4P 4-phosphatase (Thole et al. 2008) and its relatives SAC6 and SAC8
putative PI4P 4-phosphatases (Song et al. 2021). The third clade is composed of a single
member, a plant-specific protein called SAC9. SAC9 has a unique structure, with a SAC
phosphoinositide phosphatase domain at its N-terminus, immediately followed by a putative
protein/protein interaction domain (WW domain), and a long C-terminal region of 1104
amino acids where a putative coil-coiled domain is predicted (Figure 1A, (Zhong and Ye
2003). The *sac9* mutant is dwarf, it constitutively accumulates anthocyanins and it expresses
genes from stress response pathways (Williams et al. 2005). Loss-of-function alleles of *SAC9*
display a three-fold increase in $PI(4,5)P_2$ content, together with a decrease of PI4P level,
suggesting that it acts as a $PI(4,5)P_2$ 5-phosphatase in planta (Williams et al. 2005).

Using *in vivo* confocal microscopy, we found that SAC9 localizes in a subpopulation of
trans-Golgi network/early endosomes (TGN/EEs) enriched in a region close to the cell
cortex. Loss of SAC9 results into $PI(4,5)P_2$ mis-patterning at the subcellular level, leading to
the accumulation of $PI(4,5)P_2$ in subcortical compartments associated with TGN/EEs. Similar

cellular and developmental phenotypes were observed when SAC9 was mutated on its
putative catalytic cysteine, suggesting that the phosphoinositide phosphatase activity of
SAC9 is required for function. SAC9 interacts and colocalizes with the endocytic component
SH3P2. In the absence of SAC9, and therefore when the patterning of the $PI(4,5)P_2$ is
compromised, SH3P2 localization is affected and the clathrin-mediated endocytosis is
significantly reduced. Thus, SAC9 is required to maintain efficient endocytic uptake,
highlighting the importance of restricting the $PI(4,5)P_2$ pool to the plasma membrane.

RESULTS

The cysteine 459 in the catalytic domain of SAC9 is required for SAC9 function

We investigated the root phenotype in the already described mutant alleles of SAC9
(Vollmer et al. 2011; Williams et al. 2005). As previously described, *sac9-1$^{-/-}$* and *sac9-3$^{-/-}$*
knock-out mutants are two times shorter compared to the Wild-type (WT) Col-0, at 12 days
post-germination (dpg; Figure 1A to 1C). We also observed a three-time decrease in the
lateral root density of *sac9-1$^{-/-}$* and *sac9-3$^{-/-}$* compared to WT plants (Figure 1B to 1D). To
confirm that the phenotypes observed were due to the loss-of-function of SAC9, we
generated Arabidopsis lines expressing SAC9 fused to a fluorescent protein (mCIT-SAC9
and TdTOM-SAC9) under SAC9 native promoter (*SAC9pro)* or the *UBIQUITIN10* promoter
(*UBQ10pro*), respectively (Figure 1, Figure 1-figure supplement 1A, 1B). We found that both
SAC9pro:mCIT-SAC9 and *UBQ10pro:TdTOM-SAC9* rescued *sac9-3$^{-/-}$* mutant phenotypes in
two independent homozygous T3 transgenic lines for each construct (Figure 1B, 1C; Figure
1-supplemental Figure 1A). These results indicate that the root phenotypes described above
are caused by *SAC9* loss-of-function, and that N-terminally tagged SAC9 proteins are
functional.

Next, we mutated the cysteine in the conserved C-x(5)-R-[T/S] catalytic motif found in all
SAC domain-containing phosphoinositide phosphatase (Hsu and Mao, 2015, 2013; Figure 1A
and Figure 1-figure supplement 1A). Such cysteine-to-alanine substitution were shown to
block the catalytic phosphatase activity of other SAC domain-containing proteins (Manford
et al. 2010; Tani and Kuge 2014), and thus mCIT-SAC9^{C459A} is a putative catalytically dead
version of the enzyme. By contrast to wild-type mCIT-SAC9, we could not find any

transgenic lines expressing *SAC9pro:mCIT-SAC9*^{C459A} that were able to rescue the *sac9-3*
phenotype, out of 24 independent lines analyzed in T1 (Figure 1B, 1C, Figure 1-
Supplemental Figure 1A, 1B). Further analyses on two independent T3 homozygous lines
confirmed these initial results and showed that mCIT-SAC9^{C459A} fusions were stable and
accumulated to similar extent as wild-type mCIT-SAC9 (Figure 1B, 1C, Figure 1-
Supplemental Figure 1A, 1B). Thus, the putative catalytic cysteine, C459, is required for
SAC9 function but not for the stability of the protein, suggesting that the phosphatase activity
of SAC9 is participating in the observed phenotypes.

SAC9 localizes to a population of early endosomes close to the plasma membrane

Some phosphoinositide-phosphatases such as Metazoan Sac1 and yeast Sac1p, are able to
dephosphorylate *in vitro* several phosphoinositide species, but display a narrower specificity
in vivo (Hughes et al. 2000; Guo et al. 1999; Rivas et al. 1999). Also, some enzymes involved
in phosphoinositide metabolism, such as the yeast PI 4-kinases Stt4p and Pik1p, specifically
impact distinct pools of a given phosphoinositide species depending on their subcellular
localization (Roy and Levine 2004; Yoshida et al. 1994; Flanagan et al. 1993). Using
homozygous T3 transgenic lines expressing the functional *SAC9pro:mCIT-SAC9* or
Ub10pro:TdTOM-SAC9 constructs, we assessed the subcellular distribution of SAC9 in
rescued *sac9-3*$^{-/-}$ homozygous plants using live-cell fluorescence imaging. In meristematic
epidermal cells of Arabidopsis roots, mCIT-SAC9 and TdTOM-SAC9 were mainly diffused
in the cytosol and excluded from the nucleus (Figure 2A to 2C, Figure 2-figure supplement
1A). At the cortex of the cell, in close vicinity to the plasma membrane (Zi focal plane,
Figure 2A; Figure 2-Supplemental Figure 1B), mCIT-SAC9 localized to a cortical population
of mobile dotty structures (Figure 2B, Video 1). To assess whether this discrete subcellular
localization was relevant for SAC9 function, we generated transgenic lines expressing a
mCIT-SAC9 variant in which the predicted coil-coiled motif was deleted, under the native
promoter (Figure 1A, *SAC9pro:mCIT-SAC9*$^{\Delta CC}$). Analyses of 24 independent T1 revealed
that mCIT-SAC9$^{\Delta CC}$ did not complement sac9-3$^{-/-}$ dwarf phenotype (Figure 1-figure
supplement 1A). Similar to mCIT-SAC9^{C459A}, these results were confirmed on two
independents homozygous T3 lines and western blot analyses showed that mCIT-SAC9$^{\Delta CC}$
accumulated to similar extent as the wild-type mCit-SAC9 in each of these lines (Figure 1-
figure supplement 1B). However, mCIT-SAC9$^{\Delta CC}$ did not localize to endosome and remained

entirely cytosolic (Figure 2B, Figure 2-Supplemental Figure 1C). Thus, even though SAC9
localization in cortical intracellular compartments is discreet, and only slightly enriched
compared to its cytosolic localization, it appears critical for function. Furthermore, this
analysis excludes a scenario in which the mCit-SAC9-containing compartments are cytosolic
densities, because they are not seen with a cytosolic mutant of SAC9.

The putative catalytically dead version, mCIT-SAC9^{C459A}, accumulated to a lesser extent
in the cytosol compared to the native form. In addition, SAC9^{C459A} showed a three-fold
increase in the number of labeled dotty structures compared with the signal collected for the
wild-type fusion proteins mCIT-SAC9 and TdTOM-SAC9 (Figure 2C to 2D). This result
suggests that the putative catalytic cysteine of SAC9 is required for the dynamic interaction
of SAC9 with intracellular membranes. However, co-visualization of TdTOM-SAC9 and
mCIT-SAC9^{C459A} (Figure 2E) in Col-0 background demonstrated that all TdTOM-SAC9
dotty structures colocalized with mCIT-SAC9^{C459A} at the cortex of the cell.

By comparing the signal observed at the cortex of the cell (Zi) with the signal collected at
the center of the cell (Zii), we observed and quantified that mCIT-SAC9 labelled more
intracellular compartments close to the plasma membrane than at a distal position, while this
bias was less pronounced for FM4-64 labelled compartments (Figure 2F to 2H). Similarly,
when imaging cells in their cortical part close to the plasma membrane (Zi), the mCIT-SAC9
and mCIT-SAC9^{C459A} signal was concentrated in intracellular compartments at the close
vicinity with the plasma membrane (Figure 2F, 2I). These results suggest a function for
SAC9 in regulating phosphoinositide homeostasis either at or in the close vicinity of the
plasma membrane.

We next investigated the nature of the intracellular structures labeled by mCIT-SAC9.
Both mCIT-SAC9 and mCIT-SAC9^{C459A} colocalized with the amphiphilic styryl dye (FM4-
64) stained endosomal compartments at the cell's cortex, just beneath the plasma membrane,
while the soluble mCIT-SAC9$^{\Delta CC}$ did not (Figure 3A and 3B, Figure 2-Supplemental Figure
1B, 1C; Figure 3-Supplemental Figure 1A). Early endosomes/TGN (EE/TGN) are sensitive
to BFA while late endosomes (LE/MVB) are not (Takagi and Uemura 2018). We observed
that Brefeldin A (BFA) treatment led to the aggregation of mCIT-SAC9 and mCIT-
SAC9^{C459A} into BFA bodies (Figure 3C and 3D). suggesting that both functional and C459-
mutated SAC9 fusion proteins may localize to EE/TGN compartments.

To get more precise insights into the SAC9's localization at the TGN, we crossed
Arabidopsis lines expressing fluorescent tagged SAC9 and SAC9^{C459A} proteins with
endomembranes markers (Geldner et al. 2009). Using live-cell imaging in root meristematic
cells at a Zi focal plane, we observed and quantified that mCIT-SAC9 colocalized with TGN
markers (CLC2-RFP >79% colocalization; mCH-RabA1g >91%, mCH-VTI12 >67%; Figure
3E and 3F). The mCIT-SAC9 showed very weak colocalization with a late endosome
(LE/MVB) marker (mCH-RabF2a <17%, Figure 3E and 2F), and with a Golgi marker (mCH-
Got1p <1%, Figure 3E and 3F). Similarly, mCIT-SAC9^{C459A} strongly colocalized with
CLC2-RFP and mCH-RAB-A1g markers, whereas it did not colocalized with the Golgi
marker (Figure 3-supplemental Figure C, 3G and 3H). This confirmed that SAC9 and
SAC9^{C459A} fusion proteins localized to the TGN/EE, with the noticeable difference that SAC9
localization in endomembrane compartments was restricted to the cortex of the cell,
suggesting a role of SAC9 in PI(4,5)P$_2$ homeostasis between the plasma membrane and early
endosomal compartments.
201

SAC9 is required to maintain the pool of PI(4,5)P$_2$ at the plasma membrane

It was previously reported that Arabidopsis *sac9-1* loss-of-function mutant had a
diminution of PI4P and a three-fold accumulation of PI(4,5)P$_2$, suggesting that SAC9 has a
PI(4,5)P$_2$ 5-phosphatase catalytic activity (Williams et al. 2005). In order to visualize the
effect of the perturbation of the phosphoinositide metabolism in the absence of SAC9, we
introgressed two independent PI(4,5)P$_2$ biosensors (Simon et al. 2014) into *sac9-3$^{-/-}$* genetic
background. As previously described, both PI(4,5)P$_2$ biosensors labeled the plasma
membrane and were excluded form intracellular compartments in wild-type cells (Figure 4A
and 4B). In *sac9-3$^{-/-}$*, mCIT-2xPH^{PLC} and mCIT-TUBBYc not only labeled the plasma
membrane, but also decorated cortical intracellular dotty structures (Figure 4A and 4B). In
addition to the absence of complementation observed macroscopically for *sac9-3* mutants
expressing the putative catalytically-dead *SAC9^{C459A}* variant (Figure 1B and 1C, Figure 1-
figure supplement 1A), the same aberrant localization for the PI(4,5)P$_2$ biosensor mCIT-
TUBBYc was observed in these lines (Figure 4C). These results demonstrate that C459 is
required for the regulation of the pool of PI(4,5)P$_2$ and point out toward a role of SAC9 for
the restriction of the PI(4,5)P$_2$ at the plasma membrane via its catalytic phosphatase activity.
We also observed an increase in the number of intracellular compartments labeled by the

PI4P biosensors (Simon et al. 2014; Simon et al. 2016) mCIT-PH^{FAPP1} and mCIT-P4M^{SidM},

but not for mCIT-PH^{OSBP} (Figure 4A and 4B). This result is consistent with a diminution of

the PI4P pool at the plasma membrane and therefore the relocalization of the PI4P biosensor

in intracellular compartments, as previously reported (Simon et al. 2016). The subcellular

localization of PI3P sensors (mCIT-FYVE^{HRS} and mCIT-PHOX^{p40}) was identical between

Col-0 and $sac9^{-/-}$ cells (Figure 4B, Supplemental Figure 2A). We detected a slight but

significant decrease in the density of intracellular compartments decorated by mCIT-C2^{Lact}

phosphatidylserine biosensor in $sac9$-$3^{-/-}$ (Figure 4B; Figure 4-figure supplement 1). Taken

together, these results indicate that SAC9-depletion leads to a massive change in PI(4,5)P$_2$

subcellular patterning, which is present on intracellular cortical structures instead of only

being present at the plasma membrane.

Previous ultrastructural investigation using electron microscopy, reported a massive

accumulation of vesicles, presumably containing cell wall, at the close vicinity to the plasma

membrane in the $sac9$ mutant (Vollmer et al. 2011). When co-imaging 2xmCH-2xPH^{FAPP1}

together with mCIT-TUBBYc in $sac9$-$3^{-/-}$ (Figure 4D), the dotty structures decorated by the

PI(4,5)P$_2$ biosensor were observed at the cortex of the cell (Zi), in the close vicinity of the

plasma membrane (Figure 4D upper panel), whereas those structures were rarely observed in

the internal part of the cell (Figure 4D lower panel). Moreover, dotty structures containing

PI(4,5)P$_2$ were associated with PI4P-labelled compartments but they did not strictly overlap.

Confocal imaging of $sac9$-$3^{-/-}$ mutant co-expressing mCIT-TUBBYc with endosomal markers

revealed that the dotty structures containing PI(4,5)P$_2$ were found associated but strictly

overlapping with the FM4-64 endocytic tracer as well as TGN markers (Figure 4-figure

supplement 1C). Furthermore, BFA treatment efficiently induced 2xmCH-2xPH^{FAPP1}-positive

BFA bodies in $sac9$-$3^{-/-}$ but did not affect the distribution of mCIT-TUBBYc compartments in

the same cells (Figure 4-figure supplement 1D). Therefore, PI(4,5)P$_2$-containing intracellular

compartments in $sac9$-$3^{-/-}$ are devoid of any ARF GTPase activated by BFA sensitive ARF-

GEF, and are associated with TGN markers (Figure 4-figure supplement 1E). *In vivo* time-

lapse imaging of PI(4,5)P$_2$ biosensor mCIT-TUBBYc and mCIT-2xPH^{PLC} in $sac9$-$3^{-/-}$ mutant

revealed that those intracellular structures were mobile in the cortex of root epidermal cells,

hence, behaving like intracellular compartments (Figure 4-figure supplement 1F, Video 2).

Moreover, in $sac9$-$3^{-/-}$ coexpressing the PI(4,5)P$_2$ biosensor *mCIT-TUBBYc* together with the

non-functional *SAC9pro:tdTOM-SAC9*^{C459A}, mCIT-TUBBYc-labelled intracellular structures

did not strictly colocalized but were observed at the same Z plan in close association with

tdTOM-SAC9^{C459A} (Figure 4-figure supplement 1G).

We next addressed the turnover and the origin of the cortical intracellular PI(4,5)P$_2$

compartment observed in *sac9-3$^{-/-}$*. We previously showed that short-term treatment (15-30

min) with phenyl arsine oxide (PAO), a pan PI 4-kinases inhibitor, significantly depletes

PI4P (Figure 4-figure supplement 2) but not PI(4,5)P$_2$ pools at the plasma membrane of plant

cells, whereas longer treatment (> 60 min) affects the synthesis of both lipids (Figure 4-figure

supplement 2A, Platre et al., 2018; Simon et al., 2016). We used this pharmacological

approach to test the effect of the inhibition of either PI4P, or both PI4P and PI(4,5)P$_2$

synthesis, on *sac9-3$^{-/-}$* anomalous PI(4,5)P$_2$ intracellular compartments (Figure 4-figure

supplement 2A to 2E). Solubilization of mCIT-PH$^{FAPP1-E50A}$ PI4P biosensor in *sac9-3$^{-/-}$* cells

treated for either 30 or 120 min with PAO confirmed the efficient PI4P depletion in both

conditions (Figure 4-figure supplement 2A to 2E). Solubilization of mCIT-TUBBYc

PI(4,5)P$_2$ biosensor in WT after 120 min of PAO exposure, but not after 30 min, confirmed

that an efficient PI(4,5)P$_2$ depletion occurred only for the longest treatment (Figure 4-figure

supplement 2A to 2E). 30 min PAO treatment did not affect anomalous *sac9-3$^{-/-}$* PI(4,5)P$_2$

compartments, but 120 min PAO treatment significantly reduced the number of anomalous

PI(4,5)P$_2$ compartments compared to both 120 min mock treatments or 30 min short

treatment (Figure 4-figure supplement 2A to 2E), showing that intracellular PI(4,5)P$_2$

compartments in *sac9-3$^{-/-}$* are dependent on the PI4P synthesis, itself substrate for PI(4,5)P$_2$

production.

SAC9 is required for efficient endocytic trafficking

Because of the specific localization of SAC9 at the cortex of the cell and its colocalization

with TGN/EE markers, we wondered whether PI(4,5)P$_2$ defective patterning in *sac9-3$^{-/-}$*

correlates with endocytic defects. We counted the numbers of labeled endosomes following

FM4-64 endocytic tracer treatment in cells from WT and *sac9-3$^{-/-}$* seedlings (Rigal et al.

2015). We observed a significant near two-fold decrease in the number of FM4-64-labelled

endosomes per cells in *sac9-3$^{-/-}$* compared to the WT (Figure 5A and 5B, Figure 5-figure

supplement 1A), that was not caused by smaller cells in *sac9-3$^{-/-}$* as the density of FM4-64-

labelled endosomes was also strongly and significantly decreased compared to the WT

(Figure 5-figure supplement 1).We inhibited recycling with BFA, and used FM4-64 tracer to

monitor the endocytic trafficking by measuring the number of BFA bodies labeled by FM4-

64 in Col-0 and *sac9-3*[-/-]. We observed significantly less FM4-64-labelled BFA bodies per

cells in *sac9-3*[-/-] compared to the WT (Figure 5C and 5D, Figure 5-figure supplement 1B),

confirming the lower rate of endocytic trafficking in this mutant. We then assessed whether

SAC9 depletion affected the trafficking of cargo proteins. We, therefore, performed another

BFA assay, but using the integral membrane protein PIN-FORMED2 fused with GFP (PIN2-

GFP) which localizes at the plasma membrane and in intracellular organelles as it

continuously recycles (Armengot et al. 2016). We observed significantly less PIN2-GFP-

labelled BFA bodies per cell in *sac9-3*[-/-] compared to WT (Figure 5E and F). PIN2-GFP being

partially located on intracellular organelles before BFA treatment, the effect observed may,

therefore, indicate an endocytic defect as supported by the others experiments, and/or a more

general defect in trafficking in *sac9-3*[-/-].

To gain further insights into the function of SAC9 in the endocytic trafficking, we

investigated the sensitivity of the *sac9* mutant to pharmacological inhibition of endocytosis.

To this end, we used the recently described ES9-17, which is a specific inhibitor of clathrin-

mediated endocytosis (Dejonghe et al. 2019). We treated wild-type and *sac9-3*[-/-] seedlings

with ES9-17 for 180 min and labeled the plasma membrane and endosomes with FM4-64.

We observed dome-shaped plasma membrane invaginations in ES9-17 long-term treatment of

Col-0 seedlings, almost exclusively in elongating or differentiated cells (epidermal or root

cap cells), substantiating the possibility that these invaginations constitute read-outs of long

term disturb endocytic trafficking (Figure 5G and 5H, Doumane et al., 2021). Strikingly, we

observed a much higher number of dome-shaped plasma membrane invagination decorated

by the PI(4,5)P$_2$ biosensors mCIT-TUBBYc in cells from ES9-17 treated *sac9-3*[-/-] (Figure

5I), showing that SAC9 depletion causes over-sensitivity to inhibition of endocytic

trafficking. Hypersensitivity to endocytic trafficking inhibition, decreased internalization of

the bulk endocytic tracer FM4-64 and defects in PIN2 protein trafficking together indicate

that endocytic trafficking is impacted in the absence of SAC9.

SAC9 is required for SH3P2 localization at the plasma membrane.

To gain some insights into SAC9 function, we next screened for SAC9 interactors using a

yeast two hybrid assay against a universal Arabidopsis normalized dT library (ULTImate

Y2H SCREEN, hybrigenics). As a bait, we used a portion of SAC9 (AA 499 to 966), which

includes the WW domain, a putative protein-protein interaction platform (Figure 6A). Out of

59.6 million screened interaction, we recovered and sequenced 260 independent clones

corresponding to 107 different proteins. However, among these proteins, most were classified

as proven artifact or low confidence interaction and only three candidates (2.8 %) were

ranked as very high confidence in the interaction (Supplementary File 1B). Among the high

confident interaction candidate, we identified 11 clones corresponding to SH3P2 (5

independent clones). In the yeast-two-hybrid screen, the selected interaction domain

identified for SH3P2, corresponds to the C-terminal part of the proteins (aa 213-368), which

includes the SH3 domain (Figure 6A). We decided to focus on this candidate because, like

SAC9, SH3P2 is linked to both clathrin-mediated endocytosis and membrane

phosphoinositides. Indeed, SH3P2 i) colocalizes with clathrin light chain, ii) cofractionates

with clathrin-coated vesicles (Nagel et al. 2017), and iii) co-immunoprecipitates in planta

with clathrin heavy chain (Nagel et al. 2017), clathrin light chain (Adamowski et al. 2018)

and the dynamin DRP1A (Ahn et al. 2017). Furthermore, SH3P2 binds to various

phosphoinositides in vitro (Zhuang et al. 2013; Ahn et al. 2017), and its plasma membrane

localization is dependent on $PI(4,5)P_2$ in vivo (Doumane et al., 2021).

In an attempt to validate the reverse interaction, SH3P2-GFP was transiently expressed in

N. benthamiana, then immunoprecipitation followed by mass spectrometry analysis (IP-MS)

was done. Among the top interactors were two SAC9 protein isoforms in *N. benthamiana*, as

determined by log fold-change and p-value in peptide count when compared to GFP control

(Fig. 6B-C). As SH3P2 was previously shown to be degraded via the proteasome (Leong

2022) , Bortezomib (BTZ) treatment was used to inhibit SH3P2 degradation *in planta*, and

we observed a corresponding increase in the log fold change and p-value of SAC9 peptide

counts compared to control (GFP/BTZ). Notably, AUXILIN-LIKE1/2 which are already

described protein partners of SH3P2 (Adamowski et al. 2018), were found as among the top

hits in both conditions, either with or without BTZ, thereby validating the approach (Fig. 6B-

C).

Using live cell imaging we observed and quantified that fluorescently tagged SAC9 and

$SAC9^{C459A}$ colocalized with SH3P2 in a subcortical population of endomembrane

compartments (Figure 6B, 6C), probably TGN/EEs (Figure 3E and 3F). Because SH3P2 was

described to play a role in different steps of membrane trafficking, from the endocytosis at

the plasma membrane to autophagy, we next addressed more precisely which pool of SH3P2

was affected in the absence of SAC9. Using confocal imaging, we observed a diminution of

the signal corresponding to SH3P2-sGFP at the plasma membrane compared to the cytoplasm

in *sac9-3,* while the amount of SH3P2-sGFP detected via Western blot was similar between

the two genotypes (Figure 6D to 6F, Figure 6-figure supplement 1). Moreover, in the

absence of functional *SAC9*, SH3P2-TdTOM accumulates in structure closely associated with

the aberrant $PI(4,5)P_2$ pool decorated with mCIT-TUBBYc (Figure 6H). Altogether, the

trafficking defects observed in the *sac9* mutant, the impaired localization of SH3P2 in the

absence of a functional SAC9 and the formation of a SAC9-SH3P2 complex all point toward

a role of SAC9 in endocytic trafficking.

359   The endocytic protein T-PLATE has an altered dynamic in the absence of SAC9

SAC9 is observed at the close vicinity to the plasma membrane and its absence causes a

mislocalization of its protein partner SH3P2, as well as a reduction of the overall endocytosis

process. We then assessed if the clathrin-mediated endocytosis at the plasma membrane was

affected in the absence of SAC9. Using Total Internal Reflection Fluorescence (TIRF)

microscopy, we determined the density and dynamic behavior of the endocytic protein from

the T-PLATE complex, T-PLATE-GFP, in the *sac9* mutant compared with wild-type plants.

Quantitative analysis revealed that the density of T-PLATE-GFP was reduced in sac9-3[-/-]

compared with the WT while the amount of TPLATE-GFP detected via western blot was

similar between the two genotypes (Figure 7A, 7B; Figure 6-figure supplement 1; video 3).

Analysis of the dynamics of T-PLATE-GFP at the plasma membrane of etiolated hypocotyl

revealed a decrease of the dwell time of T-PLATE-GFP at the plasma membrane in the *sac9*

mutant compared to the WT (Figure 7C, 7D). This result is in line with a reduction of the

endocytic flow in the absence of SAC9, and suggests that maintaining a strict plasma

membrane accumulation of $PI(4,5)P_2$ is critical for clathrin-mediated endocytosis.

375   DISCUSSION

Here, we showed that SAC9 putative catalytic cysteine is required to complement the *sac9*
phenotype, suggesting that SAC9 phosphoinositide phosphatase activity is key for its
function. Fluorescent SAC9 protein fusions colocalize with TGN/EE markers in a
subpopulation of endosomes close to the plasma membrane. We found that the subcellular
patterning of $PI(4,5)P_2$ is defective in *sac9* mutants, consistent with the idea that SAC9 is a
$PI(4,5)P_2$ 5-phosphatase. In planta, SAC9 interacts and colocalizes with the endocytic
trafficking component SH3P2. In the absence of SAC9, and therefore when the patterning of
the $PI(4,5)P_2$ at the plasma membrane and at its close vicinity is affected, SH3P2 localization
at the plasma membrane is decreased and the rate of clathrin-mediated endocytosis is
significantly reduced. Together, these findings underlie the importance of strictly restricting
$PI(4,5)P_2$ to the plasma membrane during the endocytic process.

SAC9 and phosphoinositide interconversion along the endocytic pathway

In animal cells, a phosphoinositide conversion cascade has been described through the
successive action of phosphoinositide kinases and phosphatases. This cascade starts with
$PI(4,5)P_2$ at the plasma membrane and ends-up with PI3P in the membrane of early
endosomes (Noack and Jaillais 2017; Schmid and Mettlen 2013; Abad et al. 2017; Posor et
al. 2015; Posor et al. 2013; Shin et al. 2005). Briefly, $PI(4,5)P_2$ is dephosphorylated at the
plasma membrane or during the endocytic process by 5-phosphatase enzymes such as OCRL
and synaptojanin (De Matteis et al. 2017)(Cauvin et al. 2016; Nandez et al. 2014; Ben El
Kadhi et al. 2011; Posor et al. 2013). PI4P is then phosphorylated into $PI(3,4)P_2$ inside
clathrin-coated pits by a PI3-Kinase (PI3K C2α), before being converted into PI3P through
the action of a PI4P phosphatase on clathrin-coated vesicles (Schmid and Mettlen 2013;
Posor et al. 2013). Because both $PI(4,5)P_2$ and PI3P are essential regulators of endocytic
trafficking, this conversion cascade, and in particular the precise recruitment of dedicated
enzymes at the right place and at the right time during endocytic trafficking, is of critical
importance for clathrin-mediated endocytosis to proceed normally (Noack and Jaillais 2017;
Schmid and Mettlen 2013).

The plants TGN collects endocytic vesicles and redirects cargo proteins either to the
plasma membrane for recycling or to late endosomes/vacuole for degradation (Dettmer et al.
2006; Narasimhan et al. 2020; Rodriguez-Furlan et al. 2019). The plant TGN/EE is enriched

in PI4P, not PI3P, which instead accumulates in late endosomes (Noack and Jaillais 2020;
Simon et al. 2014). Interestingly, we found that i) SAC9 localizes to clathrin-coated
compartments close to the plasma membrane, and ii) the *sac9* mutant accumulates PI(4,5)P_2
in vesicular structures at the cortex of the cells. Together, we propose that SAC9 represent
the long sought-after enzyme which performs the PI(4,5)P_2-to-PI4P conversion during the
plant endocytic process (Figure 8). As such, SAC9 is required to erase PI(4,5)P_2 in
endosomal membranes and thereby maintains this lipid strictly at the plasma membrane.

SAC9 being a putative PI(4,5)P_2 phosphatase, we could have expected that SAC9
overexpression may have induced gain-of-function phenotypes caused by ectopic PI(4,5)P_2
dephosphorylation. However, transgenic lines that express SAC9 under the expression of the
UBQ10 promoter did not show any obvious phenotypes, and instead complemented the *sac9*
loss-of-function phenotype like the endogenous promoter. This absence of overexpression
phenotypes may be explained by several hypotheses. First, it is possible that SAC9 is not
sufficiently overexpressed when under the controlled of the *UBQ10* promoter compared to
the endogenous SAC9 promoter. Second, it is possible that SAC9 overexpressing lines
compensate for the excess SAC9 activity by adjusting their PI4P and PI(4,5)P_2 balance.
However, our data point toward a third possibility. Most lipid phosphatase studied to date
require membrane association to dephosphorylate their cognate phosphoinositide, while they
do not act on their target lipid when they are in the cytosol. We found that SAC9 strongly
accumulates in the cytoplasm when it is expressed under the control of either promoter and in
particular when it is overexpressed. Thus, SAC9 membrane association, rather than its
expression level, is likely to be the limiting factor preventing SAC9 gain-of-function
phenotype in over-expression lines.

Accumulation of PI(4,5)P_2-containing vesicles in the *sac9* mutant

In the absence of SAC9, PI(4,5)P_2 accumulates inside the cell in what appears to be abnormal
compartments that are associated with but independent from TGN/EEs. We can speculate that
the prolonged accumulation of PI(4,5)P_2 on clathrin-coated compartments, and perhaps the
lack of PI4P production on these structures, impairs the function of these vesicles. We also
observed that PI4P biosensors accumulates in TGNs in the *sac9* mutants. We previously
found that PI4P biosensors relocalize to PI4P-containing TGNs upon specific depletion of

PI4P at the plasma membrane (Simon et al., 2016, Platre et al., 2018). PI4P being the putative
product of SAC9 (Williams et al., 2009), it is possible that there is a reduced amount of PI4P
at the plasma membrane in this mutant. In that case, the balance of PI4P between the plasma
membrane and the TGN would be affected between these two compartments, which could
explain the relocalization of the PI4P sensors to TGN compartments.

Ultrastructural analyses previously revealed that *sac9* massively accumulates vesicles,
presumably containing cell wall precursors, in close proximity to the irregular cell wall
deposition (Vollmer et al. 2011). The finding of these unknown subcortical compartments in
sac9 resonates with our confocal microscopy observations of ectopic $PI(4,5)P_2$-containing
vesicles in this mutant. Importantly, we found that these vesicles are not extensively labeled
after a short-term FM4-64 treatment. This suggests that these vesicles are accumulating
slowly over time in the *sac9* mutants and are not actively exchanging membrane materials
with the TGN/EEs or the plasma membrane, at least not in the 30-minute time frame of the
FM4-64 pulse labeling experiment. Moreover, recent proteomic characterization of isolated
Arabidopsis clathrin-coated vesicles identified SAC9 as one of the most abundant clathrin-
coated vesicles proteome components (Dahhan et al. 2022). Interestingly, in that study, no
other 5-phosphatase were identified to be associated with clathrin-coated vesicles. Together,
we propose that $PI(4,5)P_2$ accumulation on endocytic vesicles after scission could lead to an
altered dynamics of these vesicles, which may be unable to fuse correctly with TGN/EEs,
ultimately leading to their accumulation in the cell cortex (Figure 8).

We observed that the endocytic pathway is partially impaired in the absence of SAC9, and
that *sac9* was oversensitive to ES9-17-mediated inhibition of endocytosis. In animal cells,
membrane protuberances have been reported in cultured cells knocked-out or knock-down for
enzymes of $PI(4,5)P_2$ metabolism (Gurung et al. 2003; Terebiznik et al. 2002; Mochizuki and
Takenawa 1999). These protuberances could therefore be a conserved process affected by
disturbed $PI(4,5)P_2$ homeostasis in both plant and animal cells leading to such plasma
membrane distortion. It is not the first observation of plasma membrane protuberances in
cells where $PI(4,5)P_2$ homeostasis is affected: upon PIP5K6 over-expression in pollen tubes,
which increases $PI(4,5)P_2$ levels, plasma membrane invaginations also occurs. The authors
elegantly demonstrated that these plasma membrane distortions are due to clathrin-mediated
endocytosis defects (Zhao et al. 2010). These structures were also reported when inducible
depletion of $PI(4,5)P_2$ at the plasma membrane of root cells was performed using the iDePP
system (Doumane et al. 2021), suggesting that the tight regulation of the $PI(4,5)P_2$

metabolism is important for the integrity of the plasma membrane. Previous study on the cellular organization of the primary roots of *sac9-1* seedlings at ultrastructural level reported the excessive membrane material either in direct contacts or close to the wall forming the protuberances (Vollmer et al. 2011). The irregular cell wall deposition in the *sac9* mutant leads to cell wall ingrowths, coined "whorl-like structures" (Vollmer et al. 2011). We propose that such protuberances could be a readout of long-term endocytic defects, for instance caused by an excess of plasma membrane due to unbalanced exocytosis. If so, the appearance of plasma membrane protuberances upon $PI(4,5)P_2$ metabolism perturbation could be an additional indication of disturbed endocytosis.

How could SAC9 regulate endocytosis?

We observed that the endocytic pathway is partially impaired in the absence of SAC9. We envision several scenarios to explain these endocytic defects. They are not mutually exclusive and ultimately, it would not be surprising if the *sac9* endocytic phenotype results from a combination of altered cellular pathways relying, directly or indirectly, on the precise spatio-temporal regulation of anionic lipid homeostasis. First, as mentioned in the paragraph above, the *sac9* mutant likely accumulates ectopic vesicle following endocytosis. These vesicles trap membrane components inside the cell, which may overall impact the endocytic process. In addition, it is possible that the localization or dynamics of $PI(4,5)P_2$ binding proteins, which regulate clathrin-mediated endocytosis, could be impacted, as we observed for SH3P2 and T-PLATE (Zhuang et al. 2013; Yperman et al. 2021b; Yperman et al. 2021a). Other proteins from the T-PLATE complex also interact with anionic lipids, including $PI(4,5)P_2$, and could be affected in *sac9* (Yperman et al. 2021b; Yperman et al. 2021a). Furthermore, the dynamics of dynamin-related proteins, which contain a lipid-binding plekstrin homology domain, and the AP2 adaptor complex, which relies on $PI(4,5)P_2$ for localization in planta (Doumane et al. 2021), could also be altered in *sac9*. It was also recently reported that perturbation of clathrin-containing vesicles at the TGN/EEs also impacts clathrin-mediated endocytosis (and conversely) (Yan et al. 2021). It is thus possible that perturbing phosphoinositides metabolism after the scission of clathrin-coated vesicles from the plasma membrane impacts the dynamics of clathrin-mediated endocytosis at the cell surface in an indirect way, for example because of slowed recycling of the clathrin pool or other endocytic components at the surface of TGN/EEs.

It is also important to note that bulk endocytosis in general and clathrin-mediated
endocytosis in particular are still going-on in the *sac9* mutant, albeit at a reduced rate. Thus,
while PI(4,5)P$_2$ strict exclusion from intracellular membranes is important for the endocytic
process, the plant membrane trafficking system appears to be extremely resilient, as it
manages to operate despite the accumulation of these membrane of mixed identity inside the
cell. Here, we can speculate that SAC9 may not be the only 5-phosphatase enzyme that can
control PI(4,5)P$_2$ homeostasis during endocytosis. In addition, while PI(4,5)P$_2$ is the main
phosphoinositide regulating the recruitment of endocytic regulators in animals, it is likely not
the case in plants (Markovic and Jaillais 2022). Indeed, PI4P, not PI(4,5)P$_2$, is the major
anionic lipids that powers the plasma membrane electrostatic field (Simon et al. 2016), and is
very likely key to recruit endocytic proteins in plants (Yperman et al. 2021b; Yperman et al.
2021a). In addition, phosphatidic acid (PA) is also important to drive the electrostatic field of
the plasma membrane (Platre et al. 2018) and PA could be involved in the localization of
AtEH1/Pan1, a component of the T-PLATE complex (Yperman et al. 2021a, Dragwidge et
al., 2022).

Limitation of the study

SAC9 was previously proposed to act as a PI(4,5)P$_2$ 5-phosphatase in planta, mainly
because the *sac9* mutant accumulates PI(4,5)P$_2$ (Williams et al. 2005). We found an
accumulation of PI(4,5)P$_2$ sensors inside the cells in *sac9*, which is fully consistent for a 5-
phosphatase enzyme. However, we could not confirm this activity in vitro. This is mainly due
to problems in purifying SAC9 catalytic domain in sufficient quantity. Yet, it represents a
weakness of our study that we should keep in mind. Indeed, although this is not the most
parsimonious explanation, it is possible that SAC9 is not a 5-phosphatase and that instead it
controls the localization or activity of another enzyme bearing this catalytic activity.
Nonetheless, the fact that the putative catalytically inactive mCIT-SAC9^{C459A} fusion protein
does not rescue *sac9* macroscopic phenotype and PI(4,5)P$_2$ localization suggests that SAC9
phosphatase activity is required for its function. We showed that the number of BFA bodies
labeled with FM4-64 in *sac9-3* vs wild-type is reduced. Such reduction may be due to i) a
lower amount of BFA bodies formed per cell (i.e., the number per cell area or volume), ii)
reduced FM4-64 internalization from the cell surface, or iii) defects in the balance between
endocytic and exocytic trafficking which alter TGN/EE function, iv) a combination of those.

In either case, such decrease suggests that membrane trafficking flux through the endosomal
system is impacted in *sac9*.

To explain the endocytic defects observed in the *sac9* mutant, we mainly focused on
PI(4,5)P$_2$ mis-patterning. However, biochemical measurements also showed that the *sac9*
mutant accumulates less PI4P than its wild-type counterpart. Given the importance of PI4P
for plant cell function (Marković & Jaillais 2022; Noack et al;, 2022; Noack et al. 2020;
Simon et al. 2016), it is possible that PI4P rather than (or in combination with) PI(4,5)P$_2$
defects are involved in the *sac9* phenotypes. Given that phosphoinositide metabolism is
highly intricate, we recognize that it is difficult to fully untangle the specific involvement of
each lipid in the observed phenotypes. In addition, the SAC9 protein may carry specific
functions outside of its catalytic activity. For example, SAC9 directly interacts with SH3P2,
yet the physiological relevance of this interaction for both SAC9 and SH3P2 function
remains unclear. Structure-function analyses aiming at dissecting this interaction will be
instrumental to clarify this point. Based on confocal imaging, we concluded that the level of
plasma membrane associated SH3P2-sGFP is reduced in the *sac9* mutant. In addition, the
overall levels of SH3P2 are not affected in the mutant. Subcellular fractionations and
quantitative colocalization analyses between SH3P2-GFP and compartment markers in the
sac9 mutant background will be needed to fully understand how SAC9 impacts SH3P2
localization. In addition, the exact function of SH3P2 in endocytosis is also elusive and it
will be important to analyze it in details in the future (Adamowski et al., 2022). Finally, one
of the clear limitations in interpreting the *sac9* phenotype, which is common to most genetic
approaches, is the accumulation of defects over a long-time period in the mutant. As such, it
is impossible to pin-point toward direct or indirect effects of PI(4,5)P$_2$ mis-patterning on the
cellular and developmental phenotypes of this mutant. Future studies, aiming at rapidly
manipulating SAC9 function and localization in an inducible manner will be instrumental in
disentangling the direct and indirect effects of SAC9 on lipid dynamics and endocytosis
regulation.

MATERIALS AND METHODS

Cloning

SAC9 promoter *(SAC9pro*; 800 bp) was amplified from Col-0 genomic DNA using *SAC9prom_fwd_gb* TTGTATAGAAAAGTTGCTATTGAAAAAAGATAGAGGCGCGTG and *SAC9prom_rev_gb* TTTTTTGTACAAACTTGCCTGAGCTCAGGACCAAGCGG primers and the corresponding PCR product was recombined into *pDONR-P1RP4* (life technologies www.lifetechnologies.com/) vector by BP reaction to give *SAC9pro/pDONR-P1RP4*. The *mCIT* and *TdTOM* containing vectors *cYFPnoSTOP/pDONR221* and *TdTOMnoSTOP/pDONR221* were described before (Simon et al., 2014; Jaillais et al., 2011).

The genomic sequence of *SAC9* (At3g59770) was amplified by PCR using 7-day old Arabidopsis seedlings gDNA as template and the *SAC9-B2R* GGGGACAGCTTTCTTGTACAAAGTGGCTATGGATCTGCATCCACCAGGTTAGT and *SAC9-B3wSTOP* GGGGACAACTTTGTATAATAAAGTTGCTCAGACACTTGAAAGGCTAGTCCAT primers. The corresponding PCR product was recombined into *pDONR-P2RP3* vector by BP reaction to give *SAC9g/pDONR-P2RP3*.

SAC9$^{\Delta CC}$ was amplified from *SAC9g/pDONR-P2R-P3* with SAC9g_deltaCC_C-F /5phos/ggaattgatccagctacc SAC9g_deltaCC_N-R /5phos/AAGCTTCTTTGCCTGTATTATG primers and then ligated to give *SAC9g$^{\Delta CC}$/pDONR-P2R-P3*.

SAC9^{C459A} mutation was obtained by site-directed mutagenesis using the partially overlapping SAC9-C459A-F TACGTTTTAACGCTGCTGATTCCTTGGATCGAACAAATGC and SAC9-C459A-R AGGAATCAGCAGCGTTAAAACGTATCACCCCATTTTGATGTG primers on *SAC9g/pDONR-P2RP3*.

The gateway expression vector SH3P2shortprom (0.4 kb):SH3P2gDNA-tdTOM was obtained from pDONR P4-P1R SH3P2 short prom amplified with the primers: attb4-sh3p2promshortF GGGGACAACTTTGTATAGAAAAGTTGCTGGTTAAGGGTCTTCTAGATGGTGTAG, attb1-sh3p2promR

GGGGACTGCTTTTTTGTACAAACTTGCTCTTCACCAGATCAAGAGCTATTCACAA

A, pDONR221/SH3P2g amplified with the primers: attB1_SH3P2g_F

GGGGACAAGTTTGTACAAAAAAGCAGGCTTAACC_ATGGATGCAATTAGAAAC

AAGCTAG, attB2_SH3P2g_R

GGGGACCACTTTGTACAAGAAAGCTGGGTA_GAAAACTTCGGACACTTTGCTAGC

AAGAAC and pDONR-P2R-P3 TdTOM together with pDEST pH7m34GW by LR GW

Final destination vectors were obtained using three fragments LR recombination system (life

technologies, www.lifetechnologies.com/) using pB7m34GW destination vector (Karimi et

al. 2007). The following reactions were set-up to generate the corresponding destination

vectors: UBQ10prom:tdTOM-SAC9g/pH, SAC9prom:mCIT-SAC9g/pB, pAtSAC9:mCIT-

SAC9g^{DEAD}/pB, SAC9prom:TdTOM-SAC9g^{DEAD}/pH , pAtSAC9:mCIT-SAC9g$^{\Delta CC}$/pB,

SH3P2shortprom:SH3P2gDNA-tdTOM, pUb10:SH3P2gDNA-tdTOM (Supplementary File

1).

Growth condition and plant materials

Arabidopsis thaliana Columbia-0 (Col-0) accession was used as wild-type (WT) reference

genomic background throughout this study. Arabidopsis seedlings *in vitro* on half Murashige

and Skoog (½ MS) basal medium supplemented with 0.8% plant agar (pH 5.7) in continuous

light conditions at 21 °C. Plants were grown in soil under long-day conditions at 21 °C and

70% humidity 16 h daylight. Wild-type Col-0 and heterozygous (or homozygous) sac9-3

were transformed using the dipping method (Clough and Bent 1998).

For each construct generated in this paper (UBQ10prom:tdTOM-SAC9g/pH,

SAC9prom:mCIT-SAC9g/pB, pAtSAC9:mCIT-SAC9g^{DEAD}/pB, SAC9prom:TdTOM-

SAC9g^{DEAD}/pH, pAtSAC9:mCIT-SAC9g$^{\Delta CC}$/pB, SH3P2shortprom:SH3P2gDNA-tdTOM,

pUb10:SH3P2gDNA-tdTOM), between 20 and 24 independent T1 were selected on

antibiotics (Basta or hygromycin) and propagated. In T2, all lines were screened using

confocal microscopy for fluorescence signal and localization. Between 3 to 5 independent

lines with a mono-insertion and showing a consistent, representative expression level and

localization were selected and grown to the next generation. Each selected line was

reanalyzed in T3 by confocal microscopy to confirm the results obtained in T2 and to select

homozygous plants. At this stage, we selected one representative line for in depth analysis of

the localization and crosses and two representative lines for in depth analysis of mutant
complementation.

FM4-64 staining and drug treatments

Seedlings were incubated in wells containing 1 µM FM4-64 (Life Technologies T-3166;
from a stock solution of 1.645 mM = 1 mg/ml in DMSO) in half Murashige and Skoog (½
MS) liquid basal medium without shaking for 30 min and in dark. Seedlings were then
mounted in the same medium and imaged within a 10 min time frame window (1 h ± 5 min).

Seedlings were incubated in wells containing Brefeldin A (BFA; Sigma B7651) applied at 50
µM (from a stock solution of 30 mM in DMSO), or a corresponding volume of DMSO as
'mock' treatment, dissolved in liquid ½ MS for one hour in dark without shaking before
mounting in the same medium and imaging. For co-treatment with 50 µM BFA and 1 µM
FM4-64, FM4-64 and BFA were added at the same time. Imaging was performed within a 14
min time frame window (1 h ± 7 min).

For PAO treatment, seedlings were incubated in wells containing 30 µM PAO (Sigma P3075,
www.sigmaaldrich.com, PAO stock solution at 60 mM in DMSO), or a volume of DMSO as
mock treatment, during the indicated time. Roots were imaged within a 10 min time frame
window around the indicated time. The PAO effects on the localization of the biosensors
were analyzed by counting manually the number of labelled compartments per cell.

ES9-17 was stored at -20°C as a 30 mM stock solution in DMSO. Seedlings were incubated
in dark without shaking for the indicated duration (± 7 min) in liquid ¼ MS (pH 5.7) in wells
containing 1 % DMSO to help solubilization of ES9-17, and either 1 µM ES9-17 or the
corresponding additional volume of DMSO (mock treatment). For 1 µM ES9-17 and 1 µM
FM4-64, FM4-64 was added 30 min after the ES9-17 (the indicated time corresponds to ES9-
17 exposure).

Western-Blot

To verify that both T-PLATE and SH3P2 fusion proteins were both identically expressed in
wild type and sac9, 10-days-old seedlings were grind in liquid nitrogen, weighed and
resuspended in 1 ml of extraction buffer (100 mM Tris–HCl pH 7.5, 150 mM NaCl,

5 mM EDTA, 5% glycerol, 10 mM DTT, 0.5% Triton X 100, 1% Igepal and 1% protease

inhibitors (Sigma) into milliQ water) and centrifuged at 5000rpm for 10 min at 4°C to obtain

the total protein extract in the supernatant. The total protein extract was then filter on column

and 100µl of the filtrate denatured with 50µl of Lamelli buffer 3x at 95°C for 5 min. 40 µL of

the total protein extract was loaded onto 7.5 % polyacrylamide gel, run for 120 min at 120 V

and blotted on nitrocellulose membranes 2hours, 100 V on ice. Membrane were then

incubated 5min in red-ponceau buffer, washed with milliQ water and imaged. Then

membranes were blocked in 5 % milk dissolved in TBST buffer (10 mM Tris-HCL, 150 mM

NaCL, 0.05 % Tween20) for 1hour. mCIT tagged proteins were revealed by using

respectively GFP monoclonal antibody (anti-GFP mouse monoclonal, Roche; at 1/1000 in 5

% milk over-night) as primary antibodies and anti-mousse IgG-HRP conjugated secondaries

antibodies (Mouse IgG, HRP conjugate W402B, Promega; 1/5000 in TBST, 4 hours). As a

loading control we used anti-tubulinα antibodies as primary antibodies (1/1000 in 5 % milk

over-night). Finally tagged proteins were detected by chemiluminescence using ECL,

substrate to HRP enzyme, revelation.

Transient Expression in *N. benthamiana*

669   The AtSH3P2-GFP constructs was described previously (Leong et al., 2022). The binary

670   plasmid was transformed into Agrobacterium tumefaciens strain C58C1 and transient

671   expression of the construct was performed by infiltration of N. benthamiana leaves at the

672   four-to six-leaf stage.

Immunoprecipitation

To verify that SAC9 fusion proteins were present in the plant cells, leaves from 1-month-

old plants were grind in liquid nitrogen, weighed and resuspended in 1 ml of extraction buffer

(100 mM Tris–HCl pH 7.5, 150 mM NaCl, 5 mM EDTA, 5% glycerol, 10 mM DTT,

0.5% Triton X 100, 1% Igepal, 100 µM MG132 and 1% protease inhibitors (Sigma) into

milliQ water) and centrifuged at 5000rpm for 10 min at 4°C to obtain the total protein extract

in the supernatant. The total protein extract was then filter on column, incubated 40 min with

50 µL of magnetic protein G Dynabeads (Millipore) fused to a GFP monoclonal antibody

(1/500 in PBS 0.002 % tween). Beads, antibodies and antibodies bund proteins were

magnetically precipitated on columns, eluted and denatured in 40 µL of Laemmli buffer 2X

at 95°C for 5 min. 20 µL of the immunoprecipitation was loaded, blotted on nitrocellulose

membranes, incubated in red-ponceau buffer and blocked in 5 % milk as previously describe.

Tagged SAC9 fusion proteins were revealed by using GFP monoclonal antibody (anti-GFP
mouse monoclonal, Roche) and detected by chemiluminescence using ECL revelation as for
T-PLATE and SH3P2 fusion proteins.

Pull-down assays in N. benthamiana using GFP-Trap assays were performed as previously
described (Leong et al., 2022). For immunoblot analysis, protein extracts were separated by
SDS-PAGE, transferred to PVDF membranes (Biorad), blocked with5% skimmed milk in
TBS, and incubated with primary antibodies anti-GFP-HRP (SantaCruz) antibody using
1:2000 dilutions in TBS containing 0.1% Tween 20. The immunoreaction was developed
using an ECL Prime Kit (GE Healthcare) and detected with Amersham Imager 680 blot and
gel imager.

Mass spectrometry data analysis
Proteins in the *N. benthamiana* proteome corresponding to the peptide hits were identified
according to the annotated proteome (Kourelis et al. 2019). The proteins were submitted to
BLAST (Sayers et al. 2021) to identify their isoform(s) in the *A. thaliana* proteome(Cheng et
al. 2017). The dataset containing the *A. thaliana* protein ID and peptide counts were
submitted to the R package IPinquiry4 (https://github.com/hzuber67/IPinquiry4) which uses a
Genewise Negative Binomial Generalized Linear Model developed by EdgeR to calculate the
log fold-change (log(FC)) and log10(p value).

NanoLC-MS/MS analysis and data processing
Proteins were purified on an NuPAGE 12% gel (Invitrogen) and Coomassie-stained gel
pieces were digested in gel with trypsin as described previously (Borchert *et al*, 2010) with a
small modification: chloroacetamide was used instead of iodoacetamide for
carbamidomethylation of cysteine residues to prevent formation of lysine modifications
isobaric to two glycine residues left on ubiquitinylated lysine after tryptic digestion. After
desalting using C18 Stage tips peptide mixtures were run on an Easy-nLC 1200 system
coupled to a Q Exactive HF-X mass spectrometer (both Thermo Fisher Scientific) as
described elsewhere (Kliza *et al*, 2017) with slight modifications: the peptide mixtures were
separated using a 87 minute segmented gradient from 10-33-50-90% of HPLC solvent B
(80% acetonitrile in 0.1% formic acid) in HPLC solvent A (0.1% formic acid) at a flow rate
of 200 nl/min. The seven most intense precursor ions were sequentially fragmented in each
scan cycle using higher energy collisional dissociation (HCD) fragmentation. In all

measurements, sequenced precursor masses were excluded from further selection for 30 s.

The target values were 105 charges for MS/MS fragmentation and 3x106 charges for the MS

scan.

Acquired MS spectra were processed with MaxQuant software package version 1.5.2.8 with

integrated Andromeda search engine. Database search was performed against a *Nicotiana*

benthamiana database containing 74,802 protein entries. Endoprotease trypsin was defined as

protease with a maximum of two missed cleavages. Oxidation of methionine,

phosphorylation of serine, threonine and tyrosine, GlyGly dipeptide on lysine residues, and N-

terminal acetylation were specified as variable modifications. Carbamidomethylation on

cysteine was set as fixed modification. Initial maximum allowed mass tolerance was set to

4.5 parts per million (ppm) for precursor ions and 20 ppm for fragment ions. Peptide, protein

and modification site identifications were reported at a false discovery rate (FDR) of 0.01,

estimated by the target-decoy approach (Elias and Gygi). The iBAQ (Intensity Based

Absolute Quantification) and LFQ (Label-Free Quantification) algorithms were enabled, as

was the "match between runs" option (Schwanhäusser et al. 2011).

Data availability

The mass spectrometry data from this publication will be made available on PRIDE archive

(https://www.ebi.ac.uk/pride/archive/) and with the identifier [PXD033585].

Live cell imaging

Most images (see exceptions below) were acquired with the following spinning disk confocal

microscope set up: inverted Zeiss microscope (AxioObserver Z1, Carl Zeiss Group,

http://www.zeiss.com/) equipped with a spinning disk module (CSU-W1-T3, Yokogawa,

www.yokogawa.com) and a ProEM+ 1024B camera (Princeton Instrument,

http://www.princetoninstruments.com/) or Camera Prime 95B (www.photometrics.com)

using a 63x Plan-Apochromat objective (numerical aperture 1.4, oil immersion). GFP and

mCITRINE were excited with a 488 nm laser (150 mW) and fluorescence emission was

filtered by a 525/50 nm BrightLine! single-band bandpass filter (Semrock,

http://www.semrock.com/), mCHERRY en TdTOM was excited with a 561 nm laser (80

mW) and fluorescence emission was filtered by a 609/54 nm BrightLine! single-band

bandpass filter (Semrock, http://www.semrock.com/). For quantitative imaging, pictures of

epidermal root meristem cells were taken with detector settings optimized for low
background and no pixel saturation. Care was taken to use similar confocal settings when
comparing fluorescence intensity or for quantification.

Colocalization experiments were performed on an inverted Zeiss CLSM710 confocal
microscope or an inverted Zeiss CLSM800 confocal microscope and inverted Zeiss
CLSM980 confocal microscope using a 63x Plan-apochromatic objective. Dual-color images
were acquired by sequential line switching, allowing the separation of channels by both
excitation and emission. In the case of colocalization, we also controlled for a complete
absence of channel crosstalk. GFP was excited with a 488 nm laser, mCIT was excited with a
515nm laser and mCH/tdTOM were excited with a 561 nm laser. Imaging was performed in
the root epidermis in cells that are at the onset of elongation. Only cells imaged at their Zi
where considered for the colocalization analysis.

Yeast two-hybrid screen

The yeast two hybrid screen was performed by hybrigenics services
(https://www.hybrigenics-services.com/contents/our-services/interaction-discovery/ultimate-
y2h-2), using the ULTImate Y2H screen against their Universal Arabidopsis Normalized
library obtained using oligo_dT. A codon optimized residue (aa 499-966) of SAC9
gttaataatcagggggggatataacgctccccttccaccgggatgggaaaaaagagctgatgccgtaactggaaaatcatattatataga
tcacaatacaaagacaacaacatggagtcatccatgtcctgataaaccatggaagagacttgacatgaggtttgaggaatttaagagat
caactatcttatctcctgtgtcagaacttgccgatcttttctgcaacaaggtgatatccatgcaaccctctatactggctcgaaagctatgc
acagccaaattctcaacatcttcagtgaagaatcaggagcatttaaacagttttctgcagcacagaaaaacatgaagattacactacaga
gaagatataaaaatgctatggttgatagttcacggcaaaaacagctcgagatgtttctgggaatgaggcttttcaagcatcttccatcaatt
cctgtccagcctttacatgtactttctcgaccatctggtttctttctgaaacctgtacctaacatgtccgaaagttccaatgatgggtccagtc
tgctgagtatcaagaggaaggacataacttggctatgtccacaagctgcagatattgttgaattatttatctatctcagtgagccttgccat
gtatgtcaacttctactgaccatatcacacggtgcggatgatttgacatgtccatccactgtggacgtgagaactggacgccacatagag
gaccttaaattagttgttgagttagttcaactggattaccgattacctgtaattatgttttctggacagggtgcttcaataccacgctgtgcaa
atggtacaaatcttctggtacccttaccagggccaattagttctgaggatatggctgttactggagctggtgcacgtcttcatgaaaaaga
tacgtcaagtctttcactgctatatgattttgaagaactagaaggacagttggatttcttaacccgtgtagttgctgttacattttatccagctg
gtgctgttagaattcctatgactcttggtcagatagaagtccttggaatttctcttccatggaaaggaatgtttacttgtgaacgtactggag
gaagattagctgaacttgcaaggaaaccagatgaagatggaagtccttttttcatcttgttctgacttgaatccgtttgctgcaacaacatctt
tacaggctgaaactgtttccacaccagtacaacagaaggatcccttttcccagtaatctgcttgacctttttgacaggagaggactcttcttct

gaccccttcccacaaccagtggtggaatgtattgcaagtggaggcaatgacatgcttgatttcttagacgaagcagttgttgaatatcgc

ggctctgacactgttcctgacgggtct was cloned in a pB66 vector (N-GAL4-bait-C fusion). The

screen was performed on 0.5 mM 3AT. 59 million interactions were analyzed, and 260

positive clones were sequenced (ATNOR_dT_hgx4515v1 _pB66, Supplementary File 1B.

TIRF imaging

3 days old etiolated seedlings were used for hypocotyl epidermal cells observations. TIRF-

VAEM imaging were made using an ilas2 TIRF microscope (Roper Scientific) with 100x

Apo NA 1.46 Oil objective and a Prime 95B camera (Photometrics,

https://www.photometrics.com/) and 1.5 coverslips were used (VWR 61-0136). Time lapses

were acquired during 300 time-points for 300s (acquisition time 500ms). Spot density were

measured using Spot_detector ImageJ macro (Bayle et al. 2017). Since the data below 5

seconds and beyond 70 seconds exceeds the typical lifetime of the clathrin coated vesicle at

the plasma membrane, we removed them from the analysis.

Because manual verification of TPLATE-GFP lifetimes is greatly limited by the number of

CCVs that can be detected, particle identification and tracking were performed using ImageJ

plugin Trackmate (https://research.pasteur.fr/fr/software/trackmate/). Trajectories were

reconstructed following a three-stage workflow: (i) detection of peaks potentially associated

with fluorescent emitters, (ii) quality test and estimation of the subpixel position and (iii)

track reconnection. To discriminate between signal and background, particular attention has

been paid to the size and shape of the observable objects. Particle of minimum size 0.5 with a

threshold of 50 and a contrast >0.04 were filtered, to capture as many spots as possible

without background. For many reasons, such as variation in fluorescent intensity, loss of

focus or photobleaching, the emitter can be missing for several time points causing the

premature stop of tracks. Therefore, the maximum number of frames separating two

detections was set to three frames (Bayle et al. 2021). As a final verification, a visual

inspection of the tracks can be performed on a reconstituted image, where all the tracks from

a movie are represented.

2000 tracks were selected per acquisition starting from frame 5 to avoid segmenting truncated

tracks. Acquisition were made on 7 hypocotyl cells from three different plants per genotype

and per replicate.

Dissociation indexes

Dissociation indexes of membrane lipid fluorescent biosensors were measured and calculated as previously described (Platre et al., 2018). Briefly, we calculated "indexNoDex", defined as the ratio between the fluorescence intensity (Mean Grey Value function of Fiji software) measured in two elliptical regions of interest (ROIs) from the plasma membrane region (one at the apical/basal plasma membrane region and one in the lateral PM region) and two elliptical ROIs in the cytosol in the mock condition. For quantification, we used FAPP-E50A as a positive control, since the delocalization of a biosensor from the plasma membrane to the cytoplasm helps us to use automatic tools.

Measures, counting and statistical analysis

Primary root length and number of lateral roots were manually measured from pictures. For comparing the primary root length and lateral root density between each genotype, we used a one-way ANOVA and post hoc Tukey HSD pairwise tests (95% confidence level).

For quantitative imaging, pictures of epidermal root meristem cells were taken with detector settings optimized for low background and no pixel saturation. Care was taken to use similar confocal settings when comparing fluorescence intensity. Pseudo-colored images were obtained using the "Green Fire Blue" look-up-table (LUT) of Fiji software (http://www.fiji.sc/). The intracellular compartments ("spots") per cell were automatically counted.

We automatically measured the density of intracellular compartments labeled per root using the procedure described in (Bayle et al. 2017) for each biosensor, and we used two-sided non-parametric Wilcoxon rank-sum tests to compare Col-0 and $sac9$-$3^{-/-}$ genotypes in Figure 4. To account for multiple testing, we used a Bonferroni correction and lowered the significance level of each test at alpha = $0.05/11 = 0.004$.

To assess the effect of the inhibitor PAO on anomalous mCIT-TUBBYc intracellular compartments, we manually counted their number per cell. We tested PAO effect and the effect of treatment duration (30 min and 2 h) on the number of marked intracellular compartments using a generalized linear mixed-effect model (Poisson family) to account for

image ID as a random factor. Two sided post-hoc tests were performed using the R package

"lsmeans" (Russell et al., 2016). We compared the number of FM4-64 positive compartments

in Col-0 and *sac9-3*[-/-] using a generalized linear mixed model (Poisson family) to account for

image ID (*id est* root) as a random factor.

The density of FM4-64 labeled compartments was also compared in Col-0 and *sac9-3*[-/-] using

a linear mixed model accounting for image ID as a random factor, followed by a Wald χ^2 test

(function *Anova* in R package "car").

To compare the effects of BFA on FM4-64 we tried to automatically count the number and

size of the BFA bodies in Col-0 and *sac9-3*[-/-] seedlings, but the analysis was not optimal to

treat the images acquired for sac9-3. We therefore manually counted the number of BFA

body per cell in multiple samples, using the same region of the root. We then compared the

results of the BFA treated Col-0 and *sac9-3*[-/-] seedlings using a generalized linear mixed

model (Poisson family) with image ID (*id est* root) as a random factor (Wald χ^2 test : $\chi^2 =$

33.8, p < 0.001). To compare the effects of BFA on Col-0 and *sac9-3*[-/-] seedings expressing

PIN2-GFP, we counted manually and compared the treatments BFA-Col-0 with BFA-Sac9-3

using a generalized linear mixed model (Poisson family) with image ID (*id est* root) as a

random factor. (Wald χ^2 test : $\chi^2 = 42.1$, p < 0.001). For the dissociation index analysis, we

performed all our statistical analyses in R (v. 3.6.1, (R Core Team, 2019), using R studio

interface and the packages ggplot2 (Wickham 2016), lme4 (Bates et al. 2014), car (Fox and

Weisberg 2011), multcomp (Hothorn et al. 2008) and lsmeans (Lenth and Lenth 2018). To

compare TPLATE-GFP density between Col-0 and sac9-3[-/-] we used a two-sided non-

parametric Kruskal Wallis rank-sum tests for each replicate and obtained each time a

statistical difference (p.value < 0.05 between the two genotypes). Graphs were obtained with

R and R-studio software using the package "ggplot2", and customized with Inkscape

(https://inkscape.org).

871   Quantification of mCIT-SAC9 compartment densities and colocalization with

872   compartments markers

Because the signal of mCIT-SAC9 is mainly diffused in the cytosol, no automatic spot

detection could be used for quantification of densities and colocalization analyses in Figure 2,

3, and 6. Therefore, for comparing the number of intracellular compartments containing

mCIT-SAC9 or mCIT-SAC9^{C459A} protein-fusions per cell across conditions, we manually

counted them and used either a generalized linear mixed-effect model (Poisson family) for

counting comparisons or a linear mixed effect model (and associated ANOVAs) for density

comparisons, accounting for image ID (id est root) as a random factor.

Since the localization of the marker for the membrane compartment was larger in z compared

to the restricted localization of SAC9 (only present close to the surface of the cell), we

counted the number of mCIT-SAC9 labeled structures which were also labelled by the

compartment markers in the cell cortex (Zi plane). The percentage of endosomes labelled by

mCIT-SAC9 or mCIT-SAC9C459A colocalizing with a given marker, counted manually are

presented in the graphs. Positive colocalization was called when the compartment marker was

present as a dotted structured overlaying the mCit-SAC9 signal.

The same approach was used to deduce the localization of the mutated version of SAC9.

After running our mixed models, we subsequently computed two-sided Tukey post hoc tests

(function glht in R package "multcomp", Horthorn et al., 2008) to specifically compare each

pair of conditions."

ACKNOWLEDGEMENTS

We are grateful to the SiCE group (RDP, Lyon, France), Dr Yohann Boutté (LBM,

Bordeaux, France), Dr Fabrice Besnard and Dr Nicolas Doll (RDP, Lyon, France), Hugo

Ducuing (INMG, Lyon, France), Sébastien This (CIRI, Lyon, France) and Augustin Le

Bouquin (IGFL, Lyon, France), Dr Sophie Piquerez (IPS2, Paris-Saclay, France) for

comments and discussions. We thank Patrice Bolland, and Alexis Lacroix from our plant

facility, and Claire Lionnet (RDP, Lyon, France). We are also grateful to Dr Daniël van

Damme (VIB, Ghent, Belgium) for sharing with us T-PLATE-GFP and for discussions. We

would like to thanks Pr Dr Erika Isono (University of Konstanz, Konstanz, Germany) for

sharing SH3P2-sGFP transgenic line with us. We thank Hybrigenics for the yeast two hybrid

screen. We are also grateful to E. Russinova (VIB, Ghent, Belgium) for kindly providing us

ES9-17; S. Bednarek for sharing markers and for discussions. We acknowledge the

contribution of SFR Biosciences (UAR3444/CNRS, US8/Inserm, ENS de Lyon, UCBL)
facilities, notably the LBI-PLATIM-MICROSCOPY for assistance with imaging. This work
was supported by two Seed Fund ENS LYON-2016 and LYON-2019 (to MCC), a Junior
Investigator grant ANR-16-CE13-0021 (to MCC), ERC no. 3363360-APPL under FP/2007–
2013 (to YJ). This work was supported by an Emmy Noether Fellowship GZ: UE188/2-1
from the Deutsche Forschungsgemeinschaft (DFG; to SÜ). MD and AL were funded by
Ph.D. fellowships from the French Ministry of Research and Higher Education.

LEGENDS

Figure 1. Structure-function analysis of SAC9. A, Schematic representation of SAC9 protein. The SAC catalytic domain,
as well as the WW domain and the Coil-coiled domain, are represented. **B,** Representative images of the macroscopic
phenotype observed in (i) wild-type (Col-0), (ii) *sac9-1⁻/⁻* and *sac9-3⁻/⁻* loss of function mutants, (iii) *sac9-3⁻/⁻* complemented
lines expressing full-length genomic DNA encoding SAC9 fused to yellow (mCIT-SAC9,line #1000-9-1) or red (TdTOM-
SAC9, line #987-5-4) fluorescent proteins and a mutated version of the putative catalytic cysteine residue within the C-x(5)-
R-[TS] catalytic motif in the SAC domain (mCIT-SAC9^C459A, line #1354-12-14; right panel). Pictures are taken 12 days post
germination (dpg). Note that a second independent transgenic line is presented for each construct in Figure 1- figure
supplement 1. **C,** Quantification of primary root length in *sac9-3⁻/⁻* homozygous mutants expressing mCIT-SAC9 and mCIT-
SAC9^C459A under the control of the native promoter (*SAC9prom*) and TdTOM-SAC9 under the expression of the Ubq10
promoter. Wild-type (Col-0) seedlings and two independent mutant alleles of *SAC9, sac9-1⁻/⁻* and *sac9-3⁻/⁻*, are used as
controls. **D,** Same as (C) but for the quantification of the lateral root density (ratio of the number of lateral roots to primary
root length). In the plots, middle horizontal bars represent the median, while the bottom and top of each box represent the
25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the interquartile range, excluding data
beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and minimum values. Details for
statistical analysis can be found in the Methods section and Supplementary File 1C. N= number of replicates; n= number of
roots.

Figure 2. SAC9 localizes in the cytosol and in intracellular compartment enriched at the cell cortex. A, Schematic
representation of two root epidermal cells (cell #1, indicated as C1 and cell #2 indicated as C2) imaged at two different focal
planes (cortical plane, close to the plasma membrane and designated as Zi and a median plane, in the middle of the cell
designated as Zii). Note that the C1/C2 and Zi/Zii notations are used consistently according throughout the figures. **B,**
Confocal images of the subcortical part (i.e., Zi) of the Arabidopsis root epidermis expressing mCIT-SAC9, mCIT-SAC9^ΔCC

mutated in the predicted coil-coiled domain, and mCIT-SAC9^{C459A} mutated in its putative catalytic cysteine, under the
control of SAC9 native promoter (*SAC9prom*). **C,** Representative images of the fluorescent signal observe in *sac9-3$^{-/-}$* mutant
expressing mCIT-SAC9, TdTOM-SAC9 and mCIT-SAC9^{C459A}. **D,** Comparison of the number of labeled intracellular
compartments per cell in *sac9-3$^{-/-}$* root epidermis expressing mCIT-SAC9 and TdTOM-SAC9, mCIT-SAC9^{C459A}. **E,**
Colocalization analysis on *Ub10pro:TdTOM-SAC9* (magenta) and *SAC9pro:CIT-SAC9^{C459A}* (green). **F,** Confocal images of
the subcortical part of the Arabidopsis root epidermis (upper panels, Zi) or the center of the cell (lower panels, Zii)
expressing *mCIT-SAC9* together with the PI(4,5)P$_2$ biosensor *2mCH-2xPH^{PLC}* (note that the cells shown in Zi and Zii are the
same, just on different focal plane). **G,** Quantification of the number of endosomes labelled by mCIT-SAC9 observed per
cell at two different focal planes (Zi and Zii). **H,** Quantification of the number of endosomes labelled by FM4-64 observed
per cell at two different focal planes (Zi and Zii). **I,** Co-visualization of *SAC9pro:mCIT-SAC9^{C459A}* (green) together with the
plasma membrane marker Lti6b-2xmCH (magenta) in epidermal root cells taken in the Zi focal plane. The inset shows a
magnification, with the yellow arrows indicating some of the intracellular compartments decorated by mCIT-SAC9^{C459A}
observed at the close vicinity to the labelled plasma membrane in cell C1. Scale bar: 10 µm in A-F; 5 µm in G (left panel)
and 2 µm in the close up of G (right panel). In the plot, middle horizontal bars represent the median, while the bottom and
top of each box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the
interquartile range, excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and
minimum values. Details of the statistical analysis could be found in Supplementary File 1D. The plane (Zi or Zii) in each
image is mentioned, and the image display is representative for the plane used for the analysis. N= number of replicates; n=
number of roots.

Figure 3. SAC9 localizes to a subpopulation of TGN/EE. A, B Confocal images of *SAC9pro:mCIT-SAC9* (A, green) or *SAC9pro:mCIT- SAC9^{C459A}* (B, green) in vivo in Arabidopsis root epidermis together with the endocytic tracer FM4-64 (magenta). **C, D** Confocal images of Arabidopsis root epidermis co-expressing *SAC9pro:mCIT-SAC9* (C, green) or *SAC9pro:mCIT- SAC9^{C459A}* (D, green) together Clathrin Light Chain 2 (CLC2) fused to RFP (*CLC2-RFP*; magenta). Upper panel: fluorescent signals observed in the mock treatment; Lower panel: fluorescent signals observed after 50 µM 60 min BFA treatment. **E,** Confocal images of Arabidopsis root epidermis co-expressing *SAC9pro:mCIT-SAC9* (green) and TGN markers *CLC2-RFP, mCH-VTI12, mCH-RabA1g*, *mCH-RabD1*, the late endosome/pre--vacuolar compartment (LE/MVB) marker *mCH-RabF2a/Rha1* and the Golgi marker *mCH-Got1p* (magenta). **F,** Percentage of colocalization between mCIT-SAC9 and a given endosomal compartment marker per cell, at z_i. N= number of roots, n = number of cells. In the plots, middle horizontal bars represent the median, while the bottom and top of each box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the interquartile range, excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and minimum values. The plane (Z_i or Z_{ii}) in each image is mentioned, and the image display is representative for the plane used for the analysis. Scale bars: 10 µm.

Figure 4. SAC9 restricts PI(4,5)P₂ at the plasma membrane. A, Confocal images of Arabidopsis root epidermis expressing mCIT-tagged sensors in WT (Col-0) and *sac9-3⁻ᐟ⁻* genetic backgrounds. Fluorescence intensity is color-coded (green fire blue scale). **B,** Quantification of density of labelled intracellular compartments (pixel⁻²) in whole roots epidermis expressing mCIT-tagged lipid sensors in wild-type (Col-0) and *sac9-3⁻ᐟ⁻*. Non-parametric Wilcoxon rank sum tests with Bonferroni correction. In the plots, middle horizontal bars represent the median, while the bottom and top of each box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the interquartile range, excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and minimum values. Details for statistical analysis can be found in the Methods section and supplementary Table 6. N= number of replicates; n= number of roots. **C,** Confocal images of mCIT-TUBBYc in root epidermal cell of *sac9-3* heterozygous, *sac9-3⁻ᐟ⁻* (left panel) or *sac9-3⁻ᐟ⁻* expressing *TdTOM-SAC9^C459A* (right panel); N=3 replicates, n=47 roots. **D,** Confocal images of *sac9-3⁻ᐟ⁻* Arabidopsis root epidermal cell (C1 and C2) expressing *mCIT-TUBBYc* (green) and *2xmCH-PH^FAPP1* (magenta) at their cortex (upper panel, Zi) and at their center (bottom panel, Zii); N =3 biological replicates, n=49 roots. Scale bar: 10 µm. The plane (Zi or Zii) in each image is mentioned, and the image display is representative for the plane used for the analysis. Note that in panel D, the same cells are shown in the upper and lower panel, but at different focal planes.

Figure5

Figure 5. The endocytic flux is perturbed in the *sac9* mutant. A, Representative images of Col-0 and *sac9-3*[-/-] seedlings treated for 30 min with FM4-64, which is endocytosed and labels endocytic intracellular compartments. Scale bar: 10 µm. **B,** Quantification from the experiment showed in (A). Violin and box plots quantifying the number of FM4-64 labeled intracellular compartments in Col-0 and *sac9-3*[-/-]. **C,** Representative images of root epidermis following BFA and FM4-64 treatment of Col-0 and *sac9-3*[-/-] seedlings. Examples of FM4-64 labeled BFA bodies are pointed out (green arrowheads). **D,** Quantification from the experiment showed in (C). For Col-0 and *sac9-3*[-/-], the proportion (%) of cells containing from none to six BFA bodies is displayed. Dotted line: means. **E,** Representative images following BFA treatment of PIN2-GFP/WT and PIN2-GFP/*sac9-3*[-/-] seedlings. Examples of PIN2-GFP labeled BFA bodies are pointed out (green arrowheads). Scale bars: 10 µm. **F,** Quantification from the experiment showed in (E). For *PIN2-GFP*/WT and PIN2-GFP/*sac9-3*[-/-], the proportion (%) of cells containing from none to six BFA bodies is displayed. Dotted line: means. **G,** Over-sensitivity of *sac9-3*[-/-] to prolonged inhibition of endocytosis. Seedlings were treated 180 min with 30 µM ES9-17 or DMSO (mock), FM4-64 being added after 30 min (150 min of exposure). The picture showed after the ES9-17 treatment are the results of the projection of a z-stack. **H,** ES9-17 treatments led to dome-shaped plasma membrane invagination. blue arrowheads: invaginations with an obvious connection to the plasma membrane; green arrowheads: invaginations without a clear connection to the plasma membrane (often connected to medullar plasma membrane). N= 2 biological replicates, n = 8 roots. **I,** in *sac9-3* mutant, ES9-17 treatments led to dome-shaped plasma membrane invagination labelled by the PI(4,5)P$_2$ biosensor mCIT-TUBBYc; N= 2 biological replicates, n = 10 roots. Blue arrowheads: invaginations with an obvious connection to the plasma membrane; green arrowheads: invaginations without a clear connection to the plasma membrane (often connected to medullar plasma membrane), Scale bars: 10 µm. Details of the statistical analysis could be found in Supplementary File 1G. N= 3 biological replicates, n = number of cells.

Figure6

Figure 6. SAC9 interact and colocalize with SH3P2 and regulates it subcellular distribution. **A,** Schematic representation of the yeast two-hybrid screen using SAC9 as a bait, where SH3P2 was found as a protein partner. The selected interaction domain (interacting with bait 1) corresponds to the amino acid sequence shared by all the eleven prey fragments matching the same reference protein. Note that one other fragment of SAC9 was intended to be screened for interacting proteins (bait 2), but the construct was autoactivate in yeast. For design purpose, the scale between SAC9 and SH3P2 was not respected. **B,** Volcano plot of SH3P2-GFP IP-MS after transient expression in *N. benthamiana*. X-axis displays the log fold-change (log(FC)) of proteins in SH3P2-GFP compared to the control (GFP). The Y axis shows the statistical significance, log10(p value). Dashed lines represent threshold for log(FC)>1 and p value<0.05. **C,** Table showing the peptide counts of the top hits obtained in the IP-MS (n = 3 independent pooled biological replicates), including the controls GFP and GFP/BTZ. Note that multiple protein isoforms of SAC9 are present in the *N. benthamiana* annotated proteome (Kourelis et al., 2013), and two were found in the IP-MS as SAC9 interactors. **D,** Representatives images of the colocalization between mCIT-SAC9 and SH3P2-tdTOM in a proximal (Zi) or distal (Zii) region from the plasma membrane of Arabidopsis epidermal root cells. Scale =5µm. **E,** Quantification of the number of SAC9-labelled endosomes colocalizing with SH3P2 signal per cell. N= number of replicates; n= number of roots. **F,** Representatives images of the SH3P2-sGFP localization (green) in WT and *sac9-3* mutant in which the plasma membrane was labelled using FM4-64 (magenta). **G,** Quantification of the dissociation index of SH3P2-sGFP in WT and *sac9-3* mutant in which the plasma membrane was labelled using FM4-64. N= number of replicates; n= number of roots. **H,** Representative images of the localization of *SH3P2pro:SH3P2-tdTOM* transformed in *sac9-3* heterozygous plant expressing the PI(4,5)P2 biosensor *mCIT-TUBBYc*. N = 2 biological replicates, n = 50 roots. In the plots, middle horizontal bars represent the median, while the bottom and top of each box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the interquartile range, excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and minimum values. Details of the statistical analysis could be found in Supplementary File 1H.

Figure7

1030

Figure 7. TPLATE dynamics and density at the plasma membrane is disturbed in the *sac9* mutant. A, Representatives images of the TPLATE-GFP localization in WT and *sac9-3* mutant at the plasma membrane observed by TIRF microscopy in etiolated hypocotyl. Scale = 5µm. **B,** Quantification of the density at the plasma membrane of TPLATE-GFP in WT and *sac9-3* mutant observed by TIRF microscopy in etiolated hypocotyl. Only one of the three replicates are represented. In the plot, middle horizontal bars represent the median, while the bottom and top of each box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the interquartile range, excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and minimum values. N=13 plants n=63 cells for the WT ; N=11, n=55 for sac9-3. **C,** Representatives kymograph of the TPLATE-GFP dynamics at the plasma membrane in WT and *sac9-3* mutant observed by TIRF microscopy in etiolated hypocotyl over 5 min; scale = 60s. **D,** Frequency distribution of the lifetimes of TPLATE-GFP tracks in WT (green) and *sac9-3* mutant (oranges). Results from an independent samples T-Test followed by Mann-Whitney U test is presented (see Supplementary File 1L for details). **E,** Histograms of median normalized fluorescence of TPLATE-GFP in WT (left) and *sac9-3* mutant (right) representing the density of tracks per track length. Details of the statistical analysis could be found in Supplementary File 1J. For WT, 21 cells from N=10 plants were used, n=33983 tracks; For sac9, N= 13 n=27142 tracks.

Figure8

Figure 8. Model for the mode of action of SAC9 in regulating PI(4,5)P$_2$ subcellular patterning. In wild type plants, SAC9 restricts the localization of PI(4,5)P$_2$ at the plasma membrane allowing endocytic processes to occur. In the absence of SAC9, endocytosis PI(4,5)P$_2$ accumulates in atypical endomembrane compartments that are no longer able to fuse with the TGN possibly because of their abnormal anionic lipid signature. SAC9 interacts with SH3P2 close to the plasma membrane. The defects in PI(4,5)P$_2$ patterning in absence of SAC9 leads to decreasing of the endocytic rate and the formation of membrane protuberances in contact with the plasma membrane. Note that in this model, the observed increase in intracellular PI4P levels as measured by probes is not included.

1059

Figure 1- figure supplement 1. A, Representative images of the complementation analysis of the *sac9-3* mutant. For each
construct, 24 primary T1 transformants were selected. Independent T2 and T3, T4 plants were obtained for subsequent
analysis. Line 1000-9-1; 987-5-4; 1354-12-14 used for the quantitative analysis of *sac9* complementation are shown in
Figure 1. Example of the phenotype observed in other independent transgenic lines 1000-10-h2; 987-14-5; 1354-15-11, as
well as two independent transgenic lines 1365-1-5-3 and 1356-4-1 are presented. **B**, immunoblot showing the expression of
the various SAC9 variants in independent transgenic lines (the number of the lines shown correspond to the line presented in
panel A and in Figure 1). mCIT-SAC9 and its respective mutants were detected by immunoprecipitation using an anti-GFP
antibody. Ponceau S staining serves as a loading control. The experiment was repeated three times with similar results. N =
number of roots, n= number of cells.
1069
1070

Figure 2- figure supplement 1. A, Confocal imaging of mCIT-SAC9 (green) together with the nuclear marker CENH3-RFP (magenta). Scale bars: 10 µm. **B,** Representative images of the signal corresponding to mCIT-SAC9 and FM4-64 at two focal planes (Zi and Zii) for a same cell (see C1 and C2 as examples). **C,** Representative images of the signal corresponding to mCIT-SAC9$^{\Delta CC}$ and FM4-64 at two focal planes (Zi and Zii) for a same cell (see C1 and C2 as examples). **C,** Quantification of the number of endosomes labelled by FM4-64 at two focal planes (Zi and Zii) for a same cell.

1078
1079
1080
1081 **Figure 3- figure supplement 1.** Quantification of the number of endosomes per cell labelled by mCIT-SAC9 variants which
1082 colocalized with FM4-64 at Zi. In the plots, middle horizontal bars represent the median, while the bottom and top of each
1083 box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the interquartile range,
1084 excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and minimum values.
1085 Details of the statistics are presented in the Supplementary File 1K. N = number of roots, n= number of cells.
1086
1087
1088

Figure3-supplFigure2

1089

1090

Figure 3- figure supplement 2. A, Confocal images of Arabidopsis root epidermis co-expressing *SAC9pro:mCIT-SAC9^{C459A}*
(green) and markers such as *mCH-RabA1g*; *mCH-RabD1*, *mCH-Got1p* (magenta). The plane (Zi or Zii) in each image is
mentioned, and the image display is representative for the plane used for the analysis. **B,** Percentage of colocalization
between mCIT-SAC9^{C459A} and a given endosomal compartment marker at Zi. Similar letters correspond to markers with
non-significantly different percentages of colocalization with mCIT-SAC9^{C459A} (one-way ANOVA and post hoc Tuckey
HSD pairwise test with a 95 % confidence level). In the plots, middle horizontal bars represent the median, while the bottom
and top of each box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the
interquartile range, excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and
minimum values. Scale bars: 10 µm. N= number of roots, n = number of cells.
1100

Figure 4- figure supplement 1. A, Representative confocal images of *Arabidopsis* root epidermis expressing mCIT-tagged lipid sensors for PI3P in wild-type (Col-0) and *sac9-3*[-/-] genetic backgrounds. **B,** Confocal images of Arabidopsis root epidermis expressing mCIT-tagged lipid sensors for phosphatidylserine (PS) in wild-type (Col-0) and *sac9-3*[-/-] genetic backgrounds. Fluorescence intensity is color-coded (green fire blue scale). **C,** Confocal images of *sac9-3*[-/-] Arabidopsis root epidermis expressing mCIT-TUBBYc (green) and *in vivo* stained with FM4-64 endocytic tracer (magenta); N=3 replicates, n=10 roots. **D,** Confocal images of *sac9-3*[-/-] root epidermal cells co-expressing mCIT-TUBBYc (green) and 2xmCH-PH[FAPP1] (magenta). Upper panel: fluorescent signals observed in the mock treatment; lower panel: fluorescent signal observed after 50 μM 60 min BFA treatment at the Zii plane. N=3 replicates, n=12 roots **E,** Confocal images of *sac9-3*[-/-] root cells where mCIT-TUBBYc (green) was covisualized together with various TGN markers (magenta) *in vivo* in transgenic lines expressing W129R (mCH-RabA1g) or W34R(mCH-RabA1e). N=3 replicates, n=15 roots **F,** Temporal-color coded XY 2D image of mCIT-2xPH[PLC] and mCIT-TUBBYc in *sac9-3*[-/-] root epidermis (Fiji Spectrum Temporal-color code). **G,** Confocal images of mCIT-TUBBYc (green) was co-visualized together with tdTOM-SAC9C[459A] (magenta) *in vivo* in Col-0 WT and *sac9-3*[-/-]; N=3 replicates, n=13 roots. Scale bar: 10 μm.

1117

Figure 4- figure supplement 2. A, Schematic representation of the mode of action of the PAO on the PI4P and PI(4,5)P$_2$
metabolism based on previously published work (Doumane et al. 2020; Platre et al. 2018). **B,** Confocal images of WT root
epidermal cells expressing mCIT-TUBBYc PI(4,5)P$_2$ biosensor that underwent mock, or 30 µM PAO treatments for 30- or
120- min. **C,** Confocal images of *sac9-3*[-/-] root epidermal cells expressing mCIT-TUBBYc PI(4,5)P$_2$ biosensor that
underwent mock, or 30 µM PAO treatments for 30- or 120- min. **D,** Quantification of the number of mCIT-TUBBYc labeled
intracellular compartments per cell in *sac9-3*[-/-] following mock, or 30 µM PAO treatments for 30- or 120- min. Generalized
linear mixed effect model (Poisson family) to account for root as a random factor, and two sided post hoc *lsmeans* pair wise
test with a 95 % confidence level. In the plot, middle horizontal bars represent the median, while the bottom and top of each
box represent the 25th and 75th percentiles, respectively. At most, the whiskers extend to 1.5 times the interquartile range,
excluding data beyond. For range of value under 1,5 IQR, whiskers represent the range of maximum and minimum values.
All statistical tests were two-sided. Details of the statistical analysis could be found in Supplementary File 1L. **E,** Confocal
images of *sac9-3*[-/-] root epidermal cells expressing mCIT-PH[FAPP1-E50A] PI4P biosensor that underwent mock or 30 µM PAO
treatments for 30- or 120-min. Scale bar: 10 µm. N: number of roots, n: numbers of cells.

1131

Figure5-supplFigure1

1132

1133

1134 **Figure 5- figure supplement 1. A,** Example of confocal images used for the number of FM4-64 labeled intracellular
1135 compartments per cell in Col-0 and *sac9-3*[-/-]. Scale bar: 75 µm **B,** Example of confocal images used for the number of cell
1136 with the corresponding number of BFA bodies stained by FM4-64 in Col-0 and *sac9-3*[-/-]. Scale bar: 10 µm

1137

Figure6-supplFigure1

1138

1139 **Figure 6- figure supplement 1**. Western blot analysis of TPLATE-GFP and SH3P2-GFP in WT and *sac9-3⁻ᐟ⁻*. The
1140 expression of the transgene was detected using an anti-GFP antibody. Ponceau S staining as well as western blot using the
1141 anti-tubulinα antibody served as a loading control. The experiment was repeated three times with similar results. Note that
1142 the narrow lanes correspond to the signal obtained with the crude extracts, for which the detection of TPLATE-GFP was not
1143 optimal, and therefore were not used to analyze the results.

1144

Supplementary File 1 containing:

Supplementary File 1A. Reagent and resources

Supplementary File 1B. Raw data from ULTImate Y2H SCREEN Arabidopsis thaliana - SAC9 (aa 499-966) vs Universal
Arabidopsis Normalized_dT

Supplementary File 1C. Details of the statistics corresponding to Figure 1C-D

Supplementary File 1D Details of the statistics corresponding to Figure 2D and 2G

Supplementary File 1E. Details of the statistics corresponding to Figure 3G-F

Supplementary File 1F. Details of the statistics corresponding to Figure 4B

Supplementary File 1G. Details of the statistics corresponding to Figure 5

Supplementary File 1H. Details of the statistics corresponding to Figure 6E, 6G

Supplementary File 1I. Details of the statistics corresponding to Figure 7B

Supplementary File 1J. Details of the statistics corresponding to Figure1-figure supplement 1

Supplementary File 1K. Details of the statistics corresponding to Figure 3- figure supplement 2

1158

1159   **Video 1.** Time-lapse imaging of mCIT-SAC9 using spinning disk confocal microscope (1 sec per frame)

1160   **Video 2.** Time-lapse imaging of mCIT-2xPH^{PLC} in sac9 mutant (1 sec per frame) using spinning disk confocal microscope.

1161   **Video 3.** Time-lapse imaging of T-PLATE-GFP at the plasma membrane of WT and in sac9-3 mutant using TIRF
1162   microscopy. Time lapses were acquired during 300 time-points for 300s (acquisition time 500ms).

1163

REFERENCES

Abad U, Sassi M, Traas J (2017) Flower development: from morphodynamics to morphomechanics. Philosophical transactions of the Royal Society of London Series B, Biological sciences 372 (1720). doi:10.1098/rstb.2015.0545

Adamowski, M., I. Matijević, M. Narasimhan and J. Friml (2022). "SH3Ps recruit auxilin-like vesicle uncoating factors into clathrin-mediated endocytosis." bioRxiv.
a

Adamowski M, Narasimhan M, Kania U, Glanc M, De Jaeger G, Friml J (2018) A Functional Study of AUXILIN-LIKE1 and 2, Two Putative Clathrin Uncoating Factors in Arabidopsis. The Plant cell 30 (3):700-716. doi:10.1105/tpc.17.00785

Ahn G, Kim H, Kim DH, Hanh H, Yoon Y, Singaram I, Wijesinghe KJ, Johnson KA, Zhuang X, Liang Z, Stahelin RV, Jiang L, Cho W, Kang BH, Hwang I (2017) SH3 Domain-Containing Protein 2 Plays a Crucial Role at the Step of Membrane Tubulation during Cell Plate Formation. The Plant cell 29 (6):1388-1405. doi:10.1105/tpc.17.00108

Armengot L, Marquès-Bueno MM, Jaillais Y (2016) Regulation of polar auxin transport by protein and lipid kinases. Journal of experimental botany 67 (14):4015-4037

Balla T (2013) Phosphoinositides: tiny lipids with giant impact on cell regulation. Physiol Rev 93 (3):1019-1137. doi:10.1152/physrev.00028.2012

Bates D, Mächler M, Bolker B, Walker S (2014) Fitting linear mixed-effects models using lme4. arXiv preprint arXiv:14065823

Bayle V, Fiche JB, Burny C, Platre MP, Nollmann M, Martiniere A, Jaillais Y (2021) Single-particle tracking photoactivated localization microscopy of membrane proteins in living plant tissues. Nat Protoc 16 (3):1600-1628. doi:10.1038/s41596-020-00471-4

Bayle V, Platre MP, Jaillais Y (2017) Automatic Quantification of the Number of Intracellular Compartments in Arabidopsis thaliana Root Cells. Bio Protoc 7 (4). doi:10.21769/BioProtoc.2145

Ben El Kadhi K, Roubinet C, Solinet S, Emery G, Carreno S (2011) The inositol 5-phosphatase dOCRL controls PI(4,5)P2 homeostasis and is necessary for cytokinesis. Current biology : CB 21 (12):1074-1079. doi:10.1016/j.cub.2011.05.030

Carim SC, Ben El Kadhi K, Yan G, Sweeney ST, Hickson GR, Carreno S, Lowe M (2019) IPIP27 Coordinates PtdIns(4,5)P2 Homeostasis for Successful Cytokinesis. Current biology : CB 29 (5):775-789 e777. doi:10.1016/j.cub.2019.01.043

Carland F, Nelson T (2009) CVP2- and CVL1-mediated phosphoinositide signaling as a regulator of the ARF GAP SFC/VAN3 in establishment of foliar vein patterns. The

Plant journal : for cell and molecular biology 59 (6):895-907. doi:10.1111/j.1365-
313X.2009.03920.x

Carland FM, Nelson T (2004) Cotyledon vascular pattern2-mediated inositol (1,4,5)
triphosphate signal transduction is essential for closed venation patterns of
Arabidopsis foliar organs. The Plant cell 16 (5):1263-1275. doi:10.1105/tpc.021030

Cauvin C, Rosendale M, Gupta-Rossi N, Rocancourt M, Larraufie P, Salomon R, Perrais D,
Echard A (2016) Rab35 GTPase Triggers Switch-like Recruitment of the Lowe
Syndrome Lipid Phosphatase OCRL on Newborn Endosomes. Current biology : CB
26 (1):120-128. doi:10.1016/j.cub.2015.11.040

Cheng CY, Krishnakumar V, Chan AP, Thibaud□Nissen F, Schobel S, Town CD (2017)
Araport11: a complete reannotation of the Arabidopsis thaliana reference genome.
The Plant Journal 89 (4):789-804

Clough SJ, Bent AF (1998) Floral dip: a simplified method for Agrobacterium□mediated
transformation of Arabidopsis thaliana. The plant journal 16 (6):735-743

1212

Dahhan DA, Reynolds GD, Cárdenas JJ, Eeckhout D, Johnson A, Yperman K, Kaufmann
WA, Vang N, Yan X, Hwang I, Heese A, De Jaeger G, Friml J, Van Damme D, Pan J,
Bednarek SY (2022) Proteomic characterization of isolated Arabidopsis clathrin-
coated vesicles reveals evolutionarily conserved and plant-specific components. Plant
Cell. doi:10.1093/plcell/koac071

De Matteis MA, Staiano L, Emma F, Devuyst O (2017) The 5-phosphatase OCRL in Lowe
syndrome and Dent disease 2. Nature Reviews Nephrology 13 (8):455-470.
doi:10.1038/nrneph.2017.83

Dejonghe W, Sharma I, Denoo B, De Munck S, Lu Q, Mishev K, Bulut H, Mylle E, De
Rycke R, Vasileva M, Savatin DV, Nerinckx W, Staes A, Drozdzecki A, Audenaert
D, Yperman K, Madder A, Friml J, Van Damme D, Gevaert K, Haucke V, Savvides
SN, Winne J, Russinova E (2019) Disruption of endocytosis through chemical
inhibition of clathrin heavy chain function. Nat Chem Biol 15 (6):641-649.
doi:10.1038/s41589-019-0262-1

Del Signore SJ, Biber SA, Lehmann KS, Heimler SR, Rosenfeld BH, Eskin TL, Sweeney ST,
Rodal AA (2017) dOCRL maintains immune cell quiescence by regulating endosomal
traffic. PLoS Genet 13 (10):e1007052. doi:10.1371/journal.pgen.1007052

Dettmer J, Hong-Hermesdorf A, Stierhof YD, Schumacher K (2006) Vacuolar H+-ATPase
activity is required for endocytic and secretory trafficking in Arabidopsis. The Plant
cell 18 (3):715-730. doi:10.1105/tpc.105.037978
Doumane M, Colin L, Lebecq A, Fangain A, Bareille J, Hamant O, Belkhadir Y, Jaillais Y,
Caillaud M-C (2020) iDePP: a genetically encoded system for the inducible depletion
of PI (4, 5) P2 in Arabidopsis thaliana. bioRxiv
Doumane M, Lebecq A, Colin L, Fangain A, Stevens FD, Bareille J, Hamant O, Belkhadir Y,
Munnik T, Jaillais Y, Caillaud MC (2021) Inducible depletion of PI(4,5)P2 by the
synthetic iDePP system in Arabidopsis. Nat Plants 7 (5):587-597.
doi:10.1038/s41477-021-00907-z
Dragwidge, J. M., Y. Wang, L. Brocard, A. De Meyer, R. Hudecek, D. Eeckhout, C.
Chambaud, P. Pejchar, M. Potocky and M. Vandorpe (2022). "AtEH/Pan1 proteins drive
phase separation of the TPLATE complex and clathrin polymerisation during plant
endocytosis." bioRxiv.
Flanagan CA, Schnieders EA, Emerick AW, Kunisawa R, Admon A, Thorner J (1993)
Phosphatidylinositol 4-kinase: gene structure and requirement for yeast cell viability.
Science 262 (5138):1444-1448. doi:10.1126/science.8248783
Fox J, Weisberg S (2011) The car package contains functions and data sets associated with
the book an R companion to applied regression. R-Project.
Geldner N, Denervaud-Tendon V, Hyman DL, Mayer U, Stierhof YD, Chory J (2009) Rapid,
combinatorial analysis of membrane compartments in intact plants with a multicolor
marker set. The Plant journal : for cell and molecular biology 59 (1):169-178.
doi:10.1111/j.1365-313X.2009.03851.x
Golani Y, Kaye Y, Gilhar O, Ercetin M, Gillaspy G, Levine A (2013) Inositol polyphosphate
phosphatidylinositol 5-phosphatase9 (At5ptase9) controls plant salt tolerance by
regulating endocytosis. Molecular plant 6 (6):1781-1794. doi:10.1093/mp/sst072
Gujas B, Cruz TMD, Kastanaki E, Vermeer JEM, Munnik T, Rodriguez-Villalon A (2017)
Perturbing phosphoinositide homeostasis oppositely affects vascular differentiation in
Arabidopsis thaliana roots. Development 144 (19):3578-3589.
doi:10.1242/dev.155788
Guo S, Stolz LE, Lemrow SM, York JD (1999) SAC1-like domains of yeast SAC1, INP52,
and INP53 and of human synaptojanin encode polyphosphoinositide phosphatases.
The Journal of biological chemistry 274 (19):12990-12995
Gurung R, Tan A, Ooms LM, McGrath MJ, Huysmans RD, Munday AD, Prescott M,
Whisstock JC, Mitchell CA (2003) Identification of a novel domain in two
mammalian inositol-polyphosphate 5-phosphatases that mediates membrane ruffle

localization. The inositol 5-phosphatase skip localizes to the endoplasmic reticulum
and translocates to membrane ruffles following epidermal growth factor stimulation.
The Journal of biological chemistry 278 (13):11376-11385.
doi:10.1074/jbc.M209991200

He K, Marsland Iii R, Upadhyayula S, Song E, Dang S, Capraro BR, Wang W, Skillern W,
Gaudin R, Ma M, Kirchhausen T (2017) Dynamics of phosphoinositide conversion in
clathrin-mediated endocytic traffic. Nature 552 (7685):410-414.
doi:10.1038/nature25146

Hothorn T, Bretz F, Westfall P, Heiberger RM (2008) Multcomp: simultaneous inference for
general linear hypotheses. R package version:1.0-3

Hughes WE, Woscholski R, Cooke FT, Patrick RS, Dove SK, McDonald NQ, Parker PJ
(2000) SAC1 encodes a regulated lipid phosphoinositide phosphatase, defects in
which can be suppressed by the homologous Inp52p and Inp53p phosphatases. The
Journal of biological chemistry 275 (2):801-808

Ischebeck T, Werner S, Krishnamoorthy P, Lerche J, Meijon M, Stenzel I, Lofke C, Wiessner
T, Im YJ, Perera IY, Iven T, Feussner I, Busch W, Boss WF, Teichmann T, Hause B,
Persson S, Heilmann I (2013) Phosphatidylinositol 4,5-bisphosphate influences PIN
polarization by controlling clathrin-mediated membrane trafficking in Arabidopsis.
The Plant cell 25 (12):4894-4911. doi:10.1105/tpc.113.116582

Karimi M, Depicker A, Hilson P (2007) Recombinational cloning with plant gateway vectors.
Plant physiology 145 (4):1144-1154. doi:10.1104/pp.107.106989

Kourelis J, Kaschani F, Grosse-Holz FM, Homma F, Kaiser M, van der Hoorn RA (2019)
Homology-guided re-annotation improves the gene models of the alloploid Nicotiana
benthamiana. bioRxiv:373506

Lenth R, Lenth MR (2018) Package 'lsmeans'. The American Statistician 34 (4):216-221

Leong JX, Raffeiner, M., Spinti, D., Langin, G., Franz-Wachtel, M., Guzman, A. R., Kim, J.-
G., Pandey, P., Minina, A. E., Macek, B., et al. (2022) A bacterial effector counteracts
host autophagy by promoting degradation of an autophagy component. The EMBO
journal

Manford A, Xia T, Saxena AK, Stefan C, Hu F, Emr SD, Mao Y (2010) Crystal structure of
the yeast Sac1: implications for its phosphoinositide phosphatase function. The
EMBO journal 29 (9):1489-1498. doi:10.1038/emboj.2010.57

Marković, V. and Y. Jaillais (2022). "Phosphatidylinositol 4‑phosphate: a key determinant
of plasma membrane identity and function in plants." New Phytologist.

1300

Mei Y, Jia WJ, Chu YJ, Xue HW (2012) Arabidopsis phosphatidylinositol monophosphate 5-
kinase 2 is involved in root gravitropism through regulation of polar auxin transport
by affecting the cycling of PIN proteins. Cell research 22 (3):581-597.
doi:10.1038/cr.2011.150

Mochizuki Y, Takenawa T (1999) Novel inositol polyphosphate 5-phosphatase localizes at
membrane ruffles. The Journal of biological chemistry 274 (51):36790-36795.
doi:10.1074/jbc.274.51.36790

Nagel MK, Kalinowska K, Vogel K, Reynolds GD, Wu Z, Anzenberger F, Ichikawa M,
Tsutsumi C, Sato MH, Kuster B, Bednarek SY, Isono E (2017) Arabidopsis SH3P2 is
an ubiquitin-binding protein that functions together with ESCRT-I and the
deubiquitylating enzyme AMSH3. Proceedings of the National Academy of Sciences
of the United States of America 114 (34):E7197-E7204.
doi:10.1073/pnas.1710866114

Nandez R, Balkin DM, Messa M, Liang L, Paradise S, Czapla H, Hein MY, Duncan JS,
Mann M, De Camilli P (2014) A role of OCRL in clathrin-coated pit dynamics and
uncoating revealed by studies of Lowe syndrome cells. Elife 3:e02975.
doi:10.7554/eLife.02975

Narasimhan M, Johnson A, Prizak R, Kaufmann WA, Tan S, Casillas-Perez B, Friml J (2020)
Evolutionarily unique mechanistic framework of clathrin-mediated endocytosis in
plants. Elife 9. doi:10.7554/eLife.52067

Noack L, Bayle V, Armengot L, Rozier F, Mamode-Cassim A, Stevens F, Caillaud M,
Munnik T, Mongrand S, Jaillais Y (2020) A nanodomain anchored-scaffolding
complex is required for PI4Kα function and localization in plants.
bioRxiv:2020.2012.2008.415711. doi:10.1101/2020.12.08.415711

Noack LC, Jaillais Y (2017) Precision targeting by phosphoinositides: how PIs direct
endomembrane trafficking in plants. Current opinion in plant biology 40:22-33.
doi:10.1016/j.pbi.2017.06.017

Noack LC, Jaillais Y (2020) Functions of Anionic Lipids in Plants. Annual review of plant
biology 71:71-102. doi:10.1146/annurev-arplant-081519-035910

Novakova P, Hirsch S, Feraru E, Tejos R, van Wijk R, Viaene T, Heilmann M, Lerche J, De
Rycke R, Feraru MI, Grones P, Van Montagu M, Heilmann I, Munnik T, Friml J
(2014) SAC phosphoinositide phosphatases at the tonoplast mediate vacuolar function

in Arabidopsis. Proceedings of the National Academy of Sciences of the United States of America 111 (7):2818-2823. doi:10.1073/pnas.1324264111

Paez Valencia J, Goodman K, Otegui MS (2016) Endocytosis and Endosomal Trafficking in Plants. Annual review of plant biology 67:309-335. doi:10.1146/annurev-arplant-043015-112242

Pirruccello M, Nandez R, Idevall-Hagren O, Alcazar-Roman A, Abriola L, Berwick SA, Lucast L, Morel D, De Camilli P (2014) Identification of inhibitors of inositol 5-phosphatases through multiple screening strategies. ACS Chem Biol 9 (6):1359-1368. doi:10.1021/cb500161z

Platre MP, Jaillais Y (2016) Guidelines for the Use of Protein Domains in Acidic Phospholipid Imaging. Methods in molecular biology 1376:175-194. doi:10.1007/978-1-4939-3170-5_15

Platre MP, Noack LC, Doumane M, Bayle V, Simon MLA, Maneta-Peyret L, Fouillen L, Stanislas T, Armengot L, Pejchar P, Caillaud MC, Potocky M, Copic A, Moreau P, Jaillais Y (2018) A Combinatorial Lipid Code Shapes the Electrostatic Landscape of Plant Endomembranes. Developmental cell 45 (4):465-480 e411. doi:10.1016/j.devcel.2018.04.011

Posor Y, Eichhorn-Gruenig M, Puchkov D, Schöneberg J, Ullrich A, Lampe A, Müller R, Zarbakhsh S, Gulluni F, Hirsch E (2013) Spatiotemporal control of endocytosis by phosphatidylinositol-3, 4-bisphosphate. Nature 499 (7457):233-237

Posor Y, Eichhorn-Grünig M, Haucke V (2015) Phosphoinositides in endocytosis Molecular and cell biology of lipids.

Rigal A, Doyle SM, Robert S (2015) Live cell imaging of FM4-64, a tool for tracing the endocytic pathways in Arabidopsis root cells. Methods in molecular biology 1242:93-103. doi:10.1007/978-1-4939-1902-4_9

Rivas MP, Kearns BG, Xie Z, Guo S, Sekar MC, Hosaka K, Kagiwada S, York JD, Bankaitis VA (1999) Pleiotropic alterations in lipid metabolism in yeast sac1 mutants: relationship to "bypass Sec14p" and inositol auxotrophy. Molecular biology of the cell 10 (7):2235-2250. doi:10.1091/mbc.10.7.2235

Rodriguez-Furlan C, Minina EA, Hicks GR (2019) Remove, recycle, degrade: regulating plasma membrane protein accumulation. The Plant cell 31 (12):2833-2854

Rodriguez-Villalon A, Gujas B, van Wijk R, Munnik T, Hardtke CS (2015) Primary root protophloem differentiation requires balanced phosphatidylinositol-4,5-biphosphate

levels and systemically affects root branching. Development 142 (8):1437-1446.
doi:10.1242/dev.118364

Roy A, Levine TP (2004) Multiple pools of phosphatidylinositol 4-phosphate detected using
the pleckstrin homology domain of Osh2p. The Journal of biological chemistry 279
(43):44683-44689. doi:10.1074/jbc.M401583200

Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau Donald C, Connor R, Funk K,
Kelly C, Kim S, Madej T, Marchler-Bauer A, Lanczycki C, Lathrop S, Lu Z,
Thibaud-Nissen F, Murphy T, Phan L, Skripchenko Y, Tse T, Wang J, Williams R,
Trawick Barton W, Pruitt Kim D, Sherry Stephen T (2021) Database resources of the
national center for biotechnology information. Nucleic Acids Research 50 (D1):D20-
D26. doi:10.1093/nar/gkab1112

Schmid SL, Mettlen M (2013) Cell biology: Lipid switches and traffic control. Nature 499
(7457):161-162

Schwanhäusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M
(2011) Global quantification of mammalian gene expression control. Nature 473
(7347):337-342

Shin H-W, Hayashi M, Christoforidis S, Lacas-Gervais S, Hoepfner S, Wenk MR,
Modregger J, Uttenweiler-Joseph S, Wilm M, Nystuen A (2005) An enzymatic
cascade of Rab5 effectors regulates phosphoinositide turnover in the endocytic
pathway. The Journal of cell biology 170 (4):607-618

Simon ML, Platre MP, Assil S, van Wijk R, Chen WY, Chory J, Dreux M, Munnik T, Jaillais
Y (2014) A multi-colour/multi-affinity marker set to visualize phosphoinositide
dynamics in Arabidopsis. The Plant journal : for cell and molecular biology 77
(2):322-337. doi:10.1111/tpj.12358

Simon ML, Platre MP, Marques-Bueno MM, Armengot L, Stanislas T, Bayle V, Caillaud
MC, Jaillais Y (2016) A PtdIns(4)P-driven electrostatic field controls cell membrane
identity and signalling in plants. Nat Plants 2:16089. doi:10.1038/nplants.2016.89

Song L, Wang Y, Guo Z, Lam SM, Shui G, Cheng Y (2021) NCP2/RHD4/SAC7, SAC6 and
SAC8 phosphoinositide phosphatases are required for PtdIns4P and PtdIns (4, 5) P2
homeostasis and Arabidopsis development. New Phytologist

Takagi J, Uemura T (2018) Use of Brefeldin A and Wortmannin to Dissect Post-Golgi
Organelles Related to Vacuolar Transport in Arabidopsis thaliana. Methods in
molecular biology 1789:155-165. doi:10.1007/978-1-4939-7856-4_12

Tani M, Kuge O (2014) Involvement of Sac1 phosphoinositide phosphatase in the
metabolism of phosphatidylserine in the yeast Saccharomyces cerevisiae. Yeast 31
(4):145-158

Tejos R, Sauer M, Vanneste S, Palacios-Gomez M, Li H, Heilmann M, van Wijk R, Vermeer
JE, Heilmann I, Munnik T, Friml J (2014) Bipolar Plasma Membrane Distribution of
Phosphoinositides and Their Requirement for Auxin-Mediated Cell Polarity and
Patterning in Arabidopsis. The Plant cell 26 (5):2114-2128.
doi:10.1105/tpc.114.126185

Terebiznik MR, Vieira OV, Marcus SL, Slade A, Yip CM, Trimble WS, Meyer T, Finlay BB,
Grinstein S (2002) Elimination of host cell PtdIns(4,5)P(2) by bacterial SigD
promotes membrane fission during invasion by Salmonella. Nature cell biology 4
(10):766-773. doi:10.1038/ncb854

Thole JM, Vermeer JE, Zhang Y, Gadella TW, Jr., Nielsen E (2008) Root hair defective4
encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair
development in Arabidopsis thaliana. The Plant cell 20 (2):381-395.
doi:10.1105/tpc.107.054304

van Leeuwen W, Vermeer JE, Gadella TW, Jr., Munnik T (2007) Visualization of
phosphatidylinositol 4,5-bisphosphate in the plasma membrane of suspension-cultured
tobacco BY-2 cells and whole Arabidopsis seedlings. The Plant journal : for cell and
molecular biology 52 (6):1014-1026. doi:10.1111/j.1365-313X.2007.03292.x

Vollmer AH, Youssef NN, DeWald DB (2011) Unique cell wall abnormalities in the putative
phosphoinositide phosphatase mutant AtSAC9. Planta 234 (5):993-1005.
doi:10.1007/s00425-011-1454-4

Wickham H (2016) ggplot2: elegant graphics for data analysis. springer,

Williams ME, Torabinejad J, Cohick E, Parker K, Drake EJ, Thompson JE, Hortter M,
Dewald DB (2005) Mutations in the Arabidopsis phosphoinositide phosphatase gene
SAC9 lead to overaccumulation of PtdIns(4,5)P2 and constitutive expression of the
stress-response pathway. Plant physiology 138 (2):686-700.
doi:10.1104/pp.105.061317

Yan X, Wang Y, Xu M, Dahhan DA, Liu C, Zhang Y, Lin J, Bednarek SY, Pan J (2021)
Cross-Talk between Clathrin-Dependent Post-Golgi Trafficking and Clathrin-
Mediated Endocytosis in Arabidopsis Root Cells. The Plant cell

Yoshida S, Ohya Y, Goebl M, Nakano A, Anraku Y (1994) A novel gene, STT4, encodes a
phosphatidylinositol 4-kinase in the PKC1 protein kinase pathway of Saccharomyces
cerevisiae. The Journal of biological chemistry 269 (2):1166-1172

Yperman K, Papageorgiou AC, Merceron R, De Munck S, Bloch Y, Eeckhout D, Jiang Q,
Tack P, Grigoryan R, Evangelidis T (2021a) Distinct EH domains of the endocytic
TPLATE complex confer lipid and protein binding. Nature communications 12 (1):1-
11

Yperman K, Wang J, Eeckhout D, Winkler J, Dai Vu L, Vandorpe M, Grones P, Mylle E,
Kraus M, Merceron R (2021b) Molecular architecture of the endocytic TPLATE
complex. Science Advances 7 (9):eabe7999

Zhang Y, Persson S, Hirst J, Robinson MS, van Damme D, Sanchez-Rodriguez C (2015)
Change your TPLATE, change your fate: plant CME and beyond. Trends in plant
science 20 (1):41-48. doi:10.1016/j.tplants.2014.09.002

Zhao Y, Yan A, Feijo JA, Furutani M, Takenawa T, Hwang I, Fu Y, Yang Z (2010)
Phosphoinositides regulate clathrin-dependent endocytosis at the tip of pollen tubes in
Arabidopsis and tobacco. The Plant cell 22 (12):4031-4044.
doi:10.1105/tpc.110.076760

Zhong R, Burk DH, Morrison WH, 3rd, Ye ZH (2004) FRAGILE FIBER3, an Arabidopsis
gene encoding a type II inositol polyphosphate 5-phosphatase, is required for
secondary wall synthesis and actin organization in fiber cells. The Plant cell 16
(12):3242-3259. doi:10.1105/tpc.104.027466

Zhong R, Burk DH, Nairn CJ, Wood-Jones A, Morrison WH, 3rd, Ye ZH (2005) Mutation of
SAC1, an Arabidopsis SAC domain phosphoinositide phosphatase, causes alterations
in cell morphogenesis, cell wall synthesis, and actin organization. The Plant cell 17
(5):1449-1466. doi:10.1105/tpc.105.031377

Zhong R, Ye ZH (2003) The SAC domain-containing protein gene family in Arabidopsis.
Plant physiology 132 (2):544-555. doi:10.1104/pp.103.021444

Zhuang X, Wang H, Lam SK, Gao C, Wang X, Cai Y, Jiang L (2013) A BAR-domain
protein SH3P2, which binds to phosphatidylinositol 3-phosphate and ATG8, regulates
autophagosome formation in Arabidopsis. The Plant cell 25 (11):4596-4615.
doi:10.1105/tpc.113.118307

1462