1.   **Horizontal connectivity in V1 :**

2.   **Prediction of coherence in contour and motion integration**

3.

4.   <u>Abbreviated Title :</u> Horizontal propagation of "network belief" in V1

5.

6.   Benoit Le Bec[1¶], Xoana G. Troncoso[1,3&], Christophe Desbois[1,2&], Yannick Passarelli[1],

7.   Pierre Baudot[1,4], Cyril Monier[1], Marc Pananceau[1&] and Yves Frégnac[1]*.

8.

9.   <u>Place of work and Affiliations :</u>

10.   [1]: NeuroPSI-UNIC, Paris-Saclay Institute of Neuroscience, CNRS, Paris-Saclay University, France.

11.   [2]: Coaffiliation: Ecole Nationale Vétérinaire d'Alfort, Maisons-Alfort, France.

12.   [3]: Present address: Institut Polytechnique, Palaiseau, France.

13.   [4]: Present address: Median Technologies, Sophia-Antipolis, France.

14.

15.   *: <u>Corresponding Author :</u>

16.   E-mail : <u>yves.fregnac@cnrs.fr</u> (YF)

17.

19.   # Abstract (279 words)

20.   This study demonstrates the functional importance of the Surround context relayed laterally in V1

21.   by the horizontal connectivity, in controlling the latency and the gain of the cortical response to

22.   the feedforward visual drive. We report here four main findings : 1) a centripetal apparent motion

23.   sequence results in a shortening of the spiking latency of V1 cells, when the orientation of the

24.   local inducer and the global motion axis are both co-aligned with the RF orientation preference; 2)

25.   this contextual effects grows with visual flow speed, peaking at 150-250°/s when it matches the

26.   propagation speed of horizontal connectivity (0.15-0.25 mm/ms); 3) For this speed range, the

27.   axial sensitivity of V1 cells is tilted by 90° to become co-aligned with the orientation preference

28.   axis; 4) the strength of modulation by the surround context correlates with the spatiotemporal

29.   coherence of the apparent motion flow. Our results suggest an internally-generated binding

30.   process, linking local (orientation /position) and global (motion/direction) features as early as V1.

31.   This long-range diffusion process constitutes a plausible substrate in V1 of the human

32.   psychophysical bias in speed estimation for collinear motion. Since it is demonstrated in the

33.   anesthetized cat, this novel form of contextual control of the cortical gain and phase is a built-in

34.   property in V1, whose expression does not require behavioral attention and top-down control

35.   from higher cortical areas. We propose that horizontal connectivity participates in the

36.   propagation of an internal "prediction" wave, shaped by visual experience, which links contour

37.   co-alignment and global axial motion at an apparent speed in the range of saccade-like eye-

38.   movements.

39.

40. **Introduction**

41.   Low-level visual perception is often described as a "pop-out" holistic process, elevated at the

42.   state of awareness, independently of behaviorally directed attention. At the perceptual level,

43.   Gestalt rules link stimuli according to feature similarity, spatial proximity or common fate (1, 2). Their

44.   neural implementation is thought to occur at various stages of visual cortical processing (3) and best

45.   captured, in terms of receptive field (RF) computation, as the combination of smooth "association

46.   field" operators in space (4, 5) and in time (6). However, the detailed synaptic implementation and

47.   the cortical regionalization of this dual space-time mechanism remain largely unknown (review in

48.   (7)). An unsolved issue, which is central to this study, is to determine to which degree "horizontal"

49.   connections, intrinsic to the primary visual cortex (V1), are instrumental to the neural

50.   implementation of Gestalt laws.

51.   Classically, the consensus is that binding in space and time are processed in separate cortical

52.   areas, respectively in V1 for local attributes of form and contour integration, and in MT for global

53.   motion (8-10). Although the functional expression of contour-related responses in V1 has been

54.   shown to depend on the behavioral attention context and to be task-specific (11), it requires the

55.   triggering of a stimulus-driven bottom-up process already detectable in the anesthetized brain (12-

56.   14). Local circuits intrinsic to primary visual cortex such as the long-range horizontal connections, are

57.   involved in contour integration, but the final binding process is gated and tuned in strength by

58.   selective top-down signals conveyed by re-entrant cortico-cortical pathways ((11, 15); review in (16)).

59.   In contrast, in terms of hypothetical cortical circuitry contribution, neural correlates of motion

60.   binding appear well established in MT (17-19) but far less documented in V1 (but see (20)).

61.   In spite of many intracellular receptive field (RF) studies in cat V1 (21-24), only a few have

62.   directly measured the synaptic nature of contextual influence originating from the "silent surround"

63.   (25-30). A striking feature from our own observations is that the strength of distal synaptic echoes

64.   exponentially decreases, while their latency linearly increases, with the relative eccentricity of the

65.   stimulus from the RF center. If intracellular studies quantify synaptic convergence of lateral input,

66.   optical imaging, in a converse way, monitors the full network depolarization pattern and dynamics in

67.   response to a local stimulus. Intrinsic imaging reveals the spatial extent of the diffusive pattern of

68.   intracortical activity, spread across several hypercolumns (up to 6° in cat, in (31)). Mesoscopic

69.   dynamic imaging, using voltage sensitive dyes with a temporal activation precision in the millisecond

70.   range, report depolarizing wavefronts propagating across the superficial layers of V1, observed both

71.   in the ongoing and visually driven states (non human primate (NHP) (32-36); cat (37-40); rat (41)).

72.   Their speed range is comparable across mammalian species and matches remarkably closely that

73.   inferred from our intracellular recordings in the cat (between 0.05 to 0.50 mm/ms in (26)).

74.   Intracellular recordings provide a unique way to dissect out the diversity of synaptic input

75.   connectivity pattern across individual V1 cells. Through selective averaging, they allow to infer the

76.   mean functional kernels of feedforward and lateral synaptic inputs composing the V1 receptive field.

77.   A prior study in our lab (29) proposed a new feature-specific averaging method, while keeping the

78.   mapping stimulus geometry invariant across cells in a receptive-field-centric framework. This was

79.   done by realigning each of the individual subthreshold maps relatively to a common origin and " 0° "

80.   reference axis, corresponding respectively to the center and the preferred orientation/direction of

81.   each recorded RF. The averaging of 25 RF maps demonstrated that, irrespective of single cell RF

82.   diversity, the mean synaptic connectivity kernel of V1 neurons receives predominantly iso-oriented

83.   inputs from neighboring collinear V1 hypercolumns. A remarkable feature is that its spatial

84.   organization pattern exactly mirrors the perceptual "Association Field" for collinear contours in

85.   human psychophysics (4, 42).

86.   The extensive intracellular study on dynamic Center-Surround interaction we present here - on a

87.   larger cell sample (n=65) - is based on the same premise. It seeks to maximize the paired neighbor-

88.   to-neighbor interactions between Surround and Center and quantify a novel form of contextual

89.   control on the spiking responsiveness of V1 cells. For this purpose, we have designed here a variety

90.   of multi-stroke apparent motion (AM) sequences of specific speed and spatial anisotropy, which,

91.   under precise timing and local feature co-alignment conditions, allowed us to control the temporal

92.   phase between horizontal and feedforward inputs (Figure 1A). Two sets of visual protocols were

93.   applied separately in two distinct sets of cells with complementary objectives. The "**cardinal**"

94.   protocol (Fig. 1B), was designed to : i) optimize the visibility of synaptic inputs originating laterally

95.   from the "Near" RF surround, outside the spiking "minimal discharge field" (MDF), but potentially

96.   overlapping with the subthreshold depolarizing receptive field (SRF); ii) probe for a possible axial bias

97.   in the preferred recruitment of contextual Surround information integration, using 2-3 stroke AM

98.   sequences along the two main (length/width) axes of the RF). The second series of protocols, termed

99.   "**radial**" (Fig. 1C), was designed to minimize the risk of spurious feedforward contamination when

100.   probing the locations closest to the RF borders (position D1 in Fig. 1A). Accordingly, the synaptic

101.   feedforward cortical imprint was no longer limited to the spiking discharge field (MDF), but included

102.   as well the surrounding depolarizing SRF evoked by point-like stimuli during sparse noise mapping.

103.   Conversely, this new requirement restricted the Surround stimulation to "Far" regions outside the

104.   SRF. To compensate for potentially weaker Surround responses, we increased the number of AM

105.   strokes and imposed radial symmetry in order to recruit simultaneously more sources in the RF

106.   periphery at the same relative Surround eccentricity.

107.   ------------------

108.   **Fig 1. Working Hypothesis and Visual Stimulation Protocols**

109.   *[A] : Working Hypothesis :* the AM sequence is designed so as to synchronize the synaptic activation time of

110.   horizontal volleys, evoked by the sequential presentation of 2 to 5 Gabor patches (GP) in the Surround, with

111.   the feedforward activation of the RF Center (shaded rectangle in the D0 position). The Surround GPs, regularly

112.   spaced from "Far" (D5) to "Near" (D1) periphery along the apparent motion path (horizontal axis), are flashed

113.   in succession at high contrast. The test reference GP, terminating the AM sequence, is flashed at low/medium

114.   contrast in the RF center (D0). Each orange arrow represents the lateral propagation of the presynaptic volley

115.   elicited by each distal GP. Other synaptic recruitment paths potentially also contribute also to Surround-Center

116.   modulation, in particular cascades of proximal neighbor-to-neighbor links (grey arrows).

117.   *[B] "Cardinal" Protocol :* the two columns illustrate the exploration of the two cardinal axes (horizontal for the

118.   RF orientation preference; vertical for the RF width axis) with 2 or 3-stroke AM sequences of iso-oriented

119.   stimuli ("collinear", first row; "parallel", second row) or cross-oriented stimuli (two bottom rows, same motion

120.   paths but the local inducer orientation is now orthogonal to the RF orientation preference).

121.   *[C]: "Radial" Protocol* : the two columns illustrate the various AM flow patterns (individual rows), either

122.   funnelled along a given motion axis (**SECTOR**, left), radially contracting (CP: centripetal) or expanding (CF:

123.   centrifugal), or across the full Surround (**FULL**, right). In the SECTOR condition, two opposite motion axes are

124.   symmetrically explored, aligned either with the RF orientation preference (horizontal axis (± 30°) in rows 1, 2

125.   and 4) or with its width axis (vertical axis (± 30°), row 3). In the FULL condition, all directional motion axes

126.   (discretized by 30° steps) are stimulated concurrently and intersect the RF center (horizontal grey icon). From

127.   top to bottom, each row illustrates one specific configuration of AM flow: 1) centripetal collinear flow (**CP-ISO**)

128.   from Surround to Center (red, row 1); 2) centrifugal collinear flow (**CF-ISO**) from Center to Surround (green,

129.   row 2), 3) centripetal cross-oriented flow (**CP-CROSS**) across the RF width axis (gold, row 3); 4) randomized

130.   collinear pattern (**RND-ISO**) where each successive GP location has been randomized in space and time (blue,

131.   bottom row). Note that in all the conditions other than CF-ISO, the last GP of the sequence is flashed with the

132.   optimal orientation in the RF center (D0). See Text for details.

133.   -----------

134.   Our working hypothesis posits that the spatio-temporal optimization of the control of cortical

135.   responsiveness should be reached where, *in space*, the local orientation of each Surround GP

136.   matches that of the recorded RF (as expected from the "Association Field" profile), and when, *in*

137.   *time,* the phase of the stimulation of each surround node compensates for the lag of the

138.   postsynaptic response, due to the dependency of the PSP latency on the eccentricity relative to the

139.   RF center (Fig. 1A, "latency basin" in Fig. 2A). Our functional prediction is that a latency shortening

140.   and an overall cortical gain amplification of the composite evoked PSP should be observed when the

141.   global flow speed of the AM sequence synchronizes the respective synaptic impacts of the lateral

142.   and feedforward drives onto the intracellularly recorded cell.

143.   This study addresses the importance of the Surround context dynamics relayed laterally in V1 by

144.   the horizontal connectivity, in controlling the timing and the responsiveness gain of the cortical

145.   response. It also gives evidence for neural correlates of "filling-in" responses at the earliest stage of

146.   visual cortical processing, i.e. the primary visual cortex. Remarkably, these responses are shown to

147.   be specific of co-alignment, and their timing anticipates the occurrence of the same orientation

148.   feature in the RF center, even in the absence of feedforward activation. We infer, from a detailed

149.   contextual synaptic interaction analysis, a conceptual framework which accounts for the neural

150.   correlates in V1 of specific Gestalt-like visual configurations, linking local (orientation) and global

151.   (contour integration, motion) features.

152.

153.   # Material and Methods

154.   ## 1. Animal preparation and electrophysiological recordings

155.   All experiments were performed in anesthetized (alfaxalone, Alfaxan®, 2.5 mg/kg/hr) and

156.   paralyzed (pancuronium Pavulon® (0.1 ml/kg/hr) or rocuronium Esmeron®, 4 mg/kg/hr) adult cats of

157.   either sex, in accordance with the American Physiological Society's Guiding Principles for the Care

158.   and Use of animals. This study was performed in institutionally approved animal breeding and

159.   experimental facilities (CNRS Gif Campus : Central Animal Care facilities ; authorization # D91-272-

160.   105). The scientific project was approved by the French Ministry of Higher Education and Research

161.   (MESR) (# 3414-201603111-3088501) and monitored by the local governmentally approved CNREEA

162.   Ethics Committee (CEEA – 059 : *Comité d'éthique en matière d'expérimentation animale*, Region Paris

163.   Centre et Sud). Experiments where continuously supervised by a veterinary DVM. Every effort was

164.   made to minimize stress and suffering. Anesthesia depth, analgesia level and the general

165.   physiological state stability were continuously monitored from the recording and analysis of EKG and

166.   extradural ECoG signals. End-tidal $CO2$ was maintained between 3.5 and 4.5%, and body

167.   temperature was regulated around 38°C throughout the experiment. Once the animal's head was

168.   fixed to the stereotaxic frame, and the craniotomy performed, we checked again the presence of a

169.   slit pupillary state, indicative of adequate anesthesia during the surgery phase. Then, atropine (1%)

170.   and phenylephrine (5%) were applied to both eyes to dilate pupils, block the accommodation and

171.   withdraw nictitating membranes. Artificial pupils and corrective lenses were used to protect the

172.   eyes and focus the backprojection of the eye fundus on the stimulation screen. At the end of the

173.   recording session (48 to 80 hours), the animal were sacrificed using either an overdosage of

174.   pentobarbital (180 mg/kg, iv) or an intra-cardiac perfusion performed after alfaxalone injection rate

175.   was increased high enough to induce a flat EEG.

176.   *In vivo* intracellular recordings of cortical cells were performed in the primary visual cortex (area

177.   17) using sharp Glass electrodes pulled from 1.5-mm-thick borosilicate capillaries (World Precision

178.   Instruments Inc.). Electrodes were filled with a 2M potassium methyl sulfate (pH ranging from 7.2 to

179.   7.4), 4 mM KCl solution for compensating the tip potential, and 1.5% biocytin. Electrode resistances

180.   before impalement ranged between 70 and 130 MΩ. Cells were recorded in current clamp bridge

181.   mode using an Axoclamp 2A (Axon Instrument inc.) or a SEC-05X (NPI inc.). The average resting

182.   potential was between -62 to -78 mV (mean ± σ = -66,8 ± 8,3 mV) with no or a slight holding current

183.   (<= -0.5 nA) when needed. Signals were filtered and amplified with a CyberAmp380 (Axon Instrument

184.   inc.) and digitized at 10 kHz by an ITC-1600 (InstruTECH®) card. Based on recording depth and the

185.   laminar location and morphological identification of fourteen biocytin labeled cells, the overall

186.   sampling was distributed from supra- to infra-granular layers. However, in view of the limited

187.   number of labelled cells (experimentally constrained by gliolysis increase with the length of

188.   recording) and a slight bias for recordings in deep layers, no laminar study was done in this reduced

189.   sample.

190.

191.   ## 2. Visual stimulation

192.   For the "cardinal" protocols, visual stimuli were generated, using the ELPHY in-house software

193.   (G. Sadoc, UNIC-CNRS), on a CRT P31 green monochrome monitor positioned at 57 cm (1 cm = 1° of

194.   visual angle) at refresh rate of 60 Hz (19 inches, 640 x 480 pixels). For the "radial" protocols, we used

195.   gamma-corrected LCD green monitor (27 inches screen, 1920 x 1080 pixels) at 28.5 cm from the cat's

196.   retinal plane (0.5 cm = 1° of visual angle) at a refresh rate of 144 Hz .

197.   ### 2.1. Initial quantification of RF optimal features :

198.   The receptive field (RF) position of each recorded neuron and its ocular dominance were initially

199.   established using moving light bars. Stimulation was done through the dominant eye for the rest of

200.   the experiment. Subthreshold ON and OFF (SRF) sub-regions and spike minimal discharge fields

201.   (MDF) were then mapped by forward correlation of visual responses to a random "sparse noise" (SN)

202.   protocol : white (40 cd/m^2) and dark (1 cd/m^2) non-overlapping squares were presented one at a

203.   time for 16-34 ms against a grey background (20 cd/m^2) in each position of the exploration grid (8x8

204.   or 20x20 pixels) centered over the coarse RF region. In some cells, an additional 1D mapping was

205.   done with an optimally oriented thin long bar, flashed randomly at various eccentricities across the

206.   RF width (Fig. 2A).

207.   Once the RF location defined, the preferred orientation, phase and spatial frequency of the cell

208.   were determined using a dense Gabor Noise (GN) stimulation protocol (43) : flashed (16-34 ms)

209.   Gabor patches (GPs, adjusted in size to the s-DF extent) were randomly presented at six different

210.   orientations (with 30° steps), four phases (0°, 90°, 180° and 270°) and five spatial periods (ranging

211.   from 0.1 to 4.9 cycles per degree of visual angle). For each cell, the parameters of the Gabor patch

212.   eliciting the strongest response in the RF center were retained to compose a template of preferred

213.   features of the inducer (independently of its position) in the subsequent apparent motion (AM)

214.   sequences. In order to evoke a sizeable synaptic "horizontal" response, the GP flashed in the

215.   Surround were always presented at high contrast (mean $\pm$ σ = 92.5 $\pm$ 11.5 %; range 75-100%),

216.   whereas the feedforward central stimulus was flashed in the MDF either at low or medium contrast

217.   (mean $\pm$ σ = 40,6 $\pm$ 24.5 %; range (25-100%)).

218. **2.2. "Cardinal" Apparent Motion (AM) Protocol :**

219.   In order to assess the dependency of the center-surround interaction on local orientation and

220.   global motion, we first designed two/three-stroke center-surround AM protocols for which the

221.   Gabor patches were flashed along the two "cardinal" RF axes defined by the RF preferred orientation

222.   and width (Fig. 1B). We looked at the modulatory effect that a two-stroke activation of the "silent"

223.   surround would produce (along each of these axis) on the response to a test stimulus flashed within

224.   the MDF (the third stroke). Randomly interleaved three-stroke stimulation sequences were applied

225.   moving away from the receptive field center ("centrifugal" (CF) condition, green arrows in Fig. 1B), or

226.   from the surround to the center ("centripetal" (CP) condition, red arrows) from one side of the RF at

227.   a time.

228.   The size of the Gabor patch mask was set at 150% of the length of the spiking discharge field

229.   while the standard deviation of its Gaussian envelope was set at 20%. The stimulus inter-node

230.   distance was of 120% of its length, often leading the more proximal GP to partially encroach on the

231.   border of the impulse SRF. The orientation of the local inducer could be defined two ways, either in

232.   relation with the RF orientation preference, or in relation with the AM motion axis. These two

233.   analysis contexts will be considered separately in the Results. Here, by sake of uniformity with the

234.   "radial" protocol, the ISO- and CROSS- conditions are defined relative to the chosen AM axis. The

235.   complete set of Surround-Only stimulations (Fig. 1B, left column) included 8 conditions [CP-

236.   ISO_main-axis (red arrows), CP-ISO_width-axis (magenta), CP-CROSS_main-axis (cyan), CP-

237.   CROSS_width-axis (gold), CF-ISO_main-axis (green), CF-ISO_width-axis, CF-CROSS_main axis, CF-

238.   CROSS_width axis (brown)] X 2 direction conditions (one side at a time)], to which 2 control

239.   conditions (1 Center-Only + 1 Blank trial, where no stimulus was presented) were added. Hence a

240.   total of 18 conditions was completed in 13 cells. Center-Surround interactions were tested in 23 cells

241.   with full AM sequences terminating in the RF center (Fig. 1B, right column), under 10 conditions [4

242.   conditions (CP-ISO_main-axis, CP-CROSS_width-axis, CF-ISO_main-axis, CF-CROSS_width-axis) X 2

243.   directions (one side at a time) + Center-Only stimulation + 1 blank]. For each study (Surround-Only;

244.   Surround-Center), the different stimulus conditions were interleaved randomly in each trial block.

245.   Recordings lasted for 20 blocks.

246.   Given the constraint of the monitor distance (57 cms), its refreshing rate (60 Hz), and the fixed

247.   duration of the static Gabor inducer presentation (16 ms), individual cell AM speed values only

248.   depended on the size of the recorded RF which was used to define the sequential interstimulus step

249.   distance. Accordingly, AM speeds were preset in the suitable range of values to compensate for the

250.   dependency of visual PSP latency with relative eccentricity (26, 29).

251. **2.3. "Radial" Apparent Motion (AM) Protocol :**

252.   This series of protocols was done with a larger (27 inches) and faster screen (144 Hz), using a

253.   shortened viewing distance (28.5 cms), which allowed a more extensive exploration of the "Far"

254.   surround. The visual field was paved with a grid composed of 5 concentric rings of increasing

255.   eccentricity centered on the RF covering up to 25° in the surround (Fig. 1C). To the difference of the

256.   "cardinal" protocol, where stimulus size and spacing were defined relatively to the spiking MDF size,

257.   the distance between each outer ring and the patch size were adjusted in this new protocol to the

258.   full extent of the subthreshold depolarizing receptive field (SRF) mapped with impulse stimuli. This

259.   change in the RF core definition, from that used in the "cardinal" protocol condition, was done to

260.   prevent, on a cell-by-cell basis, a direct feedforward contamination by the stimulation of the most

261.   proximal Surround location (position D1 in Fig. 1A).

262.   Additionally, in the "radial" protocol, we designed Apparent Motion sequences whose spatio-

263.   temporal coherence was parametrized to dissect the dependency of the inducer orientation relative

264.   to the motion path, as well as the spatio-temporal congruency within the AM flow (see Results). For

265.   each recorded cell, the AM sequence of interest was the centripetal iso-oriented configuration (**CP-**

266.   **ISO**, top "red" row in Fig. 1C), where the local inducer orientation was either co-aligned along the

267.   preferred orientation axis of the recorded RF (SECTOR configuration, left column in panel 1C) or

268.   along the global motion axes (FULL configuration, right column). Three supplementary visual AM

269.   configurations were tested, in which the spatio-temporal structure of the local Gabor patch (GP, local

270.   inducer) sequence was altered, while keeping the overall stimulus energy distribution unchanged: i)

271.   in the first control condition, namely the centrifugal iso-oriented AM sequence (**CF-ISO**, second

272.   "green" row from the top, Fig. 1C), the temporal order of the sequential presentation of the co-

273.   aligned GPs was reversed, resulting in a centrifugal (CF) flow from Center to "Far" Surround

274.   locations; ii) the second control was designed to study the contextual impact of the local GP

275.   orientation relative to the motion axis. In the centripetal cross-oriented AM condition, the local

276.   orientation of the inducer was orthogonal to the motion axis (**CP-CROSS**, third "yellow" row from the

277.   top, Fig. 1C); iii) in the third control condition, the same retinal space (SECTOR or FULL) as in the CP-

278.   ISO condition of interest was stimulated with the same GPs (i.e., with the same number, features,

279.   and energy distribution at each step of the AM) but the coherence of the flows was randomized both

280.   in space and time (**RND-ISO**, bottom "blue" row). For each block, each stimulus location was visited

281.   only once, and the last stimulus location was the RF center. Apart from the RND-ISO condition (Fig.

282.   1C, 4[th] row), the repeated sequences were identical across blocks (pseudo-randomization).

283.   The stimulus set in the "radial" protocol - where all contextual conditions were explored - was

284.   composed of 16 dynamic apparent motion sequences (4 conditions (CP-ISO (red arrows), CF-ISO

285.   (green), CP-CROSS (gold), RND-ISO (blue)) X 2 trajectories (with or without Center stimulation) X 2

286.   configurations (SECTOR and FULL), and 21 stimulation types where Gabor Patches were flashed in

287.   isolation (5 peripheral locations X 2 conditions X 2 configurations + 1 GP flashed in the Center). In

288.   each block of trials, each stimulus was presented only once. Additionally, 6 "blank" trials were also

289.   included in each block. The resulting set of 43 stimulations was presented in random order in each

290.   block. Recordings lasted for 19-50 blocks (median 50).

291.   In order to optimize synaptic summation between Surround and Center stimulation, the AM

292.   sequence speed was function of the spatial extent of the SRF and of the stroke duration of Gabor

293.   patches and adjusted cell-by-cell in order to fit the latency basin of "Far" surround response

294.   latencies. In the "radial" protocol where AM speed and stroke duration could be independently

295.   parametrized, the stroke duration was in the 30 ms range (± 9 ms). It was 16.6 ms in the "cardinal"

296.   protocol.

297.

298. 3. Intracellular Data analysis

299. 3.1. Data formatting :

300.   Spikes were threshold-detected in the raw traces and replaced by alpha functions in the membrane

301.   potential (Vm) traces ($\alpha(t)=a*t*\exp(-t/\tau)$) , where "a" is the slope at the peak of the second

302.   derivative of the rising phase, and τ is the half-width of the spike), as done in previous studies (29).

303.   Resulting traces were band-pass filtered (0.1-300 Hz) and peristimulus triggered waveform (PSTW)

304.   were averaged across trials and conditions. Prior to any statistical analysis, all traces were down-

305.   sampled to 1 kHz and smoothed using a sliding average window of 7 ms. For spikes, individual

306.   peristimulus time histograms were smoothed using a Gaussian window (σ= 3 ms). For a given trial

307.   block containing all conditions, the corresponding mean activity (average Vm and Spiking discharge

308.   rate) was subtracted from the raw PSTWs and PSTHs.

309.   **3.2. Response quantification and statistical significance** :

310.   For the "cardinal" protocol, the control Z-statistics for Vm (mean and standard deviation) were

311.   computed with a 4 kHz sampling rate over a 100-ms sample of ongoing activity preceding the

312.   stimulation. The response waveforms were thresholded in amplitude above the 95% one-sided

313.   confidence interval. The "integral response" was defined as the integral value of the thresholded

314.   depolarizing waveforms over a predefined response integration window (0-250 ms). For each cell,

315.   Wilcoxon paired signed rank (WPSR) tests were applied to compare cell-specific responses (n=23)

316.   evoked for two distinct contextual conditions. The contextual modulation was quantified by a

317.   facilitation/depression index given by the ratio between the significant integral responses observed

318.   in the center-surround vs center-only conditions (or between two different contextual conditions of

319.   interest). Similar processing was applied at the spike rate level. Twenty-three cells constitute the

320.   population used for the Results section for the "cardinal" protocol.

321.   For the "radial" protocol, since a larger spectrum of stimulus conditions was used, more extensive

322.   statistical analyses were done on the individual cell's responses, without or with selection criteria.

323.   All significance tests fell into two categories: quantifying the significance of a response to a stimulus

324.   or comparing data statistics between two conditions. To do the former, we relied on the activity

325.   during blank trials, which provides the null distribution for visual responses (i.e., the activity that

326.   would occur should the cell not respond to the stimulus, which was our "null" hypothesis).

327.   Consequently, for each cell, the average ongoing synaptic or spiking activity recorded during "blank"

328.   trials was subtracted point by point from each evoked PSTW/PSTH trial activity. This was done to

329.   remove, on a cell-by-cell basis, any experimental artifact or activity change common to the blank and

330.   stimulations, while avoiding a loss of information on the Vm distribution statistics. This preprocessing

331.   explains why the confidence intervals obtained from the permutation test are centered on 0 when

332.   evaluating significance regarding spontaneous activity (and why there are negative spikes). For

333.   studying the dependency on the stimulus context, we used cell-by-cell paired comparison between

334.   distinct tested conditions. Statistical analysis was then done using a randomization permutation test

335.   with 10^4 repetitions.

336.   From these computations, we extracted the null hypothesis waveform envelope for a given

337.   confidence interval at each point in time. Only cells with a response to the Center-Only condition

338.   significantly larger than the blank ($p < 0.01$) in the interval 0-to-120 ms ("0" being the onset of the

339.   last stimulation in the center when the RF was actually stimulated, "120"ms integrating most of the

340.   feedforward onset and offset responses) were considered for further analysis. For PSTW and PSTHs,

341.   respectively thirty-seven and twenty-two cells were retained after this initial screening and

342.   constitute the population used for the Results section for the "radial" protocol. In the remaining

343.   fifteen "nonspiking" cells, analysis was restricted to subthreshold activity.

344.   ## 3.3. Quantification of the contextual change in response gain and phase

345.   The concept of "gain control", dominant in cortical physiology or psychophysics, tends to

346.   reduce information transfer functions to simple scalar static "gain" measurements. It consequently

347.   ignores the kinetic aspects of cortical transmission modulation. The intracellular recordings approach

348.   developed here allows to compare both the gain and the phase (or advance/delay as analyzed here)

349.   of the impulse response (e.g. the transfer function in linear system theory) in various contextual

350.   conditions. When using this engineering terminology, we make the simplifying assumption that the

351.   test stimulus presentation is short enough in duration and localized enough in space, so that the

352.   feedforward test stimulus response gives an approximation of an impulse response in space and

353.   time.

354.   We quantified the time-course and the strength of the contextual control of the cortical transfer

355.   function by measuring the changes in i) subthreshold latency (i.e. the "phase") and ii) the integral

356.   value of the depolarizing postsynaptic potential (PSP) response to the complete AM sequence

357.   terminating in the recorded cell's receptive field (i.e. the "gain"). For the "cardinal" protocol, the

358.   latency measure was defined by the onset of the mean PSP reaching significance (Z-test, $p<0.05$). For

359.   the "radial" protocol, we chose to use a more conservative measure to avoid spurious detections of

360.   fluctuations in the ongoing synaptic noise. Accordingly, the latency change (Δ latency) was measured

361.   at half-height of the Center-Only response amplitude peak, separately for Vm and spikes. For the

362.   response gain (Δ response), we measured the change in the PSP depolarizing envelope integral (and,

363.   eventually, spiking integral), relative to the Center-Only response, over the whole time-course of the

364.   test-response. Here, the rationale was to integrate differences in evoked responses for all possible

365.   time relationships between feedforward and lateral input. The same statistical analysis

366.   (randomization test, 10^4 repetitions; $p<0.05$) was repeated for PSTWs and PSTHs for both criteria.

367.   The paired-analysis was replicated across conditions to measure statistical significance between the

368.   main condition of interest (CP-ISO) and the other contextual conditions.

369.   ## 3.4. Surround-Only responses*:*

370.   For the "cardinal" protocol, statistical significance was assessed using Z-test statistics on

371.   response integrals ($p<0.05$). For the "radial" protocol, we used a permutation test (10^4 repetitions;

372. $p< 0.01$). Due to the noisy nature and the weak amplitude of purely lateral responses, we only

373.   retained Vm waveforms for which the statistical significance threshold was trespassed for at least 15

374.   consecutive milliseconds in the interval 0-120 ms.

375.   ## 3.5. Linear predictions and Dynamic non-linearities:

376.   For the "radial" protocol, we determined in each individual case whether the actual Surround-

377.   Only responses to dynamic sequences were larger than (or equal to) the sum of the temporally

378.   shifted responses to each GP flashed in isolation at each location (reproducing the spatial and

379.   temporal ordering of a virtual AM sequence). This linear predictor was in a first step compared to

380.   the actual AM dynamic sequences before they reached the RF Center. A non-linearity was detected

381.   in cells where the actual build-up of anticipatory activity during AM sequences deviated significantly

382.   from the linear predictor. Because GPs, when flashed in isolation in the close vicinity of the SDF,

383.   already induced significant depolarizing responses (29), we considered Surround-Only synaptic

384.   waveforms as significantly larger than their "static" linear predictor, if they crossed the upper limit of

385.   the confidence interval for a minimum duration of 7 consecutive ms in the interval 0-120 ms

386.   (permutation test, 10^4 repetitions, p < 0.05). A second predictor was calculated by subtracting the

387.   Center-Only response from the complete AM "Surround-then-Center" response. Note that the

388.   Surround-Only component potentially included the synergetic effect arising from the interaction

389.   between the successively recruited distal input sources in the Surround. This two-term

390.   (Center/Surround) linear predictor was compared to the observed "Surround-Only" AM response to

391.   test for non-linear interactions between the integration of the AM contextual information in the

392.   "Near" Surround and the flashed feedforward drive.

393. **3.6. Population analysis:**

394.   In order to compare responses between cells, each individual flow pattern's specific geometry –

395.   defined on a cell-by-cell basis according to the preferred orientation and direction of each cell – was

396.   realigned on a common "0°" reference axis (visualized by the semi-horizontal axis pointing to the

397.   right in all Figures), regardless of the absolute orientation preference of each cell. No normalization

398.   of responses was used in the "cardinal" protocol. In the "radial" protocol, the amplitude of the

399.   PSTWs and PSTHs of each cell were first normalized to the peak of the Center-Only response. Before

400.   averaging, PSTWs and PSTHs of all the cells were also realigned with respect to the onset latency of

401.   their individual Center-Only response. This common origin was defined as the time abscissa of the

402.   first point (blue dot) of the average Center-Only response departing from the "blank" response mean

403.   by more than 3σ . This process, replicated independently for subthreshold (Vm) and spiking

404.   responses, allowed a realignment of all normalized cell-by cell profiles and the calculus of the mean

405.   population response profile. Note that the peak value of the average population PSP can be less than

406.   the norm (1.0) when the onset-to-peak rise time varies across cells.

407.

408. **Results**

409.   This intracellular study (65 cells) of Center-Surround interactions in V1 of the anesthetized cat is

410.   based on two distinct visual protocols, applied in two different sets of cells (see Methods). The first

411.   protocol, termed "**cardinal**" (Fig. 1B), recruiting both "Near" and "Far" RF surround, was applied to a

412.   first batch of 23 cells. The second series of protocols, termed "**radial**" (Fig. 1C), limiting the Surround

413.   stimulation to "Far" regions outside the SRF only, was performed on a second batch of 42 cells.

414.   Accordingly, response averages were done separately for the two visual stimulation protocols.

415. **1. Probing synaptic responses from the "silent" Surround**

416. **1.1. Spatial summation**

417.   The "silent Surround" is classically defined as the region outside the minimal discharge field

418.   (MDF), where impulse stimuli do not evoke a significant spiking response. However, the intracellular

419.   mapping of synaptic responses with sparse or dense noise (SN or DN), devoid of spatial summation,

420.   shows a broader subthreshold (non-spiking) receptive field (SRF) extending beyond the border of the

421.   MDF, and often invading the opponent contrast discharge field, even in Simple cells (43, 44). This

422.   defines a Near-Surround domain, connex to the MDF, where the contribution of purely lateral

423.   synaptic input cannot be unambiguously disentangled from subthreshold feedforward activation

424.   (25).

425.   The mapping of the minimal discharge field (MDF) with small light or dark squares (0.2-0.9°)

426.   showed on average a spiking core field of 1-3° of visual angle. Overall, the size of the impulse

427.   subthreshold depolarizing receptive field (SRF) of the cortical neurons recorded in this study ranged

428.   from 2.5° to 7.5° (5,1 ± 1,6°), for eccentricities from the area centralis between 1.1° and 8.3° (4.0 ±

429.   1.6°). This initial mapping was used to define, in a conservative way in each cell, the retinotopic

430.   imprint of the feedforward input, whether it produced a spike (MDF) or only a depolarizing

431.   subthreshold response (SRF).

432.   In contrast, elongated light bars (5-12° length) flashed across the RF width axis, which increases

433.   input spatial summation, elicited synaptic responses, in the same cell, originating much further away

434.   from the RF center (7.1° for bars against 2.5° for SN in the example cell in Fig. 2A). Using annular

435.   gratings at the optimal phase and orientation, with still larger spatial summation, our lab previously

436.   demonstrated distal synaptic responses originating beyond 10° of relative eccentricity (Fig. 2B, taken

437.   from (26)). Pooling these different observations together establishes that the recruitment efficacy of

438.   lateral connectivity evoking Surround responses, hence their recording detectability, strongly

439.   depends on the level of spatial summation, and is selective to the orientation feature of the stimulus

440.   flashed in the periphery (26, 27, 45).

441.   ------------

442.   **Fig 2. A synaptic view of visual V1 receptive fields**

443.   *[A] : Latency basin of synaptic responses relative to the RF Center :* Top left, spatial maps of the ON- and OFF-

444.   Depolarizing Fields latencies using sparse noise (SN) mapping. Heat scale: color code for latency values (from 0

445.   (black) to 200 ms (yellow)). Bottom left, 1D-mapping of ON- and OFF- PSP responses in the same cell, evoked

446.   by long bars (7.1° x 0.7°) flashed at the optimal orientation and positioned at different eccentricities (ordinates,

447.   ranging from -4° (bottom) to +6° (top)) across the RF width axis. Thick boxes (along the left ordinate axis)

448.   designate the positions which are in overlap with the SRF mapping using SN (above, top left). Red (ON) and

449.   blue (OFF) linear regressions illustrate the 1D-latency basin profiles of Surround-Only subthreshold responses.

450.   The linear fits of latency versus eccentricity give apparent horizontal speeds of propagation estimates of 0.18-

451.   0.38 mm/ms for ON- (red, left) and OFF- (blue, right) responses.

452.   *[B] Dependency on spatial summation* (adapted from (26)) : From top to bottom, phase-reversal (2 Hz)

453.   responses to disk [1] and annular gratings [2 & 3], for increasing inner border eccentricities from the

454.   RF center (0° in [1], 5.6° in [2] and 10.3° in [3]). Note the decay in response strength and the increase of onset

         latency in Surround-Only responses [2 & 3] with eccentricity (reaching 32 ms for the "Far" Surround annulus).

455.   *[C] Apparent Speed of Horizontal Propagation (ASHP) distribution* : Stacked data histograms from (26)

456.   using long bars (1D-mapping, blue) or sparse noise (green), and from this study (Gabor, light green). Note that,

457.   independently of the probe stimulus, most AHSP values inferred from Surround latency basin slopes range

458.   between 0.05 and 0.60 mm/ms.

459.   ----------------

460. **1.2. Latency dependency of Surround responses with relative eccentricity**

461.   In addition to the spatial extent of the RF, a pivotal observation, on which our working

462.   hypothesis is grounded, is that the further away from the MDF center, the latency of subthreshold

463.   surround responses progressively increases in a linear fashion with relative eccentricity. Fig. 2A

464.   illustrates the spatio-temporal latency maps of depolarizing and spiking responses of a V1 cell for

465.   sparse noise (top panel) and of synaptic subthreshold PSPs for optimally oriented long bars flashed at

466.   different eccentricities across the RF width axis (bottom panel). In this example, because spatial

467.   summation is more efficient for optimal lines than squares, surround responses become detectable

468.   on a much larger spatial extent. By measuring the onset response latencies at each tested

469.   eccentricity in the 1D-plots of ON- and OFF-responses, and applying linear regressions between the

470.   paired sets of values, one can infer estimates of the apparent speed of horizontal propagation

471.   (ASHP), ranging between 0.18 and 0.38 mm/ms. A similar observation can be done using optimally

472.   oriented and phased isotropic annular gratings centered on the RF. In the example illustrated in

473.   Figure 2B, spatial summation and response detectability still increase and the resulting latency shifts

474.   can reach up to 20-50 ms for relative eccentricity values as large as 11°.

475.   On the whole, by comparing past and present studies using a variety of stimuli (light/dark squares,

476.   long edges, Gabor patches), we conclude that the dependency of "Surround-Only" responses with

477.   relative eccentricity from the RF center remains a hallmark of lateral processing. As shown in Figure

478.   2C, whatever probe stimulus used, the inferred apparent speed of horizontal propagation (ASHP)

479.   remains qualitatively the same (SN in green; 1D mapping in blue, Gabor in light green, in spite of the

480.   fact that the detectability of "Surround-Only" responses depends strongly on spatial summation .

481. **1.3. Apparent Motion Axial sensitivity and Orientation tuning**

482.   Once the Gabor template optimizing the Center response was determined (see Methods), each

483.   RF and/or its Surround were tested with 2-stroke (Surround-Only) or 3-stroke apparent motion (AM)

484.   sequences along the cardinal RF axes (Fig. 1B). These "cardinals" axes, extending symmetrically

485.   around the RF center, correspond, one, to the "main" orientation preference axis (represented

486.   arbitrarily by the horizontal, in all the figures) and the other, to the RF width axis (vertical). For each

487.   of these axes, the orientation of the local Gabor inducer stimuli could be either the same, "**ISO-RF**",

488.   or orthogonal, "**CROSS-RF**" to that of the RF orientation preference axis.

489.   Examples of Surround-Only AM responses of the "cardinal" protocol are illustrated in Fig. 3A.

490.   Their magnitude can reach the size of the response to a low contrast input in the center : compare

491.   Surround-Only CP-ISO AM responses (red waveform) along the preferred orientation axis with the

492.   Center-Only response (black waveform). As a reference, the center insert in Fig. 3A illustrates the

493.   mapping of the SRF border (white contour) and the ON- and OFF-discharge fields (respectively, filled

494.   in red and blue). The surround source locations of GPs used in the AM sequence partially encroach

495.   on the SRF border, but not on the MDF. The AM evoked synaptic responses are much stronger than

496.   the impulse SN response, as shown by the absence of depolarizing responses to light (lower dark red

497.   trace) and dark (blue trace) pixel stimulations, presented in the center of the lesser eccentricity

498.   peripheral Gabor location. In this example, as in most recorded RFs, the centripetal Surround-Only

499.   AM response along the preferred orientation axis (CP-ISO, left and right red waveforms) is the

500.   strongest among the responses evoked by the Surround stimulation alone : compare it with the CP-

501.   CROSS conditions along the RF width axis (top and bottom, gold waveforms), for which responses are

502.   much weaker.

503.   ---------

504.   **Fig 3. Spatio-temporal and axial selectivity of "Surround-Only" responses to Centripetal AM flow**

505.   *[A] Single Cell example* **:** Central inset cartoon : ON and OFF discharge fields, respectively in red and blue.

506.   White contour delineates the subthreshold receptive field (SRF). For comparison, respective positions of the

              Gabor inducer stimuli flashed in the surround are overlaid on the RF map. Two-stroke "Surround-Only" AM

507.   responses : comparison of "collinear" (red) and "parallel" (gold) responses evoked respectively along the

508.   orientation preference axis and across the width axis of the RF, with the Center-only response (black

509.   waveform). Bottom left corner, Vm subthreshold responses for sparse noise (ON (dark red) and OFF (dark blue)

510.   waveforms).

511.   *[B] Dependency of Surround-Only responses on the orientation of the inducer* : Right, top and bottom panels

512.   illustrate the Gabor inducer features and the color codes for the motion axis explored respectively in the ISO-

513.   and CROSS-configurations. The color code conventions are : ISO-RF: red for "collinear" along the orientation

514.   preference axis; gold for "parallel" across the RF width axis; CROSS-RF: cyan for "parallel" across the

515.   orientation preference axis; magenta for "collinear" along the RF width axis. See Text for details.

516.   -------------

517.   A complete study of the dependency of the Surround-Only AM responses on the GP inducer

518.   orientation relative to the RF preferred orientation was replicated in 13 cells. The average responses

519.   of this population have been computed by realigning arbitrarily the orientation and direction

520.   preference of each recorded cell respectively with the horizontal and right pole axes. The mean

521.   population tuning responses are illustrated in Fig. 3B, where the different AM flow conditions are

522.   schematized by color-coded arrows in the two right inserts. The top insert regroups the specific

523.   centripetal AM flows where the inducer is either "collinear" (red) or "parallel" (gold)) with the RF

524.   preferred orientation (ISO-RF condition). The bottom insert details the stimulation cases where the

525.   inducer is orthogonal with the RF preferred orientation (CROSS-RF: cyan along the main axis,

526.   magenta along the RF width axis).

527.   This reveals two effects of the Surround contribution. The first one is a non-specific

528.   depolarization component found for all AM configurations, regardless of the orientation of the

529.   inducers relative to that of the RF. Globally, the recruitment of the "Near" periphery significantly

530.   rose the level of excitability of the recorded cells (Wilcoxon paired signed rank test on integral

531.   activity values between the condition of interest and the resting state, n=13, $p<0.001$). At the single

532.   cell level, a significant depolarization compared to the resting state (one sided z-score, $p<0.05$) was

533.   evoked in 76% of cells for the CP-ISO "collinear" (red) condition, and in 55% in the "parallel" (gold)

534.   condition at high AM speed. Pooling together, for each cell, responses to all CP configurations,

535.   significant spiking responses were observed in 59% of the cases. Since the classical MDF center

536.   location was kept unstimulated, the most likely explanation is that, in the "cardinal" protocol, all

537.   centripetal Surround-Only flows still recruit synaptic input originating from the feedforward SRF in

538.   addition to the proximal horizontal input.

539.   The second component of the surround contribution is <u>orientation selective</u> and its impact can be

540.   seen in Figure 3B, riding on the top of the non-specific component. It reflects the dominance of

541.   Surround responses evoked by centripetal flows of Gabor elements "co-aligned" with the RF

542.   preferred orientation (horizontal axis in Fig. 3B) : the evoked depolarizing Surround-Only

543.   subthreshold response elicited by CP-ISO collinear flow along the main RF axis (red trace, horizontal

544.   axis in all figures) is, on average (n=13)**,** 2.6 times larger than that evoked by CP-ISO- (magenta) or

545.   CP-CROSS- (gold) flows along the RF width axis, and that evoked by CP-CROSS- flows (cyan) across

546.   the RF main axis. A strong axial response heterogeneity is indeed noticeable in the polar

547.   representation of figure 3B, when considering separately the main and the RF width axes. On the one

548.   hand, the comparison of the CP-ISO and CP-CROSS mean subthreshold response profiles shows a

549.   clear orientation and axial bias of the "Surround-only" responses for Gabors co-aligned along the

550.   orientation preference axis (red waveforms). On the other hand, an absence of local feature

551.   selectivity was found for AM along the RF width axis (magenta and gold traces).

552.   Globally , the comparative analysis between each configuration of Surround-Only stimulation

553.   validates the prediction of the "Dynamic Association Field" hypothesis, a concept defined in our

554.   previous study (29): the collinear configuration promotes input collection along the orientation

555.   preference axis, resulting in the emergence of centripetal axial direction selectivity for high-speed co-

556.   aligned stimuli.

557.

558. ## 2.   Specificity of the interaction between "local" (inducer) and "global" (AM motion) features

559. **2.1. Inducer co-alignment with the RF orientation promotes centripetal synaptic integration along the global motion axis**

560.   The following contextual analysis focuses now on the modulatory effect that a two-stroke

561.   activation of the silent Surround (along each of the cardinal axes) can produce on the response to a

562.   subsequent test Gabor stimulus (the third stroke), flashed at the preferred orientation within the

563.   MDF (right column in Fig. 1B). In addition to stimulations restricted to the Surround, the contextual

564.   impact of the Surround was measured (in an interleaved way) by comparing the response to the full

565.   apparent motion sequence terminating in, or departing from the RF (3-stroke AM, centripetal (CP)

566.   "Surround-then-Center" and centrifugal (CF) "Center-Then-Surround" conditions) to the response to

567.   the test stimulus in the RF ("Center-Only" condition).

568.   The complete set of CP- and CF- configurations, with Gabor inducers either co-aligned with the

569.   motion axis along the RF preferred orientation ("collinear"), or orthogonal to the RF width axis

570.   ("parallel"), was tested in 23 cells. As expected from the study of Surround-Only responses (section

571.   1.3), both centripetal "collinear" and "parallel" centripetal AM flows invading the RF elicited a

572.   significant facilitating effect on the Center test response (WPSR, n=23, 2 conditions: AM

573.   Center+Surround vs. Center-only; $p<0.001$) both at the subthreshold (Vm) and spiking levels. Fig. 4

574.   illustrates the case of a highly selective dependency of contextual control on the flow direction,

575.   emphasizing the functional facilitatory impact of centripetal collinear (CP-ISO) flows (red profile,

576.   upper panel): in addition to the change in response integral, a significant shortening is observed

577.   both at the subthreshold (upper waveforms) and spiking (lower filled PSTHs) levels (compare with

578.   black profiles of the Center-Only response) (WPSR test, n=23, 2 conditions, $p<0.01$). In contrast, a

579.   late change in the PSTW waveform integral, but not of its latency, is observed when the AM flow is

580.   centrifugal (CF) and proceeds from Center to Surround (green and brown, bottom panel of Fig. 4).

581.   -------------

582.   **Fig. 4 : Example of contextual modulation of the Center-response by Centripetal and Centrifugal AM ("cardinal" protocol).**

583.   Example of Center-Surround AM responses, compared to the Center-Only condition. The comparison, in the

584.   same cell, between centripetal (CP: top panel) and centrifugal (CF: bottom panel) motion flows illustrates the

585.   importance of the Surround-then-Center timing in the associative effect. For each configuration, a polar

586.   representation of PSTWs (top) and PSTHs (bottom) is shown for the four AM axes. In each central inset, the

587.   relative positions of the GP inducers are indicated in relation with the receptive field (rectangle, horizontal for

588.   the preferred orientation. The contextual responses are overlaid on the Center-Only stimulation (black). Note

589.   that the contextual latency shortening and the spike discharge rate increase of the Center response are

590.   observed only in the collinear Centripetal condition (top, red). Color code conventions for AM flow (arrow): CP-

591.   ISO-RF: red for centripetal "collinear" along the orientation preference axis; CP-CROSS-RF: gold for centripetal

592.   "parallel" across the RF width axis; CF-ISO-RF: green for centrifugal "collinear" along the RF main axis, CF-

593.   CROSS-RF: brown for centrifugal "parallel" across the RF width axis . See Text for details.

594.   -------------

595.   This example illustrates our main findings at the population level : First, at the Vm level, a stronger

596.   facilitatory impact was found for centripetal (CP) over centrifugal (CF) sequences (WPSR, n=23, 2

597.   conditions, p<0.005). By calculating the ratio between the "significant" response integral (one sided

598.   Z-score, p<0.05) of centripetal vs centrigugal AM sequences, the CP/CF bias in responsiveness had a

599.   median value close to 2.0 (median ± MAD: 1.82 ± 0.42). Second, the centripetal modulatory effect of

600.   the test response was also selective of the motion axis and was stronger for collinear activation along

601.   the RF main axis (ISO-RF) than along the width axis (CROSS-RF) (WPSR, n=23; 2 conditions for each

602.   cell ; p<0.001). By calculating the ratio between the "significant" response integral (one sided Z-

603.   score, p<0.05) of CP AM sequences and the Center-Only test response, a facilitation by the Surround

604.   of the test subthreshold (Vm) response was observed respectively in 95% of occurrences for

605.   "collinear" and 69% for "parallel" conditions. A similar finding was also found at the spiking level,

606.   with respectively 84% of facilitatory effects for "collinear" and 66% for "parallel" conditions. The ISO-

607.   CROSS feature bias in responsiveness (median ± MAD) was respectively equal to 1.45 ± 0.28 for Vm

608.   and 1.36 ± 0.37 for the spiking activity. Finally, collinear centripetal AM sequences was found to be

609.   the only condition which induced a significant positive phase advance compared to the test response

610.   (paired statistics at the Vm level, WPSR test, n=23, 2 conditions, p<0.01).

611.   These differential contextual kinetic changes, modulating both response integral and PSP onset

612.   latency, fit qualitatively with those expected from a simple summation model, recapitulating the

613.   respective timing in the recruitment order of feedforward and lateral inputs : CP-AM sequences tend

614.   to impact the early phase of the response when the stimulation of the Surround precedes that of the

615.   Center, whereas CF-AM sequences tend to affect the late phase of the response when stimulation of

616.   the Center precedes that of the Surround.

617.   2.2. Differential impact of "Near" and "Far" Surround recruitment

618.   In order to better delineate the contribution of the "Far" periphery, we designed a more

619.   conservative stimulation protocol (the "radial" protocol, Fig. 1C) where the Surround stimulation was

620.   allowed only outside the impulse SRF (and not only the MDF, as done previously in the "cardinal"

621.   protocol). The spacing between GP inducers was increased from 120% of the MDF to 100% of the

622.   SRF. The absence of overlap between the nearest Surround stimulus and the SRF was checked for

623.   each of the recorded cells. Stimulation was no longer restricted to the main and cardinal axes but

624.   extended to angular SECTORS surrounding the original cardinal axes by two neighboring (± 30°)

625.   stimulation axes, so as to match the spatial sensitivity of the synaptic association field (29). We also

626.   dropped, from the stimulation set, the AM configurations used in the "cardinal" protocol for which

627.   non-specific component responses were evoked (when co-varying the local inducer orientation and

628.   the motion axis).

629.   This enabled us to focus on our main conditions of interest with GP elements ISO-oriented with

630.   the motion axis always presented along the RF preferred orientation axis (CP-ISO and CF-ISO).

631.   Elements cross-oriented to the motion axis only referred to stimulations along the width axis of each

632.   recorded cell (CP-CROSS). Note here the terminology change in ISO and CROSS conditions - imposed

633.   by the radial symmetry – from that used in the formal ("cardinal") protocol, where collinearity and

634.   cross-orientation the ISO-were defined relative to the RF preferred orientation axis (and not the

635.   motion axis). In addition, in the "radial" protocol, each of the 37 cells recorded was stimulated with

636.   an extended benchmark of contextual 5- to 6-stroke AM flows, richer than in the "cardinal" protocol.

637.   The parametric systematization of the various flows of ISO-axis or CROSS-axis GP sequences is

638.   detailed in each row of Fig. 1C.

639.   Fig. 5 illustrates important spatial differences in the Surround periphery recruitment between

640.   the two protocols ("cardinal" in A; "radial" in B), by comparing the respective time-course profiles of

641.   the intracellular responses, evoked by the sole stimulation of the surround (Surround-Only, dotted

642.   traces) and by the complete AM sequence terminating in the RF center (Surround-then-Center,

643.   continuous traces). The top row in Fig. 5 illustrates the RF maps of the spiking discharge fields (ON

644.   (red) and OFF (blue)) and of the subthreshold SRF (topological union of ON and OFF depolarizing

645.   fields) delimited by a white contour. In order to focus the comparison on the stimulation axes used in

646.   the "cardinal" protocol, we illustrate, for the "radial" protocol cell (B, right panel), only the CP-ISO

647.   and CP-CROSS Sector AM configurations (first and third row, in the left column of Fig. 1C, red and

648.   gold schemes) along respectively the RF main and width axes. A straightforward observation directly

649.   emerges from the comparison of the visual field paving with Surround GPs applied in each protocol :

650.   for the same spatial viewing field size centered on each RF (20°x20°), the two Surround sources of

651.   stimulation are visible for the "cardinal" protocol, while only one - the most proximal (to the RF

652.   center) - appears for the "radial" protocol. The closest GP location in the periphery (D1 position in

653.   Fig. 1A) partially encroaches on the border of the SRF in the "cardinal" protocol, while this is not the

654.   case (by design) for any surround sources in the "radial" protocol.

655.   This difference in the spatial overlap between the most proximal "surround" stimuli and the SRF

656.   accounts for the differences in intracellular response magnitudes observed in each protocol. In the

657.   "cardinal" case (Fig. 5A), the "Near" Surround-Only stimulation evokes sizeable depolarizing

658.   responses, both for CP-ISO and CP-CROSS stimulations (dotted red and gold waveforms). Both types

659.   of CP Surround-then-Center AM flows strongly amplify the evoked PSP amplitude, although the peak

660.   amplitude gain is the largest in the CP-ISO condition. In contrast, the typical behaviour observed in

661.   the "radial" protocol (Fig. 5B) shows a different profile. For a comparable magnitude of the Center-

662.   only response (8.0 mV ("radial") vs. 7.4 mv ("cardinal")), the purely modulatory influence of the CP-

663.   ISO AM flow restricted to the "Far" Surround was generally much weaker (2-3 times) than that seen

664.   for the "cardinal" protocol (here, 3.9 mV against 9.8 mV). In contrast to the CP-ISO condition, the

665.   CP-CROSS Surround-Only stimulation almost did not induce a contextual response (compare the

666.   dotted red and gold traces in the bottom right panel in Fig. 5B). As a consequence, in the "radial"

667.   protocol, the complete CP-ISO AM sequence (terminating in the RF center) is the only configuration

668.   inducing a phase advance and a significant amplification of the early Center-only response.

669.   -----------

670.   **Fig.5 : Differential recruitment of "Near" vs "Far" Surround between the "cardinal" vs. "radial" protocols**

671.   **[A]:** *"cardinal" protocol* (left panel), and **[B]: "radial" protocol** (right panel). Top row, receptive field maps.

672.   Same convention as in Fig. 3A. Only the Surround GPs on the RF main (horizontal) and width (vertical) axis

673.   (common to both protocols) are represented within the same viewing field (20°x20°). Note the encroachment

674.   of the most proximal GPs in the "cardinal" protocol over the SRF (white contour), whereas no spatial overlap is

675.   seen in the "radial" protocol. Bottom rows, the CP-ISO (red) and CP-CROSS (gold waveforms) are overlaid for

676.   the "Surround-Only" (dotted) and the "Surround-then-Center" (continuous trace) conditions and compared to

677.   the "Center-Only" test condition (black). Insets show respectively an expanded version of the PSPs onset on a

678.   30 ms time window (dotted grey rectangle). For each protocol, arrows indicate the change in PSP slope due to

679.   the feedforward synaptic drive triggered by the Center test stimulus in the D0 location. Note that the CP-ISO

680.   (red) and CP-CROSS (gold) response latencies are the same for the "cardinal" protocol, suggestive of a common

681.   non-specific input. In contrast, the phase advance is seen only for the CP-ISO configuration in the "radial"

682.   protocol. See Text for details.

683.   -----------

684.   A mechanistic difference between the two protocols is further attested by the comparison of

685.   the rising slopes of the time-profiles of the AM responses with that of the feedforward response

686.   (Center-Only). In the "cardinal" example cell (bottom left panel in Fig. 5A), both the CP-ISO Surround-

687.   Only and the CP-ISO Surround-then-Center evoked PSPs have the same initial rising slope as the

688.   Center-Only response. The latencies, when corrected for the 16 ms ISI interval, are the same in D1

689.   and D0, thus suggestive of a direct synaptic feedforward input signature even for the D1 position

690.   (thus overlapping with the RF). Furthermore, both the CP-ISO and CP-CROSS synaptic responses start

691.   to rise simultaneously in the "cardinal" case, which indicates that the "Near" stimulated periphery

692.   recruits feedforward synaptic drive (common to D0 and D1). In contrast, in the case of the "radial"

693.   example (bottom right inset in Fig. 5B), the early phase of the CP-CROSS waveform is

694.   indistinguishable from that of the Center-Only (gold and black) waveform. The initial rising slope of

695.   the CP-ISO response (red) occurs much earlier (15-20 ms ahead) and is initially much slower than the

696.   fast rise seen for the Center-Only response (black). These latter features are suggestive of a graded

697.   spatio-temporal diffusion process compatible with horizontal activity propagation. This qualitative

698.   comparison clearly emphasizes that, by preventing the closest surround-location from encroaching

699.   on the SRF border, the non-specific tonic component of the boosting of cortical responsiveness seen

700.   in the "cardinal" protocol is no longer recruited in the "radial" protocol.

701.   Remarkably enough, on the whole population, the "radial" protocol led to a drastic decrease of

702.   the proportion of cells showing "Surround-Only" responses (32 % ("radial": point by point

703.   permutation test regarding spontaneous activity for a minimum of 15 consecutive ms, 10^4

704.   repetitions, p<0.01)), vs. 100 % ("cardinal": Z-test above background activity, p < 0.05)). Compared to

705.   the "cardinal" protocol, it also produced weaker Surround responses, that never reached the peak

706.   amplitude of the Center-Only test response, and, accordingly, to a smaller – although significant –

707.   facilitatory modulation of the Center-response. We conclude that, in contrast to the "cardinal"

708.   protocol, the "radial" stimulation of the Surround no longer recruits as efficiently the "Near"

709.   periphery, which seems to integrate both lateral and feedforward input. Consequently, the "radial"

710.   protocol appears best tailored to detect the selective contribution of purely lateral recruitment of

711.   the "Far" periphery.

712.   ## 2.3. Anticipatory "filling-in" responses ("radial" protocol)

713.   To highlight the emergence of a lateral anticipatory wave due the sole recruitment of the distal

714.   periphery (Surround-Only wave) and quantify its functional role in the contextual gain control of the

715. response evoked by the feedforward drive, we determined - in the "radial" protocol - the individual

716. significance of each cell response to the Surround-Only condition, using a point-by-point

717. randomization test (10^4 repetitions, p < 0.01). The largest proportion of cells showing significant

718. Surround-Only responses was found for the CP-ISO condition (12 cells (32% of the population)). A

719. single cell example is illustrated in the left column of Fig. 6. A complete CP-ISO sequence invading

720. the RF Center reveals a clear boosting effect (compare continuous red (AM) and black (Center-Only)

721. traces, top and bottom panels). In spite of the fact that the closest surround location was sparing the

722. SN-mapped SRF, a sizeable significant change in membrane potential was still observed in the AM

723. sequence restricted to the surround, from D5 to D1 locations, in the absence of D0 stimulation

724. (bottom left graph in Fig. 6, dashed red trace). For this cell, the combined dynamic recruitment of

725. the GP sources along the AM motion axis shows a progressive build-up signal which anticipates, by

726. ten milliseconds, the onset of the feedforward signal evoked by the Center-stimulus. In contrast, the

727. amplitude of the linear predictor waveform - based on individual static responses to GPs in isolation

728. – remains close to the resting state, even at the onset of the Center- Only response (green curve,

729. bottom left panel in Figure 6). Note that the Surround-Only trace (in red) trespasses the upper limit

730. of the confidence interval (shaded green envelope) tested by point by point permutation between

731. the Surround Linear Predictor (SLP) and the actually recorded response to the Surround-Only

732. sequence (10^4 repetitions, (p<0.01)). Thus, in this particular single cell example, the AM sequence

733. recruits a non-linear summation process of subliminal surround sources. The significant deviation

734. from the linear prediction attests for the role of spatio-temporal synergy in maximizing the

735. functional impact of horizontal propagation.

736. ------------

737. **Fig.6 : "Filling-in" responses evoked by Surround-Only AM ("radial" protocol).**

738. The analysis here is restricted to cells where Surround-Only CP-ISO stimulation evoked a significant response

739. (n=12, see Text for criteria). Trace color code : black for Center-only; red for Surround-then-Center, dashed red

740. for Surround-Only, dotted green for Surround Linear Predictor (SLP). The left middle horizontal insets

741.   respectively show the chronograms of the stroke-by-stroke stimulation sequences for Center-Only (1 stroke,

742.   filled black box, upper line), Surround-Then-Center (6 strokes, filled red, middle line) and Surround-Only (5

743.   strokes, filled pale red, bottom line) protocols. Empty boxes indicate omitted stimuli. The time onset of each

744.   GP stroke is labeled by a thin dotted vertical line. In this figure and the following ones, the blue dot and vertical

745.   dotted blue bar on the y-axis indicate the "threshold" amplitude change of the Center-Only control curve from

746.   rest, above which statistical significance of the response is reached (p<0.01) for each individual cells. Their

747.   abscissa serves as a reference for the temporal realignment of the different contextual responses. **Left** : Single

748.   cell example. **Top left**: The confidence intervals (permutation test; 10^4 repetitions, p<0.01) for Center-Only and

749.   Surround-then-Center compared to Blank are represented respectively by gray and pale red envelopes. The

750.   complete CP-ISO AM sequence (D5 to D0) evokes a significant facilitation of the Center-Only Vm response.

751.   **Bottom Left** : The red and black curves represent the response averages across trials (shaded area: ± SEM).

752.   When omitting the D0 stroke, the recruitment by the CP-ISO AM flow, although limited to the silent surround

753.   (D5 to D1), still induces a significant depolarizing activation (Surround-Only : dashed red). The temporal profile

754.   of the lateral wave of activity matches the "predicted" invasion of the RF Center (black), had it been stimulated.

755.   The build-up of the Surround response during AM departs significantly from the sum of the static responses

756.   evoked by each distal GP stroke in isolation (Surround linear predictor (SLP) : dashed green, confidence interval

757.   of a significant difference between SLP (predicted) and Surround-Only (observed) waveform, p<0.05).

758.   **Right, top panel** : Average response profiles for Surround-then-Center, Center-Only and Surround-Only

759.   conditions (n=12). **Right, bottom panel** : the CP-ISO Surround-Only response (mean: dotted red ± SEM: shaded

760.   area) is compared (with a different ordinate scale than in the top right panel) with the "expectation" (dashed

761.   black trace) obtained by subtracting the Center-Only (black trace, top right) from the complete CP-ISO AM

762.   response (red trace, Top right).

763.   ----------------

764.   In the subpopulation of cells showing significant CP-ISO Surround-Only responses (point by point

765.   permutation test, 10^4 repetitions, p < 0.01), paired comparison between responses to Surround-only

766.   sequences and their Surround linear predictor revealed that 6/12 cells (50% of significant cases; 16%

767.   of the entire population (n=37)) showed significant anticipatory responses in the CP-ISO condition

768.   larger than that predicted by the linear summation of individual GP responses. For the remaining

769.   cells, observed responses were indistinguishable from their Surround linear predictor. Note also that

770.   the slight difference between the dotted black and red curves in the Figure 6 right bottom panel –

771.   potentially indicative of a sublinear center/surround summation - could simply be due to a shift in

772.   reversal potential during depolarization.

773.   When pooling altogether Surround-Only significant cases and realigning individual cell responses

774.   on the onset time of their respective Center-Only responses, the population average shows a mean

775.   depolarizing wave riding clearly above the noise level. This wave, depicted in the top right panel of

776.   Fig. 6, peaks at an average value corresponding to a third of the Center-Only test response peak

777.   amplitude. It corresponds to absolute values ranging between 1 to 4 mV (0.35 of normalized

778.   amplitude for the dotted red average trace vs 0.85 for the black trace). Remarkably, for the

779.   optimized AM speed, the Surround-Only average response reaches its maximum at the time of

780.   expectation of the Center-Only response. These findings strongly suggest a causal relationship

781.   between purely contextual lateral presynaptic information and its postsynaptic integration during

782.   complete apparent motion terminating in the RF center. This observation does not reflect simply a

783.   trivial non-specific summation effect : i) the gain control effect is non-linear in half of the cases

784.   (Figure 6, left); ii) the process is shown to be selective not only of the timing of the Gabor stroke

785.   sequence (optimal speed) but also of the local feature (orientation) of the Gabor inducer (CP-ISO). In

786.   spite of similar timing sequence of presentation of Gabor patches in CP-ISO and CP-CROSS

787.   configurations, the two contextual responses are actually different.

788.

789. **3. Dependency of contextual cortical control on the spatio-temporal coherence of the AM flow**

790.   We have attempted to systematically identify and quantify the key spatial and temporal factors of

791.   the CP-ISO AM sequence which condition the effectiveness of the contextual control of the cortical

792.   gain. For this purpose, we compared for each of the 37 cells, the CP-ISO configuration with AM

793.   sequences in which the spatio-temporal structure of the GP presentation was altered (Fig. 1C). The

794.   statistical significance of the measures extracted from the mean Vm responses of individual cells (top

795.   panel in Fig. 7) was assessed by comparing each AM condition to the same reference (Center-Only).

796.   Two selection criteria were used independently, based on a significant change, either i) in the onset

797.   latency, or ii) change in response strength (integral). In addition, we also used a less stringent

798.   requirement, combining one or the other or both criteria (randomization test, 10^4 repetitions, p <

799.   0.05). Both congruent and complementary trends were found across the three statistical criteria we

800.   used.

801.   ------------

802.   **Fig. 7. Statistical significance of contextual response changes**

803.   Color code, same as in Fig. 1C, schematized by icons in the middle column. SECTOR configuration : red for CP-

804.   ISO, green for CF-ISO, gold for CP-CROSS; dark blue for RND; FULL configuration : pale blue for RND-ISO.

805.   **A :** Top left panel, ranked amplitude distributions (n=37) of changes in latency (ms, left column) and in

806.   response integral value (normalized ratio, right column) of subthreshold responses (Vm), for each SECTOR AM

807.   condition of interest (first three rows). The fourth row corresponds to the FULL-RND condition (see Text for

808.   justification). Ordinates : latency shortening and integral response increase are plotted upward. Abscissa : cell

809.   rank (1 to 37) ordered independently for each AM condition. Filled bars represent "significant" individual cases

810.   (one-sided permutation test, 10^4 repetitions, p< 0.05). Ordinates : left column, latency change ("Δ Latency")

811.   measured at half-height of the Center-Only peak response; right column, change in the integral value of

812.   "significant" depolarizing responses (± 3σ from mean V_{rest}), integrated to the point of return to baseline of the

813.   Center-only test control. The integral value change ("Δ Response") is expressed as a normalized ratio relative to

814.   the Center-Only test response.

815.   **B :** Bottom left panel, each of the three histograms represents the proportions of significant cases for each AM

816.   flow condition. Left, "Δ latency" criteria. Right: "Δ Response" criteria. Center inset (BOTH): all cases showing

817.   either a significant latency advance, or response integral increase, or both.

818.   **C,** right panel : Bihistogram of the Δ(response_integral) in ordinate vs. Δ(latency_change) in abscissa, averaged

819.   across conditions. Numbers and SEM correspond to values calculated across the whole population (upper plot)

820.   or restricted to the pool of "significant" cells (lower plot). Note that the global separability between CP-ISO and

821.   all other test conditions increases when restricting the pooling to the "significant" cells.

822.   -------------

823.   The impact of each AM configuration is expressed in figure 7A by ranked amplitude distributions

824.   of latency changes (left column) and response integral changes (right column), relative to the Center-

825.   Only condition. The proportions of significant facilitatory effects correspond to the number of

826.   occurrences where significant <u>positive</u> modulations relative to this common reference were

827.   observed. Paired comparisons between each condition (condition of interest and other controls)

828.   were also done but not illustrated for sake of clarity. The left histogram in Fig. 7B represents the

829.   proportion of "significant" latency advance cases (upwards filled bars in the left column of Fig. 7A). It

830.   is three times larger for the CP-ISO condition (30%, red column) than for the other conditions. The

831.   right histogram in Fig. 7B represents the proportion of "significant" response integral changes. The

832.   facilitation effect (positive Δ Integral) is less selective across conditions, but remains specific of the

833.   collinear configuration, since most significant cases are found in both CP-ISO and CF-ISO conditions

834.   (respectively red and green). The middle histogram inset in Fig.7B shows the impact of pooling both

835.   change criteria in a non-exclusive way (logical "OR") : the proportion of significant facilitation cases

836.   (latency shortening OR Response Integral increase) peaks for the CP-ISO condition (41%, red

837.   histogram bar in the bottom center inset), while remaining two times larger than for the CF-ISO

838.   condition (19%, green bar) and three times larger than the other conditions (11%, gold bar for CP-

839.   CROSS; 14%, blue bar for RND-ISO). These different analyses show that the CP-ISO AM flow is by far

840.   the most effective configuration to promote the visibility of Surround-Center interactions.

841.   For the CF-ISO condition, no consistent latency change was observed at the PSP population level

842.   (n=37), since the rising phase of the CF-induced Vm response, as expected, was generally

843.   indistinguishable from the Center-Only response. Only 8% of the cells showed a significant latency

844.   advance of a modest magnitude in the CF-ISO condition compared to the 30% found in the CP-ISO

845.   condition (green bar vs red in the left histogram, Fig. 7B). Conversely, contextual facilitation of the

846.   Center response integral was present in a slightly larger proportion of cells (14%, green bar in the

847.   right histogram of Fig. 7B). Those observations concord with those of the "cardinal" protocol, since

848.   the rising phase of the CF-induced response, as expected, was indistinguishable from the Center-Only

849.   response. On the other hand, the increase in response integral value concords with a lasting

850.   depolarization following the center-only test peak response, as more and more Gabor patches

851.   flashed at increasing eccentricity from the RF center evoke a series of lagged desynchronized lateral

852.   inputs of exponentially decreasing amplitude. The centrifugal control demonstrates that the spatial

853.   coalignment of GPs is not enough in itself to induce the observed boosting effect. Some form of

854.   spatio-temporal ordering is required such that an anticipatory flow from the periphery is generated,

855.   which, *in fine*, will boost sensory responsiveness at the time when the test stimulus hits the RF

856.   Center.

857.   Since it is well established in most V1 cells that moving cross-oriented stimuli across the

858.   orientation preference axis of the recorded RF are ineffective in firing the cell, we focused our study

859.   of CP-CROSS AM sequences along an angular sector restricted around the width axis (Fig. 1C: left

860.   column "SECTOR"; gold condition). This choice however implies in the SECTOR configuration that the

861.   analysis relies on the comparison of response flows (CP-ISO vs. CP-CROSS) sweeping across different

862.   surround subregions, while maintaining the orientation of all GPs the same. Nevertheless, the

863.   absence of an orientation-selective component for centripetal flows of CROSS-oriented GPs along the

864.   main axis of the RF (CP-CROSS_main-axis, cyan traces in Fig. 3) justifies this simplification choice. In

865.   spite of these subtle protocol differences, only 11% of the cells showed a significant positive latency

866.   advance to the CP-CROSS stimulation, while none displayed any significant positive change in the

867.   response integral, when compared to the Center-Only test response (bottom panel, respectively left

868.   and right gold histogram bars in Fig. 7). Overall, the weak changes in latency advance or amplification

869.   of the Center-Only response found in individual cells during CP-CROSS AM flows was washed out

870.   completely when averaging across cells (RIGHT panel in Fig. 8, compare gold and black traces). A

871.   similar conclusion is reached whether the surround was partially (SECTOR) or uniformly (FULL)

872.   recruited: in both conditions (SECTOR and FULL), the contextual mean CP-CROSS responses were

873.   almost indistinguishable from the Center-Only test response.

874.   -------------

875.   **Fig. 8 : Contextual Gain Control induced by centripetal-ISO AM flow ("radial" protocol)**

876.   **Top** row, subthreshold response (Vm); **Bottom** row, firing rate response (spikes/s). The first three columns,

877.   from the left illustrate the boosting effect produced by a CP-ISO AM sequence (red curve, Surround-then-

878.   Center) when compared to the response to the test GP inducer flashed in the RF Center (black curve, Center-

879.   Only). The local orientation of the GP inducers in the Surround and the global AM axis are co-aligned with the

880.   RF orientation preference. The last (right) column focuses on contextual dependency of the effect. **From left to**

881.   **right, first column :** Single cell example, where the vertical dotted gray lines indicate the respective timing of

882.   each peripheral stoke stimulation at successive eccentricities (from D5 to D1), whereas the continuous vertical

883.   gray line indicates the onset of the "Center-Only" test stimulation (D0). The blue dot and corresponding dotted

884.   line give the latency of the first point in time where the Center-Only response departs significantly from the

885.   resting state (p < 0.01). **Second column** from the left, population averages for the CP-ISO (red) and the Center-

886.   Only (black) conditions. Averaging is done in two steps, first across trials for each cell for the same context

887.   condition, then across cells after realigning individual mean waveforms around a common onset latency, that

888.   of their Centre-Only response ("0" mark of the "Relative Time" abscissa, blue points in left panels). Response

889.   amplitudes are normalized relatively to the peak of the "Center-Only" response. The **third column** illustrates

890.   the population average profile of individual cells showing either significant positive latency advance, response

891.   integral increase, or both, compared to the center-only test control (permutation test, 10^4 repetitions, p<0.05).

892.   The difference in rising phases and onset latencies between the CP-ISO (red) and Center-Only (black) responses

893.   are shown on an expanded time basis. The mean ± SEM value envelopes are respectively illustrated by a

894.   continuous curve and a shaded area. **Right column**, the statistical significance analysis is extended to cells

895.   which showed significant "Surround-only" responses (subthreshold: n=20; spiking: n=10). Color code, same as

896.   in Fig. 1: SECTOR configuration, red for CP-ISO, green for CF-ISO, gold for CP-CROSS. FULL configuration, blue

897.   for CP-RND-ISO. The contextual averages show that CP-ISO (red) is the only condition where a change in onset

898.   latency is observed, both at the Vm (top) and spiking levels.

899.   ---------------

900.   In a third control condition, the same retinal SECTOR space as in our CP-ISO condition of interest

901.   was stimulated with the same GPs (i.e., with the same number, features, and stimulus energy

902.   distribution) but the coherence of the flows was pseudo-randomized both in space and time. Such

903.   RND-ISO condition induced a slight facilitation effect on average, although significantly smaller than

904.   the one seen in the CP-ISO condition. We explored the possibility that this partial effect could be of

905.   residual nature and come from an insufficient level of randomization achievable in the SECTOR

906.   stimulation. Indeed, the number of visited nodes (i.e. GPs) is rather low (n=26), with the additional

907.   constraint that the last Gabor patch in the sequence had to be the one flashed in the Center. Note

908.   that, in contrast to coherent-AM sequences, GPs flashed during each step of the RND-ISO AM

909.   sequence could belong to any ring of eccentricity, thereby increasing at each step of the AM

910.   sequence the probability to find D1- (or D2-) to center stimulations (see Methods). Close

911.   examination of individual cell recordings showed that the slight facilitation most likely reflected the

912.   fact that in some RND-ISO trials, the penultimate GPs were presented occasionally by chance at the

913.   D1, or even D2 eccentricity (Fig. 1A), just one stroke ahead of the RF center stimulation. We

914.   concluded that the Sector RND-ISO sequence often recruited facilitating two-stroke AM Surround-

915.   Center pairs whose effect has already been reported (see fig. 7 in (29)).

916.   To dilute the remnant impact of proximal-to-center interactions seen in the RND SECTOR

917.   condition, we used, as a more randomized control, the FULL stimulation configuration in which the

918.   probability to find such co-aligned D1 (or D2) GP followed by the RF center stimulation during the

919.   entire stimulation protocol was lowered. This was achieved by doubling the number of possible

920.   peripheral node locations (from n=26 to n=52), thereby increasing the level of randomization of the

921.   AM sequence. In the RND FULL mode, the stimulation was not restricted to retinal sectors on both

922.   sides of the RF but was extended isotropically to its entire Surround (bottom right blue icon in Fig.

923.   1C). The bi-histograms in Fig. 7C illustrate the observation that the full randomization (pale blue for

924.   FULL-RND) produces changes in latency and response strength much weaker than the partial

925.   randomization limited to the SECTOR configuration (dark blue). However, still a limited fraction of

926.   the cells response to the FULL RND stimulation displayed a slight facilitatory effect (11% for latency

927.   change and 3% for response gain) compared to the Center-Only test response (blue columns, in

928.   respectively the left and right histogram bars in the bottom panel of Fig. 7B). This mild effect

929.   disappears when averaging across cells (compare average blue and black waveforms in the right

930.   column in Fig. 8). We conclude that the contextual boosting of the cortical response was largely

931.   washed out by the reduced probability of observing collinear proximal activation in the FULL RND

932.   sequences, resulting from the more extensive randomization of the flow's spatio-temporal

933.   coherence than in the SECTOR RND condition.

934.   Let us focus more in depth on the CP-ISO condition. The left column in Fig. 8 illustrates the

935.   typical contextual behaviour observed for Vm and spiking responses at the single cell level for the CP-

936.   ISO configuration in the "radial" protocol, when compared to the Center-Only condition : the AM CP-

937.   ISO sequence produces an anticipatory depolarizing wave that shortens the onset latencies of both

938.   the rising phases of the PSP (upper row) and the spiking response (bottom row), by respectively 14.7

939.   and 16.3 ms. For that particular example, both a progressive build-up of a subthreshold

940.   depolarization and an anticipatory firing response are observed at early latencies, but this ramp-up

941.   feature was not systematically observed in all cells. At the population level, we found that centripetal

942.   flows composed of Gabor elements co-aligned with the global motion path along the RF main axis

943.   (CP-ISO) resulted in an overall boosting of the neural response compared to that evoked by the test

944.   Center stimulus alone.

945.   This finding is illustrated in the second column (from the left) of Fig. 8. Note here that the PSTW

946.   and PSTHs realignments (see Fig. 6 legend) are done separately for the averaging, since a few

947.   millisecond integration lag is found between the absolute measurements of significant onset times

948.   for Vm and spikes for the Center-Only responses (see the causal shift in the vertical dotted blue lines

949.   in the single cell example). On average (n=37), the CP-ISO stimulation, in the SECTOR condition,

950.   resulted in an overall latency shortening and a slight amplification of the depolarizing subthreshold

951.   response envelope (Vm, upper row). It also led to a reduction in latency of the spiking discharge

952.   (bottom panel) in cells where spiking activity evoked in the Center-Only condition was significant

953.   (permutation test, 10^4 repetitions, p<0.05; n=22).

954.   In order to refine the statistical significance of the effect, we first restricted the averaging to all

955.   the cells for which, individually, the CP-ISO sequence presentation led to a significant change in the

956.   response latency and/or response integral (when compared to center-only response), whatever its

957.   sign (Figure 7A). Note here that even if the dominating effect induced by the CP-ISO stimulation was

958.   an overall clear bias toward a latency reduction, it occasionally led to a latency lengthening (negative

959.   Δ latency as illustrated in Figure 7A). We observed more globally that 1) the proportion and

960.   amplitude of positive Δ-latency and Δ-response integral was much larger than the proportion of

961.   negative ones for the CP-ISO condition of interest and 2) the proportion of significant positive

962.   modulatory effects was larger in the CP-ISO condition than in all other conditions. This led us to focus

963.   on (one-sided) significant positive contextual subthreshold (Vm) changes in onset latencies and/or

964.   response integral values. For this subpopulation, the CP-ISO condition produced significantly

965.   facilitated responses in 41 % of the cells (permutation test, 10^4 repetitions, p<0.05, for 15 cells out of

966.   37). The third column in Fig.8 illustrates the selected population average profile. Note that the

967.   quenching around the "0" time abscissa, seen for the Center-Only responses, results from the

968.   realignment process applied before averaging (see Methods).

969.   To further demonstrate the contextual boosting specificity concomitantly at the synaptic and

970.   the firing levels, we tried to maximize the population size. For this last analysis, we added - to the

971.   pool of CP-ISO "significant" cells - the contingent of cells for which significant responses could be

972.   elicited by an AM flow stimulation limited to the Surround (permutation test, 10^4 repetitions, p <

973.   0.01; see Methods). This pooling increased the number of significant cases from 15 to 20 cells for

974.   subthreshold analysis, and from 6 to 10 for spiking analysis (one of the additional cells was not

975.   spiking). Once again, in congruence with the statistical analysis of Fig. 7, the largest proportion of

976.   significant Surround-Only responses was found for the CP-ISO condition (CP-ISO: 12/37 cells, 32%;

977.   CF-ISO: 22%; CP-CROSS: 8%; RND-FULL: 11%). The right column in Fig. 8 shows that the phase

978.   advance effect is selective of the CP-ISO condition : the response onset of the mean Vm waveform

979.   (n=20) was shortened by 8 ms and the PSP integral increased by 31%. A similar contextual phase shift

980.   is observed at the spiking level (bottom).

981.   Altogether, our intracellular results demonstrate that a build-up of anticipatory synaptic activity

982.   is observed by sequentially recruiting local inducers co-aligned with the RF orientation axis in a

983.   centripetal way. The efficiency with which the closest surround location is recruited impacts the

984.   strength of the contextual modulation. In particular, the "Near" periphery recruitment of the

985.   "cardinal" protocol points to a bias for CP-ISO AM flows along the RF preferred orientation axis,

986.   superimposed over a non-specific facilitation component observed, at both Vm and Spiking levels for

987.   all contextual stimulation conditions. Conversely, the more precise recruitment of the "Far"

988.   Surround, achieved in the "radial" protocol, points to the selective control of the cortical gain by the

989.   CP-ISO context. This specificity truly highlights a spatio-temporal requirement in Surround

990.   recruitment in order to boost sensory responsiveness. This further emphasizes the critical need of

991.   spatial and temporal synergy between each distal surround source synaptic impact with the synaptic

992.   feedforward volley corresponding to the Center-Only drive (Fig. 1A).

993.

994. ## 4.   Timing specificity in Predictive "filling-in" responses and Surround-then-Center interactions

995. ### 4.1. Synaptic mechanisms involved in predictive filling-in responses ("radial" protocol)

996.   The central issue addressed here is to delineate how the temporal specificity of each purely

997.   contextual surround component accounts for the contextual differences in cortical gain control when

998.   lateral input is finally combined with the feedforward flow originating from the RF center. This can be

999.   studied only in cells which displayed significant responses to the sole stimulation of the Surround

1000.   (point by point permutation test, 10^4 repetitions, p<0.01). In these 12 cells, we first checked that the

1001.   global contextual impact of surround and center interaction is the same as in the global population.

1002.   Fig. 9 allows the straightforward comparison of the impact of the CP-ISO AM SECTOR configuration

1003.   (red trace) with the three other flow configurations (light green for CF-ISO, gold for CP-CROSS and

1004.   blue for FULL RND-ISO). The top left panel confirms the leftward shift advance of the Vm response by

1005.   about 7 ms (top left panel, Fig. 9), selective of the CP-IS0 condition. Note that the population size is

1006.   too small to warrant further statistical analysis. At the spiking level, peak values are indistinguishable,

1007.   but a clear 15 ms advance only appears in spike initiation of the CP-ISO condition.

1008.   ------------

1009.   **Fig. 9 : Mechanistic analysis of the "Filling-in" contribution to contextual gain control.**

1010.   **Left,** contextual gain control analysis, restricted to the population of cells (n=12) showing significant "filling-in"

1011.   responses. Mean responses, averaged for each AM flow pattern (CP-ISO (red), CF-ISO (light green), CP-CROSS

1012.   (gold) and RND-ISO (blue)) are represented at the Vm (top panel) and spiking levels (bottom) with their

1013.   (shaded) S.E.M. envelopes. Same color code as in Fig. 1C. The selective latency advance and amplification of

1014.   the subthreshold response induced specifically by the CP-ISO condition (top, n=12) translate at the spiking level

1015.   in cells displaying spiking activity (bottom, n=7). **Right**, comparison of the contextual time course of Surround-

1016.   Only responses with the predicted contribution of the Surround according to an additive model. Top, observed

1017.   traces (n=12); Bottom, linear predictors of the Surround-Only responses obtained by subtracting the "Center-

1018.   Only" trace from the "Surround-Then-Center" response. The predicted profiles were calculated independently

1019.   for each contextual condition (same color code). The averaging process was extended to all cells showing a

1020.   significant contextual change in latency and/or response integral (n=15, see Text for detailed criteria). The

1021.   expected occurrence of the response evoked by the omitted Center stimulus is indicated by a grey shaded

1022.   temporal window. The time abscissa of the peak of the "Center-Only" stimulus response is indicated by a thin

1023.   vertical bar.

1024.   --------------

1025.   Let us focus now on the timing specificity of the Surround-Only component as a function of its

1026.   mode of spatio-temporal recruitment. An elementary mechanistic approach is to compare,

1027.   independently for each AM configuration (CP-ISO, CF-ISO, CROSS-ISO, RND), the temporal profile of

1028.   the average Surround-only response (top right panel) to the linear prediction ("Surround-then-

1029.   Center" minus "Center-Only", bottom right panel)) obtained from the 15 cells showing significant

1030.   Center-Surround modulation (top panel, third column, Figure 8). The detailed comparison of the

1031.   relative time-course of "filling-in" subthreshold responses (Fig. 9, top right panel) shows that the

1032.   synaptic (Vm) contribution of the "silent" periphery is the largest for the CP-ISO condition (red).

1033.   Most remarkably, the Surround-only component is temporally highly selective and peaks precisely at

1034.   the expected time of occurrence of the Center-Only response onset, had the RF Center been

1035.   stimulated ("zero" relative time; see the dotted red trace in top right panel of Fig. 6).

1036.   This, of course, does not mean that the other contextual flows are not effective in modifying the

1037.   subthreshold baseline, such as the random (RND-FULL) configuration, which shows a depolarizing

1038.   plateau (blue trace) preceding the onset of the expected Center-Only response. This initial bump is

1039.   compatible with the cumulative invasion (across randomized strokes) of lateral activity and

1040.   corresponds to a non-null probability of stimulation by chance of the proximal periphery of the

1041.   subthreshold SRF. However, in contrast to the CP-ISO AM sequence, it collapses towards the baseline

1042.   before the expected onset of the Center-Only response.

1043.   For the CF-ISO condition (green trace, top right panel in Fig. 9), a mirror activation pattern

1044.   regarding the Center-Only onset response is observed: starting from rest, the transient CF-ISO

1045.   Surround-Only contribution growth appears only after a delay (30-35 ms), slightly after the peak of

1046.   the Center-Only response had the test stimulus been presented, before displaying a progressive

1047.   depolarization decay lasting up to 250 ms. This decay phase is interpreted as indicative of distal input

1048.   weakening and desynchronization, the further away from the RF center (as expected from the

1049.   stimulus design).

1050.   In contrast, the CP-CROSS AM sequence synaptic integration is almost ineffective (gold traces). The

1051.   magnitude of its peak amplitude is several folds lower than that observed in the "cardinal" protocol,

1052.   reaching a peak value of roughly 10% of the Center-Only peak response : compare gold traces in Fig.

1053.   5A, where the CP-CROSS peak magnitude of the surround-only response reaches 4.4 mV, hence 59%

1054.   of the 7.4 mV center-only control peak response in the cardinal protocol vs the 1.3 mV of the

1055.   Surround-Only CP-CROSS response, hence 16 % of the 8 mV of the center-only peak response. This

1056.   argues once again for a "non-specific" component of the response, best seen in the "cardinal"

1057.   protocol, that crucially depends on whether the most proximal peripheral Gabor patch encroaches

1058.   on the SRF border or not.

1059.   An important point in this analysis is that predictors of the Surround modulatory effect, obtained

1060.   by subtracting the feedforward (FF) response to the contextual response ("Surround-then-Center"

1061.   *minus* "Center-Only", shown in the bottom right panel of Figs. 6 and 9), are computed even in cells

1062.   which did not show significant AM "Surround-Only" responses. In spite of the fact that the two

1063.   populations used for averaging differ, a striking similarity is apparent in the overall time-courses of

1064.   recorded (top) and predicted (bottom) waveforms (right panel, Fig. 9). This isomorphism globally

1065.   supports qualitatively a linear (additive) synaptic integration of the dynamic AM Surround lateral

1066.   contribution and the flashed Center drive..

1067.   In summary, the CP-ISO condition is the only stimulation context where the temporal profile of

1068.   the Surround-Only contribution matches precisely the time-course required to interact with the

1069.   feed-forward activation window (shaded area, bottom right panel in Fig. 9) and define an effective

1070.   modulatory "spiking opportunity window". In contrast, in all other flow conditions, Surround-Only

1071.   contribution is out of sync with the feedforward drive and cannot exert mechanistically some form of

1072.   additive gain control of the test feedforward response. Crucially, the CP-ISO exclusivity of this precise

1073.   time-localized concentration in depolarizing activity power of purely contextual information actually

1074.   explains the selectivity of the contextual gain during complete sequences terminating in the RF

1075.   center.

1076.   Our data clearly support the selective dependency of a Surround-only activation wave on both

1077.   spatial and temporal features of the AM stimulation. Co-alignment of local features in the surround

1078.   with the preferred orientation of the test RF and appropriate timing in the sequential surround

1079.   recruitment are both needed. If both conditions are fulfilled, the diffusion waveform, responsible for

1080.   generating subthreshold synaptic responses "at the right place, at the right time", predicts the

1081.   spatio-temporal profile of the next stimulus to come (in the RF center). We therefore interpret

1082.   those waves as neuronal correlates of a "filling-in" process, indicative of an anticipatory activity

1083.   build-up in the cortical retinotopic region for which the "expected" visual feedforward synaptic

1084.   volley is missing.

1085. **4.2. Dependency of the spiking latency advance on the temporal phase between the Feedforward and Horizontal input waves ("cardinal" protocol)**

1086.   The timing constraints of our protocols have a dual source: i) the synaptic activation from the

1087.   surround displays a temporally constrained profile in latency basin (Fig. 2), and ii) the order of the

1088.   sequential stimulation (centripetal & centrifugal) in the Surround (Fig. 4). In order to better quantify

1089.   these issues, the onset latencies of the postsynaptic potentials evoked independently in the Center-

1090.   Only and Surround-Only conditions were measured in the "cardinal" protocol and temporally

1091.   realigned on a common reference, defined as the time at which the Center-stimulus should have

1092.   been shown in the Surround-only 2-stroke AM sequence. The subtraction between the two input

1093.   onset latencies will be referred in the rest of the text as the **phase** between feedforward and

1094.   horizontal inputs. In spite of its weaker specificity, since the "cardinal" protocol almost systematically

1095.   induced significant responses for sequences limited to the surround and strong tonic responses at

1096.   both sub- and suprathreshold levels during complete sequences stimulating the center, we used the

1097.   measured onset latencies of this protocol to maximize the number of cells for this analysis. This

1098.   calculus was however restricted to instances where both "Surround-Only" and "Surround-Then-

1099.   Center" responses were tested for each condition, in the same cells. The interest of such

1100.   quantification is to give an objective measure which allows the direct comparison of the effects of

1101.   different associative sequences realized for a given cell as well as across various cells (Fig. 10).

1102.   ---------------

1103.   **Fig. 10 : Causal impact of the temporal phase between feedforward and horizontal input on the spiking latency change ("cardinal" protocol)**

1104.   **Left** panel : Pooled data from the "radial" (red), and "cardinal" (orange) dynamic AM protocols (this study).

1105.   Note that most apparent speed of horizontal propagation (ASHP) values, extracted from intracellular latency

1106.   basin slopes with relative eccentricity, range between 0.05 and 0.60 mm/ms (with a peak between 0.10 and

1107.   0.30 mm/ms). Mirror histogram (pointing downwards) : pooled data using flashed long bars (1D-mapping, blue),

1108.   sparse noise mapping (green) from (26), or Gabor patches (light green). **Right** panel: For each individual cell

1109.   recorded in the "cardinal" protocol, the latencies of Center-Only and Surround-Only Vm responses are

1110.   measured and subtracted, defining the temporal phase between the "horizontal" and "feedforward (FF)"

1111.   inputs. The scatter plot shows the relationship between the input phase and the resulting change in spiking

1112.   latency of the recorded cell produced by the AM modulation (ordinate axis). Symbols for the centripetal

1113.   conditions : "x" in gold color (CP-CROSS-parallel) and "+" symbol in red color (CP-ISO-coaligned). Symbols for

1114.   the centrifugal condition : "*" brown star (CF-CROSS-parallel) and green diamond (CF-CROSS-coaligned). In grey,

1115.   bilinear fits. See Text for details. Note a significant reduction in spiking latency when the horizontal input is

1116.   integrated post-synaptically ahead of the feedforward drive by 5-25 ms (phase advance, rightwards). The

1117.   scatter plot data are projected on the x-axis (top row) or on the y-axis (right column) as Gaussian kernel density

1118.   estimators (KDE). The color of each KDE distribution follows the convention of the "cardinal" visual stimulation

1119.   protocol (Fig. 1B).

1120.   ----------

1121.   When pooling all cells (n=23 cells, 67 measures), the distribution of input phase relationships

1122.   ranged from +25 ms of phase advance to -50 ms of phase lag between horizontal and feedforward

1123.   input waves at the subthreshold level (abscissa in the bottom-right scatter plot of Fig. 10). As

1124.   expected, a strong bias in the distribution was observed in favor of phase advance of the Surround

1125.   input for the centripetal sequences (gold "x" (CP-CROSS) and red "+" (CP-ISO) symbols) and delayed

1126.   activation for the centrifugal sequences (brown "*" (CF-CROSS-parallel) and green diamond (CF-ISO-

1127.   coaligned) symbols). At the spiking level, the latency shortening of the contextual (Surround +

1128.   Center) output, compared to the Center-Only condition, appears crucially dependent on this input

1129.   phase relationship: it was only observed for centripetal condition, i.e. when the Surround response

1130.   onset (the horizontal input), preceded in time the Center-only response onset (the feedforward (FF)

1131.   input). The intercept of the grey bi-linear fit, shown in the scatter plot of Fig. 10, has been optimized

1132.   by two half-regression lines independently of the stimulus conditions (taking all the data points

1133.   together). The optimization of explained variance by a bi-linear regression composed of two connex

1134.   linear segments results in a partition of the data cloud in two abscissa regions on each side of the

1135.   zero phase axis: one (on the left side) where the horizontal input lags in phase the feedforward drive,

1136.   and for which no significant trend is observed (r^2=0.01; n=32, p>0.5); the other (on the right side),

1137.   where the horizontal input is postsynaptically integrated ahead of the feedforward drive by 5-25 ms

1138.   (r^2= 0.356; n=37; p<10^{-5}). In this latter regression domain, the observed reduction in spiking latency

1139.   (latency advance pointing upwards on the y-axis, reflects in an almost linear way the phase advance

1140.   of the synaptic echo of the horizontal input relative to that of the feedforward drive ("0" abscissa).

1141.   The Gaussian kernel density estimator (KDE) of the input phase distribution (along the x-axis,

1142.   above the scatter plot) clearly shows that the depolarizing input wave evoked by a collinear pair of

1143.   stimuli flashed in a centripetal sequence in the "Near" periphery (CP-ISO, in red) was arriving earlier

1144.   (17 ms from the scatter plot projections; 9 ms for paired data with CP-CROSS, in gold) than when the

1145.   stimuli were oriented orthogonally to the width axis path (CP-CROSS, in gold) (Wilcoxon WPSR test

1146.   on latencies, n=39, 2 conditions, p<0.01). Note that, in some cells, the latency advance at

1147.   both Vm and Spiking levels of the CP-CROSS stimulation (gold symbols) is non-null. However, the

1148.   average 9 ms difference in subthreshold responses of CP-ISO and CP-CROSS conditions of the

1149.   "cardinal" protocol corresponds to the 8 ms latency advance (between CP-ISO and Center-Only

1150.   response), found for the CP-ISO significant subpopulation of "significant" cells in the "radial"

1151.   protocol. Note that, for this precise contingent, the latency advance component of the CP-CROSS

1152.   response was indistinguishable from the reference (Figs 8 and 9).

1153.   Only centripetal collinear CP-ISO sequences could induce significant shortening in output spiking

1154.   latency, by an advance as large as 30 ms. As shown by the KDE distribution on the y-axis (right of the

1155.   scatter plot), the median latency reduction observed for the collinear axis during a centripetal

1156.   stimulation (red line on the shaded red envelope) was 11 ms. In the other conditions, either

1157.   centrifugal or centripetal "parallel" along the RF width axis, no significant latency advance could be

1158.   seen: the three other distribution fits are all centered on a null latency change. At the spiking level,

1159.   the median difference between the onset latency of CP- and CF-conditions is of 13 ms (respectively,

1160.   red and green lines).

1161. **4.3. Synergy depends on the relative match between apparent motion and intracortical horizontal propagation speed**

1162.   A subtle difference between the two types of protocols is that the AM speeds used in the

1163.   "radial" protocol depended on the GP duration and on the size of the SRF, and not only on the size of

1164.   the MDF (as this was the case for the "cardinal" case). This finer speed tuning was adjusted to the

1165.   characteristics of each individual cell since the response latency and strength of evoked responses

1166.   both depend on the stroke duration and the distance between strokes. For the 37 cells of the

1167.   "radial" protocol, the average AM speed was of 189 ± 47 °/sec, (ranging from 72 to 312 °/s), hence

1168.   below the average speed of 329 ± 101 °/s (range 175-500 °/s) used in the "cardinal" protocol. Note

1169.   that despite this difference, the global ASHP distributions reported in the "radial" and "cardinal"

1170.   protocols overlap closely with the overall range of distributions reported in (26) and (29) (mirror

1171.   histograms in left panel of Fig. 10). A detailed comparison shows that the inferred speed of laterally

1172.   conveyed activity is slightly accelerated for "dynamic" sequences of AM flows recruiting the

1173.   periphery (top of the histogram) than for "static" flashed stimuli measures (bottom), peaking

1174.   respectively at 0.175 (n=60) and 0.125 mm/ms (n=168). The slight shift in the ASHP distribution

1175.   reflects a change in the median values (0.225 vs 0.175 mm/ms) rather than in the mean values which

1176.   remain similar (0.231 and 0.251 mm/ms). Those results point to the notion that the speed of laterally

1177.   conveyed activity is not completely hard-wired but depends partly on the type of visual stimulus

1178.   used. Therefore, higher levels of spatio-temporal coherence along the motion path slightly increases

1179.   the speed of laterally mediated activity, leading most likely to a cumulative shortening of the

1180.   response onset latency at each updated locus of stimulation of the AM path in the cortex, a

1181.   phenomenology already reproduced in simulations of nested horizontal propagation (47).

1182.   Functionally, a critical prediction of our "working hypothesis" is that the visibility of the lateral

1183.   contextual modulation of the feedforward input should therefore increase when the speed of the

1184.   AM-evoked wave in cortex grows closer to horizontal propagation speed. Focusing on the SECTOR

1185.   configuration of the "radial" protocol, we studied more in depth the impact of the CP-ISO AM kinetic

1186.   effects on synaptic integration in V1 in a subset of twelve cells, by comparing, in each of these cells,

1187.   the impact of AM sequences of various speeds. The speed of lateral connectivity recruitment was

1188.   controlled in a graded way by introducing temporal delays between GPs strokes (detailed in the

1189.   chronograms in the bottom panel of Fig. 11). We first determined independently for each cell the

1190.   "optimal" AM speed of surround recruitment, which set the 100% nominative reference speed value

1191.   (ASHP). In addition, for each of these cells, similar AM sequences were replayed at 70%, 50% and

1192.   30% of their nominal speed value, and interleaved in a pseudo random fashion. In spite of cell-by-cell

1193.   differences in their absolute optimal ASHP values (varying between 150°/s to 250°/s), population

1194.   averages shown in Figure 11 were done by grouping across cells responses obtained at 100%, 70%,

1195.   50% and 30% of their respective "optimal" ASHP.

1196.   --------------

1197.   **Fig.11 : Dependency of the Contextual effect on the speed of the AM flow :**

1198.   **Bottom,** the chronograms depict the 4 inter-stroke interval conditions used in each cell to probe the temporal

1199.   dependency of the interaction between the horizontal and the feedforward synaptic waves (color code

1200.   reddens with speed value). The exact value of the "optimal" reference speed (100%) is defined on a cell-by-cell

1201.   basis. Mean speed values, averaged across cells (n=12), are given left of each chronogram. **Top,** population

1202.   averages for the CP-ISO SECTOR condition. All individual responses are realigned with the Center-Only

1203.   response onset. The color-coded overlaid waveforms, averaged across cells, allow the comparison of the time-

1204.   course of the response for the optimal speed value (red) and its S.E.M. envelope (shaded red) with that

1205.   observed for various proportional reductions of the AM flow speed (from top to bottom: 70% (dark orange),

1206.   50% (orange) and 30% (light orange)). The contextual response modulation amplitude decreases proportionally

1207.   to the AM speed reduction from its optimal value.

1208.   --------------

1209.   The comparison of the average contextual PSP response profiles of this subpopulation for

1210.   incremental AM speeds (30%, 50%, 70% and 100% of the "optimal" value) highlights a cumulative

1211.   speed-dependent anticipatory recruitment of the periphery, with a bumpy stairway-like profile

1212.   preceding the main response peak (top panel in Fig. 11). Crucially, the overlay of response profiles

1213.   for different speeds suggests a progressive merging of lateral inputs impinging on the central target

1214.   RF, from weaker amplitude and asynchronous early responses at lower speed to larger amplitude

1215.   and synchronous "in-phase"-summation with the feedforward drive (Center-Only, black curve in Fig.

1216.   11) when approaching the "optimal" AM speed. As speed value grows (from 30% (light orange) to

1217.   100% (red) of the optimal speed), lateral influence gets progressively synchronized when it reaches

1218.   the RF center. This is attested by the presence of dissociated bumps (preceding the Center response)

1219.   at slower speed, which likely reflects sequences of asynchronous activation by the distal surround

1220.   sources. In contrast, at the "optimal" speed, the precise time-concentration increase in depolarizing

1221.   activity power enables a more efficient interaction with the center-only response, thereby boosting

1222.   in a causal way the feedforward PSP. Once more, this result clearly emphasizes the crucial timing of

1223.   synaptic recruitment in the Surround, which controls the emergence of synergy between long-

1224.   distance (orange arrows in Fig. 1A) and cascades of shorter neighbor-to-neighbor links (grey arrows

1225.   in Fig. 1A) along the AM path. We conclude that "in-phase" summation of lateral inputs with the

1226.   predicted time of the feedforward drive produces the shortening of latency and amplitude

1227.   amplification reported during CP-ISO flow at high AM speed.

1228.   Altogether, our results show that facilitatory contextual control of V1 responses is optimized

1229.   along the orientation preference axis when the global motion speed lies within the range of visual

1230.   flow speed predicted from the apparent conduction velocity of horizontal connectivity. For slower

1231.   motion, the horizontally-driven facilitation effect disappears and stimuli cross-oriented to the motion

1232.   path, presented across the width axis, remain the best drivers of cortical cells. We conclude that the

1233.   axial preference of V1 cells shifts abruptly by an angle of 90 ° when the motion speed switches from

1234.   2-20 °/s to 100-250 °/s.

1235.   In summary, the spatial congruence of the elements regarding the preferred orientation axis,

1236.   the spatio-temporal coherence of the flows regarding the RF center (centripetal), the speed of

1237.   recruitment of the periphery and the precise timing selectivity of Surround-only responses, all

1238.   suggest that some form of peripheral subthreshold "prediction" is generated internally in V1, to

1239.   interact, within the proper spatio-temporal window, with the feedforward Center-Only synaptic

1240.   volley. These highly selective features justify, to our view, the qualification for the existence of neural

1241.   correlate of "predictive filling-in" in V1, to be distinguished from a non-specific anticipatory

1242.   depolarizing spread.

1243.

1244.   Discussion

1245.   Most studies of contextual modulation in the early visual system are based on synchronous

1246.   Surround and Center stimulation, pointing to mixed suppressive and facilitatory effects (13, 14, 48-

1247.   53). We document here a dynamic regime of asynchronous Surround-Center activations, of specific

1248.   speed and spatial anisotropy, which, under precise timing and co-alignment conditions, reveals a

1249.   facilitatory form of contextual cortical gain control. This effect, which links global motion direction

1250.   and local orientation, is strong enough in the anesthetized brain to shorten spiking response latency

1251.   by a few tens of milliseconds and shift the axial preference of V1 cells by 90° for saccade-like speed,

1252.   favoring contour integration along the orientation preference axis.

1253.   1. Orientation anisotropy, Motion streaks and Speed sensitivity

1254.   The dependency on co-alignment and motion along the RF preferred orientation axis reported

1255.   here correlates with the anatomical bias for iso-oriented long range connections described in higher

1256.   mammalian visual cortex (*cat*: (54-56); *tree shrew:* (57-59); *macaque* : (60, 61); *squirrel monkey* :

1257.   (62)). At the functional level, it is consistent with the concepts of the perceptual "Association Field"

1258.   in humans (4), the neural "facilitation field" in the behaving NHP (11, 63) and the synaptic "Dynamic

1259.   Association Field" in cat V1 (29). In terms of modulatory efficiency, the requirement of maximizing

1260.   the spatio-temporal synergy from the Surround corroborates the stimulus dependency in the

1261.   recruitment of long-distance iso-tuned interactions reported with imaging techniques at the

1262.   mesoscopic level (40).

1263.   Importantly in V1 neurons, the integration axis (the RF preferred orientation) for which the

1264.   strongest responsiveness modulation by fast (150-250°/s) apparent motion is observed, is

1265.   orthogonal to the preferred directional axis of V1 neurons (the RF width axis) when probed with

1266.   slowly moving bars and full field gratings. Classically, area 17 (V1) receptive fields are best activated

1267.   by slow motion across the RF width axis (1-2 °/s in (46); 3-10 °/s in (64), but quantitative studies

1268.   reveal however a zoo of speed tuning selectivities in V1 ((65-67); review in (68)). Among these,

1269.   "parallel motion direction selective" neurons could provide a potential substrate for the effects

1270.   reported here. However such trigger features are classically found in higher visual cortical areas

1271.   dedicated to global motion (type II MT cells (69)), but not in V1. In contrast, Geisler and colleagues

1272.   hypothesized, then described, a specific class of cells in V1 responding to Gaussian blobs, named

1273.   "streak" detectors, oriented in the direction of motion (70, 71). To account for disambiguation of

1274.   motion direction, Geisler introduced a hypothetical multiplier circuit between direction selective and

1275.   "parallel" detectors, whose existence needs still to be demonstrated (70). Realistic estimates for the

1276.   speed threshold above which "motion streak" operates point to medium range values, around 20-

1277.   60°/s for cat and 10-20°/s for monkey (inferred from (72)). This "streak" effect, however, depends on

1278.   the test stimulus,i.e. small isotropic Gaussian blobs, critical to avoid recruiting side-inhibition, and is

1279.   absent for oriented plaid patterns (71). Other studies reported related findings: Orban and

1280.   colleagues showed that spot directional tuning shifted by 60-90° above a speed threshold, whereas

1281.   bar tuning remained invariant (68, 73). Intrinsic optical imaging experiments - combining textures

1282.   tiled with small bars and independent motion components – point to a complex dependency of

1283.   orientation domains recruitment on local motif length and motion speed (74). Similarly, 90° shifts in

1284.   orientation domains were described for moving dot noise above 14.5°/s in cat area 17 (75).

1285.   In contrast, the 90° shift in axial motion sensitivity reported here is found at much faster

1286.   (saccade-like) AM speeds (>50°/s), and relies on the same local inducer (oriented Gabor) shifted in

1287.   space and time. It is the precise time-lagged recruitment of Surround-then-Center synaptic inputs,

1288.   that conditions the contextual amplification and the phase advance of the Center response. Both the

1289.   binding specificity of the local (orientation) and global (motion axis) features of the AM flow, on the

1290.   one hand, and the match between the optimal AM speed and ASHP values, on the other hand,

1291.   support the view that the main player is long-range horizontal connectivity intrinsic to V1.

1292.   2. Novelty of the study

1293.   The novelty of our findings is several fold : 1) these lateral diffusion effects recorded in the

1294.   anesthetized, hence non-attentive, animal are likely to be intrinsic to V1 and do not require

1295.   behavioral attention and top-down control from higher cortical areas (weakened by anesthesia) ; 2)

1296.   they depend on the spatiotemporal coherence of the AM sequence and grow with flow speed until

1297.   matching the apparent speed of horizontal propagation (ASHP) in V1; 3) in that range of speed, they

1298.   produce a shift by 90° of axial motion sensitivity, which becomes co-aligned with the preferred

1299.   orientation RF axis; 4) they provide evidence for an internally-generated propagation process,

1300.   binding local (orientation and position) and global (motion and direction) features; 5) they constitute

1301.   a plausible neural substrate of psychophysical effects in humans reporting a perceptual bias in speed

1302.   estimate for collinear motion (76).

1303. 2.1. Intra-V1 horizontal diffusion vs. cortico-cortical feedback

1304.   Conventional models of visual cortical processing assume that complex computations, such as

1305.   motion direction disambiguation, are encoded only at higher levels in the visual cortical area

1306.   hierarchy, and are absent in V1 (9, 10, 77-81)). Most electrophysiological studies in the behaving

1307.   primate point to the late retroaction in V1 of a top-down signal gated by attention (82-85). They

1308.   attribute the delay of several tens of milliseconds of the change in firing of V1 cells after response

1309.   onset, to the late action of the cortico-cortical feedback (86-90), rather than to the slow horizontal

1310.   propagation in V1. However, puzzlingly enough, axonal feedback transmission from higher areas is

1311.   10-100 times faster than intrinsic diffusion in V1 (see also (91) for the parallel recruitment of dual

1312.   lateral-feedback kinetics).

1313.   The challenging view presented here is that, as early as V1, horizontal connections play already

1314.   an important role in the early perception of retinal flow and global motion. The visibility and spatial

1315.   spread of their functional dominance in cats & ferrets vs non-human primates (NHP) may reflect

1316.   species-specific differences in retino-cortical magnification factor. Horizontal axon length (4-8 mm) is

1317.   the same across species, but covers up to 8° of visual angle in cat-ferret (_cat_ : (54, 55, 92); _ferret_ :

1318.   (59) whereas it remains within a radius of 1.5 hypercolumn (2-3°) in primates (93), which matches

1319.   the spatial extent of collinear facilitation observed in human psychophysics (42, 94-96).

1320.   The multi-stroke AM protocols ("cardinal" and "radial") were both designed to promote the in-

1321.   phase synchronization of the synaptic impacts produced by the sequential stimulation of each GP

1322.   flashed along the motion path with the central feedforward drive (orange arrows in Fig. 1A). Our

1323.   prediction was that this synchronization would be optimized for a flow speed in the range of the

1324.   apparent propagation speed of horizontal connectivity (ASHP in (26)). This view is confirmed by the

1325.   speed-dependency tuning study reported here (see Results). However, the variability of the

1326.   intracellular anticipatory Vm profile across cells suggests that other synaptic recruitment paths

1327.   are likely to contribute, in particular the neighbor-to-neighbor links (grey arrows in Fig. 1A). It is thus

1328.   likely that the "horizontal" drive efficiency results from a combination of the asynchronous visual

1329.   activation of two classes of synaptic sources: 1) long-range monosynaptic inputs recruited at

1330.   different eccentricities, whose distributed and delayed impact is integrated in phase by the target

1331.   cell; 2) polysynaptic short-range input, reflecting chains of next-to-next neighbor activation (studied

1332.   in our previous 2-stroke AM (29)). This second effect is amplified by the re-ignition process, imposed

1333.   at each GP locus by the next feedforward input recruited on the apparent motion trajectory. Note

1334.   that this rolling wave effect is not seen for single stroke stimuli, since, otherwise, the spiking

1335.   discharge field of the recorded cell would extend far beyond the spatial region defined by the core

1336.   thalamo-cortical projection.

1337.   Since the reverberation delay imposed by cortico-cortical loops is in the order of a few

1338.   milliseconds only (97), it is plausible that most long-range interactions in the primate may operate by

1339.   nested polysynaptic feedback afferents from higher cortical areas through chains of

1340.   neighbor-to-neighbor lateral relays, whereas in the cat and ferret, a substantial part would remain

1341.   encapsulated monosynaptically within V1 (7). Our results may provide new insights on the

1342.   evolutionary complexification of cortico-cortical computation. They could also spark interest in AI

1343.   and deep learning in artificial vision, where the diffusion effects of horizontal connectivity in a single

1344.   V1-like terminal layer can be confronted with the backpropagation of "network belief" implemented

1345.   by nested feedback connections in a multi-layer hierarchical cortical-like architecture (98).

1346. ## 2.2. Dependency on spatiotemporal coherence and visual flow speed

1347.   Since 2-stroke AM provided evidence mostly for subthreshold synaptic amplification (29), we

1348.   aimed here at promoting the efficiency of centripetal AM CP-ISO sequences to produce a spiking

1349.   discharge. This was achieved by increasing the number of elementary strokes in the AM sequence,

1350.   from 3 to 6 from the "cardinal" to the "radial" protocols. In the "radial" protocol, the comparison

1351.   between FULL-RND, SECTOR-RND and SECTOR-CP-ISO conditions (Fig 7 right panel) provides

1352.   evidence that the boosting of sensory responsiveness grows with the level of spatio-temporal

1353.   coherence. The speed dependency of the effect demonstrates unambiguously that the efficiency of

1354.   the contextual control of the cortical response gain and phase is optimized when a causal coherence

1355.   is reached, i.e. when the horizontal wave predicts the arrival of the feedforward wave.

1356.   The optimal range of speed (150-250°/s) reported here contrasts with a recent study of

1357.   anticipatory responses in the fixating monkey, best seen for stimuli cross-oriented to the motion

1358.   path, in continuous slow motion across the RF width axis (6.6°-13.2°/s) (99). The authors propose a

1359.   complex mechanism, based on the interplay of the "feedforward" trace imposed by the continuous

1360.   motion of the stimulus and a medium range isotropic horizontal propagation wave. This activity

1361.   bump still allows a suboptimal slight interaction with the feedforward drive when it propagates

1362.   faster than the stimulus. Their interplay would cause a pre-activation of the cells, facilitating in turn

1363.   the feedforward reactivation. This proposed mechanism is highly dependent on the retino-cortical

1364.   magnification factor and does not account for the global dynamics of the anticipatory wave for

1365.   distances beyond monosynaptic connection lateral extent. The propagation speed across long

1366.   trajectories (> 3° from the Spiking discharge field center), well beyond the spatial extent of

1367.   monosynaptic links observed in NHP (<2°), is indeed ten times slower (<= 0.02 mm/ms) than the

1368.   apparent speed of horizontal propagation (0.05-0.50 mm/ms; see (26, 32)). The global kinetics are

1369.   more in line with a slower "rolling wave" diffusion mechanism (see also *in vitro* (100)). The speed

1370.   range of this anticipatory wave reported in awake macaque (<15°/s) is lower than in the anesthetized

1371.   cat area 17 (15-40°/s in (39)), where it has been shown that a moving square is more rapidly

1372.   processed than a flashed one. These different effects share some similarities with those reported

1373.   here, but differ significantly in their speed range (by a factor of 10-20) and preferred axial

1374.   sensitivities (CROSS-parallel vs ISO-Collinear). Furthermore, they do not require collinearity, a

1375.   hallmark of horizontally-mediated V1 computation.

1376. 2.3. Binding of global and local features as early as V1

1377.   The current consensus concerning cortical correlates of low-level perception mixes hierarchical

1378.   processing and cortico-cortical recurrency : V1 plays a pivotal role in synchronization dynamics back

1379.   and forth with high-order cortical areas (16, 101). Intra-V1 synchronization and inhibitory

1380.   interactions are hypothesized to interact through a form of "competitive coding", before the output

1381.   is further processed downstream in the cortical hierarchy. A non-exclusive alternative is that

1382.   reverberating activity in V1 already participates in non-attentive "pop-out" perception. We propose

1383.   that horizontal connections, intrinsic to V1, are instrumental to the neural implementation of low-

1384.   level perceptual Gestalt laws, in parallel or before information is broadcasted to the mosaic of higher

1385.   cortical areas. Our data suggest that, in Gestalt-like configurations, horizontal connectivity

1386.   propagates some kind of "network belief" (built-up on past activity ignition and shared input

1387.   statistics in the Surround) to the connex V1 neighborhood, the most likely to be next stimulated

1388.   (102).

1389.   Remarkably enough, the "prediction" wave, seen for "Surround-Only" AM CP-ISO sequences,

1390.   binds local information acquired along the motion path. This observation, although recorded in the

1391.   anesthetized brain, globally fits with Bayesian theoretical frameworks, where omission of the next-

1392.   to-come stimulus in repetitive and predictive activation sequences unmasks an intracortically

1393.   generated "expectation" signal mediated by horizontal connectivity. This finding has to be

1394.   distinguished from the classical "predictive coding" schema (103), where what is propagated is the

1395.   "residual error" from the expectation generated in higher cortical areas rather than the diffusion of

1396.   the network belief context (98). At optimized speed, the self-generated "Surround-Only" wave is

1397.   shown here to exhibit the same timing as the feedforward (FF) input onset evoked by the next

1398.   stimulus to come, had it been presented. More exploration, outside the scope of this paper, would

1399.   be needed to test how center-surround interaction is affected by the contrast (low/high) and the

1400.   orientation (same/mismatch) of the test Center stimulus, and how possible sub-linearities can be

1401.   interpreted in the context of the principle of redundancy reduction framework (104).

1402.   Our results suggest that V1 may solve the motion extrapolation problem just through selective

1403.   internal diffusion. Our interpretation is supported by theoretical studies of motion extrapolation

1404.   (ability to continue to predict accurately the position/speed of a moving object in the sudden

1405.   absence of visual input) : Kaplan and colleagues implemented a realistic spiking recurrent neural

1406.   network simulation of V1/MT (105), and studied cortical dynamics relaxation, when the coherent

1407.   motion of a dot was abruptly interrupted by a "blank" period for three different lateral connectivity

1408.   distributions (random, isotropic and anisotropic). The main finding, in agreement with this study, was

1409.   that anisotropy was required to allow an efficient diffusion of motion information and achieve

1410.   accurate motion/position extrapolation in the absence of FF validation (Fig. 4 in (105)).

1411. 2.4.   A neural correlate of perceptual bias in speed estimate for collinear motion

1412.   It is well established that "pop-out" and feature binding in Gestalt psychology do not depend

1413.   invariably on attentive behavior, although they can be trained during perceptual learning and

1414.   enhanced by top-down cognitive processes (11,15,16). The neural mechanisms, studied here in the

1415.   anesthetized animal, can participate in automatic bottom-up attention, a form of sensory-driven

1416.   selection that facilitates perception in the behaving state of a subset of the stimulus. Our working

1417.   hypothesis is that the built-in bias in synaptic integration we report in primary visual cortex reflects

1418.   lateral connectivity anisotropies engrammed during visuomotor development. This view is strongly

1419.   supported at the functional level by the similarities in the visual cortical "association fields" found at

1420.   the synaptic receptive field level in the anesthetized brain (our work), at the spiking discharge field

1421.   level in the awake behaving state (Li and Gilbert) and at the behavioral and perceptual level (Field

1422.   and Hess). We report here elementary changes at the subthreshold and spiking levels in V1,

1423.   detectable at the intracellular level in the anesthetized state, that may not be strong enough to

1424.   mediate an attention-guided behavior, but are efficient enough to produce an "automatic" pop-out

1425.   effect.

1426.   Our electrophysiological findings constitute a potential correlate, as early as V1, of the human

1427.   psychophysical speed-up bias in motion flow integration (47, 76). These previous psychophysical and

1428.   computational studies from our lab showed that collinear AM sequences - composed of co-aligned

1429.   ISO-Gabor patches along the global motion axis - are perceived by humans as moving "faster" than

1430.   AM sequences composed by "parallel" CROSS-oriented elements, of the same exact physical speed.

1431.   The speed bias decreased as the angle between the local GP inducer and the global motion axis

1432.   increased. This bias was highly dependent on the orientation selectivity of the inducer and

1433.   disappeared for blobs, which suggests a cortical origin. All reported features show a close link with

1434.   the synaptic association field of V1 neurons (29) and the reported data.

1435.   Crucially, the speed for which the speed-up effect is the strongest in humans is 64 °/s.

1436.   Assuming a parafoveal retino-cortical magnification factor ratio of 3 between human and cat

1437.   (*human* : (106); *cat* : (107)), the optimal speed-up range of 40-96°/s seen in human would

1438.   correspond to 120-288°/s in cat, which fits remarkably well the 100-312°/s reported here. The 64°/s

1439.   optimal velocity for humans corresponds for the cat to a speed of 192°/s on the retina, and an

1440.   apparent propagation speed of 0.19 mm/ms in cortex, in agreement with the inference made from

1441.   our intracellular recordings. Most remarkably, the boosting mechanism revealed in this study

1442.   confirms the prediction of an earlier computational model of our laboratory (47), which provided a

1443.   simple conceptual framework where synaptic summation between feedforward and lateral AM

1444.   inputs results in amplification and latency advance in V1 spiking responses.

1445.   Our intracellular study allows to go one step beyond this spike-based modeling. In (47), the

1446.   composite synaptic summation is linear and the non-linearity of the functional effect is simply due to

1447.   that of spike initiation. However, the intracellular study of the timing dependency of the association

1448.   process suggests a non-linearity in the interaction process between feedforward (FF) and horizontal

1449.   input, which becomes supra-linear when FF input lags by more than 5.5 ms the contextual signal

1450.   (Fig. 7 C in (29); see also here Fig. 10, right pane)). It remains plausible that a minimal integration

1451.   time is needed to recruit non-linear voltage-dependent mechanisms recruited by horizontal

1452.   connectivity, such as the persistent sodium current or NMDA receptor activation (25, 108).

1453.   ## 3. Concluding statements

1454.   In conclusion, the view presented here differs significantly from classical studies/models of top-

1455.   down gain control of V1 processing which ignore the contribution *per se* of horizontal connectivity

1456.   (11, 15, 16) or attribute the late enhancement of visual responses in V1 to the sole feedback from

1457.   higher cortical areas (83). We do not argue here against the consensus that top-down signals are

1458.   required, during attentive wakefulness, to express and/or amplify an integrative process intrinsic to

1459.   V1. However, our recent demonstration of a "synaptic association field" in the anesthetized

1460.   preparation demonstrates a preexisting structural bias intrinsic to V1, which could constitute the

1461.   synaptic footprint of the neural architecture needed to implement Gestalt laws (29). This finding is

1462.   further strengthened by a multiscale imaging study (40), suggesting that a critical threshold of spatial

1463.   summation has to be passed in order to make these long-range interactions detectable, while

1464.   temporal synergy conditions the effective binding of those spatial interactions (39). A likely

1465.   interpretation is that some kind of resonance regime, such as produced here by the multiple stroke

1466.   reafferent drive, is needed in the anesthetized state to propagate at the spiking level a functional

1467.   bias already present in the subthreshold impact of horizontal connectivity.

1468.   The novel conceptual focus here is on the temporal features of horizontal propagation and on

1469.   intra-V1 broadcasting. State-of-the-art modeling generally ignores the propagation distance-

1470.   dependent delay component whose impact compromises greatly the analytical solving of mean-field

1471.   equations (109); but see (110)). Interestingly, recent large-scale network simulations with

1472.   topographically-organized horizontal monosynaptic connectivity and distance-dependent axonal

1473.   conduction delays (0.10-0.60 m/s) demonstrate the consistency of spontaneous subthreshold waves

1474.   with the asynchronous-irregular regime. The multiplexing of both processes defines a sparse-wave

1475.   regime, more in line with a realistic view of a functional V1 (36). Although our interpretation of the

1476.   present results seems in agreement with some of the computational literature on predictive coding

1477.   applied to contour-through-motion extrapolation and trajectory extraction, they still remain to be

1478.   confronted with alternative models which do not require propagation of a prediction. Other classes

1479.   of probabilistic and static models (71) do not depend on propagation at finite speed and apply

1480.   primarily to isotropic stimuli which are known to be suboptimal in activating V1.

1481.   Overall, our recordings suggest that fast collinear flow facilitates the refinement of anisotropy

1482.   integration, distributed over the mesoscopic representation of trajectories across the "horizontal"

1483.   layer plane of V1. The Surround-only CP-ISO anticipatory wave, captured "at the right position, at the

1484.   right time" in a subset of our intracellular recordings, is compatible with a progressive topological

1485.   refinement of activity diffusion in cortical space. In cat visual cortex, using annular stimuli and

1486.   voltage sensitive dye imaging, Chavane et al reported correlates of "filling-in" responses (40), which

1487.   fit with our subthreshold membrane potential recordings. Using intrinsic optical imaging (which

1488.   might reflect more spiking than subthreshold activity), Chisum et al reported in the anesthetized tree

1489.   shrew that ISO-oriented collinear Gabor arrays flashed simultaneously did enhance locally V1 sensory

1490.   responsiveness (111). This latter study, which used a static input pattern of 2 second duration,

1491.   activating simultaneously all the Surround sites, detected some activity strengthening in the

1492.   areas directly (FF) activated by co-aligned local inducers. However, it failed to observe any spatial

1493.   expansion of lateral activity in the cortical regions representing stimulus gaps. In contrast, our study

1494.   predicts that the dynamic apparent motion animation of the same array, activating in succession the

1495.   co-aligned local oriented inducers, would generate a continuous anisotropic propagation wave, most

1496.   visible for AM speed faster than 50°/s. This diffusion process would, in turn, fill the gaps between

1497.   local feedforward activation zones and promote the extraction of a motion trajectory riding on an

1498.   expanding collinear contour. Evidence in V1 for an expansive "filling-in" process during apparent

1499.   motion finds some support in fMRI imaging studies in humans (112). It could be tested spatio-

1500.   temporally with more precision in cat and NHP, using voltage sensitive dye imaging and fast AM

1501.   sequences.

1502.   Irrespective of the species-dependency of the visual spatial coverage of horizontal connections

1503.   (7-8° in cats vs 1-2° in NHPs, in spite of similar axonal length), multiple stroke AM sequences appear

1504.   best suited to dynamically recruit overlapping "lateral connectional fields" along the motion

1505.   trajectory (90). The optimal separation in the visual field between the GP inducers might however be

1506.   still species-specific, since the orientation selectivity of the horizontal facilitation process seems most

1507.   prominent when co-stimulating the "Far" surround of GP-driven RFs.

1508.   This intriguing propagation of self-generated and propagated prediction at an early stage of visual

1509.   cortical processing may underlie a more profound functional role. The classically reported "edge-

1510.   detector" property of V1 neurons, as initially described by Hubel and Wiesel (46), highlights the fact

1511.   that the strongest discharge is evoked by "parallel" elements swept towards the RF center

1512.   orthogonally across its width axis. In contrast, at high AM speed (100-500°/s), the synaptic

1513.   integration of the "silent" periphery was found to depend on the motion axis. This dynamic process

1514.   unveils a spatial selectivity and anisotropy not present in the MDF mapped with static flashed

1515.   light/dark impulses. This suggests that fast retinal flow produced by eye-movements could change

1516.   radically contour integration selectivity of V1 neurons. The dynamic reconfiguration by fast retinal

1517.   slip of the RF main axis could account for the peculiar geometry of saccadic scanpaths of human

1518.   observers during the viewing of other human faces. Indeed, Alfred Lioukianovitch Yarbus long ago

1519.   reported a striking similarity between the topographic layout of the saccadic scanpaths with the

1520.   most salient long collinear contours outlining the perceptual skeleton of the face viewed by the

1521.   observer ((27, 113, 114); review in (115)).

1522.

1523. **<u>Acknowledgments :</u>**

1524.   We thank Vincent Bringuier, Frédéric Chavane, Jean Lorenceau, Peggy Seriès and Florian Gérard-

1525.   Mercier for their past collaborative contribution to the concept of the "synaptic association

1526.   field". We are indebted to Frédéric Chavane, who participated in his PhD work to some of the

1527.   early experiments. We thank Andrew Davison for helpful comments and help with the English.

1528.   We thank Gérard Sadoc for the in-house development of visual stimulation and data acquisition

1529.   software (Elphy^{TM}), Kirsty Grant and Guillaume Hucher for histology and Aurélie Daret for her

1530.   support during experimental surgery.

References :

1. Wertheimer M. Experimentelle Studien über das Sehen von Bewegung [Habilitation Frankfurt]. Leipzig: J. A. Barth; 1912. Zeitschrift für Psychologie und Physiologie der Sinnesorgane, 61;161-265.

2. Koffka K. Principles of Gestalt psychology. New York: Harcourt; 1935. 720 p.

3. Groen IIA, Silson EH, Baker CI. Contributions of low- and high-level properties to neural processing of visual scene sin the human brain. Philos Trans R Soc Lond B Biol Sci. 2017;372(1714).

4. Field DJ, Hayes A, Hess RF. Contour integration by the human visual system: evidence for a local "association field". Vision Research. 1993;33(2):173-93.

5. Hess R, Field D. Integration of contours: New insights. Trends in Cognitive Sciences. 1999;3(12):480-6.

6. Neri P, Morrone MC, Burr DC. Seeing biological motion. Nature. 1998;395:894-6.

7. Frégnac Y, Bathellier B. Cortical correlates of low-level perception: from neural circuits to percepts. Neuron. 2015;88(1):110-26.

8. Felleman DJ, Van Essen DC. Distributed hierarchical processing in the primate cerebral cortex. Cerebral Cortex. 1991;1(1):1-47.

9. Heeger DJ, Simoncelli EP, Movshon JA. Computational models of cortical visual processing. Proceedings Natlonal Academy of Science USA. 1996;93:623-7.

10. Serre T, Oliva A, Poggio T. A feedforward architecture accounts for rapid categorization. Proceedings Natlonal Academy of Sciences of the USA. 2007;104(15):6424-9.

11. Li W, Piëch V, Gilbert CD. Contour saliency in primary visual cortex. Neuron. 2006;50:951-62.

12. Polat U, Mizobe K, Pettet MW, Kasamatsu T, Norcia AM. Collinear stimuli regulate visual responses depending on cell's contrast threshold. Nature. 1998;391:580-4.

13. Mizobe K, Polat U, Pettet MW, Kasamatsu T. Facilitation and suppression of single striate-cell activity by spatially discrete pattern stimuli presented beyond the receptive field. Visual Neuroscience. 2001;18:377-91.

14. Kasamatsu T, Miller R, Zhu Z, Chang M, Ishida Y. Collinear facilitation is independent of receptive-field expansion at low contrast. Experimental Brain Research. 2010;201(3):453-65.

15. Li W, Piëch V, Gilbert CD. Learning to link visual contours. Neuron. 2008;57:442-51.

16. Gilbert CD, Li W. Top-down influences on visual processing. National Review of Neuroscience. 2013;14(5):350-63.

17. Movshon A, Adelson EH, Gizzi MS, Newsome WT. The analysis of moving visual patterns. In: Chagas C, Gattass, R.,Gross, C., editor. Pattern Recognition Mechanisms: Pontificiae Academiae Scientiarum Scripta Varia. 1985;54:117-51.

18. Castelo-Branco M, Formisano E, Backes W, Zanella F, Neuenschwander S, Singer W, et al. Activity patterns in human motion-sensitive areas depend on the interpretation of global motion. Proceedings of the Natlonal Academy of Science U S A. 2002;99:13914-9.

19. Aaen-Stockdale C, Thompson B. Visual Motion: From Cortex to Percept. IntechOpen. 2011.

20.   Aissani C, Cottereau B, Dumas G, Paradis A-L, Lorenceau J. Magnetoencephalographic signatures of visual form and motion binding. Brain Research. 2011;1408:27-41.

21.   Hirsch JA, Alonso JM, Reid RC, Martinez LM. Synaptic integration in striate cortical simple cells. The Journal of Neuroscience,. 1998;18:9517-28.

22.   Ferster D, Miller KD. Neural mechanisms of orientation selectivity in the visual cortex. Annual Review of Neuroscience. 2000;23:441-71.

23.   Douglas RJ, Martin KA. Neuronal circuits of the neocortex. Annual Review of Neuroscience. 2004;27:419-51.

24.   Priebe NJ, Ferster D. Mechanisms of neuronal computation in mammalian visual cortex. Neuron. 2012;75(2):194-208.

25.   Frégnac Y, Bringuier V. Spatio-temporal dynamics of synaptic integration in cat visual cortical receptive fields. In: Braitenberg AAaV, editor. Brain Theory: Biological basis and computational theory of Vision. 1996. p. 143-99.

26.   Bringuier V, Chavane F, Glaeser L, Fregnac Y. Horizontal propagation of visual activity in the synaptic integration field of area 17 neurons. Science. 1999;283:695-9.

27.   Baudot P. Nature is the code: high temporal precision and low noise in V1. Paris: University Paris VI; 2006.

28.   Haider B, Krause MR, Duque A, Yu Y, Touryan J, Mazer JA, et al. Synaptic and network mechanisms of sparse and reliable visual cortical activity during nonclassical receptive field stimulation. Neuron. 2010;65(1):107-21.

29.   Gerard-Mercier F, Carelli PV, Pananceau M, Troncoso XG, Fregnac Y. Synaptic Correlates of Low-Level Perception in V1. The Journal of Neuroscience. 2016;36(14):3925-42.

30.   Ozeki H, Finn IM, Schaffer ES, Miller KD, Ferster D. Inhibitory Stabilization of the Cortical Network Underlies Visual Surround Suppression. Neuron. 2016;62(4):578-92.

31.   Das A, Gilbert CD. Long-range horizontal connections and their role in cortical reorganization revealed by optical recording of cat primary visual cortex. Nature. 1995;375(6534):780-4.

32.   Grinvald A, Lieke EE, Frostig RD, Hildesheim R. Cortical point-spread function and long-range lateral interactions revealed by real-time optical imaging of macaque monkey primary visual cortex. The Journal of Neuroscience. 1994;14:2545-68.

33.   Slovin H, Arieli A, Hildesheim R, Grinvald A. Long-term voltage-sensitive dye imaging reveals cortical dynamics in behaving monkeys. Journal of Neurophysiology. 2002;88:3421-38.

34.   Reynaud A, Masson GS, Chavane F. Dynamics of Local Input Normalization Result from Balanced Short- and Long-Range Intracortical Interactions in Area V1. The Journal of Neuroscience. 2012;32:12558-69.

35.   Muller L, Reynaud A, Chavane F, Destexhe A. The stimulus-evoked population response in visual cortex of awake monkey is a propagating wave. Nature 2014;5:3675.

36.   Davis ZW, Muller L, Martinez-Trujillo J, Sejnowski T, Reynolds JH. Spontaneous travelling cortical waves gate perception in behaving primates. Nature. 2020;587(7834):432-6.

37.   Benucci A, Frazor RA, Carandini M. Standing waves and traveling waves distinguish two circuits in visual cortex. Neuron. 2007;55(1):103-17.

38.   Sato TK, Nauhaus I, Carandini M. Traveling waves in visual cortex. Neuron. 2012;2:218-29.

39.   Jancke D, Erlhagen W, SchÜner G, Dinse HR. Shorter latencies for motion trajectories than for flashes in population responses of cat primary visual cortex. Journal of Physiology. 2004;556(3):971-82.

40.   Chavane F, Sharon D, Jancke D, Marre O, Fregnac Y, Grinvald A. Lateral Spread of Orientation Selectivity in V1 is Controlled by Intracortical Cooperativity. Frontiers System Neuroscience. 2011;5:4.

41.   Xu W, Huang X, Takagaki K, Wu J. Compression and reflection of visually evoked cortical waves. Neuron. 2007;55:119-29.

42.   Polat U, Sagi D. Lateral interactions between spatial channels, suppression and facilitation revealed by lateral masking experiments. Vision Research. 1993;33:993-9.

43.   Fournier J, Monier C, Pananceau M, Frégnac Y. Adaptation of the Simple or Complex nature of V1 receptive fields to visual statistics. Nature Neuroscience. 2011;14:1053-60.

44.   Fournier Y, Monier C, Levy M, Marre O, Sari K, Kisvarday ZF, et al. Hidden Complexity of synaptic receptive fields in cat primary visual cortex. The Journal of Neuroscience. 2014;34(16):5515-28.

45.   Chavane F. Diversité d'expression fonctionnelle du champ d'intégration synaptique : une re-évaluation intracellulaire des champs récepteurs des neurones du cortex visuel primaire. Paris: University Paris VI; 1999.

46.   Hubel DH, Wiesel TN. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. Journal of Physiology (London). 1962;160:106-54.

47.   Series P, Georges S, Lorenceau J, Fregnac Y. Orientation dependent modulation of apparent speed: a model based on the dynamics of feed-forward and horizontal connectivity in V1 cortex. Vision Research. 2002;42(25):2781-97.

48.   Nelson JI, Frost BJ. Intracortical facilitation among co-oriented co-axially aligned simple cells in cat striate cortex. Experimental Brain Research. 1985;61:54-61.

49.   Li C, Li W. Extensive integration field beyond the classical receptive field of cat's striate cortical neurons: Classification and tuning properties. Vision Research. 1994;34:2337-55.

50.   Kapadia MK, Ito M, Gilbert CD, Westheimer G. Improvement in visual sensitivity by changes in local context: Parallel studies in human observers and in V1 of alert monkeys. Neuron. 1995;15:843-56.

51.   Sengpiel F, Sen A, Blakemore C. Characteristics of surround inhibition in cat area 17. Experimental Brain Research. 1997;116:216-28.

52.   Walker GA, Ohzawa I, Freeman RD. Suppression outside the classical cortical receptive field. Visual Neuroscience. 2000;17:369-79.

53.   Chen K, Song XM, Li CY. Contrast-dependent variations in the excitatory classical receptive field and suppressive nonclassical receptive field of cat primary visual cortex. Cerebral Cortex. 2013;23(23):283-92.

54.   Gilbert CD, Wiesel TN. Columnar specificity of intrinsic horizontal and corticocortical connections in cat visual corte. The Journal of Neuroscience. 1989;9:2432-42.

55.   Kisvarday ZF, Toth E, Rausch M, Eysel UT. Orientation-specific relationship between populations of excitatory and inhibitory lateral connections in the visual cortex of the cat. Cerebral cortex. 1997;7(7):605-18.

56.   Schmidt KE, Goebel R, Löwell S, Singer W. The perceptual grouping criterion of colinearity is reflected by anisotropies of connections in the primary visual cortex. European Journal of Neuroscience. 1997;9:1083-9.

57.   Fitzpatrick D. The functional organization of local circuits in visual cortex: insights from the study of tree shrew striate cortex. Cerebral Cortex. 1996;6:329-41.

58.   Chisum J, Fitzpatrick D. The contribution of vertical and horizontal connections to the receptive field center and surround in V1. Neural Networks. 2004;17:681-93.

59.   Bosking WH, Zhang Y, Chofield B, Fitzpatrick D. Orientation selectivity and the arrangement of horizontal connections in tree shrew striate corte. The Journal of Neuroscience. 1997;17(6):2112-27.

60.   McGuire BA, Gilbert CD, K. RP, Wiesel TN. Targets of horizontal connections in macaque primary visual cortex. Journal of Comparative Neurology. 1991;305:370-92.

61.   Malach R, Amir Y, Harel M, Grinvald A. Relationship between intrinsic connections and functional architecture revealed by optical imaging and in vivo targeted biocytin injections in primate striate cortex. Proceedings of the Natlonal Academy of Sciences of the USA. 1993;90:10469-73.

62.   Sincish LC, Blasdel GG. Oriented axon projections in primary visual cortex of the monkey. The Journal of Neuroscience. 2001;21:4416.

63.   Kapadia MK, Westheimer G, Gilbert CD. Spatial distribution of contextual interactions in primary visual cortex and in visual perception. Journal of Neurophysiology. 2000;84:2048-62.

64.   Pettigrew JD, Nikara T, Bishop PO. Binocular interaction on single units in cat striate cortex: simultaneous stimulation by single moving slit with receptive fields in correspondence. Experimental Brain Research. 1968;6:391-410.

65.   Orban GA, Kennedy H, Maes H. Response to movement of neurons in areas 17 and 18 of the cat: velocity sensitivity. Journal of Neurophysiology. 1981;45:1043-58.

66.   Hammond P, Smith AT. On the sensitivity of complex cells in feline striate cortex to relative motion. Experimental Brain Research. 1982;47(3):457-60.

67.   Duysens J, Orban JA, Cremieux J. Functional basis for the preference for slow movement in area 17 of the cat. Vision Research. 1984;24(1):17-24.

68.   Orban GA. Neuronal operations in the visual cortex. Studies in brain function. Berlin: Springer-Verlag; 1984.

69.   Albright TD. Direction and Orientation Selectivity of Neurons in Visual Area MT of the Macaque. Brain Research. 1984;1408:27-40.

70.   Geisler WS. Motion streaks provide a spatial code for motion direction. Nature. 1999;400(6739):65-9.

71.   Geisler WS, Albrecht DG, Crane A, Stern L. Motion direction signals in the primary visual cortex of cat and monkey. Visual Neuroscience. 2001;18:5001-516.

72.   Frazor RD, Albrecht DG, Geisler WS, Crane AM. Visual cortex neurons of monkeys and cats: Temporal dynamics of the spatial frequency response functio. Journal of Neurophysiology. 2004;91:2607-27.

73.   Crook JM, Worgötter F, Eysel UT. Velocity invariance of preferred axis of motion for single spot stimuli in simple cells of cat striate cortex. Experimental Brain Research. 1994;102:175-80.

74.   Basole A, White LE, Fitzpatrick D. Mapping multiple stimulus features in the population response of visual cortex. Nature. 2003;423:775-87.

75.   An X, Gong H, McLoughlin N, Yang Y, Wang W. The mechanism for processing random-dot motion at various speeds in early visual cortices. PLoS ONE. 2014;9(3):e93115.

76.   Georges S, Series P, Fregnac Y, Lorenceau J. Orientation dependent modulation of apparent speed: psychophysical evidence. Vision Research. 2002;42:2757-72.

77.   Mumford DB. On the computational architecture of the neocortex II: The role of cortico-cortical loops. Biological Cybernetics. 1992;66:241-51.

78.   Burr DC, Morgan MJ. Motion deblurring in human vision. Proceedings of the Royal Society London. 1997;264:431-6.

79.   Zhou H, Friedman HS, Von der Heydt R. Coding of border ownership in monkey visual cortex. The Journal of Neuroscience. 2000;20:6594-611.

80.   Born RT, Bradley DC. Structure and Function of Visual Area MT. Annual Review of Neuroscience. 2005;28:157-89.

81.   Roe AW, Chelazzi L, Connor CE, Conway B, Fujita C, Gallant JL, et al. Toward a Unified Theory of Visual Area V4. Neuron. 2012;74(1):12-29.

82.   Schwabe L, Ichida JM, Shushruth S, Mangapathy P, Angelucci A. Contrast-dependence of surround suppression in Macaque VI: experimental testing of a recurrent network model. Neuroimage. 2010;52:777-92.

83.   Poort J, Raudies F, Wannig A, Lamme VA, Neumann H, Roelfsema PR. The role of attention in figure-ground segregation in areas V1 and V4 of the visual cortex. Neuron. 2012;75(1):143-56.

84.   Nassi JJ, Lomber SG, Born RT. Corticocortical feedback contributes to surround suppression in V1 of the alert primate. The Journal of Neuroscience. 2013;33:8504-17.

85.   Bastos AM, Vezoli J, Bosman CA, Schoffelen JM, Oostenveld R, Dowdall JR, et al. Visual areas exert feedforward and feedback influences through distinct frequency channels. Neuron. 2015;85(2):390-401.

86.   Lamme VA, Supèr H, Landman R, Roelfsema PR, Spekreijse H. The role of primary visual cortex (V1) in visual awareness. Vision Research. 2000;40(10-12):1507-21.

87.   Angelucci A, Levitt JB, Walton EJ, Hupé JM, Bullier J, Lund JS. Circuits for local and global signal integration in primary visual cortex. J Neurosci. 2002;22:8633-46.

88.   Shushruth S, Ichida JM, Levitt JB, Angelucci A. Comparison of spatial summation properties of neurons in macaque V1 and V2. Journal of Neurophysiology. 2009;102(4):2069-83.

89.   Nurminen L, Angelucci A. Multiple components of surround modulation in primary visual cortex: multiple neural circuits with multiple functions? . Vision Research. 2014;104:47-56.

90.   Angelucci A, Bijanzadeh M, Nurminen L, Federer F, Merlin S, Bressloff PC. Circuits and Mechanisms for Surround Modulation in Visual Corte. Annual Review of Neuroscience. 2017;40:425-51.

91.   Roland PE, Hanazawa A, Undeman C, .Eriksson D, Tompa T, Nakamura H, et al. Cortical feedback depolarization waves: A mechanism of top-down influence on early visual areas. Proceedings of the National Academy of Sciences of the USA. 2006;103(33):12586-91.

92.   Callaway EM, Katz LC. Emergence and refinement of clustered horizontal connections in cat striate cortex. The Journal of Neuroscience. 1990;10(4):1134-53.

93.   Angelucci A, Bressloff PC. The contribution of feedforward, lateral and feedback connections to the classical receptive field center and extra-classical receptive field surround of primate V1 neurons. Progress in Brain Research. 2006;154:93-121.

94.   Polat U, Sagi D. The architecture of perceptual spatial interactions. Vision Research. 1994;34:73-8.

95.   Polat U, Sagi D. Spatial interactions in human vision, from near to far via experience-dependent cascades of connections. Proceeding of the National Academy of Sciences of the USA. 1994;91:1206-9.

96.   Polat U, Tyler CW. What pattern the eye sees best. Vision Research. 1999;39(5):887-95.

97.   Bullier J, Hupe JM, James AC, Girard P. The role of feedback connections in shaping the responses of visual cortical neurons. Progress in Brain Research. 2001;134:193-204.

98.   Friston KJ, Parr T, de Vries B. The graphical brain: belief propagation and active inference. Network Neuroscience. 2017;1(4):381-414.

99.   Benvenuti G, Chemla S, Boonman A, Perrinet L, Masson G, Chavane F. Anticipatory responses along motion trajectories in awake monkey area V1. 2020. bioRxiv 2020.03.26.010017.

100. Tanifuji M, Sugiyama T, Murase K. Horizontal propagation of excitation in rat visual cortical slices revealed by optical imaging. Science. 1994;266(5187):1057-9.

101. Gilad A, Meirovithz E, Leshem A, Arieli A, Slovin H. Collinear stimuli induce local and cross-areal coherence in the visual cortex of behaving monkeys. PLoS ONE. 2012;7(11):e49391.

102. Frégnac Y. Reading out the synaptic echoes of low-level perception in V1. Lecture Notes in Computer Science. 2012;7583:486-95.

103. Rao RPN, Ballard DH. Predictive Coding in the Visual Cortex: A Functional Interpretation of Some Extra-Classical Receptive Field Effects. Nature Neuroscience. 1999;2:79-87.

104. Martin AB, Von der Heydt R. Spike synchrony reveals emergence of proto-objects in visual corte. The Journal of Neuroscience. 2015;35:6860-70.

105. Kaplan BA, Lansner A, Masson GS, Perrinet LU. Anisotropic connectivity implements motion-based prediction in a spiking neural network. Frontiers in Computational Neuroscience. 2013;7:112

106. Cowey A, Rolls ET. Human cortical magnification factor and its relation to visual acuity. Experimental Brain Research. 1974;21:447-54.

107. Tusa RJ, Palmer LA, Rosenquist AC. The retinotopic organization of area 17 (striate cortex) in the cat. Journal of Comparative Neurology. 1978;177(2):213-35.

108. Hirsch JA, Gilbert CD. Synaptic physiology of horizontal connections in the cat's visual cortex. The Journal of Neuroscience. 1991;11(6):1800-9.

109. Deco G, Roland P. The role of multi-area interactions for the computation of apparent motion. Neuroimage. 2010;51:1018-26.

110. Veltz R. Interplay between synaptic delays and propagation delays in neural field equation. SIAM Journal on Applied Dynamical Systems. 2013;12(3):1566-612.

111. Chisum HJ, Mooser F, Fitzpatrick D. Emergent properties of layer 2/3 neurons reflect the collinear arrangement of horizontal connections in tree shrew visual corte. The Journal of Neuroscience. 2003;23(7):2947-60.

112. Muckli L, Kohler A, Kriegeskorte N, Singe rW. Primary visual cortex activity along the apparent-motion trace reflects illusory perception. PLOS Biology. 2005;3(8):e265.

113. Frégnac Y. The "dynamic association field" of visual cortex neurons: a fit between natural perception and eye-movements? . American Society for Neuroscience Symposium "Mechanisms and Functions of Visual Signals from Beyond the Classical Receptive Field in Primary Visual Cortex"; San Diego2007.

114. Frégnac Y. Searching for a fit between the silent surround of V1 receptive fields and eye-movements. Journal of Vision. 2013;13(9):1380.

115. Tatler BW, Wade NJ, Kwan H, Findlay JM, Velichkovsky BM. Yarbus, eye movements, and vision. Iperception. 2010;1(1):7-27.

1532. **Abbreviations**

1533.   AM: Apparent motion

1534.   ASHP: Apparent Speed of Horizontal Propagation

1535.   CP: Centripetal

1536.   CF: Centrifugal

1537.   CROSS-axis : cross-oriented with the motion axis

1538.   CROSS-RF : cross-oriented (orthogonal) with the RF orientation preference

1539.   FF: Feedforward

1540.   GP: Gabor Patch

1541.   DN: dense noise

1542.   ISO-axis: collinear or parallel to the motion axis

1543.   ISO-RF: collinear or parallel to the RF orientation preference

1544.   MDF: Minimal Discharge Field

1545.   NHP: Non-Human Primate

1546.   RF: Receptive Field

1547.   RND: Randomized

1548.   SRF: Subthreshold synaptic Receptive Field

1549.   SN: sparse noise

1550.   V1: Area 17 in the cat, Primary Visual Cortex

1551.   WPSR: Wilcoxon paired signed rank statistical tests

1552. **Conflict of Interest Statement :**

1553.   The authors declare no competing financial interests.

1554. **Funding :**

1555.   This work was performed at the UNIC (Unit of Neuroscience, Information and Complexity), CNRS,

1556.   UPR CNRS 3293, Gif-sur-Yvette, 91198 France. It was supported by the Centre National de la

1557.   Recherche Scientifique (CNRS) and grants to Y.F. from the Agence Nationale de la Recherche (ANR-

1558.   Horizontal-V1 (ANR-17-CE37-0006), the Paris-Saclay IDEX (NeuroSaclay and i-Code) and the FET-

1559.   Proactive Grant Agreement No 269921 (BrainScales). This project has also received postdoctoral

1560.   fellowships for X.T. under the Marie Skłodowska-Curie grant agreements No 659593 (Proaction

1561.   Perception) and No 302215 (Brain Percepts). B.L.B and Y. P. were supported by The Human Brain

1562.   Project and the ANR-Horizontal-V1.

1563.

1564. *Author contributions :*

1565.   Y.F. conceived of the study.

1566.   Y.F., P.B., X.T., B.L.B., C.D., M.P., Y.P. and C.M. designed the experiments.

1567.   P.B. and M.P ("cardinal protocol") and B.L.B. , M.P., X.T. and C.D. ("radial protocol") collected the

1568.   intracellular data. P.B. ("cardinal" protocol) and B.L.B., C.D. and X.T. ("radial" protocol) implemented

1569.   the data analysis. C.D. and M.P. designed the figures.

1570.   Y.F. and B.L.B. wrote the manuscript.

Figure 1

Figure 2 :

Figure 3 :

A Single Cell

B Population (n=13)

Figure 4 :

Centripetal

Centrifugal

8 mV
100 Spk/s
50 ms

Figure 5 :

Figure 6 :

Figure 7 :

Figure 8 :

Figure 9 :

Figure 10 :

Figure 11 :