# Combining CRISPR/Cas9 and brain imaging: from genes to proteins to networks

Sabina Marciano[1], Tudor Mihai Ionescu[1], Ran Sing Saw[1], Rachel Y. Cheong[2], Deniz Kirik[2], Andreas Maurer[1], Bernd J. Pichler[1], Kristina Herfert[1]*

[1]Werner Siemens Imaging Center, Department of Preclinical Imaging and Radiopharmacy, Eberhard Karls University Tuebingen, Germany

[2]Brain Repair and Imaging in Neural Systems, Department of Experimental Medical Science, Lund University, Sweden

*Corresponding author: Prof. Dr. Kristina Herfert (kristina.herfert@med.uni-tuebingen.de)

Address: Roentgenweg 13, 72076 Tuebingen, Germany, Tel: +49 7071 29 87680

**Abstract**

Understanding how functional connections between brain regions are arranged and influenced by genes, in the healthy and diseased states, is a major goal in neuroscience. Functional connectivity alterations are linked to molecular changes in several neurodegenerative disorders and could serve as early markers of pathology. Yet, the underlying molecular signatures driving the functional alterations remain largely unknown. Here, we combine CRISPR/Cas9 gene-editing with *in vivo* positron emission tomography (PET) and functional magnetic resonance imaging (fMRI) to investigate the direct link between genes, proteins, and the brain connectome. The extensive knowledge of the *Slc18a2* gene encoding the vesicular monoamine transporter (VMAT2), involved in the storage and release of dopamine, makes it an excellent basis for studying the gene networks relationships. We edited *Slc18a2* into the substantia nigra pars compacta of adult rats and used *in vivo* molecular imaging, behavioral, histological, and biochemical assessments to characterize the CRISPR/Cas9-mediated VMAT2 knockdown. Simultaneous PET/fMRI was performed to inspect the functional brain adaptations, beyond the predicted dopaminergic changes. Further, [$^{11}$C]flumazenil PET was carried out to investigate the dopamine-GABA interplay. We found a regional increase in postsynaptic dopamine receptor availability, preceded by a reorganization of brain networks that adapt to reduced dopamine transmission states by becoming functionally connected and organized. The hyperconnectivity within and between brain networks spreads from the contralateral thalamus and prefrontal cortical regions to the striata and hippocampi. Additionally, impaired striatal dopamine release reduces GABA-A receptor availability, complementing the increased synchrony and functional connectivity between networks observed at rest. Our study reveals that recruiting different brain networks may be an early response to the dopaminergic dysfunction preceding neuronal cell loss, postsynaptic changes, and motor impairment in neurodegenerative disorders such as Parkinson´s disease. We anticipate our combinatorial approach to be a starting point to investigate the impact of

45   specific genes on brain molecular and functional dynamics, aiding in the identification of early

46   neurobiological markers and promising therapeutic interventions.

50   **Keywords**

51   CRISPR/Cas9, Functional Connectivity, Dopamine, VMAT2, D2 receptor, [$^{11}$C]raclopride,

52   PET/fMRI.

## Introduction

The brain is a network of spatially distributed but functionally and structurally interconnected

regions that exhibit correlated activity over time. They communicate with each other via highly

specialized neuronal connections and are organized in neuronal circuits and networks.

Understanding how functional connections between regions are arranged in the healthy and

diseased brain is therefore of great interest.

Resting-state functional magnetic resonance imaging (rs-fMRI) has enabled neuroscientists to

delineate the level of functional communication between anatomically separated regions [1]. Rs-

fMRI measures the resting-state functional connectivity (rs-FC) at high spatial and temporal

resolutions based on spontaneous fluctuations of the blood oxygen level-dependent (BOLD)

signal at rest, which indirectly detects neuronal activity via hemodynamic coupling [2]. Using rs-

fMRI several brain resting-state networks in humans and rodents have been identified, such as

the default mode and sensorimotor networks (DMN, SMN) [3-9]. Alterations of these networks are

linked to neurological diseases [10, 11], and may serve as early therapeutic and diagnostic

biomarkers. However, the molecular signatures related to the functional alterations in disease

remain largely unknown.

Positron emission tomography (PET) provides a non-invasive tool to indirectly measure molecular

changes in the brain with high specificity and sensitivity. One well-characterized example is the

radioligand [$^{11}$C]raclopride, a widely used D2/D3 receptor antagonist enabling the non-invasive

determination of dopamine release and availability [12-14].

PET in combination with BOLD-fMRI has the great potential to investigate the molecular

substrate of brain functional connectivity (FC), enabling the direct spatial and temporal correlation

of both measurements [15-21]. In this context, we have recently shown that rs-FC is modulated

by intrinsic serotonin transporter and D2/3 receptor occupancy in rats [22].

Insights into functional brain circuits and their relationships to individual phenotypes can be

gained by genetic manipulations of neuronal subtypes [23]. Genome-engineering methodologies

based on clustered regularly interspaced short palindromic repeats (CRISPR)/associated RNA-

guided endonuclease (Cas9) represent a promising approach to unveil the influence of genes on

brain circuits. CRISPR/Cas9 has enabled researchers to interrogate the mammalian DNA in a

precise yet simple manner [24, 25] in several species [26-31], by editing single or multiple

genomic loci *in vitro* and *in vivo* [25, 32, 33]. However, one great hurdle is the brain delivery,

which must comply with effective nuclear access, while minimizing immunogenic reactions and

off-target editing [34]. Despite these limitations, the potential of CRISPR/Cas9 is continuously

expanding with novel nuclease variants being exploited [35-37]. Derived from *Staphylococcus*

*aureus*, SaCas9 overcomes the packaging constraints of adeno-associated viral vectors (AAVs),

allowing efficient CRISPR/Cas9 brain transfer [38-42].

Here, we use an AAV-based CRISPR/SaCas9 gene-editing approach to knock down the *Slc18a2*

gene encoding the vesicular monoamine transporter 2 (VMAT2), a key protein involved in the

storage and release of dopamine in the brain [43]. The extensive knowledge on *Slc18a2* makes it

an excellent basis for studying the gene-networks relationships. We characterize the VMAT2-

mediated dopamine signaling using *in vivo* molecular imaging, behavioral, histological, and

biochemical assessments. We investigate the impact of impaired VMAT2-dependent dopamine

neurotransmission on the DMN and SMN using a simultaneous [$^{11}$C]raclopride-PET/fMRI

protocol. Further, we dig into the dopamine-GABA interplay using [$^{11}$C]flumazenil PET. Our

results reveal that CRISPR/SaCas9-induced synaptic dysfunction prompts early network

changes, preceding motor and molecular alterations, including a regional increase in postsynaptic

dopamine receptor availability. We identify a pattern of asymmetric hyperconnectivity, and

internetwork synchronization, spreading from the contralateral thalamus (SMN), and prefrontal

cortical regions (DMN) to the striata and hippocampi, complemented by a reduced GABA-A

receptor availability.

Our findings illustrate the ability of the brain to recruit different brain networks and functionally

compensate for the dopaminergic dysfunction prior to neuronal cell loss, postsynaptic changes,

and motor impairment.

**Results**

*In vitro* **validation of CRISPR/SaCas9-induced VMAT2 knockdown in rat primary cortical**

**neurons**

To evaluate the efficiency of the AAV-based CRISPR/SaCas9 VMAT2 knockdown in rat primary

neurons, we designed AAV-SaCas9 and AAV-sgRNA targeting the first exon of the bacterial *lacZ*

gene (control) or the second exon of the *Slc18a2* gene (Fig. 1a) (sgRNAs sequences are

reported in Table 1). Seven days post-transduction, the protein expression level and mutation

rate of the harvested genomic DNA were inspected by immunofluorescence and surveyor assay

(Fig. 1b). Immunofluorescence indicated a clear reduction of VMAT2 protein expression in

neurons transduced with AAV-SaCas9 and AAV-sgRNA-*Slc18a2* (Fig. 1c). We observed 20%

editing for the digested DNA from neurons transduced with vectors for SaCas9 and sgRNA-

*Slc18a2* (Supplementary Fig.1).

**CRISPR/SaCas9-induced VMAT2 knockdown elicits postsynaptic changes but no nerve**

**terminal loss or neuroinflammation in the adult rat brain**

To test the *in vivo* efficiency of the CRISPR/SaCas9 gene-editing, we expressed SaCas9 and

sgRNA targeting *Slc18a2* to knock down the VMAT2, or targeting *lacZ* as control, by AAV-

mediated gene transfer into the right substantia nigra pars compacta (SNc). DPBS was injected

into the left SNc. [$^{11}$C]Dihydrotetrabenazine (DTBZ) PET imaging was performed 8 – 10 weeks

post-injection to quantify VMAT2 expression in the striatum (Fig. 2a).

[$^{11}$C]DTBZ binding was decreased by 30% in the right striatum of rats where the VMAT2 was

knocked down in comparison to the contralateral striatum. No changes of [$^{11}$C]DTBZ binding were

observed in the contralateral striatum, as [$^{11}$C]DTBZ binding did not differ between the left

striatum of rats injected with sgRNA targeting *lacZ* and rats injected with sgRNA targeting

*Slc18a2* (Fig. 2b,c).

We further evaluated changes of dopamine availability in the striatum using [$^{11}$C]raclopride, which

competes with dopamine for the same binding site at the D2 receptor (D2R) [13]. After 12 –

14 weeks following CRISPR/SaCas9-induced VMAT2 knockdown in nigrostriatal neurons, we

observed 17% increased binding of [$^{11}$C]raclopride in the right striatum of VMAT2 knockdown rats

and no changes in control rats (Fig. 2b,c), indicating a reduction of synaptic dopamine levels

and/or compensatory changes of D2R expression at postsynaptic medium spiny neurons. A

larger VMAT2 knockdown led to lower dopamine levels in the striatum and thus to higher D2R

binding (Fig. 2d). To explore the threshold at which the observed postsynaptic changes occur, we

calculated the [$^{11}$C]raclopride/[$^{11}$C]DTBZ binding ratio for the right and left striatum. The ratio

remained close to 1 in the DPBS-injected striatum and control rats, indicating no substantial

difference between the two hemispheres. In contrast, VMAT2 knockdown rats displayed large

[$^{11}$C]raclopride binding changes when the level of VMAT2 knockdown was ~ 20%. From this

point, a prominent increase in D2R binding was observed in the right striatum (Fig. 2e).

Therefore, this threshold was set to split the rats into *mild* (< 20%) and *moderate* (≥ 20%).

Notably, [$^{11}$C]raclopride PET imaging was able to discriminate between different degrees of

synaptic dysfunction, classified from [$^{11}$C]DTBZ binding changes (Fig. 2f).

We inspected the integrity of dopaminergic nerve terminals and the occurrence of

neuroinflammation in the striatum after the CRISPR/SaCas9-induced VMAT2 knockdown.

[$^{11}$C]methylphenidate PET imaging of the dopamine transporter and [$^{18}$F]GE-180 PET imaging of

the translocator protein, which is overexpressed on activated microglia, was performed.

CRISPR/SaCas9-induced VMAT2 knockdown did neither alter [$^{11}$C]methylphenidate binding, nor

[$^{18}$F]GE-180 uptake (Fig. 2b,c).

**CRISPR/SaCas9-induced VMAT2 knockdown impairs motor function**

To explore the motor consequences of the CRISPR/SaCas9-induced VMAT2 knockdown, we

performed several behavioral tests (Fig. 3a).

We observed a reduction in the locomotor activity of VMAT2 knockdown rats in the open field test

(Fig. 3a), but no correlation to VMAT2 expression changes ($\Delta$ [$^{11}$C]DTBZ binding), or dopamine

availability ($\Delta$ [$^{11}$C]raclopride binding) (Supplementary Fig. 2a,b).

Next, we evaluated the forelimb akinesia using the cylinder test. VMAT2 knockdown rats

displayed a preference for the right forepaw, while control rats equivalently used their right and

left forepaw (Fig. 3c). Paw use alterations correlated highly with VMAT2 knockdown ($\Delta$ [$^{11}$C]DTBZ

binding), and dopamine availability ($\Delta$ [$^{11}$C]raclopride binding) (Fig. 3d,e).

To further examine differences in motor function, coordination, and balance, rats underwent the

beam walk test. VMAT2 knockdown rats stumbled with higher frequency to the left side, while

control rats displayed equal chances to slip in each direction (Fig. 3f). However, no correlations

between gait alterations and VMAT2 knockdown ($\Delta$ [$^{11}$C]DTBZ binding), or dopamine availability

($\Delta$ [$^{11}$C]raclopride binding) were found (Supplementary Fig. 2c,d).

As previous studies suggest that body weight changes reflect striatal dopamine depletion [44], we

inspected the impact of the VMAT2 knockdown on the rats' body weight gain. VMAT2 knockdown

rats exhibited a 30% reduction in their gained weight over a period of 14 weeks, compared with

controls (Fig. 3g). Body weight gain correlated with changes in VMAT2 expression ($\Delta$ [$^{11}$C]DTBZ

binding), and dopamine availability ($\Delta$ [$^{11}$C]raclopride binding) (Fig. 3h,i).

To assess the rotational behavior, we performed the rotameter test with and without apomorphine

administration. In the spontaneous rotation test, VMAT2 knockdown rats displayed a higher

number of ipsilateral net turns compared with control rats (Fig. 3j). The number of turns did not

correlate with VMAT2 expression changes ($\Delta$ [$^{11}$C]DTBZ binding), and changes in dopamine

availability ($\Delta$ [$^{11}$C]raclopride binding) (Supplementary Fig. 2e,f). Apomorphine-induced rotations

to the contralateral side were higher in VMAT2 knockdown rats compared with control rats (Fig.

3k), and correlated with changes in VMAT2 expression ($\Delta$ [$^{11}$C]DTBZ binding) and dopamine

availability ($\Delta$ [$^{11}$C]raclopride binding) (Fig. 3l,m).

*Ex vivo* **validation of the CRISPR/SaCas9-induced VMAT2 knockdown**

Using immunofluorescence, we confirmed the concomitant expression of SaCas9 and *Slc18a2*–

targeting sgRNA 19 weeks post-transduction, and a corresponding decrease of VMAT2

expression in the SNc of the VMAT2 knockdown group (Fig. 4b).

Immunohistochemistry revealed no changes in tyrosine hydroxylase (TH) expression levels in

striatum and SN in both groups (Fig. 4c,d), and confirmed the reduction of VMAT2 expression in

the right striatum and SN in the knockdown group (Fig. 4e).

Biochemical analysis showed a large reduction of dopamine, paralleled by an increased ratio of

metabolites (DOPAC, HVA) to dopamine, in the right striatum of VMAT2 knockdown rats (Fig.

4f,g). The reduced dopamine content correlated with the *in vivo* VMAT2 expression ($\Delta$ [$^{11}$C]DTBZ

BP$_{ND}$) and postsynaptic changes ($\Delta$ [$^{11}$C]RAC BP$_{ND}$) (Fig. 4h,i). Additionally, serotonin was

unchanged in the striata of VMAT2 knockdown and control rats, suggesting dopamine

nigrostriatal pathway specificity (Fig. 4j) (Metabolites' and neurotransmitters' striatal levels are

reported in Table 3).

**Increased resting-state functional connectivity after CRISPR/SaCas9-induced VMAT2**

**knockdown**

As multiple lines of evidence suggest a broader role of dopamine in the dynamic reconfiguration

of brain networks [17, 45, 46], we next investigated the impact of unilateral dopamine depletion

on brain rs-FC. A second cohort of rats underwent longitudinal simultaneous [$^{11}$C]raclopride-

PET/BOLD-fMRI scans at baseline and 8 - 14 weeks after CRISPR/SaCas9-induced VMAT2

knockdown (Fig. 5a). [$^{11}$C]DTBZ PET scans and behavioral analysis confirmed previous findings

in the first cohort, that is, an efficient depletion of the VMAT2 (20% decrease of [$^{11}$C]DTBZ

binding) (Supplementary Fig. 3a), paralleled by motor disturbances in the cylinder test

(Supplementary Fig. 3e-g). In line with the findings of cohort 1, dopamine availability was

decreased (10% increase in [$^{11}$C]raclopride binding) and correlated to the extent of the VMAT2

knockdown (Supplementary Fig. 3b,c). An increase in D2R binding was observed in the right

striatum when the level of VMAT2 knockdown reached ~ 20% (Supplementary Fig. 3d), enabling

a subdivision into *mild* (< 20%) and *moderate* rats (≥ 20%).

We next assessed the occurrence of rs-FC changes in DMN and SMN. Our analysis focused on

identifying early biomarkers of *mild* dysfunction and patterns of spreading of synaptic dysfunction.

Figure 5b,c illustrates intraregional rs-FC group-level correlation matrices at baseline and after

VMAT2 knockdown in *mild* and *moderate* rats for the DMN and SMN, respectively. We observed

within-network rs-FC changes in rats with *moderate* VMAT2 knockdown, in both DMN and SMN.

Rats of the *mild* knockdown group revealed rs-FC changes up to 20%, in prefrontal cortical

regions of the DMN, and between the left thalamus (Th) and somatosensory cortex (SC) in the

SMN. However, these data need to be carefully interpreted as they did not survive a more

stringent *P* value selection (*$P$< 0.01) (Supplementary Fig.4a,b) (*P* values are reported in

Supplementary Tables 1,3).

Rats with *moderate* VMAT2 knockdown exhibited a 60% increase in rs-FC within the right medial

prefrontal cortex (mPFC) and the right and left hippocampus (Hipp) (Fig. 5b) (*P* values are

reported in Supplementary Table 2).

FC increase between the left thalamus and somatosensory cortex in the SMN doubled to 34% in

rats with *moderate* VMAT2 knockdown and extended throughout the left and right thalamus and

striatum (STR), respectively (Fig. 5c) (*P* values are reported in Supplementary Table 4).

Moreover, we inspected rs-FC changes between the DMN and SMN at baseline and after the

CRISPR/SaCas9-induced VMAT2 knockdown.

Figure 5d illustrates internetwork rs-FC correlation matrices in rats with *mild* (left panel) and

*moderate* (right panel) VMAT2 knockdown. Brain graphs display the nodes and edges (raw

values) that demonstrated internetwork rs-FC changes to baseline (%). Strikingly, large

alterations between DMN and SMN were already observable in the *mild* VMAT2 knockdown

group. Rats presented opposite rs-FC changes between regions of the anterior/posterior DMN

and the SMN, compared with baseline. A 30 to 60% increase in rs-FC was observed between

regions of the anterior DMN and the SMN. Specifically, rs-FC increased between the right

orbitofrontal cortex (OFC) and striatum bilaterally and the contralateral somatosensory cortex,

and between the contralateral orbitofrontal cortex and striatum. Instead, a 20% decrease in rs-FC

was found between regions of the posterior DMN and the SMN. Specifically, rs-FC decreased

between the left retrosplenial cortex (RSC) and right somatosensory cortex (Fig. 5d, left panel) (*P*

values are reported in Supplementary Table 5).

Rats with a *moderate* VMAT2 knockdown presented increased rs-FC between regions of the

anterior/posterior DMN and the SMN, compared with baseline. Of particular note, internetwork rs-

FC changes were not found between the regions of the posterior DMN and the SMN that showed

decreased rs-FC in rats with *mild* VMAT2 knockdown. Moreover, between-network rs-FC

increase extended to other regions. A 60 to 80% increase in rs-FC was found between the medial

prefrontal cortex and the right striatum, and the motor (MC) and somatosensory cortex bilaterally.

FC increased by more than 20% between the hippocampi and contralateral somatosensory

cortex. (Fig. 5d, right panel) (*P* values are reported in Supplementary Table 6). Notably, between-

network rs-FC changes did not involve the thalamus, which connectivity was however altered

within the SMN.

Further, we examined how the rs-FC changes to baseline correlated between the *mild* and

*moderate* groups (Supplementary Fig. 5a). Group level intraregional and internetwork rs-FC

changes to baseline (%) correlated linearly between the two groups (Supplementary Fig. 5b-d).

Node correlation analysis indicated a linear increase in the magnitude of the rs-FC changes to

baseline in the hippocampi (Supplementary Fig. 5e), cingulate cortices (Supplementary Fig. 5f),

and contralateral, but not ipsilateral, thalamus (Supplementary Fig. 5g). Our data suggest a

similarity in the pattern of the intraregional and internetwork rs-FC changes between *mild* and

*moderate* VMAT2 knockdown rats and a linear relationship between the magnitude of the rs-FC

changes.

To complement the results of the intraregional and internetwork rs-FC, we evaluated changes in

regional mean connection distances. Network-wise graph theoretical analysis on node level was

paralleled by whole-brain connection-wise analysis to identify the nodes that were significantly

altered in rats with *mild* and *moderate* VMAT2 knockdown, compared with baseline, for the DMN

(Supplementary Fig. 6a,b) and SMN (Supplementary Fig. 6c,d). Briefly, the network organization

did not change in rats with *mild* VMAT2 knockdown, compared with baseline, as changes in the

global mean connection distance were not found in DMN (Supplementary Fig. 6a) nor SMN

(Supplementary Fig. 6c). Interestingly, in rats with *moderate* VMAT2 knockdown network

organization changes did not influence regions of the DMN (Supplementary Fig. 6b), but occurred

in the contralateral striatum and thalamus (Supplementary Fig. 6d) (*P* values are reported in

Supplementary Table 7).

Collectively, rs-FC results highlight lateralized effects in the SMN, as opposed to the symmetric

recruitment of DMN regions.

**CRISPR/SaCas9-induced VMAT2 knockdown alters GABA signaling**

Besides dopamine, dopaminergic neurons co-release GABA via the VMAT2 [47, 48]. To

investigate if GABA neurotransmission is altered following the VMAT2 knockdown, we performed

additional [$^{11}$C]flumazenil PET scans 10 – 12 weeks after the CRISPR/Cas9-editing, and

quantified the GABA-A binding in regions of the DMN and SMN (Fig. 6). In *mild* VMAT2

knockdown rats, we observed a decrease of [11C]flumazenil binding in the ipsilateral parietal

cortex (PaC) (14%), hippocampus (4%), and somatosensory cortex (9%) (Fig. 6a,b). In *moderate*

VMAT2 knockdown rats we observed a decrease of [11C]flumazenil binding in the ipsilateral

parietal cortex (11%), and somatosensory cortex (6%) (Fig. 6a,c). Our data indicate that

[11C]flumazenil binding was altered regardless of the VMAT2 knockdown extent. This was further

evidenced by the lack of correlation between [11C]DTBZ and [11C]flumazenil binding changes in

the target regions (linear regression data not shown).

**Discussion**

Here, we show the potential of combining CRISPR/Cas9 gene-editing with molecular and

functional brain imaging to identify early adaptations of brain circuits in response to targeted gene

and protein modulations. Using CRISPR/SaCas9, we knocked down the *Slc18a2* gene, encoding

the VMAT2, which plays a key role in the storage and release of dopamine in response to

neuronal activity [43]. The CRISPR-mediated knockdown allowed us to investigate the VMAT2-

dependent dopamine signaling in the striatum, while structurally preserving neuronal integrity.

[18F]GE-180 results suggest that glial activation is not the source of dopaminergic synaptic

dysfunction, and exclude the occurrence of inflammatory responses arising from the surgical

procedure and the chosen AAV-serotype, which could influence our readout, in line with recent

reports [49]. Our data reveal that the targeted gene knockdown in the SNc leads to an expected

reduction of dopamine release in the striatum, paralleled by [11C]raclopride binding changes. It is

conceivable that the observed postsynaptic changes are the result of an increase in binding due

to the reduced dopamine concentration in the striatum. Yet, several studies have revised this

notion [50, 51]. In this regard, our results of the drug-evoked rotational behavior are in better

agreement with a D2R compensatory upregulation. Indeed, supersensitivity to apomorphine in

rats with nigrostriatal lesions or VMAT2 knockout is accompanied by an increase in striatal D2R

binding sites, but no change in affinity [52-57]. Further, the observed striatal increase in

[$^{11}$C]raclopride binding is independent of presynaptic nerve terminal loss and occurs in response

to a ~ 20% decrease of [$^{11}$C]DTBZ binding. This confirms, in line with our [$^{11}$C]methylphenidate

results, that [$^{11}$C]raclopride can be used to delineate postsynaptic changes in the absence of

dopamine transporter-mediated compensation, triggered by neuronal loss. Accordingly, increased

[$^{11}$C]raclopride binding is observed in the early but not later stages of Parkinson`s disease [58],

characterized by severe neuronal cell loss, and dopamine transporter changes (> 50%) [59].

Consistently, rats with severe denervation (> 75%) present earlier mitigation by the dopamine

transporter, followed by D2R binding changes [60]. Hence, with our method, it is feasible to study

the consequences of synaptic dopamine dysfunction without compensations triggered by

neuronal cell loss. Moreover, the observed [$^{11}$C]raclopride and [$^{11}$C]DTBZ correlations to the

motor behavior highlight that [$^{11}$C]raclopride binding remains at control levels as long as synaptic

dopamine levels are sufficient to maintain adequate motor function. Motor disturbances strongly

correlate to pre and postsynaptic changes if movements of the forelimbs, but not whole-body, are

considered, in agreement with earlier observations in dopamine-depleted rats [44]. Depletion of

VMAT2 resulted in reduced dopamine tissue levels in the ipsilateral striatum, nicely merging with

the *in vivo* data. Moreover, metabolite analysis suggested that due to the lack of VMAT2-

mediated storage in presynaptic vesicles, dopamine is quickly converted. The increased

metabolism might as well be a possible compensatory mechanism consequent to the VMAT2

knockdown, reflecting actions that residual nigrostriatal neurons undertake to maintain dopamine

homeostasis, as already speculated by others [53, 61].

To elucidate the role of VMAT2 in locomotion, reward, and Parkinson`s disease, several

investigators have deleted its coding gene in mice [62-65]. Besides the costly and time-

consuming breeding, the gene knockout was not selective for dopaminergic neurons, resulting in

the appearance of anxiety and depressive behavior phenotypes [66]. CRISPR/Cas9-editing

overcomes these limitations, allowing gene-editing in adult and aged animals and avoiding

compensatory changes occurring at early developmental stages. Since its discovery, only two

studies have successfully applied CRISPR/Cas9 in the rat brain [40, 67], where gene-editing has

been difficult to adapt. Rats are particularly advantageous for imaging studies due to the larger

brain size and limited spatial resolution and sensitivity of preclinical scanners [68]. Here, by

inducing a *mild* to *moderate* gene knockdown, we could investigate early to late resting-state

brain network adaptations prompted by presynaptic dysfunction. We show that the selective

impairment of presynaptic dopamine storage and release is followed by rs-FC alterations within

and between the DMN and SMN. Our results confirm previous findings that the DMN, associated

with ideation and mind wandering [69], and the SMN, involved in sensory processing and motor

function [70], do not function in isolation from each other, but rather synchronize [71]. The

observed internetwork synchronization may reflect compensatory brain reorganization, as already

speculated by others [72, 73]. We also identified enhanced intranetwork rs-FC in the DMN and

SMN. Rs-FC changes were observed in prefrontal cortical regions, hippocampus, thalamus, and

striatum. Our data parallel previous findings of cortico-striato-thalamic hyperconnectivity in

decreased dopamine transmission states [74-77]. In line, increased synchronous neural

oscillatory activity and functional coupling in the basal ganglia and its associated networks have

been observed in Parkinson`s disease [78-83]. The increase of cortico-striatal FC could in part be

due to dysfunctions of multiple tonic inhibitory gate actions of D2R [84]. Increased FC across the

thalamus and prefrontal cortex has been reported in drug-treated Parkinson`s disease patients

[85, 86], potentially indicating functional compensation, as the brain recruits additional anatomical

areas to aid in restoring cognitive processes. This might as well explain the engagement of the

hippocampus, functionally connected with DMN cortical regions [87, 88]. In this regard, research

has shown that the hyperconnectivity of brain circuits is a common response to neurological

dysfunction, and may reflect a protective mechanism to maintain normal brain functioning [89].

Such a mechanism has been proposed in Parkinson`s disease, mild cognitive impairment, and

Alzheimer's disease [90-93]. Collectively, our findings support this model and indicate a

reorganization of brain networks that adapt to the synaptic dysfunction through enhanced

interregional synchrony. Recruiting alternative brain regions may be an early response to the

dysfunction preceding neuronal cell loss and motor impairment. Interestingly, brain connectome

adaptations occurred symmetrically in the DMN but were more weighted towards the contralateral

hemisphere in the SMN.

Besides dopamine, dopaminergic neurons co-release GABA via the VMAT2. This hints towards

reduced GABA following the VMAT2 knockdown, and consequent imbalance in downstream

striatal projection neurons of the direct and indirect pathway [47, 48, 94]. GABA regulates the

inhibitory neurotransmission in various brain areas through GABA-A receptors [95]. We

hypothesize that decreased binding to the GABA-A receptors may be consistent with loss of

inhibitory tone in multiple cortical areas, reflected by our [$^{11}$C]flumazenil PET data, resulting in

increased localized brain connectivity, reflected in the fMRI signal. The suppression of the

GABAergic feedback circuit, mediated by D2R and external globus pallidus neurons [84],

complements the observed elevation in neuronal synchronicity. Our postulation is in line with

earlier findings of an inverse correlation of GABA with rs-FC in DMN [96] and with a putative role

of loss of inhibitory tone in hyperconnectivity [97]. Interestingly, our [$^{11}$C]flumazenil PET data did

not reveal changes in GABA-A expression in the striatum. Instead, we observed a significant

decrease in the ipsilateral hemisphere of several cortical regions in both *mild* and *moderate* rats,

supporting previous findings of GABA modulation of the internetwork FC [98]. Studies report

downregulated inhibitory neurotransmission in Parkinson`s disease, where gene expression of

GABAergic markers is low in the frontal cortex [99, 100]. Further, inverse correlations between

[$^{11}$C]flumazenil binding and gait disturbances [101], and between GABA concentration in the

motor cortex and disease severity have been reported [102]. Moreover, our [$^{11}$C]flumazenil PET

results suggest that GABA neurotransmission is disturbed already at the *mild* stage, indicating its

potential role as an early biomarker of dopaminergic presynaptic dysfunction. Of note, the

observed changes in GABA-A binding might indicate only an apparent decrease in binding due to

the reduced GABA availability, or reflect compensatory adaptation on the contralateral

hemisphere. Future research should elucidate these aspects and also perform further

investigations on glutamate, due to the crucial role of the excitatory/inhibitory-imbalance in

several psychiatric disorders [103].

**Limitations and general remarks**

We knocked down VMAT2 in dopaminergic projection neurons from the SNc to the dorsal

striatum. To achieve selective targeting of this neuronal subtype, rat Cre driver lines have been

developed [67, 104]. Although all monoamine-releasing neurons express VMAT2, in contrast to

other brain regions, SNc neurons are predominantly dopaminergic [105]. Thus, even though we

used wild-type rats, we can largely dismiss effects on other monoaminergic neurons, as also

indicated by biochemical analysis of serotonin striatal levels.

Despite its limited off-target editing [38], undesired targeting of SaCas9 on other genes cannot be

fully excluded. Nevertheless, off-target candidates with up to 4 mismatches were screened in the

whole genome of *Rattus norvegicus* (http://www.rgenome.net/cas-offinder/), consistently with

past reports [40]. To the best of our knowledge, the off-target matches (*Ndrg1*, *RGD1305938*,

*Btn2a2*, *AABR07042293.2)* have no effects on VMAT2 function, being involved in cell

differentiation, T-cell regulation, and mRNA processing, respectively

(https://www.ncbi.nlm.nih.gov/IEB/Research/Acembly/index.html).

Another limitation of the study is the relatively small sample size, related to the complex and high-

cost procedures involved in the *in vivo* imaging measurements. In addition to this, the intrinsically

high intersubject variability in rs-fMRI, and differences in knockdown efficiency contributed to

significant variance in our cohorts. Nevertheless, the variability of gene-editing efficiency was in

line with previous *in vivo* brain studies [40, 41].

**Conclusions**

This work encourages the combinatorial use of CRISPR/Cas9 and molecular and functional *in*

*vivo* brain imaging to achieve selective modulation of genes and understand the related functional

adaptations in brain networks, beyond the targeted circuitry.

We anticipate our approach to be a starting point to shed the light on the function of specific

genes and their encoded proteins on whole-brain connectivity, useful to understand the cellular

basis of functional changes, identify early neurobiological markers, and promising therapeutic

interventions.

## Methods

### Animals

Female wild-type Long Evans rats (224 ± 30 g, n = 57) (Charles River Laboratories, Sulzfeld, Germany) were kept on a 12 h day-night cycle at a room temperature of 22 °C and 40 - 60% humidity. Animals received a standard diet and tap water *ad libitum* before and during the experimental period. All animal experiments were performed according to the German Animal Welfare Act and were approved by the local ethical authorities, permit numbers R15/19M, R4/20G.

### Viral vectors

SgRNAs targeting the second exon of the *Slc18a2* gene and the first exon of the *lacZ* gene were designed based on the PAM sequence of SaCas9 (NNGRRT) (Table 1). SgRNAs were cloned into an AAV-PHP.EB expression vector containing a GFP reporter sequence, driven by the CMV promoter, for the identification of transduced neurons. A second AAV-PHP.EB construct was produced to express SaCas9, flanked by two nuclear localization sequences (NLS) to allow its translocation into the nuclei. The vector expressed the nuclease via the CAG promoter and contains three HA-tags to visualize the targeted neurons (Fig. 1a). Cloning of the sgRNAs, plasmid construction, as well as the production of concentrated and purified AAV-PHP.EB vectors, delivered at a concentration of $10^{13}$ gc/mL, were carried by SignaGen Laboratories (Frederick, Maryland, USA). A European patent application has been filed for the AAV-PHP.EB vectors and is currently pending.

The genomic mutation rate was assessed using a different set of AAVs with AAV2/1 serotype, for a conditional design, kindly provided by Matthias Heidenreich (prev. Zhang lab, Broad Institute of MIT and Harvard, Cambridge, USA).

**Rat primary cortical neuron culture**

Primary cortical neurons were obtained from rat embryos of a pregnant Sprague Dawley rat from

embryonic day 18 (E18) (Charles River, Sulzfeld, Germany). Embryos were decapitated and

quickly removed from the mother rat. Cortical dissection was performed in ice-cold HBSS

(100 mL 10 × HBSS, 870 mL $dH_2O$, 3.3% 0.3 M HEPES pH 7.3 and 1% pen/strep)

(LifeTechnologies, Massachusetts, USA). The obtained tissue was washed three times with

10 mL ice-cold HBSS PhenolRed-free (LifeTechnologies, Massachusetts, USA) and then

digested at 37 °C for 20 min in 8 mL HBSS with 2.5% trypsin (LifeTechnologies, Massachusetts,

USA). Cortices were washed 3 times with 10 mL HBSS containing 3.7% FBS, and then gently

triturated in 2 mL HBSS. For the maintenance, neurons were plated on poly-D-lysine-coated 24

well plates (BD Biosciences, Heidelberg, Germany) or coverslips (Neuvitro Corporation,

Vancouver, USA) at a density of $16 × 10^4$/well, and cultured in Neurobasal media supplemented

with 2% 1 × B27, 0.25% Glutamax, 0.125% Glutamate and 1% pen/strep (LifeTechnologies,

Massachusetts, USA) for four days. Afterward, for the immunofluorescence, AAVs carrying the

expression of the vectors for the SaCas9 and sgRNAs (1:1 ratio), were added to the culture

medium at 200,000 MOI (AAV-PHP.EB-sgRNA-*lacZ:* $1.7 × 10^{12}$ gc/mL, AVV-PHP.EB-SaCas9:

$1.4 × 10^{12}$ gc/mL, AAV-PHP.EB-sgRNA-*Slc18a2:* $2.1 × 10^{12}$ gc/mL, final viral volume

37.5 µL/well). For the Surveyor assay, conditional AAVs for SaCas9, sgRNA-*Slc18a2*, Cre-

recombinase (AAV2/1, 1:1:0.5 ratio) were used. Neurons were processed 1-week post-viral

treatment (Fig. 1b).

**Surveyor assay**

To estimate the VMAT2 knockdown efficiency of the designed sgRNA *in vitro*, we evaluated the

presence of genetic deletions in rat primary neurons with the Surveyor assay (Surveyor kit,

Integrated DNA Technologies, Coralville, USA). One week following the viral infection, the

genomic DNA was extracted using the QuickExtract DNA Extraction solution (Epicentre,

Madison, USA), according to the manufacturer´s instructions, and was normalized to 100 ng in

dH$_2$O. 18 - 25 nt primers were designed 200 - 400 bp away from either side of the SaCas9 target

site to amplify the loci of interest by touchdown PCR (oligonucleotides used for PCR are provided

in Supplementary Table 8). DNA amplification was performed using 0.5 µL Phusion Polymerase

(LifeTechologies, Massachusetts, USA), as previously reported [106]. A single band product was

visualized on 1.5% agarose gel, isolated, and purified using QIAquick Spin columns (Qiagen,

Hilden, Germany), following the supplier's protocol. 400 ng of the purified PCR product were

mixed with 2 µL Taq DNA polymerase buffer (LifeTechnologies, Massachusetts, USA) to allow

the cross-annealing of the mutated and wild-type sequences. The re-annealing process was

conducted at the following cycling conditions: 10 min at 95 °C, 95 °C to 85 °C at -2 °C/s, hold

1 min, ramp down to 75 °C at -0.3 °C/s, hold 1 min, and so on until 25 °C temperature was

reached. Samples were then stored at 4 °C. This cross-annealing procedure converts the

mutations into mismatch duplexes (heteroduplexes), which can be recognized by performing

nuclease digestion [107]. We digested the annealed products for 20 min at 42 °C using 2.5 µL

MgCl$_2$ (0.15 M), 1 µL Surveyor nuclease, and 1 µL Surveyor enhancer. Digested products were

then resolved on a 2.5% agarose gel, stained with 0.01% SYBR Gold DNA (LifeTechnologies,

Massachusetts, USA) in 1% TBE buffer. The size of the occurring bands indicated the location of

the mutation (expected DNA fragments sizes are provided in Supplementary Table 8). To

quantify the knockdown efficiency, the software ImageJ was used. Peak areas of the bands

visualized on agarose gel were selected and the percentages of the transduced neurons

acquiring the InDel mutation were calculated using the following formula [106]:

$$Editing\ (\%) = 100 \times \left(1 - \sqrt{\left(1 - \frac{b+c}{a+b+c}\right)}\right)$$

491    where $a$ is the integrated intensity of the undigested PCR product and $b$ and $c$ are the integrated

492    intensities of each cleavage fragment from the digested product.

**Immunofluorescence of rat primary neurons**

Rat primary neurons were processed1-week post-AAV-transduction. Coverslips were washed

twice with DPBS and fixed in 4% PFA in DPBS for 15 min at room temperature. Blocking and

permeabilization were performed for 30 min in DPBS with 5% donkey serum, 0.5% Triton-X100,

and 0.05% BSA. Primary antibody incubation (VMAT2, 1:50, EB06558, Everest Biotech,

Ramona, USA, HA-tag: 1:50, 682404, Biolegend, San Diego, USA) was performed for 60 min,

followed by Cy3 conjugated secondary antibody incubation (1:250, kindly provided from Birgit

Fehrenbacher, Department of Dermatology, University of Tuebingen, Germany). Coverslips were

mounted using ProLong Antifade Diamond Medium containing DAPI (LifeTechnologies,

Massachusetts, USA), and imaged with a TCS-SP2/Leica DM IRE2 confocal laser scanning

microscope. Images were processed with Leica Confocal Software LCS (Version 2.61) (original

magnification 630) (Leica Microsystems, Wetzlar, Germany).

**Study design of the *in vivo* experiments**

In a first cohort, AAVs-CRISPR/SaCas9 were stereotactically delivered into the right SNc of wild-

type rats. Afterward, *in vivo* PET scans with imaging markers of VMAT2 expression, dopamine

availability, nerve terminal integrity, and inflammatory responses were performed in VMAT2

knockdown and control rats using [$^{11}$C]DTBZ, [$^{11}$C]raclopride, [$^{11}$C]methylphenidate, and [$^{18}$F]GE-

180, respectively. Motor consequences of the CRISPR/SaCas9-induced VMAT2 knockdown

were explored in a wide spectrum of behavioral tasks. Finally, biochemical and histological

analyses were performed to corroborate the *in vivo* data (Fig. 2,3,4a).

In a second cohort, cylinder test and [$^{11}$C]raclopride-PET/BOLD-fMRI scans were performed at

baseline and after CRISPR/SaCas9-induced VMAT2 knockdown. These measurements were

paralleled by *in vivo* PET scans with imaging markers of VMAT2 and GABA-A expression, to

518   inspect the extent of the induced VMAT2 knockdown and its impact on GABA signaling, using

519   [$^{11}$C]DTBZ and [$^{11}$C]flumazenil, respectively (Fig. 5a).

**Stereotactic injections**

Rats were anesthetized by injecting a mixture (1 mL/kg) of fentanyl (0.005 mg/kg), midazolam

(2 mg/kg,) and medetomidine (0.15 mg/kg) intraperitoneally, and were placed onto a stereotactic

frame (Harvard Apparatus, Holliston, MA, USA) with the skull flat between Lambda and Bregma.

The following coordinates were used for the injections (flat skull position): AP: - 5 mm, ML:

± 2 mm, DV: - 7.2 mm, below the dural surface as calculated relative to Bregma. Rats were

injected with 3 µL AAVs into the right SNc and were divided into two groups. Control rats (n= 10)

were injected with AAV-PHP.EB-sgRNA-*lacZ* (1.7 × 10$^{12}$ gc/mL) and AVV-PHP.EB-SaCas9

(1.4 × 10$^{12}$ gc/mL) (1:1 ratio). VMAT2 knockdown rats (n= 47) were injected with AAV-PHP.EB-

sgRNA-*Slc18a2* (2.1 × 10$^{12}$ gc/mL) and AVV-PHP.EB-SaCas9 (1.4 × 10$^{12}$ gc/mL) (1:1 ratio) (see

Fig. 1a for vector constructs). DPBS (3 µL) was sham-injected into the contralateral SNc.

Solutions were infused at a rate of 0.2 µL/min using a 5 µL Hamilton syringe (Hamilton, Bonaduz,

Switzerland) and an automated microsyringe pump (Harvard Apparatus, Holliston, MA, USA). To

allow for the diffusion of AAVs into the tissue, the needle was left in place for 10 min, and then

slowly retracted at 0.2 mm/min. After the surgery, an antidote containing atipamezole

(0.75 mg/kg) and flumazenil (0.2 mg/kg) was injected subcutaneously. The rats were kept warm

in their cages until fully recovered.

**Radiotracer synthesis**

[$^{11}$C]CO$_2$ was produced on a medical cyclotron (PETtrace 860, GE Healthcare, Uppsala, Sweden)

using the $^{14}$N(p,α)$^{11}$C route and converted to either [$^{11}$C]MeI (methyl iodide) or [$^{11}$C]MeOTf (methyl

triflate) using a Tracerlab FX MeI module (GE Healthcare). A Tracerlab FX M module (GE

Healthcare) was used for automated methylation, purification, and formulation of the tracers.

[11C]Dihydrotetrabenazine (DTBZ), a VMAT2 ligand, was synthesized starting from [11C]MeI,

which was reacted with 1 mg (+)-9-O-desmethyl-dihydrotetrabenazine (ABX, Radeberg,

Germany) in 300 µl DMF in the presence of 7.5 µl 5 M NaOH for 3 min at 40 °C [108]. Afterward,

it was purified by HPLC and formulated as a sterile pyrogen-free saline solution. The total

synthesis time from the end of the beam was 45 min. The radiochemical purity of the final

radiotracer was > 95% as determined by HPLC. Molar activity at the time of injection was 96 ± 37

GBq/µmol.

D-threo-[11C]methylphenidate, a dopamine transporter ligand, was synthesized by alkylation of D-

threo-N-NPS-ritalinic acid (ABX) using [11C]MeI [109]. After acidic deprotection, purification, and

formulation, the product was obtained with a 37 ± 6% decay-corrected radiochemical yield (from

[11C]MeI). The total synthesis time was 55 min, and the radiochemical purity of the final

formulated radiotracer was > 95% as determined by HPLC analysis. The molar activity was

determined at the time of injection as 58 ± 14 GBq/µmol.

[11C]Raclopride a D2R ligand, was synthesized by alkylation of S-(+)-O-desmethyl-raclopride

(ABX) using [11C]MeOTf. After purification and formulation, the product was obtained with a

12 ± 4% decay-corrected radiochemical yield (from [11C]methyl triflate). The total synthesis time

from the end of the beam was 55 min. The radiochemical purity of the final formulated radiotracer

was > 95% as determined by HPLC. The molar activity was determined at the time of injection

and was calculated as 88 ± 41 GBq/µmol.

[11C]Flumazenil, a GABA-A ligand, was synthesized by methylation of desmethylflumazenil (ABX)

with [11C]MeI. 2 mg of the precursor were dissolved in 0.3 ml DMF with 3 µl 5 M NaOH and

reacted for 2 min at 80 °C. After the reaction, the product was purified by semi-preparative HPLC

and reformulated by solid-phase extraction (Strata-X, Phenomenex; elution with 0.5 ml ethanol,

dilution with 5 ml phosphate-buffered saline). The total synthesis time from the end of the beam

was 50 min. The radiochemical purity of the final formulated radiotracer was > 95% as

determined by HPLC. The molar activity was determined at the time of injection as

109.5 ± 39.6 GBq/μmol.

[$^{18}$F]Fluoride was produced on a PETtrace 860 medical cyclotron (GE Healthcare) using the

$^{18}$O(p,n)$^{18}$F route.

[$^{18}$F]GE-180, a translocator protein ligand [110], was synthesized on a FASTlab synthesizer (GE

Healthcare) with precursor and reagent kits in single-use cassettes (GE Healthcare) according to

the manufacturer's instructions. The radiochemical purity of the final formulated radiotracer was

> 95% as determined by HPLC. The molar activity was determined at the time of injection as

576 ± 283 GBq/μmol.

***In vivo* PET imaging and data analysis**

For the study of cohort 1, VMAT2 knockdown (n= 14) and control rats (n= 10) underwent 60 min

dynamic PET emission scans with [$^{11}$C]DTBZ (8 – 10 weeks post-injection), [$^{11}$C]methylphenidate

(10 – 12 weeks post-injection), [$^{11}$C]raclopride (12 – 14 weeks post-injection) and [$^{18}$F]GE-180

(14 – 16 weeks post-injection) (Fig. 2a). Four rats were excluded from the data analyses because

two rats from each group died during a PET acquisition. One control rat was excluded from the

[$^{11}$C]methylphenidate analysis due to a poor signal-to-noise ratio.

For the study of cohort 2, VMAT2 knockdown rats (n= 33), underwent 60 min dynamic PET

emission acquisitions with [$^{11}$C]DTBZ (8 – 10 weeks post-injection), and [$^{11}$C]flumazenil (10 –

12 weeks post-injection). The final cohort included 23 rats (see paragraph Simultaneous

PET/fMRI experiments).

Three small-animal PET scanners (Inveon, Siemens, Erlangen, Germany) and dedicated rat

brain beds (Jomatik Gmbh, Tuebingen, Germany) with stereotactic holders and temperature

feedback control units (Medres, Cologne, Germany) were used. These ensured the delivery and

removal of the anesthesia gas and stabilized the body temperature at 37 °C during the PET data

acquisition. Anesthesia was induced by placing the animals in knock-out boxes and delivering 2%

isoflurane in oxygen air. Subsequently, a 24 G catheter (BD Insyte, NJ, USA) was placed into the

tail vein for the tracer and/or *i.v.* anesthesia administration. Afterward, animals of cohort 1 were

anesthetized with 2% isoflurane vaporized in 1.0 L/min of oxygen. Animals of cohort 2 received a

medetomidine bolus injection (0.05 mg/kg) and the anesthesia was switched to constant

medetomidine infusion (0.1 mg/kg/h), and 0.5% isoflurane in air during the scan time, as adapted

from the literature [111].

The rats were placed in the center of the field of view and PET acquisitions started 5 s before the

bolus injection of the tracer. In Supplementary Tables 9 and 10, injected activity (MBq/kg) and

molar activity (GBq/µmol) at the time of injection are reported for each radioligand. The list-mode

data from the dynamic acquisitions of [$^{11}$C]DTBZ, [$^{11}$C]raclopride, [$^{11}$C]methylphenidate, and

[$^{11}$C]flumazenil were histogrammed into 39 time-frames (12x5 s, 6x10 s, 6x30 s, 5x60 s,

10x300 s), from [$^{18}$F]GE-180 into 16 time-frames (5x60 s, 5x120 s, 3x300 s, 3x600 s). PET scans

of the study cohort 1 were reconstructed using the OSEM3D map algorithm, and a matrix size of

256 x 256 x 159, resulting in a pixel size of 0.38 x 0.38 x 0.79 mm. PET scans of the study cohort

2 were reconstructed using the OSEM2D algorithm, and a matrix size of 256 x 256 x 89, resulting

in a pixel size of 0.33 x 0.33 x 0.79 mm.

Data preprocessing analysis was performed with Matlab (Mathworks, Natick, MA, USA),

Statistical Parametric Mapping 12 (SPM12, Wellcome Trust Centre for Neuroimaging, University

College London, England), and the QModeling toolbox [112]. First, realignment of all frames was

performed using SPM12 and average images were generated for every scan. The mean images

were then used for coregistration to the Schiffer rat brain atlas provided by PMOD software [113].

To generate the respective time activity curves (TAC), volumes of interest (VOIs) were defined

over the target and reference regions. VOIs were placed over the right and left striatum for

[$^{11}$C]DTBZ, [$^{11}$C]raclopride, and [$^{11}$C]methylphenidate, and over the regions reported in Table 2 for

[$^{11}$C]flumazenil. Cerebellum was used as reference region for [$^{11}$C]DTBZ, [$^{11}$C]raclopride and

[$^{11}$C]methylphenidate. Pons was used as reference region for [$^{11}$C]flumazenil.

Binding potentials for [$^{11}$C]DTBZ, [$^{11}$C]raclopride, [$^{11}$C]methylphenidate and [$^{11}$C]flumazenil were

calculated over the all frames in the regions of interest with Logan reference [114], with a

population average k2' ([$^{11}$C]DTBZ: 0.41 min$^{-1}$, [$^{11}$C]raclopride: 0.34 min$^{-1}$, [$^{11}$C]methylphenidate:

0.18 min$^{-1}$, [$^{11}$C]flumazenil: 0.27 min$^{-1}$). [$^{11}$C]DTBZ and [$^{11}$C]raclopride binding changes (%), here

expressed as $\Delta$ binding, were calculated according to the formula:

$$\Delta\,binding = \frac{1 - binding\ right\ striatum}{binding\ left\ striatum} \times 100$$

[$^{18}$F]GE-180 uptake in the right striatum was calculated over the interval between 30 – 60 min

after scan start and normalized by the uptake calculated in the left (DPBS-injected) striatum over

the same time interval.

QModeling was used to generate voxel-wise binding potential maps for [$^{11}$C]DTBZ,

[$^{11}$C]raclopride, [$^{11}$C]methylphenidate, and [$^{11}$C]flumazenil. [$^{18}$F]GE-180 average uptake images

were generated with an in-house-written script in MATLAB.

**Simultaneous PET/fMRI experiments**

Rats (n= 33) underwent longitudinal simultaneous [$^{11}$C]raclopride-PET/BOLD-fMRI scans at

baseline and 8 – 14 weeks after CRISPR/SaCas9-induced VMAT2 knockdown. Three rats had to

be excluded from the data analyses due to aliasing artifacts, local distortion, and motion during

the data acquisition. Seven rats died during a PET/BOLD-fMRI scan. The final cohort included 23

rats. Anesthesia induction and injection were performed as described above for the cohort 2 (see

paragraph *In vivo* PET imaging and data analysis).

Next, rats were placed onto a dedicated rat bed (Medres, Cologne, Germany) and a temperature

feedback control unit (Medres, Cologne, Germany), ensuring the delivery and removal of the

anesthesia gas and stabilizing the body temperature at 37 °C during the scan time. A breathing

pad and a pulse oximeter were used to observe respiration and heart rates. Scans were acquired

using a small-animal 7 T Clinscan MRI scanner, a 72 cm diameter linearly polarized RF coil

(Bruker) for transmission, and a four-channel rat brain coil for reception (Bruker Biospin MRI,

Ettlingen, Germany). Localizer scans were first acquired to accurately position the rat brains into

the center of the PET/MRI field of view. Subsequently, local field homogeneity was optimized by

measuring local magnetic field maps. Anatomical reference scans were performed using T2-

weighted Turbo-RARE MRI sequences (TR: 1800 ms, TE: 67.11 ms, FOV: 40 x 32 x 32 mm,

image size: 160 x 128 x 128 px, Rare factor: 28, averages: 1). Finally, T2*-weighted gradient

echo EPI sequences (TR: 2500 ms, TE: 18 ms, 0.25 mm isotropic resolution, FoV 25 x 23 mm,

image size: 92 x 85 x 20 px, slice thickness 0.8 mm, slice separation 0.2 mm, 20 slices) were

acquired for BOLD-fMRI.

A dedicated small-animal PET insert developed in cooperation with Bruker (Bruker Biospin MRI,

Ettlingen, Germany) was used for the [11C]raclopride acquisitions, the second generation of a

PET insert developed in-house with similar technical specifications [115]. [11C]Raclopride was

applied via a bolus injection. In Supplementary Table 10 injected activities (MBq/kg) and molar

activities (GBq/µmol) at the time of injection are reported. PET/fMRI acquisitions started

simultaneously with the tracer injection and were performed over a period of 60 min. The list-

mode files of the PET data were histogrammed into 14 time-frames (1x30 s, 5x60 s, 5x300 s,

3x600 s), the 30s between acquisition start and the injection were excluded from the analysis.

Reconstruction was performed with an in-house-written OSEM2D algorithm. Data preprocessing

and analysis were performed as described above (see paragraph *In vivo* PET imaging and data

664 analysis). A population average k2' ([$^{11}$C]raclopride baseline: 0.20 min$^{-1}$, [$^{11}$C]raclopride VMAT2

665 knockdown: 0.23 min$^{-1}$) was set for the Logan reference [114].

666 Preprocessing of the fMRI data was performed using a pipeline employing SPM12, Analysis of

667 Functional NeuroImages (AFNI, National Institute of Mental Health (NIMH), Bethesda, Maryland,

668 USA), and in-house-written scripts in MATLAB, as reported previously [22].

669 RS-FC was calculated using a seed-based approach. To this extent, 20 regions were selected

670 from the Schiffer rat brain atlas (a list of the regions is provided in Table 2). The SPM toolbox

671 Marseille Boîte À Région d'Intérêt (MarsBaR) was employed to extract fMRI time-courses from all

672 regions [116]. These were then used to calculate pairwise Pearson's r correlation coefficients for

673 each dataset, generating correlation matrices containing 20 x 20 elements. Self-correlations were

674 set to zero. The computed Pearson's r coefficients then underwent Fischer's transformation into z

675 values for group-level analysis.

676 Several rs-FC metrics were computed on different regional levels to investigate the potential

677 effects of dopamine depletion in the right striatum. Regional node strengths were computed as

678 the sum of all correlations of one seed to the regions included in one network. Interregional node

679 strengths were defined as the sum of the correlations of one node to the regions of another

680 network. Network strengths were defined as the sum of strengths of all correlations between

681 regions belonging to a network. Internetwork strengths were calculated as the sum of all

682 correlations between two sets of regions belonging to two networks [117].

684 **Behavioral analysis**

685 *Cylinder test*

686 Untrained rats were placed individually inside a glass cylinder (19 cm Ø, 20 cm height). The test

687 started immediately and lasted 5 min. During the test session rats were left undisturbed and were

688 videotaped with a camera located at the bottom-center of the cylinder to allow a 360 ° angle view.

689 Paw touches were analyzed using a slow-motion video player (VLC software, VideoLan). The

690   number of wall touches, contacts with fully extended digits, was counted. Data were analyzed as

691   follows:

$$contralateral\ paw\ touches\ (\%\ to\ total) = \frac{contralateral\ touches}{total\ touches} \times 100$$

One rat from the cohort 2 was excluded from the analysis due to issues during video recording.

*Rotameter test*

Rotational asymmetry was assessed using an automated rotameter system composed of four

hemispheres (TSE Systems GmBH, Bad Homburg, Germany) based on the design of Ungerstedt

and Arbuthnott [118]. Rats were placed into an opaque half bowl (49 cm Ø; 44 cm height) and

fixed to a moveable wire with a collar. The number of clockwise (CW) and counterclockwise

(CCW) rotations (difference of 42.3° in their position) were automatically counted. The

spontaneous rotation test lasted 5 min. Apomorphine-evoked rotational asymmetry was

evaluated for 60 min after *s.c.* administration of apomorphine hydrochloride (0.25 mg/kg)

dissolved in physiological saline containing 0.1% ascorbic acid (Sigma Aldrich, St. Louis,

Missouri, USA). Two priming injections of apomorphine (1-week interval off-drug) were necessary

to produce sensitization to the treatment. The program RotaMeter (TSE Systems GmBH, Bad

Homburg, Germany) was used to acquire the data. Data were analyzed as follows:

$$turns/min = ccw\ rotation - cw\ rotations$$

*Beam walk test*

Rats were trained for 4 days to cross a built-in-house beam (1.7 cm width, 60 cm length, 40 cm

height), and reach a cage with environmental enrichment. Rats had 5 trials to cross the beam

with the reward resting time decreasing from 30 s to 10 s. On the test day (one week apart from

the fourth day of training), rats were videotaped. The acquired videos were analyzed using a

slow-motion video player (VLC software, VideoLan) and the number of footslips (falls) was

counted. Data were analyzed as follows:

$$contralateral\ footslips\ (\%\ to\ total\ steps) = \frac{contralateral\ footslips}{total\ steps} \times 100$$

*Open Field test*

Untrained rats were set in a rectangular box (TSE Systems GmBH, Bad Homburg, Germany) for

11 min (1 min habituation, 10 min test) to evaluate the spontaneous exploratory behavior. A

frame with light sensors, as high as the animals' body center, was connected to a receiver box to

record the rats' walked distance. The program ActiMod (TSE Systems GmBH, Bad Homburg,

Germany) was used for the analysis and the experimental session was divided into 1 min time

bins. Results were averaged from the total traveled distance over 10 min (mean ± SD).

*Body weight gain*

Rats' body weight was measured before (week 0) and 14 weeks after CRISPR/SaCas9-induced

VMAT2 knockdown. Body weight gain was calculated as follows:

$$body\ weight\ gain\ (g) = \frac{weight\ at\ week\ 14 - weight\ at\ week\ 0}{weight\ at\ week\ 0} \times 100$$

**Histology of rat brain slices**

Rats of cohort 1 (n= 20) were sacrificed 19 weeks after viral vector injection via $CO_2$ inhalation

followed by intracardial perfusion with heparinized DPBS (1:50 v/v, 100 mL). After decapitation,

brains were rapidly removed and placed in a brain matrix on ice. The left and right striata were

dissected, from 2 mm thick coronal sections, flash-frozen in liquid nitrogen, and stored at – 80 °C

until further analysis (HPLC). The remaining tissue was fixed in 4% paraformaldehyde for 48 h

and then transferred to a 20% sucrose solution for cryoprotection. Brains were cut into 35 μm

thick coronal sections on a freezing microtome (Leica Biosystems, Wetzlar, Germany) and stored

in an anti-freeze solution (0.5 M phosphate buffer, 30 % glycerol, 30 % ethylene glycol) at -

20 °C. Sections were collected in 12 equally spaced series through the entire anterior-posterior

extent of the SN and striatum and stored until further analysis. Immunohistochemistry was

performed on free-floating sections. Sections were washed 3 times with TBS buffer and antigen

retrieval was carried for 30 min at 80 °C in Tris/EDTA buffer. Afterward, pre-incubation in MeOH

(10%) and $H_2O_2$ (3%) in TBS was performed for 30 min. Following the blocking in 5% normal goat

serum in TBS-X (0.05%), primary antibody incubation was performed for 24 h at room

temperature in 1% BSA in TBS-X (TH: 1:5000, P40101, Pel-freez, Arkansas, USA, VMAT2:

1:5000, 20042, Immunostar, Hudson, USA). The tissue was rinsed in TBS-X and reacted with the

respective biotinylated secondary antibody (1:200, Vector Laboratories Ltd., Peterborough, UK)

for 60 min at room temperature in 1% BSA in TBS-X. Staining was developed using 3,3'-

diaminobenzidine (DAB Substrate Kit, Vector Laboratories Ltd., Peterborough, UK) and an

immunoperoxidase system (Vectastain Elite ABC-Kit, Vector Laboratories Ltd., Peterborough,

UK). Slices were rinsed, mounted onto chromalum gelatinized slides, dehydrated in ascending

concentrations of alcohol and xylene baths, and coverslipped with DPX mounting medium (Sigma

Aldrich, St. Louis, Missouri, USA).

**Stereological analysis**

Estimates of total numbers of TH+ cells in nigral sections were obtained with an unbiased

stereological quantification method by employing the optical fractionator principle [119]. Brain

sections from 5 rats (VMAT2 knockdown n= 2, Control n= 3) were excluded from the stereological

analysis, due to weak TH immunoreactivity. First, 5x images were acquired with the automated

Metafer slide scanning platform (MetaSystems, Altlußheim, Germany). Then, ROIs were drawn

using the VSViewer program (Metasystems, Altlußheim, Germany), and a sampling fraction of

50% was defined. Afterward, 63x images were acquired in an automated fashion based on the

sampling fraction in a random orientation within the ROI. The acquired 63x images were imported

into the VIS program and cell counting was performed with the CAST module (Visiopharm A/S,

Hørsholm, Denmark, Version 2020.08.2.8800).

The number of cells estimates was obtained by applying the formula:

$$Total\ number\ of\ TH + cells = \sum \left( \frac{Number\ of\ cells\ counted\ per\ brain}{Acquisition\ fraction\ x\ counting\ frame} \right) \times number\ of\ series$$

**Immunofluorescence of rat brain slices**

Immunofluorescence was performed on SN sections from rats of cohort 1 (35 µm thick coronal

sections), mounted onto chromalum gelatinized slides, dehydrated in ascending concentrations of

alcohol and xylene baths. Sections were washed 3 times with KPBS buffer and antigen retrieval

was carried for 30 min at 80 °C in Tris/EDTA buffer. Following blocking in 5% donkey serum and

normal horse serum in KPBS-X (0.25%), primary antibody incubation was performed for 24 h at

room temperature (VMAT2: 1:5000, 20042, Immunostar, Hudson, USA, HA-tag: 1:5000, MMS-

101R, Nordic BioSite, Taby, Sweden, GFP: 1:50.000, ab13970, Abcam, Cambridge, UK). The

tissue was rinsed in KPBS-X and reacted with the respective fluorophore-conjugated secondary

antibody (1:200, Vector Laboratories Ltd., Peterborough, UK) for 2 h in 0.2% KPBS-X. Slides

were coverslipped with Vectashield mounting medium (Vector Laboratories Ltd., Peterborough,

UK).

**Biochemistry**

Dopamine, 3,4-Dihydroxyphenylacetic acid (DOPAC), homovanillic acid (HVA), and serotonin (5-

HT) striatal levels were determined by HPLC. Brain samples from 5 rats (VMAT2 knockdown

n= 3, Control n= 2) were excluded from the analysis as striatal sections did not fall in the selected

range: 1.2 – 0.4 mm.

Briefly, striatal brain lysates generated from cohort 1 were injected by a cooled autosampler into

an ESA Coulchem III coupled to a Decade Elite electrochemical detector (Antec Scientific,

Zoeterwoude, The Netherlands) set to a potential of +350 mV. Separation was facilitated by using

an Atlantis Premier BEH C18 AX column (Waters Corporation, Massachusetts, USA) and a dual

mobile phase gradient of decreasing octane sulfonic acid (OSA) and increasing MeOH content

(mobile phase A containing 100 mM $PO_4$-buffer pH 2.50 and 4.62 mM OSA and mobile phase B

containing 100 mM $PO_4$-buffer pH 2.50 and 2.31 mM OSA), delivered at a flow rate of

0.35 mL/min to an Atlantis Premier BEH C18 column (particle size 2.5 μm, 2.1 mm x 150 mm)

(Waters Corporation, Massachusetts, USA). Data was collected using the Chemstation software

(Agilent, California, USA) and then exported to Chromeleon (LifeTechnologies, Massachusetts,

USA) for data quality control, peak integration, and concentration calculations. Striatal

metabolites' content was expressed in nmol for each sample and normalized to total protein (mg).

Dopamine turnover rate was calculated according to the formula:

$$Dopamine\ turnover\ rate = \frac{HVA + DOPAC}{DA}$$

**Statistics**

Statistical analysis was performed with GraphPad Prism 9.0 (Graphpad Software) if not otherwise

stated. Results were analyzed using paired t-tests for the within-subjects comparisons, and

unpaired t-tests for the between-groups comparisons. Correlations were performed using linear

regression analyses. Synaptic dysfunction discrimination was tested with multiple-comparison

ANOVA.

## Bibliography

1. Biswal, B., et al., *Functional connectivity in the motor cortex of resting human brain using echo-planar MRI.* Magn Reson Med, 1995. **34**(4): p. 537-41.

2. Ogawa, S., et al., *Brain magnetic resonance imaging with contrast dependent on blood oxygenation.* Proc Natl Acad Sci U S A, 1990. **87**(24): p. 9868-72.

3. Lu, H., et al., *Rat brains also have a default mode network.* Proc Natl Acad Sci U S A, 2012. **109**(10): p. 3979-84.

4. Becerra, L., et al., *Robust reproducible resting state networks in the awake rodent brain.* PLoS One, 2011. **6**(10): p. e25701.

5. Fox, M.D., et al., *The human brain is intrinsically organized into dynamic, anticorrelated functional networks.* Proc Natl Acad Sci U S A, 2005. **102**(27): p. 9673-8.

6. Jonckers, E., et al., *Functional connectivity fMRI of the rodent brain: comparison of functional connectivity networks in rat and mouse.* PLoS One, 2011. **6**(4): p. e18876.

7. Wehrl, H.F., et al., *Simultaneous PET-MRI reveals brain function in activated and resting state on metabolic, hemodynamic and multiple temporal scales.* Nat Med, 2013. **19**(9): p. 1184-9.

8. Amend, M., et al., *Functional resting-state brain connectivity is accompanied by dynamic correlations of application-dependent [(18)F]FDG PET-tracer fluctuations.* Neuroimage, 2019. **196**: p. 161-172.

9. Ionescu, T.M., et al., *Elucidating the complementarity of resting-state networks derived from dynamic [(18)F]FDG and hemodynamic fluctuations using simultaneous small-animal PET/MRI.* Neuroimage, 2021. **236**: p. 118045.

10. Badhwar, A., et al., *Resting-state network dysfunction in Alzheimer's disease: A systematic review and meta-analysis.* Alzheimers Dement (Amst), 2017. **8**: p. 73-85.

11. Bluhm, R.L., et al., *Spontaneous low-frequency fluctuations in the BOLD signal in schizophrenic patients: anomalies in the default network.* Schizophr Bull, 2007. **33**(4): p. 1004-12.

12. Volkow, N.D., et al., *Imaging endogenous dopamine competition with [11C]raclopride in the human brain.* Synapse, 1994. **16**(4): p. 255-62.

13. Laruelle, M., *Imaging synaptic neurotransmission with in vivo binding competition techniques: a critical review.* J Cereb Blood Flow Metab, 2000. **20**(3): p. 423-51.

14. Patel, V.D., et al., *Imaging dopamine release with Positron Emission Tomography (PET) and (11)C-raclopride in freely moving animals.* Neuroimage, 2008. **41**(3): p. 1051-66.

15. Roffman, J.L., et al., *Dopamine D1 signaling organizes network dynamics underlying working memory.* Science advances, 2016. **2**(6): p. e1501672-e1501672.

16. McCutcheon, R.A., et al., *Mesolimbic Dopamine Function Is Related to Salience Network Connectivity: An Integrative Positron Emission Tomography and Magnetic Resonance Study.* Biol Psychiatry, 2019. **85**(5): p. 368-378.

17. Nagano-Saito, A., et al., *Posterior dopamine D2/3 receptors and brain network functional connectivity.* Synapse, 2017. **71**(11).

18. Hahn, A., et al., *Differential modulation of the default mode network via serotonin-1A receptors.* Proc Natl Acad Sci U S A, 2012. **109**(7): p. 2619-24.

19. Vidal, B., et al., *In vivo biased agonism at 5-HT(1A) receptors: characterisation by simultaneous PET/MR imaging.* Neuropsychopharmacology, 2018. **43**(11): p. 2310-2319.

20. Shiyam Sundar, L.K., et al., *Fully Integrated PET/MR Imaging for the Assessment of the Relationship Between Functional Connectivity and Glucose Metabolic Rate.* Front Neurosci, 2020. **14**: p. 252.

21. Sander, C.Y., et al., *Imaging Agonist-Induced D2/D3 Receptor Desensitization and Internalization In Vivo with PET/fMRI.* Neuropsychopharmacology, 2016. **41**(5): p. 1427-36.

| | | |
|---|---|---|
| 851 | 22. | Ionescu, T., *Striatal and prefrontal D2R and SERT distributions contrastingly correlate with default-mode connectivity.* 2021. |
| 853 | 23. | Betley, J.N. and S.M. Sternson, *Adeno-associated viral vectors for mapping, monitoring, and manipulating neural circuits.* Hum Gene Ther, 2011. **22**(6): p. 669-77. |
| 855 | 24. | Jinek, M., et al., *RNA-programmed genome editing in human cells.* Elife, 2013. **2**: p. e00471. |
| 856 | 25. | Cong, L., et al., *Multiplex genome engineering using CRISPR/Cas systems.* Science, 2013. **339**(6121): p. 819-23. |
| 858 | 26. | Whitworth, K.M., et al., *Use of the CRISPR/Cas9 system to produce genetically engineered pigs from in vitro-derived oocytes and embryos.* Biol Reprod, 2014. **91**(3): p. 78. |
| 860 | 27. | Chen, Y., et al., *Functional disruption of the dystrophin gene in rhesus monkey using CRISPR/Cas9.* Hum Mol Genet, 2015. **24**(13): p. 3764-74. |
| 862 | 28. | Chang, N., et al., *Genome editing with RNA-guided Cas9 nuclease in zebrafish embryos.* Cell Res, 2013. **23**(4): p. 465-72. |
| 864 | 29. | Platt, R.J., et al., *CRISPR-Cas9 knockin mice for genome editing and cancer modeling.* Cell, 2014. **159**(2): p. 440-55. |
| 866 | 30. | Yu, Z., et al., *Highly efficient genome modifications mediated by CRISPR/Cas9 in Drosophila.* Genetics, 2013. **195**(1): p. 289-91. |
| 868 | 31. | Musunuru, K., et al., *In vivo CRISPR base editing of PCSK9 durably lowers cholesterol in primates.* Nature, 2021. **593**(7859): p. 429-434. |
| 870 | 32. | Ousterout, D.G., et al., *Multiplex CRISPR/Cas9-based genome editing for correction of dystrophin mutations that cause Duchenne muscular dystrophy.* Nat Commun, 2015. **6**: p. 6244. |
| 872 | 33. | Swiech, L., et al., *In vivo interrogation of gene function in the mammalian brain using CRISPR-Cas9.* Nature Biotechnology, 2015. **33**(1): p. 102-U286. |
| 874 | 34. | Wilbie, D., J. Walther, and E. Mastrobattista, *Delivery Aspects of CRISPR/Cas for in Vivo Genome Editing.* Accounts of chemical research, 2019. **52**(6): p. 1555-1564. |
| 876 | 35. | Wu, W.Y., et al., *Genome editing by natural and engineered CRISPR-associated nucleases.* Nat Chem Biol, 2018. **14**(7): p. 642-651. |
| 878 | 36. | Zetsche, B., et al., *Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system.* Cell, 2015. **163**(3): p. 759-71. |
| 880 | 37. | Kleinstiver, B.P., et al., *Engineered CRISPR-Cas9 nucleases with altered PAM specificities.* Nature, 2015. **523**(7561): p. 481-5. |
| 882 | 38. | Ran, F.A., et al., *In vivo genome editing using Staphylococcus aureus Cas9.* Nature, 2015. **520**(7546): p. 186-91. |
| 884 | 39. | Nishimasu, H., et al., *Crystal Structure of Staphylococcus aureus Cas9.* Cell, 2015. **162**(5): p. 1113-26. |
| 886 | 40. | Sun, H., et al., *Development of a CRISPR-SaCas9 system for projection- and function-specific gene editing in the rat brain.* Sci Adv, 2020. **6**(12): p. eaay6687. |
| 888 | 41. | Kumar, N., et al., *The Development of an AAV-Based CRISPR SaCas9 Genome Editing System That Can Be Delivered to Neurons in vivo and Regulated via Doxycycline and Cre-Recombinase.* Front Mol Neurosci, 2018. **11**: p. 413. |
| 891 | 42. | Hunker, A.C., et al., *Conditional Single Vector CRISPR/SaCas9 Viruses for Efficient Mutagenesis in the Adult Mouse Nervous System.* Cell Rep, 2020. **30**(12): p. 4303-4316 e6. |
| 893 | 43. | Liu, Y. and R.H. Edwards, *The role of vesicular transport proteins in synaptic transmission and neural degeneration.* Annu Rev Neurosci, 1997. **20**: p. 125-56. |
| 895 | 44. | Miyanishi, K., et al., *Behavioral tests predicting striatal dopamine level in a rat hemi-Parkinson's disease model.* Neurochem Int, 2019. **122**: p. 38-46. |
| 897 | 45. | Tomasi, D., et al., *Dopamine transporters in striatum correlate with deactivation in the default mode network during visuospatial attention.* PLoS One, 2009. **4**(6): p. e6102. |

46.  Sambataro, F., et al., *DRD2 genotype-based variation of default mode network activity and of its relationship with striatal DAT binding.* Schizophr Bull, 2013. **39**(1): p. 206-16.

47.  Tritsch, N.X., J.B. Ding, and B.L. Sabatini, *Dopaminergic neurons inhibit striatal output through non-canonical release of GABA.* Nature, 2012. **490**(7419): p. 262-6.

48.  Tritsch, N.X., et al., *Midbrain dopamine neurons sustain inhibitory transmission using plasma membrane uptake of GABA, not synthesis.* Elife, 2014. **3**: p. e01936.

49.  Chatterjee, D., et al., *Enhanced CNS transduction from AAV.PHP.eB infusion into the cisterna magna of older adult rats compared to AAV9.* Gene Ther, 2021.

50.  Hume, S.P., et al., *Effect of L-dopa and 6-hydroxydopamine lesioning on [11C]raclopride binding in rat striatum, quantified using PET.* Synapse, 1995. **21**(1): p. 45-53.

51.  Mishra, R.K., A.M. Marshall, and S.L. Varmuza, *Supersensitivity in rat caudate nucleus: effects of 6-hydroxydopamine on the time course of dopamine receptor and cyclic AMP changes.* Brain Res, 1980. **200**(1): p. 47-57.

52.  Creese, I., D.R. Burt, and S.H. Snyder, *Dopamine receptor binding enhancement accompanies lesion-induced behavioral supersensitivity.* Science, 1977. **197**(4303): p. 596-8.

53.  Konieczny, J., et al., *The significance of rotational behavior and sensitivity of striatal dopamine receptors in hemiparkinsonian rats: A comparative study of lactacystin and 6-OHDA.* Neuroscience, 2017. **340**: p. 308-318.

54.  Wang, Y.M., et al., *Knockout of the vesicular monoamine transporter 2 gene results in neonatal death and supersensitivity to cocaine and amphetamine.* Neuron, 1997. **19**(6): p. 1285-96.

55.  Seeman, P., et al., *Dopamine receptors in the central nervous system.* Fed Proc, 1978. **37**(2): p. 131-6.

56.  Gnanalingham, K.K. and R.G. Robertson, *The effects of chronic continuous versus intermittent levodopa treatments on striatal and extrastriatal D1 and D2 dopamine receptors and dopamine uptake sites in the 6-hydroxydopamine lesioned rat--an autoradiographic study.* Brain Res, 1994. **640**(1-2): p. 185-94.

57.  Graham, W.C., A.R. Crossman, and G.N. Woodruff, *Autoradiographic studies in animal models of hemi-parkinsonism reveal dopamine D2 but not D1 receptor supersensitivity. I. 6-OHDA lesions of ascending mesencephalic dopaminergic pathways in the rat.* Brain Res, 1990. **514**(1): p. 93-102.

58.  Antonini, A., et al., *Long-term changes of striatal dopamine D2 receptors in patients with Parkinson's disease: a study with positron emission tomography and [11C]raclopride.* Mov Disord, 1997. **12**(1): p. 33-8.

59.  Kraemmer, J., et al., *Correlation of striatal dopamine transporter imaging with post mortem substantia nigra cell counts.* Mov Disord, 2014. **29**(14): p. 1767-73.

60.  Sossi, V., et al., *Dopamine transporter relation to levodopa-derived synaptic dopamine in a rat model of Parkinson's: an in vivo imaging study.* J Neurochem, 2009. **109**(1): p. 85-92.

61.  Zigmond, M.J., T.G. Hastings, and R.G. Perez, *Increased dopamine turnover after partial loss of dopaminergic neurons: compensation or toxicity?* Parkinsonism Relat Disord, 2002. **8**(6): p. 389-93.

62.  Takahashi, N., et al., *VMAT2 knockout mice: heterozygotes display reduced amphetamine-conditioned reward, enhanced amphetamine locomotion, and enhanced MPTP toxicity.* Proc Natl Acad Sci U S A, 1997. **94**(18): p. 9938-43.

63.  Fon, E.A., et al., *Vesicular transport regulates monoamine storage and release but is not essential for amphetamine action.* Neuron, 1997. **19**(6): p. 1271-83.

64.  Mooslehner, K.A., et al., *Mice with very low expression of the vesicular monoamine transporter 2 gene survive into adulthood: potential mouse model for parkinsonism.* Mol Cell Biol, 2001. **21**(16): p. 5321-31.

65.  Caudle, W.M., et al., *Reduced vesicular storage of dopamine causes progressive nigrostriatal neurodegeneration.* J Neurosci, 2007. **27**(30): p. 8138-48.

66. Taylor, T.N., et al., *Nonmotor symptoms of Parkinson's disease revealed in an animal model with*
*reduced monoamine storage capacity.* J Neurosci, 2009. **29**(25): p. 8103-13.

67. Back, S., et al., *Neuron-Specific Genome Modification in the Adult Rat Brain Using CRISPR-Cas9*
*Transgenic Rats.* Neuron, 2019.

68. Zimmer, E.R., et al., *MicroPET imaging and transgenic models: a blueprint for Alzheimer's disease*
*clinical research.* Trends Neurosci, 2014. **37**(11): p. 629-41.

69. Andrews-Hanna, J.R., *The brain's default network and its adaptive role in internal mentation.*
Neuroscientist, 2012. **18**(3): p. 251-70.

70. Gottlich, M., et al., *Altered resting state brain networks in Parkinson's disease.* PLoS One, 2013.
**8**(10): p. e77336.

71. Huang, S., et al., *Multisensory Competition Is Modulated by Sensory Pathway Interactions with*
*Fronto-Sensorimotor and Default-Mode Network Regions.* J Neurosci, 2015. **35**(24): p. 9064-77.

72. Wu, C.W., et al., *Synchrony Between Default-Mode and Sensorimotor Networks Facilitates Motor*
*Function in Stroke Rehabilitation: A Pilot fMRI Study.* Front Neurosci, 2020. **14**: p. 548.

73. Wang, J., et al., *Functional reorganization of intra- and internetwork connectivity in major*
*depressive disorder after electroconvulsive therapy.* Hum Brain Mapp, 2018. **39**(3): p. 1403-1411.

74. Perlbarg, V., et al., *Alterations of the nigrostriatal pathway in a 6-OHDA rat model of Parkinson's*
*disease evaluated with multimodal MRI.* PLoS One, 2018. **13**(9): p. e0202597.

75. Monnot, C., et al., *Asymmetric dopaminergic degeneration and levodopa alter functional*
*corticostriatal connectivity bilaterally in experimental parkinsonism.* Exp Neurol, 2017. **292**: p. 11-
20.

76. Honey, G.D., et al., *Dopaminergic drug effects on physiological connectivity in a human cortico-*
*striato-thalamic system.* Brain, 2003. **126**(Pt 8): p. 1767-81.

77. Kwak, Y., et al., *Altered resting state cortico-striatal connectivity in mild to moderate stage*
*Parkinson's disease.* Front Syst Neurosci, 2010. **4**: p. 143.

78. Costa, R.M., et al., *Rapid alterations in corticostriatal ensemble coordination during acute*
*dopamine-dependent motor dysfunction.* Neuron, 2006. **52**(2): p. 359-69.

79. Gatev, P., O. Darbin, and T. Wichmann, *Oscillations in the basal ganglia under normal conditions*
*and in movement disorders.* Mov Disord, 2006. **21**(10): p. 1566-77.

80. Hammond, C., H. Bergman, and P. Brown, *Pathological synchronization in Parkinson's disease:*
*networks, models and treatments.* Trends Neurosci, 2007. **30**(7): p. 357-64.

81. Eusebio, A., et al., *Resonance in subthalamo-cortical circuits in Parkinson's disease.* Brain, 2009.
**132**(Pt 8): p. 2139-50.

82. Brazhnik, E., et al., *State-dependent spike and local field synchronization between motor cortex*
*and substantia nigra in hemiparkinsonian rats.* J Neurosci, 2012. **32**(23): p. 7869-80.

83. Moran, R.J., et al., *Alterations in brain connectivity underlying beta oscillations in Parkinsonism.*
PLoS Comput Biol, 2011. **7**(8): p. e1002124.

84. Gerfen, C.R. and D.J. Surmeier, *Modulation of striatal projection systems by dopamine.* Annu Rev
Neurosci, 2011. **34**: p. 441-66.

85. Caspers, J., et al., *Differential Functional Connectivity Alterations of Two Subdivisions within the*
*Right dlPFC in Parkinson's Disease.* Front Hum Neurosci, 2017. **11**: p. 288.

86. Owens-Walton, C., et al., *Increased functional connectivity of thalamic subdivisions in patients*
*with Parkinson's disease.* PLoS One, 2019. **14**(9): p. e0222002.

87. Kahn, I., et al., *Distinct cortical anatomy linked to subregions of the medial temporal lobe revealed*
*by intrinsic functional connectivity.* J Neurophysiol, 2008. **100**(1): p. 129-39.

88. Vincent, J.L., et al., *Coherent spontaneous activity identifies a hippocampal-parietal memory*
*network.* J Neurophysiol, 2006. **96**(6): p. 3517-31.

89. Hillary, F.G., et al., *Hyperconnectivity is a fundamental response to neurological disruption.*
Neuropsychology, 2015. **29**(1): p. 59-75.

90.   Klobusiakova, P., et al., *Connectivity Between Brain Networks Dynamically Reflects Cognitive*
*Status of Parkinson's Disease: A Longitudinal Study.* J Alzheimers Dis, 2019. **67**(3): p. 971-984.
91.   Mevel, K., et al., *The default mode network in healthy aging and Alzheimer's disease.* Int J
Alzheimers Dis, 2011. **2011**: p. 535816.
92.   Gorges, M., et al., *To rise and to fall: functional connectivity in cognitively normal and cognitively*
*impaired patients with Parkinson's disease.* Neurobiol Aging, 2015. **36**(4): p. 1727-1735.
93.   Helmich, R.C., et al., *Spatial remapping of cortico-striatal connectivity in Parkinson's disease.*
Cereb Cortex, 2010. **20**(5): p. 1175-86.
94.   O'Gorman Tuura, R.L., C.R. Baumann, and H. Baumann-Vogel, *Beyond Dopamine: GABA,*
*Glutamate, and the Axial Symptoms of Parkinson Disease.* Front Neurol, 2018. **9**: p. 806.
95.   Whiting, P.J., *The GABAA receptor gene family: new opportunities for drug development.* Curr
Opin Drug Discov Devel, 2003. **6**(5): p. 648-57.
96.   Kapogiannis, D., et al., *Posteromedial cortex glutamate and GABA predict intrinsic functional*
*connectivity of the default mode network.* Neuroimage, 2013. **64**: p. 112-9.
97.   Douaud, G., et al., *Integration of structural and functional magnetic resonance imaging in*
*amyotrophic lateral sclerosis.* Brain, 2011. **134**(Pt 12): p. 3470-9.
98.   Levar, N., et al., *Anterior cingulate GABA and glutamate concentrations are associated with*
*resting-state network connectivity.* Sci Rep, 2019. **9**(1): p. 2116.
99.   Lanoue, A.C., G.J. Blatt, and J.J. Soghomonian, *Decreased parvalbumin mRNA expression in*
*dorsolateral prefrontal cortex in Parkinson's disease.* Brain Res, 2013. **1531**: p. 37-47.
100.  Lanoue, A.C., et al., *Decreased glutamic acid decarboxylase mRNA expression in prefrontal cortex*
*in Parkinson's disease.* Exp Neurol, 2010. **226**(1): p. 207-17.
101.  Ihara, M., et al., *Association of vascular parkinsonism with impaired neuronal integrity in the*
*striatum.* J Neural Transm (Vienna), 2007. **114**(5): p. 577-84.
102.  van Nuland, A.J.M., et al., *GABAergic changes in the thalamocortical circuit in Parkinson's disease.*
Hum Brain Mapp, 2020. **41**(4): p. 1017-1029.
103.  Allen, P., et al., *Extrinsic and default mode networks in psychiatric conditions: Relationship to*
*excitatory-inhibitory transmitter balance and early trauma.* Neurosci Biobehav Rev, 2019. **99**: p.
90-100.
104.  Liu, Z., et al., *Tissue Specific Expression of Cre in Rat Tyrosine Hydroxylase and Dopamine Active*
*Transporter-Positive Neurons.* PLoS One, 2016. **11**(2): p. e0149379.
105.  Anders Björklund, S.B.D., *Dopamine neuron systems in the brain: an update.* Trends in
Neuroscience, 2007. **30**(5): p. 194-202.
106.  Ran, F.A., et al., *Genome engineering using the CRISPR-Cas9 system.* Nat Protoc, 2013. **8**(11): p.
2281-2308.
107.  Vouillot, L., A. Thelie, and N. Pollet, *Comparison of T7E1 and surveyor mismatch cleavage assays*
*to detect mutations triggered by engineered nucleases.* G3 (Bethesda), 2015. **5**(3): p. 407-15.
108.  Huang, Z.R., et al., *A Novel Potential Positron Emission Tomography Imaging Agent for Vesicular*
*Monoamine Transporter Type 2.* PLoS One, 2016. **11**(9): p. e0161295.
109.  Ding, Y.S., et al., *Pharmacokinetics and in vivo specificity of [11C]dl-threo-methylphenidate for the*
*presynaptic dopaminergic neuron.* Synapse, 1994. **18**(2): p. 152-60.
110.  Wadsworth, H., et al., *[(1)(8)F]GE-180: a novel fluorine-18 labelled PET tracer for imaging*
*Translocator protein 18 kDa (TSPO).* Bioorg Med Chem Lett, 2012. **22**(3): p. 1308-13.
111.  Paasonen, J., et al., *Functional connectivity under six anesthesia protocols and the awake*
*condition in rat brain.* Neuroimage, 2018. **172**: p. 9-20.
112.  Lopez-Gonzalez, F.J., et al., *QModeling: a Multiplatform, Easy-to-Use and Open-Source Toolbox for*
*PET Kinetic Analysis.* Neuroinformatics, 2019. **17**(1): p. 103-114.
113.  Schiffer, W.K., et al., *Serial microPET measures of the metabolic reaction to a microdialysis probe*
*implant.* J Neurosci Methods, 2006. **155**(2): p. 272-84.

114.    Logan, J., *Graphical analysis of PET data applied to reversible and irreversible tracers.* Nucl Med
1047            Biol, 2000. **27**(7): p. 661-70.
1048    115.    Judenhofer, M.S., et al., *Simultaneous PET-MRI: a new approach for functional and morphological
1049            imaging.* Nat Med, 2008. **14**(4): p. 459-65.
1050    116.    Matthew Brett, J.-L.A., Romain Valabregue, Jean-Baptiste Poline, *Region of interest analysis using
1051            an SPM toolbox*, in *8th Internation Conference on Functional Mapping of the Human Brain*. June 2-
1052            6, 2002: Sendai, Japan.
1053    117.    Rubinov, M. and O. Sporns, *Complex network measures of brain connectivity: uses and
1054            interpretations.* Neuroimage, 2010. **52**(3): p. 1059-69.
1055    118.    Ungerstedt, U. and G.W. Arbuthnott, *Quantitative recording of rotational behavior in rats after 6-
1056            hydroxy-dopamine lesions of the nigrostriatal dopamine system.* Brain Res, 1970. **24**(3): p. 485-93.
1057    119.    West, M.J., *Stereological methods for estimating the total number of neurons and synapses: issues
1058            of precision and bias.* Trends Neurosci, 1999. **22**(2): p. 51-61.

1059

1060

## Acknowledgments

We are very greatful to Feng Zhang and all members of the laboratory, Broad Institute of MIT and

Harvard, for the help and support with establishing the CRISPR gene knockdown experiments.

We thank the technical assistants Sandro Aidone, Daniel Bukala, Linda Schramm, Maren Harant,

Ramona Stremme, Elena Kimmerle and Johannes Kinzler. We also thank Ulla Samuelsson,

Ulrika Sparrhult-Björk, Dr. Ulrika Schagerlöf, Anneli Josefsson and Anna Hansen at Lund

University, Sweden for their technical support. Additionally, we acknowledge Dr. Julia Mannheim,

Dr. Andreas Schmid, Dr. Rebecca Rock, Ines Herbon, Dr. Neele Hübner, Dr. Andreas Dieterich,

Hans Jörg Rahm, Dr. Carsten Calaminus, Funda Cay, Kristin Patzwaldt, Laura Kübler, Marilena

Poxleitner, Sabrina Buss and Dominik Seyfried for their administrative, technical, and

experimental support at the Department of Preclinical Imaging and Radiopharmacy, Werner

Siemens Imaging Center, Eberhard Karls University, Tuebingen. This study is also part of the

PhD theses of Sabina Marciano and Tudor Mihai Ionescu.

## Funding

- German Research Foundation to KH

- Carl-Zeiss-Foundation to KH

- Deutscher Akademischer Austauschdienst to SM, KH

- Werner Siemens Foundation to BJP

## Author contributions

- Conceptualization: KH

- Methodology: SM, TMI

- Software: SM, TMI

- Validation: SM

- Formal analysis: SM, TMI

• Investigation: SM, TMI, AM, RSS, RYC

• Resources: BJP, DK

• Data curation: SM

• Writing – original draft: SM

• Writing – review and editing: SM, TMI, AM, DK, RSS, RYC, BJP, KH

• Supervision: KH

• Project administration: KH

• Funding acquisition: KH, BJP

**Competing interests**

The authors declare no conflict of interest.

**Data availability**

The original dataset will be made available upon request.

**Figures**

1098

**Fig. 1 *In vitro* validation of CRISPR/SaCas9-induced VMAT2 knockdown in rat primary cortical**
**neurons.** (**a**) AAV-SaCas9 and AAV-sgRNA expression vectors. (**b**) Experimental design for primary
neurons isolation and transduction. (**c**) VMAT2 immunostaining (red), nuclei labeled with DAPI (blue). GFP
(green) and HA-tag (white) indicate the expression of the sgRNA and SaCas9 vectors, respectively.

VMAT2 KD (arrows) is shown in neurons transduced with AAVs carrying SaCas9 and sgRNA-*Slc18a2*. KD,
knockdown; ITR, inverted terminal repeat; CMV, cytomegalovirus promoter; GFP, green fluorescent
protein; CAG, CMV enhancer/chicken β-actin promoter; NLS, nuclear localization signal; HA-tag,
hemagglutinin tag; polyA, polyadenylation signal; polyT, polytermination signal.

1107

1108

**Fig. 2 CRISPR/SaCas9-induced VMAT2 knockdown elicits postsynaptic changes but no nerve terminal loss or neuroinflammation in the adult rat brain.** (**a**) Schematic illustration of the experimental design. (**b**) Mean binding potential and uptake maps of control and VMAT2 KD rats co-registered to a rat brain atlas. (**c**) Binding potential values of individual control and VMAT2 KD rats in the left and right striatum. For [$^{18}$F]GE-180 uptake values normalized to the left striatum are shown. (**d**) A strong correlation between Δ [$^{11}$C]RAC and Δ [$^{11}$C]DTBZ binding is shown. (**e**) Ratio of striatal [$^{11}$C]RAC and [$^{11}$C]DTBZ binding shows prominent [$^{11}$C]RAC changes when a threshold of ~ 20% Δ [$^{11}$C]DTBZ binding is reached.

This threshold was set to separate the VMAT2 KD rats into *mild* and *moderate*. (**f**) *Mild* and *moderate* rats
could be differentiated based on the postsynaptic changes. *P< 0.01, **P< 0.001, Bonferroni-Sidak
corrected. Data are shown as boxplot with the median value (central mark), the mean value (plus sign),
interquartile range (boxes edges), and the extreme points of the distribution (whiskers). Control rats n= 8;
VMAT2 KD rats n= 12. *Mild*: $\Delta$ [$^{11}$C]DTBZ binding < 20%; *Moderate*: $\Delta$ [$^{11}$C]DTBZ binding ≥ 20%. [$^{11}$C]MP,
[$^{11}$C]methylphenidate; [$^{11}$C]RAC, [$^{11}$C]raclopride; LSTR, left striatum; RSTR, right striatum; CER,
cerebellum; KD, knockdown; DAT, dopamine transporter; TSPO, translocator protein.

1123

1124
**Fig. 3 CRISPR/SaCas9-induced VMAT2 knockdown impairs motor function.** (**a**) Schematic illustration
of the behavioral tests. (**b**) In the open field test, the distance travelled (m) by VMAT2 KD rats was reduced.
(**c**) Cylinder test. VMAT2 KD rats showed a reduction in the contralateral paw touches, compared with
controls. Rats performance in the cylinder test strongly correlated with VMAT2 expression changes
($\Delta$ [$^{11}$C]DTBZ binding) and corresponding changes in dopamine availability ($\Delta$ [$^{11}$C]RAC binding) (**d,e**). (**f**) In
the beam walk test, VMAT2 KD rats displayed a higher number of footslips to the left contralateral side
compared with control rats. (**g**) Body weight assessment 14 weeks after CRISPR/SaCas9 gene-editing
showed reduced body weight gain in VMAT2 KD compared with control rats. (**h,i**) Body weight gain
correlated with VMAT2 expression changes ($\Delta$ [$^{11}$C]DTBZ binding) and corresponding changes in
dopamine availability ($\Delta$ [$^{11}$C]RAC binding). (**j**) Spontaneous rotation in a novel spherical environment
showed increased CW rotations in VMAT2 KD rats compared with control rats. (**k**) Apomorphine-evoked
rotational behavior. VMAT2 KD rats displayed a higher number of CCW rotations compared with control
rats in the rotameter test. (**l,m**) Apomorphine-evoked rotations exhibited a strong correlation with VMAT2

expression changes ($\Delta$ [$^{11}$C]DTBZ binding) and changes in dopamine availability ($\Delta$ [$^{11}$C]RAC binding).
Data are shown as boxplot with the median value (central mark), the mean value (plus sign), interquartile
range (boxes edges), and the extreme points of the distribution (whiskers). $^{+}$*P*< 0.05, *\*P*< 0.01, *\*\*P*< 0.001.
Control rats n= 8; VMAT2 KD rats n= 12. CCW, counter-clockwise; APO, apomorphine; KD, knockdown;
[$^{11}$C]RAC, [$^{11}$C]raclopride. Illustrations in the figure were created with BioRender.com.

1143

1144

**Fig. 4 *Ex vivo* validation of the CRISPR/SaCas9-induced VMAT2 knockdown.** (**a**) Schematic illustration of the *ex vivo* analyses. (**b**) Immunofluorescence of nigral sections of control and VMAT2 KD rats confirmed the concomitant expression of SaCas9 and sgRNA. Staining for GFP (AAV-sgRNAs, green), HA-tag (AAV-SaCas9, white), and VMAT2 expression (red) for two exemplary rats is shown. VMAT2 expression was largely reduced in the SNc of VMAT2 KD rats. (**c,d**) TH expression in the ipsilateral and contralateral striatum and SNc of VMAT2 KD (n= 10) and control (n= 5) rats evidenced no cell loss. (**e**) VMAT2 immunohistochemistry in the SNc and striatum confirmed large protein reduction in the ipsilateral hemisphere of VMAT2 KD rats. (**f**) Striatal dopamine, normalized to total protein concentration, was reduced in the ipsilateral striatum of VMAT2 KD (n= 9), but not control (n= 6) rats. This reduction was paralleled by increased metabolic outcome (**g**). (**h,i**) Dopamine changes correlated with the VMAT2 KD extent and postsynaptic changes, deducted from [$^{11}$C]DTBZ and [$^{11}$C]RAC, respectively. (**j**) 5-HT content was unaltered in the striata of control (n= 6) and VMAT2 KD (n= 9) rats. Data are shown as boxplot with the

1157    median value (central mark), the mean value (plus sign), interquartile range (boxes edges), and the
1158    extreme points of the distribution (whiskers). $^{+}P< 0.05$, $*P< 0.01$. GFP, green fluorescent protein; HA-tag,
1159    hemagglutinin tag; KD, knockdown; TH, tyrosine hydroxylase; SN, substantia nigra; STR, striatum;
1160    [$^{11}$C]RAC, [$^{11}$C]raclopride; DA, dopamine; 5-HT, serotonin. Metabolites' and neurotransmitters' striatal
1161    content is reported in Table 3. Illustrations in the figure were created with BioRender.com.

1162

1163

**Fig. 5 Increased resting-state functional connectivity after CRISPR/SaCas9-induced VMAT2 knockdown.** (**a**) Schematic illustration of the experimental design. Group level correlation matrices of the DMN (**b**) and SMN (**c**) at baseline and after CRISPR/SaCas9-targeting for rats with a *mild* (left panel) and *moderate* (right panel) VMAT2 KD. In the *moderate* KD group rs-FC was increased between the right mPFC and right and left Hipp in the DMN (**b**) and between the contralateral SC and right and left Th, as well as between the left STR and right and left Th, in the SMN (**c**). (**d**) Internetwork rs-FC changes in the *mild* KD group indicated increased rs-FC between anterior regions of the DMN and the SMN (between the left OFC and STR, and between the right OFC and the left SC, and the right and left STR). Conversely, rs-FC was decreased between regions of the posterior DMN and the SMN (between the left RSC and right SC). In the *moderate* KD group, DMN-SMN rs-FC was increased. Brain graphs, right to the matrices, illustrate the nodes and edges (raw values) that demonstrated rs-FC changes to baseline (%). *$P$< 0.01. *Mild*: $\Delta$ [$^{11}$C]DTBZ binding < 20%, n= 13; *Moderate*: $\Delta$ [$^{11}$C]DTBZ binding ≥ 20%, n= 10. KD, knockdown; DMN, default-mode network; SMN, sensorimotor network. mPFC, medial prefrontal cortex; Hipp, hippocampus; SC, somatosensory cortex; Th, thalamus; STR, striatum; OFC, orbitofrontal cortex; RSC, retrosplenial cortex. Abbreviations of brain regions considered for the analysis of the fMRI data, including their respective volumes, are reported in Table 2.

1180

**Fig. 6 CRISPR/SaCas9-induced VMAT2 knockdown alters GABA signaling.** (**a**) [$^{11}$C]FMZ mean binding potential maps of *mild* and *moderate* VMAT2 KD rats co-registered to a rat brain atlas. Arrows and ROIs in coronal sections indicate brain regions of the DMN and SMN with altered [$^{11}$C]FMZ binding. [$^{11}$C]FMZ binding potentials from VOI-based analysis in DMN and SMN regions of rats with *mild* (**b**) and *moderate* (**c**) VMAT2 KD. [$^{11}$C]FMZ binding was decreased in the right PaC, SC, and Hipp of *mild* KD rats (**b**) and right PaC and SC of *moderate* KD rats (**c**). [+]$P<$ 0.05, [*]$P<$ 0.01, [**]$P<$ 0.001, Bonferroni-Sidak corrected. Data are shown as mean ± SD. *Mild*: Δ [$^{11}$C]DTBZ binding < 20%, n= 13; *Moderate*: Δ [$^{11}$C]DTBZ binding ≥ 20%, n= 10. KD, knockdown; [$^{11}$C]FMZ, [$^{11}$C]flumazenil; DMN, default-mode network; SMN, sensorimotor network. PaC, parietal cortex; Hipp, hippocampus; SC; somatosensory cortex. Abbreviations of brain regions, including their respective volumes, are reported in Table 2.

1193 **Tables**

1194 **Table 1: sgRNAs sequences.**

1195

1196

| SgRNAs | DNA Target Sequences 5'- 3' | PAM (NNGRRT) 5'- 3' |
|---|---|---|
| *Slc18a2* | CGATGAACAGGATCAGTTTGC | GCGAGT |
| *lacZ* | CCTTCCCAACAGTTGCGCAGC | CTGAAT |

1197

1198

1199 **Table 2: Brain regions included in the Paxinos rat brain atlas, including their respective volumes**
1200 **and abbreviations.**

| Brain region (ROI) | Hemisphere | ROI volume [mm$^3$] | Abbreviation |
|---|---|---|---|
| Striatum | left | 43.552 | STR |
| | right | | |
| Cingulate Cortex | left | 14.480 | CgC |
| | right | | |
| Medial Prefrontal Cortex | left | 6.304 | mPFC |
| | right | | |
| Motor Cortex | left | 32.608 | MC |
| | right | | |
| Orbitofrontal Cortex | left | 18.936 | OFC |
| | right | | |
| Parietal Cortex | left | 7.632 | PaC |
| | right | | |
| Retrosplenial Cortex | left | 18.920 | RSC |
| | right | | |
| Somatosensory Cortex | left | 71.600 | SC |
| | right | | |
| Hippocampus | left | 25.064 | Hipp |
| | right | | |
| Thalamus | left | 30.712 | Th |
| | right | | |

1201

**1202 Table 3: Dopamine, 3,4-Dihydroxyphenylacetic acid (DOPAC), homovanillic acid (HVA), and**
**1203 serotonin (5-HT) striatal content (nmol/mg) determined by HPLC.**

| Neurotransmitter / Metabolite | nmol/mg (mean ± SD) | |
|---|---|---|
| DA | Control | VMAT2 KD |
| STR-L | 244 ± 109 | 351 ± 168 |
| STR-R | 308 ± 166 | 100 ± 97 |
| DOPAC | Control | VMAT2 KD |
| STR-L | 34 ± 7 | 34 ± 11 |
| STR-R | 38 ± 4 | 27 ± 5 |
| HVA | Control | VMAT2 KD |
| STR-L | 24 ± 5 | 25 ± 5 |
| STR-R | 26 ± 6 | 19 ± 4 |
| 5-HT | Control | VMAT2 KD |
| STR-L | 17 ± 2 | 19 ± 8 |
| STR-R | 19 ± 6 | 23 ± 9 |

1204