# Bacterial supergroup specific "Cost" of *Wolbachia* infections in *Nasonia vitripennis*

Alok Tiwary[1], Rahul Babu[1], Ruchira Sen[2], Rhitoban Raychoudhury[1*]

1. Department of Biological Sciences, Indian Institute of Science Education and Research (IISER) Mohali, Knowledge City, Sector 81, SAS Nagar, Manauli, PO 140306, Punjab, India.
2. Sri Guru Gobind Singh College, Sector 26, Chandigarh-160019, India.

*Corresponding author: Rhitoban Raychoudhury, Department of Biological Sciences, Indian Institute of Science Education and Research (IISER) Mohali, Sector 81, Knowledge City, SAS Nagar, Manauli, PO 140306 (Punjab), India

Email: rhitoban@iisermohali.ac.in

## Abstract

The maternally-inherited endosymbiont, *Wolbachia*, is known to alter the reproductive biology

of its arthropod hosts for its own benefit. It is also known to induce both positive and negative

fitness effects in many hosts. Here we describe the effect of maintenance of two distinct

*Wolbachia* infections, one each from supergroups A and B, on the parasitoid host *Nasonia*

*vitripennis*. We compare the effect of *Wolbachia* infections on various traits between

uninfected, single A, single B, and double infected strains. Contrary to the previous reports, our

results suggest that there is a "significant cost" associated with the maintenance of *Wolbachia*

infections where traits like progeny family size, fecundity, longevity and rates of male copulation

is compromised in *Wolbachia* infected strains. The double infection of *Wolbachia* has more

detrimental impacts on the host as compared to single infections, while the supergroup B

infections elicits the most pronounced negative effect. These findings raise important questions

on the mechanism of survival and maintenance of these reproductive parasites in arthropod

hosts.

## Introduction

*Wolbachia* are maternally-inherited, obligatory intracellular endosymbionts of the order Rickettsiales (Hertig et al. 1927) that are found widely in arthropods and filarial nematodes (Bandi et al. 1992; Rousset et al. 1992; Weinert et al. 2015). These bacteria often alter host reproductive biology with mechanisms like male-killing, feminization, parthenogenesis, and cytoplasmic incompatibility (CI), to enhance their transmission (Werren et al. 2008)). While CI leads to an increase in the number of infected individuals in the population, male-killing, and feminization shifts the offspring sex ratio towards females, which is the transmitting sex for *Wolbachia*. Thus, *Wolbachia* increases the "fitness" of the infected hosts as it increases its own rate of transmission. While these cases highlight a parasitic effect of *Wolbachia*, there are several examples where no such effect is discernible (Hoffmann et al. 1996). Moreover, there are also examples where *Wolbachia* has now become a mutualist and offers specific and quantifiable benefits to its host. The most well-known example for this is the obligatory association of *Wolbachia* with its filarial nematode hosts like *Wuchereria bancrofti* (Fenn and Blaxter, 2006) where the mutualistic effect ensures the long-term persistence of *Wolbachia*. This shift from parasitic to mutualistic effect can also happen in facultative associations as seen in *D. simulans*, where within a span of just two decades, *Wolbachia* has evolved from a parasite to a mutualist (Weeks *et al.*, 2007). However, for the majority of *Wolbachia*-host associations, no such reports of mutualistic benefits exist and instead reveal many negative effects on the host. In addition to reproductive traits, many other life-history traits like longevity and developmental time are also known to be compromised. In *D. simulans*, *Wolbachia* infected females face a 10-20% reduction in fecundity (Hoffmann et al. 1990) whereas, infected males tend to produce fewer sperm cysts (Snook et al. 2000). *Wolbachia* infections have also been known to negatively impact the lifespan of its hosts in *D. melanogaster* (Min and Benzer, 1997) and *Aedes albopictus* (Kambris et al., 2009; McMeniman et al. 2009). In *Trichogramma kaykai and T. deion,* the infected (thelytokous) line shows reduced fecundity and adult emergence rates than the antibiotically cured (arrhenotokous) lines (Hohmann et al. 2001; Tagami et al. 2001). *Leptopilina heterotoma*, a *Drosophila* parasitoid, has adult survival rates, fecundity, and locomotor performance, of both sexes, severely compromised in *Wolbachia* infected strains

(Fleury et al. 2000). Preimaginal mortality, in insecticide-resistant *Wolbachia*-infected strains,

has been observed in both sexes of *Culex pipiens* (Duron et al. 2006). *Wolbachia* infections can

also result in a range of behavioral changes in *Aedes aegypti* where a significant proportion of

them fail to obtain blood meals with increasing age. This complex behavior is associated with an

altered phenotype where the infected females have a "bendy proboscis" as compared to their

uninfected counterparts (Turley et al. 2009).

These negative effects of *Wolbachia* on their hosts are not difficult to explain.  The presence of

bacteria within a host entails sharing of nutritional and other physiological resources (Kobayashi

and Crouch, 2009), especially with *Wolbachia* which are obligate endosymbionts and cannot

survive without cellular resources derived from their hosts (Foster et al. 2005; Slatko et al.

2010). For example, *Wolbachia* is known to compete with the host for key resources like

cholesterol and amino acids (Caragata et al. 2014). Therefore, *Wolbachia* infections can impose

severe nutritional demand on the host (Ponton et al. 2014). The precise molecular mechanisms

of many of these negative effects have not been ascertained and are generally ascribed to

partitioning off-host nutrients for its benefit (Ponton et al. 2014). However, it is known that

*Wolbachia* leads to antipathogenic effects in their hosts where the host resistance or tolerance

to the infection increases (Zug and Hammerstein, 2015). For example, *Wolbachia* can utilize the

IMD and Toll pathways (Pan et al. 2018) and increased ROS (Reactive Oxygen Species) levels in

*Wolbachia*-transfected *A. aegypti* mosquitoes, inhibiting the proliferation of Dengue virus (Pan

et al. 2012). Moreover, *Wolbachia* is also known to elevate ROS levels leading to tolerance

against *Drosophila* C viruses (Wong et al. 2011). Such immune defense requires additional

allocation of resources, which can affect other life-history traits of the host. This concept of a

"cost of immunity" is a well-established notion, which suggests a trade-off between immunity

and other life-history traits (Zuk and Stoehr, 2002). For example, elevated ROS levels negatively

effect many host traits like longevity, and fecundity (Dowling and Simmons, 2009; Monaghan et

al. 2009; Selman et al. 2012; Moné et al. 2014). Thus, there is sufficient evidence to conclude

that *Wolbachia* can have substantial negative effects on the physiology, nutrition, survival, and

overall fitness of its host.

In this study, we investigate what, if any, are the negative effects on the wasp *N. vitripennis* due

to its two resident CI inducing *Wolbachia* infections? *N. vitripennis,* with its cosmopolitan

distribution, has already been used to study *Wolbachia* distribution, acquisition, spread and

*Wolbachia*-induced reproductive manipulations (Werren *et al.*, 2008; Landmann, 2019).

However, the effect of the endosymbiont on the host life-history traits of this wasp remains

poorly understood with conflicting reports. In some *N. vitripennis* strains, *Wolbachia* was

reported to be the cause of enhanced fecundity (Stolk and Stouthamer, 1996), but a similar

effect was not observed in some other strains (Bordenstein and Werren, 2000). *N. vitripennis*

harbor two *Wolbachia* supergroup infections, one each from supergroup A and supergroup B

(Perrot-Minnot et al. 1996), and the presence of these two infections have been found in all

strains of *N. vitripennis* from continental North America to Europe and Russia (Raychoudhury et

al. 2010), indicating that it has reached fixation across the distribution of its host. The two

*Wolbachia* in *N. vitripennis* cause complete CI when present together, but single infections of

supergroup A *Wolbachia* causes incomplete CI while supergroup B infections still show

complete CI (Perrot-Minnot et al. 1996). In this study, we investigate the effects of *Wolbachia*

infections in a new strain of *N. vitripennis* recently obtained from the field. . This strain, like

other *N. vitripennis* strains, has two *Wolbachia* infections with one each from the supergroups A

and B. Sequencing of the five alleles from the well-established multi-locus strain typing system

(Baldo et al. 2006) reveal no sequence dissimilarity across the two infections (Prazapati,

personal communication) indicating this new strain is also infected by the same or very similar

*Wolbachia* infections that are present across the distribution of *N. vitripennis* (Raychoudhury et

al. 2010b). To compare supergroup specific effects, these two infections have been separated

into single infected lines of supergroup A and supergroup B *Wolbachia*. A comparative analysis

between the double infected, supergroup A infected, supergroup B infected and uninfected

strains revealed a consistent pattern of decreased longevity, quicker sperm depletion, and

reduced progeny family size for infected individuals. We have found supergroup specific

negative effects on the physiological traits of the host. While supergroup B infection has sever

reduction effects on the different traits, supergroup A infection show the negative effcts only on

few of the traits. Our results also show a higher *Wolbachia* density of supergroup B and double

infection strains across the different developmental stages of *N. vitripennis* males and females

as compared to supergroup A *Wolbachia*.

**Materials and Methods**

***N. vitripennis* strains used, their *Wolbachia* infections, and strain nomenclature**

The *N. vitripennis* strains used NV PU-14 was obtained from Mohali, Punjab, India, in 2015).NV

PU-14 was cured of *Wolbachia* by feeding the females with 1 mg/ml tetracycline in 10 mg/ml

sucrose solution for at least two generations (Breeuwer and Werren, 1990). The curing was

confirmed by PCR using supergroup-specific *ftsZ* primers (Baldo et al. 2006), and CI crosses

between the infected and uninfected strains. NV PU-14 also served as the source strain for

separating the two *Wolbachia* infections into single A and single B infected strains.

To separate the strains, we utilized relaxed selection on the females by repeatedly mating them

with uninfected males which were obtained by antibiotic curing of the same NV PU-14 strain.

Uninfected males do not have any sperm modification by *Wolbachia* which results in the

removal of any selection pressure on the females to maintain their *Wolbachia* infections.

Repeated mating with uninfected males was continued for ten generations till some of the

progenies were found to be infected with single A and single B supergroup infections. The single

infection status of these strains was confirmed by using supergroup-specific *ftsZ* gene PCR

primers (Baldo et al. 2006).

The preferred method of nomenclature of *Nasonia* strains and their *Wolbachia* infections

includes information of both supergroups as well as the host genotype. For example, [*w*NvitA

*w*NvitB]V-PU14 indicates that the host species is *N. vitripennis*, with NV-PU14 as the host

genotype, which has two *Wolbachia* infections, one each from supergroup A and supergroup B.

However, since we used only *N. vitripennis* strains in this study, the nomenclature has been

simplified by removing the species name. For example, the same strain will now be denoted as

*w*A*w*B(PU), and when cured of these infections, as 0(PU). The single *Wolbachia*-infected strains

used were designated as *w*A(PU) for the supergroup A-infected strain while *w*B(PU)  for the

supergroup B-infected strain. As the cured 0(PU) lines were in culture for three years, many of

the infected lines were cured again to obtain 'recently cured' lines to minimize the effects of

any host divergence that might have accumulated within them. These 'recently cured' lines

were named 0(*w*A PU), 0(*w*B PU), and 0(*w*A*w*B PU).

All the wasp strains were raised on *Sarcophaga dux* fly pupae with a generation time of 14-15

days at 25°C, 60% humidity, and a continuous daylight cycle.

## Sequential mating and sperm depletion of the males

To test the effect of *Wolbachia* on male reproductive traits like mating ability, individual males

were assayed for the number of copulations it can perform, as well as sperm depletion. As *N.*

*vitripennis* is haplodiploid, every successful mating will result in both female and male progenies

while an unsuccessful one will result in the production of exclusively male progenies. The males

used were obtained from virgin females hosted with one fly pupa for 24 hours and were not

given any external sources of nutrition (usually a mixture of sucrose in water) before the

experiment. Each male was then mated sequentially with virgin females from the same strain.

At the first sign of a male not completing the entire mating behavior (Jachmann and Assem

1996) it was given a rest for half an hour and was subjected to mating again until it stopped

mating altogether. The mated females were then given a day and each of them were hosted

with one fly pupa for 24 hours. The females were removed after 24 hours and the offspring

were allowed to emerge and then counted. The average number of copulations and the number

of copulations, before sperm depletion, were compared using the Mann-Whitney U test with a

significance level of 0.05.

## Host longevity, progeny family size, and fecundity

To test whether the presence of *Wolbachia* has any influence on longevity, emerging adult

wasps were kept individually in ria vials at 25°C, without any additional nutrition. Survival

following emergence was measured by counting the number of dead individuals every 6 hours.

Log Rank Statistics was used to identify differences between strains with a significance level of

0.05

To test for the effect of *Wolbachia* infections on the adult progeny family size of virgin and
mated females, each female was sorted at the pupal stage and separated into individual ria
vials. To enumerate the progeny size of mated females, some of these virgins were offered
single males from the same strain and observed till mating was successful. All the females were
then hosted individually with one fly pupa for 24 hours. These were kept at 25$^{\circ}$C for the
offspring to emerge which were later counted for progeny size by randomizing the ria vials in a
double-blind assay. The differences were compared using the Mann-Whitney U test with a
significance level of 0.05.

To investigate if *Wolbachia* female the fecundity, emerged females were hosted with one host
for 24 hours. The host pupa was placed in a foam plug so that only the head portion of the pupa
was exposed and available for the females to lay eggs. The female was removed after 24 hours
and the eggs laid were counted under a stereomicroscope (Leica M205 C). The differences in
fecundity were compared using the Mann-Whitney U test with a significance level of 0.05.

## Estimation of relative density of *Wolbachia* infections across different developmental stages of *N. vitripennis*

195    
196    

To collect the different developmental stages, females were hosted for 4 hours, (instead of 24
hours in the previous experiments), with one host to narrow down the developmental stages of
the broods. The larval and pupal stages (from Day 3 to Day 13 for males and from Day 8 to Day
14 for females) were collected every 24 hours. Larval stages for females were not done to avoid
any DNA contamination from the males as the two sexes are virtually indistinguishable at the
larval stages. Three replicates of ten larvae or pupae from the three strains *w*A(PU), *w*B(PU),
and *w*A*w*B(PU) were collected for each developmental stage. DNA extraction was done using
the phenol-chloroform extraction method, where samples were crushed in 200 µl of 0.5 M Tris-
EDTA buffer with 1% sodium dodecyl sulfate (SDS) and 2 µlof 22 mg/ml Proteinase K and
incubated overnight at 37°C. DNA was purified with buffer saturated phenol and chloroform-
isoamyl alcohol solution (24:1 ratio) and the DNA was precipitated overnight with isopropanol
at -20° C. The precipitated DNA pellet was dissolved in 60 µl nuclease-free water. The DNA
concentration of the samples was measured using a spectrophotometer (Nanodrop 2000®,

Thermo Scientific). The concentrations of all the samples were normalized to 200 ng/μl across

the different male and female developmental stage samples to be used for quantitative PCR.

CFX96 C1000® Touch Real-time qRT-PCR machine (BioRad) was used to assay the relative

density of *Wolbachia* across the strains. Amplification was done for the *Wolbachia* hcpA gene

(Forward Primer: 5'-CTTCGCTCTGCTATATTTGCTGC-3', Reverse Primer: 5'-

CGAATAATCGCAACCGAACTG-3'). The primers were tested to amplify both the *Wolbachia*

supergroup A and B strains. *Nasonia* S6K was used as the control gene (Bordenstein and

Bordenstein, 2011). The total reaction volume contained 5 μl of iTaq Universal SYBR® Green

supermix (BIO-RAD), 0.05 μl each of 10 μM forward and reverse primers, and 200 ng of

template DNA for a total volume of 10 μl for each reaction. Uninfected *N. vitripennis* DNA was

used as negative control while DNase-free water was used as non-template control. Reaction

conditions included an initial denaturation step of 95°C for 3 minutes followed by 39 cycles of

95°C for 10 seconds, annealing, and amplification at 52°C for 30 seconds for 40 cycles. All the

reactions were performed in triplicates and included a melt curve to check for non-specific

amplification. The relative density of *Wolbachia* was estimated by calculating the mean delta

threshold cycle ($\Delta C_q$), using the formula:

$$\Delta C_q = 1/3 \sum_{j=1}^{3} [1/3 \sum_{i=1}^{3} hcpA - 1/3 \sum_{i=1}^{3} S6K]$$

where i = number of technical replicates and j = number of biological replicates.

$1/\Delta Cq$ was calculated and plotted to show the *Wolbachia* density across different

developmental stages of males and females. The relative *Wolbachia* density was compared

across the three different strains using the Kruskal Wallis *H* test. Mann Whitney U test was used

to compare two different groups with a significance level of 0.05.

## Results

**The presence of *Wolbachia* reduces the lifespan of both males and females**

234 *Wolbachia* can compete with the host for available nutrition increasing nutritional stress which

235 can shorten the lifespan of the many hosts (Caragata et al. 2014, Macmenimal et al. 2009).

236 Therefore, we first investigated the effect of *Wolbachia* infections on the survival of both male

and female wasps. As figure 1 (top panel) indicates, there is a significant difference in the longevity of the infected males across the three infection types. The double-infected strain, $w$A$w$B(PU), starts to die-off first and has a significantly shorter life span compared to the two single infected strains {Log Rank Test, $\chi^2$=16.8, p < 0.001 with $w$A(PU) and $\chi^2$=33.9, p < 0.001 with $w$B(PU)}. Males from the uninfected strain, 0(PU), lived the longest and showed significantly longer lifespan compared to all the other infected strains {(Log Rank Test: $\chi^2$=76.3, p < 0.001 with $w$A$w$B(PU); $\chi^2$=33.0, p < 0.001 with $w$A(PU) and $\chi^2$=16.3, p < 0.001 for $w$B(PU)}. However, there was no significant difference in the lifespan between the two single infected strains $w$A(PU) and $w$B(PU) {(Log Rank Test, $\chi^2$=3.84, p = 0.05)}. Similarly, infected females (bottom panel) also showed a distinct reduction in life span when compared with the uninfected strain. But, unlike the males, the single A infected strain, $w$A(PU), showed the shortest lifespan {Log Rank Test: $\chi^2$=11.2, p < 0.001 with $w$A$w$B(PU), $\chi^2$=56.9, p < 0.001 with $w$B(PU) and $\chi^2$=31.1, p < 0.001, with 0(PU)} followed by $w$A$w$B(PU) {(Log Rank Test: $\chi^2$=20.4, p < 0.001 with $w$B(PU) and $\chi^2$=12.9, p < 0.001 with 0(PU)}. Curiously, 0(PU) and $w$B(PU) females showed similar lifespan {(Log Rank Test: $\chi^2$=0.24, p = 0.62)}. These results indicate a sex-specific variation as $w$A$w$B(PU) show the shortest lifespan among the males but $w$A(PU) showed the shortest among the females. Moreover, the effect of single infections on longevity also varied among the sexes as $w$A(PU) and $w$B(PU) males had similar lifespans, but in the case of females, it was $w$B(PU) and 0(PU) which had similar lifespans. But what is unambiguous from these results is that the uninfected strain lived the longest, irrespective of the sex of the hosts, indicating that the presence of *Wolbachia* is associated with the reduction in lifespan and is thus, costly for *N. vitripennis* to maintain.

**The presence of *Wolbachia* reduces the number of copulations a male can perform**

*Wolbachia* is known to be associated with a reduction in the number of mating a male can perform in *Ephestia kuehniella* (Sumida et al. 2017). To test whether similar effects are seen in *N. vitripennis*, we enumerated the number of copulations an individual male can perform across the infection types. As figure 2 indicates, there is indeed a reduction in the capacity of infected males to mate. When compared with the uninfected strain 0(PU), this reduction was most

pronounced in $w$B(PU) (MWU, U=30, p < 0.01), followed by $w$A$w$B(PU) and $w$A(PU) which

showed similar copulation capacities (MWU: U=11, p = 0.49). The uninfected 0(PU) strain

produced males with the highest number of copulations {MWU: U= 32, p < 0.05 for $w$A(PU) and

U= 27, p < 0.05 for $w$A$w$B(PU)}. Thus, the presence of *Wolbachia* substantially reduced the

number of copulations that a male could perform. However, complex phenotypes like

copulation can also be affected by the host genotype. Although all these four strains were

derived from the same field-collected isofemale line, but continuous culturing in the laboratory

can fix specific alleles within them resulting in interstrain divergence. Moreover, it is also known

that in *Nasonia* the effect of *Wolbachia*-induced phenotype is influenced by the hosts' genetic

background (Raychoudhury et al. 2012). Therefore, we cured all these infections again and

tested whether the host genotype, rather than *Wolbachia*, is causing this reduction in the

number of copulations by comparing these newly cured strains back with the previously used

uninfected strain, 0(PU). As figure 2 indicates, males from most of these re-cured strains

showed a marked and significant increase in the number of copulations that they could

perform. This number in the re-cured double infected strain, 0($w$A$w$B PU), increased to similar

levels as shown by 0(PU) (MWU: U=9.5, p = 0.2), while also showing a significant increase from

its infected counterpart $w$A$w$B(PU) (MWU: U=3, p < 0.05). Similarly, the number of copulations

for the re-cured single A supergroup infected strain, 0($w$A PU), also increased to the levels of

the uninfected strain 0(PU) (MWU: U=20, p = 0.76). However, this increase (from 77.5 ± 6.3 to

83.5 ± 12.9) with its infected counterpart was not significant (MWU: U=23, p = 0.48). The re-

cured single B supergroup infection, 0($w$B PU), was the only strain which did not revert to

uninfected levels (MWU: U=22, p < 0.05) despite showing a marginal increase (from 62.8 ± 6.6

to 78.2 ± 5.1; MWU: U=1, p < 0.05). However, what is evident is that the presence of *Wolbachia*

is also associated with a reduction in the capability of a male to mate. Furthermore, by recuring

the infected strains, we showed that this decrease is not due to the host genotype but is an

effect of *Wolbachia* present in these strains.

***Wolbachia* infected males deplete their sperm reserves faster than the uninfected ones**

*N. vitripennis* males are prospermatogenic (Boivin et al. 2005), where each male emerges with

their full complement of mature sperm and usually does not produce any more during the rest

of their lifespan (Chirault et al. 2016). Thus, if a single male is mated sequentially with as many

females it can mate with, it should eventually run out of this full complement of sperm resulting

in all-male progeny even after successful copulation. Sequential mating of the males from 0(PU),

*w*A(PU), *w*B(PU), *w*A*w*B(PU), and also the recently cured single and double infection strains,

0(*w*A PU), 0(*w*B PU), 0(*w*A*w*B PU) was done with females from the same strains. As figure 2

indicates, each male did run out of sperm at the tail end of the continuous mating and hence

producing only male progenies (shown by black dots). We looked at the number of mating done

by these males before sperm depletion to see if *Wolbachia* affects the sperm production in the

males. As shown in figure 3, the average number of daughter progenies reduced with the

number of mating (shown by the primary Y-axis on the left), indicating sperm depletion. Similar

to copulation numbers, *w*B(PU) males were also the quickest to deplete the sperm reserve

{MWU: U=27.5, p < 0.05 for *w*A(PU) and U=30, p < 0.01 compared with 0(PU)}. This was

followed by *w*A*w*B(PU) and *w*A(PU) (MWU: U=13, p = 0.7). However, the uninfected males from

0(PU) were the slowest to deplete their sperm {MWU: U= 35, p < 0.01 for *w*A(PU) and U= 24, p <

0.05 for  *w*A*w*B(PU)}. We again tested whether the host genotype, rather than *Wolbachia*, is

causing this rate of sperm depletion, by comparing it with the recently cured strains. As shown

in figure 3, the number of mating before sperms depletion increased for the recently cured

0(*w*A PU) up to the levels of 0(PU) (MWU: U=30, p = 0.06). However, this increase (from 48.14 ±

4.94 to 50.57 ± 9.41) was not significantly different from their infected counterpart *w*A(PU)

(MWU: U=16, p = 0.8). Rates of depletion for 0(*w*A*w*B PU) also increased up to the levels of

0(PU) (MWU: U=21, p = 0.66). Again, the cured strain 0(*w*B PU), increased marginally from

*w*B(PU) (from 41 ± 1.67 to 47.6 ± 6.0) (MWU: U=0, p < 0.05) but was still lower in number than

0(PU) (MWU: U=23, p < 0.05). These results indicate that the presence of *Wolbachia* has a

negative impact on the sperm number of the infected strains.

***Wolbachia* infected females produce fewer offspring**

*Wolbachia* is known to have a negative impact on the progeny size of its host (Hoffmann et al.
1990; Hohmann et al. 2001). To test whether a similar effect is seen in *N. vitripennis*, we
enumerated the adult progeny family sizes for both virgin and mated females for the four
different *Wolbachia* infection strains and their recently cured counterparts.

As figure 4 (top panel) indicates, there is a significant reduction in the average family sizes of all-
male broods produced by the virgin females of the *Wolbachia* infected strains. When compared
with the uninfected strain 0(PU), this reduction was most pronounced in *w*A*w*B(PU) (MWU:
U=21151.5, p < 0.01) followed by *w*B(PU) (MWU: U=19880.5, p < 0.05). *w*B(PU) and *w*A*w*B(PU)
showed similar progeny family size (MWU: U=18582.5, p = 0.29). However, *w*A(PU) showed
similar progeny family size when compared with 0(PU) (MWU: U=17191, p = 0.39) and *w*B(PU)
(MWU: U=17284, p = 0.26) but had larger all-male broods sizes than *w*A*w*B(PU) (MWU:
U=18252, p < 0.05).   We also compared the recently cured single and double infection strains
with the infected parental strains counterparts. The recently cured strains 0(*w*B PU) and
0(*w*A*w*B PU) showed marginal increase in the progeny family size which was comparable to the
uninfected strain 0(PU) {(MWU: U=11554, p = 0.29 with 0(*w*B PU); MWU: U=10798, p = 0.21
with 0(*w*A*w*B PU)}. However this marginal increase {from 29.84 ± 12.94 to 31.44 ± 10.30 for
0(*w*B PU) and from 28.99 ± 11.60 to 30.97 ± 11.48 for 0(*w*A*w*B PU)} was not significantly
different from their infected counterparts *w*B(PU) (MWU: U=9963.5, p = 0.34) and *w*A*w*B(PU)
(MWU: U=8650.5, p = 0.09). The recently cured strain 0(*w*A PU) did not show any increase in the
progeny family size and was comparable to *w*A(PU) (MWU: U=7085.3, p = 0.63) and 0(PU)
(MWU: U=8161, p = 0.22).

Similar to the virgin females, the reduction in the progeny family size was also observed for
average family sizes of mated females as shown by figure 4 (middle panel). When compared
with the uninfected strain 0(PU), this reduction is most pronounced in *w*B(PU) (MWU: 15582, p
< 0.01) and *w*A*w*B(PU) (MWU: U=16303, p < 0.01). However, *w*B(PU) and *w*A*w*B(PU) show
similar levels of progeny family sizes (MWU: 13732.5, p = 0.55). Interestingly, *w*A(PU) strain
shows similar levels of progeny family size as 0(PU) (MWU: U=11396.5, p = 0.86) but had larger
family sizes than *w*B(PU) (MWU: U=14080, p < 0.01) and *w*A*w*B(PU) (MWU: U=14682, p < 0.05).

Upon curing, the recently cured 0(*w*A*w*B PU) reverts back to the levels of the uninfected strain

0(PU) (MWU: U= 13295, p = 0.61) showing a significant increase from the infected counterpart

*w*A*w*B(PU) (MWU:U=12023, p < 0.05). The recently cured strain 0(*w*A PU) did not show any

increase from the infected counterpart *w*A(PU) (MWU: U=8385.5 p = 0.69) and was comparable

to 0(PU) (MWU: U=9022.5, p = 0.5). However, recently cured strain 0(*w*B PU) did not increase to

the levels of the uninfected strain 0(PU) (MWU: U=16782, p < 0.05). The marginal increase in

the progeny family size (from 28.72 ± 10.46 to 29.97 ± 8.59) was not significantly different from

the parent strain *w*B(PU) (MWU: U=13854, p = 0.47).

To know whether the difference in progeny family size of the mated females is because of the

female or male progenies or both, we compared the number of female and male progenies

separately between the four strains (figure 4 bottom panel). No difference was observed in the

number of males produced by the mated females. However, the number of female progenies

produced followed the same trend as that of the progeny family size of the mated females.

When compared to the uninfected strain 0(PU), *w*B(PU) and *w*A*w*B(PU) showed the least

number of female progenies {(MWU: 15964, p < 0.01 with *w*B(PU) and U=16283, p < 0.01 with

*w*A*w*B(Pu)}. *w*B(PU) and *w*A*w*B(PU) showed similar levels of female progenies (MWU: 13392, p =

0.33). Again, *w*A(PU) strain showed similar levels of female progenies as 0(PU) (MWU: U=11543,

p = 0.98) but showed higher female progeny number than *w*B(PU) and *w*A*w*B(PU) {(MWU:

U=14201, p < 0.01 with *w*B(PU) and MWU: U=14372, p < 0.05 with *w*A*w*B(PU)}. Upon curing,

the recently cured 0(*w*A*w*B PU) reverts back to the levels of the uninfected strain 0(PU) (MWU:

U= 13545, p = 0.42) showing a significant increase from the infected counterpart *w*A*w*B(PU)

(MWU:U=12331, p < 0.039). The recently cured strain 0(*w*A PU) did not show any increase from

the infected counterpart *w*A(PU) (MWU: U=8468 p = 0.79) and was also comparable to 0(PU)

(MWU: U=9330, p = 0.84). However, recently cured strain 0(*w*B PU) did not increase to the

levels of the uninfected strain 0(PU) (MWU: U=16749.5, p < 0.01).

**_Wolbachia_ negatively impacts the fecundity of the infected females**

The difference in progeny family sizes between the different infection strains can either be at

the level of eggs being laid by the females or there can be mortality at the larval or pupal stages.

To test this, we looked at the fecundity of both virgin and mated females for the four strains.

Among the virgin females (figure 5 top panel), when compared to the uninfected strain 0(PU),

wAwB(PU) had the least fecundity (MWU: U=5042, p < 0.01). However, there was no significant

difference between 0(PU) and wA(PU) (MWU: U=2632.5, p = 0.34) , and also between 0(PU) and

wB(PU)(MWU: U=2052.5, p = 0.71).

Amon the mated females (figure 5 bottom panel), when compared to the uninfected strain

0(PU), wAwB(PU)  had the least fecundity (MWU: U=2461.5, p < 0.01). This was followed by

wB(PU) having similar fecundity as that of wAwB(PU) (MWU: U=1479.5, p = 0.24) and wA(PU)

(MWU: U=2822.5, p = 0.22). wA(PU) showed higher fecundity than wAwB(PU) (MWU: U=2388, p

< 0.05) but was similar to the uninfected strain 0(PU) (MWU: U=3799.5, p = 0.75) .

The double infection strain wAwB(PU) showing the least fecundity both in virgin and mated

females suggest an effect of *Wolbachia* on egg production in females. The fecundity assay also

establishes that the difference in adult progeny size can be because of the difference in the

fecundity of the females.

**Relative *Wolbachia* density in single and multiple *Wolbachia* infection strains**

*Wolbachia* density has a major role to play in expressing the effects of the infection on host

biology (Hoffmann et al. 1996; Min and Benzer, 1997). An increase in cellular *Wolbachia* titers

are often associated with greater expression of their effects also has an effect on the level of CI

as suggested by the bacterial density model of CI (Breeuwer and Werren, 1993). Thus to

understand the effect of *Wolbachia* density on the host, once these are present as single and

double infections, we estimated the temporal pattern of *Wolbachia* levels across different

developmental stages of *N. vitripennis* males and females. In the case of males (figure 6, top

panel) wA(PU) had the lowest *Wolbachia* density across different larval and pupal

developmental stages when compared with wB(PU) (MWU: U=11, p < 0.01) and wAwB(PU)

(MWU: U=12, p < 0.01). However, no difference was observed between wB(PU) and wAwB(PU)

(MWU: U=51, p = 0.56).

In the case of females $w$A(PU), showed lower levels of *Wolbachia* when compared to $w$A$w$B(PU)

(MWU: U=8, p < 0.05). However, no difference was observed between $w$A(PU) and $w$B(PU)

(MWU: U=12, p = 0.12) and also $w$B(PU) and $w$A$w$B(PU) (MWU: U=19, p = 0.5).

**Discussion**

*Wolbachia* is known to negatively impact the physiological traits of the vast majority of its host

population. The present study suggests similar detrimental effects, or a "cost", associated with

the maintenance of *Wolbachia* infection in *N. vitripennis*. This cost is borne by both sexes. Our

results (summarized in Supplementary Table S1) suggest that the presence of both single and

multiple *Wolbachia* infections can be detrimental for both sexes for traits such as longevity

(figure 1) where both infected males and females showed reduced lifespan. In the case of

males, the infected strains $w$A(PU), $w$B(PU), and $w$A$w$B(PU) had fewer copulations (figure 2) but

quicker sperm depletion (figure 3). In most cases, these negative effects disappeared with the

removal of *Wolbachia*, as shown by the males of the recently cured strains 0($w$A PU) and

0($w$A$w$B PU). However, 0($w$B PU) did not revert to the levels of 0(PU). Similarly, in the case of

females (figure 4) the infected strains $w$B(PU) and $w$A$w$B(PU) had lesser progeny when

compared to the 0(PU). This difference was not due to the number of males produced but in the

fewer number of daughters. However, $w$A(PU) females, as an exception, had their progeny

family sizes comparable to 0(PU). Again, the recently cured strains showed a significant increase

in the progeny family sizes suggesting that the negative effects are due to the presence of

*Wolbachia*. The differences found in the progeny sizes were at the level of the number of eggs

being laid by the infected females (figure 5) indicating that the negative effects of *Wolbachia*

manifested themselves even before the egg-laying stage. However, the egg to larval to pupal

stage mortality could also have had an effect on the progeny family sizes but these were not

assayed.

Our data also suggests an additive or synergistic effect of the presence of the two different

*Wolbachia* supergroups in the double infection strain $w$A$w$B(PU). Evidence of such effects can

be seen in traits like male longevity (additive effect) where the deficit in longevity for $w$A$w$B(PU)

is the total of the deficits of $w$A(PU) and $w$B(PU). Similarly, for traits like female longevity, virgin

female progeny family size, and female fecundity, reduction effect on strain *w*A*w*B(PU) appears to be a combined effect of *w*A(PU) and *w*B(PU) (synergistic effect).

The negative fitness effects of CI-inducing *Wolbachia* (summarized in Supplementary Table S2) have been reported in arthropod hosts like *Drosophila* and mosquitoes. These traits include reproductive traits like fecundity, sperm cyst numbers, etc., and many other physiological traits like longevity and developmental time, which are known to be compromised. In *D. simulans*, *Wolbachia*-infected females face a 10-20% reduction in fecundity especially in laboratory strains (Hoffmann et al. 1990). *D. simulans* males infected with *Wolbachia* tend to produce fewer sperm cysts (Snook et al. 2000). *Wolbachia* infections have also been known to negatively impact the lifespan of its hosts in *D. melanogaster* (Min and Benzer, 1997) and *Aedes albopictus* (Kambris et al. 2009; McMeniman et al. 2009). However, many reports also suggest positive fitness effects because of the presence of *Wolbachia* (Poinsot and Mercot, 1997; Fry et al. 2004; Weeks et al. 2007). Similar host-specific effects have been reported in some strains of *Nasonia vitripennis* where the presence of *Wolbachia* led to enhanced fecundity (Stolk and Stouthamer 1996). However, a similar effect was not observed in other strains of *N. vitripennis* (Bordenstein and Werren, 2000). This suggests that the varied effects of *Wolbachia* can also be dependent on host strains.

*Wolbachia*-induced CI in arthropod host and nutritional competition raises important questions on the maintenance of these endosymbionts over evolutionary time scales. *Wolbachia* competes with the host for nutritional benefits like essential amino acids (Caragata et al. 2014) and hence imposes severe nutritional demands on the host (Ponton et al. 2014). Theoretical studies predict that these maternally inherited reproductive parasites should eventually turn mutualistic for long-term associations with their hosts (Prout 1994; Turelli 1994). Various theories on endosymbiosis also support the vertical transmission of endosymbionts, as the spread of the symbiont depends on the reproductive success of the host itself (Anderson and May, 1982; Ewald 1987; Bull et al. 1991; Lipsitch et al. 1995). Hence mutualism is the preferred way for endosymbionts to ensure their existence. Moreover, if there are indeed some adverse effects of maintaining *Wolbachia,* then hosts would be under strong selection pressure to

develop immunity against them. Evidence suggests that there are examples of such emergence

of host genetic factors upon *Wolbachia* infection (Zug and Hammerstein, 2015) in hosts like

*Drosophila* and mosquitoes. Host suppressor alleles have been identified conferring resistance

against feminizing and male-killing *Wolbachia* (Rigaud et al. 1999; Hornett et al. 2006).

However, no such host genetic factors have been found for CI-inducing *Wolbachia*, especially in

the case of *N. vitripennis*. A possible explanation for the maintenance of these multiple

infections comes from the high efficiency of transmission of these infections in *N. vitripennis*,

where these infections also show nearly 100% transmission as well as CI (Breeuwer and Werren,

1990). For many *Wolbachia*-arthropod associations, the pathogenicity and transmission

efficiency is mediated by bacterial titers (Correa and Ballard, 2016). Our results also show higher

*Wolbachia* density for double infection strain, which can facilitate its transmission and

maintenance in the host.

Another possibility can be that these *Wolbachia* infections in *N. vitripennis* are relatively recent,

evidence of which comes from the rapid spread of *Wolbachia* across populations (Raychoudhury

et al. 2010). These recent infections, although bearing a cost on the host at present, might

eventually lead to the host immunity against the infection.

Reproductive parasites like *Wolbachia* can also facilitate their transmission by increasing the

fitness of the infected individuals in the population against uninfected individuals (Dobson et al.

2004; Fry et al. 2004). The double infection strain, which is compatible with all the other strains,

can select against single *Wolbachia* infections (Perrot-Minnot et al. 1996). This fitness

advantage because of multiple *Wolbachia* infections can also be a plausible explanation for the

persistence of the host with *Wolbachia* even though it incurs a cost for maintenance of

*Wolbachia* on the host.

Our data suggest a significant reduction in the reproductive traits of both male and female

hosts. The fitness of adults and offspring depends on the allocation of limiting environmental

factors like nutrition (Trivers 1974). Many reports suggest that lack of nutrition results in smaller

progenies, a lesser number of eggs, impaired offspring development and metabolism (Telang

and Wells, 2004; Matzkin et al. 2013). With the presence of an endosymbiont like *Wolbachia*,

competing with the host for nutritional resources, a reduction in the number of female
offspring, a reduction in the number of copulations performed by the infected males, etc, can
be one of the implications of the nutritional stress caused by *Wolbachia* infection.

The cost of *Wolbachia* maintenance is in direct correlation with the density of *Wolbachia* strains
present in the host. Thus, in the case of females, the *w*A*w*B(PU), which shows a higher bacterial
load than single infections also has more reduced fecundity and longevity. Similarly, in the case
of males, the *w*A*w*B(PU) strain shows reduced number of copulations and the number of
daughters produced in sequential mating. However, *w*A(PU) showed minimal effect on the host,
the longevity of females being an exception. *w*A(PU) females had similar family sizes and
fecundities to that of 0(PU). Similarly, in the case of males, *w*A(PU) showed copulation rates
comparable to 0(PU) males. A possible explanation for this can be the relatively low density of
*w*A(PU) across the different developmental stages (figure 6). Our results also showed relatively
low levels of *Wolbachia* in *w*A(PU) strain, as compared to the other infections, which can result
in a less severe effect on male and female traits. Moreover, supergroup A infection strains are
known to have relatively higher levels of phage density (Bordenstein et al. 2006). According to
this phage density model (Bordenstein et al. 2006), higher phage density has an impact on the
level of CI caused by supergroup A strains, where it leads to a significant reduction in its CI
phenotype (Perrot-Minnot et al. 1996; Bordenstein et al. 2006).

Our results conform with the previous reports of the positive correlation between *Wolbachia*
abundance and the level of CI induced (Ijichi et al. 2002; Kondo et al. 2002; Bordenstein et al.
2006). *w*B(PU) *Wolbachia* showed complete CI and had higher *Wolbachia* titers in both males
and females, which was also comparable to that of the *w*A*w*B(PU), whereas *w*A(PU) had the
lowest *Wolbachia* titers among the three strains and showed incomplete CI.

*w*B(PU) shows strong deleterious effects on the host, similar to the double infection strain. The
observations can again be explained by the presence of high *Wolbachia* density in the case of
both males and females, which is comparable to that of the *w*A*w*B(PU). This also suggests that
the effects of resident *Wolbachia* infection can also be specific to particular *Wolbachia*

supergroup infection. Thus, such supergroup-specific effects of *Wolbachia* can be investigated in
further studies.

## Acknowledgments

515 We thank the Indian Institute of Science Education and Research (IISER) Mohali for the funding
516 and graduate fellowship for AT.

## Author Contribution

518 AT conceived the idea and planned the study. AT did the experiments. AT and RB collected the
519 data. AT analyzed the data. AT, RS and RR contributed to data representation and statistical
520 analysis. AT and RR wrote the manuscript.

## Competing Interests

522 The authors declare no conflict of interest.

## Data Archiving

524 The raw data for all the experiments has been archieved at Dryad with DOI:
525 10.5061/dryad.w0vt4b8s9

## References

Assem J V D and Jachmann F (2009). A Causal Ethological Analysis of the Courtship Behaviour of
an Insect (The Parasitic Wasp Nasonia vitripennis , Hym., Pteromalidae). *Behaviour* 133:
1051–1075.

Anderson RM, May RM (1982). Coevolution of hosts and parasites. *Parasitology* 85: 411-426

Baldo L, Hotopp JCD, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, *et al.* (2006). Multilocus
sequence typing system for the endosymbiont Wolbachia pipientis. *Appl Environ Microbiol*
72: 7098–7110.

Bandi C, Anderson TJC, Genchi C, Blaxter ML (1998) Phylogeny of Wolbachia in filarial

nematodes *Proc. R. Soc. Lond. B* 265, 2407-2413

Boivin G, Jacob S, Damiens D (2005). Spermatogeny as a life-history index in parasitoid wasps.

*Oecologia* **143**: 198–202.

Bordenstein SR, Bordenstein SR (2011). Temperature affects the tripartite interactions between

bacteriophage WO, Wolbachia, and cytoplasmic incompatibility. *PLoS One* **6**.

Bordenstein SR, Marshall ML, Fry AJ, Kim U, Wernegreen JJ (2006). The tripartite associations

between bacteriophage, Wolbachia, and arthropods. *PLoS Pathog* **2**: 384–393.

Bordenstein SR, Werren JH (2000). Do Wolbachia influence fecundity in Nasonia vitripennis?

*Heredity (Edinb)* **84**: 54–62.

Breeuwer JAJ, Werren JH (1990). Microorganisms associated with chromosome destruction and

reproductive isolation between two insect species. *Nature* 346: 558–560.

Breeuwer JAJ, Werren JH (1993). Cytoplasmic Incompatibility and Bacterial Density in Nasonia

vitripennis *Genetics* 135 565-574.

Bull JJ, Molineux IJ, Rice WR (1991). Selection of benevolence in a host-parasite system.

*Evolution (N Y)* 45: 875–882.

Caragata EP, Rancès E, O'Neill SL, McGraw EA (2014). Competition for Amino Acids Between

Wolbachia and the Mosquito Host, Aedes aegypti. *Microb Ecol* 67: 205–218.

Chirault M, van de Zande L, Hidalgo K, Chevrier C, Bressac C, Lécureuil C (2016). The spatio-

temporal partitioning of sperm by males of the prospermatogenic parasitoid Nasonia

vitripennis is in line with its gregarious lifestyle. *J Insect Physiol* 91–92: 10–17.

Correa CC, Ballard JWO (2016). Wolbachia Associations with Insects: Winning or Losing Against a

Master Manipulator. *Front Ecol Evol* 3.

Dobson SL, Rattanadechakul W, Marsland EJ (2004). Fitness advantage and cytoplasmic

incompatibility in Wolbachia single- and superinfected Aedes albopictus. *Heredity (Edinb)*

93: 135–142.

Dowling DK, Simmons LW (2009). Reactive oxygen species as universal constraints in life-history
evolution. *Proc R Soc B Biol Sci* 276: 1737–1745.

Duron O, Labbé P, Berticat C, Rousset F, Guillot S, Raymond M, *et al.* (2006). High Wolbachia
Density Correlates With Cost of Infection for Insecticide Resistant Culex Pipiens
Mosquitoes. *Evolution (N Y)* 60: 303.

EWALD PW (1987). Transmission Modes and Evolution of the Parasitism-Mutualism Continuum.
*Ann N Y Acad Sci* 503: 295–306.

Fenn K, Blaxter M (2006). Wolbachia genomes: Revealing the biology of parasitism and
mutualism. *Trends Parasitol* 22: 60–65.

Fleury F, Vavre F, Ris N, Fouillet P, Boulétreau M (2000). Physiological cost induced by the
maternally-transmitted endosymbiont Wolbachia in the Drosophila parasitoid Leptopilina
heterotoma. *Parasitology* 121: 493–500.

Foster J, Ganatra M, Kamal I, Ware J, Makarova K, Ivanova N, *et al.* (2005). The Wolbachia
genome of Brugia malayi: Endosymbiont evolution within a human pathogenic nematode.
*PLoS Biol* 3: 0599–0614.

Fry AJ, Palmer MR, Rand DM (2004). Variable fitness effects of Wolbachia infection in
Drosophila melanogaster. *Heredity (Edinb)* 93: 379–389.

Hertig M, Wolbach S B (1924). Studies on Rickettsia-like Microorganisms in Insects. *J.Med. Res*
44: 329–374.7

Hoffmann AA, Clancy D, Duncan J (1996). Naturally-occurring Wolbachia infection in Drosophila
simulans that does not cause cytoplasmic incompatibility. *Heredity (Edinb)* 76: 1–8.

Hoffmann AA, Turelli M, Harshman LG (1990). Factors affecting the distribution of cytoplasmic
incompatibility in Drosophila simulans. *Genetics* 126: 933–948.

Hohmann CL, Luck RF, Stouthamer R (2001). Effect of Wolbachia on the Survival and
Reproduction of Trichogramma kaykai Pinto & Stouthamer (Hymenoptera:

Trichogrammatidae). *Neotropical Entomology* 30(4): 607-612

Hornett EA, Charlat S, Duplouy AMR, Davies N, Roderick GK, Wedell N, *et al.* (2006). Evolution of

male-killer suppression in a natural population. *PLoS Biol* 4: 1643–1648.

Ijichi N, Kondo N, Matsumoto R, Shimada M, Ishikawa H, Fukatsu T (2002). Internal

spatiotemporal population dynamics of infection with three Wolbachia strains in the adzuki

bean beetle, Callosobruchus chinensis (Coleoptera: Bruchidae). *Appl Environ Microbiol* 68:

4074–4080.

Kambris Z, Al E (2009). Immune Activation by Life-Shortening Wolbachia and Reduced Filarial

Competence in Mosquitoes. *Science* 326: 134–136.

Kobayashi DY, Crouch JA (2009). Bacterial/Fungal interactions: from pathogenes to mutualistic

endosymbionts. *Annu Rev Phytopathol* 47: 63–82.

Kondo N, Ijichi N, Shimada M, Fukatsu T (2002). Prevailing triple infection with Wolbachia in

Callosobruchus chinensis (Coleoptera: Bruchidae). *Mol Ecol* 11: 167–180.

Landmann F (2019). The Wolbachia Endosymbionts. *Microbial Spectrum, American Society of*

*Microbiology*, 7 (2).

Lipsitch M, Nowak MA, Ebert D, May RM (1995). The population dynamics of vertically and

horizontally transmitted parasites. *Proc. R. Soc. Lond. B* 260, 321-327

Matzkin LM, Johnson S, Paight C, Markow TA (2013). Preadult Parental Diet Affects Offspring

Development and Metabolism in Drosophila melanogaster. *PLoS One* 8: 1–6.

McMeniman CJ, Lane R V, Cass BN, Fong AWC, Sidhu M, Wang YF, *et al.* (2009). Stable

introduction of a life-shortening Wolbachia infection into the mosquito Aedes aegypti.

*Science* 323: 141–144.

Min KT, Benzer S (1997). Wolbachia, normally a symbiont of Drosophila, can be virulent, causing

degeneration and early death. *Proc Natl Acad Sci U S A* 94: 10792–10796.

Monaghan P, Metcalfe NB, Torres R (2009). Oxidative stress as a mediator of life history trade-

offs: Mechanisms, measurements and interpretation. *Ecol Lett* 12: 75–92.

Moné Y, Monnin D, Kremer N (2014). The oxidative environment: A mediator of interspecies
communication that drives symbiosis evolution. *Proc R Soc B Biol Sci* 281.

Pan X, Pike A, Joshi D, Bian G, McFadden MJ, Lu P, *et al.* (2018). The bacterium Wolbachia
exploits host innate immunity to establish a symbiotic relationship with the dengue vector
mosquito Aedes aegypti. *ISME J* 12: 277–288.

Pan X, Zhou G, Wu J, Bian G, Lu P, Raikhel AS, *et al.* (2012). Wolbachia induces reactive oxygen
species (ROS)-dependent activation of the Toll pathway to control dengue virus in the
mosquito Aedes aegypti. *Proc Natl Acad Sci U S A* 109.

Perrot-Minnot M-JJ, Guo LR, Werren JH (1996). Double Infections with Wolbachia in the
Parasitic Wasp. *Genetics* 143: 961–972.

Poinsot D, Mercot H (1997). Wolbachia infection in Drosophila simulans: Does the female host
bear a physiological cost? *Evolution (N Y)* 51: 180–186.

Ponton F, Wilson K, Holmes A, Raubenheimer D, Robinson KL, Simpson SJ (2014).
Macronutrients mediate the functional relationship between Drosophila and Wolbachia.
*Proc R Soc B Biol Sci* 282.

Prout T (1994). Some evolutionary possibilities for a microbe that causes incompatibility in its
host. *Evolution (N Y)* 48: 909–911.

Raychoudhury R, Grillenberger BK, Gadau J, Bijlsma R, Van De Zande L, Werren JH, *et al.* (2010).
Phylogeography of Nasonia vitripennis (Hymenoptera) indicates a mitochondrial-Wolbachia
sweep in North America. *Heredity (Edinb)* 104: 318–326.

Rigaud T, Moreau J, Juchault P (1999). Wolbachia infection in the terrestrial isopod Oniscus
asellus: Sex ratio distortion and effect on fecundity. *Heredity (Edinb)* 83: 469–475.

Rousset F, Bouchon D, Pintureau B, Juchault P, Solignac M (1992). Wolbachia endosymbionts
responsible for various alterations of sexuality in arthropods. *Proc R Soc B Biol Sci* 250: 91–

98.

Selman C, Blount JD, Nussey DH, Speakman JR (2012). Oxidative damage, ageing, and life-history
637           evolution: Where now? *Trends Ecol Evol* 27: 570–577.

638     Slatko BE, Taylor MJ, Foster JM (2010). The Wolbachia endosymbiont as an anti-filarial
639           nematode target. *Symbiosis* 51: 55–65.

640     Snook RR, Cleland SY, Wolfner MF, Karr TL (2000). Offsetting effects of Wolbachia infection and
641           heat shock on sperm production in Drosophila simulans: Analyses of fecundity, fertility and
642           accessory gland proteins. *Genetics* 155: 167–178.

643     Stolk C, Stouthamer R (1996).  Influence of a Cytoplasmic Incompatibility-Inducing Wolbachia on
644           the fitness of the parasitoid wasp Nasonia vitripennis. *Proc. Exper. and Appl. Entomol.*
645           *N.E.V. Amsterdam* 7

646     Sumida Y, Katsuki M, Okada K, Okayama K, Lewis Z (2017). Wolbachia induces costs to life-
647           history and reproductive traits in the moth, Ephestia kuehniella. *J Stored Prod Res* 71: 93–
648           98.

649     Tagami Y, Miura K, Stouthamer R (2001). How does infection with parthenogenesis-inducing
650           Wolbachia reduce the fitness of Trichogramma? *J Invertebr Pathol* 78: 267–271.

651     Telang A, Wells MA (2004). The effect of larval and adult nutrition on successful autogenous egg
652           production by a mosquito. *J Insect Physiol* 50: 677–685.

653     Trivers R (1974). Parent-Offspring Conflict Parent-Offspring Conflict. American Zoologist 14:249-
654           264

655     Turelli M (1994). Evolution of Incompatibility-Inducing Microbes and Their Hosts. *Evolution* 48:
656           1500–1513.

657     Turley AP, Moreira LA, O'Neill SL, McGraw EA (2009). Wolbachia infection reduces blood-
658           feeding success in the dengue fever mosquito, Aedes aegypti. *PLoS Negl Trop Dis* **3**: 1–6.

Weeks AR, Turelli M, Harcombe WR, Reynolds KT, Hoffmann AA (2007). From parasite to

mutualist: Rapid evolution of Wolbachia in natural populations of Drosophila. *PLoS Biol* 5:

0997–1005.

Weinert LA, Araujo-Jnr E V., Ahmed MZ, Welch JJ (2015). The incidence of bacterial

endosymbionts in terrestrial arthropods. *Proc R Soc B Biol Sci* 282.

Werren JH, Baldo L, Clark ME (2008). Wolbachia: master manipulators of invertebrate biology.

*Nat Rev Microbiol* 6: 741–51.

Wong ZS, Hedges LM, Brownlie JC, Johnson KN (2011). Wolbachia-mediated antibacterial

protection and immune gene regulation in Drosophila. *PLoS One* 6.

Zug R, Hammerstein P (2015). Wolbachia and the insect immune system: What reactive oxygen

species can tell us about the mechanisms of Wolbachia-host interactions. *Front Microbiol*

6: 1–16.

Zuk M, Stoehr AM (2002). Immune defense and host life history. *Am Nat* 160.

**Figure legends**

**Figure 1. *Wolbachia* infected males and females show reduced lifespan**

Top panel: Lifespan of males. Sample sizes for the four infections 0(PU), *w*A(PU), *w*B(PU) and

*w*A*w*B(PU) were n=98, n=95, n=94 and n=95 respectively.

Bottom Panel: Lifespan of females. Sample sizes for the four infections 0(PU), *w*A(PU), *w*B(PU)

and *w*A*w*B(PU) were n=73, n=96, n=95 and n=95 respectively.

Statistical significance was tested using Log Rank statistics with p < 0.05.

**Figure 2. *Wolbachia* infected males show a reduction in the number of copulations**

Males from different *Wolbachia* infections were mated sequentially until each of them stopped

mating (please see Materials and Methods for details). 0(PU) could perform the highest number

of copulations while *w*B(PU) the least. The results show that the presence of *Wolbachia* is

associated with the reduction in the number of copulations a male can perform. The figure also

shows whether the progenies of these sequential copulations produce any daughters or not, as

a measure of sperm depletion. The details of sperm depletion are shown in figure 3. Sample

sizes for the strains 0(PU), *w*A(PU), 0(*w*A PU), *w*B(PU), 0(*w*B PU), *w*A*w*B(PU) and 0(*w*A*w*B PU)

were n=7, n=7, n=7, n=6, n=5, n=6 and n=7 respectively.

**Figure 3. *Wolbachia* infected males deplete their sperm faster than the uninfected males**

The Y-axis in black on the left of each figure represents the percentage of daughters produced

for each mating. The black dots represent the average number of daughters produced for each

sequential mating by the males of different *Wolbachia* infections (detailed in figure 2). The

number of daughters produced is taken as a measure of the number of sperm transferred

during each mating. The Y-axis, in grey, on the right, for each figure tallies the average number

of copulations that yielded at least one daughter. Thus, it measures the number of mating

before a male is depleted of its sperm. The left panel shows the males from *Wolbachia* infected

strains whereas the right panel shows their respective cured versions. Data for 0(PU) is repeated

at the top for comparison. The statistical significance was tested using the Mann-Whitney U test

with $p < 0.05$.

**Figure 4. *Wolbachia* infected females produce fewer offspring**

Progeny family size produced by females when hosted as virgins (top panel) and mated (middle

panel). The difference in the progeny family size of mated females is due to the difference in

the number of female progenies (bottom panel) as there is no significant difference in the

number of males produced (grey box plots). The statistical significance is tested using the

Mann-Whitney U test with $p < 0.05$.

**Figure 5. *Wolbachia* infection reduces female fecundity**

The measure of fecundity (number of eggs laid) by females of different infections, with virgin

females shown at the top and mated females at the bottom panel. For virgin females, 0(PU)

shows higher fecundity than $w$A$w$B(PU), while $w$A(PU) and $w$B(PU) strains have intermediate

fecundity which is comparable to both 0(PU) and $w$A$w$B(PU). However, for mated females,

0(PU) and $w$A(PU) show higher fecundity than $w$B(PU) and $w$A$w$B(PU). The statistical

significance is tested with the Mann-Whitney U test, $p < 0.05$.

**Figure 6. Quantitative estimation of *Wolbachia* across different developmental stages of *N.***

***vitripennis* males (top panel) and females (bottom panel).** The statistical significance between

groups is tested using the Mann-Whitney U test, $p < 0.05$.

Male lifespan

Female lifespan

**Legend:** ■ Daughters per mating (%)   ■ Number of matings before sperm depletion