Predictors of taxonomic splitting and its role in primate conservation

Maria J.A. Creighton[a], Alice Q. Luo[a], Simon M. Reader[a] & Arne Ø. Mooers[b]

[a]Department of Biology, McGill University, Montreal, Quebec, Canada

[b]Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

*Author for correspondence: maria.creighton@mail.mcgill.ca; ph: +1-514 398 7020; McGill University, Stewart Biology Building 1205 ave Docteur Penfield, Montreal, QC Canada, H3A 1B1 ORCID: https://orcid.org/0000-0003-1337-1946

WORD COUNT: 4969

DECLARATIONS OF INTEREST: None

ABSTRACT:

Species are the main unit used to measure biodiversity, but different preferred diagnostic criteria

can lead to very different delineations. For instance, named primate species have more than

doubled in number since 1982. Such increases have been attributed to a shift away from the

'biological species concept' (BSC) in favour of less inclusive species criteria. Critics of recent

changes in primate taxonomy have suggested taxonomic splitting may be biased toward certain

clades and have unfavourable consequences for conservation. Here, we explore predictors of

taxonomic splitting across primate taxa since the initial shift away from the BSC nearly 40 years

ago. We do not find evidence that diversification rate, the rate of lineage formation over

evolutionary time, is significantly linked to splitting, contrary to expectations if new species

concepts and taxonomic methods identify incipient species. We also do not find evidence that

research effort in fields where work has been suggested to motivate splitting is associated with

increases in species numbers among genera. To test the suggestion that splitting groups is likely

to increase their perceived risk of extinction, we test whether genera that have undergone more

splitting have also observed a greater increase in their proportion of threatened species since the

initial shift away from traditional taxonomic methods. We find no cohesive signal of taxonomic

splitting leading to higher threat probabilities across primate genera. Overall, this analysis sends

a positive message: threat statuses of primate species are not being overwhelmingly affected by

splitting. Regardless, we echo warnings that it is unwise for conservation to be reliant on

taxonomic stability. Species (however defined) are not independent from one another, thus,

monitoring and managing them as such may not meet the overarching goal of conserving

biodiversity.

KEYWORDS: species; phylogenetic species concept; primates; taxonomic splitting;

conservation

INTRODUCTION:

"Species" are an integral unit of biodiversity used across many sub-disciplines of biology, yet

how scientists define species has been subject to change. Notably, in the last 40 years, the

emergence of new methods for identifying diagnostic differences between populations (e.g.,

molecular phylogenetic methods) and changes in preferred species criteria have led to large

increases in species numbers across many clades (Agapow *et al.*, 2004).

Groves (2014) provides a brief overview of popular species definitions employed by taxonomists

through the late nineteenth to twentieth century. One notable phenomenon is the considerable

decrease in diagnosed species that occurred following the rapid adoption of the polytypic species

concept beginning in the 1890's. The polytypic species concept emphasizes that species should

be inclusive and that one should delineate taxa that resemble one another as subspecies (Groves,

2014). The popularization of the polytypic species concept was eventually accompanied by the

widespread adoption of the 'biological species concept' (BSC) beginning in the early 1960's

(Groves, 2014). The BSC defines species as populations/meta-populations that do not interbreed

with other populations/meta-populations under natural conditions (Mayr, 1963; Groves, 2014).

While this definition has been subject to revisions, the central premise of the BSC is that

reproductive barriers are key to diagnosing species (Groves, 2014). The BSC was widely

accepted and layered onto the pre-existing polytypic species concept, creating a period of relative

taxonomic stability for vertebrates from the 1960's to 1980's (Isaac *et al.*, 2004). However,

various criticisms of the BSC did emerge, the most notable being the practical difficulty of

diagnosing species under the BSC because of the need for information on reproductive barriers

(Donoghue, 1985; Tattersall, 2007). Such criticisms suggested the need for new approaches to

delineating species that offered higher diagnostic power.

In the last 40 years the 'phylogenetic' or 'diagnostic' species concept (PSC) (Cracraft, 1983) has

been widely popularized in vertebrate taxonomy due to its diagnosable advantages over previous

species definitions (Isaac *et al.,* 2004; Cotton *et al.,* 2016). Under the PSC, a species is diagnosed

as the smallest population or meta-population that is distinct in heritable differences from other

populations or meta-populations (Cracraft, 1983; Groves & Grubb, 2011; Groves, 2014).

According to its proponents, the PSC's emphasis on diagnosable evidence gives it an advantage

over other species concepts because it allows users to rely on a range of data types, including

newly available molecular markers, to make distinctions (see, e.g., the variety of data types used

to describe the newest species of ape, *Pongo tapanuliensis*; Nater *et al.,* 2017). Together, new

species concepts and methodological advancements have characterized a shift away from

biological species and toward species that are delineated based on distinctive, diagnosable

differences.

Although using distinctive differences to delineate species offers advantages, this approach has

also been subject to criticism, notably, for its tendency to split species into a range of less-

inclusive units (Agapow *et al.*, 2004; Zachos *et al.*, 2013; Zachos & Lovari, 2013). Many

populations previously recognized as subspecies or morphological variants have been elevated to

the full species status, resulting in a large increase in the number of listed species. For instance,

181 species of primates were listed by Honacki *et al.* (1982), one year prior to Cracraft's (1983)

proposal of the PSC. Today the IUCN (International Union for Conservation of Nature) lists over

500 distinct primate species (Estrada *et al.*, 2017). Some families (e.g., Cheirogaleidae and

Indriidae) have more than tripled in species richness over the past ~40 years (see Figure 1).

While some new species have been added as a result of new field discoveries, a majority are

populations which were previously identified at lower taxonomic levels that have now been

elevated to species status following a shift in the accepted approach to diagnosing primate

species (Tattersall, 2007). Some attribute this taxonomic increase to the popularity of the PSC

(see, e.g., Isaac *et al.*, 2004), while others credit increased exploration and the development of

new techniques for evaluating diagnosability (see, e.g., Köhler *et al.,* 2005; Harris & Froufe,

2005; Padial & De la Riva, 2006; Sangster, 2009). The trend of increasing species numbers by

raising taxonomic statuses has been referred to as 'taxonomic inflation' by some (Isaac *et al.*,

2004; Rylands & Mittermeier, 2014) and has been criticized for being non-random, and biased

toward certain clades (Isaac *et al.,* 2004; Agapow *et al.,* 2004; Zachos *et al.,* 2013; Zachos &

Lovari, 2013; Rylands & Mittermeier, 2014).

<u>Figure 1</u>: Species numbers and percentage change in species numbers for historic primate

families recorded at three time points by Honacki *et al.* (1982), Wilson & Reeder (2005) and the

IUCN species list from 2016 (data documented in Estrada *et al.*, 2017).

Disproportional splitting among taxonomic groups could have several causes. Variation in the

number of new species described across taxa could be driven by variation in the rate at which

lineages evolve, such that new species descriptions are tracking cryptic diversity or incipient

species formation among rapidly evolving taxa. In this case, new species listings might point to

situations where there is a discordance between patterns of genetic change and the evolution of

gross morphological changes used by traditional taxonomists. This could be due to ecology – if

some lineages are diversifying along ecological axes not captured in traditional taxonomic

approaches – or due to demographics – if some lineages have ecologies and/or histories that lead

to faster local genetic coalescent times and so diagnosability. Under either of these scenarios,

variation in which clades observe the greatest amount of splitting under new taxonomic

approaches would simply reflect underlying biological reality. However, it is also possible that

splitting is driven by other factors that may be prone to bias. Critics have argued that many

increases in species numbers are artificial, reflecting major shortcomings of new species criteria

and a reliance on insufficient data (Zachos *et al.*, 2013; Zachos & Lovari, 2013). Zachos *et al.*

(2013) provides evidence suggesting unwarranted splitting in select cases, advocating that

splitting has been taken to a "molecular extreme". Many different types of molecular data are

used to justify splitting (e.g., genetic data from mitochondrial DNA barcoding) and, by this logic,

groups may continue to be split as an increasing amount of molecular data become available for

them. If true, this would lead to the continual identification of increasingly exclusive diagnostic

features such that we are likely to find new species the more we look for them.

It is also possible that conservation interest in particular groups could motivate splitting. Limited

funding for conservation research is increasingly focused on 'biodiversity hotspots' and it has

been suggested that researchers could have a vested interest in declaring taxa in these regions as

endemic species (Karl & Bowen, 1999; Isaac *et al.*, 2004). There is evidence from some

charismatic groups that taxa receive more conservation attention and funding when comprised of

multiple, small, and taxonomically distinct populations (e.g., African apes; Stanford, 2001;

Oates, 2006; Gippoliti & Amori, 2007). Species lists are often used to determine which groups

should receive conservation attention (Mace, 2004) and so changing the way we define species

may also change which groups receive action.

Along with being potentially biased toward certain taxonomic groups and related to the point

made above, taxonomic splitting has also been suggested to result in individually more imperiled

populations which could distort conservation agendas (Agapow *et al.*, 2004; Isaac *et al.*, 2004;

Morrison *et al.*, 2009; Zachos, 2015; Robuchon *et al.*, 2019). One criterion used by the IUCN to

classify species as imperiled is population size: species may be designated "Vulnerable" if there

are fewer than 1000 mature individuals found in the wild and "Endangered" if there are fewer

than 250 (Agapow *et al.*, 2004; Frankham *et al.*, 2012). Thus, splitting one species into several

new species may result in one or more receiving a (more) imperiled status (Agapow *et al.*, 2004;

Isaac *et al.*, 2007). This could lead to seemingly rare but poorly defined species being prioritized

over well-defined and perhaps biologically more distinctive species (Pillon & Chase, 2007).

Recent evidence suggests species splitting is not a driver of threat status for birds (Simkins *et al.*,

2020), however, it is unknown whether these findings are generalizable across other taxonomic

groups (Garnett & Thomson, 2020).

Here, we set out to better understand the causes and consequences of taxonomic splitting in

primates. We test (i) predictors of taxonomic splitting, that is, whether recent taxonomic

increases are associated with the amount of research being done in fields suggested to motivate

splitting or with a lineage's underlying diversification rate, and (ii) impacts of taxonomic

splitting, that is, whether rates of splitting dictate which groups are most imperiled. Because

some newly described primate species have previously been described as subspecies or

subpopulations of more than one different biological species prior to being assigned to the full

species rank (which would make quantifying rates of splitting at the species-level difficult), we

ask these questions at the genus level.

To explore our first question regarding predictors of taxonomic splitting, we consider whether

biological factors or measures of human-induced bias explain increases in species numbers. To

151 test potential human-induced biases in splitting, we consider broad estimates of research effort

152 for each taxon and predict that more research done on a given taxon may be associated with

153 more splitting. We examine research effort in two fields: molecular genetics (since molecular

154 work could cause species to be split continuously as finer molecular distinctions are made) and

155 conservation (since splitting has been suggested to be motivated by conservation interests). To

156 explore possible biological explanations for trends in taxonomic splitting across clades, we test

157 whether recent taxonomic increases are explained by diversification rate. Clades with high recent

158 diversification rates are expected to contain more incipient or cryptic species than lineages

159 diversifying at a lower rate since they will contain more closely related lineages that resemble

160 one another. Therefore, if new approaches help to identify incipient or cryptic species,

161 diversification rate should be positively correlated with splitting.

163 To explore our second question linking splitting and risk, we test whether increases in species

164 numbers are associated with a change in the number of threatened species listed in genera

165 through time. Ideally we would look at changes in a weighted measure of threat score that

166 differentiates between threat categories of varying severity (e.g., using the Red List Index (RLI);

167 Butchart *et al.*, 2007; Bubb *et al.*, 2009); however, criteria for inclusion in Red List threat

168 categories have changed considerably over time, meaning weights assigned to categories for RLI

169 calculations do not match up with categories used in the past (see, e.g., statuses in IUCN

170 Conservation Monitoring Centre, 1986). Because of these changes in Red List criteria, we

171 instead ask whether clades with species that have been split more frequently have observed a

172 greater increase in their proportion of threatened species (defined below) through time in

173 comparison to those which have been split less frequently. We predict that if splitting is driving

an increase in threatened species, we should observe a positive association between taxonomic

increases caused by splitting and change in the proportion of threatened primate species.

MATERIALS AND METHODS:

Data

To measure the number of primate species that were described before the introduction of the

PSC and new molecular techniques, we used the last pre-PSC taxonomy, that of Honacki *et al.*

(1982). This taxonomy contains 181 species and is considered a reliable estimate of the number

of species thought to exist during the popularity of the BSC (see Rylands & Mittermeier, 2014).

We then recorded if each species in this taxonomy was historically considered to be threatened

by referencing the most complete IUCN Red List published around the same time (IUCN

Conservation Monitoring Centre, 1986). Honacki *et al.* (1982) was contrasted with the IUCN

taxonomy from 2016 and attendant data documented in Estrada *et al.* (2017), which lists 503

species. For each of these 503 species we noted their taxonomic placement (genus and family),

whether or not they were considered threatened (VU=Vulnerable, EN=Endangered, or

CR=Critically Endangered), and their biogeographic region. We note that while IUCN

assessments have some shortcomings (see, e.g., Rueda-Cediel *et al.,* 2018), the IUCN provides

the largest global data on threat status and is influential in determining how most species are

managed. For each species described by the IUCN in 2016 that was not listed in Honacki *et al.*

(1982), we scored whether the species was a "*de novo*" species description (Burgin *et al.*, 2018):

there were 40 such cases where a new species had not been previously formally identified as a

subspecies or subpopulation of another species prior to splitting. These cases represented new

species descriptions where it was unclear whether a new species was a result of taxonomic

splitting or the discovery of an entirely new population.

We compiled all the species listed by the IUCN in 2016 into 12 families and 50 genera found in

Honacki *et al.* (1982). Family name "Callimiconidae" in Honacki *et al.* (1982) was not used as

this taxon has since been recognized as a genus of the larger family "Callitrichidae" (Wilson &

Reeder, 2005). We removed *Rungwecebus kipunji* from the IUCN species list from 2016 as it

represents a newly discovered genus that does not collapse into any of the genera provided by

Honacki *et al.* (1982).

Research effort in the fields of molecular genetics and conservation was estimated for each genus

through a literature review of publications in the Web of Science Core Collection published

between 1983 (when a general trend toward taxonomic splitting first began) and 2016. All search

terms for these literature reviews and further details on related methodology are documented in

the supplementary materials (Tables S1 and S2). In this study we used the genera listed in

Honacki *et al.* (1982) (n=50), many of which have since been further separated into multiple

genera. Thus, when appropriate we included new genus names in addition to those listed by

Honacki *et al.* (1982) in the literature searches (see Table S3). In total, our final sample was 688

publications on molecular genetics and 2222 on conservation.

Diversification rate was estimated with the method-of-moments approach described in Magallon

& Sanderson (2001) (i.e., ln(taxa richness)/stem age). Diversification rate estimates generated

using this method often rely on species numbers as their estimate of taxa richness, meaning that

diversification rate estimates are inherently biased by the splitting phenomenon we are studying

(i.e., frequently split genera will receive disproportionally high diversification rates). Therefore,

richness scores for our diversification rate calculations were determined as the counts of the

well-resolved "lineages" described in Creighton *et al.* (2021). These lineages were determined by

creating a time cut-off in the 10kTrees consensus primate phylogeny (Arnold *et al.*, 2010) in an

attempt to eliminate very young newly described species and obtain a consistent (unbiased)

estimate of diversity across clades. These lineages were assigned to each of the 50 genera

described in Honacki *et al.* (1982). Diversification rate was then estimated by taking the natural

log lineage richness for each genus and dividing by the stem age of that genus (Magallon &

Sanderson, 2001). Stem ages for each genus were extracted from the 10kTrees consensus

phylogeny (version 3) (Arnold *et al.*, 2010) trimmed to contain a single branch representing each

genus. During this process, there were several instances where genera described in Honacki *et al.*

(1982) were non-monophyletic within the more recent primate phylogeny we used (Arnold *et al.*,

2010), making it unclear how to assign a divergence date for these clades. We therefore removed

eight genera from the analyses where diversification rate was a variable of interest: *Presbytis*,

Lemur, *Galago*, *Cebuella*, *Cercocebus*, *Cercopithecus*, *Papio*, and *Pygathrix*.

Analysis

To test our questions about the predictors of species-splitting and its consequences for

conservation, we fit a series of linear effects and mixed effects models. We note that the response

variables used in these analyses (i.e., measures of taxonomic increase and extinction risk) are

likely to be phylogenetically clustered, and phylogenetic models could be used to account for

this influence of phylogeny; however, many genera listed by Honacki *et al.* (1982) are non-

monophyletic, making it unclear how to designate them a single branch in modern phylogenies

(see discussion above on diversification rate). Importantly, after accounting for regional

differences, family contributed little to no variance in any of our models, indicating that

phylogenetic relationships at that level were not confounding our results. We provide further

discussion on model choice in the supplementary materials.

Data were analysed using R version 4.0.5 (R Core Team, 2021).

Predictors of Taxonomic Splitting

To determine if measures of potential human bias (i.e., research effort) or diversification rate

explained discrepancies in splitting across taxa, we tested to see if these variables were

significantly associated with the number of species added to primate genera since 1982 while

controlling for the original number of described species (i.e., biological species) and region. We

fit three generalized linear mixed effects models with Poisson distributions using the lme4

package in R (Bates *et al.*, 2014), and obtained p-values using the lmerTest package (Kuznetsova

et al., 2017). In these models the response variable was the number of species assigned to a given

genus by the IUCN in 2016 that had not been previously described by Honacki *et al.* (1982).

Each model had either conservation research effort, molecular genetics research effort or

diversification rate included as a fixed effect, as well as biogeographical region and the number

of species in the genus per Honacki *et al.* (1982) to control for their effects on splitting. The

natural logarithm (ln) of the number of species listed for each genus in Honacki *et al.* (1982) was

included as both a linear and a quadratic term following inspection of raw plots and plots against

scaled residuals from the simulation output. A square-root transformation was used on molecular

genetics research effort to decrease the impact of outliers on model fit. To assist with model

stability and convergence, we scaled all continuous variables in the model to have a mean of zero

and standard deviation of one (Becker *et al.*, 1988). Mainland Africa and Asia (hereafter

mainland Africa + Asia) were grouped together and served as the baseline region in our models

based on previous studies that have shown that the taxonomy of primates from these regions has

been relatively stable compared to Madagascar and the Neotropics (e.g., Isaac & Purvis, 2004;

Isaac *et al.*, 2004; Tattersall, 2007). We also chose to group Asia and Africa together because

one genus (*Macaca*) is found in both regions. Genus ID was included as a random effect in these

models to correct for overdispersion. Family (nested within region) was originally included as a

random effect but contributed very little to model fit and created issues with convergence due to

overfitting, and thus was dropped from the final models. We tested potential interaction terms

with all variables and region to test for regional effects, but none were significant and so these

terms were also dropped from the final models. Before running models with other predictors

included, we also ran a model including only the linear and quadratic terms for the number of

species listed for each genus in Honacki *et al.* (1982) as predictors to assess their association

with taxonomic increase in the simplest model. We ran all models a second time after removing

the 40 *de novo* species from our response (Tables S4, S5 and S6). We checked model

assumptions and fit by plotting residuals versus the fitted values and versus each covariate in the

model. Residual plots and analyses done with the Diagnostics for Hierarchical Regression

Models (DHARMa) R package (Hartig, 2017) indicated acceptable model fits.

Taxonomic Splitting and Threat Score

To test whether taxonomic splitting over time is associated with a change in the proportion of

threatened species within genera, we conducted a two-step (hierarchical) analysis on species'

threat probability between 1982 to 2016. We first fit a generalized linear mixed effects model

with a binomial distribution for the number of threatened and non-threatened species within a

genus using the lme4 package in R (Bates *et al.*, 2014). In this first model, the response

estimated the proportional counts of species at risk within genera (equivalent to the per-species

probability of threat) in the periods of 1982 and 2016. Predictors for this model included fixed

effects for the time period (the baseline of 1982 and the change to 2016), the region that

encompasses each genus' natural distribution (Madagascar, Neotropics, mainland Africa + Asia),

and an interaction between region and time period to account for geographic differences in threat

probabilities through time. Genus identity was included as a random intercept, to account for

repeated measures in 1982 and 2016, and as a random slope with time period to account for

differences in changes to threat probabilities among genera. As discussed above, we included the

taxonomic rank of family (nested within region) as a random effect but found that the variance in

threat probabilities among families was minimal and that including this term also created

convergence issues; given this, we subsequently omitted family from our models. Residual plots

and analyses with the DHARMa R package (Hartig, 2017) indicated acceptable model fits for the

final model. From this first model describing genus level changes in threat probabilities, we then

used the REextract function implemented in the merTools package in R (Knowles & Frederick,

2020) to extract the genus level random slopes for time period and their associated standard

errors, giving us an estimate of the varying effect of change in threat probability among genera

(while conditioning on regional trends) between 1982 and 2016. We then fit a linear model

where varying effect of change in threat probability from 1982 to 2016 was the response, and the

proportional change in species within each genus (i.e., the number of new species in the IUCN

species list from 2016 / original number of species in Honacki *et al.* (1982) – our measure of

taxonomic increase) – was the predictor. We weighted each estimate of genus level change in

threat probability by its standard error (w = 1/SE) to propagate the error of model estimated

random effects. We ran this model a second time after removing *de novo* species descriptions

from our response (Table S7). Raw data used for these tests are visualized in Figure 2.

Plots of the raw data indicated that patterns in taxonomic increases, the proportion of species at

risk in primate genera today, and the changes in the proportion of threatened species in primate

genera between 1982 and 2016 appeared to show regional differences (Figures S1 and S2). We

thus ran a subsequent set of models to test for regional variation in the effect of splitting. In this

analysis, we removed region from the first generalized linear mixed effects model and

subsequently included region interacting with taxonomic increase as a predictor of the varying

effect of change in genus threat probability in the second-order model. However, interaction

effects were not significant in this model and thus we only considered the results of the first set

of models reported above.

RESULTS:

Predictors of Taxonomic Splitting

None of our measures – conservation research effort ($\beta = -0.220$; $p = 0.180$; Table S4), molecular

genetics research effort ($\beta = -0.188$; $p = 0.212$; Table S5) or diversification rate ($\beta = 0.132$; $p =$

0.477; Table S6) – were significantly associated with increases in species numbers across

primate genera. Removing *de novo* species did not impact this pattern of results (Tables S4, S5

and S6). Notably, in addition to sharing a significant linear relationship with taxonomic increase

as expected (Tables S4, S5 and S6), the quadratic term added for the original number of species

in 1982 was significant in the model with diversification rate, indicating a downwardly concave

association with taxonomic increase regardless of whether de novo species were included (Table

S6); this term had a p-value less than 0.1 in models that included conservation or molecular

genetics research effort as predictors (Tables S4 and S5), indicating that the positive increase in

splitting by initial genus size decelerates as genera become very large, even after accounting for

other variables. We note that this quadratic term was significant without other predictors (i.e.,

research effort or diversification rate) present in the model, both with *de novo* species included in

the response (β= -2.268; p= 0.011) and without *de novo* species included in the response (β= -

2.086; p= 0.013). In all three models without *de novo* species, taxonomic increases were

significantly higher in the Neotropics compared to Africa and Asia (Tables S4 to S6).

Taxonomic Splitting and Threat Score

Visual inspection of the raw data on taxonomic increases and proportion of species at risk for all

genera revealed no clear indication that highly split genera have observed a disproportional

increase in perceived extinction risk (Figures 2 and S3). Results from our linear model confirmed

that taxonomic increases did not predict a change in threat score through time (β= 0.001; p=

0.975; Table S7; Figure 3). Removing *de novo* species did not impact this pattern of results

(Table S7; Figure S4).

<u>Figure 2</u>: Scatterplots with trendlines showing the change in the proportion of species identified

as being threatened in primate genera in 1982 and 2016 painted by the square root of taxonomic

increase (including *de novo* species). Total number of species in each genus is indicated by point

size.

<u>Figure 3:</u> Taxonomic increase including *de novo* species descriptions versus varying effect of

change in threat probability (i.e., our measure of genus level change in threat probability between

1982 and 2016; see Methods for an explanation of how this was estimated) (β= 0.001; p= 0.975;

Table S7). Vertical bars indicate standard errors on varying effect of change in threat probability

values.

DISCUSSION:

Our results support neither a strong biological nor a strong interest-driven mechanism for

splitting across primate genera. If the naming of new species under new taxonomic approaches

captures incipient speciation, we predicted that diversification rate should predict taxonomic

increases. However, diversification rate was not a significant predictor of taxonomic increases in

our models. We also tested the prediction that splitting is motivated by conservation interest

(Karl & Bowen, 1999; Isaac *et al.*, 2004) or by increasing molecular research within certain taxa

(Zachos *et al.*, 2013; Zachos & Lovari, 2013). However, we found no evidence that research

effort in either of these areas was associated with the amount of taxonomic splitting observed

across genera. We did find that initial genus size predicts increased splitting, but this effect

decelerates as genera approach the largest sizes. This could indicate that splitting is being driven

by variables not captured by our models. For example, genera that already contained many

species in 1982 may have been more likely to have closely related populations described as

separate species under the BSC (e.g., if we knew more about their hybrid statuses compared to

other clades). In this case, we may have already discovered much diversity in these large clades

prior to applications of the PSC and new molecular techniques. Future studies could aim to

elucidate the origins of this decelerating association, however, our results indicate it is not

explained by the rate at which lineages diverge (which should capture the presence of cryptic

species) or research effort in the two fields studied here.

While inspection of the raw data suggested that genus level measures of splitting, threat

probability, and change in threat probability all varied by region, we found that there was no

cohesive signal of taxonomic splitting leading to higher threat probability across these genera

and regions. Therefore, while there are idiosyncratic regional differences in both splitting and

threat probability there was no evidence for causal links between the two. We do note that while

the act of splitting itself does not seem to be having an overwhelming effect on threat status, it is

still possible splitting may have uncaptured negative consequences downstream in conservation

management (e.g., loss of genetic variation in captive breeding programs; Zachos, 2013).

However, our analysis sends a positive message that splitting is not currently a significant

determinant of the relative conservation priority of primate clades, consistent with recent

findings in birds (Simkins *et al.*, 2020).

Importantly, we note that our analyses have some limitations. First, quantifying research effort of any kind is difficult due to the abundance of work published in different media. As such, our estimates of research effort in conservation and molecular genetics may overlook some types of research. More work could be done to determine if additional estimates of research bias can explain increases in species numbers. For instance, cumulative funding estimates from various sources (e.g., the IUCN, non-governmental organizations and regional governments) per taxon could provide an additional or alternative measure of conservation interest. Second, it was necessary to remove eight non-monophyletic genera (some of which have undergone substantial splitting) from analyses that included diversification rate, leading to a considerable reduction in statistical power for those analyses. Third, due to changes in Red List criteria, species status, and lack of information about which species had been assessed in the 1980s, we were not able to consider differences in threat status severity (e.g., Vulnerable versus Endangered) when considering associations between taxonomic splitting and changes in threat score through time. Future studies could look at the association between splitting and changes in weighted measures of threat score (e.g., RLI) over time using a more recent starting point (i.e., after 1982) where Red List criteria become stable and there is available information on which non-threatened species have been evaluated.

Overall, we do not find support for biological processes or research bias driving taxonomic splitting across primate genera. We also find no cohesive signal of splitting leading to higher threat probabilities. Generally speaking, relying on species as the central unit of conservation and primary object of biological study behooves taxonomy to remain stable, while changing ideas

about the concept of species makes taxonomy inherently unstable (Mayr, 1996). We suggest that

areas of research requiring consistent estimates of diversity (e.g., conservation, macroecology, or

evolutionary biology) may benefit from (i) weighing evolutionary distinctiveness when

determining how species are listed/treated if attempting to capture true biological diversity (see,

e.g., Redding & Mooers, 2010; Redding *et al.*, 2015); or, for applied conservation specifically

(ii) shifting more resources toward regional management efforts that are less likely to be

influenced by changing species designations. It is well-known that closely related species are

more similar to one another than they are to more distantly related taxa. Thus, treating all species

independently and of equal weights in conservation listing may not lead to desired outcomes

(Redding & Mooers, 2010). As of 2016, approximately 60 percent of all primate species were

threatened with extinction according to the IUCN Red List of Threatened Species (Estrada *et al.*,

2017), making it imperative that conservation efforts are spent wisely to ensure optimal

conservation of primate biodiversity writ large.

ACKNOWLEDGEMENTS: This work was supported by the Natural Sciences and Engineering

Research Council of Canada (NSERC) Discovery Grants Program (SMR and AOM, grants

#2017-04720 and #2019-04950) and the Canada Foundation for Innovation (CFI) (SMR, grant

#29433). MJAC was supported by awards from the Biodiversity, Ecosystem Services, and

Sustainability (BESS) program and McGill University. We thank the Reader and Guigueno labs

at McGill, the Crawford Lab for Evolutionary Studies at SFU, Brian Leung, Hans Larsson,

William Wcislo, and Isabella Capellini for feedback on portions of the manuscript and study

design. Special thanks to Aiman Hadif for his assistance in collecting and organizing data and to

Paul Sims and Dan Greenberg for statistical advice. We also thank Sandro Lovari, Frank Zachos,

and one anonymous reviewer for their useful comments on earlier drafts.

DECLARATIONS OF INTEREST: None.

Data and code accompanying the manuscript are available on Zenodo:
https://doi.org/10.5281/zenodo.6614874

REFERENCES:

Agapow, P.M., Bininda-Emonds, O.R., Crandall, K.A., Gittleman, J.L., Mace, G.M., Marshall,
J.C. & Purvis, A. (2004). The impact of species concept on biodiversity studies. *Q. Rev.*
Biol. **79**, 161–179.

Arnold, C., Matthews, L. J. & Nunn, C.L. (2010). The 10kTrees website: a new online resource
for primate phylogeny. *Evol. Anthropol.* **19**, 114–118.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2014). *lme4: Linear mixed-effects models*
using Eigen and S4. R package version 1.1.21. https://cran.r-
project.org/web/packages/lme4.

Becker, R.A., Chambers, J.M. & Wilks, A.R. (1988). *The new S language: A programming*
environment for data analysis and graphics. Wadsworth & Brooks/Cole, Pacific Grove.

Bubb, P.J., Butchart, S.H.M., Collen, B., Dublin, H., Kapos, V., Pollock, C., Stuart, S.N. & Vié,

J.–C. (2009). *IUCN Red List Index – guidance for national and regional use*. IUCN,

Gland.

Burgin, C.J., Colella, J.P., Kahn, P.L. & Upham, N.S. (2018). How many species of mammals

are there? *J. Mammal.* **99**, 1–14.

Butchart, S.H., Akçakaya, H.R., Chanson, J., Baillie, J.E., Collen, B., Quader, S., Turner, W.R.,

Amin, R., Stuart, S.N. & Hilton-Taylor, C. (2007). Improvements to the Red List Index.

PLoS One **2**, e140.

Cotton, A., Clark, F., Boubli, J.P. & Schwitzer, C. (2016). IUCN Red List of threatened primate

species. In S.A. Wich & A.J. Marshall (Eds.), *An introduction to primate conservation* (p.

31–38). Oxford University Press, Oxford.

Cracraft, J. (1983). Species concepts and speciation analysis. In R.F. Johnson (Ed.), *Current*

ornithology (p. 159–187). Plenum Press, New York.

Creighton, M.J.A., Greenberg, D.A., Reader, S.M. & Mooers, A.Ø. (2021). The role of

behavioural flexibility in primate diversification. *Anim. Behav.* **180**, 269–290

Donoghue, M.J. (1985). A critique of the biological species concept and recommendations for a

phylogenetic alternative. *Bryologist* **88**, 172–181.

Estrada, A., Garber, P.A., Rylands, A.B., Roos, C., Fernandez-Duque, E., Di Fiore, A., Nekaris,

K.A., Nijman, V., Heymann, E.W., Lambert, J.E., Rovero, F., Barelli, C., Setchell, J.M.,

Gillespie, T.R., Mittermeier, R.A., Arregoitia, L.V., de Guinea, M., Gouveia, S.,

Dobrovolski, R., Shanee, S., Shanee, N., Boyle, S.A., Fuentes, A., MacKinnon, K.C.,

Amato, K.R., Meyer, A.L.S., Wich, S., Sussman, R.W., Pan, R., Kone, I. & Li, B. (2017).
Impending extinction crisis of the world's primates: Why primates matter. *Sci. Adv.* **3**,
e1600946.

Frankham, R., Ballou, J.D., Dudash, M.R., Eldridge, M.D.B., Fenster, C.B., Lacy, R.C.,
Mendelson III, J.R., Porton, I.J., Ralls, K. & Ryder, O.A. (2012). Implications of
different species concepts for conserving biodiversity. *Biol. Conserv.* **153**, 25–31.

Garnett, S.T. & Thomson, S.A. (2020). Are the implications for conservation of a major
taxonomic revision of the world's birds' simply serendipity? *Anim. Conserv.* **23**, 355-
356.

Gippoliti, S. & Amori, G. (2007). The problem of subspecies and biased taxonomy in
conservation lists: the case of mammals. *Folia Zool.* **56**, 113–117.

Groves, C.P. (2014). Primate taxonomy: inflation or real? *Annu. Rev. Anthropol.* **43**, 27–36.

Groves, C. & Grubb, P. (2011). *Ungulate taxonomy*. JHU Press, Baltimore.

Hartig, F. (2017). *DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression*
models. R package version 0.3.0. https://cran.r-project.org/web/packages/DHARMa.

Harris, D.J., & Froufe, E. (2005). Taxonomic inflation: species concept or historical geopolitical
bias? *Trends Ecol. Evol.* **20**, 6–7.

Honacki, J.H., Kinman, K.E. & Koeppl, J.W. (1982). *Mammal species of the world; a taxonomic*
and geographic reference. Allen Press, Inc. and Association of Systematic Collections,
Lawrence.

Isaac, N.J., Mallet, J. & Mace, G.M. (2004). Taxonomic inflation: its influence on macroecology

and conservation. *Trends Ecol. Evol.* **19**, 464–469.

Isaac, N.J. & Purvis, A. (2004). The 'species problem' and testing macroevolutionary

hypotheses. *Divers. Distrib.* **10**, 275–281.

Isaac, N.J., Turvey, S.T., Collen, B., Waterman, C. & Baillie, J.E. (2007). Mammals on the

EDGE: conservation priorities based on threat and phylogeny. *PLoS One* **3**, e296.

IUCN Conservation Monitoring Centre. (1986). 1986 *IUCN Red List of threatened animals*.

IUCN, Gland and Cambridge.

Karl, S.A. & Bowen, B.W. (1999). Evolutionary significant units versus geopolitical taxonomy:

molecular systematics of an endangered sea turtle (genus *Chelonia*). *Conserv. Biol.* **13**,

990–999.

Köhler, J., Vieites, D.R., Bonett, R.M., García, F.H., Glaw, F., Steinke, D., & Vences, M.

(2005). New amphibians and global conservation: a boost in species discoveries in a

highly endangered vertebrate group. *BioScience* **55**, 693–696.

Knowles, J.E. & Frederick, C. (2020). *merTools: Tools for Analyzing Mixed Effect Regression*

Models. R package version 0.5.2. https://CRAN.R-project.org/package=merTools.

Kuznetsova, A., Brockhoff, B. & Christensen, H.B. (2017). lmerTest Package: Tests in Linear

Mixed Effects Models. *J. Stat. Softw.* **82**, 1–26.

Mace, G.M. (2004). The role of taxonomy in species conservation. *Philos. Trans. R. Soc. B:*

Biol. Sci. **359**, 711–719.

Magallon, S. & Sanderson, M.J. (2001). Absolute diversification rates in angiosperm clades.
Evolution **55**, 1762–1780.

Mayr, E. (1963). *Animal species and evolution*. Belknap Press, Cambridge.

Mayr, E. (1996). What is a species, and what is not? *Philos. Sci.* **63**, 262–277.

Morrison, W.R., Lohr, J.L., Duchen, P., Wilches, R., Trujillo, D., Mair, M. & Renner, S.S.
(2009). The impact of taxonomic change on conservation: Does it kill, can it save, or is it
just irrelevant? *Biol. Conserv.* **142**, 3201–3206.

Nater, A., Mattle-Greminger, M. P., Nurcahyo, A., Nowak, M. G., De Manuel, M., Desai, T., ...
& Krützen, M. (2017). Morphometric, behavioral, and genomic evidence for a new
orangutan species. *Curr. Biol.* **27**, 3487–3498.

Oates, J. F. (2006). Is the chimpanzee, *Pan troglodytes*, an endangered species? It depends on
what "endangered" means. *Primates* **47**, 102–112.

Padial, J.M., & De la Riva, I. (2006). Taxonomic inflation and the stability of species lists: the
perils of ostrich's behavior. *Syst. Biol.* **55**, 859–867.

Pillon, Y. & Chase, M.W. (2007). Taxonomic exaggeration and its effects on orchid
conservation. *Conserv. Biol.* **21**, 263–265.

R Core Team. (2021). *R: A Language and environment for statistical computing*. R Foundation
for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Redding, D.W. & Mooers, A.Ø. (2010). Can systematists help decide the relative worth of bits of
biodiversity? *Systematist* **32**, 4–8.

Redding, D.W., Mooers, A.Ø., Şekercioğlu, Ç.H., & Collen, B. (2015). Global evolutionary
isolation measures can capture key local conservation species in Nearctic and Neotropical
bird communities. *Philos. Trans. R. Soc. B: Biol. Sci.* **370**, 20140013.

Robuchon, M., Faith, D.P., Julliard, R., Leroy, B., Pellens, R., Robert, A., Thévenin, C., Véron,
S. & Pavoine, S. (2019). Species splitting increases estimates of evolutionary history at
risk. *Biol. Conserv.* **235**, 27–35.

Rueda-Cediel, P., Anderson, K.E., Regan, T.J., & Regan, H.M. (2018). Effects of uncertainty
and variability on population declines and IUCN Red List classifications. *Conserv. Biol.*
32, 916-925.

Rylands, A.B. & Mittermeier, R.A. (2014). Primate taxonomy: species and conservation. *Evol.*
Anthropol. **23**, 8–10.

Sangster, G. (2009). Increasing numbers of bird species result from taxonomic progress, not
taxonomic inflation. *Proc. Royal Soc. B* **276**, 3185–3191.

Simkins, A.T., Buchanan, G.M., Davies, R.G. & Donald, P.F. (2020). The implications for
conservation of a major taxonomic revision of the world's birds. *Anim. Conserv.* **23**, 345–
352.

Stanford, C.B. (2001). The subspecies concept in primatology: the case of mountain gorillas.
Primates **42**, 309–318.

Tattersall, I. (2007). Madagascar's lemurs: cryptic diversity or taxonomic inflation? *Evol.*
Anthropol. **16**, 12–23.

Wilson, D.E. & Reeder, D.M. (2005). *Mammal species of the world: a taxonomic and*
geographic reference (vol. 1). JHU Press, Baltimore.

Zachos, F.E. (2013). Species splitting puts conservation at risk. *Nature* **494**, 35–35.

Zachos, F.E. (2015). Taxonomic inflation, the phylogenetic species concept and lineages in the
tree of life–a cautionary comment on species splitting. *J. Zoolog. Syst. Evol.* **53**, 180–184.

Zachos, F.E., Apollonio, M., Bärmann, E.V., Festa-Bianchet, M., Göhlich, U., Habel, J.C.,
Haring, E., Kruckenhauser, L., Lovari, S., McDevitt, A.D., Pertoldi, C., Rössner, G.E.,
Sánchez-Villagra, M.R., Scandura, M. & Suchentrunk, F. (2013). Species inflation and
taxonomic artefacts – A critical comment on recent trends in mammalian
classification. *Mamm. Biol.* **78**, 1–6.

Zachos, F.E. & Lovari, S. (2013). Taxonomic inflation and the poverty of the Phylogenetic
Species Concept – a reply to Gippoliti and Groves. *Hystrix* **24**, 142–144.