Predictors of Taxonomic Inflation and its Role in Primate Conservation

Maria J.A. Creighton[a], Alice Q. Luo[a], Simon M. Reader[a] & Arne Ø. Mooers[b]

[a]Department of Biology, McGill University, Montreal, Quebec, Canada

[b]Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

*Author for correspondence: maria.creighton@mail.mcgill.ca; ph: +1-514 398 7020; McGill University, Stewart Biology Building 1205 ave Docteur Penfield, Montreal, QC Canada, H3A 1B1 ORCID: https://orcid.org/0000-0003-1337-1946

WORD COUNT: 4941

DECLARATIONS OF INTEREST: None

ABSTRACT:

Species are the main unit used to measure biodiversity, but different preferred diagnostic criteria can lead to very different delineations. For instance, named primate species have more than doubled since 1982. Such increases have been termed "taxonomic inflation" and have been attributed to the widespread adoption of the 'phylogenetic species concept' (PSC) in preference to the previously popular 'biological species concept' (BSC). Criticisms of the PSC have suggested taxonomic inflation may be biased toward particular taxa and have unfavourable consequences for conservation. Here, we explore predictors of taxonomic inflation across primate taxa since the initial application of the PSC nearly 40 years ago. We do not find evidence that diversification rate, the rate of lineage formation over evolutionary time, is linked to inflation, contrary to expectations if the PSC identifies incipient species. We also do not find evidence of research effort in fields where work has been suggested to motivate splitting being associated with increases in species numbers among genera. To test the suggestion that splitting groups is likely to increase their perceived risk of extinction, we test whether genera that have undergone more splitting have also observed a greater increase in their proportion of threatened species since the introduction of the PSC. We find no cohesive signal of inflation leading to higher threat probabilities across primate genera. Overall, this analysis sends a positive message that threat statuses of primate species are not being overwhelmingly affected by splitting in line with what has recently been reported for birds. Regardless, we echo warnings that it is unwise for conservation to be reliant on taxonomic stability. Species (however defined) are not independent from one another, thus, monitoring and managing them as such may not meet the overarching goal of conserving biodiversity.

KEYWORDS: species; phylogenetic species concept; primates; taxonomic inflation;

conservation

INTRODUCTION:

"Species" are an integral unit of biodiversity used across many sub-disciplines of biology, yet how

scientists define species has been subject to change. New species definitions have led to dramatic

shifts in the number of species listed in many taxonomic groups throughout time. Notably, in the

last 40 years, the widespread popularization of the 'phylogenetic' or 'diagnostic' species concept

(PSC) (Cracraft, 1983; Isaac, et al., 2004; Cotton, et al., 2016) has led to large increases in species

numbers across many clades (Agapow, et al., 2004).

Groves (2014) provides a brief overview of popular species definitions employed by taxonomists

through the late nineteenth to twentieth century. One notable phenomenon is the considerable

decrease in diagnosed species that occurred following the rapid adoption of the polytypic species

concept beginning in the 1890's. The polytypic species concept emphasizes that species should be

inclusive and that one should delineate taxa that resemble one another as subspecies (Groves,

2014). The popularization of the polytypic species concept was eventually accompanied by the

widespread adoption of the 'biological species concept' (BSC) beginning in the early 1960's

(Groves, 2014). The BSC defines species as populations/meta-populations that do not interbreed

with other populations/meta-populations under natural conditions (Mayr, 1963; Groves, 2014).

While this definition has been subject to revisions, the central premise of the BSC is that

reproductive barriers are key to diagnosing species (Groves, 2014). The BSC was widely accepted

and layered onto the pre-existing polytypic species concept, creating a period of relative taxonomic

stability for vertebrates from the 1960's to 1980's (Isaac, et al., 2004). However, various criticisms

of the BSC did emerge, the most notable being the practical difficulty of diagnosing species under

the BSC because of the need for information on reproductive barriers. Closely related lineages

sharing parapatric (i.e., separate but contiguous) distributions are especially difficult to diagnose

under the BSC often due to limited records of potential mating interactions or of hybrid forms

(Tattersall, 2007). In these cases, genetic evidence is required to show that gene pools are linked,

e.g., actively reintegrating after being geographically separated (Tattersall, 2007). Collecting such

evidence in every case of parapatry would be a daunting task, and these critiques suggested the

need for a species concept with higher diagnostic power.

In the last 40 years the phylogenetic species concept (PSC) (Cracraft, 1983; Isaac, et al., 2004) has

been widely popularized in vertebrate taxonomy due to its diagnosable advantages over the BSC.

Under the PSC, a species is diagnosed as the smallest population or meta-population that is distinct

in heritable differences from other populations or meta-populations (Cracraft, 1983; Groves &

Grubb, 2011; Groves, 2014). According to its proponents, the PSC's emphasis on diagnosable

evidence gives it an advantage over other species concepts because it allows users to establish a

scientific proposition that is testable. As a result, shortly after being proposed influential

taxonomists quickly advocated for use of the PSC over the BSC (see, e.g., Donoghue, 1985) and

thus, the PSC began being used almost immediately after its inception.

Although the PSC offers diagnostic advantages over the BSC, it has also been subject to criticism,

notably, for its tendency to split species into a range of less-inclusive units compared to previous

species concepts (Agapow, et al., 2004; Zachos, et al., 2013; Zachos & Lovari, 2013). Under the

PSC many populations previously recognized as subspecies or morphological variants have been

elevated to the full species status, resulting in a large increase in the number of listed species. For

instance, 181 species of primates were listed by Honacki et al. (1982), one year prior to Cracraft's

proposal of the PSC (Cracraft, 1983). Today the IUCN (International Union for Conservation of

Nature) lists over 500 distinct primate species (Estrada, et al., 2017). Some families (e.g.,

Cheirogaleidae and Indriidae) have more than tripled in species numbers since the introduction of

the PSC (see Figure 1). While some new species have been added as a result of new discoveries,

a majority are populations previously identified at a lower taxonomic level that have since been

elevated to species status following the application of the PSC (Tattersall, 2007). This trend of the

PSC increasing species numbers by raising taxonomic statuses has been referred to as 'taxonomic

inflation' (Isaac, et al., 2004; Rylands & Mittermeier, 2014). Taxonomic inflation has been

criticized for being non-random, and biased toward certain taxonomic groups (Isaac, et al., 2004).

Figure 1: Species numbers and percentage change in species numbers for historic primate families recorded at three time points by Honacki et al. (1982), Wilson & Reeder (2005) and the IUCN species list from 2016 (documented in Estrada, et al., 2017).

Disproportional splitting among taxonomic groups could have several causes. Variation in the number of new species described under the PSC across taxa could be driven by variation in the rate at which lineages evolve such that phylogenetic species are tracking cryptic diversity or incipient species formation among rapidly evolving taxa. In this case, new species listed under the PSC might point to situations where there is a discordance between patterns of genetic change and the evolution of gross morphological changes used by traditional taxonomists. This could be due to ecology – if some lineages are diversifying along ecological axes that we do not notice - or due to demographics – if some lineages have ecologies and/or histories that lead to faster local genetic coalescent times and so diagnosability. Under either of these scenarios, variation in inflation would simply reflect underlying biological reality. However, it is also possible that splitting is driven by other factors that may indeed be prone to bias. Critics have argued that much inflation is artificial,

reflecting major shortcomings of applying the PSC and a reliance on insufficient data (Zachos, et

al., 2013; Zachos & Lovari, 2013). Zachos et al. (2013) provides evidence suggesting unwarranted

splitting in select cases, advocating that the PSC has been taken to a "molecular extreme". In one

cited example (Zachos, et al., 2013), Sumatran tigers (previously *Panthera tigris sumatrae*, raised

to *Panthera sumatrae*) were split by Cracraft et al. (1998) and then subsequently split into an even

greater number of species by Mazák & Groves (2006), based on diagnosable mitochondrial and

craniometrics differences, respectively. By this logic, groups may continue to be split as an

increasing amount of molecular work leads to the continual identification of increasingly exclusive

diagnostic features such that we are likely to find more PSC species the more we look for them.

It is also possible that conservation interest in particular groups could motivate splitting. Limited

funding for conservation research is increasingly focused on 'biodiversity hotspots' and it has been

suggested that researchers could have a vested interest in declaring taxa in these regions as endemic

species (Karl & Bowen, 1999; Isaac, et al., 2004). There is evidence from some charismatic groups

that taxa receive more conservation attention and funding when comprised of multiple, small, and

taxonomically distinct populations (e.g., African apes; Stanford, 2001; Oates, 2006; Gippoliti &

Amori, 2007). Species lists are often used to determine which groups should receive conservation

attention (Mace, 2004) and changing the way we define species can also change which groups

receive action.

Along with being potentially biased toward certain taxonomic groups and related to the point made

above, splitting under the PSC has been suggested to result in individually more imperilled

populations which could negatively impact conservation agendas (Agapow, et al., 2004; Isaac, et

al., 2004; Morrison, et al., 2009; Zachos, 2015; Robuchon, et al., 2019). One criterion used by the

IUCN to classify species as imperiled is population size: species may be designated "Vulnerable"

if there are fewer than 1000 mature individuals found in the wild and "Endangered" if there are

fewer than 250 (Agapow, et al., 2004; Frankham, et al., 2012). Thus, splitting one species into

several new species may result in one or more receiving a (more) imperilled status (Agapow, et

al., 2004; Isaac, et al., 2007). This could lead to seemingly rare but poorly-defined species being

prioritized over well-defined and perhaps biologically more distinctive species (Pillon & Chase,

2007). Recent evidence suggests species splitting is not a driver of threat status for birds (Simkins,

et al., 2020), however, it is unknown whether these findings are generalizable across other

taxonomic groups (Garnett & Thomson, 2020).

Here, we set out to better understand the causes and consequences of taxonomic inflation in

primates. We test (i) whether taxonomic inflation is associated with the amount of research being

done in fields suggested to motivate splitting, or with a lineage's underlying diversification rate,

and (ii) whether rates of inflation dictate which groups are considered to be most imperilled.

Because many newly described primate lineages have previously been described as subspecies or

subpopulations of multiple closely related BSC species, we ask these questions at the genus level.

To explore our first question regarding predictors of inflation, we consider whether biological

factors or measures of human-induced bias explain increases in species numbers. To test potential

human-induced biases in splitting, we consider estimates of research effort for each taxon, and

predict that more research done on a given taxon may be associated with more splitting. We

examine research effort in two fields: molecular genetics (since molecular work could cause

species to be split continuously as finer molecular distinctions are made) and conservation (since

splitting has been suggested to be motivated by conservation interests). To explore possible

biological explanations for inflation trends, we test to see if inflation is explained by diversification

rate, using a diversification measure designed to be uncoupled from changes in species naming

conventions (see Methods). Lineages with high recent diversification rates are expected to contain

more incipient or cryptic species than lineages diversifying at a lower rate, such that diversification

rate might be expected to correlate positively with inflation.

To explore our second question linking inflation and risk, we test whether inflation is associated

with a change in the number of threatened species listed in genera through time. Ideally we would

look at changes in a weighted measure of threat score that differentiates between threat categories

of varying severity (e.g., using the Red List Index (RLI); Butchart, et al., 2007; Bubb, et al., 2009);

however, criteria for inclusion in Red List threat categories have changed considerably over time,

meaning weights assigned to categories for RLI calculations do not match up with categories used

when the PSC was first introduced (e.g., see statuses in IUCN Conservation Monitoring Centre,

1986). Because of these changes in Red List criteria, we instead ask whether clades with species

that have been split more frequently have observed a greater increase in their proportion of

threatened species (defined below) through time in comparison to those which have been split less

frequently. We predict that if splitting is driving an increase in threatened species, we should

observe a positive association between inflation and change in the proportion of threatened species

listed in primate genera.

METHODS:

Data

To measure the number of primate species that were described before the introduction of the PSC,

we used the last pre-PSC taxonomy, that of Honacki et al. (1982). This taxonomy contains 181

species and is considered a reliable estimate of the number of species thought to exist during the

popularity of the BSC (see Rylands & Mittermeier, 2014). We then recorded if each species in this

taxonomy was historically considered to be threatened by referencing the most complete IUCN

Red List published around the same time (IUCN Conservation Monitoring Centre, 1986). This

taxonomy was contrasted with the IUCN taxonomy from 2016 and attendant data documented in

Estrada et al. (2017), which lists 503 species. For each of these 503 species we noted their

taxonomic placement (genus and family), whether or not they were considered threatened

(VU=Vulnerable, EN=Endangered, or CR=Critically Endangered), and biogeographic region. For

each species described by the IUCN in 2016 that was not listed in Honacki et al. (1982), we scored

whether the species was a "*de novo*" species description (Burgin, et al., 2018), i.e., 40 cases where

a new species had not been previously formally identified as a subspecies or subpopulation of

another species prior to splitting.

We compiled all the species listed by the IUCN in 2016 into 12 families and 50 genera found in

Honacki et al. (1982). Family name "Callimiconidae" in Honacki et al. (1982) was not used as this

taxon has since been recognized as a genus of the larger family "Callitrichidae" (Wilson & Reeder,

2005). *Rungwecebus kipunji* was removed from the IUCN species list from 2016 as it represents a

newly discovered genus that does not collapse into any of the genera provided by Honacki et al.

(1982).

Research effort in the fields of molecular genetics and conservation were estimated for each genus

through a literature review of publications in the Web of Science Core Collection published

between 1983 (the year the PSC was first proposed) and 2016. To obtain papers in the field of

conservation we searched the genus name AND "conservation" (hereafter 'conservation research

effort'). To obtain papers in the field of molecular genetics we searched for genus name AND the

following terms: "mitochondrial DNA" OR "barcoding" OR "bar-coding" OR "cytochrome b" OR

"phylogeography" OR "microsatellites" OR "micro-satellites" OR "population genetics" OR

(species AND genetics) OR (taxonomy AND genetics) OR "SNP" (hereafter 'molecular genetics

research effort'). All hits from these searches were reviewed and irrelevant papers were eliminated

(see supplementary materials; Tables S1 and S2). In this study we used the genera listed in Honacki

et al. (1982) (n=50), many of which have since been further separated into multiple genera. Thus,

when appropriate we included new genus names in addition to those listed by Honacki et al. (1982)

in the literature search (see Table S3). Additional details on how searches were conducted are

provided in the supplementary materials.

Diversification rate was estimated with the method-of-moments approach described in Magallon

& Sanderson (2001) (i.e., ln(taxa richness)/stem age). Diversification rate estimates generated

using this method often rely on species numbers as their estimate of taxa richness, meaning that

diversification rate estimates are inherently biased by the inflation phenomenon we are studying

(i.e., highly inflated genera will receive disproportionally high diversification rates). Therefore,

richness scores for our diversification rate calculations were determined as the counts of the well-

resolved "lineages" described in Creighton et al. (2021). These lineages were determined by

creating a time cut-off in the 10kTrees consensus primate phylogeny (Arnold, et al., 2010) in an

attempt to eliminate very young PSC species descriptions and obtain a consistent (unbiased)

estimate of diversity across clades. These lineages were assigned to each of the 50 genera described

in Honacki et al. (1982). Diversification rate was then estimated by taking the natural log lineage

richness for each genus and dividing by the stem age of that genus (Magallon & Sanderson, 2001).

Stem ages for each genus were extracted from the 10kTrees consensus phylogeny (version 3)

(Arnold, et al., 2010) trimmed to contain a single branch representing each genus. During this

process, there were several instances where genera described in Honacki et al. (1982) were non-

monophyletic within the more recent primate phylogeny we used (Arnold, et al., 2010), making it

unclear how to assign a divergence date for these clades. We therefore removed eight genera from

the analyses where diversification rate was a variable of interest: *Presbytis*, *Lemur*, *Galago*,

Cebuella, *Cercocebus*, *Cercopithecus*, *Papio*, and *Pygathrix*. We discuss the limitations of having

to remove these genera from these analyses in the discussion.

Analysis

To test our questions about the predictors of species-splitting and its consequences for

conservation, we fit a series of linear effects and mixed effects models. We note that the response

variables used in these analyses (i.e., measures of inflation and extinction risk) are likely to be

phylogenetically clustered, and phylogenetic models could be used to account for this influence of

phylogeny; however, many genera listed by Honacki et al. (1982) are non-monophyletic, making

it unclear how to designate them a single branch in modern phylogenies (see discussion above on

diversification rate). Importantly, after accounting for regional differences, family contributed

little to no variance in any of our models, indicating that phylogenetic relationships at that level

were not confounding our results. We provide further discussion on model choice in the
supplementary materials.

Data were analysed using R version 4.0.5 (R Core Team, 2021).

Predictors of Inflation

To determine if measures of potential human bias (i.e., research effort) or diversification rate

explain discrepancies in inflation across taxa, we tested to see if these variables were significantly

associated with the number of species added to primate genera since the introduction of the PSC

while controlling for the original number of described species (i.e., species number prior to the

PSC) and region. We fit three generalized linear mixed effects models with Poisson distributions

using the lme4 package in R (Bates, et al., 2014), and obtained p-values using the lmerTest package

(Kuznetsova, et al., 2017). In these models the response variable was the number of species

assigned to a given genus by the IUCN in 2016 that had not been previously described by Honacki

et al. (1982). Each model had either conservation research effort, molecular genetics research

effort or diversification rate included as a fixed effect, as well as region and the number of species

in the genus per Honacki et al. (1982) to control for their effects on inflation. The natural logarithm

of the number of species listed for each genus in Honacki et al. (1982) was included as both a

linear and a quadratic term following inspection of raw plots and plots against scaled residuals

from the simulation output. A square-root transformation was used on molecular genetics research

effort to decrease the impact of outliers on model fit. To assist with model stability and

convergence, we scaled all continuous variables in the model to have a mean of zero and standard

deviation of one (Becker, et al., 1988). Mainland Africa and Asia (hereafter mainland Africa +

Asia) were grouped together and served as the baseline region in our models based on previous

studies that have shown that the taxonomy of primates from these regions has been relatively stable

compared to Madagascar and the Neotropics (e.g., Isaac & Purvis, 2004; Isaac, et al., 2004;

Tattersall, 2007). We also chose to group Asia and Africa together because one genus (*Macaca*)

is found in both regions. Genus ID was included as a random effect in these models to correct for

overdispersion. Family (nested within region) was originally included as a random effect but

contributed very little to model fit and created issues with convergence due to overfitting, and thus

was dropped from the final models. We tested potential interaction terms with all variables and

region to test for regional effects, but none were significant and so these terms were also dropped

from the final models. Before running models with other predictors included, we also ran a model

including only the linear and quadratic terms for the number of species listed for each genus in

Honacki et al. (1982) as predictors to assess their association with inflation in the simplest model.

We ran all models a second time after removing the 40 *de novo* species from our response (Tables

S4, S5 and S6). We checked model assumptions and fit by plotting residuals versus the fitted

values and versus each covariate in the model. Residual plots and analyses with the Diagnostics

for Hierarchical Regression Models (DHARMa) R package (Hartig, 2017) indicated acceptable

model fits.

Inflation and Threat Score

To test whether taxonomic inflation over time is associated with a change in the proportion of

threatened species within genera, we conducted a two-step (hierarchical) analysis on species'

threat probability between 1982 to 2016. We first fit a generalized linear mixed effects model with

a binomial distribution for the number of threatened and non-threatened species within a genus

using the lme4 package in R (Bates, et al., 2014). In this first model, the response estimated the

proportional counts of species at risk within genera (equivalent to the per-species probability of

threat) in the periods of 1982 and 2016. Predictors for this model included fixed effects for the

time period (the baseline of 1982 and the change to 2016), the region that encompasses each genus'

natural distribution (Madagascar, Neotropics, mainland Africa + Asia), and an interaction between

region and time period to account for geographic differences in threat probabilities through time.

Genus identity was included as a random intercept, to account for repeated measures in 1982 and

2016, and as a random slope with time period to account for differences in changes to threat

probabilities among genera. As discussed above, we included the taxonomic rank of family (nested

within region) as a random effect but found that the variance in threat probabilities among families

was minimal and that including this term also generated model convergent issues; given this, we

subsequently omitted family from our models. Residual plots and analyses with the DHARMa R

package (Hartig, 2017) indicated acceptable model fits for the final model. From this first model

describing genus level changes in threat probabilities, we then used the REextract function

implemented in the merTools package in R (Knowles & Frederick, 2020) to extract the genus level

random slopes for time period and their associated standard errors, giving us an estimate of the

varying effect of change in threat probability among genera (while conditioning on regional trends)

between 1982 and 2016. We then fit a linear model where varying effect of change in threat

probability from 1982 to 2016 was the response, and the proportional change in species within

each genus (i.e., the number of new species in the IUCN species list from 2016 / original number

of species in Honacki et al. (1982) – our measure of "inflation") – was the predictor. We weighted

each estimate of genus level change in threat probability by its standard error (w = 1/SE) to

propagate the error of model estimated random effects. We ran this model a second time after

removing *de novo* species descriptions from our response (Table S7). Raw data used for these tests

are visualized in Figure 2.

Plots of the raw data indicated that patterns in inflation, the proportion of species at risk in primate

genera today, and the changes the proportion of threatened species in primate genera between 1982

and 2016 show regional differences (Figures S1 and S2). We ran a subsequent set of models to test

for regional variation in the effect of taxonomic inflation. In this analysis, we removed region from

the first generalized linear mixed effects model and subsequently included region interacting with

taxonomic inflation as a predictor of the varying effect of change in genus threat probability in the

second-order model. However, interaction effects were not significant in this model and thus we

only considered the results of the first set of models reported above.

<u>RESULTS:</u>

Predictors of Inflation

None of our measures – conservation research effort ($\beta = -0.220$; $p = 0.180$; Table S4), molecular

genetics research effort ($\beta = -0.188$; $p = 0.212$; Table S5) nor diversification rate ($\beta = 0.132$; $p =$

0.477; Table S6) – were significantly associated with taxonomic inflation of primate genera.

Removing *de novo* species did not impact this pattern of results (Tables S4, S5 and S6). Notably,

in addition to sharing a significant linear relationship with inflation as expected (Tables S4, S5 and

S6), the quadratic term added for the original number of species in 1982 was significant in the

model with diversification rate, indicating a downwardly concave association with inflation ($\beta = -$

1.890; $p = 0.042$; Table S6); this term approached statistical significance in models that included

conservation or molecular genetics research effort as predictors ($\beta = -1.529$; $p = 0.072$; Table S4

and β= -1.626; p= 0.057; Table S5), indicating that the positive increase in splitting by initial genus

size decelerates as genera become very large, even after accounting for other factors. We note that

this quadratic term was also significant without other predictors (i.e., research effort or

diversification rate) present in the model, both with *de novo* species included in the response (β=

-2.268; p= 0.011) and without *de novo* species included in the response (β= -2.086; p= 0.013).

Inflation and Threat Score

Looking at the raw data on inflation and proportion of species at risk for all genera, there was no

clear indication that highly inflated genera have observed a disproportional increase in perceived

extinction risk (Figure 2).

Figure 2: Scatterplots with trendlines showing the change in the proportion of species identified as being threatened in primate genera in 1982 and 2016 painted by inflation a) including *de novo* species, and b) excluding *de novo* species. Total number of species in each genus is indicated by point size.

Results from our GLMM confirmed that inflation did not predict a change in threat score through

time (β= 0.002; p= 0.975; Table S7; Figure 3). Removing *de novo* species did not impact this

pattern of results (Table S7; Figure S3).

<u>Figure 3:</u> Inflation versus varying effect of change in threat probability (i.e., our measure of genus

level change in threat probability) (β= 0.001; p= 0.975; Table S7). Vertical bars indicate standard

errors on varying effect of change in threat probability values.

<u>DISCUSSION:</u>

Our results support neither a strong biological nor a strong interest-driven mechanism for variation

in new species designations across primate genera. If the naming of new PSC species captures

incipient speciation, we predicted that diversification rate might predict inflation. However, we

found no evidence for this as diversification rate was not a significant predictor of inflation in our

models. We also tested the prediction that splitting is motivated by conservation interest (Karl &

Bowen, 1999; Isaac, et al., 2004) or by increasing molecular research within certain taxa, with for

example, more new species designations arising as more population genetics work is done (Zachos,

et al., 2013; Zachos & Lovari, 2013). However, we found no evidence that research effort in either

of these areas was associated with inflation across genera. We did find that the positive increase

in splitting by initial genus size decelerates as genera approach the largest sizes. This could indicate

that inflation is being driven by something not captured by our models. Therefore, future studies

could aim to elucidate the origins of this decelerating association.

While inspection of the raw data suggested that genus level measures of inflation, threat

probability, and change in threat probability all varied by region, we found that there was no

cohesive signal of inflation leading to higher threat probability across these genera and regions.

Therefore, while there are idiosyncratic regional differences in both primate inflation and threat

probability there is no evidence for causal link between the two. While in some cases splitting may

lead populations to be considered more imperilled, overall, this analysis sends a positive message

that primates species threat statuses are not being overwhelmingly affected by splitting in line with

what has been reported for birds (Simkins, et al., 2020).

It should be considered that our analyses do have some limitations. First, quantifying research

effort of any kind is difficult due to the abundance of work published in different media. As such,

our estimates of research effort in conservation and molecular genetics may overlook some types

of research. More work should be done to determine if additional estimates of research bias can

explain increases in species numbers. For instance, cumulative funding estimates from various

sources (e.g., the IUCN, non-governmental organizations and regional governments) per taxon

could be used as an additional or alternative measure of conservation interest. Second, it was

necessary to remove eight non-monophyletic genera (some that have undergone substantial

splitting) from those analyses that included diversification rate, leading to a considerable reduction

in statistical power for those analyses. Third, due to changes in Red List criteria, species status,

and lack of information about which species had been assessed in the 1980s, we were not able to

consider differences in threat status severity (e.g., Vulnerable versus Endangered) when

considering associations between inflation and changes in threat score through time. This leaves

room for the possibility that change in risk could be under- or over-estimated for certain genera.

Future studies could look at the association between inflation and changes in weighted measures

of threat score (e.g., RLI) over time using a more recent starting point (i.e., after 1982) where Red

List criteria become stable and there is available information on which non-threatened species have

been evaluated.

Overall, we do not find support for biological processes (i.e., diversification rate) or human

imposed bias (i.e., measures of research effort) driving inflation across primate genera. We also

find no cohesive signal of inflation leading to higher threat probabilities across primate genera.

Generally speaking, relying on species as the central unit of conservation and primary object of

biological study behooves taxonomy to remain stable, while changing ideas about the concept of

species makes taxonomy inherently unstable (Mayr, 1996). We suggest that areas of research

requiring consistent estimates of diversity (i.e., conservation) may benefit from either (i) adopting

stable species criteria, i.e., that do not rely on increasingly fewer diagnostic distinctions; or, for

applied conservation specifically (ii) shifting more resources toward regional management efforts

that are less likely to be influenced by changing species designations. It is well-known that closely

related species are more similar to one another than they are to more distantly related taxa. Thus,

treating all species independently and of equal weights in conservation listing may not lead to

desired outcomes (Redding & Mooers, 2010). As of 2016, approximately 60 percent of all primate

species were threatened with extinction according to the IUCN Red List of Threatened Species

(Estrada, et al., 2017), making it imperative that conservation efforts are spent wisely to ensure

optimal conservation of primate biodiversity writ large.

ACKNOWLEDGEMENTS: This work was supported by the Natural Sciences and Engineering

Research Council of Canada (NSERC) Discovery Grants Program (SMR and AOM, grants #2017-

04720 and #2019-04950) and the Canada Foundation for Innovation (CFI) (SMR, grant #29433).

MJAC was supported by awards from the Biodiversity, Ecosystem Services, and Sustainability

(BESS) program and McGill University. We thank the Reader and Guigueno labs at McGill, Brian

Leung, Hans Larsson, William Wcislo, Isabella Capellini, and the Crawford Lab for Evolutionary

Studies at SFU for feedback on portions of the manuscript and study design. Special thanks to

Aiman Hadif for his assistance in collecting and organizing data and to Paul Sims and Dan

Greenberg for crucial statistical advice.

DECLARATIONS OF INTEREST: None.

Data and code accompanying the manuscript will be made available on publication through

Zenodo.

REFERENCES:

Agapow, P. M., Bininda-Emonds, O. R., Crandall, K. A., Gittleman, J. L., Mace, G. M., Marshall,

J. C., & Purvis, A. (2004). The impact of species concept on biodiversity studies. The

Quarterly Review of Biology, 79: 161–179.

Arnold, C., Matthews, L. J., & Nunn, C. L. (2010). The 10kTrees website: a new online resource

for primate phylogeny. Evolutionary Anthropology, 19: 114–118.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2014). lme4: Linear mixed-effects models

using Eigen and S4. R package version 1.1.21. https://cran.r-

project.org/web/packages/lme4.

Becker, R. A., Chambers, J. M., & Wilks, A. R. (1988). The new S language: A programming

environment for data analysis and graphics. Wadsworth & Brooks/Cole, Pacific Grove.

Bubb, P. J., Butchart, S. H. M., Collen, B., Dublin, H., Kapos, V., Pollock, C., Stuart, S. N., &

Vié, J. –C. (2009). IUCN Red List Index—guidance for national and regional use. IUCN,

Gland.

Burgin, C. J., Colella, J. P., Kahn, P. L., & Upham, N. S. (2018). How many species of mammals

are there? Journal of Mammalogy, 99: 1–14.

Butchart, S. H., Akçakaya, H. R., Chanson, J., Baillie, J. E., Collen, B., Quader, S., ... & Hilton-

Taylor, C. (2007). Improvements to the Red List Index. PLoS One, 2: e140.

Cotton, A., Clark, F., Boubli, J. P., & Schwitzer, C. (2016). IUCN Red List of threatened primate

species. In S. A. Wich & A . J. Marshall (Eds.), An introduction to primate conservation

(p. 31–38). Oxford University Press, Oxford.

Cracraft, J. (1983). Species concepts and speciation analysis. In R. F. Johnson (Ed.), Current

ornithology (p. 159–187). Plenum Press, New York.

Cracraft, J., Feinstein, J., Vaughn, J., & Helm-Bychowski, K. (1998). Sorting out tigers (*Panthera*

tigris): mitochondrial sequences, nuclear inserts, systematics, and conservation genetics.

Animal Conservation, 1: 139–150.

Creighton, M. J. A., Greenberg, D. A., Reader, S. M., & Mooers, A. Ø. (2021). The role of

behavioural flexibility in primate diversification. Animal Behaviour. Advance online

publication.

Donoghue, M. J. (1985). A critique of the biological species concept and recommendations for a

phylogenetic alternative. Bryologist, 88: 172–181.

Estrada, A., Garber, P. A., Rylands, A. B., Roos, C., Fernandez-Duque, E., Di Fiore, A., ... &

Rovero, F. (2017). Impending extinction crisis of the world's primates: Why primates

matter. Science Advances, 3: e1600946.

Frankham, R., Ballou, J. D., Dudash, M. R., Eldridge, M. D., Fenster, C. B., Lacy, R. C., ... &

Ryder, O. A. (2012). Implications of different species concepts for conserving biodiversity.

Biological Conservation, 153: 25–31.

Garnett, S. T., & Thomson, S. A. (2020). Are the implications for conservation of a major

taxonomic revision of the world's birds' simply serendipity? Animal Conservation, 23:

355-356.

Gippoliti, S., & Amori, G. (2007). The problem of subspecies and biased taxonomy in conservation

lists: the case of mammals. Folia Zoologica, 56: 113–117.

Groves, C. P. (2014). Primate taxonomy: inflation or real? Annual Review of Anthropology, 43:

27–36.

Groves, C., & Grubb, P. (2011). Ungulate taxonomy. JHU Press, Baltimore.

Hartig, F. (2017). DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression

models. R package version 0.3.0. https://cran.r-project.org/web/packages/DHARMa.

Honacki, J. H., Kinman, K. E., & Koeppl, J. W. (1982). Mammal species of the world; a taxonomic

and geographic reference. Allen Press, Inc. and Association of Systematic Collections,

Lawrence.

Isaac, N. J., Mallet, J., & Mace, G. M. (2004). Taxonomic inflation: its influence on macroecology

and conservation. Trends in Ecology & Evolution, 19: 464–469.

Isaac, N. J., & Purvis, A. (2004). The 'species problem' and testing macroevolutionary

hypotheses. Diversity and Distributions, 10: 275–281.

Isaac, N. J., Turvey, S. T., Collen, B., Waterman, C., & Baillie, J. E. (2007). Mammals on the

EDGE: conservation priorities based on threat and phylogeny. PLoS One, 3: e296.

IUCN Conservation Monitoring Centre. (1986). 1986 IUCN Red List of threatened animals.

IUCN, Gland and Cambridge.

Karl, S. A., & Bowen, B. W. (1999). Evolutionary significant units versus geopolitical taxonomy:

molecular systematics of an endangered sea turtle (genus *Chelonia*). Conservation

Biology, 13: 990–999.

Knowles, J. E., and Frederick, C. (2020). merTools: Tools for Analyzing Mixed Effect Regression

Models. R package version 0.5.2. https://CRAN.R-project.org/package=merTools.

Kuznetsova, A., Brockhoff, B., & Christensen, H. B. (2017). lmerTest Package: Tests in Linear

Mixed Effects Models. Journal of Statistical Software, 82: 1–26.

Mace, G. M. (2004). The role of taxonomy in species conservation. Philosophical Transactions of

the Royal Society B: Biological Sciences, 359: 711–719.

Magallon, S., & Sanderson, M. J. (2001). Absolute diversification rates in angiosperm clades.

Evolution, 55: 1762–1780.

Mayr, E. (1963). Animal species and evolution. Belknap Press, Cambridge.

Mayr, E. (1996). What is a species, and what is not? Philosophy of Science, 63: 262–277.

Mazák, J. H., & Groves, C. P. (2006). A taxonomic revision of the tigers (*Panthera tigris*) of

Southeast Asia. Mammalian Biology, 71: 268–287.

Morrison, W. R., Lohr, J. L., Duchen, P., Wilches, R., Trujillo, D., Mair, M., & Renner, S. S.

(2009). The impact of taxonomic change on conservation: Does it kill, can it save, or is it

just irrelevant? Biological Conservation, 142: 3201–3206.

Oates, J. F. (2006). Is the chimpanzee, *Pan troglodytes*, an endangered species? It depends on what

"endangered" means. Primates, 47: 102–112.

Pillon, Y., & Chase, M. W. (2007). Taxonomic exaggeration and its effects on orchid

conservation. Conservation Biology, 21: 263–265.

R Core Team. (2021). R: A Language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

Redding, D. W., & Mooers, A. Ø. (2010). Can systematists help decide the relative worth of bits

of biodiversity. The Systematist, 32: 4–8.

Robuchon, M., Faith, D. P., Julliard, R., Leroy, B., Pellens, R., Robert, A., ... & Pavoine, S. (2019). Species splitting increases estimates of evolutionary history at risk. Biological Conservation, 235: 27–35.

Rylands, A. B., & Mittermeier, R. A. (2014). Primate taxonomy: species and conservation. Evolutionary Anthropology, 23: 8–10.

Simkins, A. T., Buchanan, G. M., Davies, R. G., & Donald, P. F. (2020). The implications for conservation of a major taxonomic revision of the world's birds. Animal Conservation, 23: 345-352.

Stanford, C. B. (2001). The subspecies concept in primatology: the case of mountain gorillas. Primates, 42: 309–318.

Tattersall, I. (2007). Madagascar's lemurs: cryptic diversity or taxonomic inflation? Evolutionary Anthropology, 16: 12–23.

Wilson, D. E., & Reeder, D. M. (2005). Mammal species of the world: a taxonomic and geographic reference (vol. 1). JHU Press, Baltimore.

Zachos, F. E. (2015). Taxonomic inflation, the phylogenetic species concept and lineages in the tree of life–a cautionary comment on species splitting. Journal of Zoological Systematics and Evolutionary Research, 53: 180–184.

Zachos, F. E., Apollonio, M., Bärmann, E. V., Festa-Bianchet, M., Göhlich, U., Habel, J. C., ... & Pertoldi, C. (2013). Species inflation and taxonomic artefacts—A critical comment on recent trends in mammalian classification. Mammalian Biology, 78: 1–6.

536   Zachos, F. E., & Lovari, S. (2013). Taxonomic inflation and the poverty of the Phylogenetic

537   Species Concept-a reply to Gippoliti and Groves. Hystrix, 24: 142–144.