A DROSOPHILA CIRCUIT FOR HABITUATION OVERRIDE

Swati Trisal[1,2,3], Marcia Aranha[2], Ankita Chodankar [1,2], K. VijayRaghavan[1] and Mani Ramaswami[1,2†].

[1]*National Centre for Biological Sciences, TIFR, Bangalore 560065, India.*

[2]*Trinity College Institute of Neuroscience, School of Genetics and Microbiology, Trinity College Dublin, Dublin-2, Ireland.*

[3]*School of Chemical and Biotechnology (SCBT), Shanmugha Arts, Science, Technology & Research Academy (SASTRA) University, Thanjavur 613401, India*

[†] **Correspondence**: mani.ramaswami@tcd.ie

ABSTRACT:

Habituated animals retain a latent capacity for robust engagement with familiar stimuli. In

most instances, the ability to override habituation is best explained by postulating that

habituation arises from the potentiation of inhibitory inputs onto stimulus-encoding assemblies

and that habituation override occurs through disinhibition. Previous work has shown that

inhibitory plasticity contributes to specific forms of olfactory and gustatory habituation in

Drosophila. Here we analyze how exposure to a novel stimulus causes override of gustatory

(proboscis-extension reflex or "PER") habituation. While brief sucrose contact with tarsal

hairs causes naïve *Drosophila* to extend their proboscis, persistent exposure reduces PER to

subsequent sucrose stimuli. We show that in so habituated animals, either brief exposure of

the proboscis to yeast or direct thermogenetic activation of sensory neurons restores PER

response to tarsal sucrose stimulation. Similar override of PER habituation can also be induced

by brief thermogenetic activation of a population of TH (Tyrosine-Hydroxylase) positive

neurons, a subset of which send projections to the subesophagial zone (SEZ). Significantly,

sensory-neuron induced habituation override requires transmitter release from these TH-

positive cells. Treatments that cause override specifically influence the habituated state, with

no effect on the naïve sucrose response across a range of concentrations. Taken together with

other findings, these observations in female flies are consistent with a model in which novel

taste stimuli trigger activity in dopaminergic neurons which, directly or indirectly, inhibit

GABAergic cells that drive PER habituation. The implications of these findings for general

mechanisms of attentional and sensory override of habituation are discussed.

22
23

SIGNIFICANCE STATEMENT:

Habituation can be overcome when a new context requires an enhanced response to a familiar stimulus. However, the underlying mechanisms remain incompletely understood. Previous studies have provided evidence that habituation of the sucrose-induced proboscis extension reflex (PER) in Drosophila occurs through potentiation of inhibition onto the PER pathway. This work defines controlled protocols for override of PER habituation and uses them to outline the underlying circuit mechanisms. The results presented support a model in which novel taste stimuli cause dishabituation by activating a subset of tyrosine-hydroxylase-expressing neurons that inhibit GABAergic neurons whose potentiation underlies PER habituation. At a general level, these findings further highlight a central role for inhibition and disinhibition in the control of behavioral flexibility.

INTRODUCTION.

Habituation is a form of non-associative learning in which the response to a stimulus reduces after repeated or extended passive exposure. However, a latent ability to respond to the innocuous stimulus remains. "Dishabituation" is a classical, defining feature of the habituated state, distinguishing it from sensory or synaptic fatigue (Thompson and Spencer, 1966; Rankin et al., 2009; Ramaswami, 2014). In this article we often use the term "override" in place of dishabituation, which while largely synonymous, better acknowledges that behavioral responses can be reinstated by not only classic dishabituating (novel) stimuli, but also attentional mechanisms recruited during task engagement (Cooke and Ramaswami, 2020). Habituation and the associated phenomenon of dishabituation/ override are well known across different phyla of animal kingdom (Glanzman et al., 1989; Zaccardi et al., 2004; Smith et al., 2009; Ramaswami, 2014). Two broad observations are relevant here. First, mechanisms underlying very short and longer lasting forms of habituation may differ; some forms of the

latter are known to involve inhibitory potentiation (Ramaswami, 2014; Shen et al., 2020).

Second, largely due to the difficulty of necessary behavioral experiments, in most (but not all

e.g. Semelidou et al., 2018) instances, habituation override is not distinguished from a potential

confounding process of response sensitization (Castellucci et al., 1970; Hawkins et al., 1998;

Asztalos et al., 2007a; Asztalos et al., 2007b). Thus, although arguments and evidence support

a disinhibitory mechanism (Fischer et al., 1997; Das et al., 2011; Kato et al., 2015; Ogg et al.,

2018) the neural pathways and mechanisms of habituation override remain incompletely

characterized. Here we address this issue in the gustatory system of *Drosophila*.

Drosophila sample and taste potential foods via chemosensory hairs on their tarsi and their

proboscis (Stocker, 1994; Montell, 2009). Sugars detected by sensory hairs trigger a proboscis

extension reflex (PER), which enables feeding (Minnich, 1921; Dethier, 1976). The neural

circuit for PER is only partially understood. Sensory information is processed in subesophagial

zone (SEZ) and then communicated to command neurons whose activation triggers the motor

programme required for coordinated contraction and relaxation of muscles that drive proboscis

extension and retraction (Flood et al., 2013). Repeated sucrose stimulation of the tarsus under

conditions where proboscis extension is futile, leads to reduced PER response through a

process that shows several classic features of habituation (Duerr and Quinn, 1982; Le Bourg,

1983; Fois et al., 1991; Engel and Wu, 2009; Paranjpe et al., 2012). Importantly, in habituated

animals, PER to sucrose is quickly restored if the fly is presented with a strong, novel sensory

stimulus (Le Bourg, 1983; Fois et al., 1991; Paranjpe et al., 2012). Here, we investigate circuit

mechanisms that drive override of PER habituation.

We first reproduced prior experiments providing key support for increased inhibition in the

PER pathway being the core mechanism for PER habituation (Paranjpe et al., 2012).

Thereafter, we addressed mechanisms of habituation override, which we found could be

achieved by yeast stimulation of the proboscis, thermogenetic activation of yeast-responsive

or bitter responsive sensory neurons, or activation of a dopaminergic neuron subpopulation.

We show that sensory stimulation procedures induce habituation override through

dopaminergic neuron activation. Crucially, each dishabituation protocol specifically affects the

response of habituated animals; none sensitize the naïve response. While the data do not yet

conclusively define all elements of the dishabituation circuit or the mechanism by which

dopaminergic neurons trigger override, they support a model in which novel stimuli induce

dopamine release in the SEZ, which acts to directly or indirectly inhibit inhibitory neurons that

drive PER habituation. We suggest that this work: (a) circumscribes core elements of a

sensory-central circuit for habituation override; (b) provides evidence for a new disinhibitory

pathway in the Drosophila brain; and (c) supports an emerging framework in which latent

perceptions, memories and behaviors may be generally activated through disinhibition

(Sridharan and Knudsen, 2015; Barron et al., 2017; Wang and Yang, 2018).

MATERIALS AND METHODS

Experimental Design:

Drosophila stocks*:* Fly stocks were maintained on standard corn meal media. Canton S (CS)

flies were used as wild-type controls unless otherwise stated. The stocks were obtained either

from stock centres or as generous gifts from following sources: *Ir25a-Gal4* was provided by

Carlos Ribeiro (Champalimaud Centre for the Unknown, Lisbon, Portugal), *TH-C'-Gal4*, *TH-*

D'-Gal4 and *TH-C-Gal80* were generously provided by Mark Wu (Johns Hopkins University,

Baltimore, MD), *Gad1-Gal4* was from Gero Miesenbock (Oxford University, Oxford, UK),

TH-Gal4 was provided by Gaiti Hassan (National Centre for Biological Sciences, Bangalore,

India), *Gr66a-LexA, lexAop-CD4::spGFP11; UAS-CD4::spGFP1-10* and *LexAop-TRPA1*

were provided by Kristin Scott (University of California, Berkley). *rut²⁰⁸⁰* and *UAS-rut⁺* were

provided by Martin Heisenberg. *UAS-Shiᵗˢ* was obtained from Toshi Kitamoto (University of

Iowa, Iowa city, IA), *UAS-TRPA1* was provided by Paul Garrity (Brandeis University,

Waltham, MA)**.** *Wg/CyO; Gr66a-Gal4* (BL 57670), *UAS-mCD8::GFP* (BL 5130), *UAS-*

CD8::RFP, *LexAop-CD8::GFP* (BL 32229) were obtained from Bloomington Drosophila

Stock Centre. Dopamine receptors were downregulated using lines *UAS-DopR1-miR, UAS-*

DopR2-miR and *UAS-D2R-miR* obtained from Mark Wu (Johns Hopkins University,

Baltimore, MD) in *Gad1-Gal4* expressing neurons (data not shown). Tyrosine hydroxylase was

downregulated in *TH Gal4* neurons using TH-miR-G/TM6, Sb and TH-miR-9/Cyo; TH-miR-

2/TM6, Tb provided by Mark Wu (Johns Hopkins University, Baltimore, MD) (data not

shown). However, the results were inconclusive.

Proboscis extension behaviour*:* Proboscis extension behaviour was carried out as described

in (Paranjpe et al., 2012). Briefly, 3-4 days old female flies were mounted individually, ventral

side up, on a cover slip. The flies were kept in a humidified chamber for 2 hours for recovery.

Tarsal hairs were stimulated with 2% sucrose using 1ml syringe needle. The naïve response

score was determined after stimulating tarsal hairs five times. Failed PER response was scored

as zero and complete proboscis extension was scored as one. Flies that showed naïve PER

response less than three out of five times were discarded and not used for habituation

experiments.

Habituation of proboscis extension reflex: Tarsal hairs were stimulated with 10% sucrose for

10 minutes after which the legs were washed using distilled water. The habituated response or

post exposure response was recorded by stimulating tarsus with 2% sucrose five times, similar

to naïve response.

Habituation Override 10% yeast was presented to the proboscis over one minute, in a "spaced

manner" to prevent consumption of yeast and satiation. In gist, the yeast paste was applied to

labellar bristles for about a second but withdrawn rapidly before consumption could occur and

this was repeated (about 6 times) over a one minute period. After exposure to yeast, flies were

allowed to groom for 1 minute so that yeast sticking to the proboscis could be cleaned. PER

responses were subsequently recorded as usual.

For habituation override using mechanical stimulus, flies mounted on the cover slip were

placed in a petri dish (35mm) and vortexed for 1 minute on a Scientific Industries Vortex-

Genie 2 instrument at the highest setting. Flies were allowed to recover from the shock for 1

minute before testing the response.

To distinguish override from sensitization, naïve response of flies were recorded at

concentrations of sucrose lower than 2%. This was done as the PER response is maximum at

2%, any difference as a result of manipulation would have been difficult to observe. Four

concentrations of sucrose were tested- 0.1%, 0.5%, 1% and 2%. Flies were then exposed to

either stimulus used for override or heat, for 1 minute. Post response was tested after 1 minute

rest at RT.

Heat-mediated manipulation For TRPA1 and Shi[ts] experiments, flies were transferred to 32°

and 34°C respectively on a dry bath for 1 minute. In case of Shi[ts] experiments, proboscis was

stimulated with 10% yeast while at 34°C. Flies were given 1 minute rest at RT before testing

PER response.

Immunohistochemistry: Adult brains were dissected in 1X PBS and fixed in 4%

paraformaldehyde diluted in 1X PBS with 0.3% Triton-X (PTX) for 30 minutes at room

temperature. Samples were washed with 0.3% PTX for 20 minutes three times at room

temperature and incubated with primary antibody for 48 hours at 4°C on a shaker. Samples

were again washed three times with 0.3% PTX for 20 minutes each. Secondary antibodies

conjugated with Alexa Fluor-488, Alexa Fluor- 568 and Alexa Fluor- 647 (1:500, Invitrogen),

diluted in 0.3% PTX, were added and samples were incubated for 24 hours. Brain samples

were again washed repeatedly three times for 20 minutes each and mounted in Vectashield (H-

1000, Vector laboratories) on a glass slide with spacers. Images were acquired using Zeiss

LSM 510 Meta microscope and Olympus Fluoview (FV- 3000).

Following primary antibodies were used: rabbit anti-GFP (1:100, invitrogen), mouse anti-

Bruchpilot (n82) (1:20, DSHB), mouse anti-GFP (Sigma, 1:100), rabbit anti-TH (1:1000,

Invitrogen), chick anti-GFP (Abcam, 1:5000), rabbit anti-DsRed (1:100, Clontech).

Statistical analysis:

Non parametric tests were used to analyse data. Friedman test was used to analyse habituation

override experiments which had three matched groups, followed by Dunn's multiple

comparisons post hoc test to determine difference among different treatments. Wilcoxon sign

rank test was used for comparing two paired groups whereas Mann Whitney U test was used

to compare two unpaired groups in Fig1C. All the data was analysed using GraphPad Prism

v8.4.2 software.

RESULTS.

Habituation of the proboscis-extension reflex to sucrose

Two-percent sucrose applied to tarsal hairs of immobilized, naïve flies induces robust and

reproducible proboscis extension response. However, following extended 10-minute exposure

to 10% sucrose solution, PER decreases from 92.41% to 28.96% (***p<0.0001, Wilcoxon test)

(Fig 1A-B). No change in naïve PER response is seen if water is presented for 10 minutes

instead of 10% sucrose (Fig1B). A previous study concluded that plasticity in central

GABAergic neurons is required for PER habituation (Paranjpe et al., 2012). Given their

significance for the inferred inhibitory mechanism for PER habituation, and because these

observations were made exclusively in male flies, we independently repeated key experiments

to re-examine: first, the need for the *rutabaga*-encoded adenyl cyclase in PER habituation; and

second, its reported sufficiency in *GAD1-Gal4* expressing, predominantly GABAergic neurons

for this function in our hands and in female flies. Our results were consistent with and

confirmed previously reported observations (Figure 1C).

Habituation is often distinguished from sensory or muscular fatigue by demonstrating the rapid

reinstatement of the sensory response by strong, novel stimuli (Rankin et al., 2009;

Ramaswami, 2014). Consistently, PER-habituated animals retain the ability to respond

relatively robustly to sucrose, as evidenced following strong or novel sensory stimulation.

Similar to olfactory habituation, strong mechanical stimulation (vortexing in a dish) for 1

minute substantially reinstates the sucrose-induced PER in habituated flies ($p < 0.001$, F-stat =

47.08, Dunn's multiple comparisons post-hoc test, *$p = 0.01$), without affecting baseline

sucrose sensitivity in naïve animals (Figure 1D-E; (Das et al., 2011; Paranjpe et al., 2012)).

Thus, the effect mechanical stimulation on PER is selective on the habituated state and

represents a form of habitation override and not sensitization. In order to address the

underlying circuit mechanisms, we first looked to identify more precisely defined sensory

stimuli that could cause override of PER habituation.

Novel stimuli override PER habituation

Drosophila are attracted by the taste of yeast. Gustatory receptor neurons (GRNs) that respond

to yeast components have been recently identified (Fischler et al., 2007; Wisotsky et al., 2011;

Ganguly et al., 2017); (Steck et al., 2018). To examine whether the novel taste of yeast could

influence PER habituation to sucrose, we applied a 10% yeast solution to the fly labellum,

which has previously been shown to possess yeast-responsive GRNs (Steck et al., 2018), and

tested if this resulted in override of sucrose habituation. Application of yeast to the labellum

substantially reinstated the PER response to tarsal sucrose stimulation in PER-habituated

animals ($p < 0.0001$, F-stat = 31.15, Dunn's multiple comparison post-hoc test, *$p = 0.014$; Fig

2A-B). In contrast, 10% sucrose solution, which should be familiar to the flies, when applied

to the labellum of habituated flies had no effect on PER (Fig 2B). Thus, a brief experience of

a novel and in this case, attractive stimulus appears capable of inducing habituation override.

We further looked to identify a single GRN class responsive to yeast components that may be

sufficient to drive dishabituation. To do this, we expressed heat-activated cation-permeable

TRPA1 channels in yeast-responsive GRNs and tested if heat-induced activity in these cells,

which bypassed the need for normal ligand-receptor interactions, would result in dishabituation

(Hamada et al., 2008). Such experiments showed that "thermogenetic" activation of the Ir25a

class of sensory neurons is sufficient to cause PER habituation override.

TRPA1 channels open at temperatures above 25°C; thus, activation of TRPA1-expressing

neurons can be temporally controlled by exposing experimental animals to temperatures where

the channel is either closed (RT) or open (above 25°C). We expressed TRPA1 in yeast

responsive *Ir25a-Gal4* positive sensory neurons (Steck et al., 2018). These flies were

habituated to sucrose at room temperature (RT, 21°C), transferred to 32°C post-habituation for

1 minute and then tested for PER. Thermogenetic activation of Ir25a-expressing GRNs was

sufficient to cause rapid override of sucrose habituation (Fig 2C). Thus, after Ir25a activation,

PER-habituated animals showed significantly increased PER to tarsal sucrose stimulation (p

<0.0001, F-stat $= 38.66$, Dunn's multiple comparison post-hoc test, $**p=0.0012$). Similar 32°C

exposure of genetic control animals not expressing TRPA1 did not affect PER habituation (Fig

2C).

To distinguish between the novelty and attractiveness of yeast taste as being primarily

instrumental in override, we also examined, whether thermogenetic activation of bitter-

compound responsive Gr66a expressing GRNs could similarly affect override of PER

habituation. Thus, we expressed TRPA1 in *Gr66a-Gal4* positive neurons and examined how

sucrose-responsiveness of flies habituated at RT was altered after a brief 1-minute shift to 32°C.

Sucrose induced PER increased from 6.67% to 56.29% (p<0.0001, F-stat = 39.85, Dunn's

multiple comparison post-hoc test, **p = 0.0042) indicating that brief activation of bitter-taste

sensing *Gr66a*-positive neurons is sufficient to induce dishabituation. The observation that

perception of novel bitter taste as well as novel yeast taste can promote override of PER

habituation strongly indicates that it is the novelty of the dishabituating stimulus, rather than

its attractiveness, that drives habituation override (Fig 2D). We also tested the effect of

thermogenetic activation of olfactory and visual sensory neurons by expressing TRPA1 in all

Or83b-Gal4 positive olfactory sensory neurons (OSNs)*,* in subsets (*Or85a-Gal4* or *Gr21-Gal4*

positive) of OSNs that mediate strong repulsion, or *Rh1-Gal4* expressing photoreceptors, but

we did not observe an effect on PER habituation (data not shown). Therefore, we did not

explore these stimuli further.

Does novelty-induced override of PER habituation represent a specific effect on the habituated

state or a general sensitization of the sensory response?   To differentiate between these

possibilities, we tested whether and how yeast exposure, *Ir25a*-neuron activation or *Gr66a*-

neuron activation altered PER in naïve flies to a range of sucrose concentrations. If the increase

in PER response in habituated flies were due to sensitization, then we would expect an increase

in PER after the novel stimulus is applied to naïve animals. However, as shown in 3A, 3B and

3C, there is no significant difference in PER response before and after presentation of yeast or

Ir25a or *Gr66a*-GRN activation respectively. Thus, it can be concluded that the reinstatement

of PER response that we observe in our experiments is due to a process that specifically acts

on neural correlates of PER habituation, not broadly on taste perception.

Artificial activation of Tyrosine-Hydroxylase (TH) expressing neurons overrides PER

habituation.

Across species, novel stimuli trigger central release of neuromodulators that confer or enhance

their salience (Ranganath and Rainer, 2003; Kafkas and Montaldi, 2018). In particular, the

activity of dopaminergic neurons has been implicated in the novelty response both in insects

and in mammalian systems, (Hattori et al., 2017; Morrens et al., 2020). We therefore

investigated whether thermogenetic activation of *Drosophila* tyrosine-hydroxylase (TH)

expressing central dopaminergic neurons would be (a) sufficient and (b) necessary for novel-

taste induced override of PER habituation.

In PER-habituated *TH-Gal4>UAS-TRPA1* animals, 1-minute exposure to 32°C to drive

thermogenetic activation of *TH-Gal4* expressing neurons resulted in rapid habituation override

(Fig 4A).   In PER-habituated animals, sucrose-induced PER was close to 12%; in these same

animals, activation of TH expressing neurons increased PER to 60% ($p<0.0001$, F-stat = 50.54,

Dunn's multiple comparisons post-hoc test, ***p = 0.0004).   Significantly, activation of TH

Gal4 neurons had no effect on the innate response to sucrose in naïve flies (Fig 4B).   Thus,

activation of these modulatory neurons causes habituation override, not general taste

sensitization.

TH-Gal4 labels around 200 neurons in the Drosophila brain (Friggi-Grelin et al., 2003).   In

order to more tightly define *TH*-expressing cells involved in PER-habituation override, we

tested two non-overlapping subsets of *TH-Gal4* positive neurons, marked by *TH-D'-Gal4* and

TH-C'-Gal4 drivers (which labels ~54±5 and ~45±3 neurons respectively)   for their potential

roles. While 1-minute thermogenetic activation of *TH-D'* neurons had no effect on PER

habituation, similar activation of neurons labelled by *TH-C'-Gal4* significantly increased PER

response from 28.72% in control habituated flies, to 57.81% after *TH-C'* activation ($p<0.0001$,

F-stat = 69.50, Dunn's multiple comparisons post-hoc test,**p= 0.0042 Fig 4C).   Moreover,

TH-Gal4-driven habituation override required activity in *TH-C'* cell population: thus, 32°C

exposure did not trigger PER habituation override in *TH-C-Gal80; TH-Gal4> UAS TRPA1*

flies*, in which Gal80 expression prevented TRPA1 expression in the *TH-C'* subset of *TH-Gal4*

neurons (Fig 4D). This indicates that activity in *TH-C'* subset of cells is necessary and

sufficient to cause habituation override. To further support this conclusion, we checked and

confirmed that activation of *TH-C'* neurons had no significant effect on the PER of naïve flies

across a range of sucrose concentrations (Fig 4E) confirming that *TH-C'* neurons specifically

influence mechanisms of habituation, rather than general taste perception.

Significantly, a small subset of *TH-C'* neurons (~13±6) are present locally in the SEZ

(suboesophagial zone), an area which not only receives inputs from taste sensory neurons, but

also houses interneurons and motor neurons involved in proboscis extension (Gordon and

Scott, 2009; Kain and Dahanukar, 2015). Thus, some *TH-C'* neurons are well positioned to

mediate sensory-driven novelty signals that influence mechanisms of PER habituation.

A subset of TH-expressing neurons mediate novelty-induced habituation override.

In order to test whether the activity of *TH-Gal4* and *TH-C'-Gal4* neurons is necessary for

novel-taste induced override of PER habituation, we examined whether novel-taste stimulation

could cause habituation override under conditions where synaptic output from *TH-Gal4* or *TH-*

C'-Gal4 neurons was blocked. To achieve this, we expressed the temperature-sensitive,

dominant-negative Shi^{ts1} mutant form of dynamin in these neurons and tested whether a one-

minute exposure to 10% yeast at 34°C could override habituation in *TH-Gal4, UAS-Shi^{ts1}* and

TH-C'-Gal4 UAS-Shi^{ts1} at temperatures restrictive for Shi^{ts1} dynamin function.

At permissive (room) temperatures, *TH-Gal4, UAS-Shi^{ts1}* and *TH-C'-Gal4, UAS-Shi^{ts1}* flies

behaved similarly to wild-type flies, showing both robust PER habituation after 10 minutes of

tarsal sucrose exposure (Fig 5A) and significant PER dishabituation following brief (1 min)

exposure of their labella to 10% yeast solution (Fig 5A). The respective efficiency of PER in

habituated and dishabituated animals (before and after yeast exposure) were 34.05% and

77.29% (p<0.0001, F-stat = 42.47, Dunn's multiple comparisons post-hoc test, ***p = 0.0008)

and 13.46% and 48.46% (p<0.0001, F-stat = 41.01, Dunn's multiple comparisons post-hoc test,

*p = 0.045) respectively (Fig 5A).

In contrast, if after PER habituation the same flies were shifted to and exposed to 10% yeast at

34°C, where the essential function of dynamin in transmitter release would be compromised in

Shi^{ts1}-expressing cells, then habituation override was significantly impaired as compared to

controls (Fig 5A). Thus, yeast-induced dishabituation of control flies not expressing Shi^{ts1} at

34°C was substantially more efficient (p<0.0001, F-stat = 35.83, Dunn's multiple comparisons

post-hoc test,**p= 0.0015), than of *TH-Gal4, UAS-Shi^{ts1}* flies at the same temperature (Fig

5A). Similarly, dishabituation of control flies not expressing Shi^{ts1} was also substantially more

efficient at 34°C (p< 0.0001, F-stat = 46.90, Dunn's multiple comparisons post-hoc test, ***p

= 0.0001), than of *TH-C'-Gal4, UAS-Shi^{ts1}* flies, expressing temperature-sensitive mutant

dynamin in *THC'-Gal4* neurons. These data argue that presynaptic activity in *TH-C'-Gal4* and

TH-Gal4 positive cells is required for yeast-induced override of PER habituation (Fig 5A).

An important caveat to the above conclusion is that dopaminergic neuron activation may be

associated with motivation or reward prediction and therefore be fundamentally required for

high levels of PER. In such a scenario, inactivation of TH-expressing cells would be expected

to generally reduce levels of PER. To address this issue and test whether the apparently

reduced override in Fig 5A is an artifact of blocking dopaminergic neurons during yeast

exposure, we measured the innate response to sucrose in naïve flies with and without blockage

of synaptic transmission in dopaminergic neurons (Fig 5B). There was no significant

difference in the innate response to 2% sucrose, confirming a more restricted role for these TH

positive cells in the override of habituation.

Sensory neurons projections in SEZ overlap spatially with projections of TH-neurons

In the mushroom body, dopaminergic neurons potentially compute novelty by combining

excitatory inputs from olfactory sensory channels with inhibitory inputs driven by familiarity

encoding interneurons (Zhao et al., 2021) (and see Discussion). If TH-positive cells were to

play a similar role in the SEZ, then they would be predicted to receive direct or indirect inputs

from taste sensory neurons as well as input from familiarity-representing inhibitory neurons.

Previous work has shown that processes from some TH-positive neurons are present in the SEZ

which also contains presynaptic endings of taste sensory neurons, processes of local

interneurons and dendrites of motor neurons involved in proboscis extension (Figure 6A; (Liu

et al., 2012)). Therefore, we considered the possibility that taste sensory neurons make contacts

with *TH*-positive processes.

To test whether taste-sensory neurons carrying dishabituating signals form connections, direct

or indirect, with TH-neuron processes in the SEZ, we used the GFP-reconstitution across

synaptic partners (GRASP) technique, which requires the use of dual binary transcription

systems (based on LexA and Gal4 transcription factors) to separately express two

complementing fragments of GFP, one in sensory neurons and the other in TH neurons.

Limited by the immediate availability of transgenes, we were technically restricted to analyzing

TH connections with neurons in which gene expression could be controlled by LexA. This

was possible for bitter-taste responsive *Gr66a* neurons but not for the yeast-responsive *Ir25a*

class.

We first examined whether *Gr66a* axonal projections and *TH-Gal4* marked processes were

present in close proximity within the SEZ, by examining the relative localization of GFP driven

in *Gr66a*-expressing neurons with RFP in *TH*-positive cells. Processes of sensory neurons

expressing *Gr66a-LexA*-driven GFP and dopaminergic neurons expressing *TH-Gal4* driven

RFP showed close proximity (Fig6A). Further, GRASP experiments showed that two halves

of split GFP, one expressed in *Gr66a* neurons and the other in *TH*-positive cells could combine

to reconstitute GFP fluorescence within the SEZ (Fig 6B). Robust fluorescence reconstitution

was seen in 8/13 experimental animals compared to 0/15 total animals expressing only one

split-GFP component. Thus, GRASP experiments confirm that the processes of *Gr66a* axons

and tyrosine-hydroxylase expressing cells come in close proximity of each other. However,

due to the limitation of the GRASP technique used it cannot be established certainly if there is

a direct synapse between the two cell types.

These data predict that habituation override induced by thermogenetic activation of Gr66a cells

should require activity in *TH-Gal4* positive cells. We tested this prediction by creating and

analysing PER habituation and dishabituation in *Gr66a-LexA/ LexAop-TRPA1; TH-Gal4/*

UAS-Shi^{ts1} and control *Gr66a-LexA/LexAop-TRPA1; TH-Gal4/+* lines at temperatures

permissive and restrictive for Shi^{ts1} dynamin. As shown (Fig 6C), both lines showed robust

PER and PER habituation. One minute-exposure to 34°C in control flies permissive for

synaptic transmission from *TH*-positive neurons resulted in significant (p<0.0001, F-stat =

44.02, Dunn's multiple comparisons post-hoc test, *p = 0.02) override of habituation.

However, similar 34°C thermogenetic stimulation of *Gr66a* neurons in experimental flies

where synaptic transmission from TH-positive neurons is blocked did not result in override of

PER habituation. Together the anatomical and behavioral data indicate that *Gr66a*-expressing

bitter taste sensory neurons are functionally connected to *TH*-expression modulatory neurons

whose activity is required for PER habituation override.

Control experiments demonstrating that blocking synaptic output from *TH*-expressing cells

has no effect on basal PER response in naïve animals further confirm that transmitter release

from *TH*-neurons is required for override of neuronal mechanisms of habituation, not for the

innate response to sucrose (Fig 5B).

DISCUSSION

Animal behavior is profoundly flexible. Thus, at any time, multiple potential behavioral programmes remain dormant, while a subset relevant to specific contexts are active. A growing body of evidence suggests that inhibitory inputs play a major role in preserving perceptions, behaviors and memories in dormant form until required (Barron et al., 2017). Thus, a given context, by recruiting disinhibitory circuits may override inhibition to release latent perceptions, motor programmes and memories appropriate to that context. While the overarching principles are increasingly appreciated, there is still limited understanding of how these are implemented in cells and circuits. While there are many potential reasons for this, the difficulty in studying mechanisms of override has been substantially caused by the paucity of systems in which both mechanisms of habituation or cognitive silencing can be addressed as well as where robust override can be experimentally achieved.

The results we present outline essential elements within a *Drosophila* circuit that overrides habituation, in this case of the sucrose-evoked proboscis extension reflex. In doing so, they connect sensory neurons mediating override to neuromodulatory neurons projecting to the SEZ, which houses interneurons and motor neurons that control proboscis extension. When taken together with published work showing that increased inhibition underlies PER habituation, these findings are significant because they circumscribe a central circuit for habituation override and add to the growing literature on the role and mechanism of disinhibition in gating animal cognition.

Previous work in *Drosophila* concluded that PER habituation arises from increased sucrose-evoked inhibition onto neurons that drive proboscis extension (Paranjpe et al., 2012). Two findings, which closely mirror observations on olfactory habituation, provided key support for this conclusion (Das et al., 2011; Paranjpe et al., 2012). First, the *rutabaga*-encoded adenyl cyclase is required specifically in inhibitory neurons for PER habituation, an observation that

we have independently confirmed in this study (Figure 1). Second, experimental silencing of

inhibitory neurons causes override of habituation. Together these observations indicate first,

that increased GABAergic activity is required for the expression of PER habituation and

second, that disinhibition could serve as strategy for habituation override. How might such

disinhibition be biologically achieved? Our current experiments show that novel sensory

experience induces override of PER habituation through a pathway that requires activity in the

TH-C' class of dopaminergic neurons. As thermogenetic activation of *TH-C'* cells is also

sufficient to override PER habituation, the data suggest a framework in which sensory stimuli

activate *TH-C'* neurons, which directly or indirectly, inhibit GABAergic neurons responsible

for habituation.

An important question to address is why novel but not familiar taste stimuli are effective for

override? In physiological terms, how could activation of a novel subset of (*Ir25a or Gr66a*)

sensory-neurons result in strong excitation of *TH-C'* cells, while similar levels of activation of

sucrose-response sensory-neurons causes weaker excitation of the same *TH-C'* cells? We

suggest that this occurs because familiar stimuli additionally recruit higher levels of inhibition

onto *TH-C'* neurons. At a circuit level, there are multiple ways in which such a mechanism

could be implemented, the simplest of which is shown schematically in Figure 7.

The model proposes that in the naïve state, sensory neurons (SN) excite excitatory projection

neurons (PN) that activate the PER. These SNs also excite cognate inhibitory local

interneurons (iLNs) and *TH-C'* neurons (either directly or indirectly via yet to be identified

neuron). The iLNs are postulated to make weak inhibitory synapses onto PNs and *TH-C'* cells,

while also receiving strong inhibition from *TH-C'*. Thus, novel stimuli result in weak total iLN

excitation, strong PN excitation and robust PER. The imposition of a synaptic learning rule

that coincident presynaptic and postsynaptic activity at inhibitory synapses leads to inhibitory

synapse potentiation, generates the postulated habituated state, wherein PNs receive additional

increased inhibition resulting in lower net excitation and reduced PER. At the same time the

same learning rule also results in increased inhibition and therefore decreased net excitation of

TH-C' neurons by "familiar" SNs. In this habituated state, the familiar tastant, causing only

weak *TH-C'* excitation, would not disinhibit the PER pathway. However, a novel tastant would

still trigger strong *TH-C'* cell activation, effective disinhibition of PN and thereby override of

PER habituation. The above model places some constraints on SN-LN connectivity which have

yet to be examined or established: most significantly suggesting that different classes of SNs

activate different subgroups of iLNs in the SEZ.

While our experiments lead to a comprehensive model for a habituation override circuit, they

do not yet clarify several important issues. Most obviously many aspects of the proposed

connectivity are inferred, rather than demonstrated by direct anatomical or physiological

methods. The exact subset of *TH-C'* cells involved as well as the subtypes and connectivities

of relevant inhibitory neurons in the SEZ remain unknown and crucially important to establish.

The mechanism by which relevant *TH-C'* cells function has not been formally proven. While

these cells are marked by three independent dopaminergic reporter lines, *TH-Gal4*, *TH-C'* and

TH-C-Gal80, and therefore probably dopaminergic, it remains unclear whether dopamine

release is required for habituation override.   Our attempts to address this via knockdown of

tyrosine hydroxylase through RNAi, or by various genetic manipulations of dopamine receptor

expression did not yield definitive results (Materials and Methods), often because these

manipulations affected baseline levels of PER and PER habituation.

However, given that the SEZ, which contains gustatory neurons axons as well as dendrites of

motor neurons that drive PER, is likely to be numerically simple compared, the most

parsimonious model for override would posit that taste sensory neurons trigger direct excitation

of dopaminergic processes, which in turn acts within the SEZ to directly inhibit GABAergic

cells (Pimentel et al., 2016) whose potentiation drives PER habituation.

The model above is consistent with observations on habituation override in mouse and Aplysia brains (Bristol and Carew, 2005; Smith et al., 2009; Kato et al., 2015; Ogg et al., 2018). In mouse, long term auditory habituation to a passively experienced tone is accompanied by increased activity in tone-responsive SOM+ neurons that inhibit similarly tuned pyramidal cells in the auditory cortex. However, if habituated mice are coaxed to attend to the same tone (by a reward for successful engagement), then the behaving mice show overriding inhibition of SOM+ neurons and increased activity of downstream L2/3 pyramidal neurons (Kato et al., 2015). It appears likely that this disinhibition is accomplished by modulatory inputs onto upstream VIP+ neurons. A more recent analysis showed that cholinergic inputs into the mouse olfactory bulb could cause override of a fast form of olfactory habituation. Thus, electrical or optogenetically induced acetylcholine release into the bulb caused mice to override habituation and investigate a previously ignored odor (Ogg et al., 2018). While sensitization is not formally excluded here, these studies are consistent with an emerging theme wherein neuromodulators released in response to novel or meaningful stimuli (Vankov et al., 1995; Giovannini et al., 2001; Ranganath and Rainer, 2003; Hattori et al., 2017; Kafkas and Montaldi, 2018; Morrens et al., 2020) result in disinhibition which can either enhance learning or override habituation.

The experimental results described here provides multiple lines of circumstantial evidence in support of a novelty-induced dopaminergic pathway for disinhibition of sensory perception. It outlines a habituation override circuit all the way from sensory neurons that detect stimulus, to motor neurons that mediate behavioral response. In context of the increasingly widely appreciated role for disinhibition in the control of perception, cognition and behavior (Letzkus et al., 2015; Sridharan and Knudsen, 2015; Barron et al., 2017; Wang and Yang, 2018) we suggest that this work provides a valuable intellectual and biological foundation for future studies to comprehensively identify neurons and mechanisms involved in a central pathway for behaviorally important disinhibition.

ACKNOWLEDGMENTS

We thank Ali Asgar Bohra, Camilla Roselli, Tamara Boto and James Cooke for comments on the manuscript, Frederic Marion-Poll and members of Ramaswami and VijayRaghavan labs for advice, support and useful discussions. We thank Carlos Ribeiro for valuable discussions and reagents, Pushkar Paranjpe for help setting up the PER assay, as well as the Bloomington stock centre and our many colleagues (mentioned in the text) for various Drosophila stocks used. The work was supported by a Wellcome Trust Investigator Award and a Science Foundation Ireland Investigator Programme Grant to MR, by core support from the National Centre for Biological Sciences (Tata Institute of Fundamental Research) to KVR and by a CSIR postgraduate fellowship and a Biocon-Trinity Scholarship to ST.

References

Asztalos Z, Arora N, Tully T (2007a) Olfactory jump reflex habituation in Drosophila and

effects of classical conditioning mutations. Journal of neurogenetics 21:1-18.

Asztalos Z, Baba K, Yamamoto D, Tully T (2007b) The fickle mutation of a cytoplasmic

tyrosine kinase effects sensitization but not dishabituation in Drosophila

melanogaster. Journal of neurogenetics 21:59-71.

Barron HC, Vogels TP, Behrens TE, Ramaswami M (2017) Inhibitory engrams in perception

and memory. Proceedings of the National Academy of Sciences of the United States

of America 114:6666-6674.

Bristol AS, Carew TJ (2005) Differential role of inhibition in habituation of two independent

afferent pathways to a common motor output. Learning & memory 12:52-60.

Castellucci V, Pinsker H, Kupfermann I, Kandel ER (1970) Neuronal mechanisms of

habituation and dishabituation of the gill-withdrawal reflex in Aplysia. Science

167:1745-1748.

Cooke SF and Ramaswami M (2020) 17 Ignoring the Innocuous: Neural. The Cognitive

Neurosciences:197.

Das S, Sadanandappa MK, Dervan A, Larkin A, Lee JA, Sudhakaran IP, Priya R, Heidari R,

Holohan EE, Pimentel A, Gandhi A, Ito K, Sanyal S, Wang JW, Rodrigues V,

Ramaswami M (2011) Plasticity of local GABAergic interneurons drives olfactory

habituation. Proceedings of the National Academy of Sciences of the United States of

America 108:E646-654.

Dethier VG (1976) The hungry fly: A physiological study of the behavior associated with

feeding. Oxford, England: Harvard U Press.

Duerr JS, Quinn WG (1982) Three Drosophila mutations that block associative learning also

affect habituation and sensitization. Proceedings of the National Academy of Sciences

of the United States of America 79:3646-3650.

Engel JE, Wu CF (2009) Neurogenetic approaches to habituation and dishabituation in

Drosophila. Neurobiol Learn Mem 92:166-175.

Fischer TM, Blazis DE, Priver NA, Carew TJ (1997) Metaplasticity at identified inhibitory

synapses in Aplysia. Nature 389:860-865.

Fischler W, Kong P, Marella S, Scott K (2007) The detection of carbonation by the

Drosophila gustatory system. Nature 448:1054-1057.

Flood TF, Iguchi S, Gorczyca M, White B, Ito K, Yoshihara M (2013) A single pair of

interneurons commands the Drosophila feeding motor program. Nature 499:83-87.

Fois C, Medioni J, Le Bourg E (1991) Habituation of the proboscis extension response as a

function of age in Drosophila melanogaster. Gerontology 37:187-192.

Friggi-Grelin F, Coulom H, Meller M, Gomez D, Hirsh J, Birman S (2003) Targeted gene

expression in Drosophila dopaminergic cells using regulatory sequences from tyrosine

hydroxylase. Journal of neurobiology 54:618-627.

Ganguly A, Pang L, Duong VK, Lee A, Schoniger H, Varady E, Dahanukar A (2017) A

Molecular and Cellular Context-Dependent Role for Ir76b in Detection of Amino

Acid Taste. Cell reports 18:737-750.

Giovannini MG, Rakovska A, Benton RS, Pazzagli M, Bianchi L, Pepeu G (2001) Effects of

novelty and habituation on acetylcholine, GABA, and glutamate release from the

frontal cortex and hippocampus of freely moving rats. Neuroscience 106:43-53.

Glanzman DL, Mackey SL, Hawkins RD, Dyke AM, Lloyd PE, Kandel ER (1989) Depletion

of serotonin in the nervous system of Aplysia reduces the behavioral enhancement of

gill withdrawal as well as the heterosynaptic facilitation produced by tail shock. The

Journal of neuroscience : the official journal of the Society for Neuroscience 9:4200-

4213.

Gordon MD, Scott K (2009) Motor control in a Drosophila taste circuit. Neuron 61:373-384.

Hamada FN, Rosenzweig M, Kang K, Pulver SR, Ghezzi A, Jegla TJ, Garrity PA (2008) An

internal thermal sensor controlling temperature preference in Drosophila. Nature

454:217-220.

Hattori D, Aso Y, Swartz KJ, Rubin GM, Abbott LF, Axel R (2017) Representations of

Novelty and Familiarity in a Mushroom Body Compartment. Cell 169:956-969 e917.

Hawkins RD, Cohen TE, Greene W, Kandel ER (1998) Relationships between

dishabituation, sensitization, and inhibition of the gill- and siphon-withdrawal reflex

in Aplysia californica: effects of response measure, test time, and training stimulus.

Behavioral neuroscience 112:24-38.

Kafkas A, Montaldi D (2018) How do memory systems detect and respond to novelty?

Neuroscience letters 680:60-68.

Kain P, Dahanukar A (2015) Secondary taste neurons that convey sweet taste and starvation

in the Drosophila brain. Neuron 85:819-832.

Kato HK, Gillet SN, Isaacson JS (2015) Flexible Sensory Representations in Auditory Cortex

Driven by Behavioral Relevance. Neuron 88:1027-1039.

Le Bourg E (1983) Aging and habituation of the tarsal response in Drosophila melanogaster.

Gerontology 29:388-393.

Letzkus JJ, Wolff SB, Luthi A (2015) Disinhibition, a Circuit Mechanism for Associative

Learning and Memory. Neuron 88:264-276.

Liu C, Placais PY, Yamagata N, Pfeiffer BD, Aso Y, Friedrich AB, Siwanowicz I, Rubin

GM, Preat T, Tanimoto H (2012) A subset of dopamine neurons signals reward for

odour memory in Drosophila. Nature 488:512-516.

Minnich DE (1921) An experimental study of the tarsal chemoreceptors of two nymphalid

butterflies. Journal of Experimental Zoology 33:172-203.

Montell C (2009) A taste of the Drosophila gustatory receptors. Current opinion in

neurobiology 19:345-353.

Morrens J, Aydin C, Janse van Rensburg A, Esquivelzeta Rabell J, Haesler S (2020) Cue-

Evoked Dopamine Promotes Conditioned Responding during Learning. Neuron

106:142-153 e147.

Ogg MC, Ross JM, Bendahmane M, Fletcher ML (2018) Olfactory bulb acetylcholine release

dishabituates odor responses and reinstates odor investigation. Nature

communications 9:1868.

Paranjpe P, Rodrigues V, VijayRaghavan K, Ramaswami M (2012) Gustatory habituation in

Drosophila relies on rutabaga (adenylate cyclase)-dependent plasticity of GABAergic

inhibitory neurons. Learning & memory 19:627-635.

Pimentel D, Donlea JM, Talbot CB, Song SM, Thurston AJF, Miesenbock G (2016)

Operation of a homeostatic sleep switch. Nature 536:333-337.

Ramaswami M (2014) Network plasticity in adaptive filtering and behavioral habituation.

Neuron 82:1216-1229.

Ranganath C, Rainer G (2003) Neural mechanisms for detecting and remembering novel

events. Nature reviews Neuroscience 4:193-202.

Rankin CH, Abrams T, Barry RJ, Bhatnagar S, Clayton DF, Colombo J, Coppola G, Geyer

MA, Glanzman DL, Marsland S, McSweeney FK, Wilson DA, Wu CF, Thompson

RF (2009) Habituation revisited: an updated and revised description of the behavioral

characteristics of habituation. Neurobiology of learning and memory 92:135-138.

Semelidou O, Acevedo SF, Skoulakis EM (2018) Temporally specific engagement of distinct

neuronal circuits regulating olfactory habituation in Drosophila. eLife 7.

Shen Y, Dasgupta S, Navlakha S (2020) Habituation as a neural algorithm for online odor

discrimination. Proceedings of the National Academy of Sciences of the United States

of America 117:12402-12410.

Smith JJ, Shionoya K, Sullivan RM, Wilson DA (2009) Auditory stimulation dishabituates

olfactory responses via noradrenergic cortical modulation. Neural plasticity

2009:754014.

Sridharan D, Knudsen EI (2015) Selective disinhibition: A unified neural mechanism for

predictive and post hoc attentional selection. Vision research 116:194-209.

Steck K, Walker SJ, Itskov PM, Baltazar C, Moreira JM, Ribeiro C (2018) Internal amino

acid state modulates yeast taste neurons to support protein homeostasis in Drosophila.

eLife 7.

Stocker RF (1994) The organization of the chemosensory system in Drosophila

melanogaster: a review. Cell and tissue research 275:3-26.

Thompson RF, Spencer WA (1966) Habituation: a model phenomenon for the study of

neuronal substrates of behavior. Psychological review 73:16-43.

Vankov A, Herve-Minvielle A, Sara SJ (1995) Response to novelty and its rapid habituation

in locus coeruleus neurons of the freely exploring rat. The European journal of

neuroscience 7:1180-1187.

Wang X-J, Yang GR (2018) A disinhibitory circuit motif and flexible information routing in

the brain. Current opinion in neurobiology 49:75-83.

Wisotsky Z, Medina A, Freeman E, Dahanukar A (2011) Evolutionary differences in food

preference rely on Gr64e, a receptor for glycerol. Nature neuroscience 14:1534-1541.

Zaccardi ML, Traina G, Cataldo E, Brunelli M (2004) Sensitization and dishabituation of

swim induction in the leech Hirudo medicinalis: role of serotonin and cyclic AMP.

Behavioural brain research 153:317-326.

Zhao C, Widmer YF, Diegelmann S, Petrovici MA, Sprecher SG, Senn W (2021) Predictive

olfactory learning in Drosophila. Scientific reports 11:6795.

FIGURE LEGENDS

Figure 1. Gustatory habituation of PER to sweet taste.

(**A**) Behavioural protocol of PER habituation to tarsal stimulation following sucrose exposure.

(**B**) Tarsal exposure to 10% sucrose for 10 minutes leads to decrease in PER response (***p<

0.0001, Wilcoxon signed rank test) whereas exposure to water does not affect the response (p

= 0.6133, Wilcoxon signed rank test). (**C**) As previously described (Paranjpe et al, 2012),

gustatory habituation is dependent on the Rutabaga adenylate cyclase since it is impaired in

rut^{2080} mutants. Habituation can be restored by expressing wild type *rut* in inhibitory *Gad1-*

Gal4 neurons (**p < 0.006, Mann Whitney U stat = 227). (**D**) The habituated response can be

overridden by a novel mechanical stimulus like vortexing (p<0.001, F- stat = 47.08, Dunn's

multiple comparison post-hoc test, ***p <0.001, *p = 0.01). (**E**) Vortexing does not have an

effect on naïve response to sucrose at concentrations 0.1% (p = 0.0518, Wilcoxon signed rank

test), 0.5% (p= 0.6880, Wilcoxon signed rank test) and 1 % (p= 0.1626, Wilcoxon signed rank

test) as, determined by testing PER response pre and post vortexing at different sucrose

concentrations. Therefore, the enhanced PER response after vortexing habituated animals is

not a result of general sensitization but instead represents a specific override of habituation to

sucrose. Bars represent mean±SEM, triangles represent individual data points, ns represents

statistically not significant, p>0.05.

Figure 2. Habituation override mediated by novel stimulus

(**A**) Behaviour protocol of PER habituation and its override by presenting stimulus to the

labellum (**B**) Presentation of novel 10% yeast stimulus to labellum restores PER response

(p<0.0001, F-stat = 31.15, Dunn's multiple comparisons post-hoc test, ***p<0.0001, *p =

0.014) whereas 10% sucrose, which is a familiar stimulus, does not have an effect on habituated

response (p<0.0001, F-stat = 43.49, Dunn's multiple comparisons post-hoc test, ***p<0.0001,

p > 0.99). **(C1)** Expression pattern of *Ir25a-Gal4* in the adult brain. Ir25a-Gal4 is expresssed

in gustatory receptor neurons, predominantly bitter and sweet sensory neurons in the

subesophageal zone (SEZ), as well as some olfactory receptor neurons in the antennal lobe

(AL). **(C2)** Directly activating yeast responsive *Ir25a-Gal4* cells expressing heat activated

channel *TRPA1* for 1 minute after habituation at 32°C, is sufficient to override PER habituation

to sucrose (p<0.0001, F-stat = 38.66, Dunn's multiple comparisons post-hoc test, ***p<0.0001,

**p = 0.0012) whereas at permissive temperature, *Ir25a>TRPA1* flies do not show an increase

in habituated response (p<0.0001, F-stat = 22.62, Dunn's multiple comparisons post hoc test,

***p = 0.0002, p>0.99) (*Ir25a-Gal4/+* p<0.0001, F-statistic = 38.99, Dunn's multiple

comparison post-hoc test, ***p<0.0001, p > 0.99, *UAS-TRPA1/+*, p<0.0001, F-stat = 33.88,

Dunn's multiple comparisons post-hoc test, ***p<0.0001, p > 0.99. **(D1)** Expression pattern of

Gr66a-Gal4 in the subesophageal zone (SEZ) region of adult brain. The bitter sensory neurons

marked by Gr66a form a ringed structure; they project to the anterior region of the SEZ. **(D2)**

Thermogenetic activation of bitter *Gr66a*-marked GRNs for 1 minute after habituation at 32°C,

also overrides sweet-taste habituation (p<0.0001, F-stat = 39.85, Dunn's multiple comparisons

post-hoc test, ***p<0.0001, **p = 0.0042) whereas no difference is observed in controls

(Gr66a>UAS-TRPA1 at RT p<0.0001, F-stat = 17.56, Dunn's multiple comparisons post hoc

test, ***p= 0.0009, p>0.99, *Gr66a-Gal4/+* p<0.0001, F-stat = 44.60, Dunn's multiple

comparisons post-hoc test, ***p<0.0001, p = 0.97, *UAS-TRPA1/+*, p<0.0001, F-stat = 40.22,

Dunn's multiple comparison post-hoc test, p<0.0001, p > 0.99). Together the data show that

stimulus-novelty drives override of habituation. Bars represent mean±SEM, triangles represent

individual data points, ns represents statistically not significant, p>0.05.

Figure 3. Novelty induced habituation override does not induce sensitization of sensory

response

(A) Labellar exposure to 10% yeast does not increase the naïve PER response to 0.1%, 0.5%, 1% or 2% sucrose stimulation (Probabilities of PER being unchanged using the Wilcoxon sign rank test: $p = 0.6404$, $p = 0.6731$, $p = 0.1957$, and $p = 0.1208$ at the 4 respective sucrose concentrations). **(B)** Thermogenetic activation of *Ir25a-Gal4* does not sensitize response to sucrose tested at concentration 0.1% ($p > 0.99$, Wilcoxon sign rank test), 0.5% ($p = 0.5938$, Wilcoxon sign rank test), 1% ($p = 0.7588$, Wilcoxon sign rank test), 2% ($p = 0.4814$, Wilcoxon sign rank test,). **(C)** Thermogenetic activation of *Gr66a-Gal4* does not have an effect on response to 0.1% ($p = 0.5$, Wilcoxon sign rank test), 0.5% ($p > 0.99$, Wilcoxon sign rank test), 1% ($p = 0.375$, Wilcoxon sign rank test), 2% ($p = 0.3828$, Wilcoxon sign rank test) concentrations of sucrose. Points represent mean±SEM, ns represents statistically not significant, $p > 0.05$.

Figure 4. Activation of TH expressing neurons overrides habituation

(A) Thermogenetic activation of TH expressing neurons for 1 minute at 32°C after habituation results in habituation override ($p < 0.0001$, F-stat = 50.54, Dunn's multiple comparisons post-hoc test, ***$p < 0.0001$, ***$p = 0.0004$) (*TH-Gal4/+*, $p < 0.0001$, F-stat = 49.56, Dunn's multiple comparisons post-hoc test, ***$p < 0.0001$, $p > 0.99$, *UAS-TRPA1/+*, $p < 0.0001$, F-stat = 37.31, Dunn's multiple comparisons post-hoc test, ***$p < 0.0001$, $p > 0.99$, *TH-Gal4/UAS-TRPA1* temperature control, $p < 0.0001$, F-stat = 26, Dunn's multiple comparisons post-hoc test, ***$p < 0.0001$, $p = 0.8665$). **(B)** Activation of TH expressing neurons does not lead to sensitization of gustatory response tested at 0.1% ($p = 0.09$, Wilcoxon sign rank test), 0.5% ($p = 0.1$, Wilcoxon sign rank test), 1% ($p = 0.14$, Wilcoxon sign rank test), 2% ($p = 0.1367$, Wilcoxon sign rank test). **(C)** Activation of a subset of TH expressing cells marked by *TH-C'-Gal4* for 1 minute after habation at 32°C is sufficient to override habituated response to sucrose ($p < 0.0001$, F-stat = 69.50, Dunn's multiple comparisons post-hoc test, ***$p < 0.0001$,

**p = 0.0042) (*TH Gal4/+*, p<0.0001, F-stat = 59.70, Dunn's multiple comparisons post-hoc

test, ***p<0.0001, p = 0.4914 *, UAS-TRPA1/+*, p<0.0001, F-stat = 63.52**,** Dunn's multiple

comparisons post-hoc test, ***p<0.0001, p > 0.99) (*TH-C'- Gal4* temperature control,

p<0.0001, F-stat = 21.41, Dunn's multiple comparisons post-hoc test, **p = 0.0017, p > 0.99).

(D) Activation of *TH-D'* subset of neurons, which marks subset of neurons that does not

overlap with *TH-C,'* for 1 minute after habituation at 32°C does not override habituation

(p<0.0001, F-stat = 42.70, Dunn's multiple comparisons post-hoc test, ***p <0.0001, p> 0.99)

(*UAS-TRPA1/+*, p<0.0001, F-stat = 76.79, Dunn's multiple comparisons post-hoc test,

***p<0.0001, p = 0.3710). **(E)** Combining *TH-C-Gal80* along with *TH Gal4* blocks the

expression in *TH-C'* subset of neurons only, as observed by GFP expression (SEZ represents

subesophageal zone). **(F)** The flies carrying *TH-C-Gal80* along with *TH-Gal4* fail to show

habituation override when these are activated at 32°C after habituation further confirming the

role of *TH-C'* subset of neurons (p<0.0001, F-stat = 35.74, Dunn's multiple comparisons post-

hoc test, ***p<0.0001, p >0.99) (*TH-Gal4/ UAS-TRPA1*, p<0.0001, F-stat = 45.65, Dunn's

multiple comparisons post-hoc test, ***p<0.0001, **p = 0.0025) (F-stat = 35.38**,** *TH-C-*

Gal80/+, p <0.0001, F-stat = 35.38, Dunn's multiple comparisons post-hoc test, ***p<0.0001,

p = 0.9061) (*UAS-TRPA1/+*, p<0.0001, F-stat = 30.55, Dunn's multiple comparisons post-hoc

test, ***p<0.0001, p > 0.99). **(G)** Activation of *TH-C'* Gal4 neurons does not affect the naïve

response to sucrose tested at 0.1% (p = 0.5635, Wilcoxon sign rank test), 0.5% (p = 0.3748,

Wilcoxon sign rank test), 1% (p = 0.7338, Wilcoxon sign rank test), 2% (p = 0.3125, Wilcoxon

sign rank test). Bars represent mean ±SEM, triangles represent individual data points, points in

figure 4B and 4G represent mean ±SEM, ns represents statistically not significant, p>0.05.

Figure 5. Activity of *TH-C'* subset of neurons is necessary for novelty induced override

(A) At permissive temperature (RT = 21°C), presentation of 10% yeast to the labellum

overrides habituation in both *TH-C' Gal4/ UAS-Shi^{s1}* (p<0.0001, F-stat = 41.01, Dunn's

multiple comparisons post-hoc test,***p<0.0001, *p = 0.0457) and *TH-Gal4/+;UAS-Shi^{ts1}/+*

flies and (p<0.0001, F-stat = 42.47, Dunn's multiple comparisons post-hoc test, ***p<0.0001,

***p = 0.0008) respectively. However, at restrictive temperature (34°C) blocking the synaptic

transmission of *TH-Gal4* and *TH-C'-Gal4* neurons by expressing *Shi^{ts1}* during presentation of

novel yeast stimulus after habituation to sucrose, impairs habituation override (*TH-*

Gal4/+;UAS-Shi^{ts1}/+, p<0.0001, F-stat = 102.8, Dunn's multiple comparisons post-hoc test, ,

***p<0.0001, p = 0.3897) (*TH-C'-Gal4/ UAS-Shi^{ts1}*, p<0.0001, F-stat = 37.98, Dunn's multiple

comparisons post-hoc test, ***p<0.0001, p > 0.99) respectively. The genotypic controls show

override of habituation to 10% yeast presented to the labellum (*TH-Gal4/+*, p<0.0001, F-stat

= 35.83, Dunn's multiple comparisons post-hoc test, ***p<0.0001, **p = 0.0015) (*UAS-*

Shi^{ts1}/+, p<0.0001, F-stat = 56.96, Dunn's multiple comparisons post-hoc test, p<0.0001, ***p

= 0.0009), (*TH-C'-Gal4/+*, p<0.0001, F-stat = 46.90, Dunn's multiple comparisons post-hoc

test, ***p<0.0001, ***p = 0.0001) **(B)** Blocking *TH-Gal4* neurons and *TH-C'-Gal4* neurons

does not have an effect on PER itself (p = 0.1909 and p = 0.9141, respectively, Wilcoxon sign

rank). Bars represent mean ± SEM, triangles represent individual data points, ns represents

statistically not significantly, p>0.05.

Figure 6. TH Gal4 neurons are functionally connected to GRNs

(A) Expression pattern of Gr66a with respect to TH expressing neurons. **(B)** GRASP between

Gr66a and TH expressing neurons show GFP signal in animals expressing both the components

of split GFP. No signal is observed in controls lacking either of split GFP component. **(C)**

Controls do not show any significant increase in response at 34°C (*Gr66a LexA/ +; TH-Gal4/+*,

p<0.0001, F-statistic = 55.06, Dunn's multiple comparisons post-hoc test, ***p<0.0001, p=

0.9079, *LexAop-TRPA1/+; UAS-Shi^{ts1}/+*, p<0.0001, F-stat = 53.41, Dunn's multiple

comparisons post-hoc test, ***p<0.0001, p = 0.0580,). Activation of Gr66a while inhibiting

TH expressing neurons simultaneously fails to reinstate habituated response (*Gr66a LexA/*

LexAop-TRPA1; TH-Gal4/ UAS-Shi^{ts1}, p<0.0001, F-stat = 52.57, Dunn's multiple comparisons

post-hoc test, ***p<0.0001, p= 0.9494), whereas activating Gr66a solely overrides habituation

(*Gr66a LexA/ LexAop-TRPA1; TH-Gal4/ +*, p<0.0001, F-stat = 44.02, Dunn's multiple

comparisons post-hoc test, ***p<0.0001, *p = 0.0205). Control at permissive temperature does

not show any difference in habituated response (*Gr66a LexA/ LexAop-TRPA1; TH-Gal4/ +* ,

***p= 0.0004, F-stat = 15.50, *p= 0.0322, p > 0.99, Dunn's multiple comparisons post-hoc

test). Bars represent mean ± SEM, triangles represent individual data points, ns represents

statistically not significantly, p>0.05, SEZ represents subesophageal zone.

Figure 7: Circuit model for PER habituation override.

Model for habituation override in gustatory system of *Drosophila.* Green represents excitatory

neurons and connections; Red represents inhibitory neurons and connections; Cyan represents

dopamine/ neuromodulatory neurons. Arrows show the direction of activity/ signal. Dotted

lines indicate hypothetical connections, in particular between iLNs and THC' neurons, that

need to be experimental demonstrated. As shown in Paranjpe et al, 2012, plasticity of

GABAergic neurons underlies habituation. The model proposes that during habituation

override, excitation of sensory neurons responsive to novel stimulus activates *TH-C'* subset of

dopaminergic neurons either directly or indirectly via a probable intermediate neuron. As

there is weak inhibition mediated by local inhibitory interneurons (iLNs) onto *TH-C'* neurons

in this pathway, these dopaminergic neurons exhibit strong excitation and in turn inhibit iLNs

that are potentiated during habituation. As a result, disinhibition leads to strong excitation of

projection neurons (PNs) and therefore, strong PER in response to familiar stimulus. In

contrast, excitation of sensory neurons responsive to familiar stimulus activates *TH-C'* neurons

but due to strong iLN inhibition onto *TH-C'* neurons resulting from habituation in this pathway,

dopaminergic neurons fail to disinhibit. This leads to weak PN excitation and therefore, weak

PER in response to the familiar stimulus. Several aspects of the model remain speculative.

762
763
764
765

Figure1:

Figure2:

Figure3:

Unhabituated Flies

— Pre activation   — Post activation

Figure 4:

Figure 5:

A

B

Figure 6:

A

Gr66a-LexA; TH-Gal4 > UAS-mCD8:: RFP, LexAop-mCD8::GFP

B

C

Figure 7: