mTOR activation induces endolysosomal remodeling and unconventional secretion of IL-32 via

exosomes in inflammatory reactive astrocytes

Kun Leng[1,2,3]*+, Brendan Rooney[1+], Frank McCarthy[4], Wenlong Xia[5], Indigo V.L. Rose[1,6,10],

Sophie Bax[4], Marcus Chin[7,8], Saeed Fathi[3], Kari A. Herrington[9], Manuel Leonetti[4], Aimee Kao[7],

Stephen P.J. Fancy[5], Joshua E. Elias[4], Martin Kampmann[4,10,11]*

[1]Institute for Neurodegenerative Diseases, University of California, San Francisco, San

Francisco, CA, USA

[2]Biomedical Sciences Graduate Program, University of California, San Francisco, San Francisco,

CA, USA

[3]Medical Scientist Training Program, University of California, San Francisco, San Francisco,

CA, USA

[4] Chan Zuckerberg Biohub, San Francisco, CA, USA

[5]Departments of Neurology and Pediatrics, School of Medicine, University of California, San

Francisco

[6]Neuroscience Graduate Program, University of California, San Francisco, San Francisco, CA,

USA

[7]Memory and Aging Center, Department of Neurology, University of California, San Francisco,

San Francisco, CA, USA

[8]Small Molecule Discovery Center, Department of Pharmaceutical Chemistry, University of

California, San Francisco, San Francisco, CA, USA

[9]Center for Advanced Microscopy, University of California, San Francisco, San Francisco, CA,

USA

[10]Weill Institute for Neurosciences, University of California, San Francisco, San Francisco, CA,

USA

[11]Department of Biochemistry and Biophysics, University of California, San Francisco, San

Francisco, CA, USA

*e-mail: kun.leng@ucsf.edu, martin.kampmann@ucsf.edu

+ These authors contributed equally

KEYWORDS

Astrocytes, inflammatory reactive astrocytes, neuroinflammation, mTOR, endolysosomal

system, IL-32, extracellular vesicles

LIST OF ABBREVIATIONS

CRISPR: clustered regularly interspaced short palindromic repeats

CRISPRi: CRISPR-based gene expression inhibition

hiPSC: human induced pluripotent stem cell

ITC: IL-1α+TNF+C1q

ELISA: enzyme-linked immunosorbent assay

EV: extracellular vesicle

EAE: experimental autoimmune encephalitis

ABSTRACT

Astrocytes respond and contribute to neuroinflammation by adopting inflammatory reactive

states. Although recent efforts have characterized the gene expression signatures associated with

these reactive states, the cell biology underlying inflammatory reactive astrocyte phenotypes

remains under-explored. Here, we used CRISPR-based screening in human iPSC-derived

astrocytes to identify mTOR activation a driver of cytokine-induced endolysosomal system

remodeling, manifesting as alkalinization of endolysosomal compartments, decreased

autophagic flux, and increased exocytosis of certain endolysosomal cargos. Through

endolysosomal proteomics, we identified and focused on one such cargo – IL-32, a disease-

associated pro-inflammatory cytokine not present in rodents, whose secretion mechanism is not

well understood. We found that IL-32 was partially secreted in extracellular vesicles likely to be

exosomes. Furthermore, we found that IL-32 was involved in the polarization of inflammatory

reactive astrocyte states and was upregulated in astrocytes in multiple sclerosis lesions. We

believe that our results advance our understanding of cell biological pathways underlying

inflammatory reactive astrocyte phenotypes and identify potential therapeutic targets.

INTRODUCTION

Astrocytes maintain homeostasis of the central nervous system in myriad ways, for example

through phagocytosis of synapses (1,2) or exocytosis of ATP or glutamate (3–5). In the context of

central nervous system injury or disease, astrocytes respond to and amplify inflammatory

signaling cascades (6,7), adopting inflammatory reactive astrocyte states characterized by

distinct gene expression and cytokine signatures (8–10). Although recent efforts have elucidated

these signatures, the cell biological pathways underlying inflammatory reactive astrocyte

phenotypes remain under-explored.

The endolysosomal system encompasses a diverse pool of intracellular vesicles of varying

luminal pH that mediate degradative (e.g. lysosomes, phagosomes, autophagosomes) as well as

exocytic functions (e.g. recycling endosomes, multivesicular bodies) (11,12). Although a large

body of work exists on the cross-regulation of autophagy and inflammatory responses in immune

cells (13), much less is known about how this occurs in brain cell types, especially astrocytes.

Here, using human induced pluripotent stem cell (hiPSC)-derived astrocytes, we found that the

inflammatory cytokines IL-1α+TNF+C1q in combination (hereafter abbreviated as "ITC"),

which have been widely used to induce inflammatory astrocyte reactivity in vitro (14), caused

mTOR activation and mTOR-dependent remodeling of the endolysosomal system, which was

associated with increased exocytosis of certain endolysosomal cargos. Through endolysosomal

proteomics, we identified and focused on one such cargo, the pro-inflammatory cytokine IL-32.

Although cerebrospinal fluid levels of IL-32 are elevated in neuroinflammatory diseases such as

multiple sclerosis (15), the secretion mechanism of IL-32 is not well understood, as it lacks a

classical signal peptide (16,17). Furthermore, it is unclear what cell types in the central nervous

system produce IL-32 under neuroinflammatory conditions. We found that IL-32 was

upregulated in astrocytes in multiple sclerosis lesions, was likely secreted in part via exosomes,

and was involved in the polarization of inflammatory reactive astrocyte states.

RESULTS

To identify the cellular pathways responsible for the loss of homeostatic functions that

accompany inflammatory astrocyte reactivity, we reanalyzed RNA-sequencing data from

multiple hiPSC-derived astrocyte models (18–22) (Supplementary Table 1) treated with ITC or

similar treatments (Fig. 1a), focusing on downregulated genes. We found that downregulated

genes after ITC treatment were enriched in genes encoding proteins associated with the

endolysosomal system (Fig. 1b-c, Supplementary Fig. 1), suggesting a potential link to the

reported phenotype of decreased phagocytosis.

To confirm if ITC indeed perturbed endolysosomal system function, we performed downstream

experiments using the iAstrocyte model that we previously developed (9). We found that ITC

reduced LysoTracker staining without changing total endolysosomal mass as measured by the

abundance of LAMP1 or LAMP2 (Fig. 1d-i, Supplementary File 1), which tend to be enriched in

lysosomes but can also mark other endolysosomal compartments (23). Since a decrease in

LysoTracker staining can reflect a decrease in either total endolysosomal mass or acidity, we

deduced from the above results that ITC caused endolysosomal alkalinization. Indeed, using a

genetically encoded endolysosomal pH reporter (FIRE-pHLy) (24), we found that ITC decreased

the acidity of LAMP1$^+$ endolysosomal compartments (Fig. 1j). We also observed a buildup of

LC3$^+$ puncta on immunostaining after ITC (Fig. 1d-e), suggestive of perturbed autophagic flux.

Using both a genetically encoded reporter (25) (Fig. 1k) and LC3-II western blot (26) (Fig. 1l-m,

Supplementary File 1), we found that ITC decreased autophagic flux as well, consistent with the

importance of acidic lysosomes for degradation of autophagic substrates (27).

To gain a more detailed understanding of how endolysosomal function is perturbed in

inflammatory reactive astrocytes, we performed mass spectrometry-based proteomic

characterization of LAMP1$^+$ endolysosomal compartments in ITC-treated vs control iAstrocytes

(Fig. 2a, see Methods), using the total cell lysate as a reference. We found that v-ATPase

subunits and degradative enzymes were less abundant in endolysosomal compartments in ITC-

treated iAstrocytes (Fig. 2b-c), consistent with transcript-level downregulation in the RNA-seq

data (Fig. 1b). Furthermore, CST3, a potent inhibitor of lysosomal proteases (28), accumulated in

endolysosomal compartments upon ITC treatment (Supplementary Table 2). We also detected

endolysosomal accumulation of proteins involved in vesicular exocytosis (e.g. RAB27A (29),

SNAP25 (3,5)) together with inflammatory mediators (e.g. IL-32 (16,17), CCL2 (30)),

suggesting that ITC may cause rerouting of endolysosomal trafficking to facilitate

unconventional secretion of inflammatory factors.

To explore this hypothesis, we developed a flow-cytometric assay to quantify in live

unpermeabilized iAstrocytes the exposure of the LAMP1 luminal domain on the plasma

membrane (see Methods), which is known to increase in degranulating immune cells (31) and

may be useful as a marker of the exocytic activity of certain endolysosomal pathways (32). With

this approach, we found that ITC-treated iAstrocytes had ~1.8 times the amount of cell-surface

LAMP1 relative to vehicle-treated controls, despite equivalent levels of total LAMP1 (Figure 2d-

e). We were able to directly observe putative endolysosomal exocytic events from LAMP1$^+$

vesicles via total internal reflection fluorescence microscopy of iAstrocytes transduced with a

LAMP1-mCherry fusion construct and loaded with LysoTracker Green, where the green

fluorescence from LysoTracker was lost from LAMP1-mCherry$^+$ vesicles trafficked to the

plasma membrane (Supplementary File 2). Furthermore, the abundance of cell-surface LAMP1

demonstrated a positive dose-response relationship with bafilomycin A1 (Fig. 2f), an inhibitor of

v-ATPase activity and autophagosome-lysosome fusion (33) that is known to increase

endolysosomal exocytosis (34–36). Compared to cell-surface LAMP1, LysoTracker staining

demonstrated a negative dose-response relationship (Fig. 2f), likely reflecting alkalinization of

endolysosomal compartments due to v-ATPase inhibition.

To identify factors downstream of ITC that mediate the perturbations to endolysosomal function

we observed above, we proceeded to perform targeted CRISPR-based inhibition (CRISPRi)

screens against the "druggable genome" (37) in iAstrocytes treated with vehicle or ITC (see

Methods) using cell-surface LAMP1 or LysoTracker staining as proxies of endolysosomal

system function (Fig. 3a). Consistent with our observations from the bafilomycin A1 titration

curve, the phenotype scores (labeled as "gene score", see Methods) of hits from the cell-surface

LAMP1 screens were on average anti-correlated with those from the LysoTracker screens (Fig.

3b,d). Notably, *MTOR* was a top hit in both screens (Fig. 3d), and we also found a strong

enrichment for genes associated with the mTOR pathway in the top hits from both screens (Fig.

3c-d). These results corroborate in hiPSC-derived astrocytes the rich literature on the regulation

of autophagy and lysosome function by mTOR (38,39), much of which was based on

experiments in transformed human cell lines, yeast, or animal models.

To validate our screening results, we first assessed whether ITC increased the activity of

mTORC1or mTORC2, the two protein complexes formed by mTOR with distinct regulatory

subunits and downstream signaling pathways (40). We found that ITC increased mTORC1

activity as measured by increased levels of Ser235/236-phosphoryated S6 ribosomal protein,

Thr37/46-phosphorylated 4E-BP1, and Ser757-phosphorylated ULK1 (38,41) (Fig. 3e-i). ITC

also increased mTORC2 activity as measured by Ser473-phosphorylated AKT1 (42) (Fig. 3j).

To see if there was evidence of mTOR activation in inflammatory reactive astrocytes more

broadly across different contexts, models, and species, we extracted differentially expressed

genes in astrocytes from a variety of published rodent and human transcriptomic datasets and

found broad enrichment for mTOR pathway-associated genes among upregulated genes

(Supplementary Fig. 2), including in a recently published dataset demonstrating a role for

astrocyte mTOR activation in experimental autoimmune encephalitis (EAE) (43), a mouse model

of multiple sclerosis. Next, we tested the effect of mTOR inhibitors on cell-surface LAMP1 and

LysoTracker staining. Consistent with the phenotype scores observed for *MTOR* knockdown in

our CRISPRi screens, both rapamycin and Rapalink-1 (44) decreased cell-surface LAMP1 levels

and increased LysoTracker staining (Fig. 3k-l), although the effect of rapamycin on LysoTracker

staining was not statistically significant. Both drugs significantly decreased levels of Ser235/236-

phosphoryated S6 ribosomal protein (Supplementary Fig. 3), demonstrating target engagement.

Focusing on mTOR as a central regulator of endolysosomal system function, we next explored

how modulating mTORC1 or mTORC2 activity affected the exocytic activity of endolysosomal

pathways. First, we tested whether the changes in cell-surface LAMP1 and LysoTracker staining

caused by ITC depended on mTORC1 vs. mTORC2. We found that knockdown of *MTOR*

decreased cell-surface LAMP1 and increased LysoTracker staining in ITC-treated astrocytes,

which was phenocopied to a larger degree by knockdown of *RPTOR* (which encodes a subunit

unique mTORC1) (45) than *RICTOR* (which encodes a subunit unique to mTORC2) (45) (Fig.

4a), consistent with known mTORC1-dependent mechanisms regulating autophagy and

lysosome function (38,39). We also confirmed that increasing mTORC1 activity through

knockdown of *TSC1*, an upstream inhibitor of mTORC1 (41), increased cell-surface LAMP1 and

decreased LysoTracker staining (Supplementary Fig. 4a-b), as one would predict.

Next, we assessed how mTORC1 vs. mTORC2 controlled the exocytic activity of endolysosomal

pathways by measuring the extracellular levels of three different endolysosomal cargos (see

Methods): cathepsin B (CTSB), a lysosome-enriched protease known to be released

extracellularly by astrocytes (46); IL-32, which we identified as being enriched in endolysosomal

compartments from our proteomics dataset, consistent with prior observations (47); and

mitochondria-containing extracellular vesicles (mito-EVs) (see Methods and Supplementary Fig.

5), which are known to be released by astrocytes and are thought to be derived from mitophagy

(48,49). We found that ITC increased extracellular levels of CTSB, IL-32, as well as mito-EVs,

all of which were dramatically boosted by addition of bafilomycin A1 (Fig. 4b-d). Furthermore,

MTOR knockdown decreased the exocytosis of CTSB, IL-32, and mito-EVs, although for IL-32

we only observed a statistically significant effect in the presence of bafilomycin A1 (Fig. 4b-d).

RPTOR knockdown phenocopied *MTOR* knockdown to a larger degree than *RICTOR*

knockdown for CTSB and mito-EVs, whereas for IL-32 *RICTOR* knockdown had a stronger

effect than *RPTOR* knockdown (Fig. 4b-d). On the other hand, *TSC1* knockdown increased the

exocytosis of mito-EVs and CTSB (Supplementary Fig. 4c-d), as expected. Overall, the stronger

dependence on mTORC1 for the exocytosis of CTSB and mito-EVs compared to IL-32 suggests

that IL-32 exocytosis may occur via a different endolysosomal pathway compared to CTSB and

mito-EVs.

Given prior work which suggested that extracellular IL-32 is membrane-associated and possibly

a component of extracellular vesicles (47,50,51), we proceeded to further characterize the

secretion mechanism of IL-32 by isolating extracellular vesicles (EVs) from iAstrocyte

conditioned media via differential ultracentrifugation (see Methods), with or without ITC

treatment. By western blotting, we verified that our EV preparation contained high levels of

consensus EV markers such as CD63, CD81, Hsc70, and caveolin-1 (52) and was of acceptable

purity, containing undetectable levels of apo-lipoproteins such as ApoA-I (Fig. 5a,

Supplementary File 1), in accordance with the MISEV guidelines (52). On nanoparticle tracking

analysis (see Methods), we saw that the size distribution of EVs from our preparations centered

around a diameter of 100 nm, as expected (52) (Fig. 5b). Both western blotting and nanoparticle

tracking analysis suggested an increase in the concentration of EVs with ITC treatment (Fig. 5a-

b, Supplementary Fig. 6).

Having validated our EV preparation, we proceeded to blot for IL-32 and found that IL-32 was

present in EVs isolated from ITC-treated but not vehicle-treated iAstrocytes (Fig. 5c). To see if

extracellular IL-32 was exclusively EV-associated, we analyzed the EV pellet and supernatant by

western blotting and ELISA. On western blot, IL-32 was detectable only in the EV pellet, as

were EV markers (Supplementary Fig. 7a). However, by ELISA, we saw that extracellular IL-32

partitioned roughly equally between the EV pellet and supernatant (Supplementary Fig. 7b). In

comparison, CTSB exclusively partitioned into the supernatant (Supplementary Fig. 7c). The

discrepancy between the western blot and ELISA data for IL-32 could potentially be explained

by the fact that the western blot and ELISA antibodies are polyclonal antibodies likely raised

against different immunogens. Perhaps the ELISA antibodies detect an IL-32 isoform not

detected by the western blot antibody.

Since EVs can be derived from direct budding of the plasma membrane or multivesicular body

exocytosis (which would then be referred to as exosomes) (52), we performed immunostaining

of ITC-treated iAstrocytes and visualized the subcellular localization of IL-32 with confocal

microscopy. We found that IL-32 co-localized strongly with CD63 (Fig. 5d, Supplementary Fig.

8), which is also a marker of multivesicular bodies (53), suggesting that extracellular IL-32 is

derived from multivesicular body exocytosis. We proceeded to further test this hypothesis by

observing the effect of bafilomycin A1 or knockdown of genes encoding proteins involved in

multivesicular body exocytosis on extracellular IL-32 levels. In agreement with prior reports

which demonstrated increased exosome release with bafilomycin A1 (34,36,54), we found that

treatment with bafilomycin A1 dramatically increased extracellular IL-32 levels (Fig. 5e).

Furthermore, knockdown of *RAB27A/B* decreased extracellular IL-32 levels (Fig. 5e), consistent

with the known role of RAB27A/B in exosome secretion (55). Lastly, corroborating a prior

report which identified a role for PI4KIIIβ in exosome biogenesis (54), we found that inhibition

of PI4KIIIβ (using PI4KIIIβ inhibitor 3 (56)) but not PI4KIIα (using PI-273 (57)) decreased

extracellular IL-32 levels in the presence of bafilomycin A1 (Fig. 5f).

Having explored the secretion mechanism of IL-32, we subsequently characterized its function in

inflammatory astrocyte reactivity. Our prior work established two distinct polarizations of

inflammatory reactive astrocytes after ITC treatment – a VCAM1^{+} interferon/TNF-responsive

polarization associated with CXCL10 secretion vs. a VCAM1^{-}/C3^{+} IL-1/IL-6-responsive

polarization associated with GM-CSF secretion (9). After verifying that knockdown of *IL32*

resulted in robust depletion of IL-32 at the protein level (Supplementary Fig. 9a-b), we found

that *IL32* knockdown decreased the proportion of VCAM1$^+$/C3$^-$ astrocytes and CXCL10

secretion (Fig. 6a-b), suggesting that IL-32 promotes the interferon/TNF-responsive polarization.

Also, IL-32 induction by ITC was greater in VCAM1$^+$/C3$^+$ interferon-responsive astrocytes

compared to IL-1/IL-6-rresponsive VCAM1$^-$/C3$^+$ astrocytes (Supplementary Fig. 9c-e).

Furthermore, we found that IFN-β increased the upregulation of IL-32 by ITC (Supplementary

Fig. 9f), consistent with prior work demonstrating a role of IL-32 in antiviral responses (58–60).

Next, given that IL-32 levels have been reported to be elevated in the cerebrospinal fluid of

patients with multiple sclerosis and neuro-Behcet's disease (15), we wanted to see if IL-32 was

upregulated in astrocytes under neuroinflammatory conditions. Although the fact that IL-32 does

not have an ortholog in rodents precluded us from analyzing the vast amount of published

transcriptomic data on mouse models of neuroinflammation, we found that *IL32* transcript levels

were indeed upregulated in astrocytes in various types of multiple sclerosis lesions in humans

(Fig. 6c) by reanalyzing the comprehensive single-nucleus RNA-seq dataset from Macnair *et al.*

(61) at the pseudobulk level (62) (see Methods). Interestingly, *IL32* upregulation in multiple

sclerosis lesions was strongest in astrocytes compared to other cell types (Supplementary Fig.

10). Consistent with this observation, we found that IL-32 preferentially colocalized with GFAP$^+$

astrocytes in post-mortem brain tissue from pediatric patients diagnosed with hypoxic-ischemic

encephalopathy (Fig. 6d-e), a condition involving significant neuroinflammation (63) where we

had previously identified upregulation of inflammatory reactive astrocyte markers (9).

DISCUSSION

Our results establish mTOR activation as a key feature of inflammatory astrocyte reactivity

induced by IL-1α+TNF+C1q (ITC), driving endolysosomal remodeling manifesting as

alkalinization of LAMP1$^+$ compartments and reduced autophagic flux (Fig. 6e). In light of the

above, mTOR-induced endolysosomal remodeling may account for the loss of phagocytic

activity observed in inflammatory reactive astrocytes (14). Although we did not measure

phagocytic activity here, we observed in our previously published CRISPRi screens on

inflammatory reactivity that *MTOR* knockdown rescued the phagocytic deficit induced by ITC

(9)].

Here, we found that mTOR-dependent endolysosomal remodeling also resulted in the increased

exocytosis of certain endolysosomal cargos (Fig. 6e), with increased cell-surface LAMP1 likely

acting as a non-specific marker of endolysosomal exocytic activity. Interestingly, LAMP1$^+$

astrocytes have been shown to modify disease progression in EAE (64). Furthermore, all three of

the endolysosomal cargos we have characterized – CTSB, IL-32, and mito-EVs – have been

reported to be involved in neuroinflammatory conditions. CTSB levels are elevated in the CSF as

well as brain parenchyma of patients with Alzheimer's disease (65–68), and knockout of *Ctsb*

has been shown to ameliorate the neuropathology and behavioral deficits in mouse models of

Alzheimer's disease (69–71). Mito-EVs (72) have been shown to mediate the transfer of

mitochondria from astrocytes to neurons after experimentally induced stroke in mice (48). Lastly,

IL-32 levels are elevated in the cerebrospinal fluid of patients with multiple sclerosis or neuro-

Behcet's disease (15), and a polymorphism in the IL-32 promoter has been associated with

increased risk of multiple sclerosis in two independent studies (73,74).

Given that IL-32 does not have an ortholog in rodents and that its secretion mechanism is still

incompletely characterized, we focused on elucidating IL-32 secretion, taking advantage of our

in vitro hiPSC-derived astrocyte platform. We found that after ITC treatment, intracellular IL-32

colocalized with multivesicular bodies, and that extracellular IL-32 co-fractionated with EVs.

Whether extracellular IL-32 is exclusively associated with EVs remains to be seen, as we

obtained conflicting data by western blotting vs. ELISA. Given that the western blot and ELISA

antibodies are polyclonal antibodies likely raised against different immunogens, it is possible

that the ELISA antibodies recognize a non-EV-associated IL-32 isoform not detected by the

western blot antibody. Alternatively, if extracellular IL-32 consists predominantly of one

isoform, it may be cleaved in a way such that the western blot antibody recognizes only the EV-

associated fragment whereas the ELISA antibodies recognize both the EV-associated and free

fragments. Further biochemical experiments will be necessary to elucidate whether IL-32

isoforms may be differentially associated with EVs or if extracellular cleavage of IL-32 occurs.

With respect to the secretion mechanism of extracellular IL-32, we found that knockdown of

genes encoding proteins involved in multivesicular body exocytosis such as RAB27A/B

decreased extracellular levels of IL-32, as did pharmacological inhibition of exosome biogenesis.

Overall, our results corroborate previous reports demonstrating that a portion of extracellular IL-

32 is vesicle-associated (50,51), and we establish exosomes as the likely candidate. With respect

to extracellular IL-32 not contained within EVs, it is possible that IL-32 may be loaded directly

into multivesicular body lumens and subsequently exocytosed, or that intracellular IL-32 may be

released directly through plasma membrane leakage in dying or dead cells (75).

As for the role of IL-32 in neuroinflammation, we found that knockdown of *IL32* affected the

polarization of inflammatory reactive states induced by ITC (9), decreasing the abundance of the

interferon/TNF-responsive state and its associated cytokine CXCL10. A limitation of knocking

down *IL32* with DNA-targeting CRISPRi (76) is that we could not distinguish the contribution of

the many splice isoforms of IL-32, which have been reported to have distinct activities (77);

future work could elucidate this using RNA interference or RNA-targeting CRISPR-based

systems (78). In addition to its effects on the polarization of inflammatory reactive astrocyte

states, we also found that IL-32 was induced by IFN-β, and that *IL32* transcript levels were

specifically upregulated in astrocytes in various multiple sclerosis lesions. Considering the

human genetics data demonstrating the importance of IL-32 to the pathogenesis of multiple

sclerosis, studying how astrocyte IL-32 contributes to multiple sclerosis would be a worthwhile

future research direction. Lastly, consistent with the preferential upregulation of *IL32* in

astrocytes in multiple sclerosis, we found preferential colocalization of IL-32 with astrocytes in a

different neuroinflammatory condition – hypoxic-ischemic encephalopathy (HIE), suggesting

that IL-32 may play a role in HIE as well.

In conclusion, we believe that our results highlight mTOR-dependent endolysosomal remodeling

as an important and previously underappreciated aspect of inflammatory astrocyte reactivity

which can be targeted therapeutically. We also clarified the secretion mechanism and functional

role of an important disease-associated cytokine, IL-32, in astrocytes, a cell type in which IL-32

has rarely been studied (79). Since the receptor for IL-32 is still unknown (16), we believe that

our results establish a strong foundation for future studies focused on the how IL-32 mediates its

biological effects and contributes to neuroinflammation.

COMPETING INTERESTS

A.W.K is a member of the Scientific Advisory Board for Nine Square Therapeutics, Inc. J.E. is

an advisor to Seer, Inc. M. K. has filed a patent application related to CRISPRi and CRISPRa

screening (PCT/US15/40449) and serves on the Scientific Advisory Board of Engine

Biosciences, Casma Therapeutics, and Cajal Neuroscience, and is an advisor to Modulo Bio and

Recursion Therapeutics. None of the other authors declare competing interests.

FUNDING

This research was supported by National Institutes of Health (NIH) grant F30 AG066418 to

K.L.; California Institute for Regenerative Medicine grant EDUC4-12812 and NIH grant T32

NS115706 to I.V.L.R.; Chan Zuckerberg Initiative Ben Barres Early Career Acceleration Awards

and NIH/NIA R01 AG062359 to M.K.; NIH/NIND grants (R01NS097551, P01NS083513 and

R21NS119954) to S. Fancy; NIH grants R01AG057342 and R56AG057342 to A.K. and M.C.

S.Fancy is a Harry Weaver Neuroscience Scholar of the National Multiple Sclerosis Society.

J.E.E., M.L., F.M, and S.B. were supported by the Chan Zuckerberg Biohub. The funders had no

role in study design, data collection and analysis, decision to publish or preparation of the

manuscript.

AUTHOR CONTRIBUTIONS

K.L., B.R., and M.K. conceptualized and led the overall project, and wrote the manuscript with

input from all co-authors. K.L. and B.R. performed the experiments and data analysis unless

stated otherwise. In consultation with J.E., F.M aided with sample preparation for whole-cell and

lysosome-specific proteomics experiments, and performed mass spectrometry and preliminary

computational analysis. I.V.L.R. performed sample prep for the cell-surface LAMP1 CRISPRi

screens. K.A.H. guided TIRF experiments. M.L. and S.B aided with design and sample

preparation for whole-cell and lysosome-specific proteomics experiments, respectively. M.Y.C.

and A.W.K provided FIRE-pHLy reagents and guidance on lysosome pH experiments. S.F

conducted LAMP2 immunofluorescence experiments and data analysis. All authors reviewed the

manuscript.

ACKNOWLEDGEMENTS

We thank Greg Mohl, Avi Samelson, Olivia Teter, Biswa Ramani, Brandon Desousa, Emmy Li,

Vukasin Jovanovic, Zuzana Krejciova, Nawei Sun, and Carlo Condello for contributions to

preliminary studies or helpful discussions. We thank members of the Kampmann lab for

discussions and feedback on the manuscript. We thank the staff at the UCSF Center for

Advanced Light Microscopy and the UCSF Parnassus Flow Cytometry CoLab

(RRID:SCR_018206) for their assistance. A portion of the imaging data presented in this

manuscript were acquired on microscopes at the UCSF Center for Advanced Light Microscopy

and the Innovation Core at the Weill Institute, a part of the UCSF Center for Advanced Light

Microscopy.

AVAILABILITY OF DATA AND MATERIALS

The source data and code used to analyze the data presented in this study will be shared upon

request. The source data for all western blots shown in this study can be found in Supplementary

File 1.

Figure 1 | Endolysosomal function is perturbed in inflammatory reactive astrocytes. a,

Schematic of modeling inflammatory astrocyte reactivity in vitro. **b**, Heatmap of changes in the

expression of genes encoding lysosome-localized proteins in different hiPSC-derived astrocyte

models treated with ITC or similar treatments vs vehicle control. **c**, Top GO Cellular Component

terms enriched among the top 1000 downregulated genes in iAstrocytes treated with ITC;

endolysosomal system-related terms are highlighted by asterisks. **d**, Representative images of

LysoTracker staining in live iAstrocytes or immunostaining of LC3 or LAMP1 in fixed and

permeabilized iAstrocytes; scale bar = 75 um. **e**, Quantification of imaging experiments shown in

d (n = 3 wells per condition). **f-g**, Representative immunoblot bands against LAMP1 (**f**) or

LAMP2 (**g**). **h-i**, Quantification of immunoblot experiments shown in **f-g** (n = 3 wells per

condition). **j**, Measurement of the acidity of LAMP1$^+$ endolysosomal compartments using FIRE-

pHLy via flow cytometry (see Methods). **k**, Measurement of autophagic flux with the LC3ΔG-

RFP/LC3-GFP fluorescent reporter from Kaizuka *et al*. (25). **l**, Immunoblot against LC3

demonstrating LC3-I and LC3-II bands. **m**, Quantification of autophagic flux from the LC3-II

bands in **l**; error bars reflect the 95% confidence interval associated with the standard error of the

mean; individual data points not shown because the quantities of interest are differences between

means, with no biologically meaningful pairing of individual data points across conditions. *P*

values where shown were calculated using two-sided Student's t test.

Figure 2 | Perturbation of endolysosomal function is accompanied by remodeling of the

endolysosomal proteome. **a,** Schematic of endolysosomal proteomic workflow (n = 3 wells per

condition). **b,** Volcano plot of endolysosomal proteomic data. **c,** Scatterplot comparing $\log_2$-fold

change of endolysosomal vs total cell lysate protein abundance in ITC-treated iAstrocytes

compared to vehicle-treated iAstrocytes (IP – immunoprecipitation). **d,** Representative

histograms of cell-surface LAMP1 staining intensity in vehicle- or ITC-treated iAstrocytes

compared to isotype control staining intensity measured by flow cytometry. **e,** Median

fluorescence intensity (MFI) of cell-surface LAMP1 measured by flow cytometry (n = 3 wells

per condition); *P* value by two-sided Student's t test. **f,** Cell-surface LAMP1 or LysoTracker

426   staining in iAstrocytes treated with increasing doses of bafilomycin A1 (n = 3 wells per

427   condition; error bars reflect the standard error of the mean).

Figure 3 | Multi-phenotypic CRISPR-based screening identifies mTOR as a central

regulator of endolysosomal system function. a, Schematic of CRISPR-based screening

workflow. **b**, Pearson correlation of gene scores (see Methods) of hits from the LAMP1 vs

LysoTracker screens (n = 2 biological replicate screens per condition). **c**, Enrichment analysis

against MSigDB Hallmark Pathways terms of the top 20 hits from each screen; terms pertaining

to mTOR are highlighted with stars. **d**, Heatmap of gene scores of the hits overlapping with the

highlighted MSigDB terms in **c**. **e-f**, Median fluorescence intensity (MFI) of phospho-S6 (**e**) or

pospho-4E-BP1 staining in ITC- vs. vehicle-treated iAstrocytes measured by flow cytometry. **g**,

Representative immunoblot bands corresponding to mTOR, phospho-S6, total S6, phospho-

ULK1 and phospho-AKT1 in ITC- vs vehicle-treated iAstrocytes in the presence of different

mTOR inhibitors. **h**, Quantification of immunoblotting experiments (n = 6 wells per condition

for phospho-S6/S6, n = 3 wells per condition for phospho-ULK1, n = 6 wells per condition for

pospho-AKT1/pan-AKT; *P* values from two-sided Student's t test). **k-l**, Cell-surface LAMP1 (**k**)

or LysoTracker (**l**) MFI in ITC- vs. vehicle-treated iAstrocytes in the presence of mTOR

inhibitors measured by flow cytometry (n = 6 wells for DMSO treated, n = 3 wells for all other

conditions; *P* values calculated only for ITC-treated conditions by linear regression with

adjustment for multiple testing by Holm's method, shown only if significant).

Figure 4 | Cytokine-induced remodeling or pharmacological disruption of the endolyosomal

system results in increased exocytosis of endolysosomal contents. **a-d**, Cell-surface LAMP1

or LysoTracker median fluorescence intensity (MFI) measured by flow cytometry (**a**),

extracellular CTSB concentration measured by electrochemiluminescence-based immunoassay

(**b**), extracellular IL-32 concentration measured by ELISA (**c**), or abundance of extracellular

mito-EVs measured by flow cytometry (**d**) in ITC- vs. vehicle-treated iAstrocytes transduced

with non-targeting control (NTC) sgRNAs or sgRNAs targeting genes encoding common

(*MTOR*) or unique mTORC1 (*RPTOR*) vs mTORC2 (*RICTOR*) subunits, with or without co-

treatment with bafilomycin A1. *P* values were calculated by linear regression with correction for

multiple testing using Holm's method, shown only when significant.

Figure 5 | Extracellular IL-32 co-fractionates with extracellular vesicles likely derived from

multivesicular body exocytosis. **a**, Immunoblots against consensus extracellular vesicle (EV)

markers (CD63, CD81, Hsc70, Caveolin-1) or common contaminants (e.g. ApoA-I) in EVs

isolated from iAstrocyte conditioned media or total cell lysate after vehicle vs. ITC treatment. **b**,

EV size distribution measured by nanoparticle tracking analysis. **c,** Immunoblot against IL-32 in

EVs isolated from iAstrocyte conditioned media or total cell lysate. **d**, Representative images of

dual immunostaining against LAMP1 together with IL-32 or CD63 together with IL-32; scale

bar = 60 μm. **e**-**f**, Extracellular IL-32 concentration measured by ELISA in conditioned media

from ITC- vs. vehicle-treated iAstrocytes transduced with non-targeting (NTC) sgRNAs or

sgRNAs targeting genes encoding proteins involved in multivesicular body exocytosis (**e**), or

treated with small molecules known to inhibit (PI4KIIIβ inhibitor 3) or not inhibit (PI-273)

exosome biogenesis. *P* values were calculated by linear regression with correction for multiple

testing by Holm's method, shown only when significant.

Figure 6 (legend overleaf)

Figure 6 | IL-32 regulates the polarization of inflammatory reactive astrocytes and is

upregulated in astrocytes in neuroinflammatory conditions. **a**-**b**, Proportion of IL-1/IL-6-

responsive (VCAM1-/C3+) or TNF/interferon-responsive (VCAM1+/C3-, VCAM1+/C3+)

inflammatory reactive astrocyte polarizations (**a**) or their associated cytokines (**b**) in ITC- vs.

vehicle-treated iAstrocytes transduced with non-targeting (NTC) sgRNAs or sgRNAs targeting

IL32. **c**, Log-scaled *IL32* expression in astrocytes found in normal tissue vs. multiple-sclerosis

lesions derived from pseudobulk analysis of snRNA-seq data from Macnair *et al.* **d**,

Representative immunostaining of IL-32 and GFAP in white matter brain tissue from patients

with hypoxic-ischemic encephalopathy (HIE); scale bar 50 μm. e, Percent GFAP+, OLIG1+, or

NeuN+ cells among IL-32+ cells in HIE brain tissue (n = 3 patients); *P* values calculated via beta

regression. **f**, Schematic of ITC-induced, mTOR-dependent endolysosomal remodeling and

associated exocytic activity. *P* values were calculated using the Mann-Whitney U test in **a** and **c**

and using the two-sided Student's t test in **c**.

Supplementary figure 1 | Overlap analysis of downregulated genes with endolysosomal-related terms in inflammatory reactive astrocytes across different contexts and species. GO Cellular Component terms corresponding to different components of the endolysosomal system

500 were analyzed for overlap with downregulated genes in human or rodent astrocytes (all from

501 mouse except where noted) found in various neuroinflammatory models and conditions. See

502 Supplementary Table 3 for metadata corresponding to the studies analyzed. *P* values were

503 adjusted for multiple comparisons with the Benjamini-Hochberg procedure.

Supplementary figure 2 | Overlap analysis of upregulated genes with mTOR-related terms

in inflammatory reactive astrocytes across different contexts and species. mTOR-related

terms from MSigDB, KEGG, or BioPlanet were analyzed for overlap with downregulated genes

in human or rodent astrocytes (all from mouse except where noted) found in various

neuroinflammatory models and conditions. See Supplementary Table 3 for metadata

511   corresponding to the studies analyzed. *P* values were adjusted for multiple comparisons with the

512   Benjamini-Hochberg procedure.

515　**Supplementary figure 3 | Target engagement analysis of mTOR inhibitors**. Median

516　fluorescence intensity (MFI) of phospho-S6 in ITC- vs. vehicle-treated iAstrocytes co-treated

517　with rapamycin or Rapalink-1 (n = 3 per condition).

Supplementary figure 4 | Modulation of mTORC1 activity affects endolysosomal exocytosis

bidirectionally. **a**-**d**, Median fluorescence intensity (MFI) of cell-surface LAMP1 (**a**) or

LysoTracker (**b**) staining measured by flow cytometry, abundance of mito-EVs in conditioned

media (**c**), or extracellular CTSB concentration measured by electrochemiluminescence-based

immunoassay (**d**) in ITC- vs. vehicle-treated iAstrocytes transduced with non-targeting (NTC)

sgRNAs or sgRNAs targeting *RPTOR* or *TSC1* (n = 7 wells per condition for LAMP1 and

LysoTracker data, n = 8 wells per condition for mito-EV and CTSB data; *P* values calculated by

linear regression with Holm's correction for multiple testing, shown only if significant).

Supplementary figure 5 | Gating strategy for quantifying mito-EVs from iAstrocyte

conditioned media. Events were first gated on MitoTracker Green fluorescence, with MTG$^+$

events subsequently gated by Hoechst staining. Mito-EVs were defined as MTG$^+$/Hoechst$^+$

events.

543   **Supplementary figure 6 | ITC-treated astrocytes release more extracellular vesicles**.

544   Quantification of CD63 or CD81 immunoblot bands from total cell lysate or extracellular

545   vesicles from ITC- vs. vehicle-treated iAstrocytes shown in Fig. 5a (n = 3 wells per condition; *P*

546   values by two-sided Student's t test).

Supplementary figure 7 | Partition analysis of IL-32 or CTSB. Concentration of IL-32

measured by ELISA (**a**) or CTSB measured by electrochemiluminescence-based immunoassay

(**b**) in resuspended extracellular vesicle pellets (lysed with detergent) vs the supernatant (see

Methods); n = 3 per condition, *P* values calculated only for ITC-treated conditions by two-sided

Student's t test, shown only when significant.

Supplementary figure 8 | Colocalization analysis of IL-32 with LAMP1 or CD63. Pearson's

colocalization coefficient between IL-32 and LAMP1 (left) or between IL-32 and CD63 (right)

in ITC-treated iAstrocytes co-treated with vacuolin-1 or bafilomycin A1, corresponding to

images from the immunostaining experiment shown in Fig. 5d (n = 3 wells per condition; *P*

values calculated by beta regression with correction for multiple testing by Holm's method,

shown only if significant).

Supplementary Figure 9 | IL-32 levels are higher in interferon/TNF-responsive

inflammatory reactive astrocytes and after treatment with IFN-β. **a**, Immunoblot against IL-

32 in ITC-treated iAstrocytes transduced with non-targeting (NTC) sgRNAs or sgRNAs

targeting *IL32*. **b**, Quantification of IL-32 levels from immunoblot shown in **a**. **c**, Median

fluorescence intensity (MFI) of IL-32 in IL-1/IL-6-responsive (VCAM1-/C3+) or

TNF/interferon-responsive (VCAM1+/C3-, VCAM1+/C3+) inflammatory reactive astrocytes as

defined in Leng *et al.* [ref]. **d**-**e**, *IL32* transcript levels from single-cell RNA-seq data on ITC- vs.

vehicle-treated iAstrocytes from Leng *et al.* (9) visualized on uniform manifold approximation

embedding (**d**) or as violin plots (**e**). **f**, IL-32 MFI in ITC- vs. vehicle-treated iAstrocytes treated

concurrently with vehicle or IFN-β transduced with non-targeting (NTC) sgRNAs or sgRNAs

targeting *IL32*. Dotted lines in **c** and **f** reflect estimated level of non-specific staining based on

degree of IL-32 knockdown from **b.** *P* values where shown were calculated by two-sided

Student's t test.

Supplementary Figure 10 | IL32 expression in non-astrocyte cell types in snRNA-seq data

from Macnair *et al.*(61) **a-f,** Log-scaled normalized pseudobulk counts of *IL32* in excitatory

neurons (**a**), inhibitory neurons (**b**), oligodendrocytes (**c**), oligodendrocyte precursor cells

584   (OPCs) and differentiation-committed oligodendrocyte precursors (COPs) (**d**), microglia (**e**), or

585   endothelial cells (endo) and pericytes (peri) (**f**).

METHODS

hiPSC culture

Human iPSCs (male WTC11 background (80)) were cultured in Essential 8 (E8) Medium

(ThermoFisher Scientific cat. no. A1517001) on BioLite Cell Culture Treated Dishes

(ThermoFisher Scientific) coated with Growth Factor Reduced, Phenol Red-Free, LDEV-Free

Matrigel Basement Membrane Matrix (Corning cat. no. 356231) diluted 1:100 in DMEM/F12

(ThermoFisher Scientific cat. no. 11330032). Essential 8 Medium was replaced daily. When

hiPSC colonies demonstrated mature morphology, the hiPSCs were either clump passaged with

EDTA for routine maintenance or dissociated to a near single-cell suspension with Accutase Cell

Dissociation Reagent (ThermoFisher Scientific cat. no. A11105-01) for applications requiring

cell counting. For clump passaging with EDTA, hiPSCs were washed with Dulbecco's phosphate

buffered saline (DPBS; Milipore Sigma cat. no. D8537) and then incubated with Versene

(ThermoFisher Scientific cat. no. 15040066) for 5-7 min at room temperature; the Versene

solution was then aspirated and replaced with E8 + 10 nM Y-27632 dihydrochloride ROCK

inhibitor (Tocris cat. no. 125410); hiPSC colonies were then gently detached mechanically using

a cell scraper, resuspended gently, and passaged at 1:10-1:30 dilution in E8 + Y-27632, with Y-

27632 removed the next day. For near single-cell dissociation, hiPSCs were washed with DPBS,

incubated with Accutase for 5-10 min at 37 °C, and then gently triturated with a P1000 pipette

tip; the cell suspension was then diluted with PBS, collected into conical tubes, and spun down at

300 g for 3 min; hiPSCs were then resuspended in E8 + Y-27632, counted, and plated onto

Matrigel-coated plates at the desired density in E8 + Y-27632; Y-27632 would be maintained

until the hiPSC colonies reached the appropriate size (> ~40 cells). Studies with hiPSCs at

UCSF were approved by the The Human Gamete, Embryo and Stem Cell Research (GESCR)

Committee. Informed consent was obtained from the human subject when the WTC11 line was

originally derived.

Generation of iAstrocytes

iAstrocytes were generated as detailed in Leng *et al*. (9). Briefly, we generated hiPSCs (male

WTC11 background) used for iAstrocyte differentiation by inserting a CRISPRi machinery

cassette (pC13N-dCas9-BFP-KRAB, Addgene plasmid no. 127968) into the CLYBL safe-harbor

locus and a doxycycline-inducible *NFIA* and *SOX9* cDNA cassette into the AAVS1 safe-harbor

locus using TALEN-based editing. After recovering a monoclonal CRISPRi-*NFIA-SOX9* line, we

differentiated hiPSCs to the neuroectoderm lineage via an embryoid body (EB)-based neural

induction protocol. After dissociating plating EB-derived cells, we selected and expanded those

forming neural rosettes. We used fluorescence activated cell sorting (FACS) to enrich for pure

CD133$^+$/CD271$^-$ neural progenitor cell (NPC) populations for expansion.

For iAstrocyte differentiation, these NPCs were re-plated on 1:200 Matrigel-coated dishes and

cultured with ScienCell Astrocyte Media (ScienCell Research Laboratories cat. no. 1801) + 2

µg/mL doxycycline (Millipore Sigma cat. no. D9891) to initiate iAstrocyte differentiation. A full

media change was completed every other day (maintaining doxycycline at 2 µg/mL) until NPCs

reached confluency, at which time they were dissociated with Accutase and split 1:10 onto new

Matrigel coated dishes (or cryopreserved) for continued differentiation and expansion. Expansion

was continued until day 20 (d20) of differentiation, yielding iAstrocytes. See Leng *et al*. for full

details of iAstrocyte hiPSC generation and differentiation protocols.

Induction of inflammatory reactivity in iAstrocytes

For all iAstrocyte experiments in this study, d20 iAstrocytes were plated at 20,000 cells/cm^2 on

1:200 Matrigel-coated BioLite Cell Culture Treated Dishes (ThermoFisher Scientific) in

ScienCell Astrocyte Media (without doxycycline) on d0. Full media changes were completed on

d1, d3, and d5. On d5, media was supplemented with IL-1α (3 ng/mL; Peprotech cat. no. AF-

200-01A), TNF (30 ng/mL; Peprotech cat. no. AF-300-01A), and C1q (400 ng/mL; Complement

Technology cat. no. A099) to induce inflammatory reactivity according to Liddelow *et al*. (14).

All experiments were conducted 24 hours after cytokine treatment (day 6 after plating). For

experiments involving addition of additional IFN-β, 5 ng/mL of IFN-β was added concurrently

with ITC.

CRISPRi-mediated gene knockdown using lentiviral transduction of individual sgRNAs

For knocking down target genes with a single sgRNA, we cloned CRISPRi sgRNAs into

pMK1334 (81) (Addgene cat. no. 127965), as previously described (82). For knocking down

target genes using two sgRNAs to achieve greater degree of knockdown, we cloned sgRNAs into

pMK1334 with an additional hU6 insert as described in Replogle *et al.* (83). We generated

lentivirus of the resulting constructs by co-transfecting them with 3rd generation lentiviral

packaging plasmids with TransIT-Lenti Transfection Reagent (Mirus cat. no. MIR6606)

according to the manufacturer's protocol. The next day, lentivirus was precipitated using

Lentivirus Precipitation Solution (ALSTEM cat. no. VC150) according to the manufacturer's

protocol, resuspended in DPBS, and aliquoted and stored at -80 °C. iAstrocytes were transduced

with lentivirus at the time of plating. When necessary, the functional titer of the lentivirus was

determined by measuring BFP+ cells 48 hours after transducing iAstrocytes with a serial dilution

of lentivirus. The protospacer sequences of the sgRNAs used in this study were selected from

those generated in Horlbeck *et al.* (37) and are listed below (when two protospacers are listed,

the sgRNA is cloned into pMK1334 using the Replogle *et al.* method):

Non-targeting controls (NTC):

GGAGTTAAGGCCTCGTCTAG, GTGCGGGGGCATGGCCCCGC

MTOR:

GGGACAGCGGGGAAGGCGGG

RPTOR:

GACCCCAGGTCCCAAGCCAC, GTGGGGGCCGCTAGGAAATG

RICTOR:

GGCGCAGGGCGGAATGACAG, GAGCGGGCTTACCTCGTACT

TSC1:

GGGCCTTGGCCCTTTCACGA

VAMP7:

GGCCAACTGCCCGCTCCCAG, GAGGGACGCGGGTCAGTGCA

VAMP3:

GCGGCAGCGGCGACGAGAGA, GGAAGCGAAGTTGGGACCGG

SNAP23:

GAACTCGGACACCCCAACAC, GTAGGGTGCAGCGCCAGGTC

RAB27A/B:

GACCCGAGCCAGGCGGGGAC (*RAB27A*), GCCCATCCGGAGCCACAGGT (*RAB27B*)

IL32:

GGAGCTGGGTCATCTCAGGT, GCGAAGGTGAGGACCCTCTG

Generation and lentiviral transduction of reporter constructs

To subclone the original FIRE-pHLy construct (24) (pLBR10, Ubc:FIRE-pHLy) to make

pLBR11 (GfaABC1D:FIRE-pHLy), we excised the FIRE-pHLy cassette from pLBR10 and

linearized the backbone vector (pKL20; pHIV gfaABC1D:mRuby:WPRE) with BamH1-HF

(New England BioLabs, cat. no. R3136) and ClaI (New England BioLabs cat. no. R0197). We

then ran these reactions on a 1% agarose gel, gel extracted the relevant fragments, and ligated

them together with T4 DNA Ligase (New England BioLabs cat. no. M0202) per manufacturer's

protocol to generate pLBR11. Both pLBR11 (all FIRE-pHLy experiments), and GFP-LC3-RFP-

LC3ΔG (reporter cassette from Addgene plasmid #84572) (25) reporters were transduced into

iAstrocytes with lentivirus at low multiplicity of infection (MOI < 1) during plating to prevent

high expression of exogenous lysosomal proteins.

Immunofluorescence imaging

For immunofluorescence imaging experiments involving iAstrocytes, iAstrocytes plated in

Greiner μClear 96-well plates (Greiner Bio-One cat. no. 655087) were fixed with 4%

paraformaldehyde (diluted from a 16% solution in DPBS; Electron Microscopy Sciences cat. no.

15710) for 15 min at RT. After washing three times with DPBS, we blocked and permeabilized

cells with DPBS (Milipore Sigma cat. no. D8537) + 3% BSA (Milipore Sigma cat. no. A9647) +

0.1% Triton X-100 (Millipore Sigma cat. no. X100) for 30 min at RT. Primary antibodies against

LC3 (1:50; Cell Signaling Technology cat. no. 2775), LAMP1 (1:50; Abcam cat. no. ab25630),

LAMP2 (1:500; Abcam cat. no. ab25631), IL-32 (1:250; ProteinTech cat. no. 11079-1-AP), or

CD63 (1:1420; BioLegend cat. no. 353039) were added to blocking buffer and incubated with

iAstrocytes overnight at 4 °C. Afterwards, the samples were washed with DPBS three times,

incubated with pre-adsorbed secondary antibodies (1:500 goat anti-mouse IgG Alexa Fluor 488,

1:500 goat anti-rabbit IgG Alexa Fluor 555; Abcam cat. no. ab150117 and ab150086) for 1 hour

at RT, washed three times with DPBS, incubated with 1 μg/mL Hoechst 33342 (ThermoFisher

Scientific cat. no. H3570) for 10 minutes and then washed two additional times before imaging

on an IN Cell Analyzer 6000, using a 60X 0.7 NA objective, 2×2 binning, 100-400 ms exposure,

an aperture width of ∼1 Airy unit, and 9-16 fields per well, or on an ImageExpress Confocal

HT.ai (Molecular Devices), using an APO 20x 0.95 NA water-immersion objective (Nikon), 89

North LDI lasers 405 nm, 475 nm, and 555 nm with emission filters 452/32 nm 520/428 nm and

598/25 nm for Hoechst 333342, AlexaFluor 488 and 555, respectively, 1x1 binning (325 x 325

nm pixel size), 100-400 ms exposure, 60 μm pinhole spinning disk, and 12 fields per well

captured on a Zyla 4.5 sCMOS camera (Andor).

For immunofluorescence imaging experiments involving HIE samples, all human HIE tissue was

collected with informed consent and in accordance with guidelines established by the UCSF

Committee on Human Research (H11170-19113-07), as previously described (84). Immediately

after procurement, all brains were immersed in PBS with 4% paraformaldehyde for 3 days. On

day 3, the brain was cut in the coronal plane at the level of the mammillary body and immersed

in fresh 4% paraformaldehyde and PBS for an additional 3 days. After fixation, all tissue

samples were equilibrated in PBS with 30% sucrose for at least 2 days. After sucrose

equilibration, tissue was placed into molds and embedded with OCT medium for 30 minutes at

room temperature, followed by freezing in dry-ice-chilled ethanol. UCSF neuropathology staff

performed brain dissection and its evaluation. The diagnosis of HIE requires clinical and

pathological correlation; no widely accepted diagnostic criteria are present for the pathological

diagnosis of HIE. HIE cases showed consistent evidence of diffuse white matter gliosis, as

evaluated by the qualitative increase in the number of GFAP$^+$ cells in addition to the increased

intensity of GFAP staining. For immunostaining of reactive astrocyte markers, tissue slides were

bleached with UV overnight, rinsed with PBS for 10 minutes and incubated with blocking

solution (10% normal goat serum + 0.2% Triton X-100 in PBS) for 1 hour at room temperature

and then incubated with the following primary antibodies overnight at 4 °C: chicken anti-GFAP

(1:500 dilution; BioLegend cat. no. 829401), mouse anti-NeuN (1:500 dilution; Millipore Sigma

cat. no. MAB377), mouse anti-OLIG1 (1:200; Millipore Sigma cat. no. MAB2417), or rabbit

anti IL32 (1:200 dilution; ProteinTech cat. no. 11079-1-AP). Afterwards, the samples were rinsed

three times with PBS + 0.2% Triton X-100 for 10 minutes each time and then incubated with

the appropriate secondary antibodies for 1 hour at room temperature in the dark.

LysoTracker staining and imaging

iAstrocytes were washed with DPBS and incubated with a 50 nM solution of LysoTracker Green

DND-26 (ThermoFisher Scientific cat. no. L7526) or LysoTracker Red DND-99 (ThermoFisher

Scientific cat. no. L7528) in DPBS for 5 minutes at 37°C, washed two times with DPBS, and

processed for downstream experiments. For imaging, iAstrocytes were imaged at 37°C and 5%

CO_2 using an IN Cell Analyzer 6000 as above. For flow cytometry (CRISPRi screens),

iAstrocytes were dissociated with Accutase for 10 mins at 37°C after LysoTracker staining, and

resuspended in FACS buffer (DPBS + 1% BSA + 2 mM EDTA (Milipore Sigma cat. no.

324506)).

Total internal reflection fluorescence microscopy

iAstrocytes were plated at 20,000 cells/cm2 in 35 mm imaging dishes (Ibidi, cat. no. 81156) and

simultaneously transduced with pLBR11 (GfaABC1D:FIRE-pHLy). Prior to imaging, the

conditioned media of iAstrocytes was collected. iAstrocytes were then stained with LysoTracker

as detailed above and re-incubated with the collected conditioned media for the duration of

imaging. TIRF was performed on a Nikon Ti-E inverted microscope equipped with a Nikon

TIRF illuminator, Nikon motorized stage with piezo Z inset, Andor iXon Ultra DU897 camera,

and run by NIS-Elements (v5.20.00 build 1423). iAstrocytes were imaged using an Apo TIRF

100x/1.49 oil (DIC N2/ 100X I) objective, 488nm and 561nm lasers (Agilent MLC400

monolithic laser combiner) and Sutter Lambda 10-B with ET525/50m and ET600/60m emission

filters (Chroma), respectively, for 120 seconds with 100 ms exposure (per channel).

Antibody staining for flow cytometry

To measure cell-surface protein levels (i.e. LAMP1, TFRC), iAstrocytes were dissociated with

Accutase for 10 mins at 37 °C, diluted with ice-cold DPBS + 1% BSA, and spun down at 300 x g

for 5 minutes at 4 °C. After aspirating the supernatant, all cells were incubated in 1:20 Fc block

(BD Bioscience cat. no. 564220) diluted in DPBS + 3% BSA for 10 minutes on ice, followed by

the addition of 1:50 primary antibodies against LAMP1 (AF488- or BV605-conjugated;

BioLegend cat. no. 328610 and 328634) for an additional 30 minute incubation on ice in the

dark. iAstrocytes were then washed with ice-cold DPBS + 1% BSA, spun down at 300 x g for 5

mins at 4 °C, and resuspended in DPBS + 1% BSA for flow cytometry. When astrocytes were

not transduced with sgRNAs, DAPI (ThermoFisher Scientific cat. no. D1306) was added at 0.4

mg/mL to assess cell viability; when astrocytes were transduced with sgRNAs, TO-PRO-3

(ThermoFisher Scientific cat. no. T3605) was added at 1 µM.

For intracellular staining, dissociated iAstrocytes were fixed with 2% paraformaldehyde for 10

min at RT, washed twice with DPBS + 0.5% Tween 20 (Millipore Sigma cat. no. P9461),

incubated with unconjugated primary antibodies against C3 (1:500; ThermoFisher Scientific cat.

no. PA1-29715), VCAM1 (1:250; BioLegend cat. no. 305802), phospho-S6 (1:200; Cell

Signaling Technology cat. no. 2211), pospho-4E-BP1 (1:200; Cell Signaling Technology cat. no.

2855), or IL-32 (1:250; ProteinTech cat. no. 11079-1-AP) for 20 min at RT, washed with DPBS +

0.5% Tween 20, incubated with conjugated secondary antibodies (1:1000, Donkey anti mouse

IgG AlexaFluor 488 conjugated, ThermoFisher Scientific cat. no. A-21202; 1:1000 Donkey anti

rabbit IgG AlexaFluor 555 conjugated, ThermoFisher Scientific cat. no. A-31572; 1:1000

Donkey anti goat IgG AlexaFluor 647 conjugated, ThermoFisher Scientific cat. no. A-21447) for

20 min at RT, washed with DPBS + 0.5% Tween 20, and then resuspended in DPBS + 0.5%

Tween 20 for flow cytometry.

Pooled CRISPRi screening

CRISPRi screens were conducted as described in Leng *et al*. (9). Both LysoTracker and LAMP1

screens were conducted with the H1 (i.e. "druggable genome") sgRNA sub-library from our

next-generation CRISPRi library (37). We generated lentivirus for the pooled H1 sgRNA library

as previously described (81). For each screen (i.e. LysoTracker or cell-surface LAMP1, vehicle

or ITC), iAstrocytes were plated at 20,000 cells/cm^2 onto 4 Matrigel-coated 15-cm dishes,

transduced with the H1 lentiviral sgRNA library with >70% transduction efficiency, treated with

vehicle control or IL-1α+TNF+C1q for 24 hours, stained for LysoTracker or cell-surface

LAMP1, and then sorted into LysoTracker high vs. low or cell-surface LAMP1 high vs. low (top

and bottom 35% of cells on fluorescence histogram) populations. iAstrocytes were sorted with a

BD FACSAria Fusion cell sorter at 5,000-10,000 events per second, and then pelleted via

centrifugation at 300 x g for 10 minutes for genomic DNA extraction. sgRNA abundances were

then measured using next-generation sequencing as previously described (81). LysoTracker

screens were performed with two experimental replicates per condition, while LAMP1 screens

were done with one replicate. Results from all CRISPRi screens in this study can be interactively

explored at www.crisprbrain.org (85).

Extracellular vesicle isolation

For all experiments involving isolation of extracellular vesicles (EVs) from astrocyte conditioned

media, astrocytes were treated with ScienCell Astrocyte media made using exosome-depleted

FBS (ThermoFisher Scientific cat. no. A2720803). Extracellular vesicles were isolated from 10

mL of condition media from iAstrocytes plated onto 10 cm dishes using differential

ultracentrifugation as described in Patel *et al.* (86). All spins were done at 4 °C. Briefly,

conditioned media was spun down at 300 g for 10 min, then the supernatant was spun down at

2000 g for 30 min, then the supernatant was spun down at 15,600 g for 30 min, then the

supernatant was spun down at 120,000 g for 2 hours, and finally the pellet (not visible) was

resuspended in 100 µL of DPBS. The spins at 15,600 g and 120,000 g were done using a SW 41

Ti rotor with 13.2 mL UltraClear tubes (Beckman Coulter cat. no. 344059) in a Beckman Coulter

Optima XE ultracentrifuge. To increase exosome yield for western blotting against certain targets

shown in Fig. 5a (Hsc70, Caveolin-1, IL-32), 10 nM bafilomycin A1 was added together with

vehicle control or ITC. For IL-32, bands were present on blots using samples treated with or

without bafilomycin A1 (see raw blot image files in Supplementary File 1).

Measurement of EV size distribution

The size distribution of EVs isolated above was measured via nanoparticle tracking analysis

using the NanoSight 3000 following the manufacturer's instructions. 100 µL of each resuspended

EV sample was diluted to final volume of 500 µL for analysis on the Nanosight 3000.

Western blots

iAstrocytes were dissociated with Accutase for 10 mins at 37°C, spun down at 300 x g for 5

mins, washed with ice-cold DPBS, lysed in RIPA buffer (Thermo Fisher Scientific, cat. no.

89900) with protease (Roche, cat. no. 04693159001) and phosphatase inhibitors (Sigma-Aldrich,

cat. no. PHOSS-RO), and incubated on ice for 45 minutes with brief vortexing every 15 minutes.

Samples were centrifuged at 18,000 x g for 20 mins at 4°C to pellet insoluble components of

lysate. After the total protein was quantified using the Pierce BCA Protein Assay Kit

(ThermoFisher Scientific, cat. no. 23225) in the supernatant, each sample was diluted to the same

concentration, and denatured with NuPAGE LDS Sample Buffer (4X) (ThermoFisher Scientific,

cat. no. NP0007) and reduced with DTT.

Equivalent protein amounts were run on 10% NuPAGE Bis-Tris gels (ThermoFisher Scientific,

cat. no. NP0301) and transferred to nitrocellulose membranes (Bio-Rad, 1704271). For blots

shown in Fig. 1f-h, Fig. 1l, and Fig. 3g, 20-40 μg of protein was loaded; for blots shown in Fig.

5a,c, 1-2 μg of protein was loaded. We performed ponceau staining (Milipore Sigma, cat. no.

P7170) to validate equivalent loading between lanes. Membranes were blocked with Intercept

blocking buffer (LI-COR cat. no. 927-60001) on a shaker, after which they were incubated with

primary antibodies against mTOR (1:1000, Cell Signaling Technology cat. no. 2972), phospho-

S6 (1:500; Cell Signaling Technology cat. no. 2211), S6 (1:1000; Cell Signaling Technology cat.

no. 2217), phospho-ULK1 (1:500; Cell Signaling Technology cat. no. 14202), phospho-Akt

(1:500; Cell Signaling Technology cat. no. 4060), pan-Akt (1:1000; Cell Signaling Technology

cat. 2920), phospho-4E-BP1 (1:1000; Cell Signaling Technology cat. no. 2855), LAMP1

(1:1000; Abcam cat. no. ab25630), LAMP2 (1:1000; Abcam cat. no. ab25631), CTSD (1 ug/mL,

R&D Systems cat. no. AF1014), β-actin (1:1000; Cell Signaling Technology cat. no. 4970),

EGFP (1:200; Santa Cruz Biotechnology cat. no. sc-9996), LC3 (1:1000; Cell Signaling

Technology cat. no. 2775), CD81 (1:500; BioLegend cat. no. 349502), CD63 (1:5680;

BioLegend cat. no. 353039), Apo-AI (1:200; Santa Cruz Biotechnology cat. no. sc-376818),

GAPDH (1:200; Santa Cruz Biotechnology cat. no. sc-47724), Hsc70 (1:200; Santa Cruz

Biotechnology cat. no. sc-7298), TSG101 (1:500; BioLegend cat. no. 934301), ALIX (1:500;

BioLegend cat. no. 634501), or Caveolin-1 (1:1000; Cell Signaling Technology cat. no. 3267) at

4°C overnight on a shaker. Blots were washed with TBS and incubated with secondary

antibodies goat anti-mouse IRDye 800CW (1:10,000; LI-COR cat. no. 926-32210) and goat anti-

rabbit IRDye 680RD (1:10,000; LI-COR cat. no. 926-68071) in blocking buffer (see above) for 1

hour at RT on a shaker. Afterwards, blots were washed with TBS, and imaged using an Odyssey

Fc Imaging system (LI-COR, cat. no. 2800). To re-probe blots, we incubated membranes with

1X NewBlot Nitro Stripping Buffer (LI-COR, cat. no. 928-40030) for 5 minutes at RT on a

shaker before re-blocking and probing as above. Immunoblot bands were quantified with LI-

COR ImageStudio software. For blotting against CD63 and CD81, reducing agent was omitted

from the sample buffer since the antibodies used here recognize epitopes containing disulfide

bonds.

Source data for immunoblots is provided in Supplementary File 1.

mito-EV staining for flow cytometry

Conditioned media from iAstrocytes was spun down at 300 g to remove dead cells and then

incubated with 10 µg/mL Hoechst, 5 nM TMRM (ThermoFisher Scientific cat. no. T668), and 50

nM MitoTracker Green (ThermoFisher Scientific cat. no. M7514) for 5 min at 37 °C. After

incubation, the conditioned media was analyzed directly with flow cytometry.

IL-32 or CTSB immunoassays

IL-32 concentration was measured using an ELISA kit (R&D Systems cat no. DY3040-05)

following the manufacturer's instructions. For measurement of IL-32 concentration in iAstrocyte

conditioned media, the conditioned media was first spun down at 300 g to remove dead cells,

then 50 µL of conditioned media + 25 µL of RIPA buffer + 25 µL of 1% BSA in DPBS was

added per well, with the same fraction of RIPA and diluent added to the IL32 standards. For

measurement of IL-32 concentration in the exosome fraction vs. supernatant, 75 µL of sample +

25 µL of RIPA buffer was added per well, with the same fraction of RIPA and diluent added to

the IL32 standards. CTSB concentration was measured via an electrochemiluminescence

immunoassay (Mesoscale Discoveries cat. no. K151ACAR-2) following the manufacturer's

instructions.

Whole-cell proteomics sample preparation

iAstrocytes were washed and collected (detached gently using a cell scraper) in ice-cold DPBS,

spun down at 300 x g for 5 mins at 4 °C, and stored at -80 °C. Cell pellets were lysed in 2.5%

sodium dodecyl sulfate (SDS) 50mM Tris pH 8 and heated to 95 °C for 5min, then subjected to

probe sonication. Lysates were centrifuged at 14000g for 15 min before protein quantification

using BCA assay (Pierce, Thermo Fisher Scientific) according to manufacturer's instructions.

Protein lysates were acetone precipitated by adding 5 volumes of ice-cold acetone, vortexed,

incubated at -20 °C for 1 hr and pelleted by centrifugation at 21000g and 4 °C for 15 min.

Protein pellets were resuspended in 1% sodium deoxycholate (SDC) 50 mM EPPS pH 8.5 and

reduced by incubating at 37 °C for 20 min with rotation in the presence of 5mM DTT. Protein

was then alkylated by incubating in the dark at 37 °C for 20 min in the presence of 25mM

iodoacetamide before digestion overnight with LysC at an enzyme-to-protein ratio of 1:50

followed by 3 hr digestion with trypsin at an enzyme-to-protein ratio of 1:50. Digestion was

stopped by acidification to 1% trifluoroacetic acid (TFA) and placed on ice for 10min to promote

precipitation of SDC. The SDC was then pelleted and removed by centrifugation at 21000g for

10min. The supernatant containing acidified peptides were then de-salted on in-house made

Stagetips (87) packed with polystyrene divinyl benzene reverse phase sulphonate solid phase

extraction material (CDS Empore, Fisher Scientific) as previously published (87). Stagetips were

prepared by activating with 100% methanol, conditioning with 80% acetonitrile containing 0.1%

TFA, equilibrated with 0.2% TFA, followed by sample loading, washing twice with 99%

isopropanol containing 0.1% TFA, washing twice with 0.2% TFA and once with 0.1% FA.

Peptides were eluted with 60% acetonitrile (ACN) containing 0.5% ammonium hydroxide. The

desalted peptides were then flash frozen and dried by centrifugal evaporation.

Endolysosomal proteomics sample preparation

pLBR09 (gfaABC1D: LAMP1-XTEN80-mEGFP-3XHA) used for lysosome

immunoprecipitation was generated by subcloning pLBR08 (EF1α: LAMP1-XTEN80-mEGFP-

3XHA). Briefly, pLBR08 was generated via Gibson assembly of PCR-amplified LAMP1 cDNA

(from mTagRFP-T-Lysosomes-20 acquired from Nikon Imaging Center at UCSF, Addgene

plasmid #58022) and PCR-amplified XTEN80-mEGFP-3xHA immunoprecipitation tag with a

linearized backbone generated from ClaI and BspDI (New England BioLabs cat. no. R0557)

digestion of pKL017 (pHIV EF1a:Clover:WPRE). pLBR09 was then generated by linearizing

pKL020 (pHIV gfaABC1D:mRuby:WPRE) and excising the LAMP1-XTEN80-mEGFP-3XHA

cassette of pLBR08 via restriction digest with XbaI (New England Biolabs cat. no. R0145) and

ClaI. The resulting fragments were run on a 1% agarose gel and gel extracted (Zymoclean Gel

DNA Recovery Kit, cat. no. D4007). These fragments were ligated together with T4 DNA Ligase

per manufacturer protocol. iAstrocytes were plated at 20,000 cells/cm^2 per experimental replicate

and transduced with pLBR09 lentivirus during plating. We used three replicates each for vehicle

and ITC conditions.

Lysosome immunoprecipitation was performed as previously described (88). Briefly, iAstrocytes

were washed and collected (detached gently using a cell scraper) in ice-cold DPBS, spun down at

300 x g for 5 mins at 4 °C. We then resuspended the cell pellet in homogenization buffer (25 mM

Tris HCl pH 7.5, 50 mM sucrose, 0.5 mM MgCl$_2$, 0.2 mM EGTA with protease inhibitors). The

iAstrocyte cell suspension was mechanically lysed with a 23G syringe, immediately transferred

to isotonic buffer (2.5M sucrose, 0.2 mM EGTA, 0.5 mM MgCl$_2$) and spun at 3000 x g for 10

mins at 4 °C. The supernatant was transferred to anti-HA magnetic beads (Thermo Scientific cat.

no. PI88836) for the pulldown of intact lysosomes, with the final resuspension in 1X urea buffer

(2M Urea, 1 mM DTT, 12.5 mM Tris). On-bead digest was performed by sequentially incubating

the resulting protein samples with 5 mM (final concentration) IAA at 22°C and 225 rpm for 30

mins, 0.5 µg LysC at 22°C and 225 rpm overnight, 1 µg of trypsin at 22°C and 225 rpm for 4

hours, and 1% (final concentration) TFA. Acidified peptides were then de-salted according to the

same protocol as whole-cell proteome samples. For western blotting, NuPAGE LDS Sample

Buffer (4X) was added to samples collected prior to the on-bead digestion. See Western blots

section for further details.

LC-MS Analysis

Peptides were resuspended in 2% ACN with 0.1% TFA before loading onto a 25 cm x 75 µm ID,

1.6 µm C18 column (IonOpticks) maintained at 40°C. Peptides were separated with an EASY-

nLC 1200 system (Thermo Fisher Scientific, San Jose, CA) at a flow rate of 300 nL min^{-1} using a

binary buffer system of 0.1% FA (buffer A) and 80% acetonitrile with 0.1% FA (buffer B) in a

two-step gradient. The following was performed for whole cell proteome samples, 3% to 27% B

in 105 min and from 27% to 40% B in 15min. For lysosome-specific samples, 3% to 27% B in

52.5 min and from 27% to 40% B in 14.5 min. All samples were analyzed on a Fusion Lumos

mass spectrometer (Thermo Fisher Scientific, San Jose, CA) equipped with a nanoFlex ESI

source operated at 1550 volts, RF lens set to 30%, operated in data dependent acquisition mode

with a duty cycle time of 1 sec. Full MS scans were acquired with a m/z scan range of 375-1500

m/z in the Orbitrap mass analyzer (FTMS) with a resolution of 240k for whole cell proteome and

120k for lysosome-specific samples. Selected precursor ions were subjected to fragmentation

using higher-energy collisional dissociation (HCD) with a quadrupole isolation window of 0.7

m/z, and normalized collision energy of 31%. HCD fragments were analyzed in the Ion Trap

mass analyzer (ITMS) set to Turbo scan rate. Fragmented ions were dynamically excluded from

further selection for a period of 45 sec for whole cell proteome and 60 sec for lysosome-specific

samples. The AGC target was set to 1,000,000 and 10,000 for full FTMS and ITMS scans,

respectively. The maximum injection time was set to Auto for both full FTMS and ITMS scans.

Drug treatments

Vacuolin-1 (Milipore Sigma, cat. no. 673000), bafilomycin A1 (Milipore Sigma, cat. no. B1793),

rapamycin (Milipore Sigma, cat. no. 553210), PP242 (Milipore Sigma, cat. no. 475988),

RapaLink-1 (MedChemExpress cat. no. HY-111373), PI4KIII beta inhibitor 3

(MedChemExpress cat. no. HY-15679), and PI-273 (MedChemExpress cat. no. HY-103489)

were resuspended in DMSO per manufacturer's instructions, aliquoted, and stored at -80°C. All

compounds were added to iAstrocytes alongside vehicle or ITC treatments (i.e. 24 hour

treatments) with the exception of using bafilomycin A1 for measurement of autophagic flux,

which was added to astrocytes for 4 hours before sample collection.

Data analysis

1001

1002   *Analysis of published RNA-seq datasets*

1003   We used BioJupies (89) to reanalyze bulk RNA-seq data and obtain differentially expressed

1004   genes from Perriot *et al*. (90) (GSE120411; hiPSC-derived astrocytes treated with IL1β and

1005   TNF) and Barbar *et al*. (19) (syn21861229; CD49f+ astrocytes sorted from cerebral organoids

1006   treated with vehicle control or IL-1α+TNF+C1q).

1007

Overlap analysis of differentially expressed genes (DEGs)

Human or rodent astrocyte DEGs from various diseases or disease models were extracted

manually from the supplementary information provided on the respective publications or shared

by the corresponding authors upon request; see Supplementary Table 3 for a list of all

publications and datasets used for the analysis. GO Cellular Component terms relating to the

endolysosomal system and biological pathway terms related to mTOR signaling were

downloaded from Enrichr (91–93). Overlap between astrocyte DEGs and the gene sets of interest

was assessed and visualized using the R package GeneOverlap (v1.38.0); Fisher's exact test *P*

values were adjusted for multiple testing using the Benjamini-Hochberg procedure.

1017

Proteomics

Raw files were searched against the human reference proteome (Swiss-Prot, downloaded on

20^{th} of April 2020, refer to PRIDE upload for full sequence) along with the default common

contaminants list with enzyme specificity set to trypsin within MaxQuant version 2.0.1.0 using

standard settings with match between runs and the MaxLFQ algorithm activated. Whole cell

proteome (raw data file names begin with 'Proteome') and lysosome-specific (raw data file

names begin with 'LAMP1') raw files were set as separate parameter groups to ensure that the

MaxLFQ algorithm was performed only within each sample set. The resulting proteinGroups

text file was then uploaded to Perseus version 1.6.7.0 where all reverse sequences, potential

contaminants and proteins only identified by site were filtered out of the dataset. All protein

abundance values were normalized by Log 2 transformation. The lysosome-specific raw data

files were evaluated separate from whole cell proteomic files. Search results from lysosomal raw

data files were filtered to include proteins that were quantified in at least two out of three

biological replicates within at least the ITC (raw data file names include ITC) or vehicle (raw

data file names include WT) triplicate set. Missing values were then imputed within each

replicate by replacing from the normal distribution using default settings. A two sample students

t-test was performed between the ITC treated and vehicle triplicates, with permutation-based

FDR using default settings. The resulting T-test difference and significance were used to generate

the lysosome-specific volcano plot.

1037

1038 *Pathway enrichment analysis*

1039 We used Enrichr (91–93) to perform enrichment analysis of gene lists.

1040

1041 *CRISPRi screening analysis*

1042 We analyzed next generation sequencing reads generated from our CRISPRi screens as

1043 previously described (81). The "gene score" was calculated as -$\log_{10}(P$ value) times the

1044 phenotype score.

1045

Pseudobulk analysis of snRNA-seq data from Macnair et al.

The processed feature-barcode count matrix from Macnair *et al.* (61) was downloaded from

https://zenodo.org/records/8338963 (94) and loaded into R as a Seurat object.

Seurat::AggregateExpression was then used to combine raw counts across cells from each cell

type for each sample. DESeq2::estimateSizeFactors was then used to normalize the counts, and

then the normalized counts added a pseudocount of 1 followed by transformation to $\log_2$ scale.

IL-32 log-normalized counts were then plotted across samples derived from different multiple

sclerosis lesions. *P* values for differences in IL-32 log-normalized counts were calculated by

linear regression in R using lm() without adjustment for clinical covariates.

1055

Flow cytometry

Data from flow cytometry experiments were analyzed using FlowJo (version 10.7.1). Live cells

were gated by plotting SSC-A vs. FSC-A and then single cells were gated by plotting FSC-H vs.

FSC-A. For experiments involving CRISPRi knockdown, analysis was restricted to sgRNA-

transduced cells (gating on the histogram of BFP fluorescence values). For antibody staining

experiments where median fluorescence intensity (MFI) values were reported, the average MFI

of unstained control samples were subtracted from the MFI of stained samples.

1063

Fluorescence imaging analysis

We used CellProfiler (v3.15) (95) to quantify iAstrocyte immunostaining and neuron apoptosis

(as measured by cleaved caspase 3/7 fluorescent dye) imaging data. For neuron viability,

monochrome images of caspase 3/7 were thresholded ("Threshold" module, "Manual" setting)

with respect to unstained (no caspase 3/7 dye) controls. The caspase 3/7 signal was then

quantified as the total intensity for each image, using the thresholded image as a mask. Similarly,

for iAstrocyte fluorescence microscopy images, the total image intensity of LAMP1, LAMP2,

LC3, and LysoTracker signals were quantified after first thresholding images relative to either a

no-primary antibody control (immunofluorescence—LAMP1, LAMP2, LC3) or no-dye control

(LysoTracker). The total image intensity of LAMP1, LAMP2, and LC3 was then divided by the

total intensity of Hoechst to correct for variation in cell number. All frames collected were

averaged to return a single quantification per well, which we considered as a single data point for

statistical tests.

1077

Statistics and reproducibility

Sample sizes were determined by referencing existing studies in the field. Major findings were

validated using independent samples and orthogonal approaches. Numbers of replicates are listed

in each figure. Randomization was not relevant to our study because no animals or human

subjects were involved. Statistics were computed in RStudio (version 4.0.5). Simple pairwise

tests were conducted using t.test or wilcox.test. For instances involving multiple comparisons,

linear regression using the function lm was used to compute *P* values. For percentage data or

data ranged from 0 to 1, beta regression was used to calculated *P* values using the R package

betareg (v3.1-4), with logit linking function and the bias-corrected maximum likelihood

estimator. Correction for multiple testing was performed where appropriate.

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

REFERENCES

1.   Chung W-S, Clarke LE, Wang GX, Stafford BK, Sher A, Chakraborty C, et al.

Astrocytes mediate synapse elimination through MEGF10 and MERTK pathways.

Nature. 2013 Dec 19;504(7480):394–400.

2.   Morizawa YM, Hirayama Y, Ohno N, Shibata S, Shigetomi E, Sui Y, et al. Reactive
astrocytes function as phagocytes after brain ischemia via ABCA1-mediated pathway.
Nat Commun. 2017 Jun 22;8(1):28.

3.   de Ceglia R, Ledonne A, Litvin DG, Lind BL, Carriero G, Latagliata EC, et al.
Specialized astrocytes mediate glutamatergic gliotransmission in the CNS. Nature. 2023
Oct;622(7981):120–9.

4.   Zhang Z, Chen G, Zhou W, Song A, Xu T, Luo Q, et al. Regulated ATP release from
astrocytes through lysosome exocytosis. Nat Cell Biol. 2007 Aug;9(8):945–53.

5.   Bowser DN, Khakh BS. Two forms of single-vesicle astrocyte exocytosis imaged with
total internal reflection fluorescence microscopy. Proc Natl Acad Sci USA. 2007 Mar
6;104(10):4212–7.

6.   Lee H-G, Lee J-H, Flausino LE, Quintana FJ. Neuroinflammation: An astrocyte
perspective. Sci Transl Med. 2023 Nov 8;15(721):eadi7828.

7.   Patani R, Hardingham GE, Liddelow SA. Functional roles of reactive astrocytes in
neuroinflammation and neurodegeneration. Nat Rev Neurol. 2023 Jul;19(7):395–409.

8.   Hasel P, Rose IVL, Sadick JS, Kim RD, Liddelow SA. Neuroinflammatory astrocyte
subtypes in the mouse brain. Nat Neurosci. 2021 Oct;24(10):1475–87.

9.   Leng K, Rose IVL, Kim H, Xia W, Romero-Fernandez W, Rooney B, et al. CRISPRi
screens in human iPSC-derived astrocytes elucidate regulators of distinct inflammatory
reactive states. Nat Neurosci. 2022 Nov;25(11):1528–42.

10.   Kim H, Leng K, Park J, Sorets AG, Kim S, Shostak A, et al. Reactive astrocytes

transduce inflammation in a blood-brain barrier model through a TNF-STAT3 signaling

axis and secretion of alpha 1-antichymotrypsin. Nat Commun. 2022 Nov 2;13(1):6581.

11.   Klumperman J, Raposo G. The complex ultrastructure of the endolysosomal system. Cold

Spring Harb Perspect Biol. 2014 May 22;6(10):a016857.

12.   Leidal AM, Debnath J. Emerging roles for the autophagy machinery in extracellular

vesicle biogenesis and secretion. FASEB Bioadv. 2021 May;3(5):377–86.

13.   Cadwell K. Crosstalk between autophagy and inflammatory signalling pathways:

balancing defence and homeostasis. Nat Rev Immunol. 2016 Nov;16(11):661–75.

14.   Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, et al.

Neurotoxic reactive astrocytes are induced by activated microglia. Nature. 2017 Jan

26;541(7638):481–7.

15.   Hamzaoui K, Borhani-Haghighi A, Dhifallah IB, Hamzaoui A. Elevated levels of IL-32

in cerebrospinal fluid of neuro-Behcet disease: Correlation with NLRP3 inflammasome. J

Neuroimmunol. 2022 Apr 15;365:577820.

16.   Aass KR, Kastnes MH, Standal T. Molecular interactions and functions of IL-32. J

Leukoc Biol. 2021 Jan;109(1):143–59.

17.   Kim S-H, Han S-Y, Azam T, Yoon D-Y, Dinarello CA. Interleukin-32: a cytokine and

inducer of TNFalpha. Immunity. 2005 Jan;22(1):131–42.

18.   Li X, Tao Y, Bradley R, Du Z, Tao Y, Kong L, et al. Fast Generation of Functional

1146   Subtype Astrocytes from Human Pluripotent Stem Cells. Stem Cell Reports. 2018 Oct

1147   9;11(4):998–1008.

1148   19.   Barbar L, Jain T, Zimmer M, Kruglikov I, Sadick JS, Wang M, et al. CD49f Is a Novel

1149   Marker of Functional and Reactive Human iPSC-Derived Astrocytes. Neuron. 2020 Aug

1150   5;107(3):436-453.e12.

1151   20.   Perriot S, Canales M, Mathias A, Du Pasquier R. Differentiation of functional astrocytes

1152   from human-induced pluripotent stem cells in chemically defined media. STAR

1153   Protocols. 2021 Dec 17;2(4):100902.

1154   21.   Krencik R, Zhang S-C. Directed differentiation of functional astroglial subtypes from

1155   human pluripotent stem cells. Nat Protoc. 2011 Oct 13;6(11):1710–7.

1156   22.   Tcw J, Wang M, Pimenova AA, Bowles KR, Hartley BJ, Lacin E, et al. An Efficient

1157   Platform for Astrocyte Differentiation from Human Induced Pluripotent Stem Cells. Stem

1158   Cell Reports. 2017 Aug 8;9(2):600–14.

1159   23.   Cheng X-T, Xie Y-X, Zhou B, Huang N, Farfel-Becker T, Sheng Z-H. Revisiting

1160   LAMP1 as a marker for degradative autophagy-lysosomal organelles in the nervous

1161   system. Autophagy. 2018 Jul 23;14(8):1472–4.

1162   24.   Chin MY, Patwardhan AR, Ang K-H, Wang AL, Alquezar C, Welch M, et al.

1163   Genetically Encoded, pH-Sensitive mTFP1 Biosensor for Probing Lysosomal pH. ACS

1164   Sens. 2021 Jun 25;6(6):2168–80.

1165   25.   Kaizuka T, Morishita H, Hama Y, Tsukamoto S, Matsui T, Toyota Y, et al. An

1166   Autophagic Flux Probe that Releases an Internal Control. Mol Cell. 2016 Nov

1167   17;64(4):835–49.

1168   26.   Yoshii SR, Mizushima N. Monitoring and measuring autophagy. Int J Mol Sci. 2017 Aug

1169   28;18(9).

1170   27.   Coffey JW, De Duve C. Digestive activity of lysosomes. I. The digestion of proteins by

1171   extracts of rat liver lysosomes. J Biol Chem. 1968 Jun 25;243(12):3255–63.

1172   28.   Abrahamson M, Barrett AJ, Salvesen G, Grubb A. Isolation of six cysteine proteinase

1173   inhibitors from human urine. Their physicochemical and enzyme kinetic properties and

1174   concentrations in biological fluids. J Biol Chem. 1986 Aug 25;261(24):11282–9.

1175   29.   Izumi T. In vivo Roles of Rab27 and Its Effectors in Exocytosis. Cell Struct Funct. 2021

1176   Nov 6;46(2):79–94.

1177   30.   Rong Y, Ji C, Wang Z, Ge X, Wang J, Ye W, et al. Small extracellular vesicles

1178   encapsulating CCL2 from activated astrocytes induce microglial activation and neuronal

1179   apoptosis after traumatic spinal cord injury. J Neuroinflammation. 2021 Sep

1180   12;18(1):196.

1181   31.   Alter G, Malenfant JM, Altfeld M. CD107a as a functional marker for the identification

1182   of natural killer cell activity. J Immunol Methods. 2004 Nov;294(1–2):15–22.

1183   32.   Andrews NW. Detection of lysosomal exocytosis by surface exposure of lamp1 luminal

1184   epitopes. Methods Mol Biol. 2017;1594:205–11.

1185   33.   Mauvezin C, Neufeld TP. Bafilomycin A1 disrupts autophagic flux by inhibiting both V-

1186   ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-

1187   lysosome fusion. Autophagy. 2015;11(8):1437–8.

1188   34.   Wang X, Berkowicz A, King K, Menta B, Gabrielli AP, Novikova L, et al.

1189   Pharmacologic enrichment of exosome yields and mitochondrial cargo. Mitochondrion.

1190   2022 May;64:136–44.

1191   35.   Debnath J, Leidal AM. Secretory autophagy during lysosome inhibition (SALI).

1192   Autophagy. 2022 Oct;18(10):2498–9.

1193   36.   Solvik TA, Nguyen TA, Tony Lin Y-H, Marsh T, Huang EJ, Wiita AP, et al. Secretory

1194   autophagy maintains proteostasis upon lysosome inhibition. J Cell Biol. 2022 Jun

1195   6;221(6).

1196   37.   Horlbeck MA, Gilbert LA, Villalta JE, Adamson B, Pak RA, Chen Y, et al. Compact and

1197   highly active next-generation libraries for CRISPR-mediated gene repression and

1198   activation. eLife. 2016 Sep 23;5.

1199   38.   Kim J, Kundu M, Viollet B, Guan K-L. AMPK and mTOR regulate autophagy through

1200   direct phosphorylation of Ulk1. Nat Cell Biol. 2011 Feb;13(2):132–41.

1201   39.   Jung CH, Ro S-H, Cao J, Otto NM, Kim D-H. mTOR regulation of autophagy. FEBS

1202   Lett. 2010 Apr 2;584(7):1287–95.

1203   40.   Szwed A, Kim E, Jacinto E. Regulation and metabolic functions of mTORC1 and

1204   mTORC2. Physiol Rev. 2021 Jul 1;101(3):1371–426.

1205   41.   Hay N, Sonenberg N. Upstream and downstream of mTOR. Genes Dev. 2004 Aug

1206   15;18(16):1926–45.

1207   42.   Sarbassov DD, Guertin DA, Ali SM, Sabatini DM. Phosphorylation and regulation of

1208   Akt/PKB by the rictor-mTOR complex. Science. 2005 Feb 18;307(5712):1098–101.

1209   43.   Clark IC, Gutiérrez-Vázquez C, Wheeler MA, Li Z, Rothhammer V, Linnerbauer M, et

1210   al. Barcoded viral tracing of single-cell interactions in central nervous system

1211   inflammation. Science. 2021 Apr 23;372(6540).

1212   44.   Rodrik-Outmezguine VS, Okaniwa M, Yao Z, Novotny CJ, McWhirter C, Banaji A, et al.

1213   Overcoming mTOR resistance mutations with a new-generation mTOR inhibitor. Nature.

1214   2016 Jun 9;534(7606):272–6.

1215   45.   Wullschleger S, Loewith R, Hall MN. TOR signaling in growth and metabolism. Cell.

1216   2006 Feb 10;124(3):471–84.

1217   46.   Verderio C, Cagnoli C, Bergami M, Francolini M, Schenk U, Colombo A, et al. TI-

1218   VAMP/VAMP7 is the SNARE of secretory lysosomes contributing to ATP secretion

1219   from astrocytes. Biol Cell. 2012 Apr 1;104(4):213–28.

1220   47.   Zahoor M, Westhrin M, Aass KR, Moen SH, Misund K, Psonka-Antonczyk KM, et al.

1221   Hypoxia promotes IL-32 expression in myeloma cells, and high expression is associated

1222   with poor survival and bone loss. Blood Adv. 2017 Dec 26;1(27):2656–66.

1223   48.   Hayakawa K, Esposito E, Wang X, Terasaki Y, Liu Y, Xing C, et al. Transfer of

1224   mitochondria from astrocytes to neurons after stroke. Nature. 2016 Jul

1225   28;535(7613):551–5.

1226   49.   Pan K-H, Chang H, Yang WY. Extracellular release in the quality control of the

1227   mammalian mitochondria. J Biomed Sci. 2023 Oct 8;30(1):85.

1228   50.   Hasegawa H, Thomas HJ, Schooley K, Born TL. Native IL-32 is released from intestinal

1229   epithelial cells via a non-classical secretory pathway as a membrane-associated protein.

1230   Cytokine. 2011 Jan;53(1):74–83.

1231   51.   Sun Y, Qian Y, Chen C, Wang H, Zhou X, Zhai W, et al. Extracellular vesicle IL-32

1232   promotes the M2 macrophage polarization and metastasis of esophageal squamous cell

1233   carcinoma via FAK/STAT3 pathway. J Exp Clin Cancer Res. 2022 Apr 15;41(1):145.

1234   52.   Welsh JA, Goberdhan DCI, O'Driscoll L, Buzas EI, Blenkiron C, Bussolati B, et al.

1235   Minimal information for studies of extracellular vesicles (MISEV2023): From basic to

1236   advanced approaches. J Extracell Vesicles. 2024 Feb;13(2):e12404.

1237   53.   Mathieu M, Névo N, Jouve M, Valenzuela JI, Maurin M, Verweij FJ, et al. Specificities

1238   of exosome versus small ectosome secretion revealed by live intracellular tracking of

1239   CD63 and CD9. Nat Commun. 2021 Jul 19;12(1):4389.

1240   54.   Bebelman MP, Crudden C, Snieder B, Thanou E, Langedijk CJM, Viola M, et al.

1241   Luminescence-based screening for extracellular vesicle release modulators reveals a role

1242   for PI4KIIIβ in exosome biogenesis upon lysosome inhibition. BioRxiv. 2023 Feb 23;

1243   55.   Ostrowski M, Carmo NB, Krumeich S, Fanget I, Raposo G, Savina A, et al. Rab27a and

1244   Rab27b control different steps of the exosome secretion pathway. Nat Cell Biol. 2010

1245   Jan;12(1):19–30; sup pp 1.

1246   56.   WO2013034738A1 - Autoimmune and inflammatory disorder therapy   - Google

1247   Patents [Internet]. [cited 2024 Mar 23]. Available from:

1248   https://patents.google.com/patent/WO2013034738A1

1249   57.   Li J, Gao Z, Zhao D, Zhang L, Qiao X, Zhao Y, et al. PI-273, a Substrate-Competitive,

1250   Specific Small-Molecule Inhibitor of PI4KIIα, Inhibits the Growth of Breast Cancer

1251   Cells. Cancer Res. 2017 Nov 15;77(22):6253–66.

1252   58.   Li Y, Xie J, Xu X, Liu L, Wan Y, Liu Y, et al. Inducible interleukin 32 (IL-32) exerts

1253   extensive antiviral function via selective stimulation of interferon λ1 (IFN-λ1). J Biol

1254   Chem. 2013 Jul 19;288(29):20927–41.

1255   59.   Li W, Sun W, Liu L, Yang F, Li Y, Chen Y, et al. IL-32: a host proinflammatory factor

1256   against influenza viral replication is upregulated by aberrant epigenetic modifications

1257   during influenza A virus infection. J Immunol. 2010 Nov 1;185(9):5056–65.

1258   60.   Kim DH, Park E-S, Lee AR, Park S, Park YK, Ahn SH, et al. Intracellular interleukin-

1259   32γ mediates antiviral activity of cytokines against hepatitis B virus. Nat Commun. 2018

1260   Aug 16;9(1):3284.

1261   61.   Macnair W, Calini D, Agirre E, Bryois J, Jaekel S, Kukanja P, et al. Single nuclei

1262   RNAseq stratifies multiple sclerosis patients into three distinct white matter glia

1263   responses. BioRxiv. 2022 Apr 9;

1264   62.   Squair JW, Gautier M, Kathe C, Anderson MA, James ND, Hutson TH, et al.

1265   Confronting false discoveries in single-cell differential expression. Nat Commun. 2021

1266   Sep 28;12(1):5692.

1267   63.   Liu F, McCullough LD. Inflammatory responses in hypoxic ischemic encephalopathy.

1268   Acta Pharmacol Sin. 2013 Sep;34(9):1121–30.

1269   64.   Sanmarco LM, Wheeler MA, Gutiérrez-Vázquez C, Polonio CM, Linnerbauer M, Pinho-

1270   Ribeiro FA, et al. Gut-licensed IFNγ+ NK cells drive LAMP1+TRAIL+ anti-

1271   inflammatory astrocytes. Nature. 2021 Feb;590(7846):473–9.

1272   65.   Cataldo AM, Nixon RA. Enzymatically active lysosomal proteases are associated with

1273   amyloid deposits in Alzheimer brain. Proc Natl Acad Sci USA. 1990 May;87(10):3861–

1274   5.

1275   66.   Sundelöf J, Sundström J, Hansson O, Eriksdotter-Jönhagen M, Giedraitis V, Larsson A,

1276   et al. Higher cathepsin B levels in plasma in Alzheimer's disease compared to healthy

1277   controls. J Alzheimers Dis. 2010;22(4):1223–30.

1278   67.   Batkulwar K, Godbole R, Banarjee R, Kassaar O, Williams RJ, Kulkarni MJ. Advanced

1279   Glycation End Products Modulate Amyloidogenic APP Processing and Tau

Phosphorylation: A Mechanistic Link between Glycation and the Development of

Alzheimer's Disease. ACS Chem Neurosci. 2018 May 16;9(5):988–1000.

68.   Armstrong A, Mattsson N, Appelqvist H, Janefjord C, Sandin L, Agholme L, et al.

Lysosomal network proteins as potential novel CSF biomarkers for Alzheimer's disease.

Neuromolecular Med. 2014 Mar;16(1):150–60.

69.   Hook VYH, Kindy M, Reinheckel T, Peters C, Hook G. Genetic cathepsin B deficiency

reduces beta-amyloid in transgenic mice expressing human wild-type amyloid precursor

protein. Biochem Biophys Res Commun. 2009 Aug 21;386(2):284–8.

70.   Kindy MS, Yu J, Zhu H, El-Amouri SS, Hook V, Hook GR. Deletion of the cathepsin B

gene improves memory deficits in a transgenic ALZHeimer's disease mouse model

expressing AβPP containing the wild-type β-secretase site sequence. J Alzheimers Dis.

2012;29(4):827–40.

71.   Hook G, Yu J, Toneff T, Kindy M, Hook V. Brain pyroglutamate amyloid-β is produced

by cathepsin B and is reduced by the cysteine protease inhibitor E64d, representing a

potential Alzheimer's disease therapeutic. J Alzheimers Dis. 2014;41(1):129–49.

72.   Liang W, Sagar S, Ravindran R, Najor RH, Quiles JM, Chi L, et al. Mitochondria are

secreted in extracellular vesicles when lysosomal function is impaired. Nat Commun.

2023 Aug 18;14(1):5031.

73.   Morsaljahan Z, Rafiei A, Valadan R, Abedini M, Pakseresht M, Khajavi R. Association

1299   between interleukin-32 polymorphism and multiple sclerosis. J Neurol Sci. 2017 Aug

1300   15;379:144–50.

1301   74.   Parray Z, Zargar MH, Asimi R, Dar WR, Yaqoob A, Raina A, et al. Interleukin 32 gene

1302   promoter polymorphism: A genetic risk factor for multiple sclerosis in Kashmiri

1303   population. Gene. 2022 May 25;824:146261.

1304   75.   Goda C, Kanaji T, Kanaji S, Tanaka G, Arima K, Ohno S, et al. Involvement of IL-32 in

1305   activation-induced cell death in T cells. Int Immunol. 2006 Feb;18(2):233–40.

1306   76.   Qi LS, Larson MH, Gilbert LA, Doudna JA, Weissman JS, Arkin AP, et al. Repurposing

1307   CRISPR as an RNA-guided platform for sequence-specific control of gene expression.

1308   Cell. 2013 Feb 28;152(5):1173–83.

1309   77.   Kang J-W, Park YS, Lee DH, Kim MS, Bak Y, Ham SY, et al. Interaction network

1310   mapping among IL-32 isoforms. Biochimie. 2014 Jun;101:248–51.

1311   78.   Konermann S, Lotfy P, Brideau NJ, Oki J, Shokhirev MN, Hsu PD. Transcriptome

1312   Engineering with RNA-Targeting Type VI-D CRISPR Effectors. Cell. 2018 Apr

1313   19;173(3):665-676.e14.

1314   79.   Cho KS, Park SH, Joo SH, Kim S-H, Shin CY. The effects of IL-32 on the inflammatory

1315   activation of cultured rat primary astrocytes. Biochem Biophys Res Commun. 2010 Nov

1316   5;402(1):48–53.

1317   80.   Miyaoka Y, Chan AH, Judge LM, Yoo J, Huang M, Nguyen TD, et al. Isolation of

1318   single-base genome-edited human iPS cells without antibiotic selection. Nat Methods.

1319   2014 Mar;11(3):291–3.

1320   81.   Tian R, Gachechiladze MA, Ludwig CH, Laurie MT, Hong JY, Nathaniel D, et al.

1321   CRISPR Interference-Based Platform for Multimodal Genetic Screens in Human iPSC-

1322   Derived Neurons. Neuron. 2019 Oct 23;104(2):239-255.e12.

1323   82.   Gilbert LA, Horlbeck MA, Adamson B, Villalta JE, Chen Y, Whitehead EH, et al.

1324   Genome-scale CRISPR-mediated control of gene repression and activation. Cell. 2014

1325   Oct 23;159(3):647–61.

1326   83.   Replogle JM, Bonnar JL, Pogson AN, Liem CR, Maier NK, Ding Y, et al. Maximizing

1327   CRISPRi efficacy and accessibility with dual-sgRNA libraries and optimal effectors.

1328   eLife. 2022 Dec 28;11.

1329   84.   Niu J, Yu G, Wang X, Xia W, Wang Y, Hoi KK, et al. Oligodendroglial ring finger

1330   protein Rnf43 is an essential injury-specific regulator of oligodendrocyte maturation.

1331   Neuron. 2021 Oct 6;109(19):3104-3118.e6.

1332   85.   CRISPRbrain | Home [Internet]. [cited 2024 Mar 26]. Available from:

1333   https://www.crisprbrain.org/

1334   86.   Patel GK, Khan MA, Zubair H, Srivastava SK, Khushman M, Singh S, et al.

1335   Comparative analysis of exosome isolation methods using culture supernatant for

1336   optimum yield, purity and downstream applications. Sci Rep. 2019 Mar 29;9(1):5335.

1337   87.   Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-

1338   fractionation and storage of peptides for proteomics using StageTips. Nat Protoc.

1339   2007;2(8):1896–906.

1340   88.   Abu-Remaileh M, Wyant GA, Kim C, Laqtom NN, Abbasi M, Chan SH, et al.

1341   Lysosomal metabolomics reveals V-ATPase- and mTOR-dependent regulation of amino

1342   acid efflux from lysosomes. Science. 2017 Nov 10;358(6364):807–13.

1343   89.   Torre D, Lachmann A, Ma'ayan A. BioJupies: Automated Generation of Interactive

1344   Notebooks for RNA-Seq Data Analysis in the Cloud. Cell Syst. 2018 Nov 28;7(5):556-

1345   561.e3.

1346   90.   Perriot S, Mathias A, Perriard G, Canales M, Jonkmans N, Merienne N, et al. Human

1347   Induced Pluripotent Stem Cell-Derived Astrocytes Are Differentially Activated by

1348   Multiple Sclerosis-Associated Cytokines. Stem Cell Reports. 2018 Nov 13;11(5):1199–

1349   210.

1350   91.   Xie Z, Bailey A, Kuleshov MV, Clarke DJB, Evangelista JE, Jenkins SL, et al. Gene Set

1351   Knowledge Discovery with Enrichr. Curr Protoc. 2021 Mar;1(3):e90.

1352   92.   Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr:

1353   a comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids

1354   Res. 2016 Jul 8;44(W1):W90-7.

1355   93.   Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, et al. Enrichr: interactive

1356   and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics. 2013

1357   Apr 15;14:128.

1358   94.   Macnair W, Williams A. Single nuclei RNAseq stratifies multiple sclerosis patients into

1359   distinct white matter glial responses. Zenodo. 2023;

1360   95.   McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, et al.

1361   CellProfiler 3.0: Next-generation image processing for biology. PLoS Biol. 2018 Jul

1362   3;16(7):e2005970.