Parvalbumin neurons, temporal coding, and cortical noise in complex
scene analysis

Authors: Jian Carlo Nocon[1,2,3,4], Howard J. Gritton[5,6], Nicholas M. James[1,2,3,4], Xue Han[1,2,3,4],
Kamal Sen[1,2,3,4,*]

[1]Neurophotonics Center, Boston University, Boston, Massachusetts, United States of America
02215,

[2]Center for Systems Neuroscience, Boston University, Boston, Massachusetts, United States of
America, 02215

[3]Hearing Research Center, Boston University, Boston, Massachusetts, United States of America,
02215

[4]Department of Biomedical Engineering, Boston University, Boston, Massachusetts, United
States of America, 02215

[5]Department of Comparative Biosciences, University of Illinois, Urbana, Illinois, United States of
America, 61820

[6]Department of Bioengineering, University of Illinois, Urbana, Illinois, United States of America,
61820

[*]Corresponding Author: kamalsen@bu.edu

Classification: Biological Sciences/Neuroscience

Keywords: Parvalbumin, cortical code, temporal code, cortical noise, cocktail party problem,
complex scene analysis

Abstract

Cortical representations underlying many cognitive abilities emerge from underlying circuits with several different cell types. However, cell type-specific contributions to rate and timing-based cortical coding are not well-understood. Here, we investigate the role of parvalbumin (PV) neurons in cortical complex scene analysis. Many complex scenes contain sensory stimuli which are highly dynamic in time and compete with stimuli at other spatial locations. PV neurons play a fundamental role in balancing excitation and inhibition in cortex and sculpting cortical temporal dynamics; yet their specific role in encoding complex scenes via timing-based codes, and the robustness of temporal representations to spatial competition, has not been investigated. Here, we address these questions in auditory cortex using a cocktail party-like paradigm, integrating electrophysiology, optogenetic manipulations, and a family of spike-distance metrics, to dissect PV neurons' contributions towards rate and timing-based coding. We find that suppressing PV neurons degrades cortical discrimination of dynamic sounds in a cocktail party-like setting via changes in rapid temporal modulations in rate and spike timing, over a wide range of time-scales. Our findings suggest that PV neurons play a critical role in enhancing cortical temporal coding and reducing cortical noise, thereby improving cortical representations of dynamic stimuli in complex scenes.

Significance Statement

One impressive example of sensory perception by the brain is its ability to analyze complex scenes, e.g., following what a friend is saying at a party amongst other speakers. Although some humans can solve this problem with relative ease, it remains very difficult for humans with a variety of impairments, e.g., hearing impairments, ADHD, and autism. The brain mechanisms underlying complex scene analysis remain poorly understood. Here, we recorded neural activity in auditory cortex in a complex auditory scene. When we suppressed PV neuron activity in auditory cortex, cortical performance decreased, and the timing of cortical responses was degraded. Our findings suggest that PV neurons improve the brain's ability to analyze complex scenes by enhancing the timing of cortical responses while reducing cortical noise.

Main Text

Introduction

The cerebral cortex is critical for perception, attention, decision-making, memory, and motor output. Understanding cortical circuit mechanisms underlying these functions remains a central problem in systems neuroscience. One line of questioning towards solving this problem has been to investigate the cortical code; specifically, to assess whether cortical coding relies on rate or spike timing[1]. Previous studies have demonstrated both rate and spike timing-based coding in cortex[2, 3, 4]. However, a mechanistic understanding of how cortical circuits implement these codes and on what timescales is still missing. A second line of questioning towards addressing this central problem has been to utilize a combination of anatomy, physiology and optogenetics to interrogate cortical circuits and neuron types[5, 6]. This concerted approach has allowed systems neuroscience to identify key contributions of specific cell types to cortical circuits, including inhibitory neurons (e.g., parvalbumin-expressing (PV), somatostatin-expressing (SOM) and vasoactive intestinal peptide-expressing (VIP) neurons)[5]. However, the specific contributions of these diverse cell types to the cortical code remain unclear.

A potentially powerful strategy for unraveling cell type-specific contributions to cortical coding is to investigate problems where cortical processing is likely to play a central role. An important example of such a problem is complex scene analysis, e.g., recognizing objects in a scene cluttered with multiple objects at different spatial locations. The brain displays an astonishing ability to navigate such complex scenes in everyday settings, an impressive feat yet

to be matched by state-of-the-art machines. The relative contribution of specific cell-types to this
powerful computational ability remains unclear.

PV neurons are a prominent group of inhibitory neurons in cortex. Previous studies have
investigated the role played by PV neurons in the generation of oscillations[7] and spike
synchronization[8]. PV neurons play a fundamental role in balancing excitation and inhibition[9] and
determining receptive field properties in cortex[10, 11, 12]. Optogenetic manipulation of PV neurons
has provided novel insights into cortical responses, network dynamics, and behavior[13, 14, 15, 16, 17,
18]. Specifically, a study by Moore et al.[18] revealed that optogenetic suppression of PV neurons led
to a rapid rebalancing of excitation and inhibition in cortex, with the expected increase in the
activity of excitatory neurons, but a counterintuitive increase in the activity of inhibitory neurons.
As elegantly dissected in the study, this occurs because the suppression of PV neurons leads to
an increase in activity of excitatory neurons, which then drives both excitatory and inhibitory
neurons downstream, rapidly rebalancing cortical activity. This result illuminates a property of
cortical networks consistent with theoretical models but raises another question: does the
suppression of PV neurons impact cortical temporal coding? The biophysical properties of PV
neurons are well-suited for rapid temporal processing[5] and therefore may be essential in the
cortical temporal coding of dynamic stimuli present in complex scenes. This motivates several
open questions: do PV neurons play a critical role in cortical temporal coding of dynamic stimuli?
Are such temporal codes robust to competing stimuli at other locations in space? Here, we
address these questions in auditory cortex using a combination of electrophysiology, optogenetic
suppression of PV neurons, and a family of spike distance metrics[19, 20] to dissect specific
contributions of PV neurons to the cortical code.

The auditory cortex (ACx) is well suited to investigate these issues. It is thought to play a
key role in solving the "cocktail party problem"[21, 22], one of the most impressive examples of
complex scene analysis. Here, we integrate a cocktail party-like paradigm[23] with optogenetic
suppression of PV neurons to investigate the specific contribution of PV neurons to temporal
coding in mouse ACx. We find that suppressing PV neurons degrades discrimination
performance, specifically temporal coding, in ACx, and degrades performance over a wide range
of time-scales. Our results reveal that despite the rebalancing of excitation and inhibition in
cortical networks observed previously, suppression of PV neurons disrupts coding throughout
ACx suggesting an important influence of PV neurons on cortical temporal coding and cortical
noise.

105
106
107 **Results**
108
109
We recorded single units (SUs) and multi-units (MUs) using a multielectrode array with 4
shanks and 32 channels throughout different layers in ACx of PV-Arch transgenic mice (Figures
1A-C and Supplementary Figure 1). We used a semi-automated detection and sorting algorithm
to identify 124 units from n = 9 animals[24, 25]. Of these 124 units, 101 were identified as SUs (e.g.,
Figure 1C) while the remaining 23 were identified as MUs. In the results below, we focus on SUs.
The results for MUs are given in Supplementary Figure 4C. Out of the SUs, 92 were identified as
regular spiking (RS) while the remaining 9 were identified as narrow spiking (NS) based on the
trough-peak interval of their mean waveforms (Supplementary Figure 2). RS and NS units have
been found to correspond to excitatory and inhibitory neurons, respectively, in ACx[11, 26].

We first confirmed that the effects of optogenetic suppression of PV neurons in ACx were
consistent with previous studies (Figure 1E and Supplementary Figure 3). Optogenetic
suppression occurred on 50% of trials, randomly interleaved, throughout the recording sessions.
Suppression was achieved using light output from a 532nm laser that began 50ms prior to the
auditory stimulus and consisted of continuous illuminations that co-terminated with sound offset.

Within a given 800-trial session, optogenetic suppression strength remained constant (2mW,
5mW or 10mW), but was varied across sessions. We found that upon PV suppression, RS units
increased their firing rate during both spontaneous and evoked activity (Figure 1E and
Supplementary Figure 3A), as expected[17, 18]. Different intensities enhanced the firing rate of RS
neurons in a level-dependent manner consistent with previous studies (Supplementary Figure
3Ai). Counter-intuitively, but consistent with the previous study by Moore et al.[18], NS units also
increased their firing activity (Supplementary Figure 3B). As demonstrated by Moore et al.
optogenetic suppression of PV neurons also causes an increase in inhibition and a rapid
rebalancing of excitation and inhibition in cortex. Thus, the effects of optogenetic suppression of
PV neurons on firing rates in ACx we observed are consistent with previous studies and the rapid
rebalancing of excitation and inhibition. However, the effects of PV suppression on temporal
coding in ACx remain unknown. Thus, we next inquired: does PV suppression impact temporal
coding in ACx?

Investigating cortical coding in mouse ACx using a cocktail party-like paradigm. To
better understand cortical coding of complex scenes in a mouse model amenable to circuit
interrogation using genetic tools, we adopted a cocktail party-like experimental paradigm[23] while
recording from neurons in ACx. Specifically, we recorded responses to spatially-distributed sound
mixtures to determine how competing sound sources influence cortical coding of stimuli. The
recording configuration consisted of 4 speakers arranged around the mouse at 4 locations on the
azimuthal plane: directly in front (0°), two contralateral (45° and 90°) and 1 ipsilateral (-90°) to the
right auditory cortex recording area. Target stimuli consisted of white noise modulated by human
speech envelopes extracted from a speech corpus[27]. We utilized two target waveforms (target 1
and target 2) and a masker consisting of unmodulated white noise. Mice were exposed to either
target-alone trials ('Clean') or target-masker combinations ('Masked') (Figures 2A-C).

**Mouse ACx neurons show spatial configuration sensitivity between competing
auditory stimuli.** We assessed cortical coding using neural discriminability and a variety of other
quantitative response measures. Neural discriminability between the two targets (% correct) was
computed both without the masker ("Clean") (Figure 2A); and with the masker ("Masked"), for all
possible combinations of target and masker locations (Figures 2B-C). Values near 100% and
50% represent perfect discrimination performance and chance, respectively. We refer to the
matrix of performance values from all speaker configurations as "spatial grids", and positions of
high performance ($\geq$ 70%), which were also statistically significant ($p \leq 0.05$) with a relatively
large effect size ($d \geq 1$), as "hotspots". These hotspots represent locations of enhanced
discriminability between the two targets using a spike distance-based classifier to determine how
well target identity can be predicted given the spike train from that site based on dissimilarities in
spike timing and instantaneous rate[23] (Figure 2D: see Methods, 'Neural discriminability using
SPIKE-distance').

Figure 2A illustrates spike trains from an example SU that shows high discriminability
under both target-only conditions (Figures 2A, 2D, and 2E: black); and for a specific spatial
configuration in the presence of a competing noise masker (Figures 2B, 2D, and 2E: red). In the
masked condition, discriminability depends strongly on the spatial configuration of the target and
masker, indicating that the response of this neuron is spatial configuration-sensitive (Figures 2C-
E, red versus green).

Previous studies have demonstrated that neurons with the highest performance are most
strongly correlated with behavior and strongly constrain population performance[28, 29, 30, 31, 32].
Thus, we were curious to test how the performance of the best neurons in our population would
be affected by optogenetic suppression of PV neurons. To do so, we next focused on the neurons
with high discrimination performance in our population (i.e., SUs with at least one hotspot in the

clean or masked conditions). 28 SUs showed hotspots at one or more spatial configurations, and there were 60 hotspots in the clean condition and 28 hotspots in the masked condition giving a total of 88 hotspots.

At each spatial configuration we observed a broad range of performance levels, consisting of neurons with significant hotspots (Figure 2F; filled circles), as well as neurons with poor performance (open circles), reflecting that different neurons in the population had different spatial configuration sensitivities. The "upper envelope" of maximal performance was relatively high for all spatial configurations, except co-located target-masker and ipsilateral target positions. Thus, as a population, ACx neurons showed robust performance at all spatial configurations in the contralateral hemisphere, when the target and masker were spatially separated. We did not observe any statistically significant differences in performance between SUs across different layers, or SUs with different waveform types (RS vs. NS) (Supplementary Figure 4).

Suppression of PV neurons reduces discrimination performance at hotspots. To investigate the role of PV interneurons in auditory discrimination performance, we compared discrimination performance at hotspots, with and without optogenetic suppression of PV neurons in ACx. Figure 3A shows an example SU with and without suppression. Compared to the control response (Figure 3Ai), the optogenetic response (Figure 3Aii) shows an increase in spiking between the peaks of both target stimuli. Specifically, the responses exhibited an earlier onset and decreased spike timing reproducibility during suppression (Figure 3Aiii). Figure 3B shows the spatial grids for the same example SU during both conditions, with the example configuration in Figure 3A (Clean Target 45º) outlined in black. This unit showed a decrease in performance across all clean configurations, and the hotspots in the control masked condition (Target 90º, Masker -90º; Target 45º, Masker -90º; Target 45º, Masker 90º) showed a reduction in performance to below threshold. Overall, we found that performance decreased significantly in both clean ($p = 5.40e-09$) and masked ($p = 6.24e-05$) conditions during suppression (Figure 3C).

Suppression of PV neurons degrades cortical temporal coding. To determine the extent to which changes in the temporal dynamics of rapid firing rate modulation, spike timing, and average firing rate changes that occur during suppression might affect performance, we calculated different performance metrics across all hotspots. Specifically, we used inter-spike interval (ISI)-distance, rate-independent (RI)-SPIKE-distance, and spike count, as the basis for discriminability between spike trains. ISI-distance calculates the distance between two spike trains based on the average dissimilarity in instantaneous firing rate, while RI-SPIKE-distance measures spike timing dissimilarity between two trains while accounting for changes in firing rate differences[19]. Spike count distance is the absolute difference in the number of spikes between trains, effectively measuring differences in total firing rate. We found that performance based on both ISI-distance (Figure 4A) and RI-SPIKE-distance (Figure 4B) performances were relatively high. Both performances showed highly significant decreases with optogenetic suppression (ISI-distance-based performance: $p_{clean} < 1e-04$, $p_{masked} = 4.69e-04$; RI-SPIKE-distance-based performance: $p_{clean} < 1e-04$, $p_{masked} = 6.79e-04$). In contrast, performance based on spike count over the entire stimulus (Figure 4C) was close to chance level both for control and laser conditions, indicating that spike count alone was not sufficient to account for overall performance. The significant decrease in ISI distance-based performance indicates a disruption in rate-based coding, including the dynamics of instantaneous firing rate modulations. The significant decrease in RI-SPIKE based distance indicates that spike timing-based coding is also degraded by optogenetic suppression of PV neurons (Figure 4D).

Effects on components of discrimination performance with suppression. Generally, discrimination performance depends on both the dissimilarity of responses between targets, as well as the similarity of responses within a target. To assess the relationship between different components of responses with performance, we calculated three metrics sensitive to firing rate and/or timing: the average firing rate; the rate-normalized root-mean-square (RMS) difference in the responses to the targets, which captures the difference in the temporal pattern of responses to the targets; and the trial similarity[33], which captures the reproducibility of responses across trials within a target (see Methods).

We first calculated the correlation between evoked firing rate and performance during the control condition by pooling clean and masked data. Firing rate did not show a significant correlation with performance ($r = 0.0827$, $p = 0.4437$), whereas both RMS difference and trial similarity measures were highly correlated with performance (rate-normalized RMS difference: $r = 0.3341$, $p = 0.0015$; trial similarity: $r = 0.5660$, $p < 1e-04$). These results suggest that both the pattern of firing rate modulations (quantified by RMS difference) as well as the reproducibility of responses (quantified by trial similarity) contribute to discrimination performance under control conditions. When compared these measures between the control and laser conditions, we found that rate-normalized RMS difference significantly decreased with optogenetic suppression for both clean and masked trials (Figure 5A), and trial similarity significantly decreased during clean trials (Figure 5B).

Optogenetic suppression decreases performance across a wide range of time scales. The previous analyses used spike distance measures which do not require a choice of a specific time-scale for analysis. A further interesting question regarding discrimination performance is the optimal time-scale for discrimination. Thus, we next quantified the time-scale for optimal discrimination using the van Rossum spike distance measure[34] (see Methods).

We found that optimal time-scales for discrimination (τ) for most neurons was around 40 milliseconds, with a significant proportion of neurons covering even finer time scales down to ~10 ms (Figure 6A). Optimal τ was not significantly different between control and laser conditions ($p_{clean} = 0.4117$; $p_{masked} = 0.2936$), but performance decreased significantly in the laser condition across a wide range of time-scales (Figures 6B-D, Table 1). These results indicate that PV suppression did not significantly change the optimal time-scale for discrimination but rather degraded discrimination across a wide range of time-scales.

Discussion

Diversity of cortical cell types and the cortical code. One of the most striking features of the cerebral cortex is the tremendous diversity of its cell types[35]. Understanding the computational role of such diversity in cortical coding is central to systems neuroscience. Addressing this central question requires understanding cell type-specific contributions to the cortical code at both the single neuron and the population level. A small number of previous studies have demonstrated a role of specific cell types in cortical population coding; specifically, the generation of oscillations[36, 37], and synchrony across cortical layers and areas[38, 39]. However, cell type-specific contributions to the cortical code at the single unit level, a fundamental aspect of cortical encoding, remain poorly understood. In this study, we addressed this fundamental gap by investigating the role of PV neurons in cortical coding of a complex scene, i.e., a cocktail party-like setting, in mouse ACx.

Cortical discrimination & PV neurons; rate, spike timing and temporal codes. We assessed cortical coding using neural discrimination performance and other quantitative measures. There is a rich history of quantitative work on cortical discrimination[28, 30]. These studies have suggested a critical role for neurons with the highest levels of performance in a population, which correlate strongly with behavioral performance and determine the overall performance at the population level. In this study we examined cortical discrimination of dynamic stimuli in a complex scene by the highest performing neurons in ACx, extending the previous body of work in several ways: First, we assessed the impact of optogenetic suppression of PV neurons on discrimination performance. Optogenetically suppressing PV neurons resulted in increased firing rate during spontaneous and evoked activity, which is consistent with the effects of inhibitory blockade on cortical responses[40, 41, 42]. A recent study by Moore et al. employed optogenetic suppression of PV neurons to powerfully reveal an important property of cortical networks: rapid rebalancing of excitation and inhibition upon PV suppression. Our study reveals that despite such rebalancing, cortical discrimination performance is degraded across cortical layers in ACx upon PV suppression. This finding suggests that PV neurons play a role in improving discrimination of dynamic stimuli in ACx, both sounds in quiet backgrounds, as well as in the presence of competing sounds from other spatial locations. Second, we quantified the contributions of instantaneous firing rate modulations, spike timing, and spike count towards cortical discrimination, using a family of spike distance metrics. These metrics provide a powerful set of tools for dissecting different components of cortical coding. Although these metrics have been employed in previous theoretical studies, to our knowledge this is the first time they were applied to analyze cortical responses. This analysis revealed that high discrimination performance is mediated by the temporal pattern of firing rate modulations and spike timing reproducibility, and that optogenetic suppression of PV neurons degraded both components.

Previous studies have demonstrated that auditory cortical neurons can employ both rate and spike timing-based codes[3, 4]; and provided insight into the roles of inhibitory neurons in shaping frequency tuning[10, 11, 12, 43], frequency discrimination[13, 44], adaptation[14], sparseness[45], and gap encoding[26]. An influential review on neural coding also defined a precise notion of a "temporal code" as one that contains information in spike timing beyond rate modulations[46]. Temporal codes have been challenging to identify because the contributions of rate vs. spike timing are often difficult to decouple. Our results based on spike distance measures, which quantify both rate-dependent and rate-independent components of coding, suggest that PV neurons specifically contribute to temporal coding in cortical discrimination. This computational portrait of PV neurons validates the importance of their established electrophysiological specializations–namely, fast, efficient, and temporally precise responses[5].

A general cortical representation for the cocktail party problem across species. From a comparative standpoint, we found that the key features present at the cortical level within ACx of the mammalian mouse was consistent with previous findings in songbirds. Specifically, a previous study by Maddox et al. found hotspots at particular spatial configurations of target and masker on the spatial grids of cortical level neurons in songbirds[23]. Songbirds and mice have different frequency ranges of hearing and therefore the cues used for spatial processing, e.g., interaural time difference (ITD), and interaural level difference (ILD) are frequency-dependent, and the peripheral representations of these cues are likely to be different across species with different frequency ranges of hearing. This suggests the emergence of general cortical representations for solving the cocktail party problem despite different peripheral representations of acoustic cues across species.

Time scales of cortical discrimination. A further interesting question regarding cortical discrimination is: what is the optimal time scale for maximal discrimination performance? One

characteristic time scale in our stimuli arises from the slow modulation of speech envelopes on
relatively long time-scales ~100-500 ms, or equivalently, in the 2-10 Hz frequency range[47]. We
found that the optimal time-scale for most neurons in our dataset was much finer ~40 ms, with a
significant number at even finer timescales down to ~10 ms. These time-scales are well matched
to the duration of short ultrasonic vocalizations in mice (Figure 7), and finer grain structures within
these vocalizations, e.g., spectral features and frequency sweeps[48]. These time-scales are similar
to those found in a previous study of decoding sinusoidally amplitude-modulated (SAM) tones in
mouse auditory cortex[33], and integration time scales in cat auditory cortex[49]. Phonemic structures
in speech also occupy similar time-scales, which are in the beta and low gamma range of
frequencies[50, 51]. Thus, the time-scales for optimal discrimination in ACx, may be well suited for
analyzing such vocalizations and finer spectro-temporal features within.

Cortical noise. Our findings are also relevant in the context of cortical noise, which can
have a profound impact on cortical codes[52]. We found that suppressing PV neurons did not
change the optimal time-scale for discrimination, but rather degraded performance at a wide
range of time-scales (Figure 6). Additionally, we observed that suppression impacted specific
components underlying discrimination: Most notably, PV suppression decreased the difference in
the pattern of responses (quantified by RMS difference) between targets as well as decreasing
the reproducibility of responses across trials (quantified by trial similarity). Taken together, these
observations are consistent with an overall enhancement in cortical noise level across multiple
time-scales upon PV suppression. A previous study on PV suppression in ACx observed the
rapid rebalancing of excitation and inhibition, suggesting maintenance of the stability of global
cortical representations[18]. However, our results suggest that in spite of this excitatory-inhibitory
rebalancing, PV suppression also leads to an increase in cortical noise, fundamentally impacting
the fidelity of cortical coding, including temporal coding,

Within channel vs. cross-channel inhibition. Cortical inhibitory neurons can mediate
feedforward, recurrent and di-synaptic feedback inhibition in cortical circuits, illustrated in a
schematic model in Figure 8. Previous modeling studies have demonstrated that feed-forward
within-channel inhibition can improve discrimination performance[53]; whereas inhibition across
different channels can lead to the formation of hotspots and the specific pattern of spatial
configuration sensitivity[54]. Our results suggest that PV neurons mediate within-channel inhibition,
corresponding to I neurons in the schematic model. This is consistent with our finding that
although suppressing PV neurons reduced discrimination performance, it did not completely
eliminate the presence of hotspots on the spatial grids, suggesting that PV neurons alone do not
control the emergence of hotspots. Based on these observations, we hypothesize that a separate
cell-type (X neurons in Figure 8) mediates cross-channel inhibition, resulting in the generation of
hotspots and the specific pattern of spatial configuration sensitivity on spatial grids. A candidate
cell type that may correspond to X cells are somatostatin-positive (SOM) neurons, which have
been implicated in di-synaptic feedback inhibition[55, 56] and surround suppression[57, 58]. These
distinct roles may be functionally well suited for solving the cocktail party problem, with one class
of neurons (PV) enhancing the temporal coding of dynamic stimuli at a target location, and
another class of inhibitory neurons (X) suppressing competing stimuli from other spatial locations.

Limitations and future directions. Several limitations in this study should be further
addressed in the future. Although we used a cocktail party-like paradigm to probe auditory cortical
responses to dynamic stimuli our experimental paradigm had some limitations. First, the target
stimuli did not have any specific behavioral significance, unlike the case of speech recognition at
a cocktail party. Second, the masker stimuli did not contain any temporal modulations, unlike

competing speakers at a cocktail party. Future studies should address these limitations, e.g., by employing mouse communication sounds as targets and maskers. Although we were able to characterize the time-scale for optimal discrimination in ACx, we did not characterize the integration window, or the encoding window[46, 49, 59]. Future studies that characterize both the time-scale for optimal discrimination as well as the encoding window can address whether cortical neurons also employ "temporal encoding", i.e., encode information in the temporal pattern of spikes within the encoding window[46]. Within this study, mice were awake but listening passively, whereas listening in a cocktail party-type setting is an active sensing process. It will be interesting to probe cortical coding in awake, behaving mice in experiments where animals attend to a specific spatial location. A recent theoretical model of attention in auditory cortex, the AIM network model[60], suggests distinct roles for different interneuron groups in attentional sharpening of both spatial and frequency tuning which enables flexible listening in cocktail party-like settings, e.g., monitoring the entire scene, selecting a speaker at a spatial location, and switching to a speaker at a different location. Future experiments probing distinct interneuron populations (e.g., PV, SOM and VIP neurons) in behaving animals, in conjunction with testing and extending the AIM model may further unravel cortical circuits for solving the cocktail party problem.

Materials and Methods

Subjects. All procedures involving animals were approved by the Boston University Institutional Animal Care and Use Committee (IACUC). A total of 9 mice were used in this study. Original breeding pairs of parvalbumin-Cre (PV-Cre: B6;129P2-Pvalbtm1(cre)Arbr/J), and Ai40 mice (Arch: B6.Cg-*Gt(ROSA)26Sor^tm40.1(CAG-aop3/EGFP)Hze*/J) mice were obtained from Jackson Laboratory (Maine), and all breeding was done in house. Subjects consisted of both male and female PV-Arch offspring 8-12 weeks old on the day of recording.

Surgery. Mice were surgically implanted with a head-plate as described previously[61, 62]. Briefly, under isoflurane anesthesia, stereotaxic surgery was performed to install a headplate, electrode, and optical fiber. The custom head-plate was mounted anterior to bregma allowing access to ACx caudally. The headplate was anchored to the skull with 3 stainless steel screws and dental cement. A fourth screw was connected to a metal pin and placed in the skull above the contralateral cerebellum to serve as the reference. A craniotomy was made above the right auditory cortex (AP -2.3 to -3.6, ML + 4.0 to +4.5, DV). Using a stereotaxic arm, a 32-contact linear probe (Neuronexus, Ann Arbor, MI; model: a 4x8-5mm-100-400-177-CM32) with 100µm spacing between electrode contacts and 400µm spacing between shanks, was positioned into ACx, perpendicular to the cortical surface. Because of the curvature of the ACx surface, not all four shanks could be placed at precisely the same depth during each experiment. Probes were advanced until all electrode contacts were within the cortical tissue and shanks were positioned along the rostro-caudal axis of ACx (Figures 1A-C). An optical fiber, 200µm in diameter, was placed medially to the 4 shanks and positioned between the two innermost shanks terminating at the cortical surface (Figure 1A). After implantation, mice were allowed to recover for 4-7 days before undergoing habituation to being head-fixed as described below.

Habituation. Following surgery and complete recovery, mice were first handled for several days before being head-fixed to the recording apparatus. Mice were gradually exposed to longer restraint periods at the same time of day as subsequent recording sessions. Each animal received at least 6 habituation sessions prior to the first recording day. Under head-fixed conditions, mice were loosely covered with a piece of lab tissue taped down on either side (Kimwipes: Kimberly-Clark, Irving, TX) to encourage reduced movement. At the conclusion of habituation, mice underwent one day of recording sessions.

Recording sessions and data acquisition. All recordings were made with a Tucker Davis Technologies (TDT; Alachua, FL) RZ2 recording system in an electrically-shielded sound attenuation chamber. Broadband neural signals at 24414.0625 Hz were recorded for each of the 32 channels. Local field potentials (LFPs) were band-pass filtered between 1 and 300 Hz, notch-filtered at 60 Hz, and digitized at 3051.8 Hz.

Recording sessions consisted of both non-optogenetic and optogenetic trials in random order. The inter-trial interval was 5 seconds, with 3s of stimulus playback followed by 2s of silence. Mice were exposed to target-alone ("clean") trials and target-masker ("masked") combinations. 10 trials were given per target identity for all possible combinations of target location, masker location (including "clean" trials), and optogenetic suppression of PV neurons. Thus, animals received a total of 800 trials per ~60 minute recording session, with each session having a set laser power. For each animal, all three sessions were recorded on the same day.

Auditory stimuli. All auditory stimuli were generated in Matlab and consisted of either target, masker, or combination of the two stimuli played from four TDT ES-1 electrostatic speakers. Target stimuli consisted of white noise modulated in time by human speech envelopes taken from the Harvard IEEE speech corpus[27] which has been used in previous psychological studies of the cocktail party effect[63]. Masker stimuli consisted of 10 unique tokens of unmodulated white noise. Before speaker calibration, all stimuli were generated with the same RMS value, and sampling frequency was 195312 Hz to capture the frequency range of hearing in mice. Stimuli were loaded onto a custom RPvdsEx circuit on an RZ6 Multi I/O processor, which was connected to two PM2R multiplexers that controlled the location of target and masker stimuli during trials.

During recordings, the stimuli were presented 18 cm from the mouse's head using four speakers driven by two TDT ED-1 speaker drivers. The four speakers were arranged around the mouse at four locations on the azimuthal plane: directly in front (0°), two contralateral (45° and 90°) and 1 ipsilateral (-90°) to the right auditory cortex recording area. Before recording sessions, stimulus intensity was calibrated using a conditioning amplifier and microphone (Brüel and Kjær, Nærum, Denmark; amplifier model: 2690, and microphone model: 4939-A-011). For 7 of the 9 mice, all stimuli were at a measured 75dB intensity at the mouse's head. For the remaining 2 mice, stimulus intensity was set to 70dB. Stimulus playback lasted 3s with a 1ms cosine ramp at onset and offset.

Optogenetic stimulation. Laser light for optogenetic stimulation of auditory cortex was delivered through a multimode optically-shielded 200µm fiber (Thorlabs, Newton, NJ; model: BFH48-200), coupled to a 532nm DPSS laser (Shanghai Laser Ltd., Shanghai, China; model: BL532T3-200FC), with the fiber tip positioned right above the cortical surface. Laser power was calibrated to 2mW, 5mW, or 10mW at the fiber tip using a light meter calibrated for 532nm wavelength (PM100D, Thorlabs, Newton, NJ). The intensity was determined based on optogenetic cortical PV suppression studies using Archaerhodopsin from the literature[13, 64]. During optogenetic trials, the laser was turned on 50ms before stimulus onset and co-terminated with the end of the auditory stimuli (Figure 1D). Square light pulses lasting 3.05s were delivered via TTL trigger from the RZ2 recording system to the laser diode controller (ADR-1805). Optogenetic trials were randomized throughout the recording session such that animals received all stimulus/masker pairs from each location with and without laser. Recordings were done in successive blocks with constant optogenetic suppression strengths of 2mW, 5mW, or 10mW, with each block lasting ~60 minutes and having their own set of control trials. These laser strengths are similar to those used in past studies[13, 17] and did not result in epileptiform activity in cortex.

Spike extraction and clustering. Kilosort2 (https://github.com/MouseLand/Kilosort) was used to detect multi-units within the recordings[24]. Before spike detection and sorting, the broadband signal was band-passed between 300 and 5000 Hz using a 3rd-order Butterworth filter. Kilosort results were then loaded onto Phy2 (https://github.com/cortex-lab/phy), a graphical user interface for examining spike cluster information[25]. Clusters with artifact-like spike waveforms were deemed as noise, while the remaining clusters were deemed as either single or multi-units. The spikes toolbox (https://github.com/cortex-lab/spikes) was then used to import the cluster information from Phy to Matlab and extract spike waveforms from the high-passed signal[24]. Clusters were assigned to recording channels based on which site yielded the largest average spike amplitude. To remove any remaining artifacts from laser onset and offset, all spikes with waveforms above an absolute threshold of 1500 µV or a positive value above 750 µV were discarded. Clusters that still showed a high amount of remaining artifact after removal were excluded from further analysis, and the remaining clusters were deemed single-unit (SU) if less than 5% of their inter-spike intervals were below 2ms (Figure 1C), with the rest deemed multi-units (MU). Finally, units were classified as narrow-spiking if the trough-peak interval of their mean waveform was below 0.5ms, a threshold that is consistent with past findings on excitatory and inhibitory units in mouse auditory cortex[11].

Current source density estimation and layer analysis. Current source density (CSD) analysis estimates the second spatial derivative of LFP signals to determine the relative current across the cortical laminar depth. CSDs were calculated using LFPs, as described previously[62]. LFPs from control masked trials were used, as the rise time was more similar between target identities than clean stimuli. LFPs were low-passed filtered at 150 Hz before being down-sampled by a factor of 8 to 381 Hz. For each channel, LFPs were averaged across all control masked trials prior to CSD estimation, and channels that did not show an evoked response were interpolated using neighboring sites on the same shank. After this, LFPs were spatial smoothed across the eight channels in each shank:

$$\phi(z) = \frac{\phi(z + \Delta z) + 2\phi(z) + \phi(z - \Delta z)}{4}$$

where z is the depth perpendicular to the cortical surface, Δz is the electrode spacing, and Φ is the potential. CSD was then estimated as:

$$CSD(z) = -\frac{\phi(z + \Delta z) - 2\phi(z) + \phi(z - \Delta z)}{\Delta z^2}$$

To determine the granular layer in each shank, CSD sink onset times were calculated as the time when the CSD goes below -3 times the standard deviation of pre-stimulus activity. If more than one channel was found to have the earliest sink onset, the channel whose neighbors had the earliest onsets was deemed the granular layer, or L4. The width of each layer was estimated based on previous anatomical studies[65]. L1 consisted of channels at least 500µm above the input layer, L2/3 consisted of channels 200µm to 400µm above the channel with the earliest sink onset; L4 consisted of the input channel and the channel 100µm above it; L5 consisted of channels 100 to 300µm below the input layer, and L6 consisted of all channels at 400µm below the input layer.

To estimate the effect of optogenetic suppression across layers, repeated-measures ANOVA was carried out for both spontaneous and onset firing rate, with effect sizes calculated using the Measures of Effect Size toolbox[66, 67]. Spontaneous firing rate was defined as the average firing rate during the 50ms between laser onset and sound onset, while onset firing rate was defined as the average rate during the first 0.5s of sound stimulus playback. For both measures, ANOVA was separately for regular-spiking single units, with layer as the between-subjects factor and condition as the within-subjects factor. Post-hoc Tukey-Kramer tests were

carried out if the ANOVA returned a significant factor or interaction. Repeated-measures ANOVA
was not done for narrow-spiking single units due to the small sample size per layer[68].

Neural discriminability performance using SPIKE-distance. Neural discrimination
performance was calculated using a template-matching approach similar to our previous
studies[23]. Briefly, spike trains were classified to one of the two target stimuli based on whose
template, one from each stimulus, yielded a smaller spike distance. For each target-masker
configuration, 100 iterations of template matching were done. In each iteration, one of the 10
spike trains for each target was chosen as a template, and all remaining trials were matched to
each template to determine target identity. All possible pairs of templates were used across the
100 iterations to calculate an average value of neural discriminability. SPIKE-distance[19]
calculates the dissimilarity between two spike trains based on differences in spike timing and
instantaneous firing rate without additional parameters. For one spike train in a pair, the
instantaneous spike timing difference at time t is:

$$S_1(t) = \frac{\Delta t_P^{(1)}(t)x_F^{(1)} + \Delta t_F^{(1)}(t)x_P^{(1)}}{x_{ISI}^{(1)}(t)}, t_P^{(1)} \leq t \leq t_F^{(1)}$$

where Δt_P represents the distance between the preceding spike from train 1 ($t_P^{(1)}$) and the nearest
spike from train 2, Δt_F represents the distance between the following spike from train 1 ($t_F^{(1)}$) and
the nearest spike from train 2, x_F is the absolute difference between t and $t_F^{(1)}$, and x_P is the
absolute difference between t and $t_P^{(1)}$. To calculate $S_2(t)$, the spike timing difference from the
view of the other train, all spike times and ISIs are replaced with the relevant values in train 2.
The pairwise instantaneous difference between the two trains is calculated as:

$$S''(t) = \frac{S_1(t) + S_2(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t)\rangle}$$

532   Finally, $S_1(t)$ and $S_2(t)$ are locally weighted by their instantaneous spike rates to account
533   for differences in firing rate:

$$S(t) = \frac{S_1(t)x_{ISI}^2(t) + S_2(t)x_{ISI}^1(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t)\rangle^2}$$

534   For a train of length T, the distance is the integral of the dissimilarity profile across the entire
535   response interval, with a minimum value of 0 for identical spike trains:

$$D_S = \frac{1}{T}\int_0^T S(t)dt$$

cSPIKE, a toolbox used to calculate SPIKE-distance, was used to calculate all spike train
distances between all possible spike train pairs for all spatial grid configurations[19].

Spatial grid "hotspots" of high neural discriminability were determined using three criteria:
1) mean performance must be above 70% during control trials; 2) mean control performance
distribution must be significantly different from chance ($p < 0.05$), calculated using a null
distribution obtained by classifying spike trains within each target, which should result in chance
performance; and 3) the effect size given by Cohen's d between the two distributions (control vs.
null) must be greater than 1:

$$d = \frac{\bar{x}_1 - \bar{x}_0}{\sqrt{\dfrac{(n_1 - 1)s_1^2 - (n_0 - 1)s_0^2}{n_1 + n_0 - 2}}}$$

where values with subscript 0 represent the mean, standard deviation, and number of template-
matching iterations for the null performance distribution. Additionally, configurations where at
least 3 trials for one target showed zero spiking were excluded from analysis, to avoid inaccurate
estimates of performance. In the manuscript, we focus on SUs with hotspots in the control
condition. We found a small number of "emergent" hotspots (9 hotspots from 8 units) where
performance was below threshold in the control condition but just above threshold in the laser
condition (median performance 72.8% and an inter-quartile range of 3.96%).

To analyze the effects of suppression on performance, we utilized paired t-tests between
control and optogenetic values to determine statistical significance ($p \leq 0.05$), with tests done
separately for clean and masked trials. For comparisons in performance between layers and
between narrow-spiking and regular-spiking units, we separately ran repeated-measures ANOVA
and effect size calculations for clean and masked trials, with condition as the within-subjects
factor and cell type (narrow-spiking or regular-spiking) or layer as the between-subjects factor.

To determine how firing rate modulation, spike timing, and average firing rate contribute
to discriminability, we used different distance measures as inputs to the classifier. For all
hotspots, performances using inter-spike interval (ISI)-distance, rate-independent (RI)-SPIKE-
distance, and spike count distance, the absolute difference in spike count between trains, were
also calculated and compared to SPIKE-distance-based values.

ISI-distance. To determine how optogenetic suppression affects rapid temporal
modulations in firing rate, ISI-distances were calculated. The ISI-distance calculates the
dissimilarity between two spike trains based on differences in instantaneous rate synchrony. For
a given time point:

$$I(t) = \frac{\left| x_{ISI}^{(1)}(t) - x_{ISI}^{(2)}(t) \right|}{\max\left(x_{ISI}^{(1)}(t), x_{ISI}^{(2)}(t) \right)}$$

This profile is then integrated along the spike train length to give a distance value, with values of
0 obtained for either identical spike trains or pairs with the same constant firing rate and a global
phase shift difference.

RI-SPIKE-distance. To determine how optogenetic suppression affects spike timing, RI-
SPIKE-distances between spike trains were calculated. The RI-SPIKE-distance is rate-
independent, as it does not take differences in local firing rate between the two spike trains into
account. From SPIKE-distance calculations, the final step of weighing $S_1(t)$ and $S_2(t)$ by their
instantaneous spike rates is skipped, yielding:

$$S_{1,2}^{RI}(t) = \frac{S_1(t) + S_2(t)}{2\langle x_{ISI}^1(t), x_{ISI}^2(t) \rangle}$$

Like the other measures, the dissimilarity profile is integrated to give a distance value, with a
value of 0 obtained for two identical spike trains.

Rate-normalized RMS difference and trial similarity. In addition to average firing rate,
we also calculated two other measures to determine their impact on classification performance:
the similarity of responses within target, and the dissimilarity of responses across targets. To
quantify intertrial reliability of responses to target stimuli, we adopted the measure of trial
similarity from previous studies[33]. Specifically, we randomly divided the 10 trials in each
configuration into two equal groups, binned spike times with a time resolution of 25ms, and
calculated the Pearson's correlation coefficient between the two resulting PSTHs. This process
was repeated 100 times to obtain a mean correlation coefficient, or trial similarity.

We also calculated the rate-normalized RMS difference between target responses to
quantify the dissimilarity in the temporal pattern of responses between the two targets. We first
binned each target response using the same time-resolution as trial similarity (25ms) and
normalized each PSTH such that the sum of all bins over time was 1. The RMS difference
between the two rate-normalized PSTHs was then calculated. This measure quantifies the
dissimilarity in the temporal pattern of responses across the targets, accounting for differences in
mean evoked firing rate between targets.

All three response measures (average firing rate, trial similarity, and rate-normalized
RMS difference between targets) were correlated with SPIKE-distance-based performance using
Pearson's correlation coefficients, with separate calculations done for control and laser trials.

Decoding time analysis using van Rossum distances. To estimate the decoding time
of the spike trains at each hotspot, we used van Rossum distances[34]. Briefly, the van Rossum
distance between two spike trains involves convolving each response with a decaying
exponential kernel with time constant τ. The distance between two smoothed spike trains $f_1(t)$ and
$f_2(t)$ is calculated as:

$$D_{VR} = \sqrt{\frac{1}{\tau}\int_0^\infty \left(f_1(t) - f_2(t)\right)^2 dt}$$

For each spatial grid configuration, a distance matrix containing the van Rossum distances
between all possible spike train pairs was set as the input for the template-matching approach.
Performance was calculated across a range of τ values, increasing in powers of 2 from 1ms to
256ms. Finally, to determine the optimal τ value at which performance was maximized for each
configuration, we implemented a fine-grain parameter where τ was varied in steps of 1ms, with
the optimization separately done for control and laser trials.

609
610   **References**
611

1.   deCharms RC, Zador A. Neural representation and the cortical code. *Annu Rev Neurosci*
23, 613-647 (2000).
2.   Zuo Y, Safaai H, Notaro G, Mazzoni A, Panzeri S, Diamond ME. Complementary
contributions of spike timing and spike rate to perceptual decisions in rat S1 and S2
cortex. *Curr Biol* **25**, 357-363 (2015).
3.   Yao JD, Sanes DH. Temporal Encoding is Required for Categorization, But Not
Discrimination. *Cereb Cortex* **31**, 2886-2897 (2021).
4.   Lu T, Liang L, Wang X. Temporal and rate representations of time-varying signals in the
auditory cortex of awake primates. *Nat Neurosci* **4**, 1131-1138 (2001).
5.   Tremblay R, Lee S, Rudy B. GABAergic Interneurons in the Neocortex: From Cellular
Properties to Circuits. *Neuron* **91**, 260-292 (2016).

6.　Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M. Inhibition of inhibition in visual cortex:
the logic of connections between molecularly distinct interneurons. *Nat Neurosci* **16**,
1068-1076 (2013).
7.　Antonoudiou P, Tan YL, Kontou G, Upton AL, Mann EO. Parvalbumin and Somatostatin
Interneurons Contribute to the Generation of Hippocampal Gamma Oscillations. *J
Neurosci* **40**, 7668-7687 (2020).
8.　Jang HJ, Chung H, Rowland JM, Richards BA, Kohl MM, Kwag J. Distinct roles of
parvalbumin and somatostatin interneurons in gating the synchronization of spike times in
the neocortex. *Sci Adv* **6**, eaay5333 (2020).
9.　Wehr M, Zador AM. Balanced inhibition underlies tuning and sharpens spike timing in
auditory cortex. *Nature* **426**, 442-446 (2003).
10.　Li LY*, et al.* A feedforward inhibitory circuit mediates lateral refinement of sensory
representation in upper layer 2/3 of mouse primary auditory cortex. *J Neurosci* **34**, 13670-
13683 (2014).
11.　Li LY, Xiong XR, Ibrahim LA, Yuan W, Tao HW, Zhang LI. Differential Receptive Field
Properties of Parvalbumin and Somatostatin Inhibitory Neurons in Mouse Auditory
Cortex. *Cereb Cortex* **25**, 1782-1791 (2015).
12.　Moore AK, Wehr M. Parvalbumin-expressing inhibitory interneurons in auditory cortex are
well-tuned for frequency. *J Neurosci* **33**, 13713-13723 (2013).
13.　Aizenberg M, Mwilambwe-Tshilobo L, Briguglio JJ, Natan RG, Geffen MN. Bidirectional
Regulation of Innate and Learned Behaviors That Rely on Frequency Discrimination by
Cortical Inhibitory Neurons. *PLoS Biol* **13**, e1002308 (2015).
14.　Natan RG*, et al.* Complementary control of sensory adaptation by two types of cortical
interneurons. *Elife* **4**, (2015).
15.　Blackwell JM, Geffen MN. Progress and challenges for understanding the function of
cortical microcircuits in auditory processing. *Nat Commun* **8**, 2165 (2017).
16.　Seybold BA, Phillips EAK, Schreiner CE, Hasenstaub AR. Inhibitory Actions Unified by
Network Integration. *Neuron* **87**, 1181-1192 (2015).
17.　Phillips EA, Hasenstaub AR. Asymmetric effects of activating and inactivating cortical
interneurons. *Elife* **5**, (2016).
18.　Moore AK, Weible AP, Balmer TS, Trussell LO, Wehr M. Rapid Rebalancing of Excitation
and Inhibition by Cortical Circuitry. *Neuron* **97**, 1341-1355 e1346 (2018).
19.　Satuvuori E, Kreuz T. Which spike train distance is most suitable for distinguishing rate
and temporal coding? *J Neurosci Methods* **299**, 22-33 (2018).
20.　Kreuz T, Chicharro D, Houghton C, Andrzejak RG, Mormann F. Monitoring spike train
synchrony. *J Neurophysiol* **109**, 1457-1472 (2013).
21.　Narayan R*, et al.* Cortical interference effects in the cocktail party problem. *Nat Neurosci*
10, 1601-1607 (2007).
22.　Mesgarani N, Chang EF. Selective cortical representation of attended speaker in multi-
talker speech perception. *Nature* **485**, 233-236 (2012).
23.　Maddox RK, Billimoria CP, Perrone BP, Shinn-Cunningham BG, Sen K. Competing
sound sources reveal spatial effects in cortical processing. *PLoS Biol* **10**, e1001319
(2012).
24.　Pachitariu M, Steinmetz N, Kadir S, Carandini M, Harris KD. Kilosort: realtime spike-
sorting for extracellular electrophysiology with hundreds of channels. *bioRxiv*, (2016).
25.　Rossant C*, et al.* Spike sorting for large, dense electrode arrays. *Nat Neurosci* **19**, 634-
641 (2016).
26.　Keller CH, Kaylegian K, Wehr M. Gap encoding by parvalbumin-expressing interneurons
in auditory cortex. *J Neurophysiol* **120**, 105-114 (2018).
27.　Rothauser EH, Chapman, W. D., Guttman, N., Nordby, K. S., Silbiger, H. R., Urbanek, G.
E., & Weinstock, M. I.E.E.E. recommended practice for speech quality measurements.
IEEE Trans Audio Electroacoust **17**, 225-246 (1969).
28.　Britten KH, Shadlen MN, Newsome WT, Movshon JA. The analysis of visual motion: a
comparison of neuronal and psychophysical performance. *J Neurosci* **12**, 4745-4765
(1992).

29.   Parker AJ, Newsome WT. Sense and the single neuron: probing the physiology of
679   perception. *Annu Rev Neurosci* **21**, 227-277 (1998).

680   30.   Wang L, Narayan R, Grana G, Shamir M, Sen K. Cortical discrimination of complex
681   natural stimuli: can single neurons match behavior? *J Neurosci* **27**, 582-589 (2007).

682   31.   Billimoria CP, Kraus BJ, Narayan R, Maddox RK, Sen K. Invariance and sensitivity to
683   intensity in neural discrimination of natural sounds. *J Neurosci* **28**, 6304-6308 (2008).

684   32.   Downer JD, Bigelow J, Runfeldt MJ, Malone BJ. Temporally precise population coding of
685   dynamic sounds by auditory cortex. *J Neurophysiol* **126**, 148-169 (2021).

686   33.   Hoglen NEG, Larimer P, Phillips EAK, Malone BJ, Hasenstaub AR. Amplitude modulation
687   coding in awake mice and squirrel monkeys. *J Neurophysiol* **119**, 1753-1766 (2018).

688   34.   van Rossum MC. A novel spike distance. *Neural Comput* **13**, 751-763 (2001).

689   35.   Scala F*, et al.* Phenotypic variation of transcriptomic cell types in mouse motor cortex.
690   *Nature,* (2020).

691   36.   Cardin JA*, et al.* Driving fast-spiking cells induces gamma rhythm and controls sensory
692   responses. *Nature* **459**, 663-667 (2009).

693   37.   Sohal VS, Zhang F, Yizhar O, Deisseroth K. Parvalbumin neurons and gamma rhythms
694   enhance cortical circuit performance. *Nature* **459**, 698-702 (2009).

695   38.   Bruno RM, Simons DJ. Feedforward mechanisms of excitatory and inhibitory cortical
696   receptive fields. *J Neurosci* **22**, 10966-10975 (2002).

697   39.   Xiang Z, Huguenard JR, Prince DA. Cholinergic switching within neocortical inhibitory
698   networks. *Science* **281**, 985-988 (1998).

699   40.   Wang JA, McFadden SL, Caspary D, Salvi R. Gamma-aminobutyric acid circuits shape
700   response properties of auditory cortex neurons. *Brain Research* **944**, 219-231 (2002).

701   41.   Kurt S, Moeller CK, Jeschke M, Schulze H. Differential effects of iontophoretic application
702   of the GABAA-antagonists bicuculline and gabazine on tone-evoked local field potentials
703   in primary auditory cortex: interaction with ketamine anesthesia. *Brain Res* **1220**, 58-69
704   (2008).

705   42.   Chen QC, Jen PH. Bicuculline application affects discharge patterns, rate-intensity
706   functions, and frequency tuning characteristics of bat auditory cortical neurons. *Hear Res*
707   **150**, 161-174 (2000).

708   43.   Kato HK, Asinof SK, Isaacson JS. Network-Level Control of Frequency Tuning in Auditory
709   Cortex. *Neuron* **95**, 412-423 e414 (2017).

710   44.   Briguglio JJ, Aizenberg M, Balasubramanian V, Geffen MN. Cortical Neural Activity
711   Predicts Sensory Acuity Under Optogenetic Manipulation. *J Neurosci* **38**, 2094-2105
712   (2018).

713   45.   Liang F*, et al.* Sparse Representation in Awake Auditory Cortex: Cell-type Dependence,
714   Synaptic Mechanisms, Developmental Emergence, and Modulation. *Cereb Cortex* **29**,
715   3796-3812 (2019).

716   46.   Theunissen F, Miller JP. Temporal encoding in nervous systems: a rigorous definition. *J*
717   *Comput Neurosci* **2**, 149-162 (1995).

718   47.   Ding N, Patel AD, Chen L, Butler H, Luo C, Poeppel D. Temporal modulations in speech
719   and music. *Neurosci Biobehav Rev* **81**, 181-187 (2017).

720   48.   Castellucci GA, Calbick D, McCormick D. The temporal organization of mouse ultrasonic
721   vocalizations. *PLoS One* **13**, e0199929 (2018).

722   49.   Chen C, Read HL, Escabi MA. Precise feature based time scales and frequency
723   decorrelation lead to a sparse auditory code. *J Neurosci* **32**, 8454-8468 (2012).

724   50.   Ghitza O. Linking speech perception and neurophysiology: speech decoding guided by
725   cascaded oscillators locked to the input rhythm. *Front Psychol* **2**, 130 (2011).

726   51.   Teng X, Poeppel D. Theta and Gamma Bands Encode Acoustic Dynamics over Wide-
727   Ranging Timescales. *Cereb Cortex* **30**, 2600-2614 (2020).

728   52.   Shadlen MN, Newsome WT. Noise, neural codes and cortical organization. *Curr Opin*
729   *Neurobiol* **4**, 569-579 (1994).

730   53.   Narayan R, Ergun A, Sen K. Delayed inhibition in cortical receptive fields and the
731   discrimination of complex stimuli. *J Neurophysiol* **94**, 2970-2975 (2005).

54.   Dong J, Colburn HS, Sen K. Cortical Transformation of Spatial Processing for Solving the
Cocktail Party Problem: A Computational Model(1,2,3). *eNeuro* **3**, (2016).
55.   Kapfer C, Glickfeld LL, Atallah BV, Scanziani M. Supralinear increase of recurrent
inhibition during sparse activity in the somatosensory cortex. *Nat Neurosci* **10**, 743-753
(2007).
56.   Silberberg G, Markram H. Disynaptic inhibition between neocortical pyramidal cells
mediated by Martinotti cells. *Neuron* **53**, 735-746 (2007).
57.   Adesnik H, Bruns W, Taniguchi H, Huang ZJ, Scanziani M. A neural circuit for spatial
summation in visual cortex. *Nature* **490**, 226-231 (2012).
58.   Lakunina AA, Nardoci MB, Ahmadian Y, Jaramillo S. Somatostatin-Expressing
Interneurons in the Auditory Cortex Mediate Sustained Suppression by Spectral
Surround. *J Neurosci* **40**, 3564-3575 (2020).
59.   Norman-Haignere SV*, et al.* Multiscale temporal integration organizes hierarchical
computation in human auditory cortex. *Nat Hum Behav* **6**, 455-469 (2022).
60.   Chou KF, Sen K. AIM: A network model of attention in auditory cortex. *PLoS Comput Biol*
17, e1009356 (2021).
61.   Gritton HJ*, et al.* Oscillatory activity in alpha/beta frequencies coordinates auditory and
prefrontal cortices during extinction learning. *bioRxiv*, (2020).
62.   James NM, Gritton HJ, Kopell N, Sen K, Han X. Muscarinic receptors regulate auditory
and prefrontal cortical communication during auditory processing. *Neuropharmacology*
144, 155-171 (2019).
63.   Hawley ML, Litovsky RY, Culling JF. The benefit of binaural hearing in a cocktail party:
effect of location and type of interferer. *J Acoust Soc Am* **115**, 833-843 (2004).
64.   Zhu Y, Qiao W, Liu K, Zhong H, Yao H. Control of response reliability by parvalbumin-
expressing interneurons in visual cortex. *Nat Commun* **6**, 6802 (2015).
65.   Morrill RJ, Hasenstaub AR. Visual Information Present in Infragranular Layers of Mouse
Auditory Cortex. *J Neurosci* **38**, 2854-2862 (2018).
66.   Hentschke H, Stuttgen MC. Computation of measures of effect size for neuroscience
data sets. *Eur J Neurosci* **34**, 1887-1894 (2011).
67.   Hentschke H. hhentschke/measures-of-effect-size-toolbox.) (2022).
68.   Serdar CC, Cihan M, Yucel D, Serdar MA. Sample size, power and effect size revisited:
simplified and practical approaches in pre-clinical, clinical and laboratory studies.
Biochem Med (Zagreb) **31**, 010502 (2021).
69.   Wesson DW, Donahou TN, Johnson MO, Wachowiak M. Sniffing behavior of mice during
performance in odor-guided tasks. *Chem Senses* **33**, 581-596 (2008).

Acknowledgments

769   This research was supported by the National Science Foundation (#IIS-1835270) and the
770   National Institute of Health (#1R34NS111742-01). We would like to thank Alberto Cruz-Martín,
771   Michael Economo, and Conor Houghton for comments and suggestions on the manuscript. We
772   also thank Monty Escabi and Oded Ghitza for discussions on coding time-scales in auditory
773   cortex and speech.

Author Contributions

776   J.C.N. and H.J.G. performed all experiments. J.C.N. collected and analyzed the data.
777   N.M.J. provided technical support and code for data analysis. X.H. and K.S obtained funding and
778   supervised the study. J.C.N, H.J.G, X.H., and K.S. wrote the manuscript and contributed to the
779   interpretation of the results.

781 **Competing Interest Statement:** The authors declare no conflicts of interest.

Figures and Tables

Figure 1. Experimental methods. A: Illustration depicting recording electrode location and optical fiber placement. Subjects were implanted with a 4-shank, 32-channel electrode array and optogenetic fiber in right hemisphere of ACx. Each shank contained 8 sites per shank with 100μm spacing between electrode contacts. **B:** Representative local field potential (LFP) activity from one mouse. LFP was used to estimate current source density and the layer of the recording site within each shank (Supplementary Figure 1). **C:** Example mean single unit waveform and inter-spike interval (ISI) auto-correlogram. Dashed lines in mean waveform represent one standard deviation above and below the mean, while scale bars are equal to 200μV and 1ms. Dashed red lines in correlogram represent ISIs of ±2ms. **D:** Schematic for control and optogenetic trial presentation. During 50% of all trials, a 532nm laser would turn on 50ms before sound stimulus onset and turn off coincident with sound offset. **E:** Paired comparisons of mean evoked firing rate during control and laser trials. Paired t-tests yielded a significant increase in evoked firing rate during optogenetic suppression for clean trials (n = 60 configurations; p < 1e-04, d = -0.92;) but not for masked trials (n = 28 configurations; p = 0.0571, d = -0.38).

Figure 2. Cortical discrimination in a cocktail party paradigm in mouse ACx. A: Illustration depicting the stimulus configuration for clean trials originating at +90˚ azimuth (**Ai**) and responses to both target stimuli (T) (**Aii**). Auditory stimuli were presented from speakers at 4 locations. Target stimuli consisted of white noise modulated by human speech envelopes extracted from recordings of speech sentences (see Methods, 'Auditory stimuli'). As shown in Aii, responses during clean trials exhibit spike timing and rapid firing rate modulation that follow the amplitude envelope of both target stimuli. All plotted PSTHs have a bin length of 20ms. **Bi:** Stimulus configuration for trials where targets (T) played at +90˚ and a competing masking stimulus (M) played at -90˚. Masking stimuli consisted of unmodulated white noise with the same onset and offset times as target stimuli. **Bii**: Responses to masked trials shown in Di. In this configuration, spike timing and firing rate modulation follow both target stimuli, despite the presence of the competing masker. **Ci:** Stimulus configuration for trials where targets played at 0˚ and maskers played at +45˚. **Cii**: Responses to target-masker configuration shown in Ci. For this configuration, spike timing and firing rate modulation do not follow either target stimulus, resulting in similar responses between target identities. **D:** Neural discriminability performance for all possible target-masker location configurations, referred to as the "spatial grid", for example cell featured in A-C. Outlined spots indicate configurations shown in A-C, matched by the outline color. Performance is calculated using a template-matching approach based on differences in instantaneous firing rate and spike timing similarities between trains (see Methods, 'Neural discriminability using SPIKE-distance'). The top grid shows discriminability for "clean" trials on top, while the bottom grid shows discriminability for "masked" trials. All blocks are color-coded according to the color axis shown to the right of the masked grid. Configurations with high performance ($\geq$ 70%) and a large effect size ($d \geq 1$), e.g., the configurations outlined in black and red, are referred to as "hotspots". **E:** Effect sizes for each spatial grid configuration in D, with the same outlines corresponding to examples in A-C. Positive values represent an increase in performance relative to a null distribution where spike trains within each target are template-matched to each other, while negative values represent a decrease in performance relative to null. **F:** The performance of all 28 single units exhibiting at least one hotspot during control trials. The translucent yellow surface represents the "upper envelope" of best performance across all single units for each masked spatial configuration, while the translucent blue surface represents the performance threshold of 70% for hotspots. Solid gray markers represent masked configurations with performances above threshold, while unfilled gray markers represent data points with performances below threshold. Black markers represent the maximal performance used to represent the upper envelope.

Figure 3. Effects of suppressing PV neurons on cortical discrimination. A: Responses during clean stimulus trials originating at 45° from one example cell during control (**Ai**) and optogenetic (**Aii**) conditions. **Aiii:** Inset showing zoomed-in portion of the response between 0.1 and 0.4s after sound onset, as outlined in Ai and Aii. Responses during optogenetic trials show earlier onset and less spike timing reproducibility, compared to the control. **B:** Example spatial grids from the same single unit during control and optogenetic conditions, with the performances at Clean Target 45° outlined in black to correspond to responses shown in A. Performance is color-coded according to the axis shown to the right of the 'Laser' grid. The reduction in spike timing reproducibility during optogenetic suppression (seen in Aiii) contributes to the decrease in performance (80%) compared to control trials at the same configuration (95%). Additionally, performance decreased during optogenetic suppression for the rest of the clean configurations, while performance at the masked control hotspots decreased to below threshold: Target 45°, Masker 90° (75% to 55%); Target 45°, Masker -90° (79% to 64%); and Target 90°, Masker -90° (74% to 67%). **C:** Paired comparisons of SPIKE-distance-based performance from control and PV-suppressed trials at the same spatial grid location. Paired t-tests yielded a significant decrease in performance for both clean (n = 60 configurations; p < 1e-04, d = 0.88) and masked (n = 28 configurations; p < 1e-04, d = 0.89) trials during optogenetic suppression, indicating that PV suppression significantly reduced discrimination performance.

Figure 4. Effects of suppressing PV neurons on spike timing and rate-based coding measures. A: Performance based on ISI-distance, which measures differences between trains in instantaneous firing rate only (see Methods, 'ISI-distance'). Paired t-tests showed a significant decrease in performance for both clean (n = 60 configurations; p < 1e-04, d = 0.73) and masked (n = 28 configurations; p = 4.69e-04, d = 0.75) trials. **B:** Performance based on RI-SPIKE-distance, which measures differences between trains in spike timing only (see Methods, 'RI-SPIKE-distance'). Paired t-tests showed a significant decrease in performance for both clean (p < 1e-04, d = 0.78) and masked (p = 6.79e-04, d = 0.73) trials. **C:** Performance based on differences in total spike count between spike trains was near chance level, indicating that total spike count did not account for overall discrimination performance. Paired t-tests showed a significant decrease in performance for clean trials (p = 0.0279, d = 0.29) but not for masked trials (p = 0.1121, d = 0.31). **D:** Summary figure showing contributions from spike distance measures presented in Figure 3C and 4A-C on the same scale and axis. Changes in spike timing and instantaneous firing rate-based measures (RI-SPIKE and ISI, respectively) provide relatively high discrimination performance and show a significant decrease upon optogenetic suppression of PV neurons.

Figure 5. Effects of optogenetic suppression on spiking activity measures. A: Rate-
normalized RMS difference between target PSTHs during both conditions. Paired t-tests found
significant decreases between conditions during both clean trials (n = 60 configurations; p < 1e-
04, d = 0.91) and masked trials (n = 28 configurations; p = 0.0058, d = 0.57). **B:** Trial similarity
between responses to the same target during both conditions. Paired t-tests found a highly
significant decrease between conditions during clean trials (n = 60 configurations; p < 1e-04, d =
0.77) but not masked trials (n = 28 configurations; p = 0.8216, d = 0.04).

Figure 6. Decoding time analysis. A: Histogram of optimal τ for hotspots across both conditions
(control and laser) and stimulus types (clean and masked). Dashed line indicates median value of
43ms, and shaded region represents the inter-quartile range (IQR) between 28ms and 72.5ms.
Paired t-tests did not find a significant change in optimal τ within hotspots between conditions
during either clean (n = 60 configurations; p = 0.4117, d = 0.11) or masked (n = 28 configurations;
p = 0.2936, d = 0.20) trials. **B:** van Rossum-based performance with τ set at 8ms. Performance
was found to significantly decrease during both clean (p < 1e-04, d = 0.61) and masked (p =
0.0011, d = 0.69) trials. **C:** van Rossum-based performance with τ set at 32ms. Performance was
found to significantly decrease during both clean (p < 1e-04, d = 0.75) and masked (p = 8e-04, d
= 0.71) trials. **D:** van Rossum-based performance with τ set at 256ms. Performance was found to
significantly decrease during both clean (p < 1e-04, d = 0.58) and masked (p < 1e-04, d = 1.14)
trials.
886

Figure 7. Comparison of optimal τ values and other time scales. Semi-logarithmic plot showing various time-scales for spike timing in mouse ACx neurons compared to mouse vocalizations, human speech, and neural oscillations. Top: Time scales for human speech sounds, mouse vocalizations, and sniffing periods[69]. In the mouse time-scales, short and long USV bars represent the mean (black line) ± 2 s.t.d. vocalization length. Within the plot, from left to right: the refractory period for single units (red dashed line); and the distribution of optimal τ values from Figure 6 (solid black curve), with the dashed yellow line indicating the median value of 43ms and the shaded green region representing the IQR. The bottom axes show the time scale of optimal τ values in ms and frequency in Hz, with the latter decreasing from left to right. Shaded bars represent frequency bands for neural oscillations.

897   **Figure 8. Cortical circuits for complex scene analysis.** Hypothesized conceptual model of
898   cortical circuit underlying spatial grids. C and R cells represent excitatory units, I cells mediate
899   within-channel inhibition, and X cells mediate cross-channel inhibition.

τ (ms)	d_{clean}	p_{clean}	d_{masked}	p_{masked}
1	-0.13	0.3017	0.60	0.0037
2	0.02	0.9014	0.59	0.0044
4	0.28	0.0340	0.59	0.0041
8	0.61	< 1e-04	0.69	0.0011
16	0.73	< 1e-04	0.70	0.0010
32	0.75	< 1e-04	0.71	8e-04
64	0.68	< 1e-04	0.78	3e-04
128	0.61	< 1e-04	0.98	< 1e-04
256	0.58	< 1e-04	1.14	< 1e-04

901 **Table 1. Effect sizes and paired t-test results for all τ values used in van Rossum distance-**
902 **based performance calculations.**

A ACtx

32 Channel ACtx Electrode (Rt. Hemisphere)

S2

Auditory Cortex Electrode

Optical Fibers

DG CA A1

V2

Ent

B Shank 1

Auditory Cortex Electrode

Shank 2

CA1   A1   1 2 3 4

Shank 3

Shank 4

200µV

200ms

C

200 µV

1 ms

ISI auto-correlogram

ISI (ms)

-30   30

D

50%

2s ITI   3s   2 s ITI

3.05 s

Optogenetic silencing

50%

2s ITI   3s   2s ITI

E   Clean   Masked

***   n.s.

Evoked firing rate (Hz)

Control   Laser   Control   Laser

A ISI-distance performance

B RI-SPIKE-distance performance

C Spike count distance performance

D Clean / Masked