Computation of the mitochondrial age distribution along the axon length

Ivan A. Kuznetsov[a], [b] and Andrey V. Kuznetsov[c]

[a] Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

[b] Department of Bioengineering, University of Pennsylvania, Philadelphia, PA 19104, USA

[c] Department of Mechanical and Aerospace Engineering, North Carolina State University, Raleigh, NC 27695-7910, USA; e-mail: avkuznet@ncsu.edu

Abstract

We describe a compartmental model of mitochondrial transport in axons, which we apply to compute mitochondrial age at different distances from the soma. The model predicts that the average mitochondrial age at the tip of an axon with a length of 1 cm is approximately 22 hours. The mitochondria are youngest closest to the soma and their age scales approximately linearly with distance from the soma. To the best of the authors' knowledge, this is the first attempt to predict the spatial distribution of mitochondrial age within an axon. A sensitivity study of the mean age of mitochondria to various model parameters is also presented.

Keywords: neurons; axonal transport; organelles; mean age; mathematical modeling

1. Introduction

Mitochondria generate easily usable chemical power for cells. In addition to their direct involvement in cellular respiration, they are also involved in apoptosis, calcium buffering, and many other important biological functions (Fan et al. 2001; Picard and McEwen 2018). Abnormalities in mitochondrial transport occur in many neurological disorders (Zheng et al. 2019). Recent research implicates α-synuclein-induced disruptions of mitochondrial trafficking in the development of Parkinson's disease (PD) (Clague and Rochin 2016; Shahmoradian et al. 2019). Since packed mitochondria membrane fragments are one of the components found in Lewy bodies (Shahmoradian et al. 2019), investigating the transport of mitochondria is important for understanding the fundamentals of PD, for example in regard to possible energy deficits in dopaminergic neurons due to disruptions in mitochondrial axonal transport (Prots et al. 2018).

There are three pools of mitochondria in axons: anterogradely transported, retrogradely transported, and stationary (Hollenbeck and Saxton 2005). Understanding how old the mitochondria in these pools are and how their age depends on the distance from the soma is critical, as it yields insight into how mitochondria are renewed in the axon and the potential for dysregulation of mitochondrial transport in various diseases. Estimating the age of mitochondria may be important, for example, in evaluating the possible role of oxidative damage in the development of various neurodegenerative diseases (Uttara et al. 2009).

Renewal of mitochondrial proteins involves mitochondrial fusion and fission (Misgeld and Schwarz 2017). Fusion of old and new mitochondria leads to an exchange of old and new soluble and membrane-bound components, such as mitochondrial proteins (Malka et al. 2005; Patel et al. 2013). It is believed that mitochondrial fission leads to the separation of mutated copies of the mitochondrial genome into a separate mitochondrion, which is then recycled by the autophagosome (Twig et al. 2008; Safiulina and Kaasik 2013). Fission also serves as a mechanism that prevents mitochondria from becoming excessively long, so longer mitochondria are more likely to enter fission than shorter ones (Cagalinec et al. 2013). If fission events are unopposed, they can lead to complete fragmentation of the mitochondrial network (Ahola et al. 2019). Mitochondrial fusion accelerates mitochondrial fission, which is necessary to maintain a constant average length of mitochondria. Fission is important for autophagy, the mechanism of mitochondria removal (Twig et al. 2008).

Mitochondria are moved anterogradely by kinesin (in particular, by kinesin-1) motors and retrogradely by cytoplasmic dynein (Melkov and Abdu 2018). The motion of individual

mitochondria is not continuous but rather interrupted by frequent pauses. Mitochondria move with an average velocity of 0.4-0.8 μm/s (Narayanareddy et al. 2014).

For investigating the distribution of the mean age of mitochondria in various compartments surrounding demand sites, Ferree et al. (2013) used a photoconvertible construct, termed MitoTimer, that switches from green to red fluorescence over a period of hours due to slow oxidation as it ages (red corresponds to older mitochondria and green corresponds to younger mitochondria). MitoTimer exhibits green fluorescence for the first 8 hours. After 24 hours it shows a mixture of green and red fluorescence (producing a yellow color). After 48-72 hours MitoTimer exhibits red fluorescence (Ferree et al. 2013; Hernandez et al. 2013; Gottlieb and Stotland 2015; Wilson et al. 2019). Experiments in Ferree et al. (2013) reported that mitochondria residing in compartments that surround the more distal demand sites are older than those residing in compartments surrounding more proximal demand sites. The ratio of red/green fluorescence depends on the rate of mitochondria synthesis in the soma, the rate of mitochondrial transport in the axon, the rates of fusion and fission (which equilibrate the MitoTimer protein content in mitochondria and mitochondria age), and autophagy.

A computational study (Patel et al. 2013) investigated mitochondria health, which is a proxy for mitochondrial membrane potential. An initial population of 150 unfused mitochondria was simulated. Recent research (Agrawal and Koslover 2021), which simulated mitochondria health as a decaying component, developed models for two maintenance models of mitochondria: one that assumed the exchange of mitochondrial proteins between moving and stationary mitochondria due to mitochondrial fusion and fission, and the other that assumed occasional replacement of stationary mitochondria with new ones arriving from the soma.

2. Materials and models

2.1. Governing equations

We treat the axon as a series of discrete compartments (representing mitochondria demand sites) and the age as the time since the production of a mitochondrion. The model developed in this paper accounts for stationary, anterogradely, and retrogradely moving mitochondria. We simulated a terminal with N high energy demand sites (hereafter demand sites) that contain stationary mitochondria (Fig. 1). The model developed in this research accounts for N resident and $2N$ transiting (anterograde and retrograde) kinetic states and contains $3N$ ordinary differential equations (ODEs) (we used $N=10$ in this research).

We used a multi-compartment model (Anderson 1983; Jacquez 1985; Jacquez and Simon 1993) to formulate equations expressing the conservation of the total length of mitochondria in the compartments. Our rationale for representing the axon as a series of discrete compartments comes from our desire to use the methodology for computing the age distribution of cargos in compartmental systems developed in Metzler et al. (2018), Metzler and Sierra (2018). We used a similar compartmental approach to simulate the transport of dense core vesicles in *Drosophila* motoneurons (Kuznetsov and Kuznetsov 2019; Kuznetsov and Kuznetsov 2020). We numbered the demand sites in the axon terminal from the most proximal (#1) to the most distal (#*N*) (Fig. 1). The demand sites have three compartments (representing resident, transiting-anterograde, and transiting-retrograde pools of mitochondria), see Fig. 2.

Since the length of an axon is much greater than its width, we used the total length of mitochondria in the kinetic state per unit axon length, measured in (μm of mitochondria)/μm. The total length of mitochondria is defined as the sum of the lengths of all mitochondria residing in the same kinetic state (stationary, anterograde, or retrograde) in the same compartment. The total length of mitochondria in the compartment for a particular kinetic state (Fig. 2) is the conserved quantity. The dependent variables utilized in the model are summarized in Table 1.

Table 1. Dependent variables in the model of mitochondrial transport and accumulation in the axon.

Symbol	Definition	Units
l_i^s	Total length of stationary mitochondria per unit length of the axon in the compartment by the ith demand site	(μm of mitochondria)/μm
l_i^a	Total length of anterogradely moving mitochondria per unit length of the axon in the compartment by the ith demand site	(μm of mitochondria)/μm
l_i^r	Total length of retrogradely moving mitochondria per unit length of the axon in the compartment by the ith demand site	(μm of mitochondria)/μm
$j_{a \to b}$	Flux of mitochondria length from state (compartment) a to state (compartment) b	(μm of mitochondria) s^{-1}

First, the conservation of the total mitochondria length in the compartment around the most proximal demand site (# 1) is formulated. Stating this for the stationary pool of mitochondria gives the following equation (Fig. 2):

$$L_1 \frac{dl_1^s}{dt} = p_s \left(j_{soma \to 1} + j_{2 \to 1} \right) - L_1 k_w l_1^s, \tag{1}$$

where L_1 is the length of an axonal compartment around the most proximal mitochondria demand site and t is the time. We assumed that the axonal length L_{ax} is split into N compartments of equal length, for example, $L_1 = L_{ax} / N$.

The conservation of mitochondria length in the anterograde pool of mitochondria around demand site 1 gives the following equation (Fig. 2):

$$L_1 \frac{dl_1^a}{dt} = j_{soma \to 1} - p_s j_{soma \to 1} - j_{1 \to 2} + \varepsilon L_1 k_w l_1^s. \tag{2}$$

Unlike Agrawal and Koslover (2021), we did not assume that the mitochondria that reenter the motile state are split evenly between the anterograde and retrograde components. Instead, the portion of mitochondria that reenter the anterograde component is given by ε and the portion of mitochondria that reenter the retrograde component is given by $(1 - \varepsilon)$, see Fig. 2 and Table 2.

The conservation of mitochondria length in the retrograde pool of mitochondria around the most proximal demand site gives the following equation (Fig. 2):

$$L_1 \frac{dl_1^r}{dt} = j_{2 \to 1} - p_s j_{2 \to 1} - j_{1 \to soma} + (1 - \varepsilon) L_1 k_w l_1^s. \tag{3}$$

Next, the conservation of mitochondria length in the compartment around the ith demand site ($i=2,...,N-1$) is formulated. The conservation of mitochondria length in the stationary pool around the ith demand site gives the following equation (Fig. 2):

$$L_i \frac{dl_i^s}{dt} = p_s \left(j_{i-1 \to i} + j_{i+1 \to i} \right) - L_i k_w l_i^s. \tag{4}$$

The conservation of mitochondria length in the anterograde pool in the ith demand site gives the following equation (Fig. 2):

$$L_i \frac{dl_i^a}{dt} = j_{i-1 \to i} - p_s j_{i-1 \to i} - j_{i \to i+1} + \varepsilon L_i k_w l_i^s. \tag{5}$$

The conservation of mitochondria length in the retrograde pool in the ith demand site gives the following equation (Fig. 2):

$$L_i \frac{dl_i^r}{dt} = j_{i+1 \to i} - p_s j_{i+1 \to i} - j_{i \to i-1} + (1-\varepsilon) L_i k_w l_i^s .$$ (6)

Finally, the conservation of mitochondria length in the compartment around the most distal demand site (# N) is formulated. The conservation of mitochondria length in the stationary pool gives the following equation (Fig. 2):

$$L_N \frac{dl_N^s}{dt} = p_s \left(j_{N-1 \to N} + j_{N \to 3N} \right) - L_N k_w l_N^s .$$ (7)

The conservation of mitochondria length in the anterograde pool gives the following equation (Fig. 2):

$$L_N \frac{dl_N^a}{dt} = j_{N-1 \to N} - p_s j_{N-1 \to N} - j_{N \to N-1} + \varepsilon L_N k_w l_N^s .$$ (8)

The conservation of mitochondria length in the retrograde pool gives the following equation (Fig. 2):

$$L_N \frac{dl_N^r}{dt} = j_{N \to 3N} - p_s j_{N \to 3N} - j_{N \to N-1} + (1-\varepsilon) L_N k_w l_N^s .$$ (9)

Eqs. (1)-(9) must be supplemented by the following equations for the mitochondria fluxes between the demand sites. $j_{soma \to 1}$ is an input model parameter; equations for other anterograde fluxes (Fig. 2) are

$$j_{1 \to 2} = v_a l_1^a ,$$ (10)

…

$$j_{i-1 \to i} = v_a l_{i-1}^a ,$$ (11)

$$j_{i \to i+1} = v_a l_i^a ,$$ (12)

…

$$j_{N-1 \to N} = v_a l_{N-1}^a ,$$ (13)

$$j_{N \to 3N} = v_a l_N^a .$$ (14)

Equations for retrograde fluxes (Fig. 2) are

$$j_{N \to N-1} = v_r l_N^r,$$ (15)

...

$$j_{i+1 \to i} = v_r l_{i+1}^r,$$ (16)

$$j_{i \to i-1} = v_r l_i^r,$$ (17)

...

$$j_{2 \to 1} = v_r l_2^r,$$ (18)

$$j_{1 \to soma} = v_r l_1^r.$$ (19)

The model given by Eqs. (1)-(19) can also be replaced by a single set (valid between $x=0$ and $x=L_{ax}$) of three partial differential equations (PDEs) (for anterograde, retrograde, and stationary mitochondria):

$$\frac{\partial l^a}{\partial t} = -v_a \frac{\partial l^a}{\partial x} - k_s^a l^a + \varepsilon k_w l^s,$$ (20)

$$\frac{\partial l^r}{\partial t} = v_r \frac{\partial l^r}{\partial x} - k_s^r l^r + (1-\varepsilon) k_w l^s,$$ (21)

$$\frac{\partial l^s}{\partial t} = k_s^a l^a + k_s^r l^r - k_w l^s,$$ (22)

where

$$k_s^a = \frac{2 p_s v_a}{L_1}$$ (23)

and

$$k_s^r = \frac{2 p_s v_r}{L_1}.$$ (24)

Eqs. (21)-(22) must be solved with the following boundary conditions:

At $x=0$: $\qquad v_a l^a (0) = j_{soma \to 1},$ (25)

At $x = L_{ax}$: $\qquad v_a l^a (L_{ax}) = v_r l^r (L_{ax}).$ (26)

However, there is not a simple way to investigate how mitochondrial age varies throughout the axon utilizing PDEs (20)-(22). In order to more efficiently conduct our investigation, we resort to using discrete compartments (Fig. 2).

Since we seek the steady-state solution (which is of physical significance), the initial condition does not matter. We solved Eqs. (1)-(9) with zero initial conditions in all compartments and proceeded until the solution reached steady-state. The solution procedure involves solving initial value problems for three sets of ODEs. The numerical solution was obtained using MATLAB's solver, ODE45 (MATLAB R2019a, MathWorks, Natick, MA, USA). The steady-state solution could also be obtained by setting the left-hand side of Eqs. (1)-(9) to zero and solving the obtained linear system for the densities in each compartment.

Table 2. Parameters characterizing mitochondrial transport and accumulation in the axon.

Symbol	Definition	Units	Value or range	Reference(s)	Value used in computations
v_a	Average velocity of anterogradely moving mitochondria	$\mu m\,s^{-1}$	0.5	Agrawal and Koslover (2021)	0.5
v_r	Average velocity of retrogradely moving mitochondria	$\mu m\,s^{-1}$	0.5	Agrawal and Koslover (2021)	0.5
$j_{soma \to 1}$	Rate of mitochondria production in the soma	mitochondria s^{-1}	0.0375	Agrawal and Koslover (2021)	0.0375 [a]
L_{ax}	Length of the axon	μm	10^4-10^5	Agrawal and Koslover (2021)	10^4
k_w	Kinetic constant characterizing the rate of mitochondria reentry into motile states (anterograde and retrograde)	s^{-1}	$2 \times 10^{-4} - 2 \times 10^{-3}$ [b]	Agrawal and Koslover (2021)	2×10^{-4}, 5×10^{-4}
ε	Portion of mitochondria leaving the stationary state that join the anterograde pool; $(1-\varepsilon)$ join the retrograde pool		0.5	Agrawal and Koslover (2021)	0.5

f_s	Fraction of stationary mitochondria		0.5-0.6	Agrawal and Koslover (2021)	0.5
p_s	Probability that a motile mitochondrion would transition to a stationary state at a demand site		0.4-0.5 [c]	Agrawal and Koslover (2021)	0.04, 0.4
N	Number of demand sites in the axon		5-100 [d]	Agrawal and Koslover (2021)	10

[a] We estimated $j_{soma \to 1}$ using the formula suggested in Agrawal and Koslover (2021), $Mv / (2L_{ax})$, where M is the number of mitochondria in the axon and v is the average velocity of mitochondria motion in the axon. We used 1,500 mitochondria for M, 0.5 µm/s for v, and 1 cm for L_{ax}.

[b] We used Eq. (4) from Agrawal and Koslover (2021), $f_s = \dfrac{Nvp_s}{L_{ax}k_w + Nvp_s}$. By solving this equation for k_w using values given in Table 2 (for N we used the range 10-100), we obtained that k_w is in the range 2×10^{-4} to 2×10^{-3} s^{-1}.

[c] We used $p_s = \dfrac{N_s}{2N}$, where N_s is an average number of stops that a mitochondrion makes during a back and forth journey in the axon (Agrawal and Koslover 2021).

[d] Number of demand sites in a real axon can be much larger than that. Ideally, the size of a compartment should represent a size of a mitochondria demand site (and the compartments should not be of the same size), but this would result in a very large number of compartments.

2.2. Model of mean age of mitochondria in the demand sites

To compute the age distributions of mitochondria in the stationary and moving states in the compartments surrounding the demand sites, the method described by Metzler et al. (2018), Metzler and Sierra (2018) was utilized. In order to use the methodology described in the above papers, Eqs. (1)-(9) were restated as the following matrix equation:

$$\frac{d}{dt}\mathbf{C}^*(t) = \mathrm{B}\left(\mathbf{C}^*(t), t\right)\mathbf{C}^*(t) + \mathbf{u}(t). \tag{27}$$

In Eq. (27) $\mathbf{C}^*$, using the terminology of Metzler et al. (2018), is the state vector, which is defined by its components ($\mathbf{C}^*$ is a column vector):

$$C_1^* = l_1 L_1, \ldots, C_N^* = l_N L_N, C_{N+1}^* = l_1^a L_1, \ldots, C_{2N-1}^* = l_{N-1}^a L_{N-1}, C_{2N}^* = l_N^m L_N, C_{2N+1}^* = l_1^r L_1, \ldots, C_{3N}^* = l_N^r L_N,$$
(28)

where

$$\hat{l}_1 = l_1, \ldots, \hat{l}_N = l_N, \hat{l}_{N+1} = l_1^a, \ldots, \hat{l}_{2N-1} = l_{N-1}^a, \hat{l}_{2N} = l_N^m, \hat{l}_{2N+1} = l_1^r, \hat{l}_{3N} = l_N^r.$$
(29)

Rescaling the equations in the form given by Eqs. (27) and (28) is necessary because the conserved quantity is the total mitochondrial length in a certain compartment rather than its linear density.

The total length of mitochondria in stationary states is represented by the first N components of the state vector, the total length of mitochondria in anterograde states is represented by the next N components ($N+1,\ldots,2N$), and the total length of mitochondria in retrograde states is represented by the last N components ($2N+1,\ldots,3N$), see Fig. 2. Vector $\mathbf{u}(t)$ is defined below by Eqs. (60) and (61).

In our model, matrix B($3N$,$3N$) is as follows. We accounted for the internal mitochondria fluxes between compartments, the external mitochondria flux entering the terminal from the axon, and the mitochondria flux leaving the terminal to move back to the axon (Fig. 2). Equations for the following elements of matrix B were obtained by analyzing mitochondria fluxes to and from the compartments surrounding the most proximal demand site:

$$b_{1,1} = -L_1 k_w l_1^s / \left(L_1 l_1^s \right),$$
(30)

$$b_{N+1,N+1} = -\left(p_s j_{soma \to 1} + j_{1 \to 2} \right) / \left(L_1 l_1^a \right),$$
(31)

$$b_{2N+1,2N+1} = -\left(p_s j_{2 \to 1} + j_{1 \to soma} \right) / \left(L_1 l_1^r \right),$$
(32)

$$b_{N+1,1} = \varepsilon L_1 k_w l_1^s / \left(L_1 l_1^s \right),$$
(33)

$$b_{1,N+1} = p_s j_{soma \to 1} / \left(L_1 l_1^a \right),$$
(34)

$$b_{2N+1,1} = (1 - \varepsilon) L_1 k_w l_1^s / \left(L_1 l_1^s \right),$$
(35)

$$b_{1,2N+1} = p_s j_{2 \to 1} / \left(L_1 l_1^r \right),$$ (36)

$$b_{N+2,N+1} = j_{1 \to 2} / \left(L_1 l_1^a \right),$$ (37)

$$b_{2N+1,2N+2} = j_{2 \to 1} / \left(L_2 l_2^r \right).$$ (38)

The analysis of mitochondria fluxes to and from the compartments surrounding demand site i ($i=2,\ldots,N\text{-}1$) gives equations for the following elements of matrix B:

$$b_{i,i} = -L_i k_w l_i^s / \left(L_i l_i^s \right),$$ (39)

$$b_{N+i,N+i} = -\left(p_s j_{i-1 \to i} + j_{i \to i+1} \right) / \left(L_i l_i^a \right),$$ (40)

$$b_{2N+i,2N+i} = -\left(p_s j_{i+1 \to i} + j_{i \to i-1} \right) / \left(L_i l_i^r \right),$$ (41)

$$b_{N+i,i} = \varepsilon L_i k_w l_i^s / \left(L_i l_i^s \right),$$ (42)

$$b_{i,N+i} = p_s j_{i-1 \to i} / \left(L_i l_i^a \right),$$ (43)

$$b_{2N+i,i} = \left(1 - \varepsilon \right) L_i k_w l_i^s / \left(L_i l_i^s \right),$$ (44)

$$b_{i,2N+i} = p_s j_{i+1 \to i} / \left(L_i l_i^r \right),$$ (45)

$$b_{N+i,N+i-1} = j_{i-1 \to i} / \left(L_{i-1} l_{i-1}^a \right),$$ (46)

$$b_{N+i+1,N+i} = j_{i \to i+1} / \left(L_i l_i^a \right),$$ (47)

$$b_{2N+i-1,2N+i} = j_{i \to i-1} / \left(L_i l_i^r \right),$$ (48)

$$b_{2N+i,2N+i+1} = j_{i+1 \to i} / \left(L_{i+1} l_{i+1}^r \right).$$ (49)

By analyzing mitochondria fluxes to and from the compartments surrounding the most distal demand site, equations for the following elements of matrix B were obtained:

$$b_{N,N} = -L_N k_w l_N^s / \left(L_N l_N^s \right),$$ (50)

$$b_{2N,2N} = -\left(p_s j_{N-1 \to N} + j_{N \to 3N} \right) / \left(L_N l_N^a \right),$$ (51)

$$b_{3N,3N} = -\left(p_s j_{N \to 3N} + j_{N \to N-1} \right) / \left(L_N l_N^r \right),$$ (52)

$$b_{2N,N} = \varepsilon L_N k_w l_N^s / \left(L_N l_N^s \right),$$

(53)

$$b_{N,2N} = p_s j_{N-1\to N} / \left(L_N l_N^a \right),$$

(54)

$$b_{2N,2N-1} = j_{N-1\to N} / \left(L_{N-1} l_{N-1}^a \right),$$

(55)

$$b_{3N,2N} = j_{N\to 3N} / \left(L_N l_N^a \right),$$

(56)

$$b_{3N,N} = \left(1-\varepsilon\right) L_N k_w l_N^s / \left(L_N l_N^s \right),$$

(57)

$$b_{N,3N} = p_s j_{N\to 3N} / \left(L_N l_N^r \right),$$

(58)

$$b_{3N-1,3N} = j_{N\to N-1} / \left(L_N l_N^r \right).$$

(59)

All other elements of matrix B, except for those given by Eqs. (30)-(59), are equal to zero.

The anterograde flux from the axon to the compartment with anterograde mitochondria by the most proximal demand site, $j_{soma\to 1}$, is the only flux entering the terminal. Our model assumes that all mitochondria that leave the terminal (their flux is $j_{soma\to 1}$) return to the soma for degradation, and none reenter the terminal. We also assumed that mitochondria that enter the terminal (their flux is $j_{soma\to 1}$) are all newly synthesized in the soma and their age at the time of entry is equal to zero. This means that our model does not account for the time it takes mitochondria to travel from the soma to the terminal. The mitochondrial age calculated here should be understood as the time since mitochondria entered the terminal.

The following equation yields the $N+1^{th}$ element of column vector **u**:

$$u_{N+1} = j_{soma\to 1}.$$

(60)

Because no other external fluxes enter the terminal,

$$u_i = 0 \quad (i=1,\ldots,N,N+2,\ldots,3N-1).$$

(61)

The state transition matrix, Φ, which is defined in Metzler et al. (2018), may be determined by solving the following matrix equation:

$$\frac{d}{dt}\Phi(t,t_0) = B\left(C^*(t),t\right)\Phi(t,t_0),$$

(62)

where t_0 is the initial time. In our case $t_0 = 0$.

The initial condition for Eq. (62) is

$$\Phi(t_0, t_0) = I.$$ (63)

In Eq. (63) symbol I denotes an identity matrix.

We assumed that all mitochondria that enter the terminal are new. The age density of mitochondria entering the terminal after $t = 0$ can then be determined from the following equation:

$$\mathbf{p}(c, t) = 1_{[0, t - t_0)}(c) \Phi(t, t - c) \mathbf{u}(t - c),$$ (64)

where $1_{[0, t - t_0)}$ is the indicator function, which equals 1 if $0 \le c < t - t_0$ and equals 0 otherwise. The age density of mitochondria is the ratio of the total length of mitochondria in the compartment which have an age between T and $T + dT$ over the duration of the interval dT. An integral of the age density of mitochondria over any time period with respect to time gives the length of mitochondria having an age within that time range.

The mean age of mitochondria in demand sites is (Rasmussen et al. 2016):

$$\bar{c}_i(t) = \frac{\int_0^\infty c p_i(c, t) dc}{\int_0^\infty p_i(c, t) dc} \qquad (i=1,\ldots,3N).$$ (65)

$\bar{c}_i(t)$ $(i=1,\ldots,3N)$ may be found by solving the following mean age system (Metzler et al. 2018; Rasmussen et al. 2016):

$$\frac{d}{dt} \bar{\mathbf{c}}(t) = \mathbf{g}\left(\mathbf{C}^*(t), t\right).$$ (66)

The initial condition for Eq. (66) is

$$\bar{\mathbf{c}}(0) = 0,$$ (67)

where $\bar{\mathbf{c}} = (\bar{c}_1, \ldots, \bar{c}_{3N})$. Vector $\mathbf{g}$ is defined as follows:

$$g_i(t) = 1 + \frac{\sum_{j=1}^{3N} (\bar{c}_j(t) - \bar{c}_i(t)) b_{i,j}(t) C_j(t) - \bar{c}_i(t) u_i}{C_i(t)} \qquad (i=1,\ldots,3N).$$ (68)

The following vectors were then defined to represent the mean age of mitochondria in the stationary, anterograde, and retrograde states in the demand sites, respectively:

$$\mathbf{a}_s = \left(\overline{c}_1, ..., \overline{c}_N \right), \ \mathbf{a}_a = \left(\overline{c}_{N+1}, ..., \overline{c}_{2N} \right), \ \mathbf{a}_r = \left(\overline{c}_{2N+1}, ..., \overline{c}_{3N} \right). \tag{69}$$

The arrays $\mathbf{a}_s$, $\mathbf{a}_a$, and $\mathbf{a}_r$ are one-dimensional and of size N.

Numerical solutions were again obtained utilizing MATLAB's solver, ODE45. The error tolerance parameters, RelTol and AbsTol, were set to 10^{-6} and 10^{-8}, respectively. We checked that lowering RelTol and AbsTol did not influence the solutions. The results were also checked by using the formulas reported in Metzler and Sierra (2018). In particular, the solution of Eq. (27) for a steady-state can be obtained as:

$$\mathbf{C}_{ss}^* = -\mathbf{B}_{ss}^{-1} \mathbf{u}_{ss}, \tag{70}$$

where the superscript -1 on a matrix denotes the inverse of the matrix and the subscript ss denotes steady-state. The mean ages of mitochondria in various compartments displayed in Fig. 2 at steady-state can be obtained as:

$$\overline{\mathbf{c}}_{ss} = -\left(\hat{\mathbf{C}}_{ss}^* \right)^{-1} \mathbf{B}_{ss}^{-1} \mathbf{C}_{ss}^*, \tag{71}$$

where

$$\hat{\mathbf{C}}_{ss}^* = \mathrm{diag}\left(C_{1,ss}^*, C_{2,ss}^*, ..., C_{3N,ss}^* \right) \tag{72}$$

is the diagonal matrix with the components of the state vector, defined in Eq. (28), on the main diagonal. These components ($C_{1,ss}^*$, $C_{2,ss}^*$, etc.) are calculated at steady-state. Also in Eq. (71),

$$\overline{\mathbf{c}}_{ss} = \left(\overline{c}_{1,ss}, ..., \overline{c}_{3N,ss} \right)^T \tag{73}$$

is the column vector composed of the mean ages of mitochondria in various compartments (superscript T means transposed).

3. Results

3.1. Estimation of the average age of mitochondria in the most distal demand site

According to Fig. 4C in Ferree et al. (2013), at some point in time the green/red fluorescence ratio linearly decreases with an increase of the distance from the nucleus as:

$$y = -0.5863x + 2303.5 , \tag{74}$$

where x is in μm and y is in arbitrary units. From Eq. (74) it follows that

$$\frac{dy}{dx} = -0.5863 . \tag{75}$$

In Ferree et al. (2013) MitoTimer production was induced by exposure to doxycycline. Green fluorescence starts approximately 5 hours after doxycycline induction. At that point in time, red fluorescence is equal to zero (Fig. 2A in Ferree et al. 2013). According to Fig. 2A in Ferree et al. (2013), the red/green fluorescence ratio, which is the reciprocal of y, is approximately given by

$$z = \frac{1.5}{55} t_{mit} , \tag{76}$$

where t_{mit} must be given in hours ($t_{mit} = 0$ corresponds to approximately 5 hours after doxycycline induction, the time when green fluorescence starts and red fluorescence is equal to zero, see Fig. 1C from Ferree et al. (2013)).

At x=0 we assumed that all mitochondria are newly synthesized and thus their age is zero. For the estimate of the mitochondrial age at the axon terminal, we assumed a direct relation between the age of a mitochondrion, t_{mit}, and its distance from the soma (this assumption is supported by Shlevkov and Schwarz 2017):

$$t_{mit} = \lambda x . \tag{77}$$

Combining Eqs. (75), (76), and (77), it follows that

$$\frac{dy}{dx} = \frac{d}{dx}\left(\frac{1}{z}\right) = -\frac{55}{1.5}\frac{1}{\lambda x^2} = -0.5863 . \tag{78}$$

We used Eq. (78) at $x = 200$ μm (one-third of the range displayed in Fig. 4C of Ferree et al. 2013) and solved for λ. The result is

$$\lambda = 1.56 \times 10^{-3} \text{ hours } \mu m^{-1}. \tag{79}$$

Calculating the age of mitochondria at the axon tip (at $x = L_{ax}$) leads to

$$t_{mit,x=L} = \lambda L_{ax} = 15.6 \text{ hours.} \tag{80}$$

This is consistent with the expected half-life of mitochondrial proteins (days or weeks) (Misgeld and Schwarz 2017). It is interesting that data obtained in Mandal et al. (2018) using Timer fluorescent protein support rapid turnover of mitochondria in axon terminals and suggest that the

residence time of the protein in the synapse is approximately 3 hours. Mandal et al. (2021) reported that complete turnover of mitochondria in axon terminals of zebrafish neurons occurs within 24 hours.

3.2. Total length of various populations of mitochondria per unit length of the axon and mean age of mitochondria in various demand sites

Most of the parameter values given in Table 2 are extracted from Agrawal and Koslover (2021) for the case corresponding to the optimal mitochondria health. $k_w = 5 \times 10^{-4}$ s^{-1} was chosen as the base case because, as demonstrated later, the predicted mitochondrial age for this value roughly matches that observed in experimental data.

The compartmental model predicts spatially uniform densities of anterograde, retrograde, and stationary mitochondria throughout the axon (Fig. 3). The same density distributions are obtained by solving the PDE model (20)-(22) (Fig. 4), which confirms the equivalence of the two models. The result is expected because mitochondria should be distributed in an approximately uniform manner throughout the axon. This distribution allows the mitochondria satisfy the energy demand of all demand sites.

Computations with $k_w = 5 \times 10^{-4}$ s^{-1} (Table 2) yield the mean age of stationary mitochondria in demand site # 10 (the most distal demand site) to be approximately 21.6 hours (Fig. 5), which is consistent with the estimate given by Eq. (80). It is known that older mitochondria may have impaired function due to accumulated oxidative damage (Saxton and Hollenbeck 2012). For short axons, oxidative damage is unlikely to be a contributing factor in mitochondrial diseases. For example, since the age of mitochondria at the tip of an axon with a length of 1 cm is only 22 hours, they will most likely not experience oxidative damage.

The mean ages of stationary, anterograde, and retrograde mitochondria increase from the proximal to more distal demand sites. This is in agreement with results reported in Ferree et al. (2013), Shlevkov and Schwarz (2017), which found that the age of mitochondria increased approximately linearly with distance from the soma. As expected, the mean age of anterograde mitochondria is the smallest and the mean age of retrograde mitochondria is the largest.

The mean age of stationary mitochondria of 21.6 hours in the most distal demand site is severalfold greater than the time it takes for the mitochondria to travel from the soma to the most distal demand site without stopping:

$$L_{ax} / v = \left(10^4 \, \mu m\right) / \left(0.5 \, \mu m / s\right) = 2 \times 10^4 \, s = 5.56 \text{ hours} \ll 21.6 \text{ hours}. \qquad (81)$$

Some recent research (Smit-Rigter et al. 2016; Lees et al. 2020) suggest that *in vivo* mitochondria may stay in the stationary state for longer periods of time than *in vitro*. The model parameters can be adjusted to characterize transitions between anterograde, retrograde, and stationary states. This will affect the average time that mitochondria spend stopping near the demand sites in particular neural cells. Although the mean age of mitochondria depends on the values of model parameters, we believe that the important physics of mitochondria transport is captured by our model. Computations with $k_w = 2 \times 10^{-4}$ s^{-1} (2.5 times less the value used in computing Fig. 5) and $p_s = 0.4$ yield the mean age of stationary mitochondria in demand site # 10 (the most distal demand site) to be approximately 31.5 hours (Fig. 6a).

The smaller mean ages of retrograde mitochondria in proximal demand sites in Figs. 5 and 6a are explained by the capture of younger mitochondria from the anterograde state into the stationary state and then their subsequent re-release into the retrograde state (Fig. 2). If the value of parameter p_s that characterizes the probability of capture of motile mitochondria into the stationary state is decreased, the age of retrograde mitochondria increases from the distal (#10) to proximal (#1) demand sites, as mitochondria get older as they are transported back to the soma (Fig. 6b).

3.3. Investigating sensitivity of the mean age of mitochondria in demand sites to model parameters

To investigate the sensitivity of the solution to model parameters, we calculated the local sensitivity coefficients. These are first-order partial derivatives of the observables with respect to model parameters (Beck and Arnold 1977; Zadeh and Montas 2010; Zi 2011; Kuznetsov and Kuznetsov 2019). For example, the sensitivity coefficient of the mean age of resident mitochondria in demand sites to parameter k_w at steady-state (ss) can be calculated as follows:

$$\frac{\partial a_{i,ss}}{\partial k_w} \approx \frac{a_{i,ss}\left(k_w + \Delta k_w\right) - a_{i,ss}\left(k_w\right)}{\Delta k_w}\bigg|_{\text{other parameters kept constant}} \qquad (i=1,\ldots,N), \qquad (82)$$

where $\Delta k_w = 10^{-1} k_w$ (the accuracy was tested by using various step sizes).

The sensitivity coefficients were non-dimensionalized by introducing relative sensitivity coefficients (Zadeh and Montas 2010; Kacser et al. 1995), defined as (for example):

$$S_{k_w}^{\bar{a}_{i,ss}} = \frac{k_w}{a_{i,ss}} \frac{\partial a_{i,ss}}{\partial k_w} \qquad (i{=}1,\ldots,N). \tag{83}$$

The dimensionless sensitivity of the mean age of mitochondria to various model parameters is displayed in Figs. 7-9 and Fig. S1. Out of four investigated parameters, the mean age is most sensitive to parameter ε, less sensitive to parameters p_s and v, and least sensitive to parameter k_w.

The dimensionless sensitivity of the mean age of mitochondria to the restating rate, k_w, is negative in all demand sites (Fig. 7). This is because for the case with a larger restarting rate mitochondria spend less time in the stationary state. The negative sensitivity in Fig. 7 is consistent with the results in Figs. 5 and 6, which indicate that a reduction in k_w increases the mean age of mitochondria. This is because fewer mitochondria (for smaller k_w) escape from the stationary state and transition to the moving (anterograde or retrograde) states.

The dimensionless sensitivity of the mean age of mitochondria to the stopping probability, p_s, is positive in all demand sites (Fig. 8). This is because larger values of p_s correspond to more mitochondria transitioning to the stationary state.

The dimensionless sensitivity of the mean age of mitochondria to the anterograde bias, ε, is positive in all demand sites (Fig. 9). This is because mitochondria that transition to the anterograde state end up spending more time in the axon than mitochondria that transition to the retrograde state, which exit the axon sooner.

The dimensionless sensitivity of the mean age of mitochondria to mitochondria velocity, v, is negative in all demand sites (Fig. S1). This is because the faster movement of mitochondria corresponds to less time that mitochondria spend in the axon.

4. Discussion, limitations of the model and future directions

The model gives quantitative predictions for the distribution of mitochondrial mean age throughout the axon length that can be tested experimentally, indicating that the model has predictive value.

The model predicts that the mean age of stationary mitochondria in the most distal demand site is approximately 21.6 hours. This is severalfold greater than the time it takes for mitochondria to travel without stopping from the soma to the farthest demand location. To quantify the

dependence of the solutions on model parameters, we employed analysis of the dimensionless sensitivity of the mean age of mitochondria at demand sites.

Future development of the model should address the fact that mitochondria that return to the soma from the axon are usually not destroyed, but rather fuse with a resident somatic pool of mitochondria (Misgeld and Schwarz 2017). Therefore, our assumption that mitochondria have zero age at the entrance to the axon is a simplifying one.

Loss/degeneration of mitochondria via mitophagy is an important aspect that needs to be incorporated into the model in future research. However, as Agrawal and Koslover (2021) pointed out, it is difficult to do this within the continuum approach because mitophagy involves selectively removing mitochondria with low membrane potential (in our approach these could be viewed as mitochondria that reached a certain age) by engulfing them in autophagosomes and then transporting them to the soma for degradation in lysosomes. This was done in Agrawal and Koslover (2021) by using a discrete stochastic simulation model.

Acknowledgment

IAK acknowledges the fellowship support of the Paul and Daisy Soros Fellowship for New Americans and the NIH/National Institute of Mental Health (NIMH) Ruth L. Kirchstein NRSA (F30 MH122076-01). AVK acknowledges the support of the National Science Foundation (award CBET-2042834) and the Alexander von Humboldt Foundation through the Humboldt Research Award.

References

Agrawal A, Koslover EF. 2021. Optimizing mitochondrial maintenance in extended neuronal projections. PLOS Computational Biology. 17:e1009073.

Ahola S, Langer T, MacVicar T. 2019. Mitochondrial proteolysis and metabolic control. Cold Spring Harbor Perspectives in Biology. 11:a033936.

Anderson DH. 1983. Compartmental modeling and tracer kinetics. Berlin: Springer.

Beck JV, Arnold KJ. 1977. Parameter estimation in science and engineering. New York: Wiley.

Cagalinec M, Safiulina D, Liiv M, Liiv J, Choubey V, Wareski P, Veksler V, Kaasik A. 2013. Principles of the mitochondrial fusion and fission cycle in neurons. Journal of Cell Science. 126:2187-2197.

Clague MJ, Rochin L. 2016. Parkinson's disease: A traffic jam? Current Biology. 26:R332-R334.

Fan T, Xia L, Han Y. 2001. Mitochondrion and apoptosis. Acta Biochimica Et Biophysica Sinica. 33:7-12.

Ferree AW, Trudeau K, Zik E, Benador IY, Twig G, Gottlieb RA, Shirihai OS. 2013. MitoTimer probe reveals the impact of autophagy, fusion, and motility on subcellular distribution of young and old mitochondrial protein and on relative mitochondrial protein age. Autophagy. 9:1887-1896.

Gottlieb RA, Stotland A. 2015. MitoTimer: A novel protein for monitoring mitochondrial turnover in the heart. Journal of Molecular Medicine-JMM. 93:271-278.

Hernandez G, Thornton C, Stotland A, Lui D, Sin J, Ramil J, Magee N, Andres A, Quarato G, Carreira RS, Sayen MR, Wolkowicz R, Gottlieb RA. 2013. MitoTimer A novel tool for monitoring mitochondrial turnover. Autophagy. 9:1852-1861.

Hollenbeck PJ, Saxton WM. 2005. The axonal transport of mitochondria. Journal of Cell Science. 118:5411-5419.

Jacquez JA, Simon CP. 1993. Qualitative theory of compartmental-systems. SIAM Review. 35:43-79.

Jacquez JA. 1985. Compartmental analysis in biology and medicine. 2nd ed. University of Michigan Press, Ann Arbor, MI.

Kacser H, Burns J, Fell D. 1995. The control of flux. Biochemical Society Transactions. 23:341-366.

Kuznetsov IA, Kuznetsov AV. 2019. Modelling transport and mean age of dense core vesicles in large axonal arbours. Proceedings of the Royal Society A. 475:20190284.

Kuznetsov IA, Kuznetsov AV. 2020. How old are dense-core vesicles residing in en passant boutons: Simulation of the mean age of dense-core vesicles in axonal arbours accounting for resident and transiting vesicle populations. Proceedings of the Royal Society A-Mathematical Physical and Engineering Sciences. 476:20200454.

Kuznetsov IA, Kuznetsov AV. 2019. Investigating sensitivity coefficients characterizing the response of a model of tau protein transport in an axon to model parameters. Computer Methods in Biomechanics and Biomedical Engineering. 22:71-83.

Lees RM, Johnson JD, Ashby MC. 2020. Presynaptic boutons that contain mitochondria are more stable. Frontiers in Synaptic Neuroscience. 11:37.

Malka F, Guillery O, Cifuentes-Diaz C, Guillou E, Belenguer P, Lombes A, Rojo M. 2005. Separate fusion of outer and inner mitochondrial membranes. EMBO Reports. 6:853-859.

Mandal A, Pinter K, Drerup CM. 2018. Analyzing neuronal mitochondria in vivo using fluorescent reporters in zebrafish. Frontiers in Cell and Developmental Biology. 6:144.

Mandal A, Wong HC, Pinter K, Mosqueda N, Beirl A, Lomash RM, Won S, Kindt KS, Drerup CM. 2021. Retrograde mitochondrial transport is essential for organelle distribution and health in zebrafish neurons. Journal of Neuroscience. 41:1371-1392.

Melkov A, Abdu U. 2018. Regulation of long-distance transport of mitochondria along microtubules. Cellular and Molecular Life Sciences. 75:163-176.

Metzler H, Muller M, Sierra CA. 2018. Transit-time and age distributions for nonlinear time-dependent compartmental systems. Proceedings of the National Academy of Sciences of the United States of America. 115:1150-1155.

Metzler H, Sierra CA. 2018. Linear autonomous compartmental models as continuous-time markov chains: Transit-time and age distributions. Mathematical Geosciences. 50:1-34.

Misgeld T, Schwarz TL. 2017. Mitostasis in neurons: Maintaining mitochondria in an extended cellular architecture. Neuron. 96:651-666.

Narayanareddy BRJ, Vartiainen S, Hariri N, O'Dowd DK, Gross SP. 2014. A biophysical analysis of mitochondrial movement: Differences between transport in neuronal cell bodies versus processes. Traffic. 15:762-771.

Patel PK, Shirihai O, Huang KC. 2013. Optimal dynamics for quality control in spatially distributed mitochondrial networks. PLOS Computational Biology. 9:e1003108.

Picard M, McEwen BS. 2018. Psychological stress and mitochondria: A systematic review. Psychosomatic Medicine. 80:141-153.

Prots I, Grosch J, Brazdis R, Simmnacher K, Veber V, Havlicek S, Hannappel C, Krach F, Krumbiegel M, Schuetz O, Reis A, Wrasidlo W, Galasko DR, Groemer TW, Masliah E, Schloetzer-Schrehardt U, Xiang W, Winkler J, Winner B. 2018. Alpha-synuclein oligomers induce early axonal dysfunction in human iPSC-based models of synucleinopathies. Proceedings of the National Academy of Sciences of the United States of America. 115:7813-7818.

Rasmussen M, Hastings A, Smith MJ, Agusto FB, Chen-Charpentier BM, Hoffman FM, Jiang J, Todd-Brown KEO, Wang Y, Wang Y, Luo Y. 2016. Transit times and mean ages for nonautonomous and autonomous compartmental systems. Journal of Mathematical Biology. 73:1379-1398.

Safiulina D, Kaasik A. 2013. Energetic and dynamic: How mitochondria meet neuronal energy demands. PLOS Biology. 11:e1001755.

Saxton WM, Hollenbeck PJ. 2012. The axonal transport of mitochondria. Journal of Cell Science. 125:2095-2104.

Shahmoradian SH, Lewis AJ, Genoud C, Hench J, Moors TE, Navarro PP, Castano-Diez D, Schweighauser G, Graff-Meyer A, Godie KN, Sutterlin R, Huisman E, Ingrassia A, de Gier Y, Rozemuller AJM, Wang J, De Paepe A, Erny J, Staempfli A, Hoernschemeyer J, Grosserueschkamp F, Niedieker D, El-Mashtoly SF, Quadri M, Van IJcken WFJ, Bonifati

V, Gerwert K, Bohrmann B, Frank S, Britschgi M, Stahlberg H, Van de Berg WDJ, Lauer ME. 2019. Lewy pathology in Parkinson's disease consists of crowded organelles and lipid membranes. Nature Neuroscience. 22:1099-1109.

Shlevkov, E., and Schwarz, T. L. (2017). Axonal mitochondrial transport. In P. Verstreken (Ed.), *Parkinson's disease molecular mechanisms underlying pathology* (Parkinson's Disease Molecular Mechanisms Underlying Pathology ed., pp. 113-137). Amsterdam: Elsevier. doi:10.1016/B978-0-12-803783-6.00004-3

Smit-Rigter L, Rajendran R, Silva CAP, Spierenburg L, Groeneweg F, Ruimschotel EM, van Versendaal D, van der Togt C, Eysel UT, Heimel JA, Lohmann C, Levelt CN. 2016. Mitochondrial dynamics in visual cortex are limited in vivo and not affected by axonal structural plasticity. Current Biology. 26:2609-2616.

Twig G, Elorza A, Molina AJA, Mohamed H, Wikstrom JD, Walzer G, Stiles L, Haigh SE, Katz S, Las G, Alroy J, Wu M, Py BF, Yuan J, Deeney JT, Corkey BE, Shirihai OS. 2008. Fission and selective fusion govern mitochondrial segregation and elimination by autophagy. EMBO Journal. 27:433-446.

Uttara B, Singh AV, Zamboni P, Mahajan RT. 2009. Oxidative stress and neurodegenerative diseases: A review of upstream and downstream antioxidant therapeutic options. Current Neuropharmacology. 7:65-74.

Wilson RJ, Drake JC, Cui D, Zhang M, Perry HM, Kashatus JA, Kusminski CM, Scherer PE, Kashatus DF, Okusa MD, Yan Z. 2019. Conditional MitoTimer reporter mice for assessment of mitochondrial structure, oxidative stress, and mitophagy. Mitochondrion. 44:20-26.

Zadeh KS, Montas HJ. 2010. A class of exact solutions for biomacromolecule diffusion-reaction in live cells. Journal of Theoretical Biology. 264:914-933.

Zheng Y, Zhang X, Chen Z. 2019. Mitochondrial transport serves as a mitochondrial quality control strategy in axons: Implications for central nervous system disorders. CNS Neuroscience & Therapeutics. 25:876-886.

Zi Z. 2011. Sensitivity analysis approaches applied to systems biology models. IET Systems Biology. 5:336-346.

Figure captions

Fig. 1. (a) A schematic diagram of a neuron with an axon which is assumed to contain N energy demand sites. Demand sites are numbered starting with the most proximal (#1) to the most distal (#N) *Figure generated with the aid of servier medical art, licensed under a creative common attribution 3.0 generic license. http://Smart.servier.com.*

Fig. 2. A diagram of a compartmental model showing transport in the compartments surrounding the demand sites in the axon. Arrows show mitochondria exchange between the transiting (anterograde and retrograde) states in adjacent demand sites, mitochondria capture into the stationary state and re-release from the stationary state. Mitochondria entering the axon are assumed to have zero age.

Fig. 3. Steady-state values of the total length of stationary, anterogradely moving, and retrogradely moving mitochondria per unit length of the axon in the compartment by the ith demand site. $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.4$.

Fig. 4. Steady-state values of the total length of stationary, anterogradely moving, and retrogradely moving mitochondria per unit length of the axon obtained by solving PDEs (20)-(22) with boundary conditions (25), (26). $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.4$.

Fig. 5. Steady-state values of the mean age of stationary, anterogradely moving, and retrogradely moving mitochondria in the compartment by the ith demand site. $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.4$.

Fig. 6. Steady-state values of the mean age of stationary, anterogradely moving, and retrogradely moving mitochondria in the compartment by the ith demand site for the case of (a) $k_w = 2 \times 10^{-4}$ s^{-1}, $p_s = 0.4$, (b) $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.04$.

Fig. 7. Dimensionless sensitivity of the mean age of mitochondria to the kinetic constant characterizing the rate of mitochondria reentry into motile states, k_w, versus the site number. Sensitivity is analyzed around $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.4$. Computations were performed with $\Delta k_w = 10^{-1} k_w$. A close result was obtained for $\Delta k_w = 10^{-2} k_w$.

Fig. 8. Dimensionless sensitivity of the mean age of mitochondria to the probability that a motile mitochondrion would transition to a stationary state at a demand site, p_s, versus the site number. Sensitivity is analyzed for $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.4$. Computations were performed with $\Delta p_s = 10^{-1} p_s$. A close result was obtained for $\Delta p_s = 10^{-2} p_s$.

Fig. 9. Dimensionless sensitivity of the mean age of mitochondria to the portion of mitochondria leaving the stationary state that join the anterograde pool, ε, versus the site number. Sensitivity is analyzed for $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.4$. Computations were performed with $\Delta\varepsilon = 10^{-1}\varepsilon$. A close result was obtained for $\Delta\varepsilon = 10^{-2}\varepsilon$.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

(a)

(b)

Figure 6

Figure 7

Figure 8

Figure 9

Computation of the mitochondrial age distribution along the axon length

I. A. Kuznetsov[a], [b] and A. V. Kuznetsov[c]

[a]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA

[b]Department of Bioengineering, University of Pennsylvania, Philadelphia, PA 19104, USA

[c]Dept. of Mechanical and Aerospace Engineering, North Carolina State University,

Raleigh, NC 27695-7910, USA; e-mail: avkuznet@ncsu.edu

Supplemental Materials

S1. Supplementary figures

Fig. S1. Dimensionless sensitivity of the mean age of mitochondria to the average velocity of anterogradely/retrogradely moving mitochondria, v (we assumed that $v_a = v_r = v$), versus the site number. Sensitivity is analyzed around $k_w = 5 \times 10^{-4}$ s^{-1}, $p_s = 0.4$. Computations were performed with $\Delta v = 10^{-1} v$. A close result was obtained for $\Delta v = 10^{-2} v$.