Diurnal modulation of multivesicular release controls the

efficiency of information transmission at a sensory synapse

José Moya-Díaz, Ben James, Federico Esposti, Jamie Johnston and Leon Lagnado*

Sussex Neuroscience, School of Life Sciences, University of Sussex,

Brighton BN1 9QG, UK

*Corresponding author: l.lagnado@sussex.ac.uk

Summary

Neuromodulators adjust sensory circuits to changes in the external world or the

animal's internal state and synapses are key control sites for such plasticity. Less

clear is how neuromodulation alters the amount of information transmitted through the

circuit. We investigated this question in the context of the diurnal regulation of visual

processing in zebrafish, focusing on synapses of retinal bipolar cells. We demonstrate

that contrast-sensitivity peaks in the afternoon accompanied by an average four-fold

increase in the Shannon information transmitted at individual active zones. This

increase reflects higher synaptic gain, lower spontaneous "noise" and reduced

variability of evoked responses. Simultaneously, an increase in the probability of

multivesicular events with larger information content increases the efficiency of

transmission (bits per vesicle) by factors of 2-3. This study demonstrates how the

potentiation of multivesicular release by neuromodulators can increase the synaptic

transfer of information and the efficiency of the vesicle code.

Introduction

It has long been understood that the flow of signals through neural circuits is adjusted by neuromodulators[1]. Less clear is how these alter the amount of information that is transmitted through the circuit. Here we investigate this question in the context of visual processing in the retina.

The retina is highly plastic: the input-output relation can adapt within seconds to the recent history of the visual stimulus[2,3] or, on longer time-scales, to changes in the animal's internal state[4,5]. In diurnal animals, for instance, retinal sensitivity to light is regulated both by the daily light-dark cycle and by intrinsic circadian clocks[6-8] . Key to these adjustments is dopamine, a neuromodulator which is released from amacrine cells in a circadian cycle, varying from a minimum at night, increasing during the day and peaking before dusk[6,9]. But the average luminance of a visual scene is not the variable driving most behaviours related to vision: navigation, finding food and avoiding predators all depend on detection of fast modulations in light intensity. We therefore investigated the diurnal control of temporal contrast processing, focusing on the visual signal transmitted by glutamatergic synapses of bipolar cells.

Bipolar cells are the bridge between the photoreceptors and ganglion cells that deliver the results of retinal processing to downstream circuits. Their synaptic compartments are an important control point for transformations of the visual signal[10] and contribute to a number of processing tasks, from adaptive gain control to temporal filtering and the coding of motion, colour, orientation and direction[3,11-13]. Bipolar cells are similar to other sensory neurons, such as photoreceptors, sensory hair cells and electroreceptors, in transmitting information through ribbon synapses containing specialized structures that supply vesicles to the active zone[14]. These sensory synapses do not always operate as Poisson machines in which vesicles are released independently but also signal through multivesicular release (MVR), where the fusion of two or more vesicles is co-ordinated as a single synaptic event[15-17]. The importance of MVR at a number of sites in the brain is now recognized and it has been suggested

that it might contribute to more complex strategies for transmitting information than

modulation of a rate code[18-20].

It is difficult to use Shannon's information theory to measure the amount of

information transmitted at a synapse because the experimenter needs to observe the

symbols conveying the message while also observing or controlling the sensory

input[21,22]. This has recently been achieved by multiphoton imaging of the glutamate

reporter iGluSnFR[23] in bipolar cells of larval zebrafish, where it is found that the visual

message transmitted from an active zone does not use a simple binary code but is

instead composed of a number of symbols, composed of one, two, three or more

vesicles released as one event[17]. Here we demonstrate that this strategy of coding by

amplitude as well as rate is under diurnal control. Synaptic responses to temporal

contrast reach a maximum in the afternoon and are accompanied by a four-fold

increase in the Shannon information transmitted at each active zone compared to the

morning. Dopamine contributes to this increase in information transfer by reducing

several aspects of synaptic "noise" and by increasing the probability of multivesicular

events with larger information content, which in turn increases the efficiency of

transmission quantified as bits per vesicle.

Results

Differential regulation of luminance-sensitivity and contrast-sensitivity

To investigate the diurnal modulation of visual processing in larval zebrafish we began

by imaging synaptic activity in bipolar cells with SyGCaMP2[24] (Fig. 1A). When

animals were placed on a cycle of 14 hours light and 10 hours dark, no significant

synaptic responses could be detected at *Zeitgeber* times 18-0 hours, consistent with

previous observations that larvae are blind at subjective night[25]. Visual sensitivity

began to recover within 20 mins of light onset, after which responses gradually

increased in amplitude (Fig. S1A and Fig. 1B). Plotting the irradiance-response

functions (Fig. 1C) allowed the luminance sensitivity to be quantified as the inverse of

the irradiance generating a half-maximal response ($1/I_{1/2}$). Over the course of the day,

luminance-sensitivity increased gradually over a range greater than 200-fold (Fig. 1D).

As in other species, this increase could be explained largely by actions of D2

dopamine receptors because injection of the antagonist sulpiride (~2 µM) reduced

luminance-sensitivity in the afternoon to levels measured in the morning[6] (Fig. S1).

Figure 1: Differential regulation of luminance-sensitivity and contrast-sensitivity

A. Left: Retina of a Ribeye::SyGCaMP2 fish with box over the IPL. Right: expansion of the boxed region of

showing terminals of bipolar cells. **B.** Averaged responses from ON terminals to light steps of different

irradiance measured at *Zeitgeber* time 1, 10 and 16 hours. Note large variations in amplitude and

kinetics. Each light step was of 3 s (*n* = 535 terminals from 10 fish). **C.** Peak response as a function of

irradiance for ON terminals in B. The smooth lines are Hill functions of the form $R = R_{max}*(I^h/(I^h + I_{1/2}^h))$,

where R is the peak response, I is the irradiance, h is the Hill coefficient and $I_{1/2}$ is the irradiance

generating the half-maximal response. At ZT = 16 hrs: R_{max} = 0.91 ± 0.04; h = 2.0 ± 0.2; $I_{1/2}$ = 0.066 ±

0.02 nW/mm^2 (dashed blue arrow). At ZT = 10 hrs: R_{max} = 0.85 ± 0.06; h = 0.8 ± 0.1; $I_{1/2}$ = 0.65 ± 0.18

nW/mm^2. At ZT = 1 hrs: R_{max} = 0.853 ± 0.02; h = 0.9 ± 0.2; $I_{1/2}$ = 0.88 ± 0.18 nW/mm^2 (red arrow). **D.**

Variations in luminance sensitivity as a function of *Zeitgeber* time averaged across both ON and OFF

terminals (n=535 and 335, respectively). The lower bar shows the timing of the light-dark cycle. **E.**
Averaged responses to stimuli of different contrasts measured at *Zeitgeber* time 4, 7 and 13 hrs. **F.** Peak
response amplitude as a function of contrast for terminals shown in E. The smooth lines are Hill functions
used to interpolate values of $C_{1/2}$, the contrast generating the half-maximal response. Note the diurnal
variations. At ZT = 4 hrs: $C_{1/2}$ = 86 ± 2% (dashed red arrow); h = 7.0 ± 1.2. At ZT = 7 hrs: $C_{1/2}$ = 35 ± 2%
(dashed black arrow); h = 2.7 ± 0.2. At ZT = 13 hrs: $C_{1/2}$ = 72 ± 2%; h = 3.3 ± 0.2. **G.** Variations in
contrast-sensitivity as a function of *Zeitgeber* time averaged across ON and OFF terminals. Note the
peak around ZT = 7 hours which is not mirrored in the diurnal variation in luminance sensitivity (D). The
grey bars show the periods described as "morning" and "afternoon". All error bars show ± 1 SD.

The detection of modulations in light intensity (contrast) was also under diurnal

control, but with a distinctive time-course (Fig. 1E-G; 5 Hz full-field stimuli). At ZT = 4

hours, temporal contrasts below 50% were barely detected and the half-maximal

response ($C_{1/2}$) was generated by a contrast of 86 ± 2 % (Figs. 1E and F). But at ZT =

7 hours $C_{1/2}$ it fell to 35 ± 2 % with responses saturated above 50%. When contrast

sensitivity ($1/C_{1/2}$) was mapped during the course of the day it was relatively constant

at ZT 1-5 hours and ZT 9-14 hours but increased to levels ~2.4-fold higher around ZT

= 7 hours (Fig. 1G). Notably, this peak in the contrast sensitivity of the retinal circuit

occurred at a similar *Zeitgeber* time as the maximum contrast sensitivity measured

behaviourally using the optokinetic reflex[8,26]. A qualitatively similar increase in contrast

sensitivity was also observed at the retinal output projecting to the optic tectum (Fig.

S2).

Diurnal regulation of contrast gain

To measure transmission of the visual signal in terms of its elementary units –

synaptic vesicles - we expressed the reporter iGluSnFR[23] sparsely in bipolar cells (Fig.

2A). Wiener deconvolution of iGluSnFR signals allowed us to count released vesicles

(see Methods and Fig. S3; detailed evidence that these methods allow signals to be

isolated from individual active zones vesicles has been described[17]). Synaptic function

was compared over a two-hour period beginning 1 hour after light onset ("morning")

with a two-hour period beginning 6 hours later ("afternoon"; Fig. 1G).

134
135
Figure 2. Diurnal modulation of synaptic gain
A. Multiphoton section through the eye of a zebrafish larva (7 dpf) expressing iGluSnFR in a subset of
bipolar cells. **B**. Examples of iGluSnFR signals from an individual OFF synapse elicited using a stimulus of
variable contrast modulated at 5 Hz (0-100%, full field, sine wave) in the morning (ZT 1-3 hours, grey) and
afternoon (ZT 6-9 hours, black). Note the high levels of spontaneous activity in the morning (black
arrowheads). In each case the top trace shows the iGluSnFR signal and the lower trace the estimated
number of quanta composing each event (Q_e). **C.** Average contrast-response functions in OFF bipolar
cell synapses in the morning (open circles; n = 20 synapses) and afternoon (closed; n = 59), where the
response (*R*) was quantified as the average of quanta per cycle (Q_c). The smooth lines are fits of a
sigmoid used for smoothing. Note the differences in the shape of the contrast-response functions and in
the levels of spontaneous activity (zero contrast). **D.** Average contrast-response functions in ON bipolar
cell synapses in the morning (open circles; n = 12 synapses) and afternoon (closed; n = 31). There was
no significant difference in in the morning relative to afternoon (Chi-square test, p = 0.9999). **E.** The
contrast gain calculated as the derivative of the fits to the contrast-response functions in C and D. Note
that the maximum contrast discrimination is increased by a factor of 2x in the OFF channel during the
afternoon.

152
153

Examples of glutamate transients at an individual OFF active zone are shown in

Fig. 2B. Across a range of contrasts, responses were, on average, larger in the

afternoon. We began by measuring the contrast-response function (CRF) simply as

the average number of vesicles released per cycle of a 5 Hz stimulus, choosing this

frequency because the integration time of a bipolar cell is ~200 ms[27]. There was little

diurnal modulation of the CRF measured at ON synapses but in the OFF channel the

maximum rate of release measured at 100% contrast increased from 15.25 ± 2.5

vesicles/s in the morning to 25.5 ± 1.5 vesicles/s in the afternoon (Fig. 2C and D).

This increase in synaptic gain was accompanied by an increase in contrast sensitivity,

and the combined effects were assessed as the derivative of the CRF ("contrast gain";

Fig. 2E). Contrasts in natural visual scenes rarely exceed 40%[12] and in the morning

this range was signalled best through the ON channel. But in the afternoon the OFF

channel became dominant, with contrast gains increasing by factors of 2-6.

Dopamine regulates contrast gain

To test whether dopamine contributes to diurnal changes in contrast sensitivity we

injected agonists or antagonists of dopamine receptors directly into the eye. Fig. 3A

shows examples of the output from a synapse imaged in the afternoon, before and

after injection of the D1 receptor antagonist SCH 23390 (estimated final concentration

of 0.1 μM). Counteracting the actions of endogenous dopamine reduced the average

rate of vesicle release and shifted the CRF such that the maximum contrast gain was

achieved at higher contrasts (black points in Fig. 3B and C). Conversely, increasing

activation of D1 receptors in the morning by injection of the agonist ADTN (~0.2 μM)

increased response gain.

Figure 3. Diurnal changes in dopamine levels modulate synaptic transmission.
A. Examples of iGluSnFR signals recorded in the afternoon from an individual OFF (red trace) and ON (green trace) synapses elicited using a stimulus of variable contrast before and after intravitreal injection of the D1 antagonist, SCH 23390 (black traces; 5 Hz modulation). Note that SCH 23390 abolished synaptic responses at lower contrasts in ON and OFF synapses. In each case the top trace shows the iGluSnFR signal and the lower trace the estimated Q_e. **B.** Average contrast-response functions in OFF bipolar cell synapses after administration of D1 antagonist (black dots) in the afternoon and after administration of the D1 agonist ADTN in the morning (blue dots). Each point shows the mean ± s.e.m. (SCH 23390, n =12 synapses; ADTN, n = 12 synapses). Control responses observed in the morning and afternoon are superimposed in the graph (red lines, see Fig. 2C) **C.** Average contrast-response functions in ON bipolar cell synapses in three conditions: afternoon (green dots), after intravitreal injection of D1 antagonist in the afternoon (black dots) and ADTN in the morning (blue dots). Each point shows the mean ± s.e.m. (SCH 23390, n =7 synapses; ADTN, n = 5 synapses). Control responses observed in the morning and afternoon are superimposed to the graph (green lines, see Fig. 2D). **D.** Relative response gain by diurnal modulation and after manipulation of dopaminergic signalling (dashed lines). Note that diurnal modulation of synaptic gain is higher in OFF synapses, whereas dopamine modulates the dynamic range by ~16 fold-change in ON and OFF synapses.

The dynamic range over which D1 receptors adjusted synaptic gain was calculated as the ratio of the CRFs in the presence of the agonist and antagonist: in both ON and OFF channels the maximum modulation was ~16-fold, occurring at

contrasts of 20-40% (Fig. 3D). But diurnal modulation of gain was narrower than this potential range: 1.7-fold in OFF synapses and 1.1-fold in ON.   This difference reflected, at least in part, a gain in the morning that was at least 5-fold higher than that measured with D1R receptors blocked, consistent with dopamine levels that were already high enough to potentiate synaptic transmission (Fig. 3B and C). These manipulations of retinal dopamine receptors caused qualitatively similar changes in the signals that ganglion cells transmit to the optic tectum (Fig. S2).

Modulation of synaptic noise and variability

How does diurnal modulation of contrast processing affect the information transmitted to ganglion cells? In the framework of information theory[21], an increase in synaptic gain will tend to reduce uncertainty (and therefore increase information) by causing a larger change in the number of vesicles released when contrast changes. But information is degraded by "noise" that causes responses to vary when the same stimulus is repeated and synapses are a major source of such variability within neural circuits[28,29]. One cause of synaptic noise is the stochasticity of the presynaptic processes that control the fusion of vesicles[18] and this was a prominent feature of the output from synapses of bipolar cells (Figs. 2B). We distinguished four aspects of synaptic variability and investigated the diurnal modulation of each; i) spontaneous vesicle release (Fig. 4), ii) variability in the number of vesicles released by a stimulus (Fig. 5), iii) variability in the timing of release events i.e how tightly they are time-locked to the stimulus (Fig. 6) and iv) modulation of multivesicular release (Fig. 7). Finally, we calculated how these different aspects of synaptic "noise" combined with changes in contrast gain (Fig. 3) to alter the amount of visual information transferred from individual active zones (Fig. 8).

Figure 4. Diurnal modulation of spontaneous synaptic noise

A.*Top:* Example of iGluSnFR signals from an individual OFF synapse elicited using a stimulus of variable
contrast in the morning (0-100%, 5 Hz modulation). In this example, note the high levels of spontaneous
activity that were quantified as the responses elicited at zero contrast (red dashed box) . *Bottom.*
Examples of iGluSnFR signals from the same OFF synapse after intravitreal injection of ADTN. Note the
increase in amplitude and frequency of events and the reduction of spontaneous activity. In each case the
top trace shows the iGluSnFR signal and the lower trace the estimated Q_e. **B.** Quantification of
spontaneous events composed by different Q_e in OFF synapses in the morning, morning + ADTN and
afternoon (OFF morning, n = 20 synapses; OFF morning + ADTN= 12 synapses; OFF afternoon, n=24
synapses). Note the suppression of spontaneous events in OFF synapses after intravitreal injection of
ADTN in the morning. **C.** Quantification of spontaneous events composed by different Q_e in ON synapses
in the Morning, Morning + ADTN and Afternoon (ON Morning = 12 synapses; ON Morning + ADTN = 5
synapses; ON Afternoon, n =17 synapses). Note that spontaneous activity levels were not dramatically
altered after administration of ADTN.

i) Spontaneous release

Increases in synaptic gain were accompanied by a *decrease* in the spontaneous

release of vesicles in the absence of a visual stimulus. In the morning, spontaneous

release occurred at relatively high rates (Fig. 4A) composed of both univesicular and

multivesicular events (Fig. 4B and C). Integrating across events of all amplitudes, the

average rate of spontaneous release in OFF synapses was 22.5 ± 12.5 vesicles s^{-1} in

the morning, falling to 5 ± 1 vesicles s^{-1} in the afternoon (Fig. 4B). In ON synapses

these values were 9 ± 4 vesicles s^{-1} and 2.5 ± 1 vesicles s^{-1} (Fig. 4C). In both channels,

therefore, spontaneous noise was ~4 times lower in the afternoon compared to the

morning. Increased activation of D1 receptors suppressed spontaneous release in the

morning to levels close to those measured in the afternoon, but only in OFF synapses

(Fig. 4B and C).

ii) Variability in stimulus-evoked responses.

When recording neural responses as spikes, the Fano factor is measured as the ratio of

the variance-to-mean of spikes counted in a fixed time-window after a repeated

stimulus[30,31]. We calculated the Fano factor of a synapse by counting the number of

vesicles released over each cycle of a sinusoidal stimulus (Fig. 5A). In the morning, F

was ~2.6 in both ON and OFF synapses when averaged over a range of contrasts, falling

to ~1.6 in the afternoon (both significant at $p < 0.002$, KS test; Fig. 5B-C). The increase

in contrast gain and sensitivity in the afternoon (Fig. 2C-D) was therefore also associated

with increased reliability of bipolar cell synapses. Notably, the variability of synaptic output

was higher than expected for a Poisson process, for which the Fano factor is one. The

spike responses of post-synaptic RGCs are less variable, with a Fano factor as low as 0.3

at higher contrasts[30], likely reflecting the integration of signals from multiple synaptic

inputs. Activation of D1 receptors in the morning improved the reliability of synaptic

responses to levels similar to those measured in the afternoon (Figs. 5D and E).

Figure 5. Diurnal changes in the variability of stimulus-evoked responses

A. Examples of iGluSnFR signals from individual OFF synapses in the morning and afternoon. Responses

elicited by stimuli of 60% and 40% contrast varied from cycle to cycle of the 5 Hz stimulus. In each case the

top trace shows the iGluSnFR signal and the lower trace the estimated Q_e **B.** Variability in the response of

OFF synapses calculated as the Fano factor, where each response was quantified as the total number of

vesicles released over one cycle at the contrasts shown. Comparison is made between the morning (n=18),

afternoon (n=27) and the morning after injection of ADTN (n=13). **C.** As in B, but for ON synapses (n = 12, 15

and 6 synapses for respective conditions). **D.** Average Fano factor over different contrasts in OFF synapses

in the three conditions described above. Overall, the average fano factor was significantly higher in the

morning compared to afternoon or in the morning after injection of ADTN (t-test; p<0.0001). **E.** As D, but for

ON synapses. Again, the average Fano factor was significantly higher in the morning (t-test; p<0.001).

Figure 6. The temporal precision of MVR is under diurnal control in the OFF channel

A. Example recordings from two OFF synapses stimulated at 60% contrast in three conditions: afternoon (top, black trace), morning (middle, red trace) and after intravitreal injection of ADTN in the morning (bottom, blue trace). Morning and morning + ADTN synaptic responses are from the same synapse. The modulation in intensity (5 Hz, sine wave) is shown below. Arrowheads highlight events occurring at different phases of the stimulus, with less variation with events composed for 4 or more quanta in the afternoon and after administration of ADTN in the morning. In each case the top trace shows the iGluSnFR signal and the lower trace the estimated Q_e. **B.** Temporal jitter of events composed of different numbers of quanta in OFF synapses in the afternoon (red dots ; n = 24 synapses); Morning (open red dots; n = 19 synapses) and Morning + ADTN (blue dots, n = 16). Note that during the morning events composed by multiple quanta were less phase-locked to the stimuli in comparison to the afternoon. Activation of D1 receptors had a significant effect on release of multiquantal events. Events composed by 5 or more quanta jittered by ~7 ms, similar to values observed in the afternoon. The solid lines describing these relations in the three conditions are better described by a single exponential decay function of the form $y_0 + A_{exp}((-(x-x_0)/\tau)))$ with y_0= 4.23 ± 1.2 and A = 27 ± 7 in the afternoon; y_0= 9.77 ± 1.4 and A= 28.64 ± 5.6 in the morning and y_0=5.45 ± 1.3, A = 30. ± 6.1 after activation of D1 receptor in the morning. **C.** Temporal jitter of events composed by different numbers of quanta measured in ON synapses in the afternoon (green dots; n = 14 synapses) during the (open green dots; n= 10 synapses) and during Morning + ADTN, (blue dots; n=6 =synapses). Activation of D1 receptor did not have a significant effect in the temporal precision in the ON channel. The relationship observed in the morning is better described by a straight line with a = 34.7 ± 1.5 and a slope = -3.6 ± 0.5.

iii) Temporal jitter.

Retinal ganglion cells (RGCs) encode information not just in their spike count but also

in the timing of spikes[30,32]. Spike times can vary by just a few milliseconds and this

accuracy depends on the precision of excitatory inputs received from bipolar cells[33].

The standard deviation in timing of release events ("temporal jitter") was measured

relative to the phase of a 5 Hz stimulus (60% contrast; Fig. 6A) and the larger the

release events the more precise it was on average (Fig. 6B-C). In OFF synapses the

temporal jitter was 5-8 ms higher in the morning compared to the afternoon for events

composed of up to 8 vesicles (Fig. 6B; $p < 0.008$, Kolomogorov-Smirnov test). Diurnal

modulation of temporal precision was weaker in ON synapses and only significant for

events composed of 1-3 vesicles (Fig. 3H; t-test at each Q_e). Increasing activation of

D1 receptors in the morning reduced temporal jitter in events composed for multiple

quanta in OFF synapses ($p < 0.05$; KS test) but not ON (Fig. 6B and C; $p > 0.5$).

Diurnal variations in dopamine therefore modulate the temporal accuracy of vesicle

release.

iv) Changes in the distribution of multivesicular events

Previous studies quantifying the synaptic transfer of visual information have been

limited by the inability to monitor individual active zones and used the assumption that

vesicles are released according to Poisson statistics[34,35]. But we now know that bipolar

cells do not employ a simple rate-code and visual information is also contained in the

amplitude of multivesicular events[17]. We therefore tested whether modulation of

contrast gain was accompanied by changes in Q_e, the number of quanta in an event.

A comparison of the distribution of Qe in the morning and afternoon is shown in

Fig. 7 for responses to a stimulus of 60% contrast. In ON synapses, 68% of release

events in the morning were univesicular, falling to 40% in the afternoon and reflecting

a shift in the distribution towards larger events (Fig. 7B; $p < 0.05$, KS test). This shift

towards MVR was fully reversed by antagonizing D1 receptors by injection of SCH

23390 (Fig. 7C). In the morning, MVR was more prevalent in OFF synapses with only

38% of release events being univesicular but again there was a significant shift

towards larger events in the afternoon (Fig. 7D; p<0.02). Blocking the D1 actions of

endogenous dopamine had a stronger effect in OFF synapses, increasing the

proportion of univesicular events to 66% in the afternoon (Fig. 7E; p < 0.001).

Qualitatively similar modulation of MVR was observed over a range of contrasts from

20% to 80% and blocking D1 receptors in the afternoon shifted the distribution back to

univesicular release in both ON and OFF channels (Fig. 7D and E). Diurnal variations

in dopamine therefore modulate MVR.

Figure 7. Dopamine contributes to diurnal variations in the distribution of multivesicular events

A. Examples of iGluSnFR signals from individual synapses elicited using 60% contrast stimulus (5 Hz,

30 sec) in the morning (top), afternoon (middle) and afternoon + SCH 23390 (bottom). In each case the top

trace shows the iGluSnFR signal and the lower trace the estimated Q_e. **B**. Changes in Q_e in ON synapses

in the morning and afternoon. In the afternoon the distribution was shifted toward multiquantal events

(p<0.059, Chi-squared test). **C.** Changes in the distribution of Q_e in ON synapses before and after intravitreal injection of the D1 antagonist SCH23390. The distribution was shifted toward lower Q_e (p<0.001) but was not significantly different to that measured in the morning. **D.** Changes in Q_e, in OFF synapses in the morning and afternoon. In the afternoon the distribution was shifted toward multiquantal events (p<0.007). **E.** Changes in the distribution of Q_e in OFF synapses before and after intravitreal injection of SCH 23390 in the afternoon. The distribution was shifted toward uniquantal events (p<0.001).

Modulation of information encoded at the synapse

How do changes in synaptic gain (Figs. 2-3), noise (Fig. 4-6) and MVR (Fig. 7) combine to alter the amount of visual information transmitted by the synapses of bipolar cells? A larger synaptic signal relative to noise (SNR) will tend to increase the mutual information (I) between the response (q) and the stimulus generating it (S), although the size of the increase will depend on the statistical properties of both signal and noise[36]. In the simple situation where both have a Gaussian distribution I = $0.5\log_2(1+SNR)$. But how should we quantify the synaptic signal? When analyzing the spike code, all events comprise the same symbol and the response can be described as the number of spikes in each of a series of time bins[22,36]. The output from bipolar cells is qualitatively different with a visual stimulus being encoded both by the timing of release events and their amplitudes[17]. We therefore took an approach in which MVR composed of different numbers of vesicles were considered different symbols[21,37]. The mutual information between the response and stimulus was then computed as the average amount of information about the stimulus gained from observing any symbol (see Methods).

The stimulus set S comprised 12 different contrasts but these were not fixed for each synapse because the contrast sensitivity varied between synapses and between morning and afternoon (Fig. 1E-G). To make allowance for this, we used contrasts spanning ±10% around $C_{1/2}$ measured within the synapse under study immediately before delivering the stimulus set. In the absence of information about the distribution of contrasts normally experienced by a larval zebrafish, a uniform distribution of

contrasts was used for S. Each contrast step lasted 2 s (5 Hz) and they were presented

in two different pseudo-random orders, of which one is shown in Fig. 8A.

Figure 8. Diurnal changes in the efficiency with which synapses transmit visual information

A. Examples of synaptic responses over 12 different contrasts spanning ±10% around the contrast

eliciting the half-maximal response ($C_{1/2}$) in the morning (top, light red), afternoon (middle, dark red) and

after injection of D1 antagonist SCH 23390 in the afternoon (bottom, black; note the lower frequency and

amplitude of release events). In each case the top trace shows the iGluSnFR signal and the lower trace

the estimated Q_e. Each contrast step lasted 2 s (5 Hz) and each trace is from a different OFF synapse. **B.**

Mutual information $I (S;Q)$ in four conditions: (i) morning, (ii) afternoon, iii) morning after injection of ADTN,

(iv) afternoon after injection of SCH 23390. **C.** Specific information (I_2) for events of different quantal

content in OFF synapses (33 synapses). The curve describing the relation is a least-squares fit of a power

function of the form $i = AQ_e^x$, with A = 0.20, and x = 1.81. **D.** As C, but for ON synapses (n = 13). The

curve describing the relation is almost identical (A = 0.21, and x = 1.75).

In the morning, the average mutual information between stimulus and response

was almost exactly the same for synapses in the ON and OFF channels (0.445 ±

0.035 bits s^{-1} and 0.455 ± 0.03 bits s^{-1}, respectively). In the afternoon mutual

information increased through both channels although the increase in OFF synapses

(370%) was significantly larger than in ON (270%; p < 0.001; Fig. 8B). In OFF

synapses, the maximum mutual information of 2.1 bits s^{-1} was associated with average

release rate of 2.5 vesicles s^{-1} around $C_{1/2}$, equivalent to an efficiency around 0.8 bits

per vesicle.

Several of the synaptic properties we have analyzed will contribute to the

improvement in information transmission in the afternoon, including the increase in

synaptic gain (Fig. 2), the decrease in spontaneous noise (Fig. 3) and reduced

variability of stimulus-evoked responses (Figs. 5 and 6). These aspects of synaptic

transmission were all subject to modulation by dopamine and, consistent with these

changes, mutual information in the morning was increased by activation of D1

receptors while in the afternoon it was decreased by antagonizing the effects of

endogenous dopamine (Fig. 8B).

These results demonstrate that information transmission through the retina is

under diurnal control and that dopamine is a key neuromodulator controlling these

changes. Antagonizing D1 receptors did not, however, reduce mutual information to

levels measured in the morning, leaving open the possibility that other signaling

pathways also contribute.

Changes in the efficiency of the vesicle code

The transmission of information using spikes and vesicles is the major consumer of

energy in the brain with one estimate being of the order of ~24,000 ATP molecules per

bit[38,39]. The largest part of this energy consumption is taken up by synaptic

transmission so a key question becomes the effect of neuromodulation on the

efficiency with which vesicles are used to transmit information. Strikingly, the 2.7-fold

increase in information transmitted through ON synapses in the afternoon (Fig. 8B)

was *not* associated with any change in the average rate of vesicle release (Fig. 2D and

E), while the 3.7-fold increase in OFF synapses was associated with only a 2-fold

increase in the rate around $C_{1/2}$ (Fig. 2C and E). The diurnal increase in synaptic gain

was therefore associated with a 1.4- to 2.7-fold increase in the average efficiency with

which vesicles were used to encode changes in contrast. A comparison can be made

with the information transmitted by spikes in RGCs, where the most sluggish cells

transmit ~3.5 bits/spike, while those that fire most briskly encode ~2 bits/spike[40]. An

increase in spike rate is therefore associated with a *decrease* in the visual information

per spike while an increase in vesicle release rate is associated with an *increase* in

information per vesicle.

How is this increase in the efficiency of the vesicle code achieved? The

comparison of information transmission with average rates of vesicle release obscures

a key aspect of the vesicle code operating in bipolar cells: information about contrast is

represented as changes in both the rate and amplitude of release events[23]. This is

significant because the distribution of MVR events was also a function of *Zeitgeber*

time and larger events are rarer and carry more specific information[17], as shown by the

supralinear relation between the specific information carried by each synaptic symbol

and the number of vesicles it contains (Fig. 8C-D). The diurnal modulation of the

efficiency of the vesicle code therefore depends on the shift between univesicular and

multivesicular release (Fig. 7B and D).

We also considered the possibility that modulation of retinal processing might be

accompanied by changes in the amount of information carried by a given synaptic

symbol. The relation between information carried and event amplitude did not,

however, change in the afternoon compared to the morning, at least for events

composed of 1-6 vesicles (Fig. 8C-D; OFF synapses p>0.99; ON synapses p>0.98; KS

test).

Discussion

The plasticity of synapses allows the flow of information through circuits to be modulated[1] and this study provides a quantitative understanding of this idea in the context of the diurnal control of visual processing in the retina. We find that the daily light-dark cycle alters the transmission of visual information through bipolar cells by factors of ~4 during daylight hours by adjusting four synaptic properties; the number of vesicles released by a stimulus (Fig. 3), spontaneous synaptic noise (Fig. 4), the variability of stimulus-driven responses (Figs. 5-6) and the balance between univesicular and multivesicular release (Fig. 7). Crucially, the switch in emphasis from univesicular to multivesicular release also increases the amount of information transmitted per vesicle (Fig. 8). Dopamine plays a major role in regulating all these aspects of retinal function although the relative contributions of these mechanisms differed between ON and OFF pathways.

Diurnal modulation of gain

Dopamine-dependent changes in the synaptic gain of bipolar cells might be caused either by direct modulation of processes within the terminal compartment or by actions on the circuitry in which they are embedded. A direct action is strongly supported by the presence of dopamine receptors (especially D1) on the terminal compartment of bipolar cells[41,42] and electrophysiological experiments demonstrating that their activation potentiates L-type calcium channels that control vesicle fusion[4]. Dopamine also acts on D2 receptors on cone synapses to potentiate the visual drive to bipolar cells but this mechanism alone does not easily explain the *transient* increase in contrast-sensitivity in the afternoon given that luminance sensitivity, a much more direct reflection of the strength of cone input, gradually increases throughout the day (cf. Fig. 1B and Fig. 1E). The gain of bipolar cell synapses is also strongly dependent on the inhibitory inputs that the synaptic compartments receives from amacrine cells and the possibility of diurnal modulation of inhibition remains open.

Dopamine release is controlled by the internal circadian clock as well as changes

in luminance[7] or the appearance of food-related odours[4]. But other neuromodulators,

are also released from amacrine cells, including melatonin[43], Substance P[44] and

somatostatin[45] and some of these can antagonize the actions of others[44]. A large

number of different proteins control the activity of the retinal circuit and 17% of genes

in zebrafish are under circadian regulation[7]. There is therefore a good possibility that

neuromodulators other than dopamine will also act on the synaptic output of bipolar

cells, either directly or indirectly, to regulate the visual signal transmitted to ganglion

cells.

Diurnal modulation of noise

It has long been appreciated that synaptic noise can reduce the amount of information

transmitted through a circuit of neurons[28]. When the retina operates under photopic

conditions, for instance, the release of vesicles from bipolar cells adds noise to the

signal arriving from cones and therefore causes a loss of information in RGCs[46]. It has

been suggested, however, that under other circumstances the noise in synaptic

transmission might improve information transmission, such as when stochastic

resonance increases the probability of post-synaptic depolarization crossing threshold

for spike generation[29,47]. But it seems unlikely that the retina operates under such a

regime, given that diurnal increases in synaptic gain went hand-in-hand with a

reduction in several sources of noise, including spontaneous release unrelated to a

stimulus.

All the changes in synaptic function that we observed comparing periods in the

morning and afternoon were mimicked by manipulating dopamine signalling, indicating

that this neuromodulator adjusts information transmission by orchestrating changes in

both the signal and the various noise sources that cause it to vary. The balance

between modulation of signal and noise was however, strikingly different in the ON

channel, where synaptic gain was *not* under diurnal modulation, compared to the OFF

channel, where both signal and noise were regulated. The processes by which

dopamine and other neuromodulators adjust synaptic noise are also likely to involve

both direct actions on the synaptic compartment and indirect actions on other

components of the retinal circuit.

Modulation of multivesicular release

MVR is not just a property of ribbon synapses but is also a feature of synaptic

transmission in the hippocampus[48], cerebellum[49] and somatosensory cortex[19], where

arrival of a spike can often trigger release of two or more vesicles at an active zone. A

recent combination of electrophysiology with correlative light-and electron-microscopy

has even led to the suggestion that MVR may be a fundamental mode of synaptic

transmission throughout the nervous system[20]. It is also recognized that MVR can be

adjusted by neuromodulation, for instance through muscarinic acetylcholine receptors

in the striatum{Higley, 2009 #221} or GABA$_B$ receptors in the cortex{Chalifoux, 2010

#222}, athough the implications for information transmission in these contexts is not

known. Our study has demonstrated that potentiation of MVR in the retina not only

increases the amount of information that a synapse can transmit using vesicles but

also the efficiency of coding. It will be interesting to establish how far neuromodulators

acting in other parts of the brain alter the efficiency of the information transmission and

how far this involves modulation of MVR as compared to the variability and noise that

is a feature of central synapses[28,29]. A crucial aspect of these questions will be to

understand how switching from univesicular release to multivesicular release alters the

spike code generated post-synaptically.

Acknowledgements

The authors express many thanks to all the members of Lagnado laboratory for

discussion. We also thank Tom Baden for his many insightful criticisms and suggestions.

This work was supported by grants to L.L. from the Wellcome Trust (102905/Z/13/Z).

Author contributions

J. M-D. conceived, designed and executed experiments, analyzed results and prepared

the manuscript. B. J. carried out analysis and wrote code. F. E. conceived and executed

experiments and carried out analysis. J. J. conceived and executed experiments and

carried out analysis. L. L. conceived the project, designed experiments, analyzed data,

repaired equipment, wrote code and prepared the manuscript.

Supplementary Information

Supplementary Figure 1

Figure S1. Diurnal changes in luminance sensitivity are co-ordinated by dopamine

A. Left: Averaged SyGaMP2 signals at different *Zeitgeber* time. ON terminals green and OFF terminals red. Each step of light (10 nW mm^{-2}) lasted 3 s. These averages are only from responsive terminals. Right: the percentage of terminals generating a significant response to the same light step (averaged across both ON and OFF). Bars show SD. **B.** Effects of manipulating dopamine signalling on luminance sensitivity of the ON channel. Luminance vs. response plots for ON terminals. Red circles compare this function at ZT 1 hr under control conditions (solid circle) and after injection of the non-selective dopamine receptor agonist ADTN (~0.2 μM; open circles). ADTN caused a prompt change in the luminance-response function to forms measured at ZT 10 hrs (solid black circles), increasing R_{max} from 0.53 ± 0.02 to 1.02 ± 0.07, and reducing $I_{1/2}$ from 0.88 ± 0.18 nW mm^{-2} to 0.05 ± 0.02 nW mm^{-2} (± sd, as estimated from the fitted Hill function shown). The higher gain and luminance sensitivity at ZT 10 hrs could be explained as an effect of dopamine at D2 receptors, because it was completely reversed by injection of the selective D2 receptor antagonist sulpiride (~2 μM; open black circles; R_{max} = 0.57 ± 0.13, $I_{1/2}$ = 1.16 ± 1.34 nW mm^{-2}). Results collected from *n* = 535 terminals from 38 fish. **C.** Effects of manipulating dopamine signalling on luminance sensitivity of the OFF channel. Comparing control responses at ZT 1 hr and 10 hrs showed a significant reduction in $I_{1/2}$ from 3.9 ± 1.3 to 0.0128 ± 0.005, but *without* a significant change in R_{max} (0.35 ± 0.03 vs 0.30 ± 0.02). ADTN injected at ZT 1 caused a prompt increase in luminance sensitivity, reducing $I_{1/2}$ to 0.013 ± 0.005 nW mm^{-2}. The higher luminance sensitivity at ZT 10 hrs could be partly explained as an effect of dopamine at D2 receptors, because injection of sulpiride (~2 μM; open black circles) increased $I_{1/2}$ from 0.05 ± 0.01 nW mm^{-2} to 0.35 ± 0.14 nW mm^{-2}. Results collected from *n* = 355 terminals from 38 fish.

Supplementary Figure 2

Figure S2: Diurnal changes in the visual signal delivered to the optic tectum

A. *Left panel*. Multiphoton section through the tectum of a zebrafish larva islet2b::mGCaMP6f (7 dpf) expressing the calcium reporter mGCaMP6f, which labels axons and synaptic terminals of retinal ganglion cells (RGCs). *Right panel*. Blow-up of the square red shown in the left panel. The image shows the Z plane reconstruction from where the synaptic responses were recorded. **B.** mGCaMP6f signals from individual ON and OFF RGCs synapses elicited using a stimulus contrasts of 20%,60% and 100 % modulated at 1 Hz (full field, sine wave) **C.** Average contrast-response functions displayed by ON and OFF RGCs, where the response (*R*) was quantified as the average of the fluorescence peak amplitudes measured at each cycle of stimulation. Each point shows the mean ± s.e.m. Note the differences in the magnitude of the responses between OFF RGCs in the morning relative to afternoon. **D.** mGCaMP6f signals from an individual OFF RGCs before and after intravitreal injection of the D1 antagonist in the afternoon. **E.** Average contrast-response function displayed by ON and OFF RGCs before and after administration of the D1 antagonist in the afternoon. Note the dramatic decrease in the magnitude of the response in both ON and OFF channels.

590   **Supplementary Figure 3**
591
592

593
Figure S3. Decomposition of iGluSnFR signals into vesicle counts
Summary of the basic steps for quantal decomposition of iGluSnFr signals. *1*. Raw trace extracted from
individual active zones (linescan, 1 KHz). *2*. Deconvolved trace using the estimated Wiener filter. *3*.
Histogram of event amplitudes for a representative active zone (373 events accumulated using stimulus
contrasts of 20%, 60% and 100% and a frequency of 5 Hz). The black line is a fit of eight Gaussians,
identified using a Gaussian mixture model. Note that the variance of successive Gaussians did not
increase in proportion to the peak number. The first peak had a value of 0.24, and the distance between
peaks averaged 0.25, indicating the existence of a quantal event equivalent to ~0.25. *.4*. Estimation of the
number of quanta per event. For more details about analyses see James et al., 2019.
603

Methods

Zebrafish husbandry

Fish were raised and maintained under standard conditions on a 14 h light/10 h dark cycle[35]. To aid imaging, fish were heterozygous or homozygous for the casper mutation which results in hypopigmentation and they were additionally treated with1-phenyl-2-thiourea (200 µM final concentration; Sigma) from 10 hours post fertilization (hpf) to reduce pigmentation. All animal procedures were performed in accordance with the Animal Act 1986 and the UK Home Office guidelines and with the approval of the University of Sussex Animal Welfare and Ethical Review Board. More information about experimental design and reagents is available in the Life Sciences reporting Summary.

Transgenic fish

Experiments were carried out using the following transgenic lines of zebrafish:

i) *Tg(ribeye:;Zf-SyGCaMP2)* expressing the synaptically-localized fluorescent calcium reporter SyGCaMP 2.0 in retinal bipolar cells under the ribeye-A promoter[24].

ii) Tg(−1.8ctbp2:Gal4VP16_BH) fish that drive the expression of the transcriptional activator protein Gal4VP16 were generated by co-injection of I-SceI meganuclease and endofree purified plasmid into wild-type zebrafish with a mixed genetic background. A myocardium-specific promoter that drives the expression of mCherry protein was additionally cloned into the plasmid to allow for phenotypical screening of founder fish.

iii) Tg(10xUAS:iGluSnFR_MH) fish driving the expression of the glutamate sensor iGluSnFR under the regulatory control of the 10 x UAS enhancer elements were generated by co-injection of purified plasmid and tol2 transposase RNA into offspring of AB wildtype fish outcrossed to casper wildtype fish. The sequences for the myocardium-specific promoter driving the expression of enhanced green fluorescent protein (mossy heart) were added to the plasmid to facilitate the screening process.

iv) *Tg(−1.8ctbp2:SyGCaMP6)* fish were generated by co-injection of I-SceI

meganuclease and endofree purified plasmid into wild-type zebrafish with a mixed

genetic background. The GCaMP6f variant was kindly provided by L. Looger (Janelia

Farm). This variant holds a T383S mutation in comparison to the commercially available

GCaMP6-fast version (Addgene plasmid 40755).

v) *Tg(isl2b:nlsTrpR, tUAS:memGCaMP6f)* which drives the expression of

memGCaMP6f in the optic tectum was generated by co-injecting pTol2-isl2b-hlsTrpR-pA

and pBH-tUAS-memGaMP6f-pA plasmids into single-cell stage eggs. Injected fish were

out-crossed with wild-type fish to screen for founders.

Multiphoton Imaging *In Vivo*

Experiments were carried out in a total of 117 zebrafish larvae (7–9 days post-

fertilization). Fish were immobilized in 3% low melting point agarose (Biogene) in E2

medium on a glass coverslip (0 thickness) and mounted in a chamber where they were

superfused with E2. Imaging was carried out using a two-photon microscope (Scientifica)

equipped with a mode-locked titanium-sapphire laser (Chameleon, Coherent) tuned to

915 nm and an Olympus XLUMPlanFI 20x water immersion objective (NA 0.95). To

prevent eye movements, the ocular muscles were paralyzed by injection of 1 nL of α-

bungarotoxin (2 mg/mL) behind the eye. Most imaging was carried out in the dorsal the

retina.

The signal-to-noise ratio of the microscope was optimized by collecting photons

through both the objective and a sub-stage oil condenser (Olympus, NA 1.4). Emission

was filtered through GFP filters (HQ 535/50, Chroma Technology) before detection with

GaAsP photomultipliers (H7422P-40, Hamamatsu). The signal from each detector passed

through a current-to-voltage converter and then the two signals were added by a

summing amplifier before digitization. Scanning and image acquisition were controlled

under ScanImage v.3.6 software[52]. In iGluSnFR recordings images were acquired at

10 Hz (128 × 100 pixels per frame, 1 ms per line) while linescans were acquired at 1 kHz.

In GCaMP recordings images were acquired at 20 Hz (128 × 50 pixels per frame, 1 ms

per line). Full-field light stimuli were generated by an amber LED (l_{max} = 590 nm,

Thorlabs), filtered through a 590/10 nm BP filter (Thorlabs), and delivered through a light

guide placed close to the eye of the fish. These wavelengths will most effectively

stimulate red and green cones. The microscope was synchronized to visual stimulation.

Stimulation protocols

Measurements of contrast sensitivity with SyGCaMP2 were made by stimulating the fish

with a series of 10 s stimuli (full-field sinusoidal modulation at 5 Hz) around a mean

intensity of 55 nW mm$^{-2.}$ Measurements of contrast sensitivity with iGluSnFR used 2 s

stimuli. To measure the distribution of events amplitudes and the temporal precision fish

were continuously stimulated for 30 s at a given contrast.

Luminance sensitivity was assessed by stimulating the fish with a series of light steps

(4 x 3 s) at 9 different light intensities increasing in steps of 0.5 log unit steps ranging from

11 pW mm^{-2} to 110 nW mm^{-2} (equivalent to 3.3 x 10^{11} photons mm^{-2}).

Drug injections

Dopamine signalling was manipulated by injecting the antagonist of D1 receptors SCH

23390 at a final estimated concentration of 200 nM (Sigma).Finally, the long-lasting

dopamine receptor ligand [3H] 2-amino-6,7-dihydroxy 1,2,3,4-tetrahydronapthalene

(ADTN) (Sigma) was injected to a final estimated concentration of 200 nM. We confirmed

that these drugs gained access by including 1 mM Alexa 594 in the injection needle;

within 5 mins of injection the dye could be detected within the inner plexiform layer of the

retina. Vehicle injection did not affect synaptic responses to varying contrast.

Calculation of temporal jitter

In order to quantify variability in the timing of glutamatergic events, we first calculated the

vector strength, r_q, for events composed of q quanta:

687
$$r_q = \frac{1}{N_q} \sqrt{\left(\sum_{i=1}^{N_q} cos\left(\frac{2\pi t_{q_i}}{T}\right)\right)^2 + \left(\sum_{i=1}^{N_q} sin\left(\frac{2\pi t_{q_i}}{T}\right)\right)^2}$$
(**1**)

688   where t_{qi} is the time of the i^{th} q-quantal event, T is the stimulus period, and N_q is the total

689   number of events of composed of q-quanta. The temporal jitter, J_q, can then be calculated

690   as:

691
$$J_q = \frac{\sqrt{2(1 - r_q)}}{2\pi f}$$
(**2**)

where f is the stimulus frequency.

Calculations based on Information Theory

To quantify the amount of information about a visual stimulus that is contained within

the sequence of release events from an active zone we first needed to convert bipolar

cell outputs into a probabilistic framework from which we could evaluate the specific

information (I_2), a metric that quantifies how much information about one random

variable is conveyed by the observation a specific symbol of another random

variable[36].   The time series of quantal events was converted into a probability

distribution by dividing into time bins of 20 ms, such that each bin contained either zero

events or one event of an integer amplitude. We then counted the number of bins

containing events of amplitude 1, or 2, or 3 etc. By dividing the number of bins of each

type by the total number of bins for each different stimulus, we obtained the conditional

distribution of **Q** given **S**, $p(Q|S)$, where **Q** is the random variable representing the

quanta/bin and **S** is the random variable representing the *stimulus contrasts* presented

throughout the course of the experiment. In the absence of information about the

distribution of contrasts normally experienced by a larval zebrafish, a uniform

distribution of contrasts was used for S. Each contrast step lasted 2 s (5 Hz) and they

were presented in two different pseudo-random orders, of which one is shown in Fig.

4D.   The contrast sensitivity varied between synapses and between morning and

afternoon (Fig. 1E-G) so to make allowance for this the stimulus set S was adjusted for

each synapse to span contrasts ±10% around $C_{1/2}$ measured within that synapse.

We computed the joint probability distribution by the chain rule for probability (given

the experimentally defined uniform distribution of stimuli **S**):

$$p(S, Q) = p(Q|S)p(S) \tag{7}$$

717   In order to convert this distribution into the conditional distribution of S given Q, we used

718   the definition of the conditional distribution:

$$p(S|Q) = \frac{p(S, Q)}{p(Q)} \tag{8}$$

720   From these distributions we computed two metrics: the mutual information I(**S;Q**)[53] and

721   specific information I_2(**S;q**)[37]. Mutual information is defined traditionally as:

$$I(S; Q) = H(S) - H(S|Q) \tag{9}$$

$$I(S; Q) = \sum_{s \in S} \sum_{q \in Q} p(s, q) \log_2 \frac{p(s)p(q)}{p(s, q)} = I(Q; S) \tag{10}$$

724   The specific information, I_2(**S;q**), is defined as the difference between the entropy of the

725   stimulus S minus the conditional entropy of the stimulus given the observed symbol in the

726   response q:

$$I_2(S, q) = H(S) - H(S|q) \tag{11}$$

$$I_2(S, q) = -\sum_{s \in S} p(s) \log p(s) + \sum_{s \in S} p(s|q) \log p(s|q) \tag{12}$$

representing the amount of information observing each quantal event type q ∈ **Q** carries

about the stimulus distribution **S**. Note that mutual information can also be computed from

the specific information as the dot product of the specific information vector I_2 and the

vector describing the probability of an event of a given quantal size $p(q)$. This adds to the

interpretability of both metrics – the specific information is the amount of information a

single (specific) symbol gives about the stimulus, and the mutual information is the

average amount of information about the stimulus gained from observing any symbol.

Measuring entropy and mutual information from neural responses can be a

challenging problem. Estimates require sampling from an unknown discrete probability

distribution, and in many cases recording sufficient samples to observe all non-zero

probability events is neither tractable nor practical. The biases introduced by

undersampling can be a particular problem when the full support of the distribution (all

values that map to non-zero probabilities) is high. Within the past few decades, various

approaches to correcting biases in information theoretic analyses have been developed[54].

However, as the distributions of interest in this work have both a small support and are

well sampled, we have opted to use standard estimates for the quantities of interest.

Statistics

All data are given as mean ± s.e.m. unless otherwise stated in the figure legends. All

statistical tests met appropriate assumptions and were calculated using inbuilt

functions in IgorPro (Wavemetrics). When data were not normally distributed we used

non-parametric tests. Significance was defined as $p < 0.05$. Data collection was not

randomized because all experiments were carried out within one set of animals.

Delivery of different stimuli was randomized where appropriate. Data were only

excluded from the analysis if the signal-to-noise ratio (SNR) of the iGluSnFR signals

elicited at a given synapse was not sufficient to detect unitary responses to visual stimuli

with a SNR of at least three.

757
758

References

1   Bargmann, C. I. & Marder, E. From the connectome to brain function. *Nat Methods* **10**, 483-490, doi:nmeth.2451 [pii]

10.1038/nmeth.2451 (2013).

Kastner, D. B. & Baccus, S. A. Coordinated dynamic encoding in the retina using opposing forms of plasticity. *Nature neuroscience* **14**, 1317 (2011).

Johnston, J. *et al.* A Retinal Circuit Generating a Dynamic Predictive Code for Oriented Features. *Neuron* **102**, 1211-1222 e1213, doi:10.1016/j.neuron.2019.04.002 (2019).

Esposti, F., Johnston, J., Rosa, J. M., Leung, K. M. & Lagnado, L. Olfactory stimulation selectively modulates the OFF pathway in the retina of zebrafish. *Neuron* **79**, 97-110, doi:10.1016/j.neuron.2013.05.001 (2013).

Schroder, S. *et al.* Arousal Modulates Retinal Output. *Neuron* **107**, 487-495 e489, doi:10.1016/j.neuron.2020.04.026 (2020).

Ribelayga, C., Cao, Y. & Mangel, S. C. The circadian clock in the retina controls rod-cone coupling. *Neuron* **59**, 790-801, doi:10.1016/j.neuron.2008.07.017 (2008).

Li, L. Circadian Vision in Zebrafish: From Molecule to Cell and from Neural Network to Behavior. *J Biol Rhythms* **34**, 451-462, doi:10.1177/0748730419863917 (2019).

Zang, J. *et al.* Circadian regulation of vertebrate cone photoreceptor function. *eLife* **10**, e68903, doi:10.7554/eLife.68903 (2021).

Jackson, C. R. *et al.* Retinal dopamine mediates multiple dimensions of light-adapted vision. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 9359-9368, doi:10.1523/JNEUROSCI.0711-12.2012 (2012).

Euler, T., Haverkamp, S., Schubert, T. & Baden, T. Retinal bipolar cells: elementary building blocks of vision. *Nat Rev Neurosci* **15**, 507-519, doi:10.1038/nrn3783 (2014).

Nikolaev, A., Leung, K. M., Odermatt, B. & Lagnado, L. Synaptic mechanisms of adaptation and sensitization in the retina. *Nat Neurosci* **16**, 934-941, doi:10.1038/nn.3408 (2013).

Zimmermann, M. J. Y. *et al.* Zebrafish Differentially Process Color across Visual Space to Match Natural Scenes. *Curr Biol* **28**, 2018-2032.e2015, doi:10.1016/j.cub.2018.04.075 (2018).

Matsumoto, A. *et al.* Direction selectivity in retinal bipolar cell axon terminals. *Neuron* (2021).

Lagnado, L. & Schmitz, F. Ribbon Synapses and Visual Processing in the Retina. *Annu Rev Vis Sci* **1**, 235-262, doi:10.1146/annurev-vision-082114-035709 (2015).

Glowatzki, E. & Fuchs, P. A. Transmitter release at the hair cell ribbon synapse. *Nat Neurosci* **5**, 147-154, doi:10.1038/nn796 (2002).

Singer, J. H., Lassova, L., Vardi, N. & Diamond, J. S. Coordinated multivesicular release at a mammalian ribbon synapse. *Nat Neurosci* **7**, 826-833, doi:10.1038/nn1280 (2004).

James, B., Darnet, L., Moya-Diaz, J., Seibel, S. H. & Lagnado, L. An amplitude code transmits information at a visual synapse. *Nat Neurosci* **22**, 1140-1147, doi:10.1038/s41593-019-0403-6 (2019).

Lisman, J. E., Raghavachari, S. & Tsien, R. W. The sequence of events that underlie quantal transmission at central glutamatergic synapses. *Nat Rev Neurosci* **8**, 597-609, doi:10.1038/nrn2191 (2007).

Huang, C.-H., Bao, J. & Sakaba, T. Multivesicular Release Differentiates the Reliability of Synaptic Transmission between the Visual Cortex and the Somatosensory Cortex. *The Journal of Neuroscience* **30**, 11994, doi:10.1523/JNEUROSCI.2381-10.2010 (2010).

20 Holler, S., Köstinger, G., Martin, K. A. C., Schuhknecht, G. F. P. & Stratford, K. J. Structure
and function of a neocortical synapse. *Nature* **591**, 111-116, doi:10.1038/s41586-020-03134-2
(2021).

21 Shannon, C. A Mathematical Theory of Communication. *Bell System Technical Journal* **27**,
379-423 (1948).

22 Borst, A. & Theunissen, F. E. Information theory and neural coding. *Nat Neurosci* **2**, 947-957,
doi:10.1038/14731 (1999).

23 Marvin, J. S. *et al.* An optimized fluorescent probe for visualizing glutamate
neurotransmission. *Nat Methods* **10**, 162-170, doi:10.1038/nmeth.2333 (2013).

24 Dreosti, E., Odermatt, B., Dorostkar, M. M. & Lagnado, L. A genetically encoded reporter of
synaptic activity in vivo. *Nat Methods* **6**, 883-889, doi:10.1038/nmeth.1399 (2009).

25 Emran, F., Rihel, J., Adolph, A. R. & Dowling, J. E. Zebrafish larvae lose vision at night.
Proceedings of the National Academy of Sciences **107**, 6034, doi:10.1073/pnas.0914718107
(2010).

26 Nie, K. *et al.* Effects of circadian clock protein Per1b on zebrafish visual functions.
Chronobiology international **35**, 160-168 (2018).

27 Ashmore, J. & Copenhagen, D. An analysis of transmission from cones to hyperpolarizing
bipolar cells in the retina of the turtle. *The Journal of Physiology* **340**, 569-597 (1983).

28 Faisal, A. A., Selen, L. P. & Wolpert, D. M. Noise in the nervous system. *Nat Rev Neurosci* **9**,
292-303, doi:10.1038/nrn2258 (2008).

29 Rusakov, D. A., Savtchenko, L. P. & Latham, P. E. Noisy Synaptic Conductance: Bug or a
Feature? *Trends Neurosci* **43**, 363-372, doi:10.1016/j.tins.2020.03.009 (2020).

30 Berry, M. J., Warland, D. K. & Meister, M. The structure and precision of retinal spike trains.
Proceedings of the National Academy of Sciences **94**, 5411, doi:10.1073/pnas.94.10.5411 (1997).

31 Churchland, M. M. *et al.* Stimulus onset quenches neural variability: a widespread cortical
phenomenon. *Nature Neuroscience* **13**, 369-378, doi:10.1038/nn.2501 (2010).

32 Rathbun, D. L., Warland, D. K. & Usrey, W. M. Spike timing and information transmission at
retinogeniculate synapses. *Journal of Neuroscience* **30**, 13558-13566 (2010).

33 Cui, Y., Wang, Y. V., Park, S. J. H., Demb, J. B. & Butts, D. A. Divisive suppression explains
high-precision firing and contrast adaptation in retinal ganglion cells. *eLife* **5**, e19460,
doi:10.7554/eLife.19460 (2016).

34 Freed, M. A. Quantal encoding of information in a retinal ganglion cell. *J Neurophysiol* **94**,
1048-1056, doi:10.1152/jn.01276.2004 (2005).

35 Odermatt, B., Nikolaev, A. & Lagnado, L. Encoding of luminance and contrast by linear and
nonlinear synapses in the retina. *Neuron* **73**, 758-773, doi:10.1016/j.neuron.2011.12.023 (2012).

36 Stone, J. V. *Principles of Neural Information Theory: Computational Neuroscience and*
Metabolic Efficiency (Sebtel Press, 2018).

37 DeWeese, M. R. & Meister, M. How to measure the information gained from one symbol.
Network **10**, 325-340 (1999).

38 Attwell, D. & Laughlin, S. B. An energy budget for signaling in the grey matter of the brain.
Journal of Cerebral Blood Flow & Metabolism **21**, 1133-1145 (2001).

39 Harris, Julia J., Jolivet, R. & Attwell, D. Synaptic Energy Use and Supply. *Neuron* **75**, 762-
777, doi:https://doi.org/10.1016/j.neuron.2012.08.019 (2012).

40 Koch, K. *et al.* How much the eye tells the brain. *Curr Biol* **16**, 1428-1434,
doi:10.1016/j.cub.2006.05.056 (2006).

41 Mora‐Ferrer, C., Yazulla, S., Studholme, K. M. & Haak‐Frendscho, M. Dopamine D1‐
receptor immunolocalization in goldfish retina. *Journal of Comparative Neurology* **411**, 705-714
(1999).

42 Farshi, P., Fyk-Kolodziej, B., Krolewski, D. M., Walker, P. D. & Ichinose, T. Dopamine D1
receptor expression is bipolar cell type-specific in the mouse retina. *The Journal of comparative*
neurology **524**, 2059-2079, doi:10.1002/cne.23932 (2016).

43 Ribelayga, C., Wang, Y. & Mangel, S. C. A circadian clock in the fish retina regulates
dopamine release via activation of melatonin receptors. *The Journal of Physiology* **554**, 467-482,
doi:https://doi.org/10.1113/jphysiol.2003.053710 (2004).

44 Moya-Díaz, J., James, B. & Lagnado, L. Modulation of the vesicle code transmitting the
visual signal in the retina. *bioRxiv*, 2020.2004.2022.056119, doi:10.1101/2020.04.22.056119
(2020).

45 Thermos, K. Functional mapping of somatostatin receptors in the retina: a review. *Vision*
Research **43**, 1805-1815, doi:https://doi.org/10.1016/S0042-6989(03)00169-X (2003).

46 Freed, M. A. & Liang, Z. Synaptic noise is an information bottleneck in the inner retina during
dynamic visual stimulation. *The Journal of physiology* **592**, 635-651,
doi:10.1113/jphysiol.2013.265744 (2014).

47 McDonnell, M. D. & Ward, L. M. The benefits of noise in neural systems: bridging theory and
experiment. *Nature Reviews Neuroscience* **12**, 415-425, doi:10.1038/nrn3061 (2011).

48 Christie, J. M. & Jahr, C. E. Multivesicular release at Schaffer collateral-CA1 hippocampal
synapses. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **26**,
210-216, doi:10.1523/JNEUROSCI.4307-05.2006 (2006).

49 Auger, C., Kondo, S. & Marty, A. Multivesicular release at single functional synaptic sites in
cerebellar stellate and basket cells. *The Journal of neuroscience : the official journal of the*
Society for Neuroscience **18**, 4532-4547 (1998).

50 Higley, M. J., Soler-Llavina, G. J. & Sabatini, B. L. Cholinergic modulation of multivesicular
release regulates striatal synaptic potency and integration. *Nature neuroscience* **12**, 1121-1128
(2009).

51 Chalifoux, J. R. & Carter, A. G. GABAB receptors modulate NMDA receptor calcium signals
in dendritic spines. *Neuron* **66**, 101-113 (2010).

52 Pologruto, T. A., Sabatini, B. L. & Svoboda, K. ScanImage: flexible software for operating
laser scanning microscopes. *Biomed Eng Online* **2**, 13, doi:10.1186/1475-925X-2-13 (2003).

53 de Ruyter van Steveninck, R. R., Lewen, G. D., Strong, S. P., Koberle, R. & Bialek, W.
Reproducibility and variability in neural spike trains. *Science* **275**, 1805-1808,
doi:10.1126/science.275.5307.1805 (1997).

54 Pola, G., Schultz, S. R, Petersen, R. S & Panzeri, S. in *Neuroscience Databases: A Practical*
Guide (Springer, Boston, MA, 2003).