Amoxicillin-resistant *Streptococcus pneumoniae* can be resensitized by targeting

the mevalonate pathway as indicated by sCRilecs-seq

Liselot Dewachter[a], Xue Liu[a,b], Julien Dénéréaz[a], Vincent de Bakker[a], Charlotte Costa[c], Mara Baldry[c],

Jean-Claude Sirard[c], Jan-Willem Veening[a]*

[a] Department of Fundamental Microbiology, Faculty of Biology and Medicine, University of Lausanne,

Biophore Building, Lausanne 1015, Switzerland

[b] Guangdong Key Laboratory for Genome Stability and Human Disease Prevention, Department of

Pharmacology, International Cancer Center, Shenzhen University Health Science Center, 518060

Shenzhen, China

[c] Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur Lille, U1019 - UMR 9017 - CIIL - Center for Infection

and Immunity of Lille, F-59000 Lille, France

*Correspondence to: jan-willem.veening@unil.ch Twitter: @JWVeening

Abstract

Antibiotic resistance in the important opportunistic human pathogen *Streptococcus pneumoniae* is on the rise. This is particularly problematic in the case of the β-lactam antibiotic amoxicillin, which is the first-line therapy. It is therefore crucial to uncover targets that would kill or resensitize amoxicillin-resistant pneumococci. To do so, we developed a genome-wide, single-cell based, gene silencing screen using CRISPR interference called sCRilecs-seq (subsets of CRISPR interference libraries extracted by fluorescence activated cell sorting coupled to next generation sequencing). Since amoxicillin affects growth and division, sCRilecs-seq was used to identify targets that are responsible for maintaining proper cell size. Our screen revealed that downregulation of the mevalonate pathway leads to extensive cell elongation. Further investigation into this phenotype indicates that it is caused by insufficient transport of cell wall precursors across the cell membrane due to a limitation in the production of undecaprenyl phosphate (Und-P), the lipid carrier responsible for this process. The data suggest that whereas peptidoglycan synthesis continues even with reduced Und-P levels, cell constriction is specifically halted. We successfully exploited this knowledge to create a combination treatment strategy where the FDA-approved drug clomiphene, an inhibitor of Und-P synthesis, is paired up with amoxicillin. Our results show that clomiphene potentiates the antimicrobial activity of amoxicillin and that combination therapy resensitizes amoxicillin-resistant *S. pneumoniae*. These findings could provide a starting point to develop a solution for the increasing amount of hard-to-treat amoxicillin-resistant pneumococcal infections.

Keywords

sCRilecs-seq, mevalonate pathway, morphology, peptidoglycan synthesis, cell division, antibiotic resistance, amoxicillin, clomiphene, combination therapy

Introduction

Streptococcus pneumoniae (the pneumococcus) is an opportunistic human pathogen that is responsible for diseases such as pneumonia, middle ear infections, sepsis and meningitis[1-3]. *S. pneumoniae* is listed by the WHO as a priority pathogen for the development of novel antibiotics, since it is one of the leading causes of fatal bacterial infections worldwide[1,2,4] and antibiotic resistance is on the rise[5-7]. The alarming increase in prevalence of penicillin non-susceptible *S. pneumoniae* was initially dealt with by the development and roll-out of the pneumococcal conjugate vaccines PCV7, PCV10 and PCV13 in 2001, 2009 and 2010, respectively[5]. These vaccines induce protection against infection caused by 7, 10 or 13 of the most prevalent serotypes of capsular polysaccharides that surround *S. pneumoniae*[3]. Since many of the targeted serotypes are associated with penicillin non-susceptibility, the occurrence of infections with such resistant strains decreased at first[5,8,9]. However, due to the high genomic plasticity of *S. pneumoniae*[10,11], serotype switching created penicillin non-susceptible escape mutants that are not covered by the currently available vaccines and that are once again gaining in prevalence[5,8].

Penicillin non-susceptible *S. pneumoniae* strains carry a variety of mutations in their penicillin binding proteins (PBPs) that decrease their affinity for penicillin and therefore increase resistance[4,7,12]. These mutations do not only lower the susceptibility to penicillin but also to other β-lactam antibiotics[13]. However, the efficacy of aminopenicillins such as amoxicillin is usually less affected by these mutations, meaning that amoxicillin remains a viable treatment option for many penicillin non-susceptible strains and therefore became one of the frontline antibiotics to treat pneumococcal infections[13-17]. However, highly penicillin resistant strains have been found to also be amoxicillin non-susceptible[5,6], thereby now also threatening the clinical efficacy of this widely used antibiotic.

Because of the rise in antibiotic resistance, we urgently need new antibiotic targets and/or need to find ways to extend the clinical efficacy of our current antibiotic arsenal. To contribute towards this goal, we developed sCRilecs-seq (subsets of CRISPR interference libraries extracted by fluorescence activated cell sorting coupled to next generation sequencing), a high-throughput single-cell based screening approach that we exploited to find targets that could resensitize resistant *S. pneumoniae* strains towards amoxicillin. sCRilecs-seq is based upon genome-wide gene silencing using CRISPR interference (CRISPRi), which makes use of a catalytically dead form of the Cas9 enzyme, called dCas9[18]. dCas9 is guided towards its target site in the DNA by the provided sgRNA but is unable to introduce double-stranded breaks. Instead, the dCas9 enzyme acts as a roadblock for RNA polymerase, thereby halting transcription[18-20]. CRISPRi gene silencing

not only affects the target gene but influences the expression of an entire transcriptional unit[18-22]. This

technology therefore works at the operon level and generates strong polar effects that need to be

considered. Despite this limitation, CRISPRi has proven to be a very powerful genetic tool to perform

genome-wide depletion screens in various bacteria[22-27]. In the sCRilecs-seq screen used here,

subpopulations that display a phenotype of interest are collected using Fluorescence Activated Cell

Sorting (FACS), as has previously been done in eukaryotic cells[28] and for bacterial transposon libraries[29].

The abundance of sgRNAs in the sorted fractions are then compared to a defined control population to

look for genetic factors involved in the phenotype of interest. This highly flexible screening approach

allowed us to identify targets that can be exploited in our fight against bacterial infections.

Our sCRilecs-seq results highlight the importance of the mevalonate pathway for maintaining proper cell

morphology in *S. pneumoniae*. The mevalonate pathway is the only pathway used by *S. pneumoniae* to

produce isoprenoids, a highly diverse class of organic molecules that are essential to all life on earth and

that function – among others – in processes such as cell wall synthesis, electron transport, membrane

stability and protein modification[30,31]. Our screen revealed that inhibition of the mevalonate pathway in

S. pneumoniae leads to extensive cell elongation. The data suggests that this elongation is caused by

insufficient transport of cell wall precursors across the cell membrane due to a limitation in the production

of undecaprenyl phosphate (Und-P), the lipid carrier responsible for this process[32]. This shortage of

peptidoglycan precursors allows cell elongation to continue but causes a block in cell division.

Additionally, based on the mevalonate depletion phenotype characterized here, we successfully designed

a combination treatment strategy targeting *S. pneumoniae*. In this combination treatment strategy,

amoxicillin is potentiated by the simultaneous administration of clomiphene, a widely used FDA-approved

fertility drug that was shown to block Und-P production in *Staphylococcus aureus* and to synergize with

β-lactam antibiotics[33]. This combination of compounds is particularly powerful against amoxicillin-

resistant *S. pneumoniae* strains, as it is capable of resensitizing these strains to clinically-relevant

concentrations of amoxicillin *in vitro*. Our findings could therefore provide a useful starting point for the

development of treatment strategies that are effective against the rising amount of amoxicillin-resistant

S. pneumoniae infections and could extend the clinical efficacy of this important antibiotic.

| 102 | **Results** |

| 103 | **Amoxicillin treated *S. pneumoniae* are elongated** |

104	To identify new druggable pathways that could potentiate the antimicrobial activity of amoxicillin, we first
105	established the impact of amoxicillin on pneumococcal morphology and growth. We therefore performed
106	time-lapse microscopy of *S. pneumoniae* in the presence or absence of amoxicillin. As shown in Figure 1A-
107	B and Movie S1-2, amoxicillin-treated cells initially elongate before they lyse. Indeed, quantitative image
108	analysis confirmed that amoxicillin, like some other β-lactam antibiotics[34], causes cell elongation (Figure
109	1C), while growth curves in the presence of this antibiotic demonstrate that lysis occurs since OD values
110	decrease at high amoxicillin concentrations (Figure 1D). To demonstrate that cell elongation is specifically
111	caused by amoxicillin and does not generally occur upon antibiotic treatment, we also assessed cell
112	lengths in the presence of the fluoroquinolone ciprofloxacin. As expected, ciprofloxacin did not induce
113	cell elongation[35] (Figure 1C). Because it was shown that synergy between antimicrobial compounds is
114	most often detected when they target the same process[36], we hypothesized that targets whose inhibition
115	would cause cell elongation, like amoxicillin, could potentially reinforce the antibacterial effect of this
116	antibiotic.

Figure 1: Amoxicillin causes cell elongation before triggering cell lysis. A) Snapshots of a time lapse phase contrast microscopy
experiment of *S. pneumoniae* D39V growing in the absence of antibiotics (Movie S1). B) Snapshots of a time lapse analysis of *S.*
pneumoniae D39V growing in the presence of a sub-MIC concentration of amoxicillin (0.016 μg/ml) (Movie S2). C) The effect of
sub-MIC concentrations of ciprofloxacin (0.5 μg/ml) and amoxicillin (0.016 μg/ml) on the cell length of *S. pneumoniae* D39V was
tested by phase contrast microscopy after 2h of treatment. Quantitative analysis of micrographs show that cell length increases
upon treatment with amoxicillin. Data are represented as violin plots with the mean cell length of every biological repeat indicated
with orange dots. The size of these dots indicates the number of cells recorded in each repeat, ranging from 112 to 1498 cells.
Black dots represent the mean ± SEM of all recorded means, n ≥ 3. Two-sided Wilcoxon signed rank tests were performed against
wt without antibiotic as control group (dotted line) and p values were adjusted with an FDR correction; ** $p < 0.01$. D) *S.*
pneumoniae D39V was grown in the presence of different concentrations of amoxicillin and growth was followed by monitoring
OD 595 nm. Wt, wildtype; Cip, ciprofloxacin; Amx, amoxicillin.

Development of sCRilecs-seq

To identify pathways that lead to cell elongation upon inhibition, we developed a single-cell based, high-

throughput screening strategy where we combined CRISPRi gene silencing with high-throughput selection

of a phenotype of interest by Fluorescence Activated Cell Sorting (FACS). Using this approach, which we

call sCRilecs-seq, we screened for mutants that display an increase in forward scatter (FSC), which is a

rough indicator of cell size[37]. To facilitate downstream analyses, screening was performed in a *S.*

pneumoniae strain in which the DNA and the cell division protein FtsZ are marked. DNA was visualized

using the DNA-binding protein HlpA fused to GFP. HlpA, also known as HU, is a histone-like protein that

binds DNA aspecifically and is highly expressed in *S. pneumoniae*[38]. It is therefore often used as a nucleoid

marker[38,39]. FtsZ was marked by fusing it to mCherry. To this end, the native *hlpA* and *ftsZ* genes of *S.*

pneumoniae D39V were replaced by *hlpA-gfp* and *ftsZ-mCherry*, respectively. Both of these fusions were

previously shown to support viability[38,39]. Additionally, the gene encoding dCas9 was inserted into the

pneumococcal chromosome and was placed under tight control of an IPTG-inducible P*lac* promoter[21,23].

The resulting strain (VL3117: D39V *lacI* P*lac*-*dcas9 hlpA-gfp ftsZ-mCherry*) was transformed with a pool of

1499 different integrative plasmids carrying constitutively-expressed sgRNAs (under control of the P3

promoter) that together target the entire coding genome[23]. This way, we created a pooled CRISPRi library

where each cell expresses a certain sgRNA resulting in transcriptional downregulation of a specific gene

or operon upon induction of dCas9 (Figure 2A). Note that sgRNAs were designed in such a way that the

chance for off targeting effects are minimal[23].

We next grew the library in C+Y medium at 37°C for 3.5h in the presence of 1 mM IPTG to induce dCas9.

The pooled and induced library was subjected to FACS and different fractions of the population were

sorted (see Materials and Methods). Of note, *S. pneumoniae* grows as short cell chains[40] and the number

of cells present in a chain could influence the FSC read-out. To ensure that measurements were taken at

the single-cell level, chains were mechanically disrupted by vigorous vortexing before sorting took place.

We confirmed that this approach is successful at breaking up cell chains ensuring that morphology was

evaluated for single cells rather than entire chains (Figure S1A-C). Ten percent of the population with the

highest FSC values was collected. As a control, the centermost 70% of the population was sorted as well.

For both collected fractions, sgRNAs were amplified by PCR using primers that contain Illumina adapters

(see Materials and Methods). After sequencing, sgRNA read counts were compared between the sorted

fractions. This approach (Figure 2A) enabled us to identify sgRNAs that are significantly enriched in the

fraction of the population with highest FSC values. These sgRNAs point to genes or operons necessary to

maintain normal cell size.

Figure 2: sCRilecs-seq identifies operons involved in cell size regulation. A) A pooled CRISPRi library was constructed in *S.*

pneumoniae D39V *P_lac-dcas9 lacI hlpA-gfp ftsZ-mCherry* (VL3117) by transformation of a plasmid library encoding 1499

constitutively-expressed sgRNAs that together target the entire genome. This CRISPRi library was grown in the presence of IPTG
for 3.5h to induce expression of dCas9, and cultures were sorted based on forward scatter (FSC) as a proxy for cell size. 10% of
the population with the highest FSC values was sorted, as well as the centermost 70% of the population which served as a control.
sgRNAs from sorted fractions were amplified by PCR using primers that contain Illumina adapters. Amplified sgRNAs were
sequenced and mapped to the *S. pneumoniae* genome. sgRNA read counts were compared between the different sorted fractions
to identify gene depletions that lead to increases in cell size. B) A volcano plot shows the statistical significance and enrichment
for every sgRNA in the fraction of the population with high FSC values compared to the control with normal FSC values. C)
Significant sgRNA hits were validated by studying individual mutants. Flow cytometry measurements of mutants grown with and
without IPTG were compared. Note that the entire CRISPRi library was also included in this experiment ('Library') as a control.
Data are represented as mean ± SEM, n = 3. ** p < 0.01, *** p < 0.001, **** p < 0.0001, two-sided t-tests with Holm-Sidak
correction for multiple comparisons.
179

sCRilecs-seq identifies several targets involved in maintaining proper cell size
Following this strategy, we were able to identify 17 sgRNAs that were significantly enriched in the fraction
of the population with high FSC values (Figure 2B, Figure S1D-E and Table S1), indicating that depletion of
these sgRNA targets leads to an increase in cell size. The relatively low amount of significant hits isolated
here, and the absence of a couple of known cell cycle regulators, such as GpsB[41,42], indicates that false
negative results are probably quite common. This is perhaps not surprising given the high amount of
variation detected between different repeats (Figure S1D-E). We also note that some knock down mutants
that are highly sensitive to physical perturbations might be lost due to the vigorously vortexing of cells
prior to FACS. In a next step, we validated whether the 17 significant hits from our sCRilecs-seq screen
were correctly identified. The corresponding sgRNAs were individually cloned and transformed to strain
D39V P*lac*-*dcas9 lacI hlpA-gfp ftsZ-mCherry* (VL3117) and their effect was evaluated by flow cytometry. The
median FSC value for each depletion was recorded and compared to the median value of the same strain
without induction of dCas9. As shown in Figure 2C, all of the significant FSC hits from the sCRilecs-seq
screen demonstrated increased FSC levels upon depletion, meaning that no false positives were identified
by the cell sorting approach used in our sCRilecs-seq screen and that all selected sgRNAs indeed affect FSC
values and therefore likely change cell size.
196

The reliability of our screening approach is also reflected by the fact that many of the significant hits are
known to be involved in processes that influence cell size and/or morphology (Table S1). For example, 3
out of 17 hits are involved in cell division (i.e. *divIB*, *ftsL*, *ylmH*), 1 hit interferes with peptidoglycan
synthesis (*murF*) and at least 3 hits target the production of teichoic acids (*licD1*, *tacF*, *tacL*, and the operon
containing *tarQ*, *tarP* and *licD3*). These glycopolymers are an important component of the *S. pneumoniae*

cell envelope and are necessary to maintain proper cell morphology[21,43,44]. The fact that these hits are

picked up by our screen demonstrates that our strategy is capable of extracting biologically relevant

information.

Interestingly, when performing Gene Ontology (GO) enrichment[45-47] to look for biological processes that

are found among the hits more often than expected based on random chance, one biological process, i.e.

biosynthesis of isoprenoids, was found to be significantly overrepresented (Fold Enrichment = 56, P value

= 0.0019). We therefore decided to further focus on isoprenoid biosynthesis and how disturbances in this

process lead to changes in *S. pneumoniae* cell size that might be exploited as a novel antimicrobial

strategy.

Depletion of mevalonate pathway components leads to cell elongation

S. pneumoniae synthesizes isoprenoids through the mevalonate pathway[30,48]. This pathway, with

mevalonic acid (mevalonate) as intermediate, produces the C5 precursors, isopentenyl diphosphate and

dimethylallyl diphosphate (Figure 3A), which can be condensed into large and diverse isoprenoids[30]. In *S.*

pneumoniae, all proteins involved in the mevalonate pathway are encoded by two different operons[48,49],

both of which were identified as hits in our sCRilecs-seq screen. In a first step, we made non-CRISPRi-

based deletion/complementation strains for both of these operons. In an *S. pneumoniae* D39V strain that

encodes the *hlpA-gfp* and *ftsZ-mCherry* fusion proteins as well as LacI (VL3404), complementation

constructs for the mevalonate genes under control of the IPTG-inducible P_{lac} promoter were inserted

ectopically into the genome and the native mevalonate genes were deleted (Figure 3B). The resulting two

deletion/complementation strains (VL3565 and VL3567) were then used to assess the effect of

mevalonate synthesis on cell size and growth.

Phase contrast microscopy on cells depleted for the mevalonate pathway operons demonstrated large

increases in cell length (Figure 3C-D and S2A), confirming the sCRilecs-seq screen. Even though cells also

became slightly wider upon mevalonate depletion (Figure 3C), cell length was disproportionally affected.

Cells on average attained more than twice their normal length when the mevalonate operons were

depleted (Figure 3D and S2A), whereas the mean cell width only increased 1.35x for depletion of operon

1 and 1.25x for operon 2. The disproportionately large effect on cell elongation is also demonstrated by

significant increases in the length/width ratio (Figure S2A). In addition to cell elongation, depletion of the

mevalonate operons led to severe growth defects (Figure 3E). Importantly, the elongated phenotype and

observed growth defects could be complemented by ectopic induction of the mevalonate operons with

an optimized concentration of IPTG (Figure 3F-G), demonstrating that these phenotypes are not due to

polar effects of the mutations. Time lapse microscopy upon depletion of the mevalonate operons

confirmed these findings, showing that cells initially continue to grow and elongate before lysis takes place

(Figure 3J and Movie S2-3). Of note, these phenotypes were reproducible in deletion/complementation

strains that do not carry the *hlpA-gfp* and *ftsZ-mCherry* fusions (VL3708 and VL3709) (Figure S2B-E).

Although the severe growth defects associated with depletion of the genes involved in mevalonate

biosynthesis point to the importance of this pathway for *S. pneumoniae* physiology, cultures were not

fully sterilized at late time points and many survivors remained (Figure 3E). To test whether the

mevalonate pathway is essential for *S. pneumoniae* viability and that the survival of the depletion strains

at late time points is due to leaky expression and/or suppressor mutations, we attempted to delete the

mevalonate operons without providing a complementation construct. As expected for essential genes, we

were unable to delete these operons. However, when mevalonic acid – an intermediary in the mevalonate

pathway (Figure 3A) – was added to the growth medium, we were able to delete operon 1, as was also

shown previously in *S. pneumoniae* and *S. aureus*[48,50-52]. As expected based on pathway architecture,

deleting operon 2 – which encodes proteins that work downstream of mevalonic acid – remained

impossible. We next grew *S. pneumoniae* Δ*operon 1* (VL3702) with different concentrations of mevalonic

acid. Cells grew normally and attained normal cell lengths when a high concentration of mevalonic acid

was added to the growth medium (Figure 3H-I). However, decreasing the concentration of mevalonic acid

led to cell elongation and growth defects that became more pronounced at lower concentrations, thereby

phenocopying genetic depletion of operon 1 and 2. In fact, when no mevalonic acid is provided at all,

cultures are driven to full extinction (Figure 3I). These results confirm that the mevalonate pathway is

essential for *S. pneumoniae* viability and that it is the only pathway through which this bacterium can

synthesize isoprenoids[48].

Figure 3: The mevalonate pathway is essential for *S. pneumoniae* and leads to cell elongation upon depletion. A) The mevalonate pathway and its genetic organization in *S. pneumoniae* is depicted. B) A genetic representation of the mevalonate depletion strains is shown. The native mevalonate operons were deleted and replaced by an antibiotic marker (AB) and a complementation construct under control of the P*lac* promoter was inserted at the *zip* locus[53] in the *S. pneumoniae* genome. C) Phase contrast microscopy images of *S. pneumoniae* wt or upon depletion of one of the mevalonate operons for 4h in VL3565 and VL3567 are shown. Scale bar, 5 μm. D) Quantitative analysis of phase contrast micrographs shows that cell length increased when mevalonate operons were depleted. Data are represented as violin plots with the mean cell length of every biological

repeat indicated with orange dots. The size of these dots indicates the number of cells recorded in each repeat, ranging from 100
to 2626 cells. Black dots represent the mean ± SEM of all recorded means, n ≥ 3.

E) Depletion of mevalonate operons led to a severe growth defect. Data are represented as the mean, n ≥ 3. F) The elongated
phenotype upon depletion of mevalonate operons could be complemented by inducing their expression with IPTG. Data are
represented as violin plots with the mean cell length of every biological repeat indicated with orange dots. The size of these dots
indicates the number of cells recorded in each repeat, ranging from 100 to 2626 cells. Black dots represent the mean ± SEM of all
recorded means, n ≥ 3. G) The growth defect associated with depletion of mevalonate operons could be fully complemented by
inducing their expression with IPTG. Data are represented as the mean, n ≥ 3. H) A mutant in which the first mevalonate operon
was deleted (VL3702, no complementation construct) displayed increased cell length when the concentration of mevalonic acid
added to the growth medium was decreased. Data are represented as violin plots with the mean cell length of every biological
repeat indicated with orange (or grey) dots. The size of these dots indicates the number of cells recorded in each repeat, ranging
from 100 to 2626 cells. Black dots represent the mean ± SEM of all recorded means, n ≥ 3. I) Growing the mutant in which the
first mevalonate operon is deleted (no complementation construct) with decreasing concentrations of external mevalonic acid
led to an increasing growth defect, resulting in full extinction of the culture when no mevalonic acid was provided. Data are
represented as the mean, n ≥ 3. J) Snapshot images of phase contrast time lapse experiments with mutants in which the first or
second mevalonate operon was depleted (VL3565 and VL3567) are shown. Strains were grown on agarose pads of C+Y medium
without the inducer IPTG. Scale bar, 5 µm. Two-sided Wilcoxon signed rank tests were performed against wt - IPTG as control
group (dotted line) and p values were adjusted with an FDR correction; * p < 0.05, *** p < 0.001.

Depletion of mevalonate operons primarily interferes with cell division

We next asked how depletion of mevalonate pathway components results in an elongated phenotype.
Microscopic investigation of mevalonate depleted cells showed that these bacteria contain a large
number of unconstricted FtsZ rings (Figure 4A-B). High-resolution investigation of this phenotype by
Transmission Electron Microscopy (TEM) showed that depletion of the mevalonate operons resulted in
an unusually high occurrence of initiated septa that appear to be blocked for further constriction, whereas
division septa at all stages of constriction could be found in a wild-type strain (Figure 4C). The inability to
constrict and divide could explain the observed filamentation if cell elongation still occurs.

To investigate if indeed peptidoglycan synthesis for cell elongation remains active upon depletion of the
mevalonate operons, we labeled sites of active peptidoglycan synthesis with Fluorescent D-Amino Acids
(FDAAs)[54]. In a first pulse, cells were labeled with the green FDAA, sBADA, for 15 min and a second 15 min
pulse consisted of the red FDAA, RADA (see Materials and Methods). Fluorescence microscopy showed
that peptidoglycan synthesis leads to both cell elongation and division in a wt strain (Figure 4D). Likewise,
also the mevalonate depletion strains showed sites of active peptidoglycan synthesis, indicating that not
all cell wall production is halted. However, in this case, we almost exclusively detected peptidoglycan

synthesis directed at cell elongation since almost no constricted FDAA-labeled sites were found (Figure

4D-E).. Additionally, it is clear that even though peptidoglycan synthesis continues upon mevalonate

depletion, this process is slowed down since FDAA intensity is decreased under these conditions,

indicating that less peptidoglycan is being produced (Figure 4D). Quantification of the intensity of the

sBADA signal indeed confirmed that the incorporation of the label is strongly decreased upon mevalonate

depletion (Figure 4F). The same quantification was not done for the RADA label since the relatively high

aspecific background fluorescence interfered strongly with this analysis.

Taken together, these results strongly indicate that mevalonate depletion leads to filamentation because

cell division is blocked while peptidoglycan synthesis for cell elongation can still occur, albeit at a lower

rate. However, elongation does not continue indefinitely and cells eventually lyse (Figure 3J and Movie

S3-4). Since lysis usually occurs through weakening of the peptidoglycan cell wall[56], we hypothesize that

extended depletion of mevalonate operons eventually affects the integrity and/or rigidity of the

peptidoglycan mesh, leading to cell lysis. However, considering the initial phenotypes found, we conclude

that the primary effect of mevalonate depletion is a block in constriction necessary for cell division.

Figure 4: Depletion of mevalonate operons prevents cell division. A) While *S. pneumoniae* wt cells typically contain one FtsZ ring
at the cell center, depletion of either one of the mevalonate operons led to strongly elongated cells with multiple unconstricted
FtsZ rings. Images were obtained using strains VL3404, VL3565 and VL3567 that encode the *ftsZ-mCherry* fusion. B) Quantitative
analysis of microscopy pictures was used to determine the number of Z-rings per cell. Pictures from at least 6 independent repeats
were analyzed, each including over 100 cells. C) Transmission Electron Microscopy (TEM) images show that elongated cells

contained many initiated septa that appear to be blocked in further progression of constriction (white arrow heads). Images were

obtained using strains VL333, VL3708 and VL3709 that do not encode fluorescent fusion proteins. D) Pulse labeling *S. pneumoniae*

with the green FDAA, sBADA, and 15 min later with the red RADA shows sites of active peptidoglycan synthesis involved in either

elongation or constriction. Images were obtained using strains VL333, VL3708 and VL3709 that do not encode fluorescent fusion

proteins. E-F) Quantitative image analysis of sites of active peptidoglycan synthesis labeled with FDAAs shows that depletion of

mevalonate operons eliminated septal peptidoglycan synthesis since virtually no narrow sBADA or RADA bands can be found (E)

and that the intensity of FDAA labeling decreased upon mevalonate depletion, indicating that peripheral peptidoglycan synthesis

was slowed down (F). Number of sBADA bands analyzed for each condition > 400.

Depletion of mevalonate operons prevents cell division by limiting the export of peptidoglycan

precursors

The mevalonate pathway provides the precursor for the production of all isoprenoids in *S.*

pneumoniae[30,48]. Thus, a shortage in the production of any of these isoprenoids could underlie the

phenotypic effects of perturbed mevalonate production. However, given the nature of the observed

phenotype, we thought it most likely that a shortage in the production of undecaprenyl phosphate (Und-

P) would be responsible for the observed effects. Und-P is a C55 isoprenoid that acts as the lipid carrier

for transport of cell envelope precursors from the cytoplasm to the extracellular environment[32]. Und-P is

produced by the dephosphorylation of undecaprenyl pyrophosphate (Und-PP), which is in turn

synthesized by UppS (undecaprenyl pyrophosphate synthase) by the addition of 8 isopentenyl units to the

C15 isoprenoid farnesyl-PP[30,32] (Figure 5A). To test whether the observed effects of mevalonate depletion

are indeed caused by a deficiency of Und-P, we constructed a *uppS* deletion/complementation strain

(VL3584, Figure 5B). When this strain was grown in absence of *uppS* expression, cells were elongated and

growth was reduced (Figure 5C-D), thereby phenocopying mevalonate mutants. In addition, UppS

depleted cells contained multiple unconstricted Z-rings and initiated septa that failed to constrict further,

even though peptidoglycan synthesis still occurred as demonstrated by TEM and fluorescence microscopy

(Figure 5D-F). Note that *uppS* was not identified by our sCRilecs-seq screen, likely because it is in an operon

with several other essential genes not directly involved in cell division, such as *cdsA* and *proS*[49]. We thus

conclude that the negative effects caused by depletion of the mevalonate operons can be explained by

insufficient production of Und-PP and subsequently Und-P.

The isoprenoid Und-P carries precursors of the capsule, teichoic acids and peptidoglycan across the

membrane[32]. The observed phenotype of mevalonate and *uppS* mutants could therefore be caused by a

shortage in either one or a combination of these components. To test which of these pathways contribute

to the observed phenotypes upon mevalonate depletion, we tested each of them individually. First, we

deleted the *cps* operon responsible for capsule synthesis in our D39V strain[57]. As shown in Figure S3A-B,

cps mutant cells were not elongated nor was growth rate affected, as also shown before[58], demonstrating

that a shortage of capsule synthesis does not contribute to the mevalonate depletion phenotype. Next,

we focused on the possible contribution of both teichoic acid and peptidoglycan synthesis by depleting

the enzymes responsible for coupling the final cytoplasmic precursors to Und-P before transport to the

outside of the cell. For peptidoglycan, it is known that the MurNAc-pentapeptide moiety that is produced

in the cytoplasm is coupled to Und-P by MraY[32]. Based on sequence similarity, bioinformatic analysis

indicated that the enzyme that does the same for teichoic acid precursors in *S. pneumoniae* is SPV_1620[59].

Deletion/complementation strains for these genes were created and the effect of depletion of either

MraY or SPV_1620 was assessed (strain VL3585 and VL3586, respectively). Surprisingly, depletion of

neither MraY nor SPV_1620 produced cell morphologies that resembled mevalonate operon depletions

at the time we usually assess them (Figure 5G-H, time point 4h). However, since at this time cells were

severely malformed and many had already lysed, we decided to follow these depletions through time and

also record morphologies at earlier time points. Figure 5G shows the progression of these depletions in

terms of average cell length, while Figure 5H shows representative morphologies. From this analysis it

became clear that, although depletion of both MraY and SPV_1620 led to increased cell length, the

phenotype obtained upon SPV_1620 depletion differed profoundly from the phenotypes found when

depleting mevalonate operons. Depletion of MraY, on the other hand, was highly similar to the

mevalonate phenotype at early time points. We thus conclude that a deficiency in the export of

peptidoglycan precursors is mainly responsible for the elongated phenotype that is observed when

components of the mevalonate pathway are depleted. Since we previously showed that this elongated

phenotype is caused by continued peptidoglycan synthesis in the absence of cell constriction, it appears

as though a certain threshold level of peptidoglycan precursors is needed for cells to divide. Cell

elongation, on the other hand, can proceed with a lower amount of peptidoglycan precursors indicating

switch-like behavior between elongation and constriction[43].

Figure 5: Depletion of the mevalonate pathway likely prevents cell division by limiting the export of peptidoglycan precursors which preferentially inhibits cell division. A) After conversion of mevalonic acid into the basic isoprenoid building block isopentenyl-5-PP in the mevalonate pathway, this building block is condensed into the C15 molecule Farnesyl-PP. Farnesyl-PP can be used by UppS for the production of undecaprenyl pyrophosphate (Und-PP), which after dephosphorylation to Und-P by UppP acts as the lipid carrier for the transport of precursors of peptidoglycan, the capsule and teichoic acids across the cell membrane. B) A genetic overview of the UppS depletion strains is shown. The native *uppS* gene was replaced by an antibiotic marker (AB) and a complementation construct under control of the P$_{lac}$ promoter was inserted at the *zip* locus in the *S. pneumoniae* genome. C) Depletion of UppS in VL3584 caused a growth defect similar to depletion of the mevalonate operons. We confirmed that the growth observed after an initial phase of lysis was due to suppressor mutants that are no longer sensitive to UppS depletion (Figure S3C). Data are represented as averages, n ≥ 3. D) Like depletion of the mevalonate operons, depletion of UppS caused an elongated phenotype where cells contained multiple unconstricted FtsZ rings. Images were obtained using strain VL3585 which encodes *ftsZ-mCherry*. E) Transmission Electron Microscopy (TEM) images show that cells elongated due to UppS depletion contained many initiated septa that appeared to be blocked in further progression of constriction (white arrow heads). Images were obtained using strain VL3710 that does not encode fluorescent fusion proteins. F) Pulse labeling *S. pneumoniae* depleted for UppS with the green FDAA, sBADA, and subsequently with the red RADA dye showed sites of active peptidoglycan synthesis, which are in this case all directed at cell elongation. Images were obtained using strain VL3710 that does not encode fluorescent fusion proteins. G) The effect of the depletion of MraY and SPV_1620 on cell length was followed through time using quantitative image analysis. For each biological repeat (n ≥ 3), more than 50 cells were used to calculate the average cell length. Data are represented as the mean ± SEM of these averages. H) Representative morphologies of VL3585 and VL3586 corresponding to the

analysis from panel G are shown. I) A pooled CRISPRi library was constructed in VL3834 (*S. pneumoniae* D39V P*lac*-*dCas9* Δ*mvaA*-

mvaS). This CRISPRi library was grown in the presence of the dCas9 inducer, IPTG, and limiting amounts of mevalonic acid (100

µM). Cultures were sorted based on cell size (FSC); 10% of the population with the highest and lowest values were sorted, as well

as the centermost 70% of the population which served as a control. sgRNAs from the sorted fractions were sequenced and read

counts were compared to identify gene depletions that led to changes in cell size. Volcano plots show the statistical significance

and enrichment for every sgRNA in the fraction with high FSC values (FSC High) and low FSC values (FSC Low) compared to the

control. J) Quantitative analysis of microscopy images shows the changes in cell length upon single or double depletions of the

first mevalonate operon (O1, *mvaS-mvaA*), DivIVA or Pbp2x. Data are represented as violin plots with the mean ± SEM indicated,

n ≥ 3, with each repeat containing > 90 cells except for the double O1 Pbp2x depletion where not enough surviving cells could be

visualized and the threshold was put at 10 cells. Two-sided Wilcoxon signed rank tests were performed and p values were adjusted

with an FDR correction; * $p < 0.05$.

A sCRilecs-seq screen on mevalonate depleted cells to study the underlying genetic network

To identify pathways and genes that are particularly sensitive to reduced mevalonate levels, and to obtain

clues on why a lowered concentration of extracellular peptidoglycan precursors would allow elongation

but not division, we performed a second sCRilecs-seq screen based on cell size (FSC) in a genetic

background in which the first mevalonate operon was deleted (VL3834: D39V P*lac*-*dcas9 lacI* Δ*mvaS-*

mvaA). This way, we could look for gene depletions that aggravate the observed phenotype, i.e. make

cells even longer, and depletions that compensate for the mevalonate-dependent cell elongation. During

construction of this CRISPRi library and prior to selection, cultures were grown in the presence of a high

concentration of mevalonic acid (1 mM) which ensures wild-type growth and normal cell size even though

the first mevalonate operon is deleted (see Figure 3H-I). After construction of the pooled CRISPRi library,

cultures were grown with limiting amounts of external mevalonic acid (100 µm) to trigger the elongated

phenotype caused by mevalonate deficiency (Figure S4A). Simultaneously, dCas9 expression was induced

by the addition of 1 mM IPTG for 3.5h. 10% of the population with the highest and lowest FSC values were

sorted by FACS and the centermost 70% of the population was collected to serve as a control. Comparing

sgRNA read counts from the collected fractions led to the identification of many significantly enriched

targets in the populations with the highest and lowest FSC values (Figure 5I, Figure S4B and Table S2).

GO enrichment analysis identified cell division and closely related GO categories as significantly

overrepresented in elongated cells (Table S3). This suggests that depleting genes involved in cell division

will lead to additional cell elongation upon mevalonate depletion. On the other hand, sgRNAs that are

enriched in the small fraction of the population are often involved in protein expression (transcription,

translation, ribosome assembly, etc.) or energy metabolism. Halting or interfering with protein expression

or energy production therefore appears to reduce cell elongation caused by mevalonate deficiency. The

results of these high-level analyses are clear; directly interfering with cell division reinforces the division

block imposed by mevalonate depletion and inhibiting protein expression or energy production, both of

which are necessary for growth in general, prevents cell elongation.

Additionally, when looking at individual genes and operons found among the significantly enriched hits,

some interesting patterns emerge. For example, the sgRNA targeting *divIVA* was strongly depleted from

the large fraction of the population and also highly enriched in the small fraction, indicating that the

depletion of DivIVA hampers cell elongation upon mevalonate depletion. DivIVA activity was previously

shown to be necessary for cell elongation and DivIVA is known to be phosphorylated by the

serine/threonine kinase StkP that is thought to constitute a molecular switch that governs elongation and

division[43,60-63]. Indeed, we confirmed that a double depletion of the first mevalonate operon and DivIVA

leads to reduced cell lengths in comparison to the depletion of mevalonate alone (Figure 5J), indicating

that also upon mevalonate depletion DivIVA activity assists cell elongation. However, cells depleted for

both mevalonate and DivIVA are still larger than wild-type cells, indicating that a limited amount of

elongation still occurs. Elongation enforced by mevalonate depletion is therefore not fully dependent on

DivIVA activity. Additionally, CRISPRi depletion of the SEDS proteins FtsW and RodA had opposite effects

on cell size in our screen, with FtsW depletion leading to increased cell sizes while RodA depletion causing

decreases in cell size (Table S2). These proteins are thought to be the primary transglycosylases

responsible for peptidoglycan polymerization during cell division and cell elongation, respectively[64-68].

Moreover, CRISPRi depletion of Pbp2b, the transpeptidase that works in conjunction with RodA to

perform peripheral peptidoglycan synthesis[43,67,69], showed up in the shorter cell fraction (Table S2). These

results are in line with the hypothesis that cell elongation in mevalonate depleted cells is caused by

ongoing peripheral peptidoglycan synthesis while septal synthesis is inhibited. If this is true, then

depletion of Pbp2x, the essential pneumococcal PBP required for cell division and the prime target of the

penicillins including amoxicillin[70], should show a synthetic effect upon mevalonate depletion. As *pbp2x* is

located in an operon together with *mraY*, this could not be directly assessed in the sCRilecs-seq screen.

Therefore, we constructed a clean *pbp2x* depletion strain. As shown in Figure 5J, depletion of Pbp2x

indeed led to an increase in cell length, as reported before[34], and this phenotype was augmented under

low mevalonate levels.

The mevalonate pathway as druggable target in *S. pneumoniae*

Our results so far demonstrate that depletion of the mevalonate pathway leads to cell elongation and these morphological effects can be enhanced by targeting other cell division pathways. Since amoxicillin also causes elongation in *S. pneumoniae*, we set out to design a combination treatment strategy using amoxicillin and exploiting the mevalonate depletion phenotype characterized here. In a first step, we confirmed that the antibacterial effect of amoxicillin – but not that of the fluoroquinolone ciprofloxacin – is increased upon mevalonate depletion, since amoxicillin could inhibit growth already at sub-MIC concentrations when mevalonate became limiting (Figure S5A-B). These results demonstrate that, in principle, it is possible to design a treatment strategy in which amoxicillin is potentiated by targeting the mevalonate pathway.

In a second step, we looked for chemical compounds that could block either the mevalonate pathway or the downstream production of Und-P. Three compounds were selected; simvastatin and farnesol are expected to block the HMG-CoA reductase MvaA that catalyzes one of the first steps of the mevalonate pathway[71-73], while clomiphene was shown to inhibit UppS in *S. aureus*[33]. To confirm the specificity of these compounds for the mevalonate pathway or downstream production of Und-P, we first tested whether they could induce the expected cell elongation in *S. pneumoniae*. Since only the addition of clomiphene resulted in filamentation under our conditions (Figure 6A), we selected this compound for further testing. Indeed, time lapse analysis showed considerable cell elongation in the presence of clomiphene, albeit less than upon depletion of the mevalonate operons potentially due to the lag in severe effects upon gradual genetic depletion (Figure 6B and Movie S5). Additionally, clomiphene most likely targets the production of Und-P, since sub-MIC concentrations of clomiphene led to severe growth defects in cultures that were slightly depleted for mevalonate (Figure S5C).

We next combined clomiphene with either amoxicillin or ciprofloxacin at several different concentrations and monitored the growth of an *S. pneumoniae* D39V wild-type strain by measuring OD. The success of growth was quantified by calculating the area under the curve for a growth period of 10 hours. As can be seen in Figure 6C, clomiphene did not affect the efficacy of ciprofloxacin. However, for amoxicillin, potentiation by clomiphene was observed (Figure 6D). This is consistent with time lapse analyses that show that survival of *S. pneumoniae* was much more affected when both compounds were combined (Figure 6E and Movie S6). Indeed, in the presence of clomiphene, the concentration of amoxicillin necessary to block pneumococcal growth decreased by a factor 2 to 8 (Table 1).

Resensitizing antibiotic-resistant *S. pneumoniae* strains using clomiphene

We next tested whether the potentiation by clomiphene is also present in clinically-relevant amoxicillin-resistant *S. pneumoniae*. We therefore assessed the growth of a panel of resistant strains in the presence of different concentrations of amoxicillin and clomiphene. The strains we tested are clinical isolates of strains 19F, Spain-23F and 11A[5,74]. Results are presented in Figure 6F-H and Table 1 and show that clomiphene potentiated the antimicrobial activity of amoxicillin also against clinical resistant strains. Moreover, the potentiation was even stronger in these resistant genetic backgrounds. Indeed, the concentration of amoxicillin necessary to block growth of these resistant strains decreased by a factor 16 to 64 in the presence of 8 µg/ml of clomiphene (Table 1), thereby reducing resistance below the EUCAST clinical breakpoint for sensitivity (0.5 µg/ml)[75]. Our rationally designed combination treatment strategy can thus resensitize amoxicillin-resistant *S. pneumoniae* strains so that they once again become fully susceptible to amoxicillin *in vitro*. This treatment strategy could therefore prove useful in the fight against amoxicillin-resistant *S. pneumoniae* infections.

Figure 6: Clomiphene, an inhibitor of Und-P production, potentiates amoxicillin. A) The effect of several potential inhibitors of Und-P production on the cell length of *S. pneumoniae* D39V (VL333) was tested (Sim 4 µg/ml, Far 4 µg/ml, Clom 8 µg/ml). Quantitative analysis of microscopy images shows that cell length increased upon treatment with clomiphene. Data are represented as violin plots with the mean cell length of every biological repeat indicated with orange dots. The size of these dots indicates the number of cells recorded in each repeat, ranging from 177 to 6464 cells. Black dots represent the mean ± SEM of all recorded means, n ≥ 3. Two-sided Wilcoxon signed rank tests were performed against wt without treatment as control group

(dotted line) and p values were adjusted with an FDR correction; ** p < 0.01. B) Snapshot images of phase contrast time lapse

microscopy of *S. pneumoniae* D39V (VL333) in the presence of clomiphene (8 µg/ml). Scale bar, 5 µm. C-D) OD_{595nm} growth curves

were constructed for *S. pneumoniae* D39V (VL333) in the presence of different concentrations of clomiphene and ciprofloxacin

(C) or amoxicillin (D). Heatmaps of the area under the resulting growth curves are shown. Number of biological repeats for all

experiments, n ≥ 3. E) Snapshot images of phase contrast time lapse microscopy of *S. pneumoniae* D39V (VL333) in the presence

of clomiphene (8 µg/ml) and amoxicillin (0.016 µg/ml). Scale bar, 5 µm. F-H) OD_{595nm} growth curves were constructed for *S.*

pneumoniae 19F (F), Spain-23F (G) and 11A (H) in the presence of different concentrations of clomiphene and amoxicillin.

Heatmaps of the area under the resulting growth curves are shown. Number of biological repeats for all experiments, n ≥ 3. I)

The effect of the combination treatment with amoxicillin and clomiphene was tested *in vivo* using a pneumonia superinfection

model with a clinical isolate of *S. pneumoniae* serotype 19F. Mice (n=8 per group) were infected intranasally first with H3N2 virus

and then 7 days later with pneumococcus 19F. Mice were treated at 8 h and 12h with clomiphene, amoxicillin, combination of

both, or left untreated. Lungs were collected 24 h post-infection to measure the bacterial load. CFU counts for individual mice

are shown, and the bars represent the median value. The data were compared in a Kruskall-Wallis test (One-Way ANOVA), ** p

< 0.01. Wt, wildtype; Sim, simvastatin; Far, farnesol; Clom, clomiphene; Amx, amoxicillin.

Table 1: MIC values for amoxicillin/clomiphene and ciprofloxacin/clomiphene drug combinations (µg/ml).

		Clomiphene	
	0 µg/ml	4 µg/ml	8 µg/ml
S. pneumoniae D39V			
Ciprofloxacin	1	1	1
Amoxicillin	0.032	0.016	0.004
S. pneumoniae 19F			
Amoxicillin	2	1	0.125
S. pneumoniae Spain-23F			
Amoxicillin	2	1	0.032
S. pneumoniae 11A			
Amoxicillin	4	2	0.125

We next tested whether this treatment strategy would prove useful in an *in vivo* setting, using a murine

pneumonia disease model (see Materials and Methods)[76]. Mice were infected on day 1 with influenza A

virus and superinfected on day 7 with *S. pneumoniae* 19F. Animals were treated with 5 mg of clomiphene

and/or 1 mg of amoxicillin at 8 and 12 hours post-pneumococcal infection, respectively. Twenty-four

hours post infection, the bacterial load in the lung was determined. Whereas treatment with amoxicillin

alone significantly lowered bacterial counts, administering only clomiphene did not influence the bacterial

load (Figure 6I). Treating mice with both clomiphene and amoxicillin did not show a stronger effect than

treatment with amoxicillin alone, meaning that the potentiation of amoxicillin we observed *in vitro* could

not be detected *in vivo* (Figure 6I). The same was true for CFU counts in the spleen (Figure S5D). To figure

out why our combination treatment strategy was not effective *in vivo*, we tried to estimate the active

concentration of clomiphene in the lung. Our *in vitro* tests showed that potentiation occurs at

concentrations of clomiphene of 4 µg/ml and higher, but these concentrations might not be achieved *in*

vivo. We therefore tested the growth of *S. pneumoniae* D39V *in vitro* in the presence of bronchoalveolar

lavage (BAL) fluid collected from mice treated with clomiphene. This experiment indicated that the active

concentration of clomiphene in the lung was lower than 4 µg/ml, which could explain the absence of

amoxicillin potentiation (Figure S5E).

Discussion

Here, we rationally designed a combination treatment strategy that resensitizes resistant *S. pneumoniae*

strains towards clinically relevant concentrations of amoxicillin, one of the most widely used antibiotics

to fight this human opportunistic pathogen[15-17]. By combining amoxicillin with the FDA-approved fertility

drug clomiphene[77-79], we were able to reduce amoxicillin MIC values of resistant strains 16 to 64-fold *in*

vitro, thereby decreasing them below the EUCAST clinical breakpoint for sensitivity.

sCRilecs-seq identifies several targets that regulate S. pneumoniae cell size

This combination treatment strategy was based on the results of the sCRilecs-seq screen developed here.

sCRilecs-seq is a high-throughput single-cell-based screening approach that relies on genome-wide

CRISPRi depletion combined with sorting of cells that display a phenotype of interest. We here screened

for CRISPRi gene depletions that, like amoxicillin treatment, result in cell elongation. The rationale behind

this strategy is that synergy is most often detected between compounds that target the same process[36].

Although we used the sCRilecs-seq method to screen for increases in cell size, this approach could easily

be adapted to asses any phenotype of interest that can be measured by flow cytometry. In contrast to

'classic' CRISPRi-seq screens[22-27], our approach is therefore not limited to measuring changes in fitness but

can be used to evaluate a wide array of different phenotypes. Moreover, we show that sCRilecs-seq hits

are highly reliable since no false positives were detected in our screen. Interestingly, significant hits

include a number of genes of unknown function that we can now implicate in the maintenance of proper

cell size. Our sCRilecs-seq approach is thus able to uncover novel gene functions and can help to expand

knowledge on any process of interest that can be evaluated using flow cytometry. On the other hand,

several genes known to be involved in cell size regulation were not identified as significant hits. Such false

negatives could be caused by a variety of reasons. For example, simultaneous repression of all genes in

an operon can obscure phenotypes expected upon repression of individual genes. Additionally, depletion

of genes that lead to cell lysis cause the corresponding sgRNAs to disappear from the library thereby

complicating their identification as hits. Depletion strains sensitive to mechanical vortexing might also be

missed. Finally, the high variation detected across different repeats led to large P-values for many targets,

even when they were very highly enriched. This high variation thereby limits the number of significant

hits that could be detected.

Depletion of mevalonate pathway genes leads to cell elongation through inhibition of cell division

The sCRilecs-seq screen performed here revealed an important role for the mevalonate pathway in

maintaining proper cell size. Indeed, depletion of the genes involved in this pathway led to a very strong

increase in cell length. Results obtained here strongly indicate that this cell elongation is due to a

deficiency in the production of Und-P, the lipid carrier that translocates all different cell envelope

precursors across the cell membrane[32], and subsequent limitation of the amount of peptidoglycan

building blocks that is being transported. The limiting amount of peptidoglycan precursors available for

cell wall synthesis led to an interesting phenotype. Whereas the peptidoglycan synthesis rate in general

decreased, peptidoglycan synthesis for cell elongation continued long after constriction for cell division

was inhibited. This observation points towards a difference in affinity and/or regulation of elongation and

constriction, where it appears that a certain threshold concentration of peptidoglycan precursors needs

to be exceeded in order for cell division to take place. Several potential explanations for this observation

can be put forward. First, it is possible that peptidoglycan synthesis enzymes dedicated to cell division

have a lower affinity for peptidoglycan precursors than their counterparts that function in cell elongation.

When cell wall precursors become limiting, the enzymes involved in septal ring closure would be

outcompeted and cell division would cease while elongation still occurs. On the other hand, it has

previously been suggested that a more complex regulatory mechanism decides whether cells elongate or

divide. The serine/threonine kinase StkP and its cognate phosphatase PhpP were suggested to constitute

a molecular switch that coordinates septal and peripheral peptidoglycan synthesis through

phosphorylation and dephosphorylation of its targets[43,60-62]. This switch is regulated by GpsB, which is

necessary for maximal phosphorylation in *S. pneumoniae* and is essential for viability[41,42,80]. Interestingly,

depletion of GpsB leads to an elongated phenotype with unconstricted cell division sites[41,42] that is highly

reminiscent of the mevalonate depletion phenotype, hinting at possible crosstalk between both

pathways. Results presented here indicate that the decision between elongation or division depends on

the amount of peptidoglycan precursors that is exported. Potentially, the concentration of precursor

molecules influences the activity of GpsB, StkP and/or PhpP and thereby activates this switch. Since StkP

contains PASTA domains that are known to bind peptidoglycan precursors[43,60,61,81], it has been suggested

that its activity is indeed regulated in response to the concentration of peptidoglycan building blocks

present[43,60,61]. However, it was previously shown that even though PASTA domains are necessary for StkP

activation, peptidoglycan binding is not required[81], arguing against StkP being a sensor for peptidoglycan

precursor levels. Moreover, the depletion of DivIVA upon reduced mevalonate conditions still allowed a

modest amount of cell elongation to occur. Since DivIVA is thought to be an important part of the

molecular switch between elongation and division instigated by StkP[41,62,63], it appears as though

elongation upon mevalonate depletion is at least partially independent of this molecular switch.

Therefore, further research will be necessary to determine why elongation is favored over division when

peptidoglycan precursors become limiting. Finally, it is possible that depletion of mevalonate pathway

components and probable subsequent peptidoglycan precursor shortage induces transcriptional or post-

transcriptional regulatory pathways that differentially affect the activity of protein complexes involved

cell elongation and division, thereby leading to filamentation.

Inhibition of Und-P synthesis by clomiphene resensitizes resistant S. pneumoniae to amoxicillin

We successfully exploited the knowledge gained on the mevalonate depletion phenotype to fight

amoxicillin-resistant *S. pneumoniae* strains. To do so, we confirmed that clomiphene, a compound known

to block Und-P production in *S. aureus* and potentiate β-lactam antibiotics, even towards MRSA[33], elicits

the elongated mevalonate depletion phenotype in *S. pneumoniae* and therefore most likely also inhibits

Und-P production in this bacterium. We next combined clomiphene with the β-lactam antibiotic

amoxicillin that preferentially interferes with cell division[69,70]. Results demonstrated that clomiphene can

enhance the antimicrobial effect of amoxicillin in a D39V lab strain, although this effect was rather limited.

However, potentiation by clomiphene becomes much stronger in amoxicillin-resistant strains such as 19F,

Spain-23F and 11A. In *S. pneumoniae*, resistance towards β-lactam antibiotics is mostly caused by mosaic

penicillin binding proteins (PBPs). These resistance-conferring mosaic PBPs are formed by recombination

events following horizontal gene transfer from β-lactam resistant donor strains[4,7]. We speculate that these

mosaic PBPs display suboptimal activity which allows them to remain active during amoxicillin treatment

but fail to carry out their task if the amount of peptidoglycan precursors available to them becomes

limiting due to a deficiency in Und-P production.

Whereas this combination treatment strategy works remarkably well *in vitro*, *in vivo* results using a murine

pneumonia disease model did not show benefits of combining amoxicillin treatment with clomiphene,

likely because the *in vivo* concentration of active clomiphene was too low in the lungs. Nonetheless, we

believe that clomiphene represents a promising starting point for the development of an optimized

antibacterial compound that can be used in combination with amoxicillin and potentially other β-lactams.

Clomiphene is an FDA-approved prodrug that is administered as a racemic mixture of two stereoisomers

and is metabolized by the liver into active compounds that stimulate ovulation in anovulatory women[77-

79,82]. However, since we observe antibacterial effects and amoxicillin potentiation *in vitro*, it seems likely

that one or both of the prodrug isomers exert the desired effect. Determining which clomiphene

stereoisomer has the highest antibacterial activity and designing non-metabolizable derivatives that are

active at lower concentrations would be the first step to further optimize the here proposed combination

treatment strategy for *in vivo* use. We hope that such an optimized antibacterial compound can be

exploited for the eradication of amoxicillin-resistant *S. pneumoniae* infections and could potentially also

target other species and/or strains with a different resistance profile towards β-lactam antibiotics. We

therefore believe that further investigation into this combination treatment strategy holds much promise

in combatting the ever-increasing amount of antibiotic-resistant bacterial infections.

Materials & Methods

Bacterial strains and growth conditions

All pneumococcal strains used in this work are listed in Table 2. Unless specified otherwise, strains used throughout this work are derivatives of *S. pneumoniae* D39V[49]. In general, genomic changes were introduced by homologous recombination after transformation of a linear DNA molecule into *S. pneumoniae*. This linear DNA was either obtained through a one-step PCR reaction starting from a different strain that already carried the desired transformation product (e.g. *hlpA-gfp* and *ftsZ-mCherry*) or through golden gate cloning in which three different PCR products were ligated. The first and last of these PCR products were homologous to the up- and downstream regions of the genome where a deletion or insertion had to be made. In case of insertion of expression cassettes into the *zip* locus, these fragments also contained a P_{lac} promoter and a spectinomycin (or trimethoprim) resistance marker through amplification of fragments of the pPEPZ plasmid[53]. In case of insertion of expression cassettes into the *bgaA* locus, these fragments also contained a P_{tet} promoter and a tetracycline resistance marker through amplification of fragments from strain VL2212[23]. The third, middle fragment contained an antibiotic marker or a sequence to be inserted into the genome. Golden gate cloning sites were introduced into the PCR fragments as part of the primers. PCR fragments were digested with either BsaI, Esp3I or SapI, followed by ligation and transformation. In case individual sgRNAs were cloned into *S. pneumoniae*, the sgRNA sequences were first inserted into the pPEPZ-sgRNAclone plasmid in *E. coli*, as described previously[23]. These integrative plasmids were then transformed into *S. pneumoniae*. Transformation was performed with cells that were made competent by addition of the Competence Stimulating Peptide (CSP-1) using a previously published protocol[83]. Primers that were used for cloning are listed in Table 3. *hlpA-gfp* was amplified from VL2226 with primers OVL47&48. *ftsZ-mCherry* was amplified from VL1630 with primers OVL898&901. Both PCR products were transformed to VL1998 to create VL3117. Complementation constructs for mevalonate operon 1 (*mvaS-mvaA*) and operon 2 (*mvk-mvaD-mvaK2-fni*) were made by amplification of the corresponding operons using primer pairs OVL3962&63 and OVL3958&59, respectively. Up- and downstream regions for insertion into the *zip* locus were amplified using primer pairs OVL3493&2181 and OVL2182&3496. For the complementation constructs of the *uppS* and *mraY* genes that were amplified using primers OVL4583&84 and OVL4595&96, respectively, up- and downstream regions of the *zip* locus were created using OVL3493&4341 and 2182&3496. The complementation construct for SPV_1620 was created by amplification of the gene using OVL3671&72 and up- and downstream *zip* regions were produced using OVL3493&3650 and 3649&3496. Deletion of

the corresponding native genes was performed by replacing them with a promoter-less kanamycin

resistance cassette that was expressed using the native promoter of the deleted genes. The kanamycin

cassette was amplified from the pPEPY plasmid[53] with primers OVL3981&82. The up- and downstream

regions of the genes to be deleted were amplified using OVL4069&70 and OVL4071&72 for mevalonate

operon 1 (*mvaS-mvaA*), OVL4585&86 and OVL4587&88 for *uppS*, OVL4597&98 and OVL4599&4600 for

mraY and OVL3677&4601 and OVL 4602&3680 for *SPV_1620*. Due to low expression levels [84], mevalonate

operon 2 (*mvk-mvaD-mvaK2-fni*) was replaced with a Km cassette that carried its own constitutive

promoter. This fragment was amplified with OVL3983&82 from pPEPY[53] and up- and downstream regions

for this operon were amplified with primer pairs OVL4061&62 and OVL4063&64. A complementation

construct for *divIVA* was made by amplification of this gene using primers OVL5707 &08. Up- and

downstream regions for insertion into the *bgaA* locus were amplified using primer pairs

OVL2077&OVL5705 and OVL5706&1369. Deletion of *divIVA* was performed by replacement with a

promoter-less chloramphenicol resistance cassette that was expressed using the *divIVA* promoter. The

chloramphenicol cassette was amplified from VL3117 with primers OVL5727&28. The up- and

downstream regions of *divIVA* were amplified using OV5729&30 and OVL5731&32, respectively. A

complementation construct for *pbp2x* was made by amplification of this gene using primers OVL6276&77.

Up- and downstream regions for insertion into the *bgaA* locus were amplified using primer pairs

OVL2077&OVL5717 and OVL5718&1369. Deletion of *pbp2x* was performed by replacement with a

promoter-less erythromycin resistance cassette that was expressed using the *pbp2x* promoter. The

cassette was amplified from pJWV502[23] with primers OVL2933&34. The up- and downstream regions of

pbp2x were amplified using OV6214&15 and OVL6216&17, respectively.

Strains were grown in liquid C+Y medium (pH = 6.8)[85] at 37°C without shaking under normal atmospheric

conditions. Plating was performed inside Columbia agar with 3% sheep blood and plates were incubated

at 37°C in a controlled atmosphere containing 5% CO_2.

Antibiotics for selection were used at the following concentrations: chloramphenicol (Cm) 3 µg/ml,

erythromycin (Ery) 0.5 µg/ml, gentamicin (Gm) 40µg/ml, kanamycin (Km) 150 µg/ml, spectinomycin (Spec)

100 µg/ml, tetracycline (Tc) 0.5 µg/ml, Trimethoprim (Trm) 10 µg/ml. When necessary, P$_{lac}$ promoters

were induced with various amounts of IPTG; 1 mM IPTG for induction of dCas9 for the sCRilecs-seq

screens, 20 µM IPTG for complementation of mevalonate operon 1 (*mvaS*, *mvaZ*), 100 µM for

complementation of *uppS* and 1 mM IPTG for complementation of mevalonate operon 2 (*mvk*, *mvaD*,

mvaK2, *fni*), *mraY* or *SPV_1620*. When necessary, P$_{tet}$ promoters were induced with 500 ng/ml

anhydrotetracycline (aTc). Where appropriate, mevalonic acid was added to cultures at the concentration

indicated in the text (ranging from 100 μM to 1 mM). When strains lacking the first mevalonate operon

had to be grown under non-limiting conditions, 1 mM mevalonic acid was used. The concentrations of

antibiotics and other compounds that were used to assess their antibacterial activity are indicated in the

main text and/or figures and/or figure legends.

Table 2: *S. pneumoniae* strains used in this study.

Strain	Genome	Reference
D39V	Serotype 2, wildtype	49
19F	Serotype 19F, clinical isolate	This work
Spain-23F	Serotype 23F, PMEN1, clinical isolate	German National Reference Center for Streptococci
11A	Serotype 11A, PMEN3, clinical isolate	German National Reference Center for Streptococci
VL333	D39V *prs1*::*lacI-tetR-Gm*	Lab collection
VL567	D39V Δ*cps*::*Cm*	86
VL1313	11A	5
VL1630	D39V *ftsZ-mCherry-Ery bgaA*::P$_{zn}$-*gfp-stkP*	Lab collection
VL1998	D39V *prs1*::F6-*lacI-Gm bgaA*::P$_{lac}$-*dCas9-Tc*	21
VL2226	D39V *hlpA-gfp-Cm*	38
VL3117	VL1998 *hlpA-gfp-Cm ftsZ-mCherry-Ery*	This work
VL3404	VL333 *hlpA-gfp-Cm ftsZ-mCherry-Ery*	This work
VL3565	VL3404 *zip*::P$_{lac}$-*mvk-mvaD-mvaK2-fni-Spec* Δ*mvk-mvaD-mvaK2-fni*::*Km*	This work
VL3567	VL3404 *zip*::P$_{lac}$-*mvaS-mvaA-Spec* Δ*mvaS-mvaA*::*Km*	This work
VL3584	VL3404 *zip*::P$_{lac}$-*uppS-Spec* Δ*uppS*::*Km*	This work
VL3585	VL3404 *zip*::P$_{lac}$-*mraY-Spec* Δ*mraY*::*Km*	This work
VL3586	VL3404 *zip*::P$_{lac}$-*SPV_1620-Trm* Δ*SPV_1620*::*Km*	This work
VL3671	VL3117 *zip*::P3-*sgRNA035*	This work
VL3672	VL3117 *zip*::P3-*sgRNA036*	This work
VL3673	VL3117 *zip*::P3-*sgRNA046*	This work
VL3674	VL3117 *zip*::P3-*sgRNA087*	This work
VL3675	VL3117 *zip*::P3-*sgRNA100*	This work
VL3676	VL3117 *zip*::P3-*sgRNA121*	This work
VL3677	VL3117 *zip*::P3-*sgRNA135*	This work
VL3678	VL3117 *zip*::P3-*sgRNA355*	This work
VL3679	VL3117 *zip*::P3-*sgRNA432*	This work
VL3680	VL3117 *zip*::P3-*sgRNA455*	This work
VL3681	VL3117 *zip*::P3-*sgRNA460*	This work
VL3682	VL3117 *zip*::P3-*sgRNA461*	This work
VL3683	VL3117 *zip*::P3-*sgRNA503*	This work
VL3684	VL3117 *zip*::P3-*sgRNA573*	This work

VL3685	VL3117 *zip*::P3-*sgRNA583*	This work
VL3686	VL3117 *zip*::P3-*sgRNA590*	This work
VL3687	VL3117 *zip*::P3-*sgRNA628*	This work
VL3688	VL3117 *zip*::P3-*sgRNA673*	This work
VL3689	VL3117 *zip*::P3-*sgRNA757*	This work
VL3690	VL3117 *zip*::P3-*sgRNA780*	This work
VL3691	VL3117 *zip*::P3-*sgRNA785*	This work
VL3692	VL3117 *zip*::P3-*sgRNA796*	This work
VL3693	VL3117 *zip*::P3-*sgRNA813*	This work
VL3694	VL3117 *zip*::P3-*sgRNA822*	This work
VL3695	VL3117 *zip*::P3-*sgRNA824*	This work
VL3696	VL3117 *zip*::P3-*sgRNA900*	This work
VL3697	VL3117 *zip*::P3-*sgRNA1019*	This work
VL3699	VL3117 *zip*::P3-*sgRNA1064*	This work
VL3700	VL3117 *zip*::P3-*sgRNA1236*	This work
VL3701	VL3117 *zip*::P3-*sgRNA1240*	This work
VL3702	VL333 Δ*mvaS-mvaA*::Km	This work
VL3708	VL333 *zip*::P*lac*-*mvk-mvaD-mvaK2-fni-Spec* Δ*mvk-mvaD-mvaK2-fni*::Km	This work
VL3709	VL333 *zip*::P*lac*-*mvaS-mvaA-Spec* Δ*mvaS-mvaA*::Km	This work
VL3710	VL333 *zip*::P*lac*-*uppS-Spec* Δ*uppS*::Km	This work
VL3711	VL333 *zip*::P*lac*-*mraY-Spec* Δ*mraY*::Km	This work
VL3712	VL333 *zip*::P*lac*-*SPV_1620-Trm* Δ*SPV_1620*::Km	This work
VL3834	VL1998 Δ*mvaS-mvaA*::Km	This work
VL4273	VL333 *bgaA*::P*tet*-*pbp2x-Tc* Δ*pbp2x*::Ery	This work
VL4274	VL3709 *bgaA*::P*tet*-*pbp2x-Tc* Δ*pbp2x*::Ery	This work
LD0001	VL3709 *bgaA*::P*tet*-*divIVA-Tc* Δ*divIVA*::Cm	This work

735 **Table 3: Primers used in this study.**

Primer	Sequence
OVL47	GATTGTAACCGATTCATCTG
OVL48	GGAATGCTTGGTCAAATCTA
OVL898	CCAACAAGCTTCACAAAATAAACCG
OVL901	CTTATCCGTTGCACGCTGACTC
OVL1369	GTCTTCTTTTTTACCTTTAGTAACTACTAATCCTGCAC
OVL2077	ATTCCTTCTTAACGCCCCAAGTTC
OVL2181	GCGTCACGTCTCAGCATTATTTTTCCTCCTTATTTAT
OVL2182	GCGTCACGTCTCACGGATCCCTCCAGTAACTCGAGAA
OVL2933	GATCGGTCTCGAGGAATTTTCATATGAACAAAAATATAAAATATTCTCAA
OVL2934	GATCGGTCTCGTTATTTCCTCCCGTTAAATAATAGATAACTATTAAAAAT
OVL3493	GCCAATAAATTGCTTCCTTGTTTT
OVL3496	ATGACACGGATTTTAAGAATAATTCTTTC
OVL3649	TGTGTGGCTCTTCGAGAACTCGAGAAAAAAAAACCGCGCCC
OVL3650	TGTGTGGCTCTTCGGTTTCATTATTTTTCCTCCTTATTTATTTAGATCTTAATTGTGAGC
OVL3671	CTGGTAGCTCTTCCAACATGCTGAAATGGGAAGACTTGCCTG

OVL3672	CTGGTAGCTCTTCCTCTTTATTTTAGTACCTCAAACACGGTT
OVL3677	GTATAGTAAGCTGGCAGAGAATATC
OVL3680	ATACTTTTTAGGGACAGGATCAC
OVL3958	GCGTCACGTCTCAATGCTCGTCTAGTAAAAGGAAAAAATGACAAAAAAA
OVL3959	GCGTCACGTCTCATCCGTTACGCCTTTTTCATCTGATCATTTG
OVL3962	GCGTCACGTCTCAATGCAGTATAGAACGATTTTTTACATGAATGATAAAACAG
OVL3963	GCGTCACGTCTCATCCGTTATGATCTTAAATTTTCGAGATAGCGCT
OVL3981	GCGTCACGTCTCAATGGCTAAAATGAGAATATCACCGG
OVL3982	GCGTCACGTCTCACTAAAACAATTCATCCAGTAAAATATAATATTTTATTTTCTCC
OVL3983	GCGTCACGTCTCAGAGGACGCGCAAGCTG
OVL4061	CACTACCAATTGGTGAAGTTGCT
OVL4062	GCGTCACGTCTCACCTCTTTTTCCTTTTACTAGACGAAAAAACGTC
OVL4063	GCGTCACGTCTCATTAGGGCGTAACCAGCGCC
OVL4064	TACAGGTACGATGATTTTGGTCGT
OVL4069	AGCTGAAGATAAAGCCTGTAACCA
OVL4070	GCGTCACGTCTCACCATGTAAAAAATCGTTCTATACTATTTTATCACAAATGG
OVL4071	GCGTCACGTCTCATTAGCATAAAAACTCAGACGAATCGGTCT
OVL4072	ACAGCGCCGATTATTTCCTTTG
OVL4341	GCGTCACGTCTCAATTTATTTAGATCTTAATTGTGAGCGCTC
OVL4583	GCGTCACGTCTCAAAAATTTTTGAATAGGAATAAGATCATGTTTGGATTTT
OVL4584	GCGTCACGTCTCATCCGCTAAACTCCTCCAAATCGGCG
OVL4585	TCCAGATTTTTCTTATGAGGAAACCTTATT
OVL4586	GCGTCACGTCTCACCATGATCTTATTCCTATTCAAAAATCTATCGTTTCATT
OVL4587	GCGTCACGTCTCATTAGGGAGGAGTTTAGGAGGAAATATGACC
OVL4588	CTGTACTGTCAACTATCATAAAGATAATGGT
OVL4595	GCGTCACGTCTCAAAAATATTAACTTTAGGAGACTAATATGTTTATTTCCATCAG
OVL4596	GCGTCACGTCTCATCCGTTACATCAAATACAAAATTGCGAGGGT
OVL4597	AGATTGCTGACGAGAAAAATGGTG
OVL4598	GCGTCACGTCTCACCATATTAGTCTCCTAAAGTTAATGTAATTTTTTTAATGTCC
OVL4599	GCGTCACGTCTCATTAGGAATGGCACCCTGATGTTTCA
OVL4600	AATAAATCATCCATGTTGTTAAAATTATTAAAATTGTTGT
OVL4601	GCGTCACGTCTCACCATGCTGTTCTCCTTTGTTTTTATTATAC
OVL4602	GCGTCACGTCTCATTAGAGTAGTCATAAGAAAATGAGTACAG
OVL5705	GCGTCAGGTCTCAATTTATTTAGATCTACTCTATCAATGATAGAGTTATTATACTCT
OVL5706	GCGTCAGGTCTCAGCGTAAGGAAATCCATTATGTACTATTTCTG
OVL5707	GCGTCAGGTCTCAAAAATTTAAGTAAGTGAGGAATAGAATGCCAATTACA
OVL5708	GCGTCAGGTCTCAACGCCTACTTCTGGTTCTTCATACATTGGG
OVL5717	GCGTCACGTCTCAATTTATTTAGATCTACTCTATCAATGATAGAGTTATTATACTCT
OVL5718	GCGTCACGTCTCAGCGTAAGGAAATCCATTATGTACTATTTCTG
OVL5727	GCGTCAGGTCTCAATGAACTTTAATAAAATTGATTTAGACAATTGGAAGAG
OVL5728	GCGTCAGGTCTCATTATAAAAGCCAGTCATTAGGCCTATCT
OVL5729	CTCCTTTTTTAACTCCTTTTATCAATCCTCA
OVL5730	GCGTCAGGTCTCATCATTCTATTCCTCACTTACTTAATAATAACTGGACG
OVL5731	GCGTCAGGTCTCAATAACTCCAGTGCATCCGACAGG
OVL5732	ACCAAGTCCATTTCTTTACGTTTGAC
OVL6214	GCGCGTAAGATTGAGCAA

OVL6215	GATCGGTCTCATCCTATCTTACTCCGCTATTCTAATATTTTCA
OVL6216	GATCGGTCTCGATAAATCAAGGACATTAAAAAAATTACATTAACTT
OVL6217	ACATCACCCATAAAGACCTTG
OVL6276	GCGTCACGTCTCAAAATTTAGAATAGCGGAGTAAGATATGAAGTGG
OVL6277	GCGTCACGTCTCAACGCTTAGTCTCCTAAAGTTAATGTAATTTTTTAATGTCC

Mechanical disruption of *S. pneumoniae* cell chains

Cultures were diluted 3x in PBS to a final volume of 1 ml in screw cap tubes. Tubes were placed into a FastPrep-24™ 5G Instrument (MP Biomedicals) with QuickPrep-3 adapter and shaken with a speed of 6.0 m/s for 30 s. This protocol was validated using VL3117 that was grown for 3.5h in C+Y medium. Samples were either subjected to mechanical chain disruption using the FastPrep-24™ 5G or not, after which both samples were analyzed by microscopy and flow cytometry.

Construction of the CRISPRi libraries

CRISPRi libraries were constructed by transformation of the desired strains with a pool of 1499 different pPEPZ integrative plasmids carrying constitutively-expressed sgRNA sequences that together target the entire genome[23]. sgRNAs are under control of the constitutive P3 promoter, while the *dcas9* gene is inserted chromosomally under control of the inducible P$_{lac}$ promoter. The 1499 sgRNAs were designed to each target a specific operon. All sgRNA sequences together with their targets and potential off-targets were published previously[23]. For the initial screen, D39V P$_{lac}$-*dcas9 hlpA-gfp ftsZ-mCherry* (VL3117) was transformed. Our second sCRilecs-seq screen was performed after transformation of D39V P$_{lac}$-*dcas9* Δ*mvaS* Δ*mvaA* (VL3834) in the presence of 1 mM mevalonic acid. For both libraries, at least 10^5 individual transformants were obtained and collected.

sCRilecs-seq screen

The transformed D39V P$_{lac}$-*dcas9 hlpA-gfp ftsZ-mCherry* (VL3117) library was grown for 3.5h in C+Y medium supplemented with 1 mM IPTG. The D39V P$_{lac}$-*dcas9* Δ*mvaS* Δ*mvaA* (VL3834) library was grown for 3.5h in C+Y medium supplemented with 1 mM IPTG and 100 µM mevalonic acid. Cell chains were mechanically disrupted (see "Mechanical disruption of *S. pneumoniae* cell chains") and further diluted into PBS 3-10x based on culture density.

Cells were sorted using a FACSAria™ IIIu Cell Sorter (BD Biosciences) equipped with violet, blue and red lasers and a 70 µm nozzle. In case of the D39V P$_{lac}$-*dcas9 hlpA-gfp ftsZ-mCherry* (VL3117) library, cells were gated based on FSC, SSC and GFP fluorescence. For the D39V P$_{lac}$-*dcas9* Δ*mvaS* Δ*mvaA* (VL3834) library, cells were gated based on FSC and SSC only. For both libraries, cells with the 10% highest and lowest FSC

values were collected, as well as 70% of the population located around the median FSC value. For the

D39V P$_{lac}$-*dcas9 hlpA-gfp ftsZ-mCherry* (VL3117) library, the same fractions for GFP and mCherry values

were also sorted. Flow rates and dilutions were adjusted to keep the efficiency of sorting as high as

possible (and certainly above 85%) while not exceeding a sorting time of 60 min per sample. For every

fraction, 1.5×10^6 cells were collected and 6 different biological repeats were performed.

Cells were collected into 2 ml tubes, centrifuged at 18000 g for 5 min and pellets were stored at -20°C.

Cells were lysed by dissolving pellets in 10 μl H$_2$O + 0.025% DOC + 0.05% SDS and incubating 20 min at

37°C, followed by 5 min incubation at 80°C. After samples were allowed to cool off, a colony PCR was

performed using primers that contain index and adapter sequences necessary for Illumina sequencing. 10

μl of the lysed cell mixture was added to the PCR reaction as input DNA and 30 PCR cycles were performed.

The amplicons were purified from a 2% agarose gel and Illumina sequenced on a MiniSeq according to

manufacturer's instructions. Sequencing was performed with a custom sequencing protocol[23].

Data analysis was performed as described previously[23]. The sgRNA sequences were recovered from the

resulting reads using Trimmomatic[87] and mapped onto the *S. pneumoniae* D39V genome using Bowtie2[88].

The mapped sgRNAs were counted using featureCounts[89] and the DESeq2 R package was used to define

enrichments and associated p values for every sgRNA[90]. For the initial sCRilecs-seq screen, we defined

significant hits as sgRNAs from the fraction with the highest or lowest FSC/GFP/mCherry values with an

adjusted p value < 0.1 and Log$_2$FC > 1 compared to counts from the corresponding control population. For

the second sCRilecs-seq screen performed with a Δ*mvaS-mvaA* mutant limits were set at an adjusted p

value < 0.05 and Log$_2$FC > 1.

786   **Flow cytometry**

Flow cytometry experiments were performed using NovoCyte 2100YB (ACEA Biosciences) flow cytometer

equipped with violet, blue and red lasers. To validate sCRilecs-seq hits, strains were grown for 3.5h in C+Y

medium with or without 1 mM IPTG. Cultures were diluted in PBS and cell chains were disrupted as

described above. Next, cells were incubated for 30 min at room temperature to mimic the average waiting

time in the 60 min sorting step before FSC was measured. Cell gatings were chosen based on FSC, SSC and

GFP values, as was done during sorting.

794   **Gene ontology enrichment analyses**

Gene ontology enrichment analyses were performed using the online Gene Ontology Resource platform

that is coupled to the PANTHER classification system analysis tool[45-47]. Validated significant sgRNA hits

were translated into the spr identifiers that correspond to the targeted genes, which were used as the

input gene set that was compared to the *S. pneumoniae* reference list provided by the platform. A

PANTHER overrepresentation analysis was performed to identify biological processes that are

overrepresented as defined by a Fisher's exact test using FDR-corrected p values. The significance cut-off

was set at adjusted p value < 0.05.

OD growth curves

Cultures were grown until OD_{595nm} 0.1 in C+Y medium under non-limiting conditions

(deletion/complementation strains were grown with the appropriate amount of IPTG, aTc or mevalonic

acid, as listed in "Bacterial strains and growth conditions"). Cultures were diluted 100x into C+Y medium

supplemented or not with IPTG, aTc or mevalonic acid and with or without the addition of antibacterial

compounds, as indicated in the text. 300 μl of cell suspension was transferred into 96-well plates. When

growth in the presence of BAL fluid was tested, cultures were diluted 100x into C+Y medium

supplemented with ¼ BAL fluid. 200 μl of cell suspension was transferred into 96-well plates. In both cases,

addition of compounds such as amoxicillin or clomiphene occurred at this stage by diluting stock

concentrations into the wells of the 96-well plates (dilutions were chosen to always keep the DMSO

concentrations in the wells ≤ 1%). Growth was monitored by measuring $OD_{595\ nm}$ every 10 min using a

TECAN Infinite F200 Pro. OD_{595nm} values were normalized so that the lowest value measured during the

first hour of growth was 0.001, the initial OD_{595nm} value of the inoculum. In case the area under the curve

needed to be calculated, values were log-transformed before this parameter was determined using

GraphPad Prism 9.

Minimal Inhibitory Concentration (MIC) measurements

MIC values were determined by constructing growth curves in C+Y medium (see above). The MIC was

taken to be the concentration of the tested compound where the maximum OD_{595nm} value obtained was

less than 10% of the maximal OD_{595nm} value obtained in the absence of the compound.

Phase contrast and fluorescence microscopy

For the determination of cell morphology and FtsZ localization, cultures were grown until $OD_{595\ nm}$ 0.1 in

C+Y medium under non-limiting conditions (deletion/complementation strains were grown with the

appropriate amount of IPTG or mevalonic acid, as listed in "Bacterial strains and growth conditions").

Cultures were then diluted 100x into C+Y medium supplemented or not with IPTG or mevalonic acid and

grown until the $OD_{595\,nm}$ of the wild-type strain reached 0.2. At this point, cultures were diluted to $OD_{595\,nm}$

of 0.1 and incubated for 45 min at 30°C in order to slow down growth prior to imaging. 1 ml of cell

suspension was spun down (10000 g for 2 min) and pellets were dissolved in 40 µl PBS. Cells were kept on

ice prior to imaging.

To investigate peptidoglycan production using FDAAs, cultures were grown until $OD_{595\,nm}$ 0.1 in C+Y

medium under non-limiting conditions (deletion/complementation strains were grown with the

appropriate amount of IPTG, as listed in "Bacterial strains and growth conditions"). Cultures were then

diluted 100x into C+Y medium without IPTG and grown until the $OD_{595\,nm}$ of the wild-type strain reached

0.1. At this point, 250 µM sBADA was added and cultures were incubated for 15 min at 37°C. Cells were

washed 3 times with cold PBS (centrifuge at 10000 g for 1 min at 4°C) and pellets were dissolved in C+Y

medium containing 250 µM RADA. After incubating cells for 15 min at 37°C, the three wash steps were

repeated. Pellets were dissolved in PBS and kept on ice prior to imaging.

Cells were imaged by placing 0.4 µl of cell suspension on pads made of PBS containing 1% agarose. Imaging

was performed using a Leica DMi8 microscope with a sCMOS DFC9000 (Leica) camera and a SOLA light

engine (Lumencor). Phase contrast images were acquired using transmission light with 100 ms exposure

time. Leica DMi8 filter sets were used as follows: mCherry (Chroma 49017, Ex: 560/40 nm, BS: LP 590 nm,

Em: LP 590 nm) with exposure time 700 msec, sBADA (Ex: 470/40 nm Chroma ET470/40x, BS: LP 498 Leica

11536022, Em: 520/40 nm Chroma ET520/40m) with exposure time 200 msec, and RADA (Chroma 49017,

Ex: 560/40 nm, BS: LP 590 nm, Em: LP 590 nm) with exposure time 100 msec. Images were processed

using ImageJ and deconvolution was performed using Huygens software (Scientific Volume Imaging) with

standard settings using 15 iterations for FtsZ-mCherry and sBADA and 25 iterations for RADA.

Quantification of cell length and other properties was done using MicrobeJ[91] and BactMAP[92].

For the analysis and statistical comparison of cell size across different mutants and conditions (Figure 1C;

Figure 3D, F and H; Figure S2A-C; Figure 5J; Figure 6A), the following approach was used[93]. For every

mutant and/or condition, cell size was recorded for at least 3 biologically independent repeats. The one

exception is the depletion of *mraY* at time point 4.5h where only 2 independent repeats could be analyzed.

At this late time point, the culture is most often taken over by suppressor mutants with normal

morphology. Despite many attempts, we failed to obtain a third repeat in which the *mraY* depletion

phenotype was still apparent. For every repeat, at least 100 cells were recorded unless mentioned

otherwise and their average cell length was determined. These average cell lengths were used to calculate

the mean and SEM values that are shown in the main figures. The average cell lengths of each repeat were

also used to determine any statistically significant differences using a Wilcox test with FDR corrected p

values.

Transmission electron microscopy

Cultures were grown until $OD_{595\ nm}$ 0.1 in C+Y medium under non-limiting conditions

(deletion/complementation strains were grown with the appropriate amount of IPTG, as listed in

"Bacterial strains and growth conditions"). Cultures were then diluted 100x into C+Y medium without IPTG

and grown until the $OD_{595\ nm}$ of the wild-type strain reached 0.2. At this point, cultures were diluted to

$OD_{595\ nm}$ of 0.1 and incubated for 45 min at 30°C in order to slow down growth prior to imaging. 4 ml of

cell suspension was spun down at 10000 g and pellets were fixed in glutaraldehyde solution 2.5% (EMS)

and in osmium tetroxide 1% (EMS) with 1.5% of potassium ferrocyanide (Sigma) in phosphate buffer (PB

0.1 M [pH 7.4]) for 1h at room temperature. Samples were washed twice with H_2O and pellets were

embedded in agarose 2% (Sigma) in water, dehydrated in acetone solution (Sigma) at graded

concentrations (30% - 40 min; 70% - 40 min; 100% - 2x1h). This was followed by infiltration in Epon resin

(EMS) at graded concentrations (Epon 33% in acetone - 2h; Epon 66% in acetone - 4h; Epon 100% - 2x8h)

and finally polymerized for 48h at 60°C. Ultrathin sections of 50 nm thick were cut using a Leica Ultracut

(Leica Mikrosysteme GmbH), picked up on a copper slot grid 2x1mm (EMS) coated with a polystyrene film

(Sigma). Sections were post-stained with uranyl acetate (Sigma) 4% in H_2O for 10 min, rinsed several times

with H_2O followed by Reynolds lead citrate in H_2O (Sigma) for 10 min and rinsed several times with H_2O.

Micrographs were taken with a transmission electron microscope FEI CM100 (FEI) at an acceleration

voltage of 80 kV with a TVIPS TemCamF416 digital camera (TVIPS GmbH).

***In vivo* experiments using a murine pneumonia disease model**

Male C57BL/6JRj mice (8 weeks old) (Janvier Laboratories, Saint Berthevin, France) were maintained in

individually ventilated cages and were handled in a vertical laminar flow cabinet (class II A2, ESCO,

Hatboro, PA). All experiments complied with national, institutional and European regulations and ethical

guidelines, were approved by our Institutional Animal Care and Use guidelines (D59-350009, Institut

Pasteur de Lille; Protocol APAFIS#16966 201805311410769_v3) and were conducted by qualified,

accredited personnel.

Mice were anesthetized by intraperitoneal injection of 1.25 mg (50 mg/kg) ketamine plus 0.25 mg (10

mg/kg) xylazine in 200 µl of PBS. Mice were infected intranasally with 30 µl of PBS containing 50 plaque-

forming units (PFUs) of the pathogenic murine-adapted H3N2 influenza A virus strain Scotland/20/74[76].

Seven days later, mice were inoculated intranasally with 10^5 CFU of *S. pneumoniae* strain 19F in 30 µl of

PBS. Mice were treated intragastrically with 5 mg of clomiphene (Clomid, Sanofi-Aventis, France) in 200

µl of water and or 1 mg of amoxicillin (Clamoxyl for injection, GlaxoSmithKline) in 200 µl of water at 8 and

12 hours post-infection, respectively. Mice were sacrificed 24 hours post-infection by intraperitoneal

injection of 5.47 mg of sodium pentobarbital in 100 µl PBS (Euthasol, Virbac, France). Bronchoalveolar

lavage fluids (BAL) were sampled after intratracheal injection of 1 mL of PBS and centrifugation at 1400

rpm for 10 min. Lungs and spleen were sampled to determine the bacterial load. Tissues were

homogenized with an UltraTurrax homogenizer (IKA-Werke, Staufen, Germany) and serial dilutions were

plated on blood agar plates and incubated at 37°C. Viable counts were determined 24h later. Statistical

significance between groups was calculated by the Kruskall-Wallis test (One-Way ANOVA). Analyses were

performed with Prism software (version 9, GraphPad Software, La Jolla, CA).

Acknowledgements

We thank Mark van der Linden, German National Reference Center for Streptococci for providing us with

the *S. pneumoniae* clinical isolates of Spain-23F and 11A strains and Frédéric Wallet, Regional Reference

Center for pneumococci, Lille, for providing the *S. pneumoniae* strain 19F. We are thankful to the electron

microscopy facility of the university of Lausanne for their assistance in obtaining TEM images. We thank

Emanuele Cattani for technical assistance, construction of strains VL3710, VL3711 and VL3712, and

microscopy data used in Figure S2B-C. Work in the Veening lab is supported by the Swiss National Science

Foundation (SNSF) (project grants 310030_200792 and 310030_192517), SNSF JPIAMR grant

(40AR40_185533), SNSF NCCR 'AntiResist' (51NF40_180541) and ERC consolidator grant 771534-

PneumoCaTChER. LD is the recipient of a Marie Skłodowska Curie Actions – Individual Fellowship (MSCA-

IF, proposal number 837923). JCS, MB, and CC received funding from Région Hauts de France STaRS, and

the European Union's Horizon 2020 research and innovation programme under grant agreement No

847786.

References

1 Weiser, J. N., Ferreira, D. M. & Paton, J. C. *Streptococcus pneumoniae*: transmission, colonization
and invasion. *Nat Rev Microbiol* **16**, 355-367, doi:10.1038/s41579-018-0001-8 (2018).
2 O'Brien, K. L. *et al.* Burden of disease caused by *Streptococcus pneumoniae* in children younger
than 5 years: global estimates. *Lancet* **374**, 893-902, doi:10.1016/S0140-6736(09)61204-6
(2009).
3 Bogaert, D., De Groot, R. & Hermans, P. W. *Streptococcus pneumoniae* colonisation: the key to
pneumococcal disease. *Lancet Infect Dis* **4**, 144-154, doi:10.1016/S1473-3099(04)00938-7
(2004).
4 Jensen, A., Valdorsson, O., Frimodt-Moller, N., Hollingshead, S. & Kilian, M. Commensal
streptococci serve as a reservoir for beta-lactam resistance genes in *Streptococcus pneumoniae*.
Antimicrob Agents Chemother **59**, 3529-3540, doi:10.1128/AAC.00429-15 (2015).
5 Camara, J. *et al.* Evolution of the beta-lactam-resistant *Streptococcus pneumoniae* PMEN3 clone
over a 30 year period in Barcelona, Spain. *J Antimicrob Chemother* **73**, 2941-2951,
doi:10.1093/jac/dky305 (2018).
6 Schrag, S. J. *et al.* Emergence of *Streptococcus pneumoniae* with very-high-level resistance to
penicillin. *Antimicrob Agents Chemother* **48**, 3016-3023, doi:10.1128/AAC.48.8.3016-3023.2004
(2004).
7 Hakenbeck, R., Bruckner, R., Denapaite, D. & Maurer, P. Molecular mechanisms of beta-lactam
resistance in *Streptococcus pneumoniae*. *Future Microbiol* **7**, 395-410, doi:10.2217/fmb.12.2
(2012).
8 Kaur, R., Casey, J. R. & Pichichero, M. E. Emerging *Streptococcus pneumoniae* strains colonizing
the nasopharynx in children after 13-valent pneumococcal conjugate vaccination in comparison
to the 7-valent era, 2006-2015. *Pediatr Infect Dis J* **35**, 901-906,
doi:10.1097/INF.0000000000001206 (2016).
9 Brueggemann, A. B., Pai, R., Crook, D. W. & Beall, B. Vaccine escape recombinants emerge after
pneumococcal vaccination in the United States. *PLoS Pathog* **3**, e168,
doi:10.1371/journal.ppat.0030168 (2007).
10 Hanage, W. P., Fraser, C., Tang, J., Connor, T. R. & Corander, J. Hyper-recombination, diversity,
and antibiotic resistance in pneumococcus. *Science* **324**, 1454-1457,
doi:10.1126/science.1171908 (2009).
11 Domenech, A. *et al.* Proton motive force disruptors block bacterial competence and horizontal
gene transfer. *Cell Host Microbe* **27**, 544-555 e543, doi:10.1016/j.chom.2020.02.002 (2020).
12 Chi, F., Nolte, O., Bergmann, C., Ip, M. & Hakenbeck, R. Crossing the barrier: evolution and
spread of a major class of mosaic *pbp2x* in *Streptococcus pneumoniae*, *S. mitis* and *S. oralis*. *Int J
Med Microbiol* **297**, 503-512, doi:10.1016/j.ijmm.2007.02.009 (2007).
13 Prieto, J., Calvo, A. & Gomez-Lus, M. L. Antimicrobial resistance: a class effect? *J Antimicrob
Chemother* **50 Suppl S2**, 7-12, doi:10.1093/jac/dkf508 (2002).
14 Baquero, F., Baquero-Artigao, G., Canton, R. & Garcia-Rey, C. Antibiotic consumption and
resistance selection in *Streptococcus pneumoniae*. *J Antimicrob Chemother* **50 Suppl S2**, 27-37,
doi:10.1093/jac/dkf504 (2002).
15 Jacobs, M. R. Building in efficacy: developing solutions to combat drug-resistant *S. pneumoniae*.
Clin Microbiol Infect **10 Suppl 2**, 18-27, doi:10.1111/j.1470-9465.2004.00862.x (2004).
16 Bradley, J. S. *et al.* The management of community-acquired pneumonia in infants and children
older than 3 months of age: clinical practice guidelines by the Pediatric Infectious Diseases
Society and the Infectious Diseases Society of America. *Clin Infect Dis* **53**, e25-76,
doi:10.1093/cid/cir531 (2011).

17   Thiem, U., Heppner, H. J. & Pientka, L. Elderly patients with community-acquired pneumonia:
optimal treatment strategies. *Drugs Aging* **28**, 519-537, doi:10.2165/11591980-000000000-
00000 (2011).
18   Qi, L. S. *et al.* Repurposing CRISPR as an RNA-guided platform for sequence-specific control of
gene expression. *Cell* **152**, 1173-1183, doi:10.1016/j.cell.2013.02.022 (2013).
19   Peters, J. M. *et al.* A comprehensive, CRISPR-based functional analysis of essential genes in
bacteria. *Cell* **165**, 1493-1506, doi:10.1016/j.cell.2016.05.003 (2016).
20   Bikard, D. *et al.* Programmable repression and activation of bacterial gene expression using an
engineered CRISPR-Cas system. *Nucleic Acids Res* **41**, 7429-7437, doi:10.1093/nar/gkt520
(2013).
21   Liu, X. *et al.* High-throughput CRISPRi phenotyping identifies new essential genes in
Streptococcus pneumoniae. *Mol Syst Biol* **13**, 931, doi:10.15252/msb.20167449 (2017).
22   Cui, L. *et al.* A CRISPRi screen in *E. coli* reveals sequence-specific toxicity of dCas9. *Nat Commun*
9, 1912, doi:10.1038/s41467-018-04209-5 (2018).
23   Liu, X. *et al.* Exploration of bacterial bottlenecks and *Streptococcus pneumoniae* pathogenesis by
CRISPRi-Seq. *Cell Host Microbe* **29**, 107-120 e106, doi:10.1016/j.chom.2020.10.001 (2021).
24   de Wet, T. J., Gobe, I., Mhlanga, M. M. & Warner, D. F. CRISPRi-Seq for the identification and
characterisation of essential Mycobacterial genes and transcriptional units. *BioRxiv*, 358275,
doi:https://doi.org/10.1101/358275 (2018).
25   Lee, H. H. *et al.* Functional genomics of the rapidly replicating bacterium *Vibrio natriegens* by
CRISPRi. *Nat Microbiol* **4**, 1105-1113, doi:10.1038/s41564-019-0423-8 (2019).
26   Rousset, F. *et al.* Genome-wide CRISPR-dCas9 screens in *E. coli* identify essential genes and
phage host factors. *PLoS Genet* **14**, e1007749, doi:10.1371/journal.pgen.1007749 (2018).
27   Wang, T. *et al.* Pooled CRISPR interference screening enables genome-scale functional genomics
study in bacteria with superior performance. *Nat Commun* **9**, 2475, doi:10.1038/s41467-018-
04899-x (2018).
28   Park, J. S. *et al.* A FACS-based genome-wide CRISPR screen reveals a requirement for COPI in
Chlamydia trachomatis invasion. *iScience* **11**, 71-84, doi:10.1016/j.isci.2018.12.011 (2019).
29   Smith, L. M., Jackson, S. A., Gardner, P. P. & Fineran, P. C. SorTn-seq: a high-throughput
functional genomics approach to discovering regulators of bacterial gene expression. *Nat*
Protoc, doi:10.1038/s41596-021-00582-6 (2021).
30   Boronat, A. & Rodriguez-Concepcion, M. Terpenoid biosynthesis in prokaryotes. *Adv Biochem*
Eng Biotechnol **148**, 3-18, doi:10.1007/10_2014_285 (2015).
31   Heuston, S., Begley, M., Gahan, C. G. M. & Hill, C. Isoprenoid biosynthesis in bacterial pathogens.
Microbiology (Reading) **158**, 1389-1401, doi:10.1099/mic.0.051599-0 (2012).
32   Bouhss, A., Trunkfield, A. E., Bugg, T. D. & Mengin-Lecreulx, D. The biosynthesis of peptidoglycan
lipid-linked intermediates. *FEMS Microbiol Rev* **32**, 208-233, doi:10.1111/j.1574-
6976.2007.00089.x (2008).
33   Farha, M. A. *et al.* Antagonism screen for inhibitors of bacterial cell wall biogenesis uncovers an
inhibitor of undecaprenyl diphosphate synthase. *Proc Natl Acad Sci U S A* **112**, 11048-11053,
doi:10.1073/pnas.1511751112 (2015).
34   Tsui, H. T. *et al.* Pbp2x localizes separately from Pbp2b and other peptidoglycan synthesis
proteins during later stages of cell division of *Streptococcus pneumoniae* D39. *Mol Microbiol* **94**,
21-40, doi:10.1111/mmi.12745 (2014).
35   Sorg, R. A. & Veening, J. W. Microscale insights into pneumococcal antibiotic mutant selection
windows. *Nat Commun* **6**, 8773, doi:10.1038/ncomms9773 (2015).
36   Brochado, A. R. *et al.* Species-specific activity of antibacterial drug combinations. *Nature* **559**,
259-263, doi:10.1038/s41586-018-0278-9 (2018).

37　Shi, H. *et al.* Deep phenotypic mapping of bacterial cytoskeletal mutants reveals physiological
robustness to cell size. *Curr Biol* **27**, 3419-3429 e3414, doi:10.1016/j.cub.2017.09.065 (2017).
38　Kjos, M. *et al.* Bright fluorescent *Streptococcus pneumoniae* for live-cell imaging of host-
pathogen interactions. *J Bacteriol* **197**, 807-818, doi:10.1128/JB.02221-14 (2015).
39　Beilharz, K., van Raaphorst, R., Kjos, M. & Veening, J. W. Red fluorescent proteins for gene
expression and protein localization studies in *Streptococcus pneumoniae* and efficient
transformation with DNA assembled via the Gibson Assembly method. *Appl Environ Microbiol*
81, 7244-7252, doi:10.1128/AEM.02033-15 (2015).
40　Massidda, O., Novakova, L. & Vollmer, W. From models to pathogens: how much have we
learned about *Streptococcus pneumoniae* cell division? *Environ Microbiol* **15**, 3133-3157,
doi:10.1111/1462-2920.12189 (2013).
41　Rued, B. E. *et al.* Suppression and synthetic-lethal genetic relationships of Δ*gpsB* mutations
indicate that GpsB mediates protein phosphorylation and penicillin-binding protein interactions
in *Streptococcus pneumoniae* D39. *Mol Microbiol* **103**, 931-957, doi:10.1111/mmi.13613 (2017).
42　Land, A. D. *et al.* Requirement of essential Pbp2x and GpsB for septal ring closure in
Streptococcus pneumoniae D39. *Mol Microbiol* **90**, 939-955, doi:10.1111/mmi.12408 (2013).
43　Vollmer, W., Massidda, O. & Tomasz, A. The cell wall of *Streptococcus pneumoniae*. *Microbiol*
Spectr **7**, GPP3-0018-2018, doi:10.1128/microbiolspec.GPP3-0018-2018 (2019).
44　Brown, S., Santa Maria, J. P., Jr. & Walker, S. Wall teichoic acids of Gram-positive bacteria. *Annu*
Rev Microbiol **67**, 313-336, doi:10.1146/annurev-micro-092412-155620 (2013).
45　Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. *Nat Genet* **25**, 25-29,
doi:10.1038/75556 (2000).
46　Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P. D. PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic*
Acids Res **47**, 419-426, doi:10.1093/nar/gky1038 (2019).
47　Consortium, T. G. O. The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic Acids*
Res **47**, 330-338, doi:10.1093/nar/gky1055 (2019).
48　Wilding, E. I. *et al.* Identification, evolution, and essentiality of the mevalonate pathway for
isopentenyl diphosphate biosynthesis in Gram-positive cocci. *J Bacteriol* **182**, 4319-4327,
doi:10.1128/jb.182.15.4319-4327.2000 (2000).
49　Slager, J., Aprianto, R. & Veening, J. W. Deep genome annotation of the opportunistic human
pathogen *Streptococcus pneumoniae* D39. *Nucleic Acids Res* **46**, 9971-9989,
doi:10.1093/nar/gky725 (2018).
50　Balibar, C. J., Shen, X. Y. & Tao, J. S. The mevalonate pathway of *Staphylococcus aureus*. *J*
Bacteriol **191**, 851-861, doi:10.1128/Jb.01357-08 (2009).
51　Yu, W. Q. *et al.* The mevalonate auxotrophic mutant of *Staphylococcus aureus* can adapt to
mevalonate depletion. *Antimicrob Agents Ch* **57**, 5710-5713, doi:10.1128/Aac.00726-13 (2013).
52　Reichert, S. *et al.* Genetic adaptation of a mevalonate pathway deficient mutant in
Staphylococcus aureus. *Front Microbiol* **9**, 1539, doi:ARTN 1539
10.3389/fmicb.2018.01539 (2018).
53　Keller, L. E., Rueff, A. S., Kurushima, J. & Veening, J. W. Three new integration vectors and
fluorescent proteins for use in the opportunistic human pathogen *Streptococcus pneumoniae*.
Genes (Basel) **10**, 394, doi:10.3390/genes10050394 (2019).
54　Kuru, E., Tekkam, S., Hall, E., Brun, Y. V. & Van Nieuwenhze, M. S. Synthesis of fluorescent D-
amino acids and their use for probing peptidoglycan synthesis and bacterial growth *in situ*. *Nat*
Protoc **10**, 33-52, doi:10.1038/nprot.2014.197 (2015).

55   Wheeler, R., Mesnage, S., Boneca, I. G., Hobbs, J. K. & Foster, S. J. Super-resolution microscopy
reveals cell wall dynamics and peptidoglycan architecture in ovococcal bacteria. *Mol Microbiol*
82, 1096-1109, doi:10.1111/j.1365-2958.2011.07871.x (2011).
56   Yao, Z., Kahne, D. & Kishony, R. Distinct single-cell morphological dynamics under beta-lactam
antibiotics. *Mol Cell* **48**, 705-712, doi:10.1016/j.molcel.2012.09.016 (2012).
57   Yother, J. Capsules of *Streptococcus pneumoniae* and other bacteria: paradigms for
polysaccharide biosynthesis and regulation. *Annu Rev Microbiol* **65**, 563-581,
doi:10.1146/annurev.micro.62.081307.162944 (2011).
58   Barendt, S. M. *et al.* Influences of capsule on cell shape and chain formation of wild-type and
pcsB mutants of serotype 2 *Streptococcus pneumoniae*. *J Bacteriol* **191**, 3024-3040,
doi:10.1128/JB.01505-08 (2009).
59   Denapaite, D., Bruckner, R., Hakenbeck, R. & Vollmer, W. Biosynthesis of teichoic acids in
Streptococcus pneumoniae and closely related species: lessons from genomes. *Microb Drug*
Resist **18**, 344-358, doi:10.1089/mdr.2012.0026 (2012).
60   Beilharz, K. *et al.* Control of cell division in *Streptococcus pneumoniae* by the conserved Ser/Thr
protein kinase StkP. *Proc Natl Acad Sci U S A* **109**, E905-913, doi:10.1073/pnas.1119172109
(2012).
61   Fleurie, A. *et al.* Mutational dissection of the S/T-kinase StkP reveals crucial roles in cell division
of *Streptococcus pneumoniae*. *Mol Microbiol* **83**, 746-758, doi:10.1111/j.1365-
2958.2011.07962.x (2012).
62   Fleurie, A. *et al.* Interplay of the serine/threonine-kinase StkP and the paralogs DivIVA and GpsB
in pneumococcal cell elongation and division. *PLoS Genet* **10**, e1004275,
doi:10.1371/journal.pgen.1004275 (2014).
63   Novakova, L. *et al.* Identification of multiple substrates of the StkP Ser/Thr protein kinase in
Streptococcus pneumoniae. *J Bacteriol* **192**, 3629-3638, doi:10.1128/JB.01564-09 (2010).
64   Cho, H. *et al.* Bacterial cell wall biogenesis is mediated by SEDS and PBP polymerase families
functioning semi-autonomously. *Nat Microbiol* **1**, 16172, doi:10.1038/nmicrobiol.2016.172
(2016).
65   Meeske, A. J. *et al.* SEDS proteins are a widespread family of bacterial cell wall polymerases.
Nature **537**, 634-638, doi:10.1038/nature19331 (2016).
66   Taguchi, A. *et al.* FtsW is a peptidoglycan polymerase that is functional only in complex with its
cognate penicillin-binding protein. *Nat Microbiol* **4**, 587-594, doi:10.1038/s41564-018-0345-x
(2019).
67   Tsui, H. C. *et al.* Suppression of a deletion mutation in the gene encoding essential PBP2b reveals
a new lytic transglycosylase involved in peripheral peptidoglycan synthesis in *Streptococcus*
pneumoniae D39. *Mol Microbiol* **100**, 1039-1065, doi:10.1111/mmi.13366 (2016).
68   Perez, A. J. *et al.* Movement dynamics of divisome proteins and PBP2x:FtsW in cells of
Streptococcus pneumoniae. *Proc Natl Acad Sci U S A* **116**, 3211-3220,
doi:10.1073/pnas.1816018116 (2019).
69   Berg, K. H., Stamsas, G. A., Straume, D. & Havarstein, L. S. Effects of low PBP2b levels on cell
morphology and peptidoglycan composition in *Streptococcus pneumoniae* R6. *J Bacteriol* **195**,
4342-4354, doi:10.1128/JB.00184-13 (2013).
70   Kocaoglu, O., Tsui, H. C., Winkler, M. E. & Carlson, E. E. Profiling of beta-lactam selectivity for
penicillin-binding proteins in *Streptococcus pneumoniae* D39. *Antimicrob Agents Chemother* **59**,
3548-3555, doi:10.1128/AAC.05142-14 (2015).
71   Masadeh, M., Mhaidat, N., Alzoubi, K., Al-Azzam, S. & Alnasser, Z. Antibacterial activity of
statins: a comparative study of atorvastatin, simvastatin, and rosuvastatin. *Ann Clin Microbiol*
Antimicrob **11**, 13, doi:10.1186/1476-0711-11-13 (2012).

72 Bergman, P. *et al.* Studies on the antibacterial effects of statins - *in vitro* and *in vivo*. *PLoS One* **6**,
e24394, doi:10.1371/journal.pone.0024394 (2011).
73 Kaneko, M., Togashi, N., Hamashima, H., Hirohara, M. & Inoue, Y. Effect of farnesol on
mevalonate pathway of *Staphylococcus aureus*. *J Antibiot (Tokyo)* **64**, 547-549,
doi:10.1038/ja.2011.49 (2011).
74 Imohl, M., Reinert, R. R. & van der Linden, M. Serotype-specific penicillin resistance of
Streptococcus pneumoniae in Germany from 1992 to 2008. *Int J Med Microbiol* **300**, 324-330,
doi:10.1016/j.ijmm.2009.11.004 (2010).
75 EUCAST. Breakpoint tables for interpretation of MICs and zone diameters, version 11.0.
http://www.eucast.org/clinical_breakpoints/ (2021).
76 Matarazzo, L. *et al.* Therapeutic synergy between antibiotics and pulmonary Toll-like receptor 5
stimulation in antibiotic-sensitive or -resistant pneumonia. *Front Immunol* **10**, 723,
doi:10.3389/fimmu.2019.00723 (2019).
77 Dickey, R. P. & Holtkamp, D. E. Development, pharmacology and clinical experience with
clomiphene citrate. *Hum Reprod Update* **2**, 483-506, doi:10.1093/humupd/2.6.483 (1996).
78 Huppert, L. C. Induction of ovulation with clomiphene citrate. *Fertil Steril* **31**, 1-8,
doi:10.1016/s0015-0282(16)43749-0 (1979).
79 Nasseri, S. & Ledger, W. L. Clomiphene citrate in the twenty-first century. *Hum Fertil (Camb)* **4**,
145-151, doi:10.1080/1464727012000199212 (2001).
80 Cleverley, R. M. *et al.* The cell cycle regulator GpsB functions as cytosolic adaptor for multiple
cell wall enzymes. *Nat Commun* **10**, 261, doi:10.1038/s41467-018-08056-2 (2019).
81 Zucchini, L. *et al.* PASTA repeats of the protein kinase StkP interconnect cell constriction and
separation of *Streptococcus pneumoniae*. *Nat Microbiol* **3**, 197-209, doi:10.1038/s41564-017-
0069-3 (2018).
82 Murdter, T. E. *et al.* Genetic polymorphism of cytochrome P450 2D6 determines oestrogen
receptor activity of the major infertility drug clomiphene via its active metabolites. *Hum Mol*
Genet **21**, 1145-1154, doi:10.1093/hmg/ddr543 (2012).
83 Synefiaridou, D. & Veening, J.-W. Harnessing CRISPR-Cas9 for genome editing in *Streptococcus*
pneumoniae D39V. *Appl Environ Microbiol* **87**, e02762-02720, doi:10.1101/2020.06.13.149682
%J bioRxiv (2020).
84 Aprianto, R., Slager, J., Holsappel, S. & Veening, J. W. High-resolution analysis of the
pneumococcal transcriptome under a wide range of infection-relevant conditions. *Nucleic Acids*
Res **46**, 9990-10006, doi:10.1093/nar/gky750 (2018).
85 Domenech, A., Slager, J. & Veening, J. W. Antibiotic-induced cell chaining triggers pneumococcal
competence by reshaping quorum sensing to autocrine-like signaling. *Cell Rep* **25**, 2390-2400
e2393, doi:10.1016/j.celrep.2018.11.007 (2018).
86 Moreno-Gamez, S. *et al.* Quorum sensing integrates environmental cues, cell density and cell
history to control bacterial competence. *Nat Commun* **8**, 854, doi:10.1038/s41467-017-00903-y
(2017).
87 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170 (2014).
88 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-
359, doi:10.1038/nmeth.1923 (2012).
89 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930,
doi:10.1093/bioinformatics/btt656 (2014).
90 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).

91   Ducret, A., Quardokus, E. M. & Brun, Y. V. MicrobeJ, a tool for high throughput bacterial cell
detection and quantitative analysis. *Nat Microbiol* **1**, 16077, doi:10.1038/nmicrobiol.2016.77
(2016).
92   van Raaphorst, R., Kjos, M. & Veening, J. W. BactMAP: An R package for integrating, analyzing
and visualizing bacterial microscopy data. *Mol Microbiol* **113**, 297-308, doi:10.1111/mmi.14417
(2020).
93   Lord, S. J., Velle, K. B., Mullins, R. D. & Fritz-Laylin, L. K. SuperPlots: Communicating
reproducibility and variability in cell biology. *J Cell Biol* **219**, e202001064,
doi:10.1083/jcb.202001064 (2020).
1164