Gene duplication and rate variation in the evolution of non-photosynthetic pathways in plastids

Alissa M. Williams[1,2,*], Olivia G. Carter[1], Evan S. Forsythe[1], Hannah K. Mendoza[1], Daniel B. Sloan[1]

[1]Department of Biology, Colorado State University, Fort Collins, Colorado 80523

[2]Program in Cell and Molecular Biology, Colorado State University, Fort Collins, Colorado 80523

*corresponding author: Alissa.Williams@colostate.edu

Abstract:

While the chloroplast (plastid) is known for its role in photosynthesis, it is also involved in many

other biosynthetic pathways essential for plant survival. As such, plastids contain an extensive

suite of enzymes required for non-photosynthetic processes. The evolution of the associated

genes has been especially dynamic in flowering plants (angiosperms), including examples of

gene duplication and extensive rate variation. We examined the role of ongoing gene duplication

in two key plastid enzymes, the acetyl-CoA carboxylase (ACCase) and the caseinolytic protease

(Clp), responsible for fatty acid biosynthesis and protein turnover, respectively. In plants, there

are two ACCase complexes—a homomeric version present in the cytosol and a heteromeric

version present in the plastid. Duplications of the nuclear-encoded homomeric ACCase gene and

retargeting to the plastid have been previously reported in multiple species. We find that these

retargeted copies of the homomeric ACCase gene exhibit elevated rates of sequence evolution,

consistent with neofunctionalization and/or relaxation of selection. The plastid Clp complex

catalytic core is composed of nine paralogous proteins that arose via ancient gene duplication in

the cyanobacterial/plastid lineage. We show that further gene duplication occurred more recently

in the nuclear-encoded core subunits of this complex, yielding additional paralogs in many

species of angiosperms. Moreover, in six of eight cases, subunits that have undergone recent

duplication display increased rates of sequence evolution relative to those that have remained

single copy. We also compared rate patterns between pairs of Clp core paralogs to gain insight

into post-duplication evolutionary routes. These results show that gene duplication and rate

variation continue to shape the plastid proteome.

Introduction:

The plastid is a dynamic proteomic environment in which key photosynthetic and non-photosynthetic biochemical reactions occur. Major non-photosynthetic functions of plastids include the reaction catalyzed by the acetyl-CoA carboxylase (ACCase) enzyme and protein degradation performed by the caseinolytic protease (Clp) complex (Caroca et al., 2021; Green, 2011; Konishi et al., 1996; Nishimura et al., 2017; Nishimura and van Wijk, 2015). Both of these functions are essential in plants and thus the genes involved are generally highly conserved; however, these genes have undergone rapid evolution in multiple angiosperm species (Barnard-Kubow et al., 2014; Erixon and Oxelman, 2008; Jansen et al., 2007; Park et al., 2017; Sloan et al., 2014, 2014; Wicke et al., 2011; Williams et al., 2019, 2015; Zhang et al., 2014). While many hypotheses about these patterns of accelerated evolution have been posited, the underlying evolutionary mechanisms, causes, and consequences remain largely unknown.

The ACCase enzyme catalyzes the first committed step of fatty acid biosynthesis, the carboxylation of acetyl-CoA to malonyl-CoA (Salie and Thelen, 2016; Sasaki and Nagano, 2004). This step requires four different enzyme domains—one biotin carboxylase, one biotin carboxyl carrier, and two (α and β) carboxyltransferases (Salie and Thelen, 2016; Sasaki and Nagano, 2004; Schulte et al., 1997). In plants, there are two forms of the ACCase enzyme. The homomeric version, present in the cytosol, is encoded by a single nuclear gene (Konishi et al., 1996; Konishi and Sasaki, 1994). The heteromeric version, present in the plastid, is encoded by five genes in *Arabidopsis thaliana*; each functional domain is represented by a single gene except for the biotin carboxyl carrier domain, which is encoded by two genes (Konishi et al., 1996; Konishi and Sasaki, 1994; Salie and Thelen, 2016). Four of these genes are in the nuclear genome while the fifth (*accD*) is in the plastid genome (Caroca et al., 2021; Sasaki and Nagano, 2004). In a few angiosperm lineages, including the Brassicaceae, Caryophyllaceae, Geraniaceae, and Poaceae, there have been duplications of the homomeric ACCase gene with subsequent retargeting of one copy to the plastid (Babiychuk et al., 2011; Konishi and Sasaki, 1994; Park et al., 2017; Parker et al., 2014; Rockenbach et al., 2016; Schulte et al., 1997).

The Clp complex is one of the most abundant stromal proteases and degrades a variety of targets

(Apitz et al., 2016; Bouchnak and van Wijk, 2021; Majeran et al., 2000; Montandon et al., 2019;

Nishimura et al., 2017; Nishimura and van Wijk, 2015; Welsch et al., 2018). This complex

consists of many types of subunits. Adapters bind proteins targeted for degradation and deliver

them to chaperones, which use ATP to unfold the targeted proteins into the proteolytic core of

the complex (Nishimura and van Wijk, 2015). The core consists of 14 subunits that are encoded

by nine different paralogous genes (Olinares et al., 2011a; Peltier et al., 2004; Sjögren et al.,

2006; Stanne et al., 2007). Eight of these genes reside in the nuclear genome (*CLPP3-6, CLPR1-*

4), while the ninth is encoded in the plastid genome (*clpP1*) (Nishimura et al., 2017; Olinares et

al., 2011b). The ClpP subunits contain a catalytically active Ser-His-Asp triad, whereas the ClpR

subunits do not (Nishimura and van Wijk, 2015; Porankiewicz et al., 1999). These nine paralogs

are the results of gene duplications throughout cyanobacterial and plastid evolution and are

shared by all land plants (Olinares et al., 2011a). Ongoing gene duplication of individual

subunits has been noted in a handful of angiosperm lineages (Rockenbach et al., 2016; Williams

et al., 2021, 2019).

Thus, the evolutionary trajectory of both of these essential plastid pathways is characterized by

gene duplication at both ancient and recent timescales. Gene duplication is common in land

plants, in part due to the frequency with which whole genome duplication (polyploidization)

occurs in this lineage (Clark and Donoghue, 2018; De Bodt et al., 2005; del Pozo and Ramirez-

Parra, 2015; Flagel and Wendel, 2009; Panchy et al., 2016; Wendel et al., 2018). Nearly all

species of land plants have polyploidization events in their evolutionary histories (Clark and

Donoghue, 2018; Leebens-Mack et al., 2019; Panchy et al., 2016). Angiosperms in particular

seem to have a propensity for whole genome duplication; the entire clade shares an ancient

polyploidization event and many lineages have undergone subsequent rounds of whole genome

duplication (Clark and Donoghue, 2018; Panchy et al., 2016; Renny-Byfield and Wendel, 2014;

Soltis et al., 2009). While every gene is initially affected by whole genome duplication, only 10-

30% of those duplicates are maintained in the genome longer-term (Hahn, 2009; Maere et al.,

2005; Paterson et al., 2006). Though polyploidy is likely a main contributor to gene duplication

in plants, other forms of gene duplication are also prevalent (Flagel and Wendel, 2009). For

instance, tandem duplication has been shown to be common in both *Arabidopsis thaliana* and

Oryza sativa, where tandemly arrayed gene clusters make up 15-20% of genic content.

Additionally, multiple studies have shown that transposon-mediated gene duplication is

prevalent in plants (Flagel and Wendel, 2009; Freeling et al., 2008; Rizzon et al., 2006; Wang et

al., 2006).

Gene duplication is an important evolutionary process and is thought to be a major source of

evolutionary novelty (Hahn, 2009; Ohno, 1970; Taylor and Raes, 2004; Zhang, 2003). The most

common evolutionary fate of paralogs is retention of one copy and pseudogenization and loss of

the other copy (Lynch and Conery, 2000; Zhang, 2003; Zhang et al., 2003). However, several

evolutionary mechanisms have been described in which retention of both gene duplicates is

favored. The increased gene-dosage advantage model describes a scenario in which increased

amount of gene product produced by the two identical gene copies is beneficial and thus both

copies retain ancestral function (Hahn, 2009; Ohno, 1970; Pegueroles et al., 2013; Zhang, 2003).

The neofunctionalization model posits that one paralog acquires new functions while the other

retains ancestral functionality (Hahn, 2009; Ohno, 1970; Pegueroles et al., 2013; Zhang, 2003).

In the subfunctionalization model, an ancestral function is split between the two duplicates

(Hahn, 2009; Ohno, 1970; Pegueroles et al., 2013; Zhang, 2003), in some cases creating the

possibility for each paralog to optimize a subset of the ancestral function in a process known as

escape from adaptive conflict (Des Marais and Rausher, 2008; Huang et al., 2015; Sikosek et al.,

2012).

To distinguish between these evolutionary fates, many studies have employed evolutionary rate

comparisons (Hahn, 2009; Pegueroles et al., 2013). These comparisons involve both paralogs as

well as their common ancestor (Pegueroles et al., 2013). Under both the gene-dosage advantage

and subfunctionalization models, gene duplicates are expected to evolve at approximately the

same rate as each other (Pegueroles et al., 2013). The difference in evolutionary rates predicted

by these two models is found in comparisons to the common ancestor; with a gene-dosage

advantage, the expectation is that the paralogs will evolve at the same rate as the common

ancestor, while with subfunctionalization, the expectation is that the paralogs will evolve at an

increased rate relative to the common ancestor (though this assumption has been challenged)

(Force et al., 1999; Hahn, 2009; He and Zhang, 2005; Lynch and Force, 2000; Pegueroles et al.,

2013; Zhang, 2003). By contrast, under the neofunctionalization model, asymmetry between

evolutionary rates of paralogs is expected, where one paralog retains the ancestral evolutionary

rate while the other experiences rate acceleration after being freed from selective constraints

(Hahn, 2009; Pegueroles et al., 2013; Zhang, 2003). The proportion of paralogs with asymmetric

rates of evolution has been estimated at anywhere from 5% to 65% in a variety of studies

(Conant and Wagner, 2003; Dermitzakis and Clark, 2001; Kondrashov et al., 2002; Panchin et

al., 2010; Pegueroles et al., 2013; Van de Peer et al., 2001). This wide range of estimates is likely

due to differences in study systems, definitions and identifications of paralogs, gene types, and

time since duplication. Despite the varying estimates of evolutionary rate asymmetry, it is clear

that paralogs evolve under a mixture of evolutionary regimes.

Here, we characterize recent gene duplication events and subsequent changes in evolutionary

rate in ACCase and Clp core subunits. We show that ACCase genes exhibit patterns of

duplication, retargeting, and accelerated protein evolution consistent with neofunctionalization

and/or relaxed selection. Additionally, we examine duplications of nuclear-encoded plastid Clp

core subunits and demonstrate that duplication leads to significant changes in the rate of

evolution in most cases but that patterns differ across Clp subunits, meaning multiple post-

duplication evolutionary routes are represented across pairs of paralogs. This work provides

additional insights into the interplay between gene duplication and evolutionary rate in the

molecular evolution of plastid proteins.

<u>Materials and Methods:</u>

Compilation and curation of *ACC* nucleotide sequences

Previous work identified duplications of the homomeric ACCase gene *ACC* and subsequent

retargeting of one copy to the plastid in the angiosperm families Poaceae (Konishi and Sasaki,

1994; Park et al., 2017; Rockenbach et al., 2016), Brassicaceae (Babiychuk et al., 2011; Park et

al., 2017; Parker et al., 2014; Schulte et al., 1997), Caryophyllaceae (Rockenbach et al., 2016),

and Geraniaceae (Park et al., 2017). *ACC* genes were obtained for multiple species in each of

these families. All cytosol-targeted *ACC* genes were designated *ACC1* while all plastid-targeted

ACC genes were designated *ACC2* per established conventions (Babiychuk et al., 2011; Sasaki

and Nagano, 2004); thus, sharing the same identifier does not necessarily indicate orthology

because of the multiple independent origins of plastid-targeted *ACC2* genes. *Amborella*

trichopoda, which has a single *ACC* gene that we designated *ACC1*, was used as an outgroup.

Trimmed *ACC1* and *ACC2* coding sequences (CDSs) were obtained from Rockenbach et al.

(2016) for *Amborella trichopoda, Arabidopsis thaliana, Agrostemma githago*, *Silene noctiflora*,

Silene paradoxa, and *Triticum aestivum*. The trimming in Rockenbach et al (2016) was codon-

guided and included removal of the target peptide. *ACC1* and *ACC2* CDSs from the following

species were compiled using gene identifiers from Table S4 in Park et al. (2017): Geraniaceae:

California macrophylla, Erodium texanum, Geranium incanum, Geranium maderense,

Geranium phaeum, Monsonia emarginata, Pelargonium cotyledonis; Brassicaceae: *Capsella*

rubella; Poaceae: *Oryza sativa*, *Sorghum bicolor*. Duplications of *ACC* were additionally

identified in two Poaceae species—*Aegilops tauschii* and *Zea mays*—by performing BLAST

searches against these organisms on NCBI and Phytozome v13, respectively (Camacho et al.,

2009; Goodstein et al., 2012).

All *ACC1* and *ACC2* sequences were included in a single file and aligned using the MAFFT *einsi*

option (Katoh and Standley, 2013) in codon space using the *align_fasta_with_mafft_codon*

subroutine in the sloan.pm Perl module (https://github.com/dbsloan/perl_modules). 5′ trimming

was conducted according to the trimming performed in Rockenbach et al. (2016). Additional

trimming of poorly aligned regions was performed manually in a codon-based manner.

Compilation and curation of Clp core subunit amino acid and nucleotide sequences

To identify Clp core subunit amino acid sequences, a custom Python script

(https://github.com/alissawilliams/Gene_duplication_ACCase_Clp/scripts /local_blast5.py) was

used to reciprocally blast (blastp v2.2.29) *Arabidopsis thaliana* amino acid sequences against

predicted protein sequences from each of 22 other angiosperm species in the dataset. These 22

species were the same set used in Williams et al. (2019) with the exclusion of *Silene latifolia* and

Silene noctiflora, since Clp core subunit duplications have been previously studied in *Sileneae*

(Rockenbach et al., 2016; Williams et al., 2021, 2019). This sampling was chosen to represent

both the diversity of angiosperms and the range of rate variation in Clp complex evolution

(Williams et al., 2019; see Table S3).

Compiled amino acid sequences for each subunit were aligned using the *einsi* option in MAFFT

v7.222 (Katoh and Standley, 2013) and trimmed using GBLOCKS v0.91b (Castresana, 2000)

with parameter *-b1* set to the default value of *-b2* and parameter *-b5* set to *h*. All alignments were

examined manually to confirm homology. Sequences were also screened to prevent inclusion of

multiple splice variants from a single gene. In cases where genomic data were used, only one

transcript per gene was used. In cases where transcriptomic data were used, sequences were

eliminated when alternative splicing was obvious (i.e. inclusion of an intron where the other

sequence had a gap or variation only in one short piece of the transcript at either end). Catalytic

site status and length were determined using the amino acid sequence data.

Nucleotide sequences for each identified Clp core subunit protein sequence were compiled from

the corresponding CDS or transcript sequence file. For non-CDS sequences, ORFfinder

(Wheeler et al., 2003) was used to identify the coding sequence. Compiled CDS sequences for

each subunit were aligned with the MAFFT *einsi* option (Katoh and Standley, 2013) in codon

space as above. 5′ and 3′ end trimming was performed manually in a codon-based manner.

Generating constraint trees for the *ACC* and Clp subunit alignments for use in PAML

A constraint tree stipulates a fixed topology (branching order) that is used by a phylogenetic

program (in this case, PAML) when calculating branch lengths. To generate a constraint tree for

the *ACC* alignment, RAxML v8.2.12 (Stamatakis, 2014) was used on the trimmed nucleotide

alignment with parameters *-m = GTRGAMMA, -p = 12345, -f = a, -x = 12345,* and *-# = 100*.

The resultant topology confirmed that there were independent *ACC* duplications at the base of

each family (Park et al., 2017; Rockenbach et al., 2016).

To construct constraint trees for Clp core subunits, each trimmed amino acid alignment was

analyzed with ProtTest v3.4.2 (Darriba et al., 2011) to choose a model of sequence evolution.

The top model based on the Bayesian Information Criterion was chosen for use in PhyML v3.3

(Guindon et al., 2010), which was run with 1000 bootstrap replicates and 100 random starts. The

resultant phylogenetic trees were used to determine whether duplication events were lineage-

specific or shared among species in the dataset. In almost all cases, paralogs from a single

species were sister to one another in the trees, indicating lineage-specific duplications. There

were a few cases in which paralogs from a single species were not sister to one another.

However, given low bootstrap support and the difficulty of resolving species relationships using

a single gene with highly variable rates of evolution, we proceeded under the assumption that

these duplications were lineage-specific as well. Thus, the constraint trees for each individual

Clp core subunit were constructed using the known species tree (The Angiosperm Phylogeny

Group et al., 2016), with duplications encoded as species-specific (mapped to terminal branches

of the species tree).

Running PAML for ACCase and Clp core subunit genes

For each alignment, PAML v4.9j (Yang, 2007) was used to infer d_N/d_S values for all branches

using the free ratios model (model = 1) and parameters *CodonFreq = 2* and *cleandata = 0*.

Additionally, *model = 0* and *model = 2* runs were conducted for all alignments, again using

CodonFreq = 2 and *cleandata = 0*. The *model = 0* runs forced all branches to have the same

d_N/d_S ratio, while the *model = 2* runs allowed different d_N/d_S values for specified groups of

branches.

For the *ACC* alignment, one *model = 2* run was conducted with plastid-targeted branches as the

foreground. The resultant tree had one d_N/d_S value for plastid-targeted (*ACC2*) branches and a

second d_N/d_S value for cytosol-targeted (*ACC1*) branches (including all internal pre-duplication

branches). This output was compared with the *model = 0* run to determine whether allowing two

d_N/d_S ratios (one for each of those groups) was a better fit to the data than allowing just a single

d_N/d_S value. For the Clp subunit alignments, *model = 2* was used twice. In the first run, all

terminal branches (and in the case of two subunits, internal post-duplication branches) were

designated as the foreground. In the second run, there were three classes of branches, where all

branches were categorized the same as in the first run except that post-duplication branches

(internal or terminal) were placed in a third category. The three-partition and two-partition

models were compared to determine whether allowing an additional d_N/d_S ratio for post-

duplication branches was a better fit to the data than just separating terminal from internal

branches. The models were compared using likelihood ratio tests.

For the *ACC* alignment, a branch-site test (Yang, 2007; Yang and Nielsen, 2002) was also

conducted to test for evidence of positive selection on branches for plastid-targeted genes, which

were set as the foreground branches for this analysis. A null model and an alternative model both

used the parameters *model = 2, NSsites = 2, CodonFreq = 2*, and *cleandata = 0*. The alternative

model otherwise used all default values, while the null model additionally used *fix_omega = 1*

and *omega = 1*. The models were compared using a likelihood ratio test.

Running HyPhy for *ACC*

In addition to running a PAML branch-site test on the *ACC* alignment (Yang, 2007; Yang and

Nielsen, 2002), tests for positive and relaxed selection were implemented in HyPhy v2.5.32

(Kosakovsky Pond et al., 2020). Positive selection was tested for using the aBSREL and

BUSTED methods (Murrell et al., 2015; Smith et al., 2015). The RELAX method was used to

test for relaxed vs. intensified selection (Wertheim et al., 2015). As with the PAML runs, the

constraint tree used for HyPhy methods had the branches separated into two categories (*ACC1*

and *ACC2*).

Comparisons between *ACC1* and *ACC2* genes

To compare d_N and d_S between cytosolic-targeted and plastid-targeted *ACC* genes (*ACC1* and

ACC2, respectively), a mean root-to-tip distance was calculated for each family in the tree. The

base of each duplication event was used as the root for each family. For both d_N and d_S, the four

mean distances for *ACC1* were compared to those of *ACC2* using a paired t-test in R. Because of

the *a priori* prediction that retargeting to the plastid would be associated with accelerated protein

sequence evolution, a one-sided test (*ACC2 > ACC1*) was used for d_N, while a two-sided test was

used for d_S.

Fisher's exact test on Clp subunit paralogs

Using the output from the free ratios (model = 1) PAML runs, Fisher's exact test was used to test for asymmetry in the ratio of the estimated numbers of nonsynonymous and synonymous substitutions (Pegueroles et al., 2013). Nonsynonymous and synonymous substitution estimates were entered into the *fisher.test()* function in R with default parameters. For each pair of duplicates, a test between paralog 1 and paralog 2 was performed (**Figure 1**). If the paralogs were found to be evolving symmetrically, their combined numbers of substitutions were compared to those of the ancestral branch (**Figure 1**). If the paralogs were found to be evolving asymmetrically, each one was compared individually against the ancestral branch (**Figure 1**). The four cases in which there were more than two species-specific paralogs (*Soja max* and *Gossypium raimondii CLPP5*; *Musa acuminata* and *Vitis vinifera CLPR4*) were excluded from this analysis.

Data availability

Scripts, untrimmed and trimmed alignments, PAML output, and HyPhy output are provided for both *ACC* and Clp subunits at https://github.com/alissawilliams/Gene_duplication_ACCase_Clp.

Results:

Plastid-targeted ACCases evolve more rapidly than cytosol-targeted ACCases across angiosperms

Across the sampled clades (Geraniaceae, Caryophyllaceae, Brassicaceae, and Poaceae), nearly all plastid-targeted *ACC2* genes have higher d_N/d_S values than their cytosol-targeted *ACC1* counterparts (**Figure 2, Figure S1**). The single-partition model assigned all branches a d_N/d_S value of 0.1266, while the two-partition model assigned *ACC1* branches a value of 0.0883 and *ACC2* branches a value of 0.1936 ($\chi^2 = 466.84$, p << 0.0001). This pattern is true for both terminal and internal branches. The increase in in d_N/d_S ratios in *ACC2* branches is generally

driven by increases in d_N rather than reductions in d_S (t = 4.48, p = 0.01 for d_N; t = 0.72, p =

0.5249 for d_S; **Figure 2, Figure S2**), suggesting changes in selective pressure.

Using a branch-sites test in PAML (Yang, 2007), we did not find a significant signature of

positive selection spanning the alignment ($\chi^2 = 0$, p = 1), although there were multiple individual

sites found to be under positive selection (**Table S1**). Two HyPhy methods found limited, though

significant, evidence for positive selection—the aBSREL run (Smith et al., 2015) detected one

branch under positive selection (p = 0.04) and the BUSTED run (Murrell et al., 2015) assigned

0.12% of sites in foreground (*ACC2*) branches to the positive selection class relative to 0.05% of

sites in background (*ACC1*) branches (p = 0.0026). The HyPhy RELAX method (Wertheim et

al., 2015) found significant evidence for relaxed selection in the *ACC2* branches relative to the

rest of the tree (K = 0.09, p <<0.001).

Characterizing ongoing duplication of nuclear-encoded Clp core subunit genes in

angiosperms

Of the 23 angiosperm species in our dataset, 11 had one or more duplications of nuclear genes

encoding Clp core subunits, and all eight of these genes were duplicated in at least one species

(**Figure 3**). Most of these duplications were represented by two paralogs, but in four cases, we

identified more than two paralogs for a particular subunit in a particular species. For *CLPP5*,

Soja max and *Gossypium raimondii* have five and seven copies, respectively, and for *CLPR4*,

both *Musa acuminata* and *Vitis vinifera* have four copies.

Soja max had duplications of the largest number of subunits (six of eight), followed by *Plantago*

maritima and *Populus trichocarpa* with duplications of five subunits. Of the 11 species with

duplications, *Eucalyptus grandis* and *Oenothera biennis* were the only species that had

duplications of just one subunit. Across subunits, *CLPP5* had the highest number of paralogs (37

in 23 species) and *CLPR2* had the lowest (24 in 23 species).

In total, we identified 72 gene copies of Clp core subunits resulting from duplication events,

including 40 catalytic subunits (*CLPP3-CLPP6*) (**Figure 3**). Of the 40 catalytic paralogs, we

found evidence of loss of one or more catalytic sites in multiple genes (**Table S2**). Across all 72

paralogs, we also found evidence of truncation of multiple different gene copies (including some

with catalytic site loss) (**Table S2, Table S3**).

Recent paralogs of Clp core subunits tend to have higher rates of protein sequence

evolution than their single-copy counterparts

Out of the eight nuclear-encoded Clp core subunit trees (**Figures S3-S10**), seven showed

statistically significant differences between a model that allowed for different d_N/d_S rates in gene

duplicates vs. single-copy genes (the three-partition model) and one that forces the same d_N/d_S

rate on these two types of branches (the two-partition model) based on an uncorrected

significance threshold of $p = 0.05$. (**Figure 4, Table 1**). In six of those cases, duplicated terminal

branches had a higher d_N/d_S rate than non-duplicated terminal branches, while in the remaining

case, the reverse was true. We separated internal branches from terminal branches to account for

the fact that terminal branches will, on average, have higher d_N/d_S estimates than internal

branches because selection has had more time to act on older deleterious mutations (Hasegawa et

al., 1998; Ho et al., 2005). Further, terminal branches represent both interspecific divergence and

intraspecific polymorphism, which is important because the latter inflates evolutionary rate

calculations (Ho et al., 2005; Moilanen and Majamaa, 2003; Nielsen and Weinreich, 1999).

We also compared the evolutionary rates of paralogs to one another as well as to their common

ancestor, again using an uncorrected significance threshold of p = 0.05 (**Table 2**). Of the 26 pairs

of paralogs, 13 (50%) showed statistically significant rate asymmetry relative to each another. In

10 (77%) of those cases, only one paralog had a significantly different evolutionary rate than the

common ancestor (and in all 10 of those cases, that paralog was evolving at a faster rate than the

common ancestor). Of the 13 pairs with symmetric evolutionary rates, five (38%) were

asymmetric relative to the common ancestor. In three of those cases, the combined paralog

evolutionary rate was significantly faster than that of the ancestor.

Discussion:

Neofunctionalization and accelerated evolution of duplicated *ACC* genes in multiple clades of flowering plants

Independent duplications of *ACC* and subsequent retargeting events have been previously reported in multiple angiosperm clades (Babiychuk et al., 2011; Konishi and Sasaki, 1994; Park et al., 2017; Parker et al., 2014; Rockenbach et al., 2016; Schulte et al., 1997). The process of retargeting of a paralog is inherently a form of neofunctionalization because the newly retargeted protein functions in a different cellular compartment than it did ancestrally. A hallmark of neofunctionalization is evolutionary rate asymmetry between paralogs due to selection associated with gaining a new function (Hahn, 2009; Pegueroles et al., 2013; Zhang, 2003). We found that branches of our *ACC* tree representing paralogs targeted to the plastid had statistically significantly higher d_N/d_S values than branches representing paralogs targeted to the cytosol (**Figure 2, Figure S1**), consistent with the predictions under neofunctionalization. These results were based on a trimmed alignment lacking the target peptide, which we excluded because target peptides exhibit fast rates of evolution and reduced constraints on primary amino acid sequence (Bruce, 2001, 2000; Jarvis, 2008). Thus, our results show that *ACC* genes retargeted to the plastid are undergoing evolutionary rate increases unrelated to the target peptide, suggesting that other functional domains are also evolving rapidly.

Retargeting of the cytosolic, homomeric ACCase protein to the plastid is somewhat unexpected given that a heteromeric ACCase complex already exists in plastids. Whether the retargeted homomeric ACCases functionally replaces or coexists with the heteromeric version appears to vary across clades. In some angiosperm groups, the two complexes coexist, including in *Arabidopsis thaliana* and likely in other members of the Brassicaceae (Babiychuk et al., 2011; Rousseau-Gueutin et al., 2013). In other clades, the homomeric ACCase has replaced the heteromeric version, as was reported in the Poaceae (Konishi and Sasaki, 1994). The duplication found in *Silene noctiflora* and *Silene paradoxa* may also represent a replacement event given that both species lack at least one heteromeric ACCase gene each, where *S. noctiflora* lacks all of them (Rockenbach et al., 2016). In some cases, including *Monsonia emarginata* in the Geraniaceae, the plastid-encoded *accD* gene of the heteromeric complex has been transferred to the nuclear genome, again suggesting that the heteromeric version is still functional (Park et al.,

2017; Rousseau-Gueutin et al., 2013). These contrasting histories of replacement vs. coexistence

may mean that duplicates in different clades are evolving under different selection regimes.

Variation in post-duplication fates could confound tests of selection conducted across the entire

ACC tree. Using PAML and HyPhy (Murrell et al., 2015; Smith et al., 2015; Wertheim et al.,

2015; Yang, 2007), we tested for positive selection and relaxed selection in *ACC2* genes relative

to *ACC1* genes, both of which can contribute to increased rates of protein sequence evolution.

The results were mixed; there is some evidence for relaxed selection across all *ACC2* branches as

well as for positive selection in a small number of branches and sites (**Table S1**). Across the four

families in our sample, the smallest ratio between mean *ACC2* d_N and mean *ACC1* d_N was found

in the Poaceae (1.5 vs. 2.2-2.6 for the other three families). Since the heteromeric ACCase is

completely absent in the Poaceae (Konishi and Sasaki, 1994), we would expect stronger

purifying selection on the plastid homomeric ACCase in this clade compared to clades in which

the two versions coexist. Thus, these results are consistent with the hypothesis that relaxed

selection is contributing to rate accelerations and that there is greater relaxation of selection

when homomeric and heteromeric ACCases functions redundantly in the plastid, though the

evidence is still limited. The potential for positive selection on retargeted ACCases is intriguing

given that these proteins are thought to perform the same function as the ancestral protein; it is

possible that retargeted proteins are adapting to specific biochemical and/or osmotic conditions

within the new destination. Increased evolutionary rates after subcellular retargeting have been

previously noted, though we do not fully understand their underlying causes (Byun-McKay and

Geeta, 2007; Marques et al., 2008).

Ongoing duplication of nuclear-encoded Clp core subunit genes is common in angiosperms

Across green plants, duplication of the plastid-encoded Clp core subunit gene *clpP1* has only

been found in a handful of lineages (Williams et al., 2019). While other studies have identified

recent duplications of nuclear-encoded Clp core subunit genes (Rockenbach et al., 2016;

Williams et al., 2021), our work shows that duplications of these nuclear-encoded subunits are

pervasive across angiosperms (**Figure 3**). Because we used a mix of transcriptomic and genomic

data, we took into consideration the possibility of misidentifying transcript variants as paralogs

but our use of primary transcripts only and manual curation to remove hits that appeared to be

splice variants (see Materials and Methods) minimizes the risk of this type of error.

The prevalence of whole genome duplication in plants may partially explain the prevalence of

Clp core subunit duplication (Clark and Donoghue, 2018; De Bodt et al., 2005; del Pozo and

Ramirez-Parra, 2015; Flagel and Wendel, 2009; Panchy et al., 2016; Wendel et al., 2018). For

instance, *Soja max* is a partially diploidized tetraploid, meaning that this lineage underwent a

polyploidization event very recently and has only just started the subsequent process of genome

reduction (Shultz et al., 2006). *Soja max* had the largest number of duplicated subunits across our

sample, which is consistent with this history of whole genome duplication. Similarly, *Populus*

trichopoda, which tied for the second largest number of duplicated subunits, only recently

underwent genome reduction after whole genome duplication (Tuskan et al., 2006). In these

cases, we may simply be observing the short-term effects of polyploidization prior to returning to

a single copy of each of these genes.

Subunit stoichiometry and subfunctionalization in the evolution of the plastid Clp complex

Clp core subunit ratios have been studied in *Arabidopsis thaliana* (Olinares et al., 2011a). The

core consists of two rings—a ClpP1/ClpR1-4 ring with a 3:1:1:1:1 subunit ratio, respectively,

and a ClpP3-6 ring with a 1:2:3:1 subunit ratio, respectively (Olinares et al., 2011a). Despite the

high degree of structural similarity amongst the plastid Clp core subunits, core composition (i.e.

the number of each type of core subunit) does not appear to vary in *A. thaliana* (Olinares et al.,

2011a; Peltier et al., 2004). Due to the stability of subunit interactions in *A. thaliana*, Clp

complexes in other angiosperms are typically assumed to have the same ratios of core subunits,

but our results suggest that varied numbers of core subunit paralogs may lead to varied

stoichiometry across species. Additional work has shown that loss of catalytic activity in ClpP5

(present in three copies in *A. thaliana*) is lethal while loss of catalytic activity in ClpP3 (present

in one copy in *A. thaliana*) is tolerated, suggesting that core subunit composition may be flexible

given a threshold number of catalytic subunits (Liao et al., 2018).

In fact, core subunit composition has been dynamic throughout the evolutionary history of the green lineage. The Clp complex is widely conserved across bacteria; in most bacteria, including *E. coli*, the Clp core consists of 14 identical subunits (Nishimura and van Wijk, 2015; Yu and Houry, 2007). However, in cyanobacteria, several duplications have produced four core subunits—three catalytic ClpP subunits and one catalytically inactive ClpR subunit (Andersson et al., 2009; Stanne et al., 2007). In green lineage (Viridiplantae) plastids, which are descended from ancient cyanobacteria, gene duplication has continued to expand the number of genes incorporated into the Clp core to yield nine genes (*CLPP1,3-6*, *CLPR1-4*) (Nishimura and van Wijk, 2015). Interestingly, ClpR subunits are incorporated into the core despite their lack of catalytic activity; they are thought to play a structural role in the complex, including chaperone docking (Nishimura and van Wijk, 2015; Olinares et al., 2011b, 2011a; Sjögren and Clarke, 2011). In *A. thaliana*, the Clp chaperone is believed to bind only to the ClpP1/ClpR1-4 ring, whereas chaperone proteins bind to both rings of the Clp core in bacteria (Peltier et al., 2004; Yu and Houry, 2007). This ClpP/ClpR division of function (catalytic activity vs chaperone binding) is indicative of subfunctionalization. Further, though the plastid Clp core subunit genes share common ancestry and are structurally similar, knockouts of individual subunits tend to produce severe phenotypes, including lethality in several cases (Kim et al., 2009; Koussevitzky et al., 2007; Rudella et al., 2006).

Possible subfunctionalization in recent paralogs of Clp core subunits

Given that subfunctionalization has likely played a major role in plastid Clp complex evolution, we were particularly interested in whether we could identify subfunctionalization after more recent duplication events. Taken to an extreme, subfunctionalization would involve having one gene for each of the 14 core subunits, which would lead to further expansion of the typical nine core subunit genes. The total number of core subunits after including recent paralogs and the plastid-encoded ClpP1 was less than 14 in most species. *Musa acuminata*, *Plantago maritima*, and *Populus trichocarpa* had 14 each, *Soja max* had 17, and *Gossypium raimondii* had 15 (**Figure 3**). The numbers larger than 14 were driven in both cases by multiple paralogs of *CLPP5*, with five and seven copies, respectively. ClpP5 has the largest number of subunits stoichiometrically among the eight nuclear-encoded subunits, so the fact that the two largest

numbers of paralogs were both found in *CLPP5* could potentially suggest that some species are

moving toward a 1:1 relationship between genes and core subunits. However, this explanation is

not supported by other evidence. For example, the other cases of >2 paralogs were found for

CLPR4, which encodes a protein that is present in just a single copy in the core in *Arabidopsis*.

Further, it is not clear that all of these paralogs are capable of producing functional proteins

given truncations and loss of catalytic sites (**Table S2, Table S3**).

We tested for signatures of subfunctionalization by looking at evolutionary rate asymmetry.

Under subfunctionalization, we would expect paralogs to evolve at symmetric rates relative to

one another but asymmetrically relative to their common ancestor (Pegueroles et al., 2013). We

found five of these cases in our dataset: *Plantago maritima CLPP3, Geranium maderense*

CLPP4, Medicago truncatula CLPP5, Populus trichocarpa CLPP5, and *Soja max CLPR1*

(**Table 2**). In cases of subfunctionalization, we would expect the paralogs to evolve more quickly

than the common ancestor because of relaxed selection due to their more limited functional roles,

which was only the case for the former three. In those three cases, the evidence is consistent with

subfunctionalization, particularly given that all six involved paralogs are full length. Further, the

P. maritima CLPP3 paralogs share the same substitutions in all three catalytic sites, which

indicates duplication after the loss of catalytic activity, and the *G. maderense CLPP4* and *M.*

truncatula CLPP5 paralogs all have fully retained catalytic triads (**Table S2**).

Possible pseudogenization or neofunctionalization in recent paralogs of Clp core subunits

Predictions about evolutionary rates under neofunctionalization are similar to predictions under

the degeneration/gene loss model—one paralog will maintain the ancestral evolutionary rate

while the other undergoes evolutionary rate acceleration (Hahn, 2009; Pegueroles et al., 2013;

Zhang, 2003). Previous work in this complex has shown that even ClpP1 subunits demonstrating

massive accelerations in evolutionary rate can still be functional, meaning that high evolutionary

rates alone do not necessarily indicate pseudogenization (Barnard-Kubow et al., 2014; Williams

et al., 2019, 2015). Other sequence features can help us differentiate between pseudogenization

and neofunctionalization. For example, truncation of a sequence can be evidence that it is no

longer producing a functional protein; additionally, for ClpP subunits, loss of catalytic sites may

also be an indication of degeneration/pseudogenization (though there may be exceptions,

including the *P. maritima CLPP3* paralogs mentioned above). In our dataset, the paralogs of

Musa acuminata CLPP4 and *P. trichocarpa CLPP4* follow these patterns (**Table 2**). In each of

these pairs, the paralogs are evolving asymmetrically, and the paralog with a faster rate of

evolution is truncated and lacking all three catalytic sites, suggesting loss of function (**Table S2**).

Another example of probable pseudogenization is found for the second copy of *M. acuminata*

CLPR1. This paralog was annotated as two separate genes due to an internal stop codon, which

would lead to a truncation in the resultant protein.

As for neofunctionalization, there are other cases in our dataset where paralogs evolving

asymmetrically both have retained catalytic sites and are full length (for instance, *Geranium*

maderense CLPP5 and *Oenothera biennis CLPP5*). There are no known instances of retargeting

of plastid Clp core subunits; thus, evolutionary drivers of neofunctionalization of duplicated

subunits are unknown. It is possible that neofunctionalization in this complex could involve

recruiting additional interacting proteins—the ClpT proteins, for instance, are involved in

assembly of the core and are a recently evolutionary innovation specific to green plants

(Colombo et al., 2014; Kim et al., 2015; Nishimura and van Wijk, 2015; Sjögren and Clarke,

2011). Additionally, ongoing work has identified potential new adapter proteins in the plastid

Clp complex (Montandon et al., 2019; Nishimura et al., 2015). Another possibility is tissue-

specific expression of paralogs, which has not been documented in the Clp complex but has been

identified in mitochondrial complexes (Boss et al., 1997; Guerrero-Castillo et al., 2017; Sinkler

et al., 2017).

Possible retention of Clp core paralogs under the gene dosage advantage hypothesis

We also have eight cases of symmetrically evolving paralogs that are also evolving

symmetrically relative to the common ancestor (**Table 2**). Under our initial predictions, these

would represent paralogs retained under the gene dosage advantage hypothesis (Hahn, 2009;

Ohno, 1970; Pegueroles et al., 2013; Zhang, 2003). Of these eight paralog pairs, four are from

Soja max (which had six total pairs of paralogs), and three are from *Populus trichocarpa* (which

had five total pairs of paralogs). As described above, both of these species are in the process of

rediploidization after a recent whole genome duplication (Shultz et al., 2006; Tuskan et al.,

2006). It is possible that these results reflect the fact that the gene duplications happened so

recently that the paralogs have not had time to diverge. This possibility is further supported given

that the estimates of numbers of substitutions for many of these paralogs were so low that there

was virtually no power to detect significant asymmetry.

Alternative hypotheses and future directions

While we based our analyses on established expectations for evolutionary rates under different

post-duplication fates (gene dosage advantage, neofunctionalization, and subfunctionalization),

other work has challenged the universality of these predictions. He and Zhang (2005) outline the

subneofunctionalization model, in which gene duplicates undergo rapid subfunctionalization

followed by prolonged neofunctionalization. Asymmetric evolutionary rates are often assumed to

be the result of either neofunctionalization or degeneration, but subfunctionalization can also

occur in an asymmetric fashion (He and Zhang, 2005). This hypothesis could relate to some of

our results; cases of asymmetric evolutionary rates could be due to subfunctionalization rather

than neofunctionalization. Additionally, functional constraint can also exist under

neofunctionalization, leading to lower substitution rates and possibly symmetric rates of

evolution, meaning that symmetrically evolving paralogs could represent cases of

neofunctionalization rather than subfunctionalization or gene dosage advantage (He and Zhang,

2005).

Regardless, our results demonstrate that post-duplication evolutionary fates of paralogs vary

widely across clades, even when the same genes are involved. Duplications of the homomeric

ACCase complex gene (*ACC*) and subsequent retargeting of one copy to the plastid have been

previously reported (Babiychuk et al., 2011; Konishi and Sasaki, 1994; Park et al., 2017; Parker

et al., 2014; Rockenbach et al., 2016; Schulte et al., 1997). Our results show that the retargeted

duplicates almost universally have increased d_N/d_S rates (**Figure 2, Figure S1**). As for plastid

Clp core subunit duplications, duplication has clearly shaped this complex over the course of

Viridiplantae evolution. We provide evidence of all possible post-duplication routes of recent

paralogs amongst the different subunits and different species in our dataset. Overall, our results

are compelling evidence that subunit ratios and stoichiometry may be dynamic across

angiosperm lineages. Isolation of plastid Clp complexes and analyses of subunit composition

have been performed in a handful of species (Moreno et al., 2017; Olinares et al., 2011a;

Williams et al., 2019); future work could determine these compositions in other angiosperms,

including those that have undergone recent gene duplications. Our work demonstrates that gene

duplication has been and continues to be an important force in plastid evolution.

<u>Acknowledgements</u>

This work was supported by a National Science Foundation (NSF) grant (MCB-1733227),

graduate fellowships from NSF (DGE-1321845) and the National Institutes of Health (T32-

GM132057), and the Wolves to Rams undergraduate research program (NSF Grant Numbers

1930150 and 19300092, NIH Grant Number 1T34GM137861-01).

<u>References</u>

Andersson, F.I., Tryggvesson, A., Sharon, M., Diemand, A.V., Classen, M., Best, C., Schmidt,
R., Schelin, J., Stanne, T.M., Bukau, B., Robinson, C.V., Witt, S., Mogk, A., Clarke,
A.K., 2009. Structure and function of a novel type of ATP-dependent Clp protease. J.
Biol. Chem. 284, 13519–13532. https://doi.org/10.1074/jbc.M809588200

Apitz, J., Nishimura, K., Schmied, J., Wolf, A., Hedtke, B., Wijk, K.J. van, Grimm, B., 2016.
Posttranslational Control of ALA Synthesis Includes GluTR Degradation by Clp Protease
and Stabilization by GluTR-Binding Protein. Plant Physiology 170, 2040–2051.
https://doi.org/10.1104/pp.15.01945

Babiychuk, E., Vandepoele, K., Wissing, J., Garcia-Diaz, M., Rycke, R.D., Akbari, H., Joubès,
J., Beeckman, T., Jänsch, L., Frentzen, M., Montagu, M.C.E.V., Kushnir, S., 2011.
Plastid gene expression and plant development require a plastidic protein of the
mitochondrial transcription termination factor family. PNAS 108, 6674–6679.
https://doi.org/10.1073/pnas.1103442108

Barnard-Kubow, K.B., Sloan, D.B., Galloway, L.F., 2014. Correlation between sequence
divergence and polymorphism reveals similar evolutionary mechanisms acting across
multiple timescales in a rapidly evolving plastid genome. BMC Evol Biol 14.
https://doi.org/10.1186/s12862-014-0268-y

Boss, O., Samec, S., Paoloni-Giacobino, A., Rossier, C., Dulloo, A., Seydoux, J., Muzzin, P.,
Giacobino, J.P., 1997. Uncoupling protein-3: a new member of the mitochondrial carrier
family with tissue-specific expression. FEBS Lett 408, 39–42.
https://doi.org/10.1016/s0014-5793(97)00384-0

Bouchnak, I., van Wijk, K.J., 2021. Structure, Function and Substrates of Clp AAA+ protease
systems in cyanobacteria, plastids and apicoplasts; a comparative analysis. Journal of
Biological Chemistry 100338. https://doi.org/10.1016/j.jbc.2021.100338
Bruce, B.D., 2001. The paradox of plastid transit peptides: conservation of function despite
divergence in primary structure. Biochimica et Biophysica Acta (BBA) - Molecular Cell
Research 1541, 2–21. https://doi.org/10.1016/S0167-4889(01)00149-5
Bruce, B.D., 2000. Chloroplast transit peptides: structure, function and evolution. Trends in Cell
Biology 10, 440–447. https://doi.org/10.1016/S0962-8924(00)01833-X
Byun-McKay, S.A., Geeta, R., 2007. Protein subcellular relocalization: a new perspective on the
origin of novel genes. Trends in Ecology & Evolution 22, 338–344.
https://doi.org/10.1016/j.tree.2007.05.002
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L.,
2009. BLAST+: architecture and applications. BMC Bioinformatics 10, 421.
https://doi.org/10.1186/1471-2105-10-421
Caroca, R., Howell, K.A., Malinova, I., Burgos, A., Tiller, N., Pellizzer, T., Annunziata, M.G.,
Hasse, C., Ruf, S., Karcher, D., Bock, R., 2021. Knock-down of the plastid-encoded
acetyl-CoA carboxylase gene uncovers functions in metabolism and development. Plant
Physiology. https://doi.org/10.1093/plphys/kiaa106
Castresana, J., 2000. Selection of Conserved Blocks from Multiple Alignments for Their Use in
Phylogenetic Analysis. Molecular Biology and Evolution 17, 540–552.
https://doi.org/10.1093/oxfordjournals.molbev.a026334
Clark, J.W., Donoghue, P.C.J., 2018. Whole-Genome Duplication and Plant Macroevolution.
Trends in Plant Science 23, 933–945. https://doi.org/10.1016/j.tplants.2018.07.006
Colombo, C.V., Ceccarelli, E.A., Rosano, G.L., 2014. Characterization of the accessory protein
ClpT1 from Arabidopsis thaliana: oligomerization status and interaction with Hsp100
chaperones. BMC Plant Biology 14, 228. https://doi.org/10.1186/s12870-014-0228-0
Conant, G.C., Wagner, A., 2003. Asymmetric Sequence Divergence of Duplicate Genes.
Genome Res. 13, 2052–2058. https://doi.org/10.1101/gr.1252603
Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2011. ProtTest 3: fast selection of best-fit
models of protein evolution. Bioinformatics 27, 1164–1165.
https://doi.org/10.1093/bioinformatics/btr088
De Bodt, S., Maere, S., Van de Peer, Y., 2005. Genome duplication and the origin of
angiosperms. Trends in Ecology & Evolution 20, 591–597.
https://doi.org/10.1016/j.tree.2005.07.008
del Pozo, J.C., Ramirez-Parra, E., 2015. Whole genome duplications in plants: an overview from
Arabidopsis. Journal of Experimental Botany 66, 6991–7003.
https://doi.org/10.1093/jxb/erv432
Dermitzakis, E.T., Clark, A.G., 2001. Differential selection after duplication in mammalian
developmental genes. Mol Biol Evol 18, 557–562.
https://doi.org/10.1093/oxfordjournals.molbev.a003835
Des Marais, D.L., Rausher, M.D., 2008. Escape from adaptive conflict after duplication in an
anthocyanin pathway gene. Nature 454, 762–765. https://doi.org/10.1038/nature07092
Erixon, P., Oxelman, B., 2008. Whole-Gene Positive Selection, Elevated Synonymous
Substitution Rates, Duplication, and Indel Evolution of the Chloroplast clpP1 Gene.
PLOS ONE 3, e1386. https://doi.org/10.1371/journal.pone.0001386

Flagel, L.E., Wendel, J.F., 2009. Gene duplication and evolutionary novelty in plants. New
Phytologist 183, 557–564. https://doi.org/10.1111/j.1469-8137.2009.02923.x
Force, A., Lynch, M., Pickett, F.B., Amores, A., Yan, Y., Postlethwait, J., 1999. Preservation of
Duplicate Genes by Complementary, Degenerative Mutations. Genetics 151, 1531–1545.
Freeling, M., Lyons, E., Pedersen, B., Alam, M., Ming, R., Lisch, D., 2008. Many or most genes
in Arabidopsis transposed after the origin of the order Brassicales. Genome Res 18,
1924–1937. https://doi.org/10.1101/gr.081026.108
Goodstein, D.M., Shu, S., Howson, R., Neupane, R., Hayes, R.D., Fazo, J., Mitros, T., Dirks, W.,
Hellsten, U., Putnam, N., Rokhsar, D.S., 2012. Phytozome: a comparative platform for
green plant genomics. Nucleic Acids Research 40, D1178–D1186.
https://doi.org/10.1093/nar/gkr944
Green, B.R., 2011. Chloroplast genomes of photosynthetic eukaryotes. The Plant Journal 66, 34–
44. https://doi.org/10.1111/j.1365-313X.2011.04541.x
Guerrero-Castillo, S., Cabrera-Orefice, A., Huynen, M.A., Arnold, S., 2017. Identification and
evolutionary analysis of tissue-specific isoforms of mitochondrial complex I subunit
NDUFV3. Biochim Biophys Acta Bioenerg 1858, 208–217.
https://doi.org/10.1016/j.bbabio.2016.12.004
Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New
Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the
Performance of PhyML 3.0. Syst Biol 59, 307–321.
https://doi.org/10.1093/sysbio/syq010
Hahn, M.W., 2009. Distinguishing among evolutionary models for the maintenance of gene
duplicates. J Hered 100, 605–617. https://doi.org/10.1093/jhered/esp047
Hasegawa, M., Cao, Y., Yang, Z., 1998. Preponderance of slightly deleterious polymorphism in
mitochondrial DNA: nonsynonymous/synonymous rate ratio is much higher within
species than between species. Mol Biol Evol 15, 1499–1505.
https://doi.org/10.1093/oxfordjournals.molbev.a025877
He, X., Zhang, J., 2005. Rapid Subfunctionalization Accompanied by Prolonged and Substantial
Neofunctionalization in Duplicate Gene Evolution. Genetics 169, 1157–1164.
https://doi.org/10.1534/genetics.104.037051
Ho, S.Y.W., Phillips, M.J., Cooper, A., Drummond, A.J., 2005. Time Dependency of Molecular
Rate Estimates and Systematic Overestimation of Recent Divergence Times. Molecular
Biology and Evolution 22, 1561–1568. https://doi.org/10.1093/molbev/msi145
Huang, Y., Kendall, T., Forsythe, E.S., Dorantes-Acosta, A., Li, S., Caballero-Pérez, J., Chen,
X., Arteaga-Vázquez, M., Beilstein, M.A., Mosher, R.A., 2015. Ancient Origin and
Recent Innovations of RNA Polymerase IV and V. Mol Biol Evol 32, 1788–1799.
https://doi.org/10.1093/molbev/msv060
Jansen, R.K., Cai, Z., Raubeson, L.A., Daniell, H., dePamphilis, C.W., Leebens-Mack, J.,
Müller, K.F., Guisinger-Bellian, M., Haberle, R.C., Hansen, A.K., Chumley, T.W., Lee,
S.-B., Peery, R., McNeal, J.R., Kuehl, J.V., Boore, J.L., 2007. Analysis of 81 genes from
64 plastid genomes resolves relationships in angiosperms and identifies genome-scale
evolutionary patterns. PNAS 104, 19369–19374.
https://doi.org/10.1073/pnas.0709121104
Jarvis, P., 2008. Targeting of nucleus-encoded proteins to chloroplasts in plants. New
Phytologist 179, 257–285. https://doi.org/10.1111/j.1469-8137.2008.02452.x

Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. Mol Biol Evol 30, 772–780. https://doi.org/10.1093/molbev/mst010

Kim, J., Kimber, M.S., Nishimura, K., Friso, G., Schultz, L., Ponnala, L., Wijk, K.J. van, 2015. Structures, Functions, and Interactions of ClpT1 and ClpT2 in the Clp Protease System of Arabidopsis Chloroplasts. The Plant Cell 27, 1477–1496. https://doi.org/10.1105/tpc.15.00106

Kim, J., Rudella, A., Rodriguez, V.R., Zybailov, B., Olinares, P.D.B., Wijk, K.J. van, 2009. Subunits of the Plastid ClpPR Protease Complex Have Differential Contributions to Embryogenesis, Plastid Biogenesis, and Plant Development in Arabidopsis. The Plant Cell 21, 1669–1692. https://doi.org/10.1105/tpc.108.063784

Kondrashov, F.A., Rogozin, I.B., Wolf, Y.I., Koonin, E.V., 2002. Selection in the evolution of gene duplications. Genome Biol 3, RESEARCH0008. https://doi.org/10.1186/gb-2002-3-2-research0008

Konishi, T., Sasaki, Y., 1994. Compartmentalization of two forms of acetyl-CoA carboxylase in plants and the origin of their tolerance toward herbicides. PNAS 91, 3598–3601. https://doi.org/10.1073/pnas.91.9.3598

Konishi, T., Shinohara, K., Yamada, K., Sasaki, Y., 1996. Acetyl-CoA Carboxylase in Higher Plants: Most Plants Other Than Gramineae Have Both the Prokaryotic and the Eukaryotic Forms of This Enzyme. Plant and Cell Physiology 37, 117–122. https://doi.org/10.1093/oxfordjournals.pcp.a028920

Kosakovsky Pond, S.L., Poon, A.F.Y., Velazquez, R., Weaver, S., Hepler, N.L., Murrell, B., Shank, S.D., Magalis, B.R., Bouvier, D., Nekrutenko, A., Wisotsky, S., Spielman, S.J., Frost, S.D.W., Muse, S.V., 2020. HyPhy 2.5—A Customizable Platform for Evolutionary Hypothesis Testing Using Phylogenies. Molecular Biology and Evolution 37, 295–299. https://doi.org/10.1093/molbev/msz197

Koussevitzky, S., Stanne, T.M., Peto, C.A., Giap, T., Sjögren, L.L.E., Zhao, Y., Clarke, A.K., Chory, J., 2007. An <Emphasis Type="Italic">Arabidopsis thaliana</Emphasis> virescent mutant reveals a role for ClpR1 in plastid development. Plant Mol Biol 63, 85–96. https://doi.org/10.1007/s11103-006-9074-2

Leebens-Mack, J.H., Barker, M.S., Carpenter, E.J., Deyholos, M.K., Gitzendanner, M.A., Graham, S.W., Grosse, I., Li, Z., Melkonian, M., Mirarab, S., Porsch, M., Quint, M., Rensing, S.A., Soltis, D.E., Soltis, P.S., Stevenson, D.W., Ullrich, K.K., Wickett, N.J., DeGironimo, L., Edger, P.P., Jordon-Thaden, I.E., Joya, S., Liu, T., Melkonian, B., Miles, N.W., Pokorny, L., Quigley, C., Thomas, P., Villarreal, J.C., Augustin, M.M., Barrett, M.D., Baucom, R.S., Beerling, D.J., Benstein, R.M., Biffin, E., Brockington, S.F., Burge, D.O., Burris, J.N., Burris, K.P., Burtet-Sarramegna, V., Caicedo, A.L., Cannon, S.B., Çebi, Z., Chang, Y., Chater, C., Cheeseman, J.M., Chen, T., Clarke, N.D., Clayton, H., Covshoff, S., Crandall-Stotler, B.J., Cross, H., dePamphilis, C.W., Der, J.P., Determann, R., Dickson, R.C., Di Stilio, V.S., Ellis, S., Fast, E., Feja, N., Field, K.J., Filatov, D.A., Finnegan, P.M., Floyd, S.K., Fogliani, B., García, N., Gâteblé, G., Godden, G.T., Goh, F. (Qi Y., Greiner, S., Harkess, A., Heaney, J.M., Helliwell, K.E., Heyduk, K., Hibberd, J.M., Hodel, R.G.J., Hollingsworth, P.M., Johnson, M.T.J., Jost, R., Joyce, B., Kapralov, M.V., Kazamia, E., Kellogg, E.A., Koch, M.A., Von Konrat, M., Könyves, K., Kutchan, T.M., Lam, V., Larsson, A., Leitch, A.R., Lentz, R., Li, F.-W., Lowe, A.J., Ludwig, M., Manos, P.S., Mavrodiev, E., McCormick, M.K., McKain, M., McLellan, T.,

McNeal, J.R., Miller, R.E., Nelson, M.N., Peng, Y., Ralph, P., Real, D., Riggins, C.W.,
     Ruhsam, M., Sage, R.F., Sakai, A.K., Scascitella, M., Schilling, E.E., Schlösser, E.-M.,
     Sederoff, H., Servick, S., Sessa, E.B., Shaw, A.J., Shaw, S.W., Sigel, E.M., Skema, C.,
     Smith, A.G., Smithson, A., Stewart, C.N., Stinchcombe, J.R., Szövényi, P., Tate, J.A.,
     Tiebel, H., Trapnell, D., Villegente, M., Wang, C.-N., Weller, S.G., Wenzel, M.,
     Weststrand, S., Westwood, J.H., Whigham, D.F., Wu, S., Wulff, A.S., Yang, Y., Zhu, D.,
     Zhuang, C., Zuidof, J., Chase, M.W., Pires, J.C., Rothfels, C.J., Yu, J., Chen, C., Chen,
     L., Cheng, S., Li, J., Li, R., Li, X., Lu, H., Ou, Y., Sun, X., Tan, X., Tang, J., Tian, Z.,
     Wang, F., Wang, J., Wei, X., Xu, X., Yan, Z., Yang, F., Zhong, X., Zhou, F., Zhu, Y.,
     Zhang, Y., Ayyampalayam, S., Barkman, T.J., Nguyen, N., Matasci, N., Nelson, D.R.,
     Sayyari, E., Wafula, E.K., Walls, R.L., Warnow, T., An, H., Arrigo, N., Baniaga, A.E.,
     Galuska, S., Jorgensen, S.A., Kidder, T.I., Kong, H., Lu-Irving, P., Marx, H.E., Qi, X.,
     Reardon, C.R., Sutherland, B.L., Tiley, G.P., Welles, S.R., Yu, R., Zhan, S., Gramzow,
     L., Theißen, G., Wong, G.K.-S., One Thousand Plant Transcriptomes Initiative, 2019.
     One thousand plant transcriptomes and the phylogenomics of green plants. Nature 574,
     679–685. https://doi.org/10.1038/s41586-019-1693-2
Liao, J.-Y.R., Friso, G., Kim, J., Wijk, K.J. van, 2018. Consequences of the loss of catalytic
     triads in chloroplast CLPPR protease core complexes in vivo. Plant Direct 2, e00086.
     https://doi.org/10.1002/pld3.86
Lynch, M., Conery, J.S., 2000. The Evolutionary Fate and Consequences of Duplicate Genes.
     Science 290, 1151–1155. https://doi.org/10.1126/science.290.5494.1151
Lynch, M., Force, A., 2000. The Probability of Duplicate Gene Preservation by
     Subfunctionalization. Genetics 154, 459–473.
Maere, S., Bodt, S.D., Raes, J., Casneuf, T., Montagu, M.V., Kuiper, M., Peer, Y.V. de, 2005.
     Modeling gene and genome duplications in eukaryotes. PNAS 102, 5454–5459.
     https://doi.org/10.1073/pnas.0501102102
Majeran, W., Wollman, F.-A., Vallon, O., 2000. Evidence for a Role of ClpP in the Degradation
     of the Chloroplast Cytochrome b6f Complex. The Plant Cell 12, 137–149.
     https://doi.org/10.1105/tpc.12.1.137
Marques, A.C., Vinckenbosch, N., Brawand, D., Kaessmann, H., 2008. Functional
     diversification of duplicate genes through subcellular adaptation of encoded proteins.
     Genome Biol 9, R54. https://doi.org/10.1186/gb-2008-9-3-r54
Moilanen, J.S., Majamaa, K., 2003. Phylogenetic network and physicochemical properties of
     nonsynonymous mutations in the protein-coding genes of human mitochondrial DNA.
     Mol Biol Evol 20, 1195–1210. https://doi.org/10.1093/molbev/msg121
Montandon, C., Friso, G., Liao, J.-Y.R., Choi, J., van Wijk, K.J., 2019. In Vivo Trapping of
     Proteins Interacting with the Chloroplast CLPC1 Chaperone: Potential Substrates and
     Adaptors. J. Proteome Res. 18, 2585–2600.
     https://doi.org/10.1021/acs.jproteome.9b00112
Moreno, J.C., Tiller, N., Diez, M., Karcher, D., Tillich, M., Schöttler, M.A., Bock, R., 2017.
     Generation and characterization of a collection of knock-down lines for the chloroplast
     Clp protease complex in tobacco. J Exp Bot 68, 2199–2218.
     https://doi.org/10.1093/jxb/erx066
Murrell, B., Weaver, S., Smith, M.D., Wertheim, J.O., Murrell, S., Aylward, A., Eren, K.,
     Pollner, T., Martin, D.P., Smith, D.M., Scheffler, K., Kosakovsky Pond, S.L., 2015.

Gene-Wide Identification of Episodic Selection. Molecular Biology and Evolution 32,
1365–1371. https://doi.org/10.1093/molbev/msv035
Nielsen, R., Weinreich, D.M., 1999. The age of nonsynonymous and synonymous mutations in
animal mtDNA and implications for the mildly deleterious theory. Genetics 153, 497–
506.
Nishimura, K., Apitz, J., Friso, G., Kim, J., Ponnala, L., Grimm, B., van Wijk, K.J., 2015.
Discovery of a Unique Clp Component, ClpF, in Chloroplasts: A Proposed Binary ClpF-
ClpS1 Adaptor Complex Functions in Substrate Recognition and Delivery. Plant Cell 27,
2677–2691. https://doi.org/10.1105/tpc.15.00574
Nishimura, K., Kato, Y., Sakamoto, W., 2017. Essentials of Proteolytic Machineries in
Chloroplasts. Molecular Plant 10, 4–19. https://doi.org/10.1016/j.molp.2016.08.005
Nishimura, K., van Wijk, K.J., 2015. Organization, function and substrates of the essential Clp
protease system in plastids. Biochimica et Biophysica Acta (BBA) - Bioenergetics, SI:
Chloroplast Biogenesis 1847, 915–930. https://doi.org/10.1016/j.bbabio.2014.11.012
Ohno, S., 1970. Evolution by Gene Duplication. Springer Science & Business Media.
Olinares, P.D.B., Kim, J., Davis, J.I., van Wijk, K.J., 2011a. Subunit stoichiometry, evolution,
and functional implications of an asymmetric plant plastid ClpP/R protease complex in
Arabidopsis. Plant Cell 23, 2348–2361. https://doi.org/10.1105/tpc.111.086454
Olinares, P.D.B., Kim, J., van Wijk, K.J., 2011b. The Clp protease system; a central component
of the chloroplast protease network. Biochimica et Biophysica Acta (BBA) -
Bioenergetics, Regulation of Electron Transport in Chloroplasts 1807, 999–1011.
https://doi.org/10.1016/j.bbabio.2010.12.003
Panchin, A.Y., Gelfand, M.S., Ramensky, V.E., Artamonova, I.I., 2010. Asymmetric and non-
uniform evolution of recently duplicated human genes. Biol Direct 5, 54.
https://doi.org/10.1186/1745-6150-5-54
Panchy, N., Lehti-Shiu, M., Shiu, S.-H., 2016. Evolution of Gene Duplication in Plants. Plant
Physiology 171, 2294–2316. https://doi.org/10.1104/pp.16.00523
Park, S., Ruhlman, T.A., Weng, M.-L., Hajrah, N.H., Sabir, J.S.M., Jansen, R.K., 2017.
Contrasting Patterns of Nucleotide Substitution Rates Provide Insight into Dynamic
Evolution of Plastid and Mitochondrial Genomes of Geranium. Genome Biol Evol 9,
1766–1780. https://doi.org/10.1093/gbe/evx124
Parker, N., Wang, Y., Meinke, D., 2014. Natural Variation in Sensitivity to a Loss of Chloroplast
Translation in Arabidopsis. Plant Physiology 166, 2013–2027.
https://doi.org/10.1104/pp.114.249052
Paterson, A.H., Chapman, B.A., Kissinger, J.C., Bowers, J.E., Feltus, F.A., Estill, J.C., 2006.
Many gene and domain families have convergent fates following independent whole-
genome duplication events in Arabidopsis, Oryza, Saccharomyces and Tetraodon. Trends
Genet 22, 597–602. https://doi.org/10.1016/j.tig.2006.09.003
Pegueroles, C., Laurie, S., Albà, M.M., 2013. Accelerated Evolution after Gene Duplication: A
Time-Dependent Process Affecting Just One Copy. Molecular Biology and Evolution 30,
1830–1842. https://doi.org/10.1093/molbev/mst083
Peltier, J.-B., Ripoll, D.R., Friso, G., Rudella, A., Cai, Y., Ytterberg, J., Giacomelli, L., Pillardy,
J., Wijk, K.J. van, 2004. Clp Protease Complexes from Photosynthetic and Non-
photosynthetic Plastids and Mitochondria of Plants, Their Predicted Three-dimensional
Structures, and Functional Implications. J. Biol. Chem. 279, 4768–4781.
https://doi.org/10.1074/jbc.M309212200

Porankiewicz, J., Wang, J., Clarke, A.K., 1999. New insights into the ATP-dependent Clp
protease: Escherichia coli and beyond. Molecular Microbiology 32, 449–458.
https://doi.org/10.1046/j.1365-2958.1999.01357.x
Renny-Byfield, S., Wendel, J.F., 2014. Doubling down on genomes: polyploidy and crop plants.
Am J Bot 101, 1711–1725. https://doi.org/10.3732/ajb.1400119
Rizzon, C., Ponger, L., Gaut, B.S., 2006. Striking Similarities in the Genomic Distribution of
Tandemly Arrayed Genes in Arabidopsis and Rice. PLOS Computational Biology 2,
e115. https://doi.org/10.1371/journal.pcbi.0020115
Rockenbach, K., Havird, J.C., Monroe, J.G., Triant, D.A., Taylor, D.R., Sloan, D.B., 2016.
Positive Selection in Rapidly Evolving Plastid–Nuclear Enzyme Complexes. Genetics
204, 1507–1522. https://doi.org/10.1534/genetics.116.188268
Rousseau-Gueutin, M., Huang, X., Higginson, E., Ayliffe, M., Day, A., Timmis, J.N., 2013.
Potential Functional Replacement of the Plastidic Acetyl-CoA Carboxylase Subunit
(accD) Gene by Recent Transfers to the Nucleus in Some Angiosperm Lineages. Plant
Physiology 161, 1918–1929. https://doi.org/10.1104/pp.113.214528
Rudella, A., Friso, G., Alonso, J.M., Ecker, J.R., Wijk, K.J. van, 2006. Downregulation of ClpR2
Leads to Reduced Accumulation of the ClpPRS Protease Complex and Defects in
Chloroplast Biogenesis in Arabidopsis. The Plant Cell 18, 1704–1721.
https://doi.org/10.1105/tpc.106.042861
Salie, M.J., Thelen, J.J., 2016. Regulation and structure of the heteromeric acetyl-CoA
carboxylase. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of
Lipids, Plant Lipid Biology 1861, 1207–1213.
https://doi.org/10.1016/j.bbalip.2016.04.004
Sasaki, Y., Nagano, Y., 2004. Plant Acetyl-CoA Carboxylase: Structure, Biosynthesis,
Regulation, and Gene Manipulation for Plant Breeding. Bioscience, Biotechnology, and
Biochemistry 68, 1175–1184. https://doi.org/10.1271/bbb.68.1175
Schulte, W., Töpfer, R., Stracke, R., Schell, J., Martini, N., 1997. Multi-functional acetyl-CoA
carboxylase from Brassica napus is encoded by a multi-gene family: Indication for
plastidic localization of at least one isoform. PNAS 94, 3465–3470.
https://doi.org/10.1073/pnas.94.7.3465
Shultz, J.L., Kurunam, D., Shopinski, K., Iqbal, M.J., Kazi, S., Zobrist, K., Bashir, R., Yaegashi,
S., Lavu, N., Afzal, A.J., Yesudas, C.R., Kassem, M.A., Wu, C., Zhang, H.B., Town,
C.D., Meksem, K., Lightfoot, D.A., 2006. The Soybean Genome Database (SoyGD): a
browser for display of duplicated, polyploid, regions and sequence tagged sites on the
integrated physical and genetic maps of Glycine max. Nucleic Acids Res 34, D758-765.
https://doi.org/10.1093/nar/gkj050
Sikosek, T., Chan, H.S., Bornberg-Bauer, E., 2012. Escape from Adaptive Conflict follows from
weak functional trade-offs and mutational robustness. Proc Natl Acad Sci U S A 109,
14888–14893. https://doi.org/10.1073/pnas.1115620109
Sinkler, C.A., Kalpage, H., Shay, J., Lee, I., Malek, M.H., Grossman, L.I., Hüttemann, M., 2017.
Tissue- and Condition-Specific Isoforms of Mammalian Cytochrome c Oxidase Subunits:
From Function to Human Disease [WWW Document]. Oxidative Medicine and Cellular
Longevity. https://doi.org/10.1155/2017/1534056
Sjögren, L.L.E., Clarke, A.K., 2011. Assembly of the chloroplast ATP-dependent Clp protease in
Arabidopsis is regulated by the ClpT accessory proteins. Plant Cell 23, 322–332.
https://doi.org/10.1105/tpc.110.082321

Sjögren, L.L.E., Stanne, T.M., Zheng, B., Sutinen, S., Clarke, A.K., 2006. Structural and Functional Insights into the Chloroplast ATP-Dependent Clp Protease in Arabidopsis. The Plant Cell 18, 2635–2649. https://doi.org/10.1105/tpc.106.044594

Sloan, D.B., Triant, D.A., Forrester, N.J., Bergner, L.M., Wu, M., Taylor, D.R., 2014. A recurring syndrome of accelerated plastid genome evolution in the angiosperm tribe Sileneae (Caryophyllaceae). Molecular Phylogenetics and Evolution 72, 82–89. https://doi.org/10.1016/j.ympev.2013.12.004

Smith, M.D., Wertheim, J.O., Weaver, S., Murrell, B., Scheffler, K., Kosakovsky Pond, S.L., 2015. Less Is More: An Adaptive Branch-Site Random Effects Model for Efficient Detection of Episodic Diversifying Selection. Molecular Biology and Evolution 32, 1342–1353. https://doi.org/10.1093/molbev/msv022

Soltis, D.E., Albert, V.A., Leebens-Mack, J., Bell, C.D., Paterson, A.H., Zheng, C., Sankoff, D., Depamphilis, C.W., Wall, P.K., Soltis, P.S., 2009. Polyploidy and angiosperm diversification. Am J Bot 96, 336–348. https://doi.org/10.3732/ajb.0800079

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30, 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Stanne, T.M., Pojidaeva, E., Andersson, F.I., Clarke, A.K., 2007. Distinctive Types of ATP-dependent Clp Proteases in Cyanobacteria. J. Biol. Chem. 282, 14394–14402. https://doi.org/10.1074/jbc.M700275200

Taylor, J.S., Raes, J., 2004. Duplication and Divergence: The Evolution of New Genes and Old Ideas. Annual Review of Genetics 38, 615–643. https://doi.org/10.1146/annurev.genet.38.072902.092831

The Angiosperm Phylogeny Group, Chase, M.W., Christenhusz, M.J.M., Fay, M.F., Byng, J.W., Judd, W.S., Soltis, D.E., Mabberley, D.J., Sennikov, A.N., Soltis, P.S., Stevens, P.F., 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Botanical Journal of the Linnean Society 181, 1–20. https://doi.org/10.1111/boj.12385

Tuskan, G.A., Difazio, S., Jansson, S., Bohlmann, J., Grigoriev, I., Hellsten, U., Putnam, N., Ralph, S., Rombauts, S., Salamov, A., Schein, J., Sterck, L., Aerts, A., Bhalerao, R.R., Bhalerao, R.P., Blaudez, D., Boerjan, W., Brun, A., Brunner, A., Busov, V., Campbell, M., Carlson, J., Chalot, M., Chapman, J., Chen, G.-L., Cooper, D., Coutinho, P.M., Couturier, J., Covert, S., Cronk, Q., Cunningham, R., Davis, J., Degroeve, S., Déjardin, A., Depamphilis, C., Detter, J., Dirks, B., Dubchak, I., Duplessis, S., Ehlting, J., Ellis, B., Gendler, K., Goodstein, D., Gribskov, M., Grimwood, J., Groover, A., Gunter, L., Hamberger, B., Heinze, B., Helariutta, Y., Henrissat, B., Holligan, D., Holt, R., Huang, W., Islam-Faridi, N., Jones, S., Jones-Rhoades, M., Jorgensen, R., Joshi, C., Kangasjärvi, J., Karlsson, J., Kelleher, C., Kirkpatrick, R., Kirst, M., Kohler, A., Kalluri, U., Larimer, F., Leebens-Mack, J., Leplé, J.-C., Locascio, P., Lou, Y., Lucas, S., Martin, F., Montanini, B., Napoli, C., Nelson, D.R., Nelson, C., Nieminen, K., Nilsson, O., Pereda, V., Peter, G., Philippe, R., Pilate, G., Poliakov, A., Razumovskaya, J., Richardson, P., Rinaldi, C., Ritland, K., Rouzé, P., Ryaboy, D., Schmutz, J., Schrader, J., Segerman, B., Shin, H., Siddiqui, A., Sterky, F., Terry, A., Tsai, C.-J., Uberbacher, E., Unneberg, P., Vahala, J., Wall, K., Wessler, S., Yang, G., Yin, T., Douglas, C., Marra, M., Sandberg, G., Van de Peer, Y., Rokhsar, D., 2006. The genome of black cottonwood, Populus

934   trichocarpa (Torr. & Gray). Science 313, 1596–1604.
935   https://doi.org/10.1126/science.1128691
936 Van de Peer, Y., Taylor, J.S., Braasch, I., Meyer, A., 2001. The Ghost of Selection Past: Rates of
937   Evolution and Functional Divergence of Anciently Duplicated Genes. J Mol Evol 53,
938   436–446. https://doi.org/10.1007/s002390010233
939 Wang, W., Zheng, H., Fan, C., Li, J., Shi, J., Cai, Z., Zhang, G., Liu, D., Zhang, J., Vang, S., Lu,
940   Z., Wong, G.K.-S., Long, M., Wang, J., 2006. High Rate of Chimeric Gene Origination
941   by Retroposition in Plant Genomes. The Plant Cell 18, 1791–1802.
942   https://doi.org/10.1105/tpc.106.041905
943 Welsch, R., Zhou, X., Yuan, H., Álvarez, D., Sun, T., Schlossarek, D., Yang, Y., Shen, G.,
944   Zhang, H., Rodriguez-Concepcion, M., Thannhauser, T.W., Li, L., 2018. Clp Protease
945   and OR Directly Control the Proteostasis of Phytoene Synthase, the Crucial Enzyme for
946   Carotenoid Biosynthesis in Arabidopsis. Mol Plant 11, 149–162.
947   https://doi.org/10.1016/j.molp.2017.11.003
948 Wendel, J.F., Lisch, D., Hu, G., Mason, A.S., 2018. The long and short of doubling down:
949   polyploidy, epigenetics, and the temporal dynamics of genome fractionation. Curr Opin
950   Genet Dev 49, 1–7. https://doi.org/10.1016/j.gde.2018.01.004
951 Wertheim, J.O., Murrell, B., Smith, M.D., Kosakovsky Pond, S.L., Scheffler, K., 2015. RELAX:
952   Detecting Relaxed Selection in a Phylogenetic Framework. Molecular Biology and
953   Evolution 32, 820–832. https://doi.org/10.1093/molbev/msu400
954 Wheeler, D.L., Church, D.M., Federhen, S., Lash, A.E., Madden, T.L., Pontius, J.U., Schuler,
955   G.D., Schriml, L.M., Sequeira, E., Tatusova, T.A., Wagner, L., 2003. Database resources
956   of the National Center for Biotechnology. Nucleic Acids Res 31, 28–33.
957 Wicke, S., Schneeweiss, G.M., dePamphilis, C.W., Müller, K.F., Quandt, D., 2011. The
958   evolution of the plastid chromosome in land plants: gene content, gene order, gene
959   function. Plant Mol Biol 76, 273–297. https://doi.org/10.1007/s11103-011-9762-4
960 Williams, A.M., Friso, G., Wijk, K.J. van, Sloan, D.B., 2019. Extreme variation in rates of
961   evolution in the plastid Clp protease complex. The Plant Journal 98, 243–259.
962   https://doi.org/10.1111/tpj.14208
963 Williams, A.M., Itgen, M.W., Broz, A.K., Carter, O.G., Sloan, D.B., 2021. Long-read
964   transcriptome and other genomic resources for the angiosperm Silene noctiflora. G3
965   Genes|Genomes|Genetics. https://doi.org/10.1093/g3journal/jkab189
966 Williams, A.V., Boykin, L.M., Howell, K.A., Nevill, P.G., Small, I., 2015. The Complete
967   Sequence of the Acacia ligulata Chloroplast Genome Reveals a Highly Divergent clpP1
968   Gene. PLOS ONE 10, e0125768. https://doi.org/10.1371/journal.pone.0125768
969 Yang, Z., 2007. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Mol Biol Evol 24,
970   1586–1591. https://doi.org/10.1093/molbev/msm088
971 Yang, Z., Nielsen, R., 2002. Codon-Substitution Models for Detecting Molecular Adaptation at
972   Individual Sites Along Specific Lineages. Molecular Biology and Evolution 19, 908–917.
973   https://doi.org/10.1093/oxfordjournals.molbev.a004148
974 Yu, A.Y.H., Houry, W.A., 2007. ClpP: A distinctive family of cylindrical energy-dependent
975   serine proteases. FEBS Letters 581, 3749–3757.
976   https://doi.org/10.1016/j.febslet.2007.04.076
977 Zhang, J., 2003. Evolution by gene duplication: an update. Trends in Ecology & Evolution 18,
978   292–298. https://doi.org/10.1016/S0169-5347(03)00033-8

979   Zhang, P., Gu, Z., Li, W.-H., 2003. Different evolutionary patterns between young duplicate
980   genes in the human genome. Genome Biology 4, R56. https://doi.org/10.1186/gb-2003-4-
981   9-r56
982   Zhang, Y., Ma, J., Yang, B., Li, R., Zhu, W., Sun, L., Tian, J., Zhang, L., 2014. The complete
983   chloroplast genome sequence of Taxus chinensis var. mairei (Taxaceae): loss of an
984   inverted repeat region and comparative analysis with related species. Gene 540, 201–209.
985   https://doi.org/10.1016/j.gene.2014.02.037
986

Table 1: Differences in evolutionary rates between duplicated and non-duplicated plastid Clp core subunits. Reported *p*-values are based on likelihood ratio tests for 2-partition vs. 3-partition PAML models (see Materials and Methods). Log-likelihood (lnL) values are reported for each model.

Subunit	lnL 2-partition model	lnL 3-partition model	*p*-value	Class with higher d_N/d_S
CLPP3	-10058.01	-10047.93	7.12e-06	Duplicated
CLPP4	-9606.41	-9552.26	<1.00e-10	Duplicated
CLPP5	-8121.80	-8070.90	<1.00e-10	Duplicated
CLPP6	-7697.62	-7691.31	3.82e-04	Non-duplicated
CLPR1	-14534.75	-14517.85	6.07e-09	Duplicated
CLPR2	-12018.69	-12002.74	1.63e-08	Duplicated
CLPR3	-10556.37	-10556.05	0.42	n/a
CLPR4	-10395.60	-10337.35	<1.00e-10	Duplicated

Table 2: *p*-values for asymmetries between paralogs and between paralogs and their common ancestor.

Species	Gene	Paralog 1 vs. paralog 2	Paralog 1 vs. ancestor	Paralog 2 vs. ancestor	Paralogs 1+2 vs. ancestor
P. maritima	*CLPP3*	0.83			1.25e-04*
S. max	*CLPP3*	1			0.20
P. maritima	*CLPP4*	1.28e-07	1.07e-09*	0.71	
M. acuminata	*CLPP4*	0.04	0.15	2.12e-04*	
S. max	*CLPP4*	0.57			1
P. trichocarpa	*CLPP4*	3.31e-04	3.44e-12*	1.76e-13*	
G. maderense	*CLPP4*	0.08			1.72e-03*
M. truncatula	*CLPP5*	1			3.61e-05*
P. trichocarpa	*CLPP5*	1			0.04^
O. biennis	*CLPP5*	9.12e-3	0.10	5.37e-4*	
G. maderense	*CLPP5*	1.71e-04	0.42	6.23e-4*	
S. max	*CLPP6*	0.27			1
P. trichocarpa	*CLPP6*	0.46			1
G. raimondii	*CLPP6*	0.71			1
M. acuminata	*CLPR1*	2.20e-16	0.82	2.20e-16*	
S. max	*CLPR1*	0.47			0.05^
P. trichocarpa	*CLPR1*	0.77			0.42
V. vinifera	*CLPR1*	4.45e-3	0.17	1.38e-09*	
M. guttatus	*CLPR1*	0.05	0.59	4.70e-04*	
P. maritima	*CLPR1*	4.70e-05	0.53	2.54e-05*	
P. maritima	*CLPR2*	0.01	0.18	1.63e-06*	
S. max	*CLPR3*	1			0.55
P. trichocarpa	*CLPR3*	0.34			0.35
M. truncatula	*CLPR4*	2.29e-03	0.21	0.11	
E. grandis	*CLPR4*	2.51e-05	0.68	5.42e-11*	
P. maritima	*CLPR4*	9.73e-05	9.53e-04	4.17e-14*	

* denotes that paralog(s) has/have significantly higher evolutionary rate than ancestor branch
^ denotes that paralog(s) has/have significantly lower evolutionary rate than ancestor branch

Figure 1: Expectations under different post-duplication models. For each pair of paralogs (n = 26), we first determined whether they were evolving symmetrically relative to one another using N and S estimates from PAML output. Paralogs evolving asymmetrically are predicted to represent neofunctionalization or pseudogenization events. For paralogs evolving symmetrically (n = 8), combined N and S values were compared to those of the immediate ancestor branch. Pairs evolving symmetrically relative to the common ancestor (n = 8) are predicted to represent gene dosage advantage while those evolving asymmetrically relative to the common ancestor (n = 5) are predicted to represent subfunctionalization.

Figure 2: *ACC* genes across the Brassicaceae, Caryophyllaceae, Geraniaceae, and Poaceae, with the single copy of *ACC* in *Amborella trichopoda* as an outgroup. Branch lengths represent d_N values and branch colors represent d_N/d_S ratios.

Figure 3: Copy numbers of the nuclear-encoded subunits of the plastid Clp core across angiosperms. Boxes without numbers indicate single-copy genes.

Figure 4: d_N/d_S ratios of duplicated and non-duplicated plastid Clp core subunits across angiosperms. The values were calculated using a PAML branch test with three groups, where each group was assigned its own d_N/d_S value: non-duplicated terminal branches, duplicated terminal branches (and in the cases of *CLPP5* and *CLPR4*, internal post-duplication branches), and internal branches. Significant differences (p<<0.001) are indicated with ***.

Table S1: Sites inferred to be under positive selection in *ACC2* branches based on a branch-sites test in PAML using the trimmed alignment for *ACC*.

Site	Amino acid	Probability
6	L	1.000
92	G	0.965
125	T	0.998
153	V	0.967
182	L	0.983
200	V	0.982
212	L	0.951
333	E	0.993
387	E	0.952
425	E	0.999
428	S	0.982
429	L	0.999
475	S	1.000
480	R	0.968
533	T	0.968
539	S	0.991
546	V	0.977
562	V	0.989
597	L	0.994
652	L	0.999
660	H	0.982
663	M	0.995
687	R	0.961
693	H	0.996
699	L	0.991
700	G	1.000
713	F	0.999
715	A	0.973
739	L	0.995
740	N	0.999
744	S	0.993
753	Q	0.980
766	D	0.967
769	N	0.990
776	K	0.986

794	L	0.967
797	G	0.996
837	R	0.997
849	S	0.996
859	Q	0.986
865	R	1.000
868	L	0.999
872	K	0.995
922	T	0.979
948	Q	0.985
973	T	0.984
981	T	0.999
982	P	0.999
985	K	0.998
989	N	0.999
991	R	0.997
1015	P	0.999
1027	R	0.999
1042	Q	0.991
1043	W	1.000
1044	H	1.000
1045	R	0.995
1048	L	0.978
1078	E	0.995
1085	W	0.980
1096	L	0.984
1108	T	0.999
1110	H	0.958
1147	M	0.990
1151	Q	0.994
1160	Q	0.998
1161	E	0.957
1168	K	1.000
1177	S	0.999
1197	R	0.998
1200	M	0.986
1212	Y	0.989
1243	A	0.995
1326	A	0.963

1342	I	0.981
1344	R	0.983
1507	S	0.952
1580	K	0.956
1602	R	0.971
1655	S	0.966
1735	L	1.000
1887	V	0.984
1898	A	0.999
1904	Q	0.985
1928	E	0.982
2021	P	0.958
2027	S	0.976
2063	E	1.000
2131	K	0.998
2135	E	0.962
2137	A	0.998
2164	G	0.965
2194	E	1.000
2230	P	1.000
2235	Q	0.992
2241	R	0.985
2245	G	0.976

Table S2: Loss of catalytic sites and truncation of nuclear-encoded plastid ClpP core subunits

Species	Protein	Serine	Histidine	Aspartate	Length
Eucalyptus grandis	ClpP3		Replaced with R		
Musa acuminata	ClpP3		Replaced with R		
Plantago maritima 1	ClpP3	Replaced with Y	Replaced with R	Replaced with N	
Plantago maritima 2	ClpP3	Replaced with Y	Replaced with R	Replaced with N	
Lathyrus sativus	ClpP4		Replaced with T		
Medicago truncatula	ClpP4		Replaced with A		
Musa acuminata 2	ClpP4	Gap	Replaced with R	Gap	Truncated
Plantago maritima 2	ClpP4			Gap	
Populus trichocarpa 2	ClpP4	Replaced with M	Gap	Gap	Truncated
Gossypium raimondii 3	ClpP5	Gap			
Gossypium raimondii 4	ClpP5				Truncated
Gossypium raimondii 5	ClpP5	Replaced with N			Truncated
Gossipium Raimondii 6	ClpP5	Replaced with N		Replaced with N	Truncated
Gossypium raimondii 7	ClpP5	Replaced with N		Replaced with F	
Plantago maritima	ClpP5	Gap	Gap	Gap	Truncated
Lathyrus sativus	ClpP6		Replaced with G		
Lobelia siphilitica	ClpP6	Replaced with N			
Medicago truncatula	ClpP6		Replaced with N		
Plantago maritima	ClpP6	Replaced with G	Replaced with E	Replaced with F	
Populus trichocarpa 2	ClpP6				Truncated

*Empty cell indicates presence of catalytic site or full length.

Table S3: Truncation of nuclear-encoded plastid ClpR core subunits

Truncated ClpR1 Subunits
Vitis vinifera 2
Populus trichocarpa 2
Musa acuminata 2 (internal stop codon)
Mimulus guttatus 2
Truncated ClpR2 Subunits
Plantago maritima 2
Truncated ClpR3 Subunits
Populus trichocarpa 1
Populus trichocarpa 2
Truncated ClpR4 Subunits
Eucalyptus grandis 2
Medicago truncatula 2
Vitis vinifera 2
Vitis vinifera 3
Vitis vinifera 4
Musa acuminata 3
Musa acuminata 4

Figure S1. *ACC* tree. Branch labels are d_N/d_S values. *ACC1* represents cytosolic-targeted genes while *ACC2* represents plastid-targeted genes.

Figure S2. *ACC* tree. A) Branch labels are d_N values. B) Branch labels are d_S values.

Figure S3. *CLPP3* tree. Branch labels are d_N/d_S values.

Figure S4. *CLPP4* tree. Branch labels are d_N/d_S values.

Figure S5. *CLPP5* tree. Branch labels are d_N/d_S values.

Figure S6. *CLPP6* tree. Branch labels are d_N/d_S values.

Figure S7. *CLPR1* tree. Branch labels are d_N/d_S values.

Figure S8. *CLPR2* tree. Branch labels are d_N/d_S values.

Figure S9. *CLPR3* tree. Branch labels are d_N/d_S values.

Figure S10. *CLPR4* tree. Branch labels are d_N/d_S values.