Therapeutic efficacy of an oral nucleoside analog of remdesivir against SARS-CoV-2 pathogenesis in mice.

Alexandra Schäfer [1, 4], David R. Martinez [1, 4], John J. Won [1], Fernando R. Moreira [1], Ariane J. Brown [1], Kendra L. Gully [1], Rao Kalla [2], Kwon Chun [2], Venice Du Pont [2], Darius Babusis [2], Jennifer Tang [2], Eisuke Murakami [2], Raju Subramanian [2], Kimberly T Barrett [2], Blake J. Bleier [2], Roy Bannister [2], Joy Y. Feng [2], John P. Bilello [2], Tomas Cihlar [2], Richard L. Mackman [2], Stephanie A. Montgomery [3], Ralph S. Baric [1], Timothy P. Sheahan [1, *]

[1] Department of Epidemiology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA,

[2] Gilead Sciences, Inc, Foster City, CA, USA,

[3] Department of Pathology and Laboratory Medicine, University of North Carolina School of Medicine, Chapel Hill, NC, USA,

[4] These authors contributed equally to this manuscript.

[*] Timothy P. Sheahan. sheahan@email.unc.edu

Abstract

The COVID-19 pandemic remains uncontrolled despite the rapid rollout of safe and

effective SARS-CoV-2 vaccines, underscoring the need to develop highly effective antivirals. In

the setting of waning immunity from infection and vaccination, breakthrough infections are

becoming increasingly common and treatment options remain limited. Additionally, the

emergence of SARS-CoV-2 variants of concern with their potential to escape therapeutic

monoclonal antibodies emphasizes the need to develop second-generation oral antivirals

targeting highly conserved viral proteins that can be rapidly deployed to outpatients. Here, we

demonstrate the in vitro antiviral activity and in vivo therapeutic efficacy of GS-621763, an

orally bioavailable prodrug of GS-441524, the parental nucleoside of remdesivir, which targets

the highly conserved RNA-dependent RNA polymerase. GS-621763 exhibited significant

antiviral activity in lung cell lines and two different human primary lung cell culture systems.

The dose-proportional pharmacokinetic profile observed after oral administration of GS-621763

translated to dose-dependent antiviral activity in mice infected with SARS-CoV-2. Therapeutic

GS-621763 significantly reduced viral load, lung pathology, and improved pulmonary function

in COVID-19 mouse model. A direct comparison of GS-621763 with molnupiravir, an oral

nucleoside analog antiviral currently in human clinical trial, proved both drugs to be similarly

efficacious. These data demonstrate that therapy with oral prodrugs of remdesivir can

significantly improve outcomes in SARS-CoV-2 infected mice. Thus, GS-621763 supports the

exploration of GS-441524 oral prodrugs for the treatment of COVID-19 in humans.

Introduction

SARS-CoV-2 emerged in December 2019 and has caused 223 million infections and 4.6

million deaths worldwide as of September 2021 (*1-3*). While there are multiple effective

vaccines, vaccination rates have lagged in the United States (U.S.) due to vaccine hesitancy and

public mistrust thus delaying the generation of herd immunity required to significantly diminish

community spread. In addition, outside of the U.S., many countries do not have equitable access

to vaccines and/or have been slow to vaccinate (*4-7*). This collective constellation of events is

fueling the generation of viral variants that are increasingly transmissible and are ever evolving

to escape human immunity. Therefore, there is an immediate unmet need for oral antivirals that

can be rapidly disseminated to treat COVID-19 cases in the unvaccinated, the

immunocompromised and in vaccine breakthrough cases. Next-generation oral coronavirus

(CoV) antivirals, if widely disseminated and given early in infection, could curtail the duration of

disease, reduce long-term sequelae of COVID-19, minimize household transmissions, and lessen

hospitalizations, thus having a broad impact on public health.

There are multiple direct-acting antiviral (DAA) therapies in use to treat COVID-19 (*8-*

12), including Emergency Use Authorization (EUA)-approved monoclonal antibodies (mAbs)

and FDA-approved remdesivir (RDV, GS-5734). Monoclonal antibodies have demonstrated

efficacy for treating active COVID-19 cases in outpatients (Regeneron outpatient studies) but

currently, all mAbs must be administered via injection, limiting their use to those with ready

access to healthcare (*13, 14*). In addition, several SARS-CoV-2 variants of concern (VOCs) have

evolved that are resistant to first-line mAb therapies (*15, 16*). Currently, RDV is the only FDA-

approved small-molecule direct-acting antiviral to treat COVID-19, but there are several other

DAAs currently in human clinical trials including nucleoside analogs molnupiravir (MPV,

EIDD-2801) and AT-527, as well as MPro inhibitor PF-07321332 (*17-25*) . Unlike mAbs which

specifically target the virion surface exposed spike protein of SARS-CoV-2, nucleoside analog

drugs target a highly conserved viral enzyme among CoV, the RNA-dependent RNA polymerase

(RdRp) nsp12, rendering them broadly active against multiple emerging, endemic and enzootic

CoV. Moreover, due to its high degree of conservation among CoV, the RdRp likely does not

have the same capacity for mutational change as spike, which may translate into RdRp having a

higher barrier to resistance (*18-20*). Despite demonstrated therapeutic efficacy of RDV against

SARS-CoV-2 in animal models (*17, 23, 26*) and in human clinical trials (*8*), the requirement of

intravenous administration has limited its widespread use during this pandemic. The orally

bioavailable nucleoside prodrug GS-621763, is designed for optimal delivery of the parent

nucleoside GS-441524 into systemic circulation, which is then metabolized inside cells into the

same active nucleoside triphosphate formed by RDV (*27*). Here, we detail the *in vitro* antiviral

activity in various cell models and *in vivo* therapeutic efficacy of oral GS-621763 in a mouse

model of SARS-CoV-2 pathogenesis.

Results

GS-621763 has antiviral activity against SARS-CoV-2 in cell lines and human primary cell

cultures.

85   GS-441524 is the parental adenosine nucleoside analog (Fig. 1A) of both

monophosphoramidate prodrug RDV (GS-5734, Fig. 1B), and triester prodrug GS-621763 (Fig.

1C). All three molecules are metabolized to the same active nucleotide triphosphate in cells, but

through different activation pathways. GS-621763 is rapidly metabolized during oral absorption

to GS-441524, then intracellularly converted by cellular kinases to the analog monophosphate

metabolite before further metabolism to the active nucleoside triphosphate. In contrast, the intact

phosphoramidate prodrug, RDV, is broken down inside cells directly to the same monophosphate

metabolite, effectively bypassing the rate-limiting first phosphorylation step of GS-441524 (*27*).

To determine if GS-621763 could inhibit replication of SARS-CoV-2 in cellular assays, we first

evaluated its antiviral activity against a SARS-CoV-2 reporter virus expressing nanoluciferase

(SARS-CoV-2 nLUC) in A549-hACE2 cells stably expressing the human entry receptor

angiotensin-converting enzyme 2 (ACE2) (*28*). With GS-621763, we observed a dose-dependent

antiviral effect on SARS-CoV-2 nLUC replication with an average half-maximum effective

concentration (EC_{50}) of 2.8 µM (Fig. 1D, Fig.1H, and Supplementary Figure 1A). In the same

assay, we measured EC_{50} values for the control compound RDV of 0.28 µM, similar to those

reported previously in these cells and reflective of the enhanced ability of the phosphoramidate

prodrug to rapidly and efficiently generate active triphosphate by bypassing the slower initial

phosphorylation step (Fig. 1D, Fig. 1H, Supplementary Figure 1A) (*29*). As was observed in

other cell systems, the parental nucleoside, GS-441524, was less potent (EC_{50} = 3.3 µM) than

RDV in our assay and was similar in potency to GS-621763. This suggests that the tri-isobutyryl

esters of GS-621763 are efficiently cleaved in the assay to release GS-441524 (Fig. 1D, Fig. 1H,

Supplementary Figure 1A). Importantly, we did not observe any measurable cytotoxicity of any

of the inhibitors in A549-hACE2 cells at concentrations up to 10 µM (Figure 1F, Supplementary

Figure 1B). Human primary airway epithelial (HAE) cell cultures model the cellular complexity

and architecture of the human conducting airway and are often used to determine if drugs are

transported and metabolized in the cells targeted by emerging CoV in vivo (*18*). In HAE cells

infected with WT SARS-CoV-2 (*28*) and treated with GS-621763, we observed a dose-

dependent and significant reduction in infectious virus production as compared to DMSO-

vehicle treated cultures (Fig. 1F). In similarly infected, control compound (i.e. RDV or GS-

441524) treated cultures, a significant and dose-dependent reduction in viral titers was also

observed (Fig. 1F). GS-621763, RDV, and GS-441524 inhibited reporter SARS-CoV-2

expressing Firefly luciferase (SARS-CoV-2 Fluc) replication in normal human bronchial

epithelial (NHBE) cultures with EC_{50} values of 0.125, 0.0371, and 2.454 µM, respectively (Fig.

1G, Fig. 1H). All together, these data show that GS-621763 is transported, metabolized and

potently antiviral human primary cell systems that model the tissues targeted by SARS-CoV-2 in

humans.

Dose-dependent therapeutic efficacy of GS-621763 in mouse models of COVID-19 disease.

123   We have previously performed multiple studies describing the therapeutic efficacy of

subcutaneously administered RDV in mice (*Ces1c*$^{-/-}$ C57BL/6J) genetically deleted for a secreted

plasma carboxylesterase 1c (*Ces1c*) absent in humans but dramatically reduces drug half-life in

wild-type mice (*17-19, 26, 30*).

However, the prodrug GS-621763 is designed to be rapidly cleaved pre-systemically in

vivo to release GS-441524 into circulation, with no or very minimal intact ester observed in

plasma. Therefore, GS-621763 can be studied in wild-type mice where it should also be rapidly

converted to parent GS-441524. Plasma pharmacokinetics following a single oral administration

of GS-621763 at either 5 or 20 mg/kg were first determined in uninfected BALB/c mice (Fig.

2A). Doses were selected to provide high plasma exposures of GS-441524 that would support

active triphosphate formation in the lung and to confirm pharmacokinetic dose proportionality

needed to project exposures in efficacy studies. Previous studies had shown that parent

nucleoside was at least 10-fold less efficient at generating lung triphosphate than RDV, on a

molar basis, thus requiring higher plasma exposures of parental GS-441524 to account for the

reduced metabolic efficiency (27). No exposure of intact ester prodrug, within the limit of

detection, was observed in mice. GS-441524 was both rapidly absorbed and then cleared from

systemic circulation, exhibiting a short plasma half-life of approximately 1 hr. Dose

proportional increases in both maximal plasma concentrations (C_{max}) and exposures (AUC_{0-24h})

at the two doses were observed (Fig. 2A).

To better understand the pharmacokinetic and pharmacodynamic relationship for GS-

621763, we performed a series of dose-finding studies in BALB/c mice infected with mouse-

adapted SARS-CoV-2 (SARS-CoV-2 MA10) (30). In young adult BALB/c mice infected with

10^4 plaque forming units (PFU) SARS-CoV-2 MA10, virus replicates to high titers in the

respiratory tract, mice lose 15-20% of their body weight by 4 days post-infection (dpi), and acute

lung injury/loss of pulmonary function is typically observed after virus replication peaks on 2 dpi

(30). We first defined the minimum dosage sufficient for maximal therapeutic efficacy in

BALB/c mice initiating twice daily (i.e. bis in die, BID) oral treatment with either vehicle control

or 3 mg/kg, 10 mg/kg, or 30 mg/kg GS-621763 beginning 8 hours post infection (hpi) with 10^4

PFU SARS-CoV-2 MA10 (Fig. 2B). Unlike vehicle or 3 mg/kg GS-621763 treated animals,

mice receiving either 10 or 30 mg/kg GS-621763 were completely protected from weight loss

thus demonstrating that early oral antiviral therapy can prevent the progression of disease (Fig.

2B). Congruent with the weight loss phenotype, both 10 and 30 mg/kg GS-621763 treated

animals had significantly reduced viral lung titers as compared to both the vehicle and 3 mg/kg

treated groups (Fig. 2C). To monitor the effect of drug treatment on pulmonary function, we

performed daily whole-body plethysmography (WBP) with a subset of mice from each group

(N=4 per treatment group). As shown with the WBP metric PenH, whose elevation is associated

with airway resistance or obstruction (*18*), we observed a drug dose-dependent reduction in

PenH with the maximal effect seen in the 30 mg/kg GS-621763 dose group which was

completely protected from the loss of pulmonary function observed in the other treatment groups

and vehicle (Fig. 2D). Mice treated with 3 and 10 mg/kg GS-621763 had impaired lung function

at days 2 and 3 post infection, but lung function returned to baseline by 4 dpi for all GS-621763

treated animals (Fig. 2D). Consistent with weight loss, virus titer, and pulmonary function data,

mice treated with 10 or 30 mg/kg had significantly reduced lung congestion, a gross pathologic

feature characteristic of severe lung damage (Fig. 2E). We then scored lung tissue sections for

the histologic features of acute lung injury (ALI) using two complementary semiquantitative

tools. First, using an ALI scoring tool created by the American Thoracic Society (ATS), we

blindly evaluated three diseased fields per lung section for several features of ALI including

alveolar septal thickening, neutrophils in the interstitium and in air spaces, proteinaceous debris

in airspaces, and the presence of hyaline membranes. Only mice treated with 30 mg/kg had

significantly reduced ALI scores (Fig. 2F). Second, we used a complementary tool measuring

the pathologic hallmark of ALI, diffuse alveolar damage (DAD). Mice in all treated groups

showed reduced DAD scores, but only mice receiving 30 mg/kg had significantly decreased

DAD in their lungs (Fig. 2G). Together, these data demonstrate that the oral delivery of the

nucleoside analog GS-621763 can significantly diminish SARS-CoV-2 virus replication and

associated pulmonary disease in a dose-dependent manner.

Extended therapeutic protection against COVID-19 disease by GS-621763 in mice

To determine if the potent therapeutic efficacy of GS-621763 observed with early

intervention (8 hr after infection) would extend to later times post infection, we designed a

therapeutic efficacy study with six arms where we varied both time of oral therapy initiation and

dose level in BALB/c mice infected with SARS-CoV-2 MA10 (Fig.3). As done previously, a

control group of animals received vehicle twice daily beginning at 12 hours post infection (hpi).

The next three arms of the study were dedicated to the 30 mg/kg GS-621763 dose level, with two

of the three arms receiving twice daily dosing initiated at either the 12 hpi ("30 mg/kg BID 12

hr" group) or the 24 hpi ("30 mg/kg BID 24 hr" group). The third 30 mg/kg arm was designed to

determine if dose frequency could be reduced to once daily (quaque die, QD) if initiated early at

12 hpi ("30 mg/kg QD 12 hr" group). In the last two arms, we wanted to evaluate if an increased

dose of 60 mg/kg given QD beginning at 12 hr or 24 hr ("60 mg/kg QD 12 hr" and "60 mg/kg

QD 24 hr" groups) would improve outcomes over the 30 mg/kg groups. Initiation of 30 mg/kg

BID therapy at either 12 or 24 hrs offered significant protection from weight loss (Fig. 3A),

extending the robust therapeutic phenotype observed for this dose level when initiated at very

early times (at 8 hr) (Fig 2). Interestingly, when we decreased the frequency of 30 mg/kg

treatment initiated at 12 hr to once daily ("30 mg/kg QD 12 hr" group), we also observed a

significant prevention of body weight loss (Fig 3A), thus levels of drug when administered once

a day and begun early (at 12 hr) in the course of infection were sufficient to prevent disease

progression. Increasing the dose to 60 mg/kg QD initiated at either 12 hr or 24 hr offered similar

protection from weight loss observed with vehicle treatment as the 30 mg/kg groups (Fig. 3A).

200   As body weight loss is a crude marker of viral pathogenesis, we next measured multiple

virological, physiologic, and pathologic metrics of disease. First, we measured the levels of

infectious virus present in lung tissue on 4 dpi. Unlike vehicle-treated animals who harbored an

average titer of 5.6×10^4 PFU per lung lobe, all GS-621763 dose groups significantly reduced the

levels of infectious virus in lung tissue with the average titers of most groups falling below the

limit of detection (90 PFU). Interestingly, 30 mg/kg delivered QD had significantly elevated

viral lung titers (mean titer $= 4 \times 10^2$ PFU) as compared to its BID counterpart (mean titer $= < 90$

PFU). A similar trend among treatment groups was observed when measuring the levels of

SARS-CoV-2 subgenomic and genomic nucleocapsid (N) RNA in parallel lung tissues (Fig. 3C).

All 30 mg/kg groups significantly reduced levels of SARS-CoV-2 RNA in lung tissue as

compared to vehicle-treated animals. As observed for infectious titers, 30 mg/kg given QD daily

beginning at 12 hr had elevated levels of N RNA as compared to 30 mg/kg given BID beginning

at 12 hr suggesting that trough and/or daily exposure levels of drug with QD dosing are

insufficient to suppress replication similarly to BID dosing. Increasing the once-daily dose to 60

mg/kg to raise the daily exposure and trough levels offered similar reductions in SARS-CoV-2 N

RNA as compared to 30 mg/kg BID when initiated at 12 hr, but the levels of viral RNA in the

higher dose 60mg/kg group initiated at 24 hr were not different than vehicle (Fig. 3C). Although

vehicle-treated animals exhibited significant loss of pulmonary function as measured by WBP,

this was largely prevented with GS-621763 therapy (Fig. 3D). All therapy groups initiated at 12

hr were equally protected from loss of pulmonary function as measured by the PenH metric (Fig.

3D). Of all groups actively receiving GS-621763, animals in the 30 mg/kg BID 24 hr group had

a measurable loss of lung function on 3 dpi which resolved by 4 dpi but this phenotype was not

extended to other groups. We then scored lung tissue sections for the histologic features of acute

lung injury and alveolar damage. Only 30 mg/kg BID initiated at 12 hr significantly reduced ALI

scores as compared to those in vehicle-treated animals. In addition, the 30 mg/kg BID 12 hr-

group had significantly lower ALI scores as compared to the 60 mg/kg QD 24 hr-group (Figure

3E). Only 30 mg/kg given twice a day at 12 hr most dramatically reduced DAD scores but this

protection from lung pathology was lost if given once per day or if initiated at 24 hr (Fig. 3G).

The high dose of 60 mg/kg QD when initiated at 12 hr improved DAD scores as compared to

similarly treated animals that began treatment at 24 hr. Collectively, these data demonstrate that

GS-621763 therapy can improve both virologic and pathogenic metrics, but the degree of

improvement was dependent on time of initiation and dose frequency.

The therapeutic efficacy of GS-621763 is similar to molnupiravir (MPV, EIDD-2801)

MPV is an oral nucleoside analog prodrug antiviral currently in Phase 3 clinical trial to

treat COVID-19 with demonstrated antiviral efficacy in mice against several emerging CoV

including SARS-CoV, MERS-CoV, and SARS-CoV-2 (*18, 19, 21, 22*). Like GS-621763, MPV

is a prodrug which is metabolized in vivo into a parental nucleoside (β-D-N4-hydroxycytidine,

NHC) in its metabolic progression towards the antiviral active triphosphate (*20*). To determine if

GS-621763 would provide similar protection as MPV, we then designed comparative therapeutic

efficacy studies in the mouse model of SARS-CoV-2 pathogenesis described above. Pre-efficacy

pharmacokinetic studies in BALB/c mice (30 mg/kg or 100 mg/kg) were performed with MPV

and showed dose proportional increases in NHC plasma exposures (Supplemental Fig. 2).

Pharmacokinetic modeling then determined that a daily 120 mg/kg dose (given 60 mg/kg BID)

would result in exposures similar to that observed in humans receiving 800 mg BID, a dose

being evaluated in a human clinical trial (22). The comparative efficacy study included a vehicle

group and 5 additional groups receiving two doses of MPV or GS-621763 per day 12 hrs apart

(BID). Three arms of the study began dosing at 12 hr: 30 mg/kg GS-621763, 30 mg/kg MPV

(0.5× human equivalent dose) or 60 mg/kg MPV (1× human equivalent dose). At 24 hr, we

began dosing of two additional groups: 60 mg/kg GS-621763 or 60 mg/kg MPV. While SARS-

CoV-2 MA10 infection caused rapid weight loss in vehicle control animals, all animals receiving

either GS-621763 or MPV beginning at either 12 or 24 hr were protected from weight loss (Fig.

4A). Similarly, upon titration of lung tissues at 4 dpi for infectious virus by plaque assay,

vehicle-treated animals had expectedly high levels of infectious virus which was significantly

reduced in all treatment groups, independent of drug type or initiation time (Fig. 4B). When

treatment was initiated at 12 hr, a moderate yet significant elevation in infectious titers was

observed in the 30mg/kg MPV group, inferior to either equivalently dosed GS-621763 animals

or those receiving the higher dose (60 mg/kg) of MPV (Fig. 4B). To understand the relationship

between levels of infectious virus and viral RNA in lung tissue, we performed qRT-PCR on total

RNA for SARS-CoV2 N RNA in parallel tissues utilized for plaque assay. The trend observed

with infectious virus is mirrored in the qRT-PCR data where all groups receiving antiviral

therapy had significantly reduced levels of viral RNA (Fig. 4C). In addition, animals receiving

30 mg/kg MPV (0.5× human equivalent dose) had a measurable increase in N RNA as compared

to equivalently dosed GS-621763 animals. Similar to weight loss data, vehicle-treated animals

had a significant loss of pulmonary function as measured by WBP on both 3 and 4 dpi which was

prevented in all groups receiving antiviral treatment (Fig. 4D). We then blindly evaluated lung

tissue sections for the pathological manifestations of ALI and DAD using two complementary

histologic tools described above. Congruent with the above data, ALI scores in all antiviral

therapy groups were significantly reduced as compared to vehicle controls (Fig. 4E). In

agreement with ALI scores, the DAD histologic scores were similarly reduced in all antiviral

therapy treated groups as compared to those treated with vehicle (Fig. 4F). All together, these

data show that antiviral therapy with GS-621763 and MPV when initiated early or at the peak of

virus replication (~24 hr) can both significantly diminish virus replication and improve disease

outcomes.

Discussion

Three novel human CoVs have emerged in the past 20 years, first with SARS-CoV in

2002-2003, MERS-CoV in 2012 and most recently, SARS-CoV-2 in 2019 (*1, 2, 31*). Vaccine

availability, vaccine hesitancy, and the ongoing evolution and emergence of VOCs are

collectively delaying the global control of pandemics and potential achievement of global herd

immunity and may prevent it altogether (*15, 32-34*). As such, there is an acute need for broad-

spectrum antivirals to treat COVID-19 in the unvaccinated as well as increasingly common

breakthrough infections in those vaccinated, driven by immune evading VOCs. In addition, due

to the emergence potential of the CoV family, we must also actively develop broadly acting

therapies effective the CoVs of today to prepare for those that may emerge in the future. Both

RDV and MPV are examples of nucleoside analogs with broad activity against genetically

diverse CoV (*35*). Here, we show that GS-621763, a prodrug that improves the oral delivery of

parent nucleoside GS-441524, is yet another example of an orally bioavailable nucleoside analog

prodrug that is effective against SARS-CoV-2.

Oral broadly acting antiviral therapies that target conserved viral proteins with a

diminished capacity for evolutionary change will maximize therapeutic utility against future

emerging CoV. SARS-CoV-2 has undergone a considerable amount of genetic evolution since

its emergence in 2019; each time a globally dominant SARS-CoV-2 VOC has emerged, it has

been replaced by a new VOC harboring more concerning characteristics like replicative capacity

or transmissibility (*36*). The majority of genetic changes have been localized to viral proteins

decorating the surface of the virus particle like the viral spike which has mutated to eradicate

epitopes targeted by early and promising monoclonal antiviral therapies (e.g. Eli Lily,

Regeneron) rendering them less active against newer VOCs (*37*). Approved antivirals like RDV,

those in clinical trials like MPV, AT-527, and PF-07321332 (Pfizer, oral protease inhibitor)

target highly conserved enzymes required for virus replication which likely have a diminished

capacity for change as compared to the spike protein (*38*). The widespread use of both RDV and

mAb therapies in the U.S. and globally has been limited by the necessity of delivery by

intravenous infusion which in turn requires access to qualified health care staff and facilities.

Thus, effective oral antiviral therapy or combination therapies which can be procured at a

pharmacy and self-administered by the patient would help facilitate wide-spread global access

and could have profound positive impacts on global public health.

In this study, we utilized a mouse adapted SARS-CoV-2 variant, SARS-CoV-2 MA10, in

wild-type BALB/c mice. (*17, 30*). Mice infected with this virus develop severe lung disease

reminiscent of that seen with severe COVID-19 including the development of ALI and

respiratory failure (*30*). It is important to note that the disease resulting from SARS-CoV-2

MA10 infection in mice is compressed as compared to that observed in humans with virus titer

peaking in the mouse lung between 24-48 hrs after infection, rapid loss of pulmonary function

beginning 2-3 dpi, rapid weight loss within the first 4 days of infection, lung pathology

consistent with ALI peaking 4-6 dpi and virus induced mortality within a week of infection. This

is markedly different than the COVID-19 in humans where virus titers peak in the upper airway

within the first week, but viral RNA shedding can be observed for as long as 24 days and

symptoms can take weeks to months to resolve (*39-41*). Because of this caveat associated with

our mouse model, the time in which to intervene with a direct acting antiviral and sufficiently

improve outcomes is curtailed in mice as compared to humans. Our recent study with RDV

exemplifies this where we found the degree of therapeutic benefit in mice infected with SARS-

CoV-2 MA10 was dependent on the time of initiation (*17*). Here, we show that oral

administration with GS-621763 prevents body weight loss, loss of pulmonary function, severe

lung pathology and virus replication when administered at 12 and 24 hrs after infection. While

we observed improvement in some metrics with RDV initiated at 24 hr in our prior studies (*17*),

GS-621763 therapy initiated at a similar time comparatively improved all metrics assessed.

Thus, herein we provide proof-of-concept preclinical data that the orally bioavailable ester

analog of RDV GS-621763 can exert potent antiviral effects in vivo during an ongoing SARS-

CoV-2 infection. Lastly, we show that GS-621763 therapy provides similar levels of protection

from SARS-CoV-2 pathogenesis as MPV, an oral nucleoside analog prodrug effective against

SARS-CoV-2 in mice that is currently in human clinical trials. Future directions are focused on

extending these studies to evaluate the efficacy of combinations of antivirals in our models of

SARS-CoV-2 pathogenesis and in other models that can evaluate the blockade of transmission

such as hamster and ferret.

In summary, we provide preclinical data demonstrating the in vitro antiviral activity and

in vivo therapeutic efficacy of an orally bioavailable nucleoside analog prodrug, GS-621763. The

data provided herein supports the future evaluation of orally bioavailable prodrugs of GS-441524

in humans with COVID-19. If safe and effective, this class of RdRp inhibitors could become part

of the arsenal of existing oral antivirals that are desperately needed to address a global unmet

need for the COVID-19 pandemic and CoV pandemics of the future.

Figure Legends

Figure 1. Chemical Structure and in vitro potency of GS-621763 in comparison to RDV

(GS-5734) and GS-441524.

(A) Chemical structure of the parental adenosine nucleoside analog GS-441524.

(B) Chemical structure of the monophosphoramidate prodrug RDV.

(C) Chemical structure of GS-621763, the tri-isobutyryl ester of GS-441524.

(D) Mean percent inhibition of SARS-CoV-2 replication by GS-621763, in comparison to the

prodrug RDV - and the parental nucleoside GS-441524 in A459-hACE2 cells (done in

triplicates).

(E) Cytotoxicity in A459-hACE2 cells treated with GS-621763, RDV, and GS-441524 in A459-

hACE2 cells (done in triplicates, CellTiter-Glo – CTG).

(F) Inhibition of SARS-CoV-2 replication by GS-621763, in comparison to the prodrug RDV

and the parental nucleoside GS-441524 in human primary airway epithelial cells (HAE, done in

duplicates).

(G) Inhibition of SARS-CoV-2-Fluc replication by GS-621763, RDV -, and the parental

nucleoside GS-441524 in normal human bronchial epithelial (NHBE) cultures (done in

duplicates).

(H) In vitro EC_{50} values for inhibition of viral replication by GS-621763, RDV, and the parental

nucleoside GS-441524 in A459-hACE2 and NHBE cells.

Figure 2. Dose-dependent therapeutic protection against COVID-19 disease by GS-621763

in mice.

(A) Plasma Pharmacokinetics of GS-441524 in uninfected Balb/c mice following a single oral

administration of GS-621763 at either 5 or 20 mg/kg. Plasma concentrations of GS-621763 were

below the limit of quantification at all time points.

(B) Depicted is the % starting weight in therapeutically treated mice with vehicle (n=19) or 3

mg/kg (n=9), 10 mg/kg (n=10), and 30 mg/kg (n=10) GS-621763 at 8 hr. All mice were infected

with $1x10^4$ PFU SARS-CoV-2 MA10.

(C) Lung viral titers in therapeutically treated mice with vehicle (n=19) or 3 mg/kg (n=9), 10

mg/kg (n=10), and 30 mg/kg (n=10) GS-621763 at 8 hr. All mice were infected with $1x10^4$ PFU

SARS-CoV-2 MA10. Limit of detection (LoD)

(D) Pulmonary function in therapeutically treated mice with vehicle (n=4) or 3 mg/kg (n=4), 10

mg/kg (n=4), and 30 mg/kg (n=4) GS-621763 at 8 hr. All mice were infected with $1x10^4$ PFU

SARS-CoV-2 MA10.

(E) Lung congestion score in therapeutically treated mice with vehicle (n=19) or 3 mg/kg (n=9),

10 mg/kg (n=10), and 30 mg/kg (n=10) GS-621763 at 8 hr. All mice were infected with $1x10^4$

PFU SARS-CoV-2 MA10.

(F and G) Lung pathology in the therapeutically treated mice with vehicle (n=19) or 3 mg/kg

(n=9), 10 mg/kg (n=10), and 30 mg/kg (n=10) GS-621763 at 8 hr. All mice were infected with

1×10^4 PFU SARS-CoV-2 MA10.

Data were analyzed using two-way ANOVA (weight loss and lung function) and Kruskal-Wallis

test (lung titer, congestion score, and pathology scores), *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$,

****$p < 0.0001$

Figure 3. Extended therapeutic protection of mice against COVID-19 disease by oral GS-

621763 in mice.

(A) Depicted is the % starting weight in therapeutically treated mice with either 30 mg/kg or 60

mg/kg BID or QD GS-621763 at 12 and 24 hrs (n=10 for all treatment groups, except n=8 for 60

mg/kg, 12 hr QD treatment group). All mice were infected with 1×10^4 PFU SARS-CoV-2

MA10.

(B) Lung viral titers in therapeutically treated mice with either 30 mg/kg or 60 mg/kg BID or QD

GS-621763 at 12 and 24hrs (n=10 for all treatment groups, except n=8 for 60 mg/kg, 12 hr QD

treatment group). All mice were infected with 1×10^4 PFU SARS-CoV-2 MA10. Limit of

detection (LoD).

(C) Viral N RNA in therapeutically treated mice with either 30 mg/kg or 60 mg/kg BID or QD

GS-621763 at 12 and 24 hrs (n=10 for all treatment groups, except n=8 for 60 mg/kg, 12 hr QD

treatment group). All mice were infected with 1×10^4 PFU SARS-CoV-2 MA10.

(D) Pulmonary function in therapeutically treated mice with either 30 mg/kg or 60 mg/kg BID or

QD GS-621763 at 12 and 24 hrs (n=4 for all treatment groups). All mice were infected with

1×10^4 PFU SARS-CoV-2 MA10.

(n=4 for all treatment groups).

(F and G) Lung pathology in the therapeutically treated mice with either 30 mg/kg or 60 mg/kg

BID or QD GS-621763 at 12 and 24 hrs (n=10 for all treatment groups, except n=8 for 60 mg/kg,

12 hr QD treatment group, and n=7 for vehicle group).

Data were analyzed using two-way ANOVA (weight loss and lung function) and Kruskal-Wallis

test (lung titer, lung viral RNA, and pathology scores), $*p<0.05$, $**p<0.005$, $***p<0.0005$,

$****p<0.0001$.

Figure 4. Evaluation of therapeutic intervention of GS-621763 in comparison to

molnupiravir (MPV).

(A) Depicted is the % starting weight in therapeutically treated mice with either 30 mg/kg or 60

mg/kg GS-621763 or MPV at 12 or 24 hrs (n=10 for all treatment groups). All mice were

infected with $1x10^4$ PFU SARS-CoV-2 MA10.

(B) Lung viral titers in therapeutically treated mice with either 30 mg/kg or 60 mg/kg GS-

621763 or MPV at 12 or 24 hrs (n=10 for all treatment groups). All mice were infected with

$1x10^4$ PFU SARS-CoV-2 MA10. Limit of detection (LoD).

(C) Viral N RNA in therapeutically treated mice with either 30 mg/kg or 60 mg/kg GS-621763

or MPV at 12 or 24 hrs (n=10 for all treatment groups). All mice were infected with $1x10^4$ PFU

SARS-CoV-2 MA10.

(D) Pulmonary function in therapeutically treated mice with either 30 mg/kg or 60mg/kg GS-

621763 or MPV at 12 or 24 hrs (n=4 for all treatment groups). All mice were infected with $1x10^4$

PFU SARS-CoV-2 MA10.

(F and G) Lung pathology in the therapeutically treated mice with either 30 mg/kg or 60 mg/kg

GS-621763 or MPV at 12or 24 hrs (n=10 for all treatment groups). All mice were infected with

$1x10^4$ PFU SARS-CoV-2 MA10.

Data were analyzed using two-way ANOVA (weight loss and lung function) and Kruskal-Wallis

test (lung titer, lung viral RNA, and pathology scores), $*p<0.05$, $**p<0.005$, $***p<0.0005$,

$****p<0.0001$

Supplemental Figure 1. In vitro potency and toxicity of GS-621763, RDV, and GS-441524

in A549-hACE2 cells

(A) Raw data for the inhibition of SARS-CoV-2 replication by GS-621763, RDV, and GS-

441521 in A459-hACE2 cells measured through quantitation of SARS-CoV-2 expressed nano

luciferase (nLuc), measured in triplicates.

(B) Raw data for cytotoxicity in A459-hACE2 cells treated with GS-621763, RDV, and GS-

441521 in A459-hACE2 cells measured via CellTiter-Glo, (measured in triplicates).

C) Raw data for the inhibition of SARS-CoV-2-Fluc replication by GS-621763, RDV, and GS-

441521 in NHBE cultures measured through quantitation of SARS-CoV-2 expressed firefly

luciferase (Fluc), measured in duplicates, repeated twice.

Supplemental Figure 2. Molnupiravir mouse plasma pharmacokinetics

Plasma Pharmacokinetics of N-hydroxycytidine (NHC) in uninfected Balb/c mice following

daily oral administration of molnupiravir at either 60 or 200 mg/kg (as either 30 or 100

mg/kg/dose given BID; molnupiravir at all timepoints).

Material and Methods

Small molecule drug synthesis and formulation

GS-621763, RDV, and GS-441524 were synthesized at Gilead Sciences Inc., and their chemical

identity and purity were determined by nuclear magnetic resonance, and high-performance

liquid chromatography (HPLC) analysis (*27*). Molnupiravir was purchased from

MedChemExpress LLC (NJ, USA) with a purity of 95% based on HPLC analysis.

Small molecules were solubilized in 100% DMSO for in vitro studies and in vehicle

containing 2.5% DMSO; 10% Kolliphor HS-15; 10% Labrasol; 2.5% Propylene glycol; 75%

Water (final formulation pH 2) (for GS-621763) and in vehicle containing 2.5% Kolliphor RH-

40, 10% Polyethylene glycol 300, 87.5% Water (for MPV) for in vivo studies. GS-621763, GS-

441524, GS-5734 were made available to the University of North Carolina (UNC) at Chapel Hill

under a materials transfer agreement with Gilead Sciences.

In vivo plasma pharmacokinetic analysis of GS-621763 and molnupiravir (MPV)

Mice were orally administered either a single dose of GS-621763 (in vehicle containing

2.5% DMSO; 10% Kolliphor HS-15; 10% Labrasol; 2.5% Propylene glycol; 75% Water (final

formulation pH 2) or two doses of molnupiravir (in vehicle containing 2.5% Kolliphor RH-40,

10% Polyethylene glycol 300, 87.5% Water) (BID, 12 hours apart). GS-621763 was given at

either 5 or 20 mg/kg and MPV at either 30 or 100 mg/kg. Plasma was serially isolated from 4

mice at 0.25, 1, 2, 8 and 24 hrs post GS-621763 administration. Plasma was isolated from

alternating groups of 4 mice per timepoint at 0.5, 2, 6, 12 (pre-second dose), 12.5, 18 and 24

hrs post MPV administration. 20 µl of plasma was added to a mixture containing 250 µl of

methanol and 25 µL of internal standard solution and centrifuged. 250 µl of resulting

supernatant was then transferred, filtered (Agilent Captiva 96, 0.2 μm) and dried under a stream

of nitrogen at 40 °C. Following reconstitution in a mixture of 5% acetonitrile and 95% water, a

10 μl aliquot was injected onto an LC-MS/MS system. Plasma concentrations of either GS-

621763 and GS-441524 or MPV and N-hydroxycytidine (NHC) were determined using 8 to 10-

point calibration curves spanning at least 3 orders of magnitude with quality control samples to

ensure accuracy and precision, prepared in normal mouse plasma. Analytes were separated by a

50 mm × 3.0 mm, 2.55 μm Synergi Polar-RP column (Phenomenex) using a multi-stage linear

gradient from 5% to 95% acetonitrile in mobile phase A at a flow rate of 1 ml/min.

Quantitation of GS-441524 metabolites in the lung following oral GS-621763

administration in *Balb/c* mice

Lungs from all mice administered GS-621763 were quickly isolated at 24 hrs post-dose

and immediately snap frozen in liquid nitrogen. On dry ice, frozen lung samples were

pulverized and weighed. Dry ice-cold extraction buffer containing 0.1% potassium hydroxide

and 67 mM ethylenediamine tetraacetic acid (EDTA) in 70% methanol, containing 0.5μM

chloro-adenosine triphosphate as internal standard was added and homogenized. After

centrifugation at 20,000 × g for 20 minutes, supernatants were transferred and dried in a

centrifuging evaporator. Dried samples were then reconstituted with 60 μL of mobile phase A,

containing 3 mM ammonium formate (pH 5) with 10 mM dimethylhexylamine (DMH) in

water, centrifuged at 20,000 × g for 20 minutes and final supernatants transferred to HPLC

injection vials. An aliquot of 10 μl was subsequently injected onto an API 6500 LC/MS/MS

system for analysis of GS-441524 and its phosphorylated metabolites, performed using a

similar method as described previously (*18*).

Viruses and plaque assay

493   Recombinant SARS-CoV-2 MA10 virus was generated as described previously (*30*). For

494   virus titration by plaque assay, the caudal lobe of the right lung was homogenized in PBS, and

495   the resulting homogenate was serial-diluted and inoculated onto confluent monolayers of Vero

496   E6 cells, followed by agarose overlay. Plaques were visualized with overlay of neutral red dye

497   on day 3 after infection (*30*).

In vitro assays for antiviral activity

A549-hACE2 cells were plated at a density of 20,000 cells/well/100 µl in black-walled

clear-bottom 96-well plates 24 hrs prior to infection. Compounds GS-621763, GS-5734, GS-

441524, were diluted in 100% DMSO (1:3) resulting in a 1000X dose response from 10 to 0.002

mM (10 to 0.002 µM final). All conditions were performed in triplicate. At BSL3, medium was

removed, and cells were infected with 100 µl SARS-CoV-2 nLUC (MOI 0.008) for 1 h at 37 ˚C

after which virus was removed, wells were washed (150 µl) with infection media (DMEM, 4%

FBS, 1X antibiotic/antimycotic) and infection media (100 µl) containing a dose response of drug

was added. Plates were incubated at 37 ˚C for 48 hrs. NanoGlo assay was performed 48 hpi.

Sister plates were exposed to drug but not infected to gauge cytotoxicity via CellTiter-Glo assay

(CTG, Promega, Madison, WI), 48 hrs post treatment.

Normal human bronchial epithelial (NHBE) cells (donor 41219) were purchased from

Lonza (Walkersville, MD Cat# CC-2540) and maintained in Bronchial Epithelial Cell Growth

Medium (BEGM) (Lonza, Walkersville, MD, Cat# CC-3170) with all provided supplements in

the BulletKit. Cells are passaged 2-3 times per week to maintain sub-confluent densities and are

used for experiments at passages 2-4. NHBE cells were seeded in 24-well plates at 1×10^5 cells in

a final volume of 0.5 ml BEGM^{TM} Bronchial Epithelial Cell Growth Medium (BulletKit^{TM};

Lonza, Basel, SW). Cultures were incubated overnight 37 °C with 5% CO_2. On the following

day, media was replaced with 0.5 ml growth medium. Cultures were treated with 1:3 serial

dilutions of compound using the HP D300e digital dispenser with normalization to the highest

concentration of DMSO in all wells (<1% final volume). The cells were then infected with 0.1

ml SARS-CoV-2-Fluc diluted in BEGM media at MOI = 5. Uninfected and untreated wells were

included as controls to determine compound efficacy against SARS-CoV-2-Fluc. Following

incubation with compound and virus for 24 hrs at 37 °C with 5% CO_2, culture supernatant was

removed from each well and replaced with 0.3 ml of ONE-Glo luciferase reagent (Promega,

Madison, WI). The plates were shaken at 400 rpm for 10 min at room temperature. 0.2 ml of

supernatant from each well was transferred to a 96-well opaque plate (Corning) and

luminescence signal was measured using an EnVision plate reader (PerkinElmer). Values were

normalized to the uninfected and infected DMSO controls (0% and 100% infection,

respectively). Data was fit using a four-parameter non-linear regression analysis using Graphpad

Prism. EC_{50} values were then determined as the concentration reducing the firefly luciferase

signal by 50%. The compiled data was generated based on least two independent experimental

replicates, each containing four technical replicates for each concentration.

Mouse studies and in vivo infections

All mouse studies were performed at the University of North Carolina (Animal Welfare

Assurance #A3410-01) using protocols (#20-059) approved by the University of North Carolina

Institutional Animal Care and Use Committee. All animal work was approved by the

Institutional Animal Care and Use Committee at University of North Carolina at Chapel Hill

according to guidelines outlined by the Association for the Assessment and Accreditation of

Laboratory Animal Care and the US Department of Agriculture. All work was performed with

approved standard operating procedures and safety conditions for SARS-CoV-2. Our

institutional BSL3 facilities are designed to conform to the safety requirements recommended by

Biosafety in Microbiological and Biomedical Laboratories, the US Department of Health and

Human Services, the Public Health Service, the Centers for Disease Control and Prevention, and

the National Institutes of Health. Laboratory safety plans have been submitted, and the facility

has been approved for use by the University of North Carolina Department of Environmental

Health and Safety and the Centers for Disease Control and Prevention.

546   Even groups (n=10, or less as indicated) of 10-week-old female BALB/c mice (Envigo;

#047) were used in all in vivo efficacy studies. For infection, mice were anesthetized with a

mixture of ketamine/xylazine and infected with 10^4 PFU of SARS-CoV-2 MA10 in 50 µl PBS

intranasally. Vehicle or GS-621673 was administered orally at the dosages and timepoints as

indicated. Mice were monitored daily for body weight changes and for lung function by whole-

body plethysmography. At 4 dpi, mice were euthanized, and lung tissue was harvested for viral

titer analysis, RNA and histology, and lung congestion scores were estimated (*30*). Samples for

viral load determination and for RNA isolation were stored at −80 °C until used; histology

samples were inactivated in 10% NBF and stored at 4 °C until further processing.

Histology and lung pathology scoring

557   Two separate lung pathology scoring scales, Matute-Bello and Diffuse Alveolar Damage

558   (DAD), were used to quantify acute lung injury (ALI) (*19*).

559   For Matute-Bello scoring samples were blinded and three random fields of lung tissue

were chosen and scored for the following: (A) neutrophils in alveolar space (none = 0, 1–5 cells =

1, > 5 cells = 2), (B) neutrophils in interstitial space (none = 0, 1–5 cells = 1, > 5 cells = 2), (C)

hyaline membranes (none = 0, one membrane = 1, > 1 membrane = 2), (D) Proteinaceous debris

in air spaces (none = 0, one instance = 1, > 1 instance = 2), (E) alveolar septal thickening (< 2Å~

mock thickness = 0, 2–4Å~ mock thickness = 1, > 4Å~ mock thickness = 2). Scores from A–E

were put into the following formula score = [(20x A) + (14 x B) + (7 x C) + (7 x D) + (2 x E)]/100

to obtain a lung injury score per field and then averaged for the final score for that sample.

In a similar way, for DAD scoring, three random fields of lung tissue were scored for the

in a blinded manner for: 1= absence of cellular sloughing and necrosis, 2= uncommon solitary cell

sloughing and necrosis (1–2 foci/field), 3=multifocal (3+foci) cellular sloughing and necrosis with

uncommon septal wall hyalinization, or 4=multifocal (>75% of field) cellular sloughing and

necrosis with common and/or prominent hyaline membranes. To obtain the final DAD score per

mouse, the scores for the three fields per mouse were averaged.

RNA isolation and RT-qPCR

Mouse tissue from SARS-CoV-2 infected mice was homogenized using glass beads in

TRIzol Reagent (Invitrogen). Equal volume of 100% EtOH was mixed with the TRIzol

homogenate and processed using Direct-Zol RNA MiniPrep Kit (Zymo) to extract viral RNA.

Optional DNase I treatment was conducted to ensure the adequate removal of unwanted DNA.

Eluted RNA was coupled with TaqMan Fast Virus 1-step Master Mix (Applied Biosystems) and

nCOV_N1 primers/probe (IDT) to quantify viral load via reverse transcription – quantitative

polymerase chain reaction (RT-qPCR). Samples were plated on a MicroAmp EnduraPlate

(Applied Biosystems) and run using a QuantStudio 6 Real-Time PCR System (Applied

Biosystems) to obtain viral titers. The following PCR program was run: 50 °C for 5 minutes, 95

°C for 20 seconds, followed by 45 cycles of 95 °C for 3 seconds and 60°C for 30 seconds. The

sequences of the 2019-nCOV_N1 primers and probe were as follows: Forward primer: GAC

CCC AAA ATC AGC GAA AT, Reverse primer: TCT GGT TAC TGC CAG TTG AAT CTG,

Probe: AC CCC GCA TT ACG TTT GGT GGA CC (CDC N1 qRT-PCR assay (*42*). SARS-

CoV-2 standard curve RNA was produced by PCR amplification of SARS-CoV-2 nucleocapsid

by which a 5' T7 polymerase promoter was introduced. This amplicon was used as template to

generate in vitro transcribed RNA which was then quantified and serially diluted (108 – 101

copies/µl).

Acknowledgement

D.R.M. is funded by a Burroughs Wellcome Fund Postdoctoral Enrichment Program Award and a Hanna

H. Gray Fellowship from the Howard Hughes Medical Institute and was supported by an NIH NIAID T32

AI007151 and an NIH F32 AI152296. This work was supported by NIAID R01 AI132178 and

R01AI132178-04S1 (T.P.S. and R.S.B.) and an NIH animal models contract (HHSN272201700036I) to

R.S.B. This project was supported in part by the North Carolina Policy Collaboratory at University of

North Carolina at Chapel Hill with funding from the North Carolina Coronavirus Relief Fund established

and appropriated by the North Carolina General Assembly.

Animal histopathology services were performed by the Animal Histopathology & Laboratory Medicine

Core at the University of North Carolina, which is supported in part by an NCI Center Core Support

Grant (5P30CA016086-41) to the UNC Lineberger Comprehensive Cancer Center.

Competing interest

These authors are employees of Gilead Sciences and hold stock in Gilead Sciences: Rao Kalla,

Kwon Chun, Venice Du Pont, Darius Babusis, Jennifer Tang, Eisuke Murakami, Raju

Subramanian, Kimberly T Barrett, Blake J. Bleier, Roy Bannister, Joy Y. Feng, John P. Bilello,

Tomas Cihlar, and Richard L. Mackman

Literature

1.   P. Zhou, X. L. Yang, X. G. Wang, B. Hu, L. Zhang, W. Zhang, H. R. Si, Y. Zhu, B. Li,
C. L. Huang, H. D. Chen, J. Chen, Y. Luo, H. Guo, R. D. Jiang, M. Q. Liu, Y. Chen, X.
R. Shen, X. Wang, X. S. Zheng, K. Zhao, Q. J. Chen, F. Deng, L. L. Liu, B. Yan, F. X.
Zhan, Y. Y. Wang, G. F. Xiao, Z. L. Shi, A pneumonia outbreak associated with a new
coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).
2.   N. Zhu, D. Zhang, W. Wang, X. Li, B. Yang, J. Song, X. Zhao, B. Huang, W. Shi, R. Lu,
P. Niu, F. Zhan, X. Ma, D. Wang, W. Xu, G. Wu, G. F. Gao, W. Tan, I. China Novel
Coronavirus, T. Research, A Novel Coronavirus from Patients with Pneumonia in China,
2019. *N Engl J Med* **382**, 727-733 (2020).
3.   J. H. University. (2020).
4.   A. Binagwaho, K. Mathewos, S. Davis, Time for the ethical management of COVID-19
vaccines. *Lancet Glob Health*, (2021).
5.   E. Aryeetey, E. Engebretsen, A. Gornitzka, P. Maassen, S. Stolen, A step backwards in
the fight against global vaccine inequities. *Lancet* **397**, 23-24 (2021).
6.   A. Asundi, C. O'Leary, N. Bhadelia, Global COVID-19 vaccine inequity: The scope, the
impact, and the challenges. *Cell Host Microbe* **29**, 1036-1039 (2021).
7.   J. R. Barrett, S. Belij-Rammerstorfer, C. Dold, K. J. Ewer, P. M. Folegatti, C. Gilbride,
R. Halkerston, J. Hill, D. Jenkin, L. Stockdale, M. K. Verheul, P. K. Aley, B. Angus, D.
Bellamy, E. Berrie, S. Bibi, M. Bittaye, M. W. Carroll, B. Cavell, E. A. Clutterbuck, N.
Edwards, A. Flaxman, M. Fuskova, A. Gorringe, B. Hallis, S. Kerridge, A. M. Lawrie, A.
Linder, X. Liu, M. Madhavan, R. Makinson, J. Mellors, A. Minassian, M. Moore, Y.
Mujadidi, E. Plested, I. Poulton, M. N. Ramasamy, H. Robinson, C. S. Rollier, R. Song,
M. D. Snape, R. Tarrant, S. Taylor, K. M. Thomas, M. Voysey, M. E. E. Watson, D.
Wright, A. D. Douglas, C. M. Green, A. V. S. Hill, T. Lambe, S. Gilbert, A. J. Pollard, C.
V. T. G. Oxford, Phase 1/2 trial of SARS-CoV-2 vaccine ChAdOx1 nCoV-19 with a
booster dose induces multifunctional antibody responses. *Nat Med* **27**, 279-288 (2021).
8.   J. Grein, N. Ohmagari, D. Shin, G. Diaz, E. Asperges, A. Castagna, T. Feldt, G. Green,
M. L. Green, F. X. Lescure, E. Nicastri, R. Oda, K. Yo, E. Quiros-Roldan, A.
Studemeister, J. Redinski, S. Ahmed, J. Bernett, D. Chelliah, D. Chen, S. Chihara, S. H.
Cohen, J. Cunningham, A. D'Arminio Monforte, S. Ismail, H. Kato, G. Lapadula, E.
L'Her, T. Maeno, S. Majumder, M. Massari, M. Mora-Rillo, Y. Mutoh, D. Nguyen, E.
Verweij, A. Zoufaly, A. O. Osinusi, A. DeZure, Y. Zhao, L. Zhong, A. Chokkalingam, E.
Elboudwarej, L. Telep, L. Timbs, I. Henne, S. Sellers, H. Cao, S. K. Tan, L.
Winterbourne, P. Desai, R. Mera, A. Gaggar, R. P. Myers, D. M. Brainard, R. Childs, T.
Flanigan, Compassionate Use of Remdesivir for Patients with Severe Covid-19. *N Engl J*
Med **382**, 2327-2336 (2020).

9.   J. H. Beigel, K. M. Tomashek, L. E. Dodd, A. K. Mehta, B. S. Zingman, A. C. Kalil, E.
647   Hohmann, H. Y. Chu, A. Luetkemeyer, S. Kline, D. Lopez de Castilla, R. W. Finberg, K.
648   Dierberg, V. Tapson, L. Hsieh, T. F. Patterson, R. Paredes, D. A. Sweeney, W. R. Short,
649   G. Touloumi, D. C. Lye, N. Ohmagari, M. D. Oh, G. M. Ruiz-Palacios, T. Benfield, G.
650   Fatkenheuer, M. G. Kortepeter, R. L. Atmar, C. B. Creech, J. Lundgren, A. G. Babiker,
651   S. Pett, J. D. Neaton, T. H. Burgess, T. Bonnett, M. Green, M. Makowski, A. Osinusi, S.
652   Nayak, H. C. Lane, A.-S. G. Members, Remdesivir for the Treatment of Covid-19 - Final
653   Report. *N Engl J Med* **383**, 1813-1826 (2020).
654   10.   D. F. Robbiani, C. Gaebler, F. Muecksch, J. C. C. Lorenzi, Z. Wang, A. Cho, M.
655   Agudelo, C. O. Barnes, A. Gazumyan, S. Finkin, T. Hagglof, T. Y. Oliveira, C. Viant, A.
656   Hurley, H. H. Hoffmann, K. G. Millard, R. G. Kost, M. Cipolla, K. Gordon, F. Bianchini,
657   S. T. Chen, V. Ramos, R. Patel, J. Dizon, I. Shimeliovich, P. Mendoza, H. Hartweger, L.
658   Nogueira, M. Pack, J. Horowitz, F. Schmidt, Y. Weisblum, E. Michailidis, A. W.
659   Ashbrook, E. Waltari, J. E. Pak, K. E. Huey-Tubman, N. Koranda, P. R. Hoffman, A. P.
660   West, Jr., C. M. Rice, T. Hatziioannou, P. J. Bjorkman, P. D. Bieniasz, M. Caskey, M. C.
661   Nussenzweig, Convergent antibody responses to SARS-CoV-2 in convalescent
662   individuals. *Nature* **584**, 437-442 (2020).
663   11.   S. J. Zost, P. Gilchuk, R. E. Chen, J. B. Case, J. X. Reidy, A. Trivette, R. S. Nargi, R. E.
664   Sutton, N. Suryadevara, E. C. Chen, E. Binshtein, S. Shrihari, M. Ostrowski, H. Y. Chu,
665   J. E. Didier, K. W. MacRenaris, T. Jones, S. Day, L. Myers, F. Eun-Hyung Lee, D. C.
666   Nguyen, I. Sanz, D. R. Martinez, P. W. Rothlauf, L. M. Bloyet, S. P. J. Whelan, R. S.
667   Baric, L. B. Thackray, M. S. Diamond, R. H. Carnahan, J. E. Crowe, Jr., Rapid isolation
668   and profiling of a diverse panel of human monoclonal antibodies targeting the SARS-
669   CoV-2 spike protein. *Nat Med* **26**, 1422-1427 (2020).
670   12.   D. Pinto, Y. J. Park, M. Beltramello, A. C. Walls, M. A. Tortorici, S. Bianchi, S. Jaconi,
671   K. Culap, F. Zatta, A. De Marco, A. Peter, B. Guarino, R. Spreafico, E. Cameroni, J. B.
672   Case, R. E. Chen, C. Havenar-Daughton, G. Snell, A. Telenti, H. W. Virgin, A.
673   Lanzavecchia, M. S. Diamond, K. Fink, D. Veesler, D. Corti, Cross-neutralization of
674   SARS-CoV-2 by a human monoclonal SARS-CoV antibody. *Nature* **583**, 290-295
675   (2020).
676   13.   A.-T. L.-C. S. Group, J. D. Lundgren, B. Grund, C. E. Barkauskas, T. L. Holland, R. L.
677   Gottlieb, U. Sandkovsky, S. M. Brown, K. U. Knowlton, W. H. Self, D. C. Files, M. K.
678   Jain, T. Benfield, M. E. Bowdish, B. G. Leshnower, J. V. Baker, J. U. Jensen, E. M.
679   Gardner, A. A. Ginde, E. S. Harris, I. S. Johansen, N. Markowitz, M. A. Matthay, L.
680   Ostergaard, C. C. Chang, V. J. Davey, A. Goodman, E. S. Higgs, D. D. Murray, T. A.
681   Murray, R. Paredes, M. K. B. Parmar, A. N. Phillips, C. Reilly, S. Sharma, R. L. Dewar,
682   M. Teitelbaum, D. Wentworth, H. Cao, P. Klekotka, A. G. Babiker, A. C. Gelijns, V. L.
683   Kan, M. N. Polizzotto, B. T. Thompson, H. C. Lane, J. D. Neaton, A Neutralizing
684   Monoclonal Antibody for Hospitalized Patients with Covid-19. *N Engl J Med* **384**, 905-
685   914 (2021).
686   14.   M. P. O'Brien, E. Forleo-Neto, B. J. Musser, F. Isa, K. C. Chan, N. Sarkar, K. J. Bar, R.
687   V. Barnabas, D. H. Barouch, M. S. Cohen, C. B. Hurt, D. R. Burwen, M. A. Marovich, P.
688   Hou, I. Heirman, J. D. Davis, K. C. Turner, D. Ramesh, A. Mahmood, A. T. Hooper, J.
689   D. Hamilton, Y. Kim, L. A. Purcell, A. Baum, C. A. Kyratsous, J. Krainson, R. Perez-
690   Perez, R. Mohseni, B. Kowal, A. T. DiCioccio, N. Stahl, L. Lipsich, N. Braunstein, G.
691   Herman, G. D. Yancopoulos, D. M. Weinreich, T. Covid-19 Phase 3 Prevention Trial,

692   Subcutaneous REGEN-COV Antibody Combination to Prevent Covid-19. *N Engl J Med*,
693   (2021).
694   15.   R. E. Chen, X. Zhang, J. B. Case, E. S. Winkler, Y. Liu, L. A. VanBlargan, J. Liu, J. M.
695   Errico, X. Xie, N. Suryadevara, P. Gilchuk, S. J. Zost, S. Tahan, L. Droit, J. S. Turner,
696   W. Kim, A. J. Schmitz, M. Thapa, D. Wang, A. C. M. Boon, R. M. Presti, J. A.
697   O'Halloran, A. H. J. Kim, P. Deepak, D. Pinto, D. H. Fremont, J. E. Crowe, Jr., D. Corti,
698   H. W. Virgin, A. H. Ellebedy, P. Y. Shi, M. S. Diamond, Resistance of SARS-CoV-2
699   variants to neutralization by monoclonal and serum-derived polyclonal antibodies. *Nat*
700   *Med* **27**, 717-726 (2021).
701   16.   P. Wang, M. S. Nair, L. Liu, S. Iketani, Y. Luo, Y. Guo, M. Wang, J. Yu, B. Zhang, P. D.
702   Kwong, B. S. Graham, J. R. Mascola, J. Y. Chang, M. T. Yin, M. Sobieszczyk, C. A.
703   Kyratsous, L. Shapiro, Z. Sheng, Y. Huang, D. D. Ho, Antibody resistance of SARS-
704   CoV-2 variants B.1.351 and B.1.1.7. *Nature* **593**, 130-135 (2021).
705   17.   D. R. Martinez, A. Schäfer, S. R. Leist, D. Li, K. Gully, B. Yount, J. Y. Feng, E. Bunyan,
706   D. P. Porter, T. Cihlar, S. A. Montgomery, B. F. Haynes, R. S. Baric, M. C.
707   Nussenzweig, T. P. Sheahan, Prevention and therapy of SARS-CoV-2 and the B.1.351
708   variant in mice. *Cell Rep* **36**, 109450 (2021).
709   18.   T. P. Sheahan, A. C. Sims, R. L. Graham, V. D. Menachery, L. E. Gralinski, J. B. Case,
710   S. R. Leist, K. Pyrc, J. Y. Feng, I. Trantcheva, R. Bannister, Y. Park, D. Babusis, M. O.
711   Clarke, R. L. Mackman, J. E. Spahn, C. A. Palmiotti, D. Siegel, A. S. Ray, T. Cihlar, R.
712   Jordan, M. R. Denison, R. S. Baric, Broad-spectrum antiviral GS-5734 inhibits both
713   epidemic and zoonotic coronaviruses. *Sci Transl Med* **9**, (2017).
714   19.   T. P. Sheahan, A. C. Sims, S. R. Leist, A. Schafer, J. Won, A. J. Brown, S. A.
715   Montgomery, A. Hogg, D. Babusis, M. O. Clarke, J. E. Spahn, L. Bauer, S. Sellers, D.
716   Porter, J. Y. Feng, T. Cihlar, R. Jordan, M. R. Denison, R. S. Baric, Comparative
717   therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon
718   beta against MERS-CoV. *Nat Commun* **11**, 222 (2020).
719   20.   T. P. Sheahan, A. C. Sims, S. Zhou, R. L. Graham, A. J. Pruijssers, M. L. Agostini, S. R.
720   Leist, A. Schafer, K. H. Dinnon, 3rd, L. J. Stevens, J. D. Chappell, X. Lu, T. M. Hughes,
721   A. S. George, C. S. Hill, S. A. Montgomery, A. J. Brown, G. R. Bluemling, M. G.
722   Natchus, M. Saindane, A. A. Kolykhalov, G. Painter, J. Harcourt, A. Tamin, N. J.
723   Thornburg, R. Swanstrom, M. R. Denison, R. S. Baric, An orally bioavailable broad-
724   spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and
725   multiple coronaviruses in mice. *Sci Transl Med* **12**, (2020).
726   21.   A. Wahl, L. E. Gralinski, C. E. Johnson, W. Yao, M. Kovarova, K. H. Dinnon, 3rd, H.
727   Liu, V. J. Madden, H. M. Krzystek, C. De, K. K. White, K. Gully, A. Schafer, T. Zaman,
728   S. R. Leist, P. O. Grant, G. R. Bluemling, A. A. Kolykhalov, M. G. Natchus, F. B. Askin,
729   G. Painter, E. P. Browne, C. D. Jones, R. J. Pickles, R. S. Baric, J. V. Garcia, SARS-
730   CoV-2 infection is effectively treated and prevented by EIDD-2801. *Nature* **591**, 451-457
731   (2021).
732   22.   W. Fischer, J. J. Eron, W. Holman, M. S. Cohen, L. Fang, L. J. Szewczyk, T. P. Sheahan,
733   R. Baric, K. R. Mollan, C. R. Wolfe, E. R. Duke, M. M. Azizad, K. Borroto-Esoda, D. A.
734   Wohl, A. J. Loftis, P. Alabanza, F. Lipansky, W. P. Painter, Molnupiravir, an Oral
735   Antiviral Treatment for COVID-19. *medRxiv*, (2021).
736   23.   B. N. Williamson, F. Feldmann, B. Schwarz, K. Meade-White, D. P. Porter, J. Schulz, N.
737   van Doremalen, I. Leighton, C. Kwe Yinda, L. Perez-Perez, A. Okumura, J. Lovaglio, P.

738   W. Hanley, G. Saturday, C. M. Bosio, S. Anzick, K. Barbian, T. Cihlar, C. Martens, D. P.
739   Scott, V. J. Munster, E. de Wit, Clinical benefit of remdesivir in rhesus macaques
740   infected with SARS-CoV-2. *bioRxiv*, (2020).
741 24. D. R. Owen, C. M. N. Allerton, A. S. Anderson, L. Aschenbrenner, M. Avery, S. Berritt,
742   B. Boras, R. D. Cardin, A. Carlo, K. J. Coffman, A. Dantonio, L. Di, H. Eng, R. Ferre, K.
743   S. Gajiwala, S. A. Gibson, S. E. Greasley, B. L. Hurst, E. P. Kadar, A. S. Kalgutkar, J. C.
744   Lee, J. Lee, W. Liu, S. W. Mason, S. Noell, J. J. Novak, R. S. Obach, K. Ogilvie, N. C.
745   Patel, M. Pettersson, D. K. Rai, M. R. Reese, M. F. Sammons, J. G. Sathish, R. S. P.
746   Singh, C. M. Steppan, A. E. Stewart, J. B. Tuttle, L. Updyke, P. R. Verhoest, L. Wei, Q.
747   Yang, Y. Zhu, An Oral SARS-CoV-2 M^{pro} Inhibitor Clinical Candidate for
748   the Treatment of COVID-19. *medRxiv*, 2021.2007.2028.21261232 (2021).
749 25. S. S. Good, J. Westover, K. H. Jung, X. J. Zhou, A. Moussa, P. La Colla, G. Collu, B.
750   Canard, J. P. Sommadossi, AT-527, a Double Prodrug of a Guanosine Nucleotide
751   Analog, Is a Potent Inhibitor of SARS-CoV-2 In Vitro and a Promising Oral Antiviral for
752   Treatment of COVID-19. *Antimicrob Agents Chemother* **65**, (2021).
753 26. A. J. Pruijssers, A. S. George, A. Schafer, S. R. Leist, L. E. Gralinksi, K. H. Dinnon, 3rd,
754   B. L. Yount, M. L. Agostini, L. J. Stevens, J. D. Chappell, X. Lu, T. M. Hughes, K.
755   Gully, D. R. Martinez, A. J. Brown, R. L. Graham, J. K. Perry, V. Du Pont, J. Pitts, B.
756   Ma, D. Babusis, E. Murakami, J. Y. Feng, J. P. Bilello, D. P. Porter, T. Cihlar, R. S.
757   Baric, M. R. Denison, T. P. Sheahan, Remdesivir Inhibits SARS-CoV-2 in Human Lung
758   Cells and Chimeric SARS-CoV Expressing the SARS-CoV-2 RNA Polymerase in Mice.
759   *Cell Rep* **32**, 107940 (2020).
760 27. R. L. Mackman, H. C. Hui, M. Perron, E. Murakami, C. Palmiotti, G. Lee, K. Stray, L.
761   Zhang, B. Goyal, K. Chun, D. Byun, D. Siegel, S. Simonovich, V. Du Pont, J. Pitts, D.
762   Babusis, A. Vijjapurapu, X. Lu, C. Kim, X. Zhao, J. Chan, B. Ma, D. Lye, A.
763   Vandersteen, S. Wortman, K. T. Barrett, M. Toteva, R. Jordan, R. Subramanian, J. P.
764   Bilello, T. Cihlar, Prodrugs of a 1'-CN-4-Aza-7,9-dideazaadenosine C-Nucleoside
765   Leading to the Discovery of Remdesivir (GS-5734) as a Potent Inhibitor of Respiratory
766   Syncytial Virus with Efficacy in the African Green Monkey Model of RSV. *J Med Chem*
767   **64**, 5001-5017 (2021).
768 28. Y. J. Hou, K. Okuda, C. E. Edwards, D. R. Martinez, T. Asakura, K. H. Dinnon, 3rd, T.
769   Kato, R. E. Lee, B. L. Yount, T. M. Mascenik, G. Chen, K. N. Olivier, A. Ghio, L. V.
770   Tse, S. R. Leist, L. E. Gralinski, A. Schafer, H. Dang, R. Gilmore, S. Nakano, L. Sun, M.
771   L. Fulcher, A. Livraghi-Butrico, N. I. Nicely, M. Cameron, C. Cameron, D. J. Kelvin, A.
772   de Silva, D. M. Margolis, A. Markmann, L. Bartelt, R. Zumwalt, F. J. Martinez, S. P.
773   Salvatore, A. Borczuk, P. R. Tata, V. Sontake, A. Kimple, I. Jaspers, W. K. O'Neal, S. H.
774   Randell, R. C. Boucher, R. S. Baric, SARS-CoV-2 Reverse Genetics Reveals a Variable
775   Infection Gradient in the Respiratory Tract. *Cell* **182**, 429-446 e414 (2020).
776 29. X. Xie, A. E. Muruato, X. Zhang, K. G. Lokugamage, C. R. Fontes-Garfias, J. Zou, J.
777   Liu, P. Ren, M. Balakrishnan, T. Cihlar, C. K. Tseng, S. Makino, V. D. Menachery, J. P.
778   Bilello, P. Y. Shi, A nanoluciferase SARS-CoV-2 for rapid neutralization testing and
779   screening of anti-infective drugs for COVID-19. *Nat Commun* **11**, 5214 (2020).
780 30. S. R. Leist, K. H. Dinnon, 3rd, A. Schäfer, L. V. Tse, K. Okuda, Y. J. Hou, A. West, C.
781   E. Edwards, W. Sanders, E. J. Fritch, K. L. Gully, T. Scobey, A. J. Brown, T. P. Sheahan,
782   N. J. Moorman, R. C. Boucher, L. E. Gralinski, S. A. Montgomery, R. S. Baric, A

Mouse-Adapted SARS-CoV-2 Induces Acute Lung Injury and Mortality in Standard
Laboratory Mice. *Cell* **183**, 1070-1085.e1012 (2020).
31.   R. L. Graham, E. F. Donaldson, R. S. Baric, A decade after SARS: strategies for
controlling emerging coronaviruses. *Nat Rev Microbiol* **11**, 836-848 (2013).
32.   L. Rosenbaum, Escaping Catch-22 - Overcoming Covid Vaccine Hesitancy. *N Engl J
Med* **384**, 1367-1371 (2021).
33.   B. Adhikari, P. Y. Cheah, Vaccine hesitancy in the COVID-19 era. *Lancet Infect Dis*,
(2021).
34.   Z. Wang, F. Schmidt, Y. Weisblum, F. Muecksch, C. O. Barnes, S. Finkin, D. Schaefer-
Babajew, M. Cipolla, C. Gaebler, J. A. Lieberman, T. Y. Oliveira, Z. Yang, M. E.
Abernathy, K. E. Huey-Tubman, A. Hurley, M. Turroja, K. A. West, K. Gordon, K. G.
Millard, V. Ramos, J. Da Silva, J. Xu, R. A. Colbert, R. Patel, J. Dizon, C. Unson-
O'Brien, I. Shimeliovich, A. Gazumyan, M. Caskey, P. J. Bjorkman, R. Casellas, T.
Hatziioannou, P. D. Bieniasz, M. C. Nussenzweig, mRNA vaccine-elicited antibodies to
SARS-CoV-2 and circulating variants. *Nature* **592**, 616-622 (2021).
35.   A. J. Brown, J. J. Won, R. L. Graham, K. H. Dinnon, 3rd, A. C. Sims, J. Y. Feng, T.
Cihlar, M. R. Denison, R. S. Baric, T. P. Sheahan, Broad spectrum antiviral remdesivir
inhibits human endemic and zoonotic deltacoronaviruses with a highly divergent RNA
dependent RNA polymerase. *Antiviral Res* **169**, 104541 (2019).
36.   W. T. Harvey, A. M. Carabelli, B. Jackson, R. K. Gupta, E. C. Thomson, E. M. Harrison,
C. Ludden, R. Reeve, A. Rambaut, C.-G. U. Consortium, S. J. Peacock, D. L. Robertson,
SARS-CoV-2 variants, spike mutations and immune escape. *Nat Rev Microbiol* **19**, 409-
424 (2021).
37.   D. C. Montefiori, P. Acharya, SnapShot: SARS-CoV-2 antibodies. *Cell Host Microbe* **29**,
1162-1162 e1161 (2021).
38.   R. Martin, J. Li, A. Parvangada, J. Perry, T. Cihlar, H. Mo, D. Porter, E. Svarovskaia,
Genetic conservation of SARS-CoV-2 RNA replication complex in globally circulating
isolates and recently emerged variants from humans and minks suggests minimal pre-
existing resistance to remdesivir. *Antiviral Res* **188**, 105033 (2021).
39.   Y. Liu, L. M. Yan, L. Wan, T. X. Xiang, A. Le, J. M. Liu, M. Peiris, L. L. M. Poon, W.
Zhang, Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* **20**, 656-
657 (2020).
40.   S. Zheng, J. Fan, F. Yu, B. Feng, B. Lou, Q. Zou, G. Xie, S. Lin, R. Wang, X. Yang, W.
Chen, Q. Wang, D. Zhang, Y. Liu, R. Gong, Z. Ma, S. Lu, Y. Xiao, Y. Gu, J. Zhang, H.
Yao, K. Xu, X. Lu, G. Wei, J. Zhou, Q. Fang, H. Cai, Y. Qiu, J. Sheng, Y. Chen, T.
Liang, Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in
Zhejiang province, China, January-March 2020: retrospective cohort study. *BMJ* **369**,
m1443 (2020).
41.   F. Zhou, T. Yu, R. Du, G. Fan, Y. Liu, Z. Liu, J. Xiang, Y. Wang, B. Song, X. Gu, L.
Guan, Y. Wei, H. Li, X. Wu, J. Xu, S. Tu, Y. Zhang, H. Chen, B. Cao, Clinical course
and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a
retrospective cohort study. *Lancet* **395**, 1054-1062 (2020).
42.   X. Lu, L. Wang, S. K. Sakthivel, B. Whitaker, J. Murray, S. Kamili, B. Lynch, L.
Malapati, S. A. Burke, J. Harcourt, A. Tamin, N. J. Thornburg, J. M. Villanueva, S.
Lindstrom, US CDC Real-Time Reverse Transcription PCR Panel for Detection of
Severe Acute Respiratory Syndrome Coronavirus 2. *Emerg Infect Dis* **26**, (2020).

Figure 1

A GS-441524

B RDV

C GS-621763

D A549-hACE2

E A549-hACE2

F Antiviral Assay in Primary HAE

G Normal Human Brochial Epithelial cells

H In vitro potency in human airway cells

Compound	Average EC$_{50}$ (nM)	
	A549-hACE2	NHBE
GS-621763	2801	125 ± 22
RDV	293.8	37.1 ± 0.4
GS-441524	3368	2454 ± 63

Figure 4

A. Weight Loss

B. Viral Lung Titer 4dpi

C. Nucleocapsid RNA Lung 4dpi

Group/dosage
- ▽ vehicle
- ● GS-621763; 30mg/kg BID, 12hr
- △ MPV; 30mg/kg; BID, 12hr
- ▼ MPV; 60mg/kg; BID, 12hr
- ■ GS-621763; 60mg/kg; BID, 24hr
- ▲ MPV; 60mg/kg; BID, 24hr

D. Pulmonary Function

E. ATS Acute Lung Injury 4dpi

F. Diffuse Alveolar Damage 4dpi

Figure 3

A. Weight Loss

B. Viral Lung Titer 4dpi

C. Nucleocapsid RNA Lung 4dpi

Group/dosage

D. Pulmonary Function

E. ATS Acute Lung Injury 4dpi

F. Diffuse Alveolar Damage 4dpi

Figure 2

A. Plasma pharmacokinetics

B. Weight loss

C. Viral Titer Lung 4dpi

D. Pulmonary Function

E. Congestion Score 4dpi

F. ATS Acute Lung Injury 4dpi

G. Diffuse Alveolar Damage 4dpi