Quantitative prediction of variant effects on alternative splicing using endogenous pre-messenger RNA structure probing

Authors: Jayashree Kumar[a,b], Lela Lackey[a,c], Justin M. Waldern[a], Abhishek Dey[a], David H. Mathews[d], Alain Laederach[a,b,1]

Affiliations: [a] Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC

[b] Curriculum in Bioinformatics and Computational Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC

[c] Department of Genetics and Biochemistry, Center for Human Genetics, Clemson University, Greenwood, SC

[d] Department of Biochemistry & Biophysics and Center for RNA Biology, 601 Elmwood Avenue, Box 712, School of Medicine and Dentistry, University of Rochester, Rochester, NY 14642

Abstract:

Splicing is a highly regulated process that depends on numerous factors. It is particularly challenging to quantitatively predict how a mutation will affect precursor messenger RNA (mRNA) structure and the subsequent functional consequences. Here we use a novel Mutational Profiling (-MaP) methodology to obtain highly reproducible endogenous precursor and mature mRNA structural probing data in vivo. We use these data to estimate Boltzmann suboptimal ensembles, and predict the structural consequences of mutations on precursor mRNA structure. Together with a structural analysis of recent cryo-EM spliceosome structures at different stages of the splicing cycle, we determined that the footprint of the B^{act} complex on precursor mRNA is best able to predict splicing outcomes for exon 10 inclusion of the alternatively spliced *MAPT* gene. However, structure alone only achieves 74% accuracy. We therefore developed a β-regression weighting framework that incorporates splice site strength, structure and

30   exonic/intronic splicing regulatory elements which together achieves 90% accuracy for

31   47 known and six newly discovered splice-altering variants. This combined

32   experimental/computational framework represents a path forward for accurate

33   prediction of splicing related disease-causing variants.

35   Introduction

Precursor messenger RNA (pre-mRNA) splicing is a highly regulated process in

eukaryotic cells (Z. Wang and Burge 2008). Numerous factors control splicing including

trans-acting RNA-binding proteins (RBPs), components of the spliceosome, and the

pre-mRNA itself. Pre-mRNA structure is a key attribute that directs splicing, particularly

alternative splicing, but we have only a poor understanding of pre-mRNA structure-

mediated splicing mechanisms (Taylor and Sobczak 2020). In addition, it is particularly

challenging to develop quantitative models capable of predicting splicing outcome,

specifically the Percent Spliced In (PSI) for alternatively spliced junctions. This difficulty

is especially true for predicting the effects of genetic variation at exon-intron junctions.

Indeed, mutations may affect not only the binding specificity of RBPs but also may alter

pre-mRNA structure (Tazi, Bakkour, and Stamm 2009).

Similar to the challenge of predicting PSI outcomes, the consequences of mutations on

pre-mRNA structure are difficult to predict. First and foremost, little is known about

native pre-mRNA structure because pre-mRNAs are relatively short-lived in cells

(Herzel et al. 2017). Only recently has pre-mRNA structure determination become

amenable to high resolution in vivo experimental characterization (Mustoe et al. 2018).

Second, it is not clear what structures of a pre-mRNA control spliceosome assembly

and activity. Finally, we lack quantitative measures for the relative weighting of RBPs'

affinity for specific motifs in pre-mRNA to the importance of pre-mRNA structure.

Several technical developments address these issues and enable us to propose an

integrated, RNA structure based-framework that accurately predicts the percent of

splicing. In this study, we used a combination of endogenous pre-mRNA chemical

structure probing (Homan et al. 2014), an RNA structure model that considers multiple

alternative structures in equilibrium (Dethoff et al. 2012; Lai et al. 2018), quantitative

analysis of exonic and intronic splicing enhancers/silencers (Fairbrother et al. 2002; Z.

Wang et al. 2004; Yang Wang, Ma, et al. 2012; Yang Wang, Xiao, et al. 2012), and a β-

regression weighting (Ferrari and Cribari-Neto 2004).

In this work we measure endogenous pre-mRNA structure in vivo by combining recent

developments in RNA structure Mutational Profiling (so-called -MaP approaches) with

targeted amplification of specific exon-intron junctions. This novel approach enables us

to obtain single-nucleotide RNA structure probing data for endogenous pre- and mature

mRNAs in the same cell. The high reproducibility of these data also makes it possible to

use Boltzmann suboptimal sampling guided by the data (Spasic et al. 2018) to predict

free energies of unfolding for an ensemble of structures. In addition, we can now

leverage recent high resolution cryo-Electron Microscopy (cryo-EM) structures of

various stages of the spliceosome during the splicing cycle to reveal the effective

spliceosomal footprint on pre-mRNA (L. Zhang et al. 2019).

As a model system to validate our framework, we study the effects of 47 experimentally

measured mutations at the Exon10-Intron10 junction of the human Microtubule-

Associated Protein Tau gene, *MAPT* (Park, Ahn, and Gallo 2016; Catarina Silva and

Haggarty 2020). Exon 10 is a cassette exon that is alternatively spliced resulting in a

Tau protein with either four microtubule binding repeats (4R) or three repeats (3R). The

ratio of 3R to 4R isoforms is approximately 1:1 (Hefti et al. 2018). This is highly unusual

for a splicing event as single-cell RNA-seq analysis demonstrates that this type of

event, where alternative isoforms are expressed equally, comprises less than 20% of all

splicing events (Song et al. 2017). The Exon10-Intron10 junction has 29 clinically

validated disease-causing mutations (Stenson et al. 2003) that impair the function of

Tau protein and are implicated in many neurodegenerative diseases (Spillantini et al.

1998; Hutton et al. 1998; Clark et al. 1998; Rizzu et al. 1999; Goedert et al. 1999).

Although some mutations alter the Tau protein sequence (Mirra et al. 1999; Iseki et al.

2001), 20 disease-associated mutations are known that deregulate *MAPT* pre-mRNA

splicing by altering the 1:1 ratio of 3R to 4R *MAPT* isoforms (Hutton et al. 1998;

D'Souza et al. 1999; Hasegawa et al. 1999; Jiang et al. 2000). An additional 27

mutations were previously experimentally tested to measure Exon 10 PSI with splicing

assays (D'Souza and Schellenberg 2000; Tan et al. 2019; Grover et al. 1999), making

this junction the most experimentally characterized junction of clinical importance in the

human genome and an excellent system for developing forward predictive models of

splicing. Our work thus provides a framework for integrating endogenous pre-mRNA

structure probing data with our current structural understanding of spliceosome

assembly and *trans*-acting RBPs to achieve unprecedented quantitative prediction

accuracy of the effect of mutations at structured exon-intron junctions.

Results

Median ratio of individual and tissue 3R to 4R MAPT mRNA isoforms is 1:1

Splicing of *MAPT* Exon 10 yields a 1:1 ratio of alternatively spliced isoforms (Goedert et

al. 1989; Andreadis 2005). To corroborate the 1:1 isoform ratio among tissues and

individuals, we analyzed RNA-sequencing data from the Genotype-Tissue Expression

(GTEx) database (Lonsdale et al. 2013). We selected tissue types with median *MAPT*

transcripts per million greater than 10 (Figure 1-figure supplement 1A) and calculated

the Percent Spliced In (PSI) value for Exon 10 for each sample (Figure 1A-source data

1; Materials and methods). We examined the distribution of PSIs for each tissue type

over 2,315 tissue samples in 375 individuals of median age 61 (Figure 1A; Figure 1-

figure supplement 1B). A PSI of 0 indicated that none of the *MAPT* transcripts in a

sample had Exon 10 spliced in (3R isoform), whereas a PSI of 1 corresponded to all

MAPT transcripts having Exon 10 spliced in (4R isoform). We found variation in Exon 10

PSI both within and between different tissue types; the pituitary gland had the largest

variation among brain tissues, and the cerebellum had the least variation but the

difference between the two standard deviations was 0.04. Also, while the pituitary gland

and caudate had the lowest and highest median Exon 10 PSI respectively among

individual samples, the distance between the two values was only 0.25. Interestingly,

although *MAPT*'s function in breast tissue is not understood compared with its function

in the brain, for breast tissue, individuals had greater variation in Exon 10 PSI and a

lower median PSI compared with the pituitary gland (Figure 1-figure supplement 1B).

We also discovered a large amount of variation within tissues of an individual (Figure 1-

figure supplement 1C), although there was significantly greater variation between

individuals than within a single individual (see Supplementary file 1 for ANOVA table).

Overall, 75% of samples were within a standard deviation of the median PSI of 0.54,

which confirmed that the 3R to 4R isoform ratio was approximately 1:1 among

individuals and within different tissue types. The consistency of this isoform ratio,

despite the likely presence of different levels of RBPs, suggest that inherent sequence

and structural features regulate splicing at this exon-intron junction. RNA structure

regulates alternative splicing around exon-intron junctions (Warf and Berglund 2010;

Buratti and Baralle 2004) and a hairpin structure at the exon 10-intron 10 junction is

implicated in establishing the 3R to 4R 1:1 isoform ratio (Hutton et al. 1998; Varani et al.

1999; Grover et al. 1999; Donahue et al. 2006). Hence, we next used high-throughput

chemical mapping techniques to interrogate the endogenous in vivo structure of the

MAPT junction.

Structure of 3R and 4R MAPT mature mRNA isoforms is open and accessibility of

exons is similar for the two isoforms

Although the structure of the *MAPT* pre-mRNA was previously studied computationally

and in vitro (Varani et al. 1999; Lisowiec et al. 2015; Tan et al. 2019; Chen et al. 2019)*,*

the structures of the mature 3R and 4R isoforms and *MAPT* pre-mRNA have not been

assessed in their endogenous in vivo context. We used dimethyl sulfate (DMS) to

chemically probe RNA structure in T47D and neuronal SH-SY5Y cells and primer-

amplified the Exon 9-Exon 11 and Exon 9-Exon 10-Exon 11 junctions during library

preparation for Mutational-Profiling (-MaP) (Figure 1B; Materials and methods). This

approach leverages the read-through aspect of MaP technology to probe the structure

of two alternatively spliced isoforms in the same cells. DMS reactivities for replicates,

and between T47D and SH-SY5Y MAPT mRNAs were highly correlated (Figure 1-figure

supplement 2A; Figure 1-figure supplement 2B; Figure 1-figure supplement 2D; Figure

1-figure supplement 2E).

We also collected in vivo DMS data for the small ribosomal RNA (SSU) whose

secondary structure is known from X-ray crystallography (Petrov et al. 2014) (Figure 1-

figure supplement 3A). The DMS reactivities for unpaired nucleotides in the SSU were

significantly higher than for paired nucleotides (Figure 1-figure supplement 3B),

confirming our probing strategy accurately recapitulates RNA secondary structure. We

used the SSU in vivo data to calibrate the estimation of equilibrium ensembles as

guided by MaP technology (Methods and materials), and we validated that structure

prediction guided by experimental DMS reactivities yielded more accurate estimation of

the SSU structure (Figure 1-figure supplement 3C). The median DMS reactivity of the

mature *MAPT* isoforms was 0.22, significantly greater than the median DMS reactivity of

the SSU, 0.0083 (Figure 1-figure supplement 3D); these results suggested that the

nucleotides of the mature *MAPT* isoforms were more accessible and unpaired

compared with the highly structured SSU, indicating that our endogenous in vivo

probing strategy reveals important differences in the structure of cellular RNAs.

Reactivities of Exon 9 and Exon 11 were highly correlated between the 3R and 4R

isoforms (Figure 1-figure supplement 2C). Additionally, computed base-pairing

probabilities guided by the experimental data for the two isoforms revealed that,

although there were some long-range interactions, 66% of base pairs spanned less than

50 nucleotides and were contained within the exon units (Figure 1B). This result

suggested that the mature exons function as their own structural unit. However, the

mature isoform structures did not suggest how they might regulate splicing of Exon 10.

Hence, we next chemically probed the MAPT pre-mRNA.

Figure 1

Figure 1: In vivo DMS-MaP structure probing data for 3R and 4R mature *MAPT* transcripts that are expressed in a 1:1 ratio.

A) Ratio of 3R and 4R MAPT transcripts is approximately 1:1 among brain tissues. There are 14 exons alternatively spliced in *MAPT*. Exons 4A, 6, and 8 are not found in brain mRNA. The four exons highlighted in color are repeat regions that form the microtubule binding domain in the Tau protein. Exon 10 is alternatively spliced to form the 3 repeat (3R) or 4 repeat (4R) isoform. The six canonical transcripts found in the central nervous system can be separated into 3R and 4R transcripts. Percent Spliced In (PSI) of Exon 10 was calculated from RNA-seq data for 2315 tissue samples representing 12 central nervous system tissue types and collected from 375 individuals in GTEx v8 database. The violin plot for each tissue type and the corresponding region on the brain diagram is colored by the median PSI for all samples of a given type. The patterned regions on the brain diagram indicate tissue types with no data. Tissue types Spinal cord and

Nucleus accumbens are not visualized on the brain diagram. The median PSI of

0.54 among all tissue samples is indicated by the red dotted line.

B) In vivo DMS-MaP structure probing data across exon9-exon11 junction of 3R

mature MAPT transcript. T47D cells were treated with DMS. Structure probing

data for junctions of interest were obtained using primers (Supplementary file 4)

following RT of extracted RNA. DMS reactivity is plotted for each nucleotide

across spliced junctions. Each value is shown with its standard error and colored

by reactivity based on color scale. High DMS reactivities correspond to

unstructured regions, whereas low DMS reactivities correspond to structured

regions. The base pairs of the predicted secondary structure guided by DMS

reactivities are shown in the arcs colored by pairing probabilities.

C) In vivo DMS-MaP structure probing data across exon9-exon10-exon11 junction

of 4R mature MAPT transcript

MAPT pre-mRNA Exon 10-Intron 10 junction is more structured compared with the mature isoforms

Existence of a hairpin at the *MAPT* Exon 10-Intron 10 junction, implicated in regulating Exon 10 splicing, was established by NMR and in vitro chemical probing (Varani et al. 1999; Lisowiec et al. 2015); however, the endogenous in vivo structure of this region has yet to be determined. While collecting data for mature *MAPT* isoform junctions, we simultaneously obtained data for the pre-mRNA Exon 10-Intron 10 junction (Figure 2A; Materials and methods). Replicates were highly correlated (Figure 2-figure supplement 1A). Surprisingly, although Exon 10 was still being spliced, the reactivities for Exon 10 in pre-mRNA and the mature 4R isoform were highly correlated (Figure 2-figure supplement 1B). Again, base pairing between nucleotides appeared to be contained within exons, independent of introns. The reactivities were highly correlated between data collected in SH-SY5Y and T47D cells (Figure 2-figure supplement 1C); thus, despite likely differences in RBP concentrations, the structure of the pre-spliced region is the same between cell lines. Additionally, we found lower DMS reactivities for the pre-mRNA Exon 10-Intron 10 junction compared with the mature isoform junctions (Figure 2-figure supplement 1D), which suggests that pre-mRNA is more structured than mature mRNA. We uncovered strong evidence for the previously in vitro identified hairpin structure in the DMS reactivity data; pairing probabilities were greater than 0.8 for the entire hairpin stem (Figure 2A).

Shifts in structural ensemble of MAPT Exon 10-Intron 10 junction associated with disease mutations correlate with changes in splicing level of Exon 10

Many RNAs inhabit multiple conformations in vivo to form a structural ensemble instead of a single rigid structure (Halvorsen et al. 2010; Adivarahan et al. 2018). We posit that a structural ensemble at the *MAPT* Exon 10-Intron 10 junction regulates Exon 10 splicing and disease mutations alter the composition of the structural ensemble to disrupt splicing.

We used Boltzmann sampling of RNA structures guided by DMS reactivity data (Spasic

et al. 2018) (Materials and methods) to sample 1000 structures for the wild type and two

mutant intronic sequences, +15A>C and +19C>G. The two mutations alter, in opposite

directions, the isoform ratio at this junction (Tan et al. 2019). We visualized the

structural ensemble for the 3000 structures using t-Distributed stochastic neighbor

embedding (t-SNE) (Van Der Maaten and Hinton 2008) (Figure 2B; Materials and

methods). Each structure is a dot and is colored by the $\Delta G^{\ddagger}$ of unfolding of the 5' splice

site defined as the last three nucleotides of Exon 10 and the first six nucleotides of

Intron 10 (Yeo and Burge 2004). The lower the unfolding free energy, the easier to

unfold the structure. Overall, although there was a range of unfolding free energies for

the three ensembles, there were three distinct populations of free energies for the three

sequences (Figure 2-figure supplement 2A). We used k-means clustering to identify

representative structures for each cluster (Figure 2B; Figure 2-figure supplement 2B;

Materials and methods). We quantified and visualized the density of the clusters (Figure

2C; Materials and methods) and revealed distinct regions in the structure space

occupied by each sequence. More than 55% of structures in the ensemble of the

+19C>G mutation, which shifts the isoform balance entirely 3R (3R mutation) (Figure

2C inset), clustered in the lower left quadrant with larger unfolding free energies for the

splice site. This result was evidenced by the highly base-paired exon-intron junction in

the representative structure for the cluster. Hence, in the presence of the 3R mutation,

the structural ensemble of the junction shifted towards more closed structures.

Conversely, greater than 50% of structures in the ensemble of the +15 A>C mutation,

which shifts the isoform balance entirely 4R (4R mutation) (Figure 2C inset), were

clustered in the upper left quadrant with lower unfolding free energies for the splice site.

The representative structure for this region was more open and accessible around the

exon-intron junction. Correspondingly, the wild-type sequence had structures distributed

across the entire space consistent with an ensemble of structures. The exon-intron

junction of the representative structure for this region was not as accessible with the 4R

mutation, but it had fewer base-pairs than with the 3R mutation, a result recapitulated by

the two other representative structures in the right quadrants (Figure 2-figure

supplement 2B).

Figure 2

A

B

C

Figure 2: The 4R and 3R mutations shift DMS reactivity-guided structural ensemble of
Exon 10-Intron 10 junction to more open and closed structures, respectively.

A) In vivo DMS-MaP structure probing data across Exon 10-Intron 10 junction of
precursor MAPT transcript. T47D cells were treated with DMS. Structure probing
data for junctions of interest were obtained using primers (Supplementary file 4)
following RT of extracted RNA. DMS reactivity is plotted for each nucleotide.
Each value is shown with its standard error and colored by reactivity based on
the color scale. High DMS reactivities correspond to unstructured regions,
whereas low DMS reactivities correspond to structured regions. Base pairs of
predicted secondary structure guided by DMS reactivities are shown by arcs
colored by pairing probabilities. Strongly predicted hairpin structure near exon-
intron junction is highlighted by dotted box.

B) t-SNE Visualization of structural ensemble of wildtype (WT) and, +19C>G (3R)
and +15A>C (4R) mutations. Structures were predicted using Boltzmann
suboptimal sampling and guided by DMS reactivity data generated in A. Data
were visualized using t-Distributed Stochastic Neighbor Embedding (t-SNE).
Shown are 3000 structures corresponding to 1000 structures per sequence.
Each dot represents a single structure and is colored by calculated unfolding free
energy of splice site at exon-intron junction (3 exonic, 6 intronic bases). Data
were clustered by k-means clustering and representative structures for three of
the clusters are shown. The exon-intron junction is marked by EIJ on each
structure. Positions of 3R and 4R mutations are marked by a red asterisk on their
respective representative structures.

C) Density contour plots of structural ensemble of WT and, 3R and 4R mutations.
Contour plots were derived from the distribution of points on the t-SNE plot in B.
The darker the blue, the higher the density of structures. Contour lines
additionally represent density of points. Color scales for the three plots are
identical. Gel insets of RT-PCR products from splicing assays in HEK293 cells
for 3R and 4R mutation are in their respective density plots.

Unfolding mRNA within the spliceosome B^(act) complex yields best prediction of Exon 10

splicing level

RNA structure controls alternative splicing by hindering or aiding accessibility of key

regulatory regions to spliceosome components (McManus and Graveley 2011; Warf and

Berglund 2010). The 5' splice site, defined as the last 3 nucleotides of the exon and first

6 nucleotides of the intron, is the minimum region of RNA that must be accessible for

base pairing with the U1snRNA (Blanchette and Chabot 1997; Singh, Singh, and

Androphy 2007). However, the splicing cycle, orchestrated by the spliceosome,

traverses multiple stages to prepare the pre-mRNA and catalyze the two-step splicing

reaction (Matera and Wang 2014) (Figure 3A). The RNA itself adopts many

conformations as different components of the spliceosome bind to it (L. Zhang et al.

2019). In addition to the 5' splice site, a larger segment of RNA likely needs to unpair to

accommodate the changing conformations induced by the spliceosome. We analyzed

high resolution Cryo-EM structures of the human spliceosome Pre-B, B, Pre-B^(act), and

B^(act) complexes (Charenton, Wilkinson, and Nagai 2019; Bertram et al. 2017; Townsend

et al. 2020; X. Zhang et al. 2018) to quantify the number of nucleotides around the 5'

splice site for which sufficient density was observed in the cryo-EM structure and which

were unpaired (Materials and methods). As can be seen in Figure 3A, the number of

unpaired pre-mRNA nucleotides observed in each structure increased through the

splicing cycle. Thus, it is likely that RNA structures outside the U1snRNA binding site

have to be unfolded to accommodate splicing.

To evaluate the footprint of the spliceosome that best predicts splicing outcome, we

initially focus on predicting 20 synonymous and intronic mutations as a training set

(Figure 3-figure supplement 1A). These mutations are most likely to have a structural

component to their function (Sharma et al. 2019; Lin, Taggart, and Fairbrother 2016).

The distribution of $\Delta G^{\ddagger}$ of unfolding of the splice sites in the presence of these mutations

was correlated with Exon 10 PSI (Figure 3-figure supplement 1B). We calculated the

$\Delta G^{\ddagger}$ of unfolding of the RNA near the 5' splice site in the four splicing stages' footprints.

Features of the unfolding free energy distribution including mean and standard deviation

were then used in a beta regression to predict Exon 10 PSI (Materials and methods; Eq.

1). Unfolding larger regions of the exon-intron mRNA junction improved the predictive

power of the model, and the B^{act} complex footprint yielded the best prediction accuracy

(R^2 = 89%; Figure 3B). Crucially, we found that using features of the distribution of

unfolding free energies in the structural ensemble produced greater predictive power

than simply using the unfolding free energy of a single minimum free energy (MFE)

structure (Figure 3-figure supplement 1C). We performed bootstrapping cross-validation

and confirmed that unfolding the RNA within the B^{act} spliceosome complex yielded the

best prediction (Figure 3C). We tested the structural ensemble-based model on 24 non-

synonymous and compensatory mutations. Although the model performed well for

compensatory mutations (median bootstrapped R^2=0.76), it yielded significantly less

accurate predictions for non-synonymous mutations (median bootstrapped R^2=-0.21)

(Figure 3-figure supplement 1D). One possible reason this structure-only model has

limited performance is that it does not account for the effects of mutations on potential

splicing regulatory elements (SREs) in the sequence.

Figure 3

A

B

C

Figure 3: The best predictor of Exon 10 PSI for intronic and synonymous mutations was

the unfolding free energy of pre-mRNA during the B^{act} stage of splicing

A) Spliceosome footprint on pre-mRNA during splicing cycle. Structure in the center

of the cycle is the WT representative structure from Fig 2B. The dotted box

indicates the zoomed-in region at each stage of interest. Cryo-EM structures of

the human spliceosome complex at stages Pre-B (PDB ID: 6QX9), B (PDB ID:

5O9Z), Pre-B^{act} (PDB ID: 7ABF) and B^{act} (PDB ID: 5Z56) are available in the

Protein Data Bank. The region around the 5' splice site of pre-mRNA within the

spliceosome at each stage is highlighted in blue on the zoomed-in representative

structure. The number of nucleotides for each stage is as follows: Pre-B (2

exonic, 8 intronic); B (10 exonic, 17 intronic); Pre-B^{act} (9 exonic, 20 intronic); B^{act}

(12 exonic, 31 intronic). These values represent the minimum number of

nucleotides required to be unfolded to be accessible to the spliceosome. The

mean free energy and standard error to unfold RNA within the spliceosome at

each stage is calculated for the WT structural ensemble and indicated under the

zoomed-in structure.

B) Exon 10 PSIs of synonymous and intronic mutations predicted with the unfolding

free energy of pre-mRNA within the spliceosome in B, Pre-B, Pre-B^{act}, B^{act} stages

versus corresponding experimental PSIs measured in splicing assays. Exon 10

PSIs were predicted using Eq. 1. Grey line represents the best fit with dotted

lines indicating the 95% confidence interval. Pearson correlation coefficients (R^2)

of experimental to predicted PSIs were calculated for each stage. Violin plots

(inset) show R^2s calculated for each mutation category by training and testing on

subsets of all mutations by non-parametric bootstrapping; Synonymous (n=6),

Intronic (n=14), Wildtype (n=1).

C) Overall Pearson correlation coefficients (R^2) calculated for experimental versus

predicted Exon 10 PSIs by nonparametric bootstrapping of mutations. Subsets of

mutations were randomly sampled 10 times, trained and tested using unfolding

free energy of the exon-intron junction region of pre-mRNA within the

spliceosome for the respective splicing stage. Pearson's R^2 was calculated by

comparing predicted PSIs to experimental PSIs. A two-tailed Wilcoxon Rank

Sum test was used to determine significance between B^{act} complex and the other

three complexes. Level of significance: ***p-value $< 10^{-6}$, **p-value < 0.001, * p-

value < 0.01

Effect of exonic non-synonymous mutations was best predicted by motif strength

changes of splicing regulatory elements

Exon 10 splicing is highly regulated by differential binding of RBPs to *cis*-SREs within

exon 10 and intron 10 (Qian and Liu 2014). While our structure-only model performs

moderately well for 47 mutations (R^2=0.74) (see Supplementary file 2 for further details

about mutations), *MAPT* Exon 10 PSIs of non-synonymous mutations were poorly

predicted (median bootstrapped R^2 = -0.21, Figure 4-figure supplement 1B). Hence, we

investigated whether these non-synonymous mutations are predicted better by

incorporating changes to the strength of adjacent SREs. We identified SREs by

similarity to reported general enhancer and silencer hexamer motifs (Fairbrother et al.

2002; Z. Wang et al. 2004; Yang Wang, Ma, et al. 2012; Yang Wang, Xiao, et al. 2012)

(Materials and methods). We calculated the changes to splice site, enhancer, and

silencer motif strengths in the presence of a mutation (Materials and methods) and

visualized the motif strength changes in a heatmap (Figure 4A). We found that using

splice site strength as the sole predictor yielded poor prediction of Exon 10 PSI in all

mutation categories (Figure 4B; Eq. 3) because most mutations were outside the splice

site. We quantified a weak positive correlation between PSI and enhancer strength and

a significant negative correlation between PSI and silencer strength (Figure 4A; Figure

4-figure supplement 1C). We modeled Exon 10 PSI with the changes to the motif

strength of all splicing regulatory elements (Eq. 4) and found an increase in prediction

accuracy (R^2=0.51; Figure 4C) compared with using only splice site strength (R^2=0.29).

Non-synonymous mutations were predicted more accurately using SRE strength with a

median bootstrapped R^2 of 0.75.

Many RBPs have been identified that regulate *MAPT* Exon 10 splicing (Qian et al. 2011;

Ian D'Souza and Schellenberg 2006; Kondo et al. 2004; J. Wang et al. 2004; L. Gao et

al. 2007; S. Ding et al. 2012; Broderick, Wang, and Andreadis 2004; Yan Wang et al.

2010; Kar et al. 2006, 2011; P. Ray et al. 2011). To determine whether these proteins

specific to Exon 10 splicing would improve the model's accuracy, we calculated

changes to the strength of their RBP motifs obtained from high throughput sequencing

of bound RNAs (Dominguez et al. 2018; D. Ray et al. 2013) (Materials and methods).

Unlike SRE motifs, there was no clear pattern or correlation between motif strength

change and PSI (Figure 4-figure supplement 2A, B). Subsequently, the model's

prediction accuracy was lower (R^2=0.08, Figure 4-figure supplement 2C), and changes

to the strength of general SRE motifs were better predictors of Exon 10 PSI.

Figure 4

A

Figure 4: Combining the strength of all splicing regulatory elements improves prediction of Exon 10 PSI by 75% compared with using only splice site strength

A) Heatmap of splicing regulatory element (SRE) relative strength for 47 mutations compared with wildtype (WT). A value of 0 indicates mutation does not change WT SRE strength, positive values indicate SRE strength is greater than WT, and negative values indicate SRE strength is weaker than WT. Splice site strengths were calculated using MaxEntScan; a splice site was defined as the last 3 nucleotides of the exon and first 6 nucleotides of the intron. Enhancer and silencer strengths were calculated from position weight matrices of known motifs derived from cell-based screens (Materials and methods).

B) Exon 10 PSIs of 47 mutations predicted from change in splice site strength and plotted against experimental PSIs measured in splicing assays. Exon 10 PSIs predicted using Eq. 3. Each point on the scatterplot represents a mutation and is colored by mutation category. Grey line represents the best fit with dotted lines indicating the 95% confidence interval. Pearson correlation coefficient (R^2) calculated of experimental to predicted PSIs. Violin plot shows R^2s calculated for each category by training and testing on subsets of all mutations by non-

502   parametric bootstrapping; Exonic non-synonymous (n=11), Exonic synonymous

503   (n=7), Intronic (n=15), Compensatory (n=14), Wildtype (n=1).

504  C) Exon 10 PSIs of 47 mutations predicted by combining change in splice site,

505   enhancer, and silencer strength and plotted against experimental PSIs measured

506   in splicing assays. Exon 10 PSIs predicted using Eq. 4.

Model with both structural and SRE motif changes yields best prediction of Exon 10 PSI

Our quantitative models showed that, although SRE motif changes accurately predicted

the effects of non-synonymous mutations, structural changes were a better predictor of

splicing outcomes of intronic and synonymous mutations. Combining all features (Eq. 6)

yielded the highest prediction accuracy (R^2 = 0.89) (Figure 5A). This combined

interactive model consistently produced more accurate predictions of Exon 10 PSI

compared with a structure-only model and an SRE-only model for all mutation

categories (Figure 5B). An additive model (Eq. 7) had lower prediction accuracy (R^2=

0.80) (Figure 5-figure supplement 1A), and this lower accuracy resulted primarily from

less accurate PSI predictions of non-synonymous mutation effects (Figure 5-figure

supplement 1B).

To determine whether structure or SRE changes were responsible for the splicing

changes from each mutation, we hierarchically clustered the four primary features for

the 47 experimentally validated mutations (Materials and methods). Six categories

emerged from the clustering of features (Figure 5C) where approximately 80% of

mutations modified both structure and silencer strength (Figure 5-figure supplement

1C). Further, we found that for more than 50% of mutations both structure and SRE

motif strength were altered in the same direction and accordingly promoted Exon 10

splicing in that direction (Figure 5D). For the remaining mutations in which structure and

SRE strength changed in opposite directions, structure dominated the direction of

splicing for 18% of mutations, and SRE strength was dominant for 20% (Figure 5D).

Overall, these results supported our conclusion that both structure and SREs have

equally important effects in regulating splicing at this exon-intron junction.

Figure 5

Figure 5: Combining structure and SRE strength into a unified model is the best

predictor of Exon 10 PSI

A) Exon 10 PSIs of 47 mutations predicted from combined model using structure

and SRE strength and fit to experimental PSIs measured in splicing assays.

Exon 10 PSIs predicted using Eq. 6. Each point on scatterplot represents a

mutation and is colored by mutation category. Grey line represents the best fit

with dotted lines indicating the 95% confidence interval. Pearson correlation

coefficient (R^2) calculated of experimental to predicted PSIs.

B) Violin plots of correlation coefficients for each mutation category for structure

model, SRE model, and combined model. R^2s calculated for each mutation

category by training and testing on subsets of all mutations by non-parametric

bootstrapping 10 times. Structure model uses unfolding free energy of pre-mRNA

within spliceosome at B^{act} stage as predictor. SRE strength model uses relative

change in SRE strength as predictor. Combined model using both structure and

SRE strength and weighs the features based on if mutation is

intronic/synonymous or non-synonymous (Materials and methods).

C) Heatmap of the normalized changes in structure and SRE strength for each

mutation clustered by affected features. Features were normalized such that a

value of 1 implied that change in the feature should result in Exon 10 being

spliced in (4R isoform, blue), whereas a value of 0 implies Exon 10 should be

spliced out (3R isoform, red). Mutations were clustered using hierarchal

clustering on normalized features (Materials and methods). Experimental PSIs

are plotted for each mutation with a PSI of 1 colored as blue, PSI of 0.5 colored

as white and PSI of 0 colored as red.

D) Pie chart showing the features that regulate Exon 10 splicing for the 47

experimentally validated mutations. The pie chart was generated based on the

heatmap in C. Exon 10 splicing for 51.1% of mutations is supported by changes

in both structure and SRE, which implies that structure, at least one SRE, and

PSI are either all blue or all red. Exon 10 splicing for 23.4% of mutations is

supported by changes in SRE wherein one of the SREs is the same color as PSI.

For 17.0% of mutations, structural changes support Exon 10 splicing wherein

structure and PSI are the same color. For 4 mutations (8.5%), the colors of none

of the features match the color of PSI.

Mutations around the MAPT Exon 10-Intron 10 junction skew to Exon 10 inclusion

Having established that our quantitative models accurately predicted Exon 10 PSIs for

experimentally validated mutations, we interrogated the model by performing a

systematic mutagenic analysis spanning a 100-nucleotide window of the exon-intron

junction (Figure 6A). Our model predicts that more mutations result in the inclusion of

Exon 10 (4R isoform). This is consistent with the observation that a majority (75%) of

known disease associated mutations (Figure 6B) are also 4R; this result is also

consistent when categorized by all substitution types (Figure 6-figure supplement 1A).

We found that a significantly greater proportion of disease mutations (76.4%) resulted in

changes to both structure and SRE compared with non-disease mutations (36.0%)

(Figure 6C) suggesting that mutations which affect both structure and SREs have a

greater likelihood of causing disease compared with mutations that alter only one of the

two factors. Intriguingly, mutations overall caused a slight skew towards a structured

exon-intron junction, which would result in decreased inclusion of Exon 10 (Figure 6A,

Figure 6-figure supplement 1B). However, changes to SRE strength skewed towards

increased inclusion of Exon 10 (Figure 6-figure supplement 1C), which suggested that

SREs were acting to counter the effect of structural changes. Our model reveals how a

complex balance of structure and SRE RBP binding sites effectively results in the

observed 50:50 ratio of the 3R and 4R isoforms.

To assess the general applicability of our model beyond our mutation training set, we

predicted Exon 10 PSIs for 55 variants of unknown significance (VUSs) found in dbSNP

(see Supplementary file 3 for further details of VUSs). These are mutations observed in

the human population but are not currently associated with disease. The mean Exon 10

PSI for VUSs was 0.66, and 70% were within a standard deviation of the mean (Figure

6D). We observed that only a few mutations were predicted to have a PSI of zero (3R)

(Figure 6D red bar). We therefore experimentally verified with splicing assays (Materials

and methods) 6 VUSs: 3 VUSs predicted to be 3R, 1 VUS predicted to be 4R and 2

VUSs predicted to maintain the WT splicing ratio (Figure 6D). We found these 6

predictions were correct (Figure 6E). The three 3R VUSs made the region around the

exon-intron junction more structured while the 4R VUS made the region less structured

compared to WT (Figure 6-figure supplement 1D) matching the direction of Exon 10

splicing change. Though we see changes to SRE strength match up to Exon 10 splicing

direction for +30U>C and -6G>A, this was not the case for +25C>G and +23U>C

(Figure 6-figure supplement 1E). For +23U>C and +26G>A, we observed changes in

structured-ness around the exon-intron junction and silencer strengths in diverging

directions (Figure 6-figure supplement 1D, E) suggesting that these opposing changes

would preserve the WT 3R/4R ratio.

Figure 6

Figure 6: Mutations around Exon 10-Intron 10 junction skew towards inclusion of Exon 10

A) Heatmap of predicted Exon 10 PSIs for every possible mutation around 100 nucleotide window of Exon 10-Intron 10 junction. Combined model was trained using 47 mutations with experimental PSIs measured from splicing assays and used to predict PSIs for all mutation combinations for 100 nucleotides around the junction. Tiles with sequence indicate the wild type nucleotide at the position. Heatmap of mean PSI per position and mean relative change in unfolding free energy of pre-mRNA within spliceosome at B[act] stage compared with wild type is shown below the gene diagram.

B) Violin plot of predicted PSIs for all possible mutations around Exon 10-Intron 10 junction and only disease mutations. All possible mutations (n=300), disease

mutations (n=17). A two-tailed Wilcoxon Rank Sum test was used to determine

significance between the two categories. Level of significance: ***p-value $< 10^{-6}$,

**p-value < 0.001, * p-value < 0.01

C) Pie chart showing features that drive Exon 10 splicing for disease and non-

disease mutations. The pie chart was generated by quantifying the number of

mutations for which the direction of predicted Exon 10 PSI matched the direction

of structure or SRE change. Exon 10 splicing for 76.4% of disease mutations is

supported by changes to both structure and SRE compared with only 36.0% of

non-disease mutations. The difference in proportions was tested with a one-tailed

Fisher's exact test.

D) Histogram displaying the distribution of predicted PSIs using the combined model

for 55 variants of unknown significance (VUSs) found in dbSNP. Density curve

was overlaid on top of histogram showing that predicted PSIs skew away from

3R. Dotted line shows mean predicted PSI of 0.66. VUSs tested in splicing

assays are indicated by their dbSNP RS IDs.

E) Representative gel of RT-PCR data for splicing assay in the presence of VUSs.

Splicing reporter was transfected into HEK293 cells. The mean Exon 10 PSI

displayed for each variant was calculated from three replicates and standard

error is shown in brackets below. Structure diagram on left displays the location

of the VUSs tested.

Discussion

In vivo DMS chemical probing of endogenous MAPT Exon 10-Intron 10 junction

Splicing specificity is complex (Baralle and Giudice 2017). The spliceosome does not rely on sequence alone to correctly identify 5' and 3' splice sites; other cues ensure correct binding to appropriate locations. In addition, the 5' splice site must be accessible to permit base pairing with the U1snRNA to initiate splicing (Roca et al. 2012). The *MAPT* Exon 10-Intron 10 junction is a well-studied example of the effect of 5' splice site secondary structure in splicing regulation. A hairpin was hypothesized initially because disease mutations close to the exon-intron junction (Hutton et al. 1998; Grover et al. 1999) shifted the isoform balance to either completely exclude or include Exon 10. Although NMR, in vitro chemical probing, and computation confirmed the presence of the hairpin (Varani et al. 1999; Chen et al. 2019; Lisowiec et al. 2015), recent studies showed that most RNAs were less structured in vivo and in the nucleus compared with in vitro conditions (Sun et al. 2019; Rouskin et al. 2014). However, our results revealed that this is not the case for the Exon10-Intron10 junction: in vivo chemical probing of the endogenous junction showcased strong evidence of structure.

In this study we observed that, in vivo, endogenous exons are less structured than introns, as found by Sun et al (Sun et al. 2019). Mature *MAPT* 3R and 4R exon-exon junctions are less structured compared with the pre-mRNA Exon 10-Intron 10 junction. The high correlation of structure we observed between the same exons found in different *MAPT* isoforms corroborates results observed with yeast ribosomal protein genes (Zubradt et al. 2016), which suggests that RNA folding in both pre- and post-spliced human exons is highly local and modular in exons.

Changes to structural ensemble around the 5' splice site are strong predictors of Exon 10 splicing

We showed that structural ensembles have an important function at the Exon 10-Intron 10 junction. If the 5' SS was always paired, only one isoform lacking Exon 10 would

result. However, the simultaneous presence of 3R and 4R isoforms implies that the

junction is accessible in a subset of the structures. Unlike transfer RNAs and ribosomal

RNAs that have single structures (Petrov et al. 2014), most RNAs are dynamic,

unfolding and refolding within a landscape (Cruz and Westhof 2009; Giegé et al. 2012).

We found disease mutations produced distinct shifts in the ensemble of the *MAPT* Exon

10-Intron 10 junction; the shifts corresponded to changes in the 3R:4R isoform ratio and

confirmed that ensembles are essential at this junction. The activity of ensembles was

corroborated by our quantitative model; including free energy features of the structural

ensemble produced 1.5 times more accurate prediction of Exon 10 PSI compared with

the unfolding free energy of the minimum free energy (MFE) structure.

Considering a larger spliceosome footprint on pre-mRNA produced more accurate

prediction of Exon 10 PSI

The U1snRNA base pairs with the nine nucleotide sequence around the exon-intron

junction (Roca et al. 2012). However, our analysis of the Cryo-EM structures of the

human spliceosomal assembly cycle revealed that a larger region of the pre-mRNA

interacts with the spliceosome during the splicing cycle and is therefore unfolded. Like

the other main cellular ribonucleoprotein complex, the ribosome (Ingolia 2016), there is

likely a spliceosomal footprint on the pre-mRNA and a minimum span around the

splicing signals (5' splice site, 3' splice site and branch point) must be single-stranded

for splicing to occur. Accordingly, our structural model performed most accurately when

we used the unfolding free energy of 43 nucleotides around the 5' exon-intron junction

that exists within the spliceosome B[act] complex. This suggests that structures distal to

the exon-intron junction regulate Exon 10 splicing, a finding that corroborates evidence

that RNA structure near this exon-intron junction is more extended than previously

determined (Tan et al. 2019). This result, combined with our use of Boltzmann

suboptimal sampling demonstrates the key role of pre-cursor mRNA structure in splicing

outcome.

RNA structure and SREs have complementary functions in MAPT Exon 10 regulation

Considerable evidence supports a function for either splicing regulatory elements and

their corresponding RBPs or RNA structure in alternative splicing of *MAPT* Exon 10 at

the 5' splice site (Andreadis 2012). However, there was no consensus as to which of the

two factors is dominant. The regression model we developed established the relative

importance of RNA structure vs. SREs at the exon-intron junction. We discovered a

cooperative relationship between SREs and RNA structure whereby exonic non-

synonymous mutations promoted splicing changes primarily by SRE motifs and exonic

synonymous and intronic mutations by RNA structure around the exon-intron junction. A

combined model that accounted for both structure and SREs was the most accurate

predictor of Exon 10 PSI, and most experimentally validated mutations altered RNA

structure and SRE motif strength around the Exon 10-Intron 10 junction in the same

direction (Figure 5D). The model further suggested that the overall region favored

increased Exon 10 inclusion (Figure 6 A,B), which confirmed previous experimental

findings that inclusion is Exon 10's typical splicing mode (Q. S. Gao et al. 2000). This

preference was proposed to be due to a weak 5' splice site (Ian D'Souza and

Schellenberg 2005), and, indeed, we found that almost all experimentally validated

mutations strengthened the splice site to increase inclusion of Exon 10 (Figure 4A).

However, interestingly, our model revealed that structural changes caused by the

mutations resulted in a more structured exon-intron junction, which would imply

decreased Exon 10 inclusion. However, SRE strength alterations overall skewed more

towards increased Exon 10 inclusion, which suggest that SREs and the RBPs that bind

them buffer the effects of RNA structure to maintain the 1:1 isoform ratio at this junction.

Our work revealed that structure and splicing regulatory elements most often have

opposite effects on splicing outcomes. However, disease variants were the exception to

this rule and resulted in a synergistic effect on splicing outcome (Fig. 6E), leading to a

greater disruption of splicing, and therefore increased pathogenicity. The combined

model was finally validated by accurate prediction of the effects of six previously

untested VUSs on Exon 10 splicing (Figure 6E). As was the case with the complete

mutagenesis, there were few VUSs predicted to completely alter the ratio of isoforms to

entirely 3R: only 5 VUSs had PSIs less than 0.25. However, our model accurately

predicted the effect of the three 3R VUSs tested. Interestingly, the systematic

computational mutagenesis revealed a hotspot of 3R mutations around 25-30

nucleotides downstream of the exon-intron junction (Figure 6A) and indeed the 3R

VUSs experimentally validated were located in this region.

Quantitative modeling of splicing regulation at exon-intron junctions

Predictive models can measure the contribution of individual factors to an outcome.

Structure around the 5' splice site and SRE motifs were excellent predictors of Exon 10

splicing in cells. The use of general SRE motifs enables this splicing framework to

extend to other exon-intron junctions. By using a common dependent variable of Exon

10 PSI, we could use experimentally validated mutation data from disparate sources.

Although our model provided an exact PSI prediction for each mutation, its principal

utility was in predicting the direction in which the 3R:4R isoform ratio shifted from the

wild type balance. On the basis of RNA-sequencing of brain tissue from healthy

individuals, we find a range of Exon 10 PSIs between individuals and between tissues

within an individual (Figure 1A). Even in individuals with progressive supranuclear palsy,

a tauopathy in which *MAPT* variants are implicated, there is variability in Exon 10 PSIs

between different brain tissues (Majounie et al. 2013). Ultimately, although it is likely

that what is considered the correct ratio for normal brain function varies between

tissues, our model provides a means to determine the baseline change of Exon 10

splicing simply based on sequence features. Many neurodegenerative diseases are

caused by mutations around the *MAPT* Exon 10-Intron 10 junction, and there are no

approved therapeutics that target this junction. Our work suggests that it is crucial to

consider the larger structural context of the Exon 10-Intron 10 junction and the interplay

between structure and SREs when considering the consequences of mutations on

splicing regulation and the design of potential therapeutics to alter this ratio.

Materials and methods

809

810 *MAPT* **Exon 10 PSIs for GTEx tissue types**

811 Aligned BAM files of individual samples from the Genotype-Tissue Expression (GTEx)

812 v8 project, for tissue types with *MAPT* TPM greater than 10, were accessed in the

813 AnVIL/Terra environment (Kumar 2020a). Reads aligning to *MAPT* were extracted in

814 Terra (Kumar 2020b) and downloaded. Exon 10 PSIs were quantified per BAM file with

815 Outrigger (Song et al. 2017) using *MAPT* transcriptome reference from Ensembl

816 GRCh38. Only samples with at least 10 reads mapping across the Exon 10-Intron 10

817 junction were considered. Median values for each tissue type were calculated and then

818 visualized on the brain diagram with R package, CerebroViz (Bahl, Koomar, and

819 Michaelson 2017). Source file for Figure 1 provides Exon 10 PSI values for the 2,962

820 samples. An ANOVA test was run in R to test significance in variation between

821 individuals versus within an individual (for individuals with MAPT expression in more

822 than 7 tissues) (Supplementary file 1).

823

824 **Culture of T47D and SH-SY5Y cells**

825 Mammary gland carcinoma cells (T47D) were cultured in RPMI 1640 medium,

826 supplemented with 10% Fetal Bovine Serum (FBS) and 0.2 units/mL of human insulin at

827 37°C and 5% CO_2. Bone marrow neuroblastoma SH-SY5Y cells were cultured in 1:1

828 mixture of 1X Minimum Essential Medium (MEM) and 1X F12 medium, supplemented

829 with 10% FBS at 37 °C and 5% CO_2.

830

831 **In vivo DMS-MaP probing for** *MAPT* **RNA**

832 Approximately 20 million T47D cells and 30 million SHSY-5Y cells were harvested by

833 centrifugation and resuspended in bicine buffered medium (300 mM Bicine pH 8.3, 150

834 mM NaCl, 5 mM $MgCl_2$) followed by treatment with DMS (1:10 ethanol diluted) for 5 min

835 at 37°C. For the negative control (unmodified RNA), instead of DMS, an equivalent

836 amount of ethanol was added to the same number of T47D and SH-SY5Y cells. After

837 incubation, the reactions were neutralized by addition of 200 μl of 20% by volume β-

mercaptoethanol. Total RNA was extracted by Trizol (ThermoFisher Scientific), treated

with TURBODNase (ThermoFisher Scientific), purified using Purelink RNA mini kit

(ThermoFisher Scientific) and quantified with NanoDrop™ spectrophotometer.

DMS-MaP cDNA synthesis, library construction and sequencing for *MAPT* RNA

Purified RNA (9 µg) was reverse transcribed using Random Primer 9 (NEB) and

SuperScript II reverse transcriptase under error prone conditions as described in Smola

et al., 2015. The resultant cDNA was purified using G50 column (GE healthcare) and

subjected to second strand synthesis (NEBNext Second Strand Synthesis Module).

Supplementary file 4 lists PCR primers used for library generation. The cDNA was

amplified by the NEB Q5 HotStart polymerase (NEB). Secondary PCR was performed

to introduce TrueSeq barcodes (Smola et al. 2015). All samples were purified using the

Ampure XP (Beckman Coulter) beads and quantification of the libraries was performed

with Qubit dsDNA HS Assay kit (ThermoFisher Scientific). Final libraries were run on

Agilent Bioanalyzer for quality check. TrueSeq libraries were then sequenced as

necessary for their desired length as paired end 2×151 and 2×301 read multiplex runs

on MiSeq platform (Illumina) for pre-cursor and mature MAPT isoforms respectively.

Sequenced reads have been uploaded to the NCBI SRA database under BioProject ID

PRJNA762079.

In vivo DMS-MaP probing for *SSU ribosome*

For in vivo ribosomal structure data, we used approximately 10 million T47D cells in 10

cm plates for each condition. We removed the growth media, added 1.8 mL of bicine

buffered growth medium (200 mM Bicine pH 8.3) followed by treatment at 37°C with 200

uL of DMS diluted in ethanol (1.25% final DMS) for 5 min. For the negative control

(unmodified RNA), instead of DMS, an equivalent amount of ethanol was added to the

same number of T47D cells. After incubation, all reactions were neutralized by

addition of ice cold 20% by volume β-mercaptoethanol and kept on ice for 5 minutes.

Total RNA was extracted by Trizol (ThermoFisher Scientific), chloroform and isoamyl

alcohol using phase lock heavy tubes (5PRIME Phase Lock Gel). RNA was purified

using Purelink RNA mini kit (ThermoFisher Scientific), treated with TURBODNase

(ThermoFisher Scientific) and quantified with NanoDrop™ spectrophotometer.

DMS-MaP cDNA synthesis, library construction and sequencing for *SSU*

ribosome

Purified RNA (5 ug) was reverse transcribed using Random Primer 9 (NEB) and

SuperScript II reverse transcriptase under error prone conditions as described Smola et

al., 2015. The resultant cDNA was purified using G50 column (GE healthcare) and

subjected to second strand synthesis (NEBNext Second Strand Synthesis Module).

Standard Nextera DNA library protocol (Illumina) was used to fragment the cDNA and

add sequencing barcodes. All samples were purified using Ampure XP (Beckman

Coulter) beads and quantification of the libraries was performed with Qubit dsDNA HS

Assay kit (ThermoFisher Scientific). Final libraries were run on Agilent Bioanalyzer for

quality check. Libraries were sequenced as paired end 2×151 read multiplex runs on

MiSeq platform (Illumina). Sequenced reads have been uploaded to the NCBI SRA

database under BioProject ID PRJNA762079.

DMS-MaP analysis

Sequenced reads were analyzed using the ShapeMapper pipeline(Busan and Weeks

2018), version (v2.1.4) which calculates the DMS reactivity of each nucleotide *i* as

follows:

$$R_i = mutr_S - mutr_U$$

where *mutr$_S$* is the mutation rate in the sample treated with DMS, *mutr$_U$* is the mutation

rate in the untreated control. DMS reactivities were normalized within a sample and per

nucleotide type (A, C, U, G) using the normalization method described in Busan and

Weeks, 2018. DMS probing data were collected for the Exon 9-Exon 11 and Exon 9-

Exon 10-Exon 11 junctions using a single pair of primers listed in Supplementary file 4.

The ShapeMapper pipeline ran for the two junctions in a single run with reference

sequences for both junctions entered in one FASTA file. For the SSU, sequenced reads

were first aligned to the SSU ribosome sequence using Bowtie2 parameters from Busan

and Weeks, 2018. Using samtools, alignments with MAPQ score greater than 10 were

kept, sorted, and converted back into FASTQ files after which the ShapeMapper

pipeline was executed.

Updating DMS parameters for RNAstructure using SSU ribosome data from T47D

cells

To use DMS data to guide secondary structure prediction by the Rsample (Spasic et al.

2018) component of RNAstructure (Reuter and Mathews 2010), we calibrated the

expected DMS reactivities per nucleotide. Using the SSU ribosome mapping data and

the known secondary structure (Petrov et al. 2014), we determined histograms for DMS

reactivities for unpaired nucleotides, nucleotides paired at helix ends, and nucleotides

paired in base pairs stacked between two other base pairs. These DMS histograms can

be invoked by Rsample with the "--DMS" command line switch as part of RNAstructure

6.4 or later. The histograms had long tails to relatively high reactivities. We empirically

found that limiting reactivities in the histograms and in the input data to a reactivity of 5

(where higher values are set to 5) gave the best performance at improving SSU rRNA

secondary structure. The "--max 5" parameter is used with Rsample to apply this

limitation. Pre-release of Rsample code including in vivo DMS parameters is included

as a zip file for review, and will be included in RNAstructure 6.4.

Base-pairing probabilities for SSU

The partition function for the SSU was generated using Rsample, using either the

sequence or using the sequence and the DMS reactivities. All possible i-j base pairing

probabilities were summed for each nucleotide i to generate a base pairing probability

per nucleotide i.

ROC curves for predicting SSU base pairs

Using the known secondary structure of the SSU, we assigned a nucleotide as either 0

or 1 if it was paired or unpaired. DMS reactivities were used to predict whether a

nucleotide was paired; the higher the DMS reactivity, the more likely a nucleotide is

unpaired. Base pairing probabilities were subtracted from 1 to obtain the probability that

a base was unpaired, with 0 implying base was paired and 1 implying that base was

unpaired. ROC curves and AUC values were generated using the plotROC (Sachs

2017) R package.

Arc plots

Arc plots were generated using Superfold (Siegfried et al. 2014) modified to process

DMS reactivity data.

Generating structural ensemble of Exon 10-Intron 10 *MAPT* junction

The partition function of the Exon 10-Intron 10 MAPT junction for wild type (WT) and

mutations was calculated with DMS reactivities as restraints using Rsample (Spasic et

al. 2018). The DMS reactivities, which were collected for the WT sequence, were also

used for the mutations to restrain the structural space with the reactivity made NA at the

nucleotide where the mutation occurred. The program stochastic (Reuter and Mathews

2010) was used to sample 1000 structures (in CT format) from the Boltzmann

distribution wherein the likelihood a structure is sampled was proportional to the

probability that it occurred in the distribution (Y. Ding and Lawrence 2003).

t-SNE visualization of structural ensembles of WT, 3R and 4R mutations of Exon

10-Intron 10 *MAPT* junction

Structural ensembles were generated as described above for WT, 3R, and 4R

mutations. For each sequence, the 1000 structures in CT format were converted to dot-

bracket (db) format with ct2dot (Reuter and Mathews 2010), after which the db structure

was transformed into the element format using rnaConvert in the Forgi package

(Kerpedjiev, Höner Zu Siederdissen, and Hofacker 2015). In the element format, every

base is represented by the subtype of RNA structure in which it is found: stem (s),

hairpin (h), loop(m), 5'end(f), and 3'end(t). Hence, each db structure is a string of

characters. These characters were digitized (f, t:0, s:1, h:2, m:3) to create a numerical

matrix with 1000 rows and 234 columns, the length of the exon-intron junction.

Combining the matrices for the three sequences resulted in a 3000×234 matrix. This

matrix was entered into the tSNE function from the scikit-learn python package

(Pedregosa et al. 2011) and dimensionality was reduced to a 3000×2 matrix which was

then plotted with ggplot2 (Wickham 2016) in R. The $\Delta G^{\ddagger}$ of unfolding of the splice site

was calculated for each of the 3000 structures as described below. Source file for

Figure 3B lists t-SNE reduced data with corresponding free energies.

Determining representative structures for clusters in t-SNE plot

The 3000×2 matrix, the result of t-SNE dimensionality reduction, was clustered using k-

means clustering with the k-means function from the scikit-learn python package

(Pedregosa et al. 2011). The value of k was set to 5 as determined visually. A custom

python script was used to deduce the representative structure for each cluster by first

calculating the most common RNA structure subtype at each position. The actual

structure in the ensemble, most similar to the RNA structure with the most common

subtypes at each position, was then determined and deemed to be the representative

structure of that cluster.

Visualizing density of structures in t-SNE plot

A custom python script was written. For the WT and, 3R, and 4R mutant sequences, a

meshgrid was created for the three matrices using a 1000-point interpolation and

NumPy (Harris et al. 2020) meshgrid function which returns two two-dimensional arrays

that represent all the possible x-y coordinates for the three matrices. A gaussian kernel

was next fit and evaluated for each 1000×2 matrix with SciPy gaussian_kde function

(Virtanen et al. 2020) to smoothen over the meshgrid. Contour lines were generated for

the smoothed data with Matplotlib contour function (Hunter 2007) and contourf was

used to plot the data.

Quantifying nucleotides around the 5' splice site in cryo-EM structure

The Protein Databank (PDB) files for Pre-B (PDB ID: 6QX9), B (PDB ID: 5O9Z), Pre-

B^{act} (PDB ID: 7ABF) and B^{act} (PDB ID: 5Z56) complexes were downloaded from the

PDB website. A custom python script was used to extract pre-mRNA from each PDB

file. The number of nucleotides were counted for mRNA found near the 5' splice site.

The result was visually confirmed by visualizing the PDB on PyMol.

Calculating $\Delta G^{\ddagger}$ of unfolding of a region of interest

The $\Delta G^{\ddagger}$ of a structure was calculated using the efn2 program in RNAstructure (Reuter

and Mathews 2010). This represents the non-equilibrium unfolding energy of the region

as the sequence is not allowed to refold after unfolding (Mustoe et al. 2018). The base

pairs within a region of interest were removed using a custom python script. The $\Delta G^{\ddagger}$ of

the "unfolded" structure was next re-calculated with efn2. The $\Delta G^{\ddagger}$ of unfolding of a

region was the subtraction of the $\Delta G^{\ddagger}$ of the original structure from the $\Delta G^{\ddagger}$ of the

unfolded structure. For example, for determining the $\Delta G^{\ddagger}$ of unfolding of the splice site,

we removed all base pairs within the last 3 nucleotides of the exon and the first 6

nucleotides of the intron.

1002

Calculating the change in strength of SRE motifs

Splice Site: Strength of the WT splice site was calculated with MaxEntScan (Yeo and

Burge 2004). Strength was recalculated in the presence of splice site mutations either in

the last 3 bases of Exon 10 or first 6 bases of Intron 10. WT strength was subtracted

from the mutant strength: a 0 implied no change in splice site strength, positive values

implied that a mutation made splice site stronger, resulting in increased inclusion of

Exon 10, and negative values implied that a mutation made splice site weaker and

decreased inclusion of Exon 10.

Enhancers and Silencers: Overrepresented hexamers in cell-based screens of general

exonic and intronic splicing enhancers (ESEs, ISEs) and silencers (ESSs, ISSs) were

obtained from Fairbrother et al., 2002, Wang et al., 2004, Wang, Ma et al., 2012 and

Wang, Xiao et al., 2012. Position weight matrices (PWMs) of hexamers for each

category were re-calculated as described in Fairbrother et al., 2002 and collated in

Supplementary file 5. There were 8 clusters of ESE motifs, 7 of ESS motifs, 7 of ISE

motifs, and 8 clusters of ISS motifs; each cluster had an associated PWM. For each

PWM, a threshold strength was found by taking the 95th percentile value of strength of

all possible k-mers of PWM length. This threshold was used to determine whether there

was a valid SRE motif at a particular position. The strength of the PWM motif was

calculated across the exon-intron junction using a sliding window for both WT sequence

and per mutation. The only windows that differed were around the location of the

mutation, and only windows with strength above the threshold were considered. The

WT strength was subtracted from the mutation strength for each window, and all

windows were then summed to yield a ∆strength for every PWM per mutation. The

average of the non-zero ∆strengths was calculated for ESE, ESS, ISE and ISS

categories. The ESE and ISE ∆strengths were summed to obtain an enhancer strength,

and the ESS and ISS ∆strengths were summed to obtain a silencer strength.

Supplementary file 6 presents all SRE ∆strengths for the 47 mutations and 55 VUSs.

1030

Calculating the change in strength of RBP motifs

Position Frequency Matrices (PFMs) were available from Ray et al. 2013 for the

following RBPs: SRSF1, SRSF2, SRSF7, SRSF9, SRSF10, PCBP2, RBM4 and SFPQ.

PFMs were converted into PWMs by normalizing frequencies to 0.25 (Prior probability

for nucleotide frequency) and calculating the log2 value. Overrepresented hexamers

were available from Dominguez et al., 2018 for the following RBPs: SRSF11, SRSF4,

SRSF5 and SRSF8. PFMs for those RBPs were calculated as described in Fairbrother

et al., 2002. ∆strength in RBP motifs were calculated the same way as SRE motifs. The

average of non-zero values of RBPs implicated in either the inclusion or exclusion of

Exon 10 was computed separately. All RBP ∆strengths for the 47 mutations are found in

Supplementary file 6.

1042

Models and bootstrapping

Exon 10 PSI was limited to values between 0 and 1 with 0 signifying that no transcripts

had Exon 10 and 1 implying that all transcripts had Exon 10. Hence, standard linear

regression was no longer appropriate and features were fit with a beta regression model

to Exon 10 PSI. Regression parameters were determined using the betareg package

(Cribari-Neto and Zeileis 2010) in R. Bootstrapping was performed by sampling without

replacement 70% of the mutations to train and test the model and calculating the

Pearson correlation coefficient (R^2) between true values and predictions of the sample.

This bootstrapping was executed 10 times resulting in a range of R^2s, ensuring that no

subset of mutations skewed model performance. Since there were only 4 mutations that

maintained the wildtype 3R to 4R ratio in our training set, we added 3 additional variants

of unknown significance (VUSs) from the Single Nucleotide Polymorphism database

(dbSNP) which we experimentally verified preserved the wildtype splicing pattern (see

Supplementary file 7 for gel). WT VUSs tested and added to the training set were

assigned a PSI of 0.5 to indicate equivalence to the WT sequence. Eq. **1**, the structure

ensemble model, uses four characteristics describing ***X***, the ∆G° of unfolding of the

region of interest around the exon-intron junction for 1000 structures in the ensemble.

Eq. **2**, the minimum free energy (MFE) model, uses just ***Y***, the ∆G° of unfolding of the

exon-intron junction found within the spliceosome at the B^{act} stage for the single MFE

structure. Eq. **3**, the splice site model, uses the difference in splice site strength

between WT sequence and a mutation where *SS* represents splice site. Eq. **4**, the

combined SRE model, uses the difference in SRE strength between WT sequence and

a mutation where *SS* represents splice site, *E* represents enhancer, and *S* represents

silencer. Eq. **5**, the RBP model, uses the difference in RBP motif strength between WT

sequence and a mutation where *Ex* represents RBPs involved in the exclusion of Exon

10 and *In* represents RBPs involved in the inclusion of Exon 10. Eq. **6** is the interactive

model between structure and SRE, and Eq.**7** is the additive model. *isNonSynonymous,*

isSynonymous and *isIntronic* represent the category of mutation and is either 0 or 1.

Supplementary file 6 summarizes the performance of the models and features utilized.

1072

1073 $$PSI \sim Mean(X) + SD(X) + Skew(X) + Kurtosis(X) \qquad [\mathbf{1}]$$

1074

1075 $$PSI \sim Y \quad [\mathbf{2}]$$

1076

1077 $$PSI \sim \Delta SS \quad [\mathbf{3}]$$

1078

$$PSI \sim \Delta E + \Delta S + \Delta SS \quad [\mathbf{4}]$$

1080

$$PSI \sim \Delta Ex + \Delta In \quad [\mathbf{5}]$$

1082

$$PSI \sim [Mean(X) + SD(X) + Skew(X) + Kurtosis(X)] * [isSynonymous + isIntronic]$$
$$+ [\Delta E + \Delta S + \Delta SS] * [isNonSynonymous] \quad [\mathbf{6}]$$

1085

$$PSI \sim [Mean(X) + SD(X) + Skew(X) + Kurtosis(X)] + [\Delta E + \Delta S + \Delta SS] \quad [\mathbf{7}]$$

1087

Clustering changes in structural and SRE features

For each feature, non-zero values greater than the 95th percentile value were set to the 95th percentile or, if less than the 5th percentile value, were set to the 5th percentile for visualization, after which all values were normalized to the maximum absolute value. Silencer Δstrength and mean ΔG° of unfolding of exon-intron junction of ensemble were inverted to follow the visualization such that values closer to 1 would result in greater Exon 10 inclusion and values closer to 0 would result in lower Exon 10 inclusion. Features were then assigned values 0 or 1 depending on whether the feature changed at all in the presence of the mutation. These digitized features were first clustered by hierarchal clustering resulting in 6 clusters. Each individual cluster was clustered again by hierarchal clustering but using the normalized feature values instead of 0s and 1s.

Splicing Assays

HEK-293 cells (ATCC CRL-1573) were grown at 37°C in 5% CO_2 in Dulbecco's Modified Eagle Medium (Gibco) supplemented with 10% FBS (Omega Scientific) and 0.5% Penicillin Streptomycin (Gibco). The wild type splicing reporter plasmid was generously provided by the Roca lab and is described in Tan et al., 2019. Single-nucleotide point mutations were generated using a Q5 site-directed mutagenesis kit (NEB) and confirmed by Sanger sequencing, or custom ordered directly from GenScript. Reporter plasmids (2 μg) were transfected into HEK-293 cells in 6-well plates at ~60-

90% confluency using Lipofectamine 3000 (ThermoFisher Scientific). Cells were harvested after 1 day by aspirating the media and resuspending the cells in 1 mL Trizol reagent (ThermoFisher Scientific). RNA was isolated using the PureLink RNA Isolation Kit (ThermoFisher Scientific) with on-column DNase treatment, following manufacturer's instructions. RNA (1 µg) was reverse transcribed to cDNA using Superscript VILO reverse transcriptase (ThermoFisher Scientific). Reverse transcriptions were performed by annealing (25°C 10 minutes), extension (50°C 10 minutes), and inactivation (85°C 10 min) steps. Heat-inactivated controls were prepared by heating the reaction without RNA at 85°C for 10 minutes prior to adding RNA, then following the described reaction conditions. The cDNA was PCR amplified with NEB Q5 HotStart polymerase (NEB) using splicing assay primers from IDT (AGACCCAAGCTGGCTAGCGTT forward, GAGGCTGATCAGCGGGTTTAAAC reverse) with 25 cycles. PCR product was purified and concentrated using the PureLink PCR micro clean up kit (ThermoFisher Scientific), following manufacturer's instructions. Splicing products were visualized by loading ~200 ng of DNA on a 2% agarose gel in 1X tris-acetate EDTA (TAE) buffer and staining with ethidium bromide. Gel images were quantified with ImageJ.

Supplementary files, figure source files, SNRNASMs and code are available at GitHub repository: https://git.io/JuSW8

Acknowledgements

This work was supported by the US National Institutes of Health R01 HL111527 and R35 GM 140844 to A.L. and R01 GM076485 to D.M. The authors wish to thank the Roca Lab for providing wildtype splicing reporter plasmids, Dr. Zefeng Wang for intronic splicing enhancer and silencer motifs, and Drs. Peter Castaldi, John Platig and Kevin Weeks for insightful discussions.

Competing Interests

The authors have declared that no competing interests exist.

References

Adivarahan, Srivathsan, Nathan Livingston, Beth Nicholson, Samir Rahman, Bin Wu, Olivia S. Rissland,
and Daniel Zenklusen. 2018. "Spatial Organization of Single MRNPs at Different Stages of the Gene
Expression Pathway." *Molecular Cell* 72 (4): 727-738.e5.
https://doi.org/10.1016/J.MOLCEL.2018.10.010.

Andreadis, Athena. 2005. "Tau Gene Alternative Splicing: Expression Patterns, Regulation and
Modulation of Function in Normal Brain and Neurodegenerative Diseases." *Biochimica et Biophysica
Acta - Molecular Basis of Disease* 1739 (2): 91–103. https://doi.org/10.1016/j.bbadis.2004.08.010.

———. 2012. "Tau Splicing and the Intricacies of Dementia." *Journal of Cellular Physiology* 227 (3):
1220–25. https://doi.org/10.1002/jcp.22842.

Bahl, Ethan, Tanner Koomar, and Jacob J Michaelson. 2017. "CerebroViz: An R Package for Anatomical
Visualization of Spatiotemporal Brain Data." *Bioinformatics (Oxford, England)* 33 (5): 762–63.
https://doi.org/10.1093/bioinformatics/btw726.

Baralle, Francisco E, and Jimena Giudice. 2017. "Alternative Splicing as a Regulator of Development and
Tissue Identity." *Nature Reviews Molecular Cell Biology* 18 (7): 437–51.
https://doi.org/10.1038/nrm.2017.27.

Bertram, Karl, Dmitry E Agafonov, Olexandr Dybkov, Berthold Kastner, Reinhard Lü, and Holger Stark
Correspondence. 2017. "Cryo-EM Structure of a Pre-Catalytic Human Spliceosome Primed for
Activation." *Cell* 170: 701-706.e11. https://doi.org/10.1016/j.cell.2017.07.011.

Blanchette, Marco, and Benoit Chabot. 1997. "A Highly Stable Duplex Structure Sequesters the 5' Splice
Site Region of HnRNP A1 Alternative Exon 7B." *RNA*, no. 3: 405–19.

Broderick, Jennifer, Junning Wang, and Athena Andreadis. 2004. "Heterogeneous Nuclear
Ribonucleoprotein E2 Binds to Tau Exon 10 and Moderately Activates Its Splicing." *Gene* 331 (1–2):
107–14. https://doi.org/10.1016/j.gene.2004.02.005.

Buratti, Emanuele, and Francisco E Baralle. 2004. "Influence of RNA Secondary Structure on the Pre-
MRNA Splicing Process." *MOLECULAR AND CELLULAR BIOLOGY* 24 (24): 10505–14.
https://doi.org/10.1128/MCB.24.24.10505–10514.2004.

Busan, Steven, and Kevin M Weeks. 2018. "Accurate Detection of Chemical Modifications in RNA by
Mutational Profiling (MaP) with ShapeMapper 2." *RNA* 24 (2): 143–48.
https://doi.org/10.1261/rna.061945.117.

Catarina Silva, M., and Stephen J. Haggarty. 2020. "Tauopathies: Deciphering Disease Mechanisms to
Develop Effective Therapies." *International Journal of Molecular Sciences* 21 (23): 1–49.
https://doi.org/10.3390/ijms21238948.

Charenton, Clément, Max E. Wilkinson, and Kiyoshi Nagai. 2019. "Mechanism of 5' Splice Site Transfer
for Human Spliceosome Activation." *Science* 364 (6438): 362–67.

https://doi.org/10.1126/science.aax3289.

Chen, Jonathan L., Walter N. Moss, Adam Spencer, Peiyuan Zhang, Jessica L. Childs-Disney, and

Matthew D. Disney. 2019. "The RNA Encoding the Microtubule-Associated Protein Tau Has

Extensive Structure That Affects Its Biology." *PLOS ONE* 14 (7): e0219210.

https://doi.org/10.1371/journal.pone.0219210.

Clark, Lorraine N, Parvoneh Poorkaj, Zbigniew Wszolek, Daniel H Geschwind, Ziad S Nasreddine, Bruce

Miller, Diane Li, et al. 1998. "Pathogenic Implications of Mutations in the Tau Gene in Pallido-Ponto-

Nigral Degeneration and Related Neurodegenerative Disorders Linked to Chromosome 17."

Proceedings of the National Academy of Sciences of the United States of America 95 (22): 13103–

7. https://doi.org/10.1073/pnas.95.22.13103.

Cribari-Neto, Francisco, and Achim Zeileis. 2010. "Beta Regression in {R}." *Journal of Statistical Software*

34 (2): 1–24. https://doi.org/10.18637/jss.v034.i02.

Cruz, José Almeida, and Eric Westhof. 2009. "The Dynamic Landscapes of RNA Architecture." *Cell* 136

(4): 604–9. https://doi.org/10.1016/j.cell.2009.02.003.

D'Souza, I, P Poorkaj, M Hong, D Nochlin, V. M.- Y. Lee, T D Bird, and G D Schellenberg. 1999.

"Missense and Silent Tau Gene Mutations Cause Frontotemporal Dementia with Parkinsonism-

Chromosome 17 Type, by Affecting Multiple Alternative RNA Splicing Regulatory Elements."

Proceedings of the National Academy of Sciences 96 (10): 5598–5603.

https://doi.org/10.1073/pnas.96.10.5598.

D'Souza, Ian, and Gerard D. Schellenberg. 2005. "Regulation of Tau Isoform Expression and Dementia."

Biochimica et Biophysica Acta - Molecular Basis of Disease, January 3, 2005.

https://doi.org/10.1016/j.bbadis.2004.08.009.

———. 2006. "Arginine/Serine-Rich Protein Interaction Domain-Dependent Modulation of a Tau Exon 10

Splicing Enhancer: Altered Interactions and Mechanisms for Functionally Antagonistic FTDP-17

Mutations Δ280K and N279K." *Journal of Biological Chemistry* 281 (5): 2460–69.

https://doi.org/10.1074/jbc.M505809200.

D'Souza, Ian, and Gerard David Schellenberg. 2000. "Determinants of 4-Repeat Tau Expression.

Coordination between Enhancing and Inhibitory Splicing Sequences for Exon 10 Inclusion." *Journal*

of Biological Chemistry 275 (23): 17700–709. https://doi.org/10.1074/jbc.M909470199.

Dethoff, Elizabeth A., Jeetender Chugh, Anthony M. Mustoe, and Hashim M. Al-Hashimi. 2012.

"Functional Complexity and Regulation through RNA Dynamics." *Nature*. Nature Publishing Group.

https://doi.org/10.1038/nature10885.

Ding, Shaohong, Jianhua Shi, Wei Qian, Khalid Iqbal, Inge Grundke-Iqbal, Cheng Xin Gong, and Fei Liu.

2012. "Regulation of Alternative Splicing of Tau Exon 10 by 9G8 and Dyrk1A." *Neurobiology of*

Aging 33 (7): 1389–99. https://doi.org/10.1016/j.neurobiolaging.2010.11.021.

Ding, Ye, and Charles E Lawrence. 2003. "A Statistical Sampling Algorithm for RNA Secondary Structure

Prediction." *Nucleic Acids Research* 31 (24): 7280–7301. https://doi.org/10.1093/nar/gkg938.

Dominguez, Daniel, Peter Freese, Maria S. Alexis, Amanda Su, Myles Hochman, Tsultrim Palden,

Cassandra Bazile, et al. 2018. "Sequence, Structure, and Context Preferences of Human RNA

Binding Proteins." *Molecular Cell* 70(5): 854–867.e9. https://doi.org/10.1016/j.molcel.2018.05.001.

Donahue, Christine P, Christina Muratore, Jane Y Wu, Kenneth S Kosik, and Michael S Wolfe. 2006.

"Stabilization of the Tau Exon 10 Stem Loop Alters Pre-MRNA Splicing." *Journal of Biological*

Chemistry 281 (33): 23302–6. https://doi.org/10.1074/jbc.C600143200.

Fairbrother, William G., Ru-Fang Yeh, Phillip A. Sharp, and Christopher B. Burge. 2002. "Predictive

Identification of Exonic Splicing Enhancers in Human Genes." *Science* 297: 1007–13.

http://dx.doi.org/10.1126/science.1073774.

Ferrari, Silvia L.P., and Francisco Cribari-Neto. 2004. "Beta Regression for Modelling Rates and

Proportions." *Journal of Applied Statistics* 31 (7): 799–815.

https://doi.org/10.1080/0266476042000214501.

Gao, Lei, Junning Wang, Yingzi Wang, and Athena Andreadis. 2007. "SR Protein 9G8 Modulates Splicing

of Tau Exon 10 via Its Proximal Downstream Intron, a Clustering Region for Frontotemporal

Dementia Mutations." *Molecular and Cellular Neuroscience* 34 (1): 48–58.

https://doi.org/10.1016/j.mcn.2006.10.004.

Gao, Qing Sheng, John Memmott, Robert Lafyatis, Stefan Stamm, Gavin Screaton, and Athena

Andreadis. 2000. "Complex Regulation of Tau Exon 10, Whose Missplicing Causes Frontotemporal

Dementia." *Journal of Neurochemistry* 74 (2): 490–500. https://doi.org/10.1046/j.1471-

4159.2000.740490.x.

Giegé, Richard, Frank Jühling, Joern Pütz, Peter Stadler, Claude Sauter, and Catherine Florentz. 2012.

"Structure of Transfer RNAs: Similarity and Variability." *Wiley Interdisciplinary Reviews: RNA* 3 (1):

37–61. https://doi.org/10.1002/WRNA.103.

Goedert, M., M. G. Spillantini, R. A. Crowther, S. G. Chen, P. Parchi, M. Tabaton, D. J. Lanska, et al.

1999. "Tau Gene Mutation in Familial Progressive Subcortical Gliosis." *Nature Medicine* 5 (4): 454–

57. https://doi.org/10.1038/7454.

Goedert, M., M G. Spillantini, R. Jakes, D Rutherford, and R A Crowther. 1989. "Multiple Isoforms of

Human Microtubule-Associated Protein Tau: Sequences and Localization in Neurofibrillary Tangles

of Alzheimer's Disease." *Neuron* 3 (4): 519–26. https://doi.org/10.1016/0896-6273(89)90210-9.

Grover, Andrew, Henry Houlden, Matt Baker, Jennifer Adamson, Jada Lewis, Guy Prihar, Stuart

Pickering-Brown, Karen Duff, and Mike Hutton. 1999. "5' Splice Site Mutations in Tau Associated

with the Inherited Dementia FTDP-17 Affect a Stem-Loop Structure That Regulates Alternative

Splicing of Exon 10*." *Journal of Biological Chemistry* 274 (21): 15134–43.

http://www.jbc.org/content/274/21/15134.full.pdf.

Halvorsen, Matthew, Joshua S Martin, Sam Broadaway, and Alain Laederach. 2010. "Disease-Associated

Mutations That Alter the RNA Structural Ensemble." *PLoS Genetics* 6 (8).

https://doi.org/10.1371/journal.pgen.1001074.

Harris, Charles R, K Jarrod Millman, Stéfan J van der Walt, Ralf Gommers, Pauli Virtanen, David

Cournapeau, Eric Wieser, et al. 2020. "Array Programming with {NumPy}." *Nature* 585 (7825): 357–

62. https://doi.org/10.1038/s41586-020-2649-2.

Hasegawa, Masato, Michael J Smith, Masaaki Iijima, Takeshi Tabira, and Michel Goedert. 1999. "FTDP-

17 Mutations N279K and S305N in Tau Produce Increased Splicing of Exon 10." *FEBS Letters* 443

(2): 93–96. https://doi.org/10.1016/S0014-5793(98)01696-2.

Hefti, Marco M, Kurt Farrell, Soong Ho Kim, Kathryn R Bowles, Mary E Fowkes, Towfique Raj, and John

F Crary. 2018. "High-Resolution Temporal and Regional Mapping of MAPT Expression and Splicing

in Human Brain Development." *PLoS ONE* 13 (4). https://doi.org/10.1371/journal.pone.0195771.

Herzel, Lydia, Diana S. M. Ottoz, Tara Alpert, and Karla M. Neugebauer. 2017. "Splicing and

Transcription Touch Base: Co-Transcriptional Spliceosome Assembly and Function." *Nature*

Reviews Molecular Cell Biology. https://doi.org/10.1038/nrm.2017.63.

Homan, Philip J, Oleg V Favorov, Christopher A Lavender, Olcay Kursun, Xiyuan Ge, Steven Busan,

Nikolay V Dokholyan, and Kevin M Weeks. 2014. "Single-Molecule Correlated Chemical Probing of

RNA." *Proceedings of the National Academy of Sciences* 111 (38): 13858–63.

https://doi.org/10.1073/pnas.1407306111.

Hunter, J D. 2007. "Matplotlib: A 2D Graphics Environment." *Computing in Science \& Engineering* 9 (3):

90–95. https://doi.org/10.1109/MCSE.2007.55.

Hutton, M., C. L. Lendon, P. Rizzu, M. Baker, Susanne Froelich, H. H. Houlden, S. Pickering-Brown, et al.

1998. "Association of Missense and 5'-Splice-Site Mutations in Tau with the Inherited Dementia

FTDP-17." *Nature* 393 (6686): 702–4. https://doi.org/10.1038/31508.

Ingolia, Nicholas T. 2016. "Ribosome Footprint Profiling of Translation throughout the Genome." *Cell* 165

(1): 22–33. https://doi.org/10.1016/j.cell.2016.02.066.

Iseki, Eizo, Takehiko Matsumura, Wami Marui, Hiroaki Hino, Toshinari Odawara, Naoya Sugiyama,

Kyoko Suzuki, Hajime Sawada, Tetsuaki Arai, and Kenji Kosaka. 2001. "Familial Frontotemporal

Dementia and Parkinsonism with a Novel N296H Mutation in Exon 10 of the Tau Gene and a

Widespread Tau Accumulation in the Glial Cells." *Acta Neuropathologica* 102 (3): 285–92.

http://dx.doi.org/10.1007/s004010000333.

Jiang, Zhihong, Jocelyn Cote, Jennifer M Kwon, Alison M Goate, and Jane Y Wu. 2000. "Aberrant

Splicing of Tau Pre-MRNA Caused by Intronic Mutations Associated with the Inherited Dementia

Frontotemporal Dementia with Parkinsonism Linked to Chromosome 17." *Molecular and Cellular*

Biology 20 (11): 4036–48. https://doi.org/10.1128/MCB.20.14.5360-5360.2000.

Kar, Amar, Kazuo Fushimi, Xiaohong Zhou, Payal Ray, Chen Shi, X. Chen, Zhiren Liu, She Chen, and

Jane Y Wu. 2011. "RNA Helicase P68 (DDX5) Regulates Tau Exon 10 Splicing by Modulating a

Stem-Loop Structure at the 5' Splice Site." *Molecular and Cellular Biology* 31 (9): 1812–21. https://doi.org/10.1128/MCB.01149-10.

Kar, Amar, Necat Havlioglu, Woan Yuh Tarn, and Jane Y Wu. 2006. "RBM4 Interacts with an Intronic Element and Stimulates Tau Exon 10 Inclusion." *Journal of Biological Chemistry* 281 (34): 24479–88. https://doi.org/10.1074/jbc.M603971200.

Kerpedjiev, Peter, Christian Höner Zu Siederdissen, and Ivo L Hofacker. 2015. "Predicting RNA 3D Structure Using a Coarse-Grain Helix-Centered Model." *RNA* 21 (6): 1110–21. https://doi.org/10.1261/rna.047522.114.

Kondo, Shinichi, Noriaki Yamamoto, Tomohiko Murakami, Masayo Okumura, Akila Mayeda, and Kazunori Imaizumi. 2004. "Tra2β, SF2/ASF and SRp30c Modulate the Function of an Exonic Splicing Enhancer in Exon 10 of Tau Pre-MRNA." *Genes to Cells* 9 (2): 121–30. https://doi.org/10.1111/j.1356-9597.2004.00709.x.

Kumar, Jayashree. 2020a. "AnVIL_GTEx_V8_hg38_JKversion." Terra. 2020. https://app.terra.bio/#workspaces/fccredits-iridium-pear-1617/AnVIL_GTEx_V8_hg38_JKversion.

———. 2020b. "ExtractReadsFromBamFileForAGivenRegion_SortAndIndex." Terra. 2020. https://app.terra.bio/#workspaces/jk-billing-terra87/AnVIL_GTEx_V8_hg38_JKversion_NewWS/workflows/JayashreeKumar_UNCChapelHill/https://app.terra.bio/#workspaces/fccredits-iridium-pear-1617/AnVIL_GTEx_V8_hg38_JKversion/workflows/JayashreeKumar_UNCChapelHil.

Lai, Wan Jung C., Mohammad Kayedkhordeh, Erica V. Cornell, Elie Farah, Stanislav Bellaousov, Robert Rietmeijer, Enea Salsi, David H. Mathews, and Dmitri N. Ermolenko. 2018. "MRNAs and LncRNAs Intrinsically Form Secondary Structures with Short End-to-End Distances." *Nature Communications* 9 (1). https://doi.org/10.1038/s41467-018-06792-z.

Lin, Chien Ling, Allison J Taggart, and William G Fairbrother. 2016. "RNA Structure in Splicing: An Evolutionary Perspective." *RNA Biology* 13 (9): 766–71. https://doi.org/10.1080/15476286.2016.1208893.

Lisowiec, Jolanta, Dorota Magner, Elzbieta Kierzek, Elzbieta Lenartowicz, and Ryszard Kierzek. 2015. "Structural Determinants for Alternative Splicing Regulation of the MAPT Pre-MRNA." *RNA Biology* 12 (3): 330–42. https://doi.org/10.1080/15476286.2015.1017214.

Lonsdale, John, Jeffrey Thomas, Mike Salvatore, Rebecca Phillips, Edmund Lo, Saboor Shad, Richard Hasz, et al. 2013. "The Genotype-Tissue Expression (GTEx) Project." *Nature Genetics* 45 (6): 580–85. https://doi.org/10.1038/ng.2653.

Maaten, Laurens Van Der, and Geoffrey Hinton. 2008. "Visualizing Data Using T-SNE." *Journal of Machine Learning Research* 9: 2579–2605. https://lvdmaaten.github.io/publications/papers/JMLR_2008.pdf.

Majounie, Elisa, William Cross, Victoria Newsway, Allissa Dillman, Jana Vandrovcova, Christopher M.

Morris, Michael A. Nalls, et al. 2013. "Variation in Tau Isoform Expression in Different Brain Regions and Disease States." *Neurobiology of Aging* 34 (7): 1922.e7-1922.e12. https://doi.org/10.1016/j.neurobiolaging.2013.01.017.

Matera, A. Gregory, and Zefeng Wang. 2014. "A Day in the Life of the Spliceosome." *Nature Reviews Molecular Cell Biology* 15 (2): 108–21. https://doi.org/10.1038/nrm3742.

McManus, C. Joel, and Brenton R. Graveley. 2011. "RNA Structure and the Mechanisms of Alternative Splicing." *Current Opinion in Genetics and Development.* https://doi.org/10.1016/j.gde.2011.04.001.

Mirra, Suzanne S ;, Jill R ; Murrell, Marla ; Gearing, and Maria G Spillantini. 1999. "Tau Pathology in a Family with Dementia and a P301L Mutation in Tau." *Journal of Neuropathology and Experimental Neurology* 58 (4): 335. https://search.proquest.com/docview/229718949/fulltextPDF/AA5FABCB40F44E3FPQ/1?accountid =14244.

Mustoe, Anthony M., Steven Busan, Greggory M. Rice, Christine E. Hajdin, Brant K. Peterson, Vera M. Ruda, Neil Kubica, Razvan Nutiu, Jeremy L. Baryza, and Kevin M. Weeks. 2018. "Pervasive Regulatory Functions of MRNA Structure Revealed by High-Resolution SHAPE Probing." *Cell* 173 (1): 181-195.e18. https://doi.org/10.1016/j.cell.2018.02.034.

Park, Sun Ah, Sang Il Ahn, and Jean Marc Gallo. 2016. "Tau Mis-Splicing in the Pathogenesis of Neurodegenerative Disorders." *BMB Reports* 49 (8): 405–13. https://doi.org/10.5483/BMBRep.2016.49.8.084.

Pedregosa, F, G Varoquaux, A Gramfort, V Michel, B Thirion, O Grisel, M Blondel, et al. 2011. "Scikit-Learn: Machine Learning in Python." *Journal of Machine Learning Research* 12: 2825–30.

Petrov, Anton S., Chad R. Bernier, Burak Gulen, Chris C. Waterbury, Eli Hershkovits, Chiaolong Hsiao, Stephen C. Harvey, et al. 2014. "Secondary Structures of RRNAs from All Three Domains of Life." *PLoS ONE* 9 (2): e88222. https://doi.org/10.1371/journal.pone.0088222.

Qian, Wei, Khalid Iqbal, Inge Grundke-Iqbal, Cheng Xin Gong, and Fei Liu. 2011. "Splicing Factor SC35 Promotes Tau Expression through Stabilization of Its MRNA." *FEBS Letters* 585 (6): 875–80. https://doi.org/10.1016/j.febslet.2011.02.017.

Qian, Wei, and Fei Liu. 2014. "Regulation of Alternative Splicing of Tau Exon 10." *Neuroscience Bulletin* 30 (2): 367–77. https://doi.org/10.1007/s12264-013-1411-2.

Ray, Debashish, Hilal Kazan, Kate B Cook, Matthew T Weirauch, Hamed S Najafabadi, Xiao Li, Serge Gueroussov, et al. 2013. "A Compendium of RNA-Binding Motifs for Decoding Gene Regulation." *Nature* 499 (7457): 172–77. https://doi.org/10.1038/nature12311.

Ray, Payal, Amar Kar, Kazuo Fushimi, Necat Havlioglu, Xiaoping Chen, and Jane Y. Wu. 2011. "PSF Suppresses Tau Exon 10 Inclusion by Interacting with a Stem-Loop Structure Downstream of Exon 10." *Journal of Molecular Neuroscience* 45 (3): 453–66. https://doi.org/10.1007/s12031-011-9634-z.

Reuter, Jessica S, and David H Mathews. 2010. "RNAstructure: Software for RNA Secondary Structure

Prediction and Analysis." *BMC Bioinformatics 2010 11:1* 11 (1): 1–9. https://doi.org/10.1186/1471-
2105-11-129.

Rizzu, Patrizia, John C. Van Swieten, Marijke Joosse, Masato Hasegawa, Martijn Stevens, Aad Tibben,
Martinus F. Niermeijer, et al. 1999. "High Prevalence of Mutations in the Microtubule-Associated
Protein Tau in a Population Study of Frontotemporal Dementia in the Netherlands." *American*
Journal of Human Genetics 64 (2): 414–21. https://doi.org/10.1086/302256.

Roca, Xavier, Martin Akerman, Hans Gaus, Andrés Berdeja, C. Frank Bennett, and Adrian R. Krainer.
2012. "Widespread Recognition of 5′ Splice Sites by Noncanonical Base-Pairing to U1 SnRNA
Involving Bulged Nucleotides." *Genes and Development* 26 (10): 1098–1109.
https://doi.org/10.1101/gad.190173.112.

Rouskin, Silvi, Meghan Zubradt, Stefan Washietl, Manolis Kellis, and Jonathan S Weissman. 2014.
"Genome-Wide Probing of RNA Structure Reveals Active Unfolding of MRNA Structures in Vivo. TL
- 505." *Nature* 505 VN- (7485): 701–5. https://doi.org/10.1038/nature12894.

Sachs, Michael C. 2017. "{plotROC}: A Tool for Plotting ROC Curves." *Journal of Statistical Software,*
Code Snippets 79 (2): 1–19. https://doi.org/10.18637/jss.v079.c02.

Sharma, Yogita, Milad Miladi, Sandeep Dukare, Karine Boulay, Maiwen Caudron-Herger, Matthias Groß,
Rolf Backofen, and Sven Diederichs. 2019. "A Pan-Cancer Analysis of Synonymous Mutations."
Nature Communications 10 (1): 1–14. https://doi.org/10.1038/s41467-019-10489-2.

Siegfried, Nathan A, Steven Busan, Greggory M Rice, Julie A E Nelson, and Kevin M Weeks. 2014. "RNA
Motif Discovery by SHAPE and Mutational Profiling (SHAPE-MaP)." *Nature Methods* 11 (9): 959–65.
https://doi.org/10.1038/nmeth.3029.

Singh, Natalia N, Ravindra N Singh, and Elliot J Androphy. 2007. "Modulating Role of RNA Structure in
Alternative Splicing of a Critical Exon in the Spinal Muscular Atrophy Genes." *Nucleic Acids*
Research 35 (2): 371–89. https://doi.org/10.1093/nar/gkl1050.

Smola, Matthew J, Greggory M Rice, Steven Busan, Nathan A Siegfried, and Kevin M Weeks. 2015.
"Selective 2′-Hydroxyl Acylation Analyzed by Primer Extension and Mutational Profiling (SHAPE-
MaP) for Direct, Versatile and Accurate RNA Structure Analysis." *Nature Protocols* 10 (11): 1643–
69. https://doi.org/10.1038/nprot.2015.103.

Song, Yan, Olga B Botvinnik, Michael T Lovci, Boyko Kakaradov, Patrick Liu, Jia L Xu, and Gene W Yeo.
2017. "Single-Cell Alternative Splicing Analysis with Expedition Reveals Splicing Dynamics during
Neuron Differentiation." *Molecular Cell* 67 (1): 148-161.e5.
https://doi.org/10.1016/j.molcel.2017.06.003.

Spasic, Aleksandar, Sarah M Assmann, Philip C Bevilacqua, and David H Mathews. 2018. "Modeling
RNA Secondary Structure Folding Ensembles Using SHAPE Mapping Data." *Nucleic Acids*
Research 46 (1): 314–23. https://doi.org/10.1093/nar/gkx1057.

Spillantini, M. G., Jill R Murrell, Michel Goedert, Martin R Farlow, Aaron Klug, and Bernardino Ghetti.

1998. "Mutation in the Tau Gene in Familial Multiple System Tauopathy with Presenile Dementia."

Proceedings of the National Academy of Sciences 95 (13): 7737–41.

https://doi.org/10.1073/pnas.95.13.7737.

Stenson, Peter D, Edward V Ball, Matthew Mort, Andrew D Phillips, Jacqueline A Shiel, Nick S T Thomas,

Shaun Abeysinghe, Michael Krawczak, and David N Cooper. 2003. "Human Gene Mutation

Database (HGMD): 2003 Update." *Human Mutation* 21 (6): 577–81.

https://doi.org/10.1002/humu.10212.

Sun, Lei, Furqan M. Fazal, Pan Li, James P. Broughton, Byron Lee, Lei Tang, Wenze Huang, Eric T.

Kool, Howard Y. Chang, and Qiangfeng Cliff Zhang. 2019. "RNA Structure Maps across Mammalian

Cellular Compartments." *Nature Structural and Molecular Biology* 26 (4): 322–30.

https://doi.org/10.1038/s41594-019-0200-7.

Tan, Jiazi, Lixia Yang, Alan Ann Lerk Ong, Jiahao Shi, Zhensheng Zhong, Mun Leng Lye, Shiyi Liu, et al.

2019. "A Disease-Causing Intronic Point Mutation C19G Alters Tau Exon 10 Splicing via RNA

Secondary Structure Rearrangement." *Biochemistry*. https://doi.org/10.1021/acs.biochem.9b00001.

Taylor, Katarzyna, and Krzysztof Sobczak. 2020. "Intrinsic Regulatory Role of RNA Structural

Arrangement in Alternative Splicing Control." *International Journal of Molecular Sciences* 21 (14): 1–

35. https://doi.org/10.3390/ijms21145161.

Tazi, Jamal, Nadia Bakkour, and Stefan Stamm. 2009. "Alternative Splicing and Disease." *Biochimica et*

Biophysica Acta (BBA) - Molecular Basis of Disease 1792 (1): 14–26.

https://doi.org/10.1016/J.BBADIS.2008.09.017.

Townsend, Cole, Majety N. Leelaram, Dmitry E. Agafonov, Olexandr Dybkov, Cindy L. Will, Karl Bertram,

Henning Urlaub, Berthold Kastner, Holger Stark, and Reinhard Lührmann. 2020. "Mechanism of

Protein-Guided Folding of the Active Site U2/U6 RNA during Spliceosome Activation." *Science* 370

(6523). https://doi.org/10.1126/science.abc3753.

Varani, Luca, Masato Hasegawa, Maria Grazia Spillantini, Michael J Smith, Jill R Murrell, Bernardino

Ghetti, Aaron Klug, Michel Goedert, and Gabriele Varani. 1999. "Structure of Tau Exon 10 Splicing

Regulatory Element RNA and Destabilization by Mutations of Frontotemporal Dementia and

Parkinsonism Linked to Chromosome 17 (Alternative MRNA Splicing∖intronic Mutation∖stem-Loop

RNA Structure)." *Neurobiology* 96: 8229–34. https://doi.org/10.1073/pnas.96.14.8229.

Virtanen, Pauli, Ralf Gommers, Travis E Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau,

Evgeni Burovski, et al. 2020. "{SciPy} 1.0: Fundamental Algorithms for Scientific Computing in

Python." *Nature Methods* 17: 261–72. https://doi.org/10.1038/s41592-019-0686-2.

Wang, Junning, Qing Sheng Gao, Yingzi Wang, Robert Lafyatis, Stefan Stamm, and Athena Andreadis.

2004. "Tau Exon 10, Whose Missplicing Causes Frontotemporal Dementia, Is Regulated by an

Intricate Interplay of Cis Elements and Trans Factors." *Journal of Neurochemistry* 88 (5): 1078–90.

https://doi.org/10.1046/j.1471-4159.2003.02232.x.

Wang, Yan, Lei Gao, Sze Wah Tse, and Athena Andreadis. 2010. "Heterogeneous Nuclear
Ribonucleoprotein E3 Modestly Activates Splicing of Tau Exon 10 via Its Proximal Downstream
Intron, a Hotspot for Frontotemporal Dementia Mutations." *Gene* 451 (1–2): 23–31.
https://doi.org/10.1016/j.gene.2009.11.006.
Wang, Yang, Meng Ma, Xinshu Xiao, and Zefeng Wang. 2012. "Intronic Splicing Enhancers, Cognate
Splicing Factors and Context-Dependent Regulation Rules." *Nature Structural & Molecular Biology*
19 (10): 1044–52. https://doi.org/10.1038/nsmb.2377.
Wang, Yang, Xinshu Xiao, Jianming Zhang, Rajarshi Choudhury, Alex Robertson, Kai Li, Meng Ma,
Christopher B Burge, and Zefeng Wang. 2012. "A Complex Network of Factors with Overlapping
Affinities Represses Splicing through Intronic Elements." *Nature Structural & Molecular Biology* 20
(1): 36–45. https://doi.org/10.1038/nsmb.2459.
Wang, Zefeng, and Christopher B Burge. 2008. "Splicing Regulation: From a Parts List of Regulatory
Elements to an Integrated Splicing Code." *RNA* 14: 802–13. https://doi.org/10.1261/rna.876308.
Wang, Zefeng, Michael E. Rolish, Gene Yeo, Vivian Tung, Matthew Mawson, and Christopher B. Burge.
2004. "Systematic Identification and Analysis of Exonic Splicing Silencers." *Cell* 119 (6): 831–45.
https://doi.org/10.1016/j.cell.2004.11.010.
Warf, M Bryan, and J Andrew Berglund. 2010. "Role of RNA Structure in Regulating Pre-MRNA Splicing."
Trends in Biochemical Sciences. https://doi.org/10.1016/j.tibs.2009.10.004.
Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
https://ggplot2.tidyverse.org.
Yeo, Gene, and Christopher B Burge. 2004. "Maximum Entropy Modeling of Short Sequence Motifs with
Applications to RNA Splicing Signals." *Journal of Computational Biology* 11 (2–3): 377–94.
https://doi.org/10.1089/1066527041410418.
Zhang, Lingdi, Anne Vielle, Sara Espinosa, and Rui Zhao. 2019. "RNAs in the Spliceosome: Insight from
CryoEM Structures." *Wiley Interdisciplinary Reviews: RNA* 10 (3): 1–11.
https://doi.org/10.1002/wrna.1523.
Zhang, Xiaofeng, Chuangye Yan, Xiechao Zhan, Lijia Li, Jianlin Lei, and Yigong Shi. 2018. "Structure of
the Human Activated Spliceosome in Three Conformational States." *Cell Research* 28 (3): 307–22.
https://doi.org/10.1038/cr.2018.14.
Zubradt, Meghan, Paromita Gupta, Sitara Persad, Alan M Lambowitz, Jonathan S Weissman, and Silvi
Rouskin. 2016. "DMS-MaPseq for Genome-Wide or Targeted RNA Structure Probing in Vivo."
Nature Methods 14 (1): 75–82. https://doi.org/10.1038/nmeth.4057.

1456
1457
1458

Supplementary Figures

Figure 1-figure supplement 1: Distribution of Exon 10 PSIs calculated for RNA-seq

data from GTEx database.

A) Distribution of TPM values of *MAPT* gene expression for tissues in the GTEx

database sorted by median TPM. Dotted box indicates tissues with median TPM

greater than 10. MAPT is highly expressed in the brain, and there is little

expression in other tissues. Figure was downloaded from the GTEx website.

B) Distribution of Exon 10 PSI for 12 central nervous system, muscle-skeletal,

colon-sigmoid, and breast-mammary tissue types. Percent Spliced In (PSI) of

Exon 10 was calculated from RNA-seq data for 2,962 tissue samples among 15

tissue types collected from 818 individuals in GTEx v8 database. The violin plot

for each tissue type and the corresponding region on the brain diagram is colored

by the median PSI for all samples of that type. The median PSI of 0.56 for all

tissue samples is indicated by the red dotted line.

C) Distribution of Exon 10 PSI for tissues per individual. Only individuals with *MAPT*

expression data in 8 or more tissues were plotted. Median PSI for each individual

is labelled by red dot on box plot.

Figure 1-figure supplement 2: DMS structure probing data for mature *MAPT* 3R and

4R isoforms

A) DMS reactivity data from T47D cells for two biological replicates for Exon 9-Exon

11 junction (3R isoform). Structure probing data for junctions of interest were

obtained using primers (Supplementary file 4) following RT of extracted RNA.

DMS reactivity is plotted for each nucleotide across spliced junctions for both

replicates overlaid in plot on the left. For scatter plot on the right, DMS reactivity

for Rep 1 vs Rep 2 is plotted per nucleotide with Pearson's correlation coefficient

displayed.

B) DMS reactivity data from T47D cells for two biological replicates for Exon 9-Exon

10-Exon 11 junction (4R isoform).

C) Comparison of DMS reactivity data for 3R vs 4R isoforms. Replicates 1 and 2

were pooled for each isoform. Top plot shows DMS reactivity plotted for each

nucleotide with isoforms overlaid. No data were collected for Exon 10 for the 3R

isoform because Exon 10 is spliced out. Bottom left scatter plot shows DMS

reactivities for Exon 9 in the 3R vs 4R context, whereas bottom right scatter plot

shows DMS reactivities for Exon 11 in the 3R vs 4R context. Pearson's

correlation coefficient is shown for each comparison.

D) DMS reactivity data from T47D and SH-SY5Y cells for Exon 9-Exon 11 junction

(3R isoform). Replicates from T47D cells were pooled. DMS reactivity is plotted

for each nucleotide across spliced junctions for both replicates overlaid in plot on

the left. For scatter plot on the right, DMS reactivity for T47D vs SH-SY5Y is

plotted per nucleotide with Pearson's correlation coefficient displayed.

E) DMS reactivity data from T47D and SH-SY5Y cells for Exon 9-Exon 10-Exon 11

junction (4R isoform).

1503

1504

Figure 1-figure supplement 3: DMS structure probing data for human small subunit

ribosome RNA (SSU) collected from T47D cells.

A) T47D cells were treated with DMS. Data for SSU were extracted by aligning reverse

transcribed RNA-seq data against the SSU sequence, after which reactivities were

calculated. DMS reactivities are plotted for each of the four sub-domains of the SSU.

Each value is shown with its standard error and colored by reactivity based on color

scale. Reactivities were cut off at 2. High DMS reactivities correspond to

unstructured regions, whereas low DMS reactivities correspond to structured

regions. The secondary structure of the SSU was downloaded from Loren William's

1514 lab Ribosome Gallery website

1515 (http://apollo.chemistry.gatech.edu/RibosomeGallery/eukarya/H%20sapiens/SSU/ind

1516 ex.html).

1517 B) Violin plots showing distribution of DMS reactivities for adenines and cytosines

1518 partitioned by paired versus unpaired nucleotides. Pairing status of nucleotides was

1519 determined from the known secondary structure of the SSU. Median DMS reactivity

1520 is indicated by thick horizontal black line on violin plot.

1521 C) ROC curves for predicting whether a nucleotide in the SSU is paired. Three different

1522 parameters were used for each of the three curves: DMS reactivities, base pairing

1523 probabilities predicted from SSU sequence, and base pairing probabilities for SSU

1524 sequence that were guided by DMS reactivities. The area under the curve (AUC) for

1525 each curve was calculated with AUCs closer to 1 corresponding to higher accuracy

1526 of predictions. Dotted line indicates AUC of 0.5 which corresponds to a model

1527 making random predictions.

1528 D) Comparison of distribution of DMS reactivities between SSU, *MAPT* 3R and 4R

1529 isoforms. Larger plot shows a density histogram of the DMS reactivities for each

1530 RNA. Inset boxplots display distribution of DMS reactivities. Level of significance:

1531 ***p-value $< 10^{-6}$

Figure 2-figure supplement 1: DMS structure probing data for precursor *MAPT* Exon

10-Intron 10 junction

A) DMS reactivity data from T47D cells for two biological replicates for Exon 10-

Intron 10 junction. Structure probing data for junctions of interest were obtained

using primers following RT of extracted RNA. DMS reactivity is plotted for each

nucleotide across spliced junctions for both replicates overlaid in plot on the left.

For scatter plot on the right, DMS reactivity for Rep 1 vs Rep 2 is plotted per

nucleotide with Pearson's correlation coefficient displayed.

B) DMS reactivity data comparing Exon 10 in precursor vs mature transcript.

Replicates 1 and 2 were pooled for each transcript. Right plot shows DMS

reactivity plotted for each nucleotide with mature and precursor RNAs overlaid.

DMS data for all of Exon 10 could not be collected for the precursor RNA due to

the position of primers chosen for amplification. Scatter plot on the left shows

DMS reactivities for Exon 10 in the precursor vs mature mRNA context with

Pearson's correlation coefficient shown for the comparison.

C) DMS reactivity data from T47D and SH-SY5Y cells for Exon 10-Intron 10

junction. Replicates from T47D cells were pooled. DMS reactivities are plotted for

each nucleotide across exon-intron junctions for both cell types overlaid in plot on

the left. For scatter plot on the right, DMS reactivity for T47D vs SH-SY5Y is

plotted per nucleotide with Pearson's correlation coefficient displayed.

D) Boxplots of distribution of DMS reactivities between SSU, *MAPT* 3R isoform, 4R

isoform and pre-cursor mRNA.

1555

Figure 2-figure supplement 2: The 3R and 4R mutations shift the WT structural

ensemble towards less and more accessible exon-intron junctions, respectively.

A) Density histogram showing the distribution of unfolding free energies of the splice

site (defined as last 3 nucleotides of exon, first 6 nucleotides of intron) for all

structures in the ensemble for WT, 3R and 4R mutated sequence. Distributions

for each sequence are colored according to the legend.

B) t-SNE Visualization of structural ensemble of wild type (WT) and, 3R (+19C>G)

and 4R (+15A>C) mutations. Structures were predicted using Boltzmann

suboptimal sampling and guided by DMS reactivity data (in Figure 2A). Data

were visualized using t-Distributed Stochastic Neighbor Embedding (t-SNE).

Shown are 3000 structures corresponding to 1000 structures per category. Each

dot represents a single structure and was colored by calculated unfolding free

energy of splice site at exon-intron junction (3 exonic bases, 6 intronic bases).

Data were clustered by k-means clustering and representative structures for five

clusters are shown. Bar plots next to the representative structure show the

proportion of the cluster in WT, 3R and 4R. The exon-intron junction is marked by

EIJ on each structure. Position of 3R and 4R mutations are marked by a red

asterisk on their respective representative structures. There are two additional

representative structures shared by WT and 4R sequence which have similar

structural contexts around the EIJ as the representative WT structure in Figure

2B.

1578

1579

Figure 3-figure supplement 1: Structure is a poor predictor of Exon 10 PSI for exonic

non-synonymous mutations

    A) Positions of intronic and synonymous experimentally validated mutations, used in

        training the structure model, are shown on the wild type representative structure

from Figure 2B. Each nucleotide is colored by its corresponding reactivity value

based on the color scale.

B) Violin plots showing the distribution of unfolding free energy of the exon-intron

pre-mRNA in the spliceosome B^{act} complex for the 1000 structures in the

ensembles of the 22 intronic and synonymous mutations. Each violin plot is

colored by whether the mutation promotes the 3R or 4R isoform ratio or the ratio

remains 50:50.

C) Exon 10 PSIs of 22 mutations predicted using unfolding free energy of the exon-

intron pre-mRNA in B^{act} complex of the spliceosome for the single minimum free

energy (MFE) structure and plotted against experimental PSIs measured in

splicing assays. Exon 10 PSIs predicted using Eq. 2. Each point on the

scatterplot represents a mutation and is colored by mutation category. Dotted

diagonal line is the x=y line, and the closer the points are to the diagonal, the

more accurate the prediction. Pearson correlation coefficient (R^2) of experimental

to predicted PSIs was calculated.

D) Exon 10 PSIs of non-synonymous and compensatory mutations predicted using

the unfolding free energy of pre-mRNA within the spliceosome B^{act} stage plotted

against corresponding experimental PSIs measured in splicing assays. Exon 10

PSIs were predicted using Eq. 1.

E) Violin plots show R^2s calculated for each mutation category by training and

testing on subsets of all mutations by non-parametric bootstrapping; Non-

synonymous (n=10), Compensatory (n=14).

 Figure 4-figure supplement 1: RBP binding motif strength is a poor predictor of Exon

 10 PSI for all mutations

 A) Exon 10 PSIs of 47 mutations predicted from structural change and plotted

 against experimental PSIs measured in splicing assays. Exon 10 PSIs predicted

 using Eq. 1. Each point on the scatterplot represents a mutation and is colored

 by mutation category. Grey line represents the best fit with dotted lines indicating

 the 95% confidence interval. Pearson correlation coefficient (R^2) of experimental

 to predicted PSIs. Violin plot shows R^2s calculated for each category by training

 and testing on subsets of all mutations by non-parametric bootstrapping; Exonic

 non-synonymous (n=11), Exonic synonymous (n=7), Intronic (n=15),

 Compensatory (n=14), Wildtype (n=1).

 B) Scatter plot of change in enhancer or silencer strength versus Exon 10 PSI. Each

 point represents a mutation. Blue line represents the line of best fit with dotted

1620   lines indicating the 95% confidence interval. Pearson correlation coefficient (R^2)

1621   is shown. The negative correlation between silencer strength and Exon 10 PSI is

1622   statistically significant with a p-value of 0.004.

1623

1624

1625

Figure 4-figure supplement 2: RBP binding motif strength is a poor predictor of Exon

10 PSI for all mutations

A) Heatmap of relative RBP binding motif strengths compared to wild type for 44

mutations. A value of 0 indicates that the mutation does not change RBP binding

motif strength, a positive value indicates increase in RBP binding motif strength,

and a negative value indicates weaker strength. RBPs implicated in the

regulation of Exon 10 splicing were collected from Qian & Liu, 2014 and the RBP

binding motifs were from Dominguez et al., 2018 and Ray et al., 2013. RBPs on

the left, implicated in the splicing inclusion of MAPT Exon 10, are highlighted in

pink, and RBPs involved in the exclusion of Exon 10 are highlighted in green.

Mutations are marked based on whether they promote the 3R or 4R isoform ratio

or the ratio remains 50:50.

B) Scatter plot displaying change in RBP motif strength versus Exon 10 PSI,

categorized based on whether the RBP is implicated in exclusion or inclusion of

Exon 10. Neither correlation coefficient is statistically significant.

C) Exon 10 PSIs of 44 mutations and wild type predicted using change in RBP motif

strength and plotted against experimental PSIs measured in splicing assays.

Exon 10 PSIs predicted using Eq. 5. Each point represents a mutation and is

colored by category of mutation. Dotted diagonal line is the x=y line, and the

closer the points are to the diagonal, the more accurate the prediction. Pearson

correlation coefficient (R^2) of experimental to predicted PSIs was calculated.

1648

1649

Figure 5-figure supplement 1: Additive model of structure and SRE has poorer

predictive performance compared with an interactive model specifically for synonymous

and non-synonymous mutations

A) Exon 10 PSIs of 47 mutations and wild type predicted using addition between

structure and SRE strength and fit to experimental PSIs measured in splicing

assays. Exon 10 PSIs are predicted using Eq. 7. Each point on scatterplot

represents a mutation and is colored by category of mutation. Grey line

represents the best fit with dotted lines indicating the 95% confidence interval.

Pearson correlation coefficient (R^2) of experimental to predicted PSIs was

calculated.

B) Violin plots showing correlation coefficients for each mutation category for

structure and SRE additive model. R^2s calculated for each mutation category by

training and testing on subsets of all mutations by non-parametric bootstrapping

10 times.

C) Pie chart of number and proportion of experimentally validated mutations in each

1665   cluster for heatmap in Fig 5B. Color of segment of pie chart matches up to the

1666   color of dendrogram branch in Fig 5B.

Figure 6-figure supplement 1: Complete mutagenesis of 100-nucleotide window

spanning Exon 10-Intron 10 junction

A) Heatmap of mean predicted Exon 10 PSIs grouped by wild type and mutant

nucleotide. Mutations were grouped by wild type and mutant nucleotide, and

mean predicted PSIs were calculated by group and colored according to color

scale. Violin plots of the distribution of PSI per group are shown in tile

corresponding to group. On each tile, mean PSI is indicated by dot and labeled

within violin plot.

B) Violin plot of the distribution of normalized change in unfolding free energy of the

exon-intron pre-mRNA in the spliceosome B^{act} complex from WT for all mutations

around a 100-nucleotide window of exon-intron junction. Mean of -0.67 is

indicated by dot. Dotted line represents the 0 value where there is no difference

between WT and mutant unfolding free energy. Positive values imply region

becomes less structured and has increased inclusion of Exon 10 (4R isoform);

negative values are interpreted as more structured and decreased inclusion of

Exon 10 (3R isoform).

C) Violin plots showing the distribution of normalized change in splice site,

enhancer, and silencer strength compared with WT for all mutations spanning a

100-nucleotide window of exon-intron junction. Mean is indicated by large black

dots on violin plots. Dotted lines represent the 0 value where there is no

difference from WT strength for mutation. Positive values suggest increased

inclusion of Exon 10 (4R isoform), whereas negative values are interpreted as

decreased inclusion of Exon 10 (3R isoform).

D) Violin plot shows the distribution of unfolding free energy of the exon-intron pre-

mRNA in the spliceosome B^{act} complex for the 1000 structures in the ensembles

of wild type and the 6 VUSs experimentally tested. Each violin plot is colored by

whether the mutation promotes the 3R or 4R isoform ratio or the ratio remains

50:50. The dotted line indicates the median unfolding free energy of the WT

ensemble.

E) Bar plots display the change in enhancer and silencer strength of the 6 VUSs
compared with WT.
F) Quantification of Exon 10 PSI of three replicates for splicing assay gels for 6
VUSs. One tailed Wilcoxon Rank Sum test was used to calculate significance of
Exon 10 PSI of VUS of interest compared to WT.

Supplementary Files

Supplementary file 1: ANOVA table for between individuals and within individuals Exon 10 PSI comparison

Supplementary file 2: Details on 47 experimentally tested *MAPT* mutations used in training model

Supplementary file 3: Details on 55 variants of unknown significance (VUSs) in *MAPT* from dbSNP

Supplementary file 4: Primers used for amplification of exon-exon or exon-intron junctions

Supplementary file 5: Re-calculated Position Weight Matrices for ESEs, ESSs, ISEs, ISSs

Supplementary file 6: Details on beta regression model results and features used for each training and test set

Supplementary file 7: Gel of RT-PCR data for splicing assay for new WT VUSs