Inference of disease-associated microbial gene modules based on metagenomic and metatranscriptomic data

Zhaoqian Liu[1], Qi Wang[1], Anjun Ma[2], Dongjun Chung[2], Jing Zhao[2], Qin Ma[2,#], Bingqiang Liu[1,3,#]

[1] School of Mathematics, Shandong University, Jinan, Shandong, 250100, China.

[2] Department of Biomedical Informatics, The Ohio State University, Columbus, OH, 43210, USA.

[3] Shandong National Center for Applied Mathematics, Jinan, Shandong, 250100, China.

[#] To whom correspondence should be addressed: Dr. Qin Ma, qin.ma@osumc.edu; Dr. Bingqiang Liu, bingqiang@sdu.edu.cn

Abstract

The identification of disease-associated microbial characteristics is crucial for disease diagnosis and therapy. However, the heterogeneity, high dimensionality, and large amounts of microbial data present tremendous challenges for the discovery of key microbial features. In this paper, we present IDAM, a novel computational method for disease-associated gene module inference from metagenomic and metatranscriptomic data. This method integrates gene context conservation (uber-operon) and regulatory mechanisms (gene co-expression patterns) to explore gene modules associated with specific phenotypes using a mathematical graph model, without relying on prior meta-data. We applied IDAM to publicly available datasets from inflammatory bowel disease, melanoma, type 1 diabetes mellitus, and irritable bowel syndrome and demonstrated the superior performance of IDAM in disease-associated characteristics inference compared to popular tools. We also showed high reproducibility of the inferred

gene modules of IDAM using independent cohorts with inflammatory bowel disease. We

believe that IDAM can be a highly advantageous method for exploring

disease-associated microbial characteristics. The source code of IDAM is freely available

at https://github.com/OSU-BMBL/IDAM.

Introduction

Trillions of microbes colonize the human body and play critical roles in multiple

fundamental physiological processes, and changes in microbial composition and

functions are intimately interwoven with multiple diseases, ranging from obesity to cancer

and autism[1,2]. With the rapid development of sequencing technologies, the Human

Microbiome Project[3], integrative Human Microbiome Project[4], and Metagenomics of the

Human Intestinal Tract project[5] have generated large amounts of microbial data related to

numerous diseases. Considerable effort has been aimed at using these data to produce

clinical insights and deepen the understanding of the mechanisms responsible for

disease progression and treatment[6]. Some pathogenic organisms have been found to be

closely related to disease states and can be used as potential markers of disease

occurrence and development[7,8]. These findings hint at a promising avenue for clinical

application—disease diagnosis and therapy—via the use of key microbial taxonomic or

functional characteristics.

One of the classical paradigms of key microbial characteristics identification is to detect

taxa and metabolic pathways that can statistically significantly distinguish between two or

more groups[9-11]. Additionally, given the success of machine learning in multiple scenarios,

there have been several attempts to use machine learning in the identification of

microbial characteristics[12,13]. Such methods include random forest and deep feedforward

networks[12,13]. However, these existing studies face several challenges. *First*, most of

them focus on 16S ribosome RNA and metagenomic (MG) data. 16S ribosome RNA data

suffers from low taxonomic resolution and an absence of functional information[14], which

may induce spurious characteristic identification with diseases since diverse microbial

communities from different patients can have very similar functional capabilities[14]. While

MG data can provide information at all taxonomic levels, researchers have found that a

large proportion of reads cannot be successfully mapped to existing reference genomes

during taxonomic identification[15,16]. Furthermore, only a minority of genes are

well-annotated with biochemical functions[15]. The limitations of reference genome

databases and annotation systems will lead to the potential loss of valuable information

while taxonomic or metabolic pathways analysis using MG data[15,16]. Meanwhile, the

functional analyses produced by MG analyses are not necessarily equivalent to the true

activities within the microbiome. Multiple metagenomically abundant genes in the human

gut microbiome are significantly down-regulated at the transcriptional level[17]. *Second*,

existing methods are designed based on the strong assumption that data with sufficiently

large sample sizes and accurate metadata information are available to design groups or

train models. However, the current metadata related to many sequencing samples is

incomplete, misleading, or not publicly available[18], which may lead to these methods

being infeasible or causing bias in key characteristic inference. These challenges drive

the need to develop an easy-to-use and effective method that keeps up with current data

for disease-associated microbial characteristic inference.

In this study, we proposed a novel method to **I**nfer **D**isease-**A**ssociated **M**icrobial gene

modules (**IDAM**), which to a large extent overcomes these limitations. The data we used

was matched MG and metatranscriptomic (MT) data sequenced from microbial samples

for assessing active functions within communities[19]. To make full use of the available data,

we focused on a community-level analysis that goes beyond species level[15], so that the

reads do not need to be mapped to reference genomes, and provided a comprehensive

overview of functions performed by either individual or different microbial assemblages.

We investigated disease-associated key characteristics by detecting gene modules with

specific co-expression patterns across certain subsets of samples[19], without requiring

prior metadata of samples. This can be a complement to existing supervised methods in

the absence of high-quality metadata. Here, we noted that one cannot fully characterize

complex functional activities based solely on ensemble co-expression, by which gene

expression is assessed using reads from various species (**Fig. 1a**) and genes with no

biological relevance may be clustered together (**Fig. 1b**). Fortunately, previous studies

have suggested that functionally associated genes are generally highly conserved during

evolution[20,21], forming uber-operon structures (**Supplementary Fig. S1**). This finding

motivated us to integrate gene context conservation to study cross-species

functional-related mechanisms, thereby alleviating the bias in key microbial

characteristics inference. This idea has not been used by anyone else so far. We used

uber-operon structures as the evolutionary footprints of cross-species, functionally

related genes (**Fig. 1c**). We believe that genes are more likely to be functionally related if

they have similar expression patterns and share the same uber-operon during evolution.

We applied IDAM to published matched MG and MT datasets of inflammatory bowel

diseases (IBD) (813 samples; including ulcerative colitis (UC) and Crohn's disease

(CD))[22]. The results suggested that IDAM performed better in key disease-associated

characteristic inference than popular tools, including both supervised and unsupervised

approaches. We further demonstrated the generalizability, reproducibility, and good

performance of IDAM through independent cohorts with IBD and other publicly available

datasets associated with melanoma[23], type 1 diabetes mellitus (T1DM)[24], and irritable

bowel syndrome (IBS)[25]. Additionally, we found the obtained gene modules are likely to

correspond to accessory genomic elements such as operons[15]. Hence, we believe that

IDAM can be a highly advantageous method for disease-associated microbial

characteristic inference based on MG and MT data. This approach will be valuable for

investigating the relationships between microbiome and human diseases.

Fig. 1. Key ideas of IDAM. a. Sequencing data from four samples (p_1 and p_2 from

healthy cohorts, and p_3 and p_4 from disease cohorts). Since MG shotgun sequencing

does not isolate microorganisms, the reads from different species (represented by purple,

green, and blue) were jumbled together. We therefore could not distinguish them easily at

the species level without alignment to reference genomes, but we could observe the

community-level expression of each gene based on the merged reads from all species.

The microbial community transcript number (MT data) can be corrected using the

underlying genomic copy number (MG data) in practice. Here, we simply show the

number of transcriptomic reads as expressions. The colors of the circles (grey, blue, and

red) indicate the expression level of the genes within each sample. It is assumed that g_1

and g_2 are functionally related in healthy samples, while g_4 and g_5 are functionally related

in disease samples. **b**. Co-expression modules. There were three co-expression modules,

which are defined as genes with coordinated expression patterns under sample subsets.

They were not equivalent to the known functional relationships. **c**. Gene modules by

integrating uber-operon structures and expression similarity. Each ellipse, including

multiple genes, represents an uber-operon. We searched for modules consisting of

genes with both co-expressed patterns and shared uber-operons, which resulted in two

identified biclusters. Therefore, we inferred there were functional-related relationships

between g_1 and g_2, as well as g_4 and g_5, which were consistent with the ground truth.

Results

IDAM: a metagenomic and metatranscriptomic analysis framework for

disease-associated microbial gene module inference

We first present a formulation of the problem addressed by this article, and then provide

an algorithmic overview of IDAM.

Computational formulation for gene module inference

The inference of gene modules can be mathematically formulated as an optimization

problem. For matched MG and MT data, a gene-sample expression matrix $A = (a_{ij})_{m \times n}$

can be constructed, which includes m genes $(g_1, g_2, \cdots, g_m)$ and n samples

$(p_1, p_2, \cdots, p_n)$. a_{ij} $(1 \leq i \leq m,\ 1 \leq j \leq n)$ represents the expression of gene g_i within

sample p_j, which is calculated by normalizing the RNA abundance (MT data) against the

DNA abundance (MG data) of g_i within sample p_j [26]. Based on the matrix, the

co-expression patterns present in gene and sample subsets can be detected by

identifying local low-rank submatrices[27]. Given the constraints of gene context

conservation, we integrated the submatrices with a graph model to maximize the number

of genes connecting to a single uber-operon. Specifically, we define a bipartite graph

$G = (V_1, V_2, E)$, where each node $u \in V_1$ represents an uber-operon. Each node $v \in V_2$

represents a gene in A, and the edge $e \in E$ connecting a node $u \in V_1$ to a node $v \in V_2$

exists if the gene v belongs to uber-operon u (**Methods**). Using this graph and matrix

A, the gene module inference problem can be formulated as the identification of a set of

local low-rank submatrices within A that maximizes the number of connected

components between the corresponding gene subsets and V_1 on G. We maximize the

number of connected components between gene subsets and V_1, to minimize the

heterogeneity of the gene subsets within the uber-operons, thereby ensuring that the

genes within each identified submatrix are as functionally related as possible. Based on

the identified submatrices within A (each contains a subset of genes and samples and

we referred them to as biclusters hereinafter), the gene modules, i.e., the gene subsets,

can be regarded as the characteristics of the shared phenotypes of the sample sets.

Such an idea greatly alleviates the dependency on metadata in disease-associated

characteristic inference. However, this problem is theoretically intractable (NP-hard,

Supplementary Information). Hence, instead of solving this problem in a globally

optimized way, we developed a heuristic algorithm, IDAM, as an approximate optimal

solution.

Overview of IDAM

The main goal of IDAM is to identify disease-associated microbial gene modules (**Fig. 2a**).

First, the community-level expression within each sample is assessed using HUMAnN2

based on matched MG and MT data[28]. This step provides a gene-sample expression

matrix, each entry of which represents the expression of a gene within a sample. Second,

IDAM combines the expression similarity and gene distribution within uber-operons to

measure the likelihood that each pair of genes has the same function. This step provides

a list based on the combined assessment for gene module initialization (**Fig. 2b**). Third,

the gene pairs in the list are used as seeds, and each seed is expanded by iteratively

adding other functionally related genes, to generate a bicluster until the new bicluster

starts to become smaller than previously identified biclusters (**Fig. 2c**). Finally, the

biclusters consisting of gene modules and corresponding sample sets are considered as

the final output of the algorithm. We regard the gene modules as functional

characteristics of the most common phenotype or state of the corresponding sample sets.

Details are available in the Methods section.

Fig. 2. Workflow of IDAM. a. The flow chart. We consider the matched MG and MT data

of multiple samples as the input, from which we obtain an expression matrix. By module

initialization and expansion, the biclusters will be generated as the output. **b.** Module

initialization. For each gene pair, we assess both expression similarity and context

conservation to produce a combined score. Let us use g_3g_4 as an example. Based on

known uber-operon structures, the two genes belong to u_2. We assign a reward U for g_3g_4

as a context conservation measurement. The expression similarity is assessed according

to the number of samples in which the rows are identical, and we obtain a score, S

(Methods). Finally, U and S are combined via a function C (w is a tuning parameter),

generating a score, C_{34}. All gene pairs are sorted in a decreasing order based on the

combined scores. **c.** Module expansion. We use gene pairs in the list and the samples

under which the two genes are co-expression to obtain the initial biclusters and expand

them gradually. We iteratively add a new gene with the highest combined score into a

bicluster until the updated bicluster does not get bigger. Let us use g_3g_4 as an example.

First, we add three genes (g_1, g_2, and g_5) into g_3g_4, forming three new biclusters (denoted

as $g_3g_4g_1$, $g_3g_4g_2$, and $g_3g_4g_5$). Then, the combined score of each bicluster is assessed

(**Methods**). Since bicluster $g_3g_4g_5$ has the highest score, we replace the previous

bicluster with this one. Next, we iteratively add a new gene (g_1 and g_2) to the current

bicluster and implement the same processes. After comparing the combined scores, we

find $g_3g_4g_5g_2$ to have the highest score, but the bicluster is of size 4×2, which is smaller

than $g_3g_4g_5$ (3×3). Therefore, we stop here and the bicluster $g_3g_4g_5$ will be provided as the

final output.

The gene modules inferred by IDAM distinguished different phenotypes and

enabled the discovery of disease subtypes

To assess the usefulness of IDAM, we assessed the phenotype classification

performance of inferred gene modules. We applied IDAM to a publicly available dataset

of patients with IBD and controls (referred to as non-IBD, **Methods**). The following

analyses are based on the IBD-associated dataset unless otherwise specified. By

assessing the community-level expression of each sample in the IBD-associated dataset,

we obtained a matrix consisting of the expression of 917,127 genes across 734 samples.

Based on this matrix, IDAM found 39 biclusters containing 47 to 5,452 genes

(**Supplementary Table 1**), totally covering 29,127 genes. Ideally, samples with the same

gene module should have highly consistent phenotypes. We used Z-tests to evaluate the

phenotypic consistency of samples in each bicluster (**Methods**). Results suggested that

33 gene module associated sample sets were consistently enriched for a particular

phenotype, with 20 enriched for IBD and 13 for non-IBD (**Fig. 3a and Supplementary**

Table 2).

We recognized that microbial communities can be influenced by multiple factors, such as

age, race, and diet. These confounding factors can lead to high inter-individual

heterogeneity, make disease-associated microbial characteristics less clear, and

increase the risk of false positives or negatives in gene module inference. Hence, we

further assessed whether the gene modules identified using IDAM had reliable

associations with phenotypes, by applying a propensity score-based matching for each

gene module to confirm its association with phenotypes. We constructed paired samples

by matching the propensity scores estimated from other covariates, such as age, gender,

and comorbidities. The Wilcoxon signed-rank test based on the paired samples indicated

that 79.49% (31 of 39) gene modules were statistically significantly related to a specific

phenotype, with q-values < 0.20 (**Fig. 3b and Supplementary Table 3**), demonstrating

the high reliability of the gene modules identified using IDAM.

We specifically observed the enriched phenotypes of samples within each bicluster, and

found 12 sample sets were with consistent phenotype UC and eight with CD, indicating

the gene modules identified could recognize the subtypes of IBD (**Fig. 3c**). We here gave

an example of three biclusters to show the phenotype specificity, namely Bicluster 4,

Bicluster 15, and Bicluster 28 (**Fig. 3d**). Because of space limitations, we show the

expression of the top 3% of genes in each bicluster: 35/1,162, 17/552, and 29/1264 for

Biclusters 4, 15, and 28, respectively. IDAM detected the checker-board substructures

from the original expression data. Gene module 4, corresponding to Bicluster 4, is

specific to non-IBD, while Gene modules 15 and 28, corresponding to Biclusters 15 and

28, are specific to UC and CD, respectively, suggesting the capability of IDAM to uncover

gene modules of disease subtypes. These gene modules were associated with 117

pathways (with q-values < 0.05 based on the hypergeometric test), of which 23 were also

identified as differentially abundant across samples with different phenotypes based on

pathway abundance from HUMAnN2 (**Methods and Supplementary Table 4**). For

example, there was a significant enrichment in short-chain fatty acid metabolism (e.g.,

butyrate metabolism) in non-IBD-associated gene modules. The L-methionine

biosynthesis, superpathway of L-citrulline metabolism, and superpathway of L-lysine,

L-threonine, and L-methionine biosynthesis were enriched in non-IBD-associated gene

modules, while preQ$_0$ biosynthesis was enriched in IBD. The superpathway of

lipopolysaccharide biosynthesis was enriched in CD-associated gene modules, and

methylphosphonate degradation was enriched in UC-associated gene modules. These

findings are in keeping with previous observations[29-32], indicating that the gene modules

from IDAM can provide functional insights into different phenotypes.

We also compared IDAM with several decomposition-based and biclustering-based

approaches, including ICA[33], QUBIC[34], ISA[35], and FABIA[36]. These methods have not

been applied to microbial data for gene module inference, but they may reveal the

functional characteristics of different phenotypes without the need for prior metadata. The

results suggested that there were 37, 30, and 13 gene modules identified by QUBIC, ICA,

and FABIA, respectively, while no gene module was detected by ISA. We further

evaluated the phenotype consistency of gene module-associated sample sets using

Z-tests. While 84.62% (33/39) sample sets from IDAM had a consistent phenotype, this

number was only 64.86% (24/37), 26.67% (8/30), and 30.77% (4/13) for QUBIC, ICA, and

FABIA, respectively (**Fig. 3e**). This finding indicated that the gene modules inferred by

IDAM can cluster samples with the same phenotypes more effectively than other tools.

This finding was not surprising, since IDAM took both gene expression and gene context

conservation into consideration, thereby identifying more biologically meaningful gene

modules, while others have used only gene expression. We also compared the running

time of these methods (**Fig. 3f**). IDAM took seven hours, QUBIC and FABIA took 12 and

20 hours, respectively, while ISA was terminated after 55 hours without results. ICA took

seven minutes; however, the application power of ICA was limited by the relatively low

accuracy; only less than a third of the gene modules could distinguish between

phenotypes. Collectively, the gene modules identified by IDAM performed best in

distinguishing the phenotypes and can provide leads for new disease subtypes or states

for future clinical testing.

Fig. 3. Performance of gene modules inferred by IDAM in distinguishing phenotypes. a. The values of ln(1+*q*-value) of non-IBD, UC, and CD, shown as pink, green, and purple, respectively. Each column represents a gene module. For modules 1-13 (except module 9), the ln(1+q-value) of UC and CD overlapped. For modules 14-22, the ln(1+q-value) of CD and non-IBD overlapped. We only showed the values

corresponding to non-IBD in these cases. The yellow horizontal line indicates the critical

value of *q*-value 0.20. The color of the circles below this line means that the

corresponding phenotype exhibited strong enrichment within the gene

module-associated sample sets. **b.** *q*-values of each gene module with respect to

different phenotypes. Each column represents a gene module, and the color indicates the

q-value. * indicates a *q*-value of < 0.20. **c.** The number of gene modules identified for

each phenotype. **d.** An example of the gene modules associated with phenotypes. The

color of each cell in the heatmap indicates the log2 value of the expression. Grey

represents that the gene is not expressed within the corresponding sample. **e.** Percent of

gene modules that were associated with samples enriched in certain phenotypes,

identified by IDAM and competing algorithms. NA means no results from ISA. **f.** Running

time of IDAM and competing algorithms.

Species-level analysis suggested that the gene modules inferred by IDAM were

biologically associated with phenotypes

Since differences in community-level functions ultimately imply species-level functional

differences, we traced genes within each gene module back to individual species. These

genes were from 133 species, which were dominated by Bacteroidetes and Firmicutes,

followed by Proteobacteria at the phylum level (**Fig. 4a**). These were consistent with

previous research into IBD-associated taxa[32]. To provide the genomic and functional

context of the identified gene modules, we aligned the genes comprising gene modules to

the reference genomes of these species in RefSeq[37] and observed the distribution of

genes on the reference genomes. We found the genes located on a genome tend to form

several operons, instead of being spread out over the whole genome randomly. For

example, the genes aligned to the representative genome of *Escherichia coli* make up

several operons from RegulonDB[38], such as operon *glnALG*, *ptsA-fsaB-gldA*, and

dcuB-fumB (**Supplementary Fig. S2**). The operon *glnALG* encodes propanediol

dehydratase, which has been demonstrated of higher levels in individuals with CD

compared to that in healthy individuals[39]. This suggested the gene modules from IDAM

were able to provide possible mechanisms for diseases that are suitable for further

investigation.

We further classified the species according to gene module-associated sample

phenotypes from Z-tests and obtained 43, 63, and 34 species associated with non-IBD,

UC, and CD, respectively (**Methods and Supplementary Table 5**). Seven of these

species overlapped between UC and CD. We collected IBD-related species (34 for

non-IBD, 32 for UC, and 90 for CD) from Peryton, a database of microbe-disease

associations, and previous taxonomic studies into IBD[40-43] (**Supplementary Table 6**).

Among them, a total of 31 species were captured by IDAM (**Fig. 4a**), including 17 for

non-IBD, 12 for UC, and 9 for CD (shown in red in Supplementary Table 6). The seven

overlapped species from IDAM have been reported to be associated with both UC and

CD[40-42], suggesting that the IDAM results are reliable. We compared IDAM with QUBIC[34],

ICA[33], FABIA[36], and LEfSe[11]. ICA, QUBIC, and FABIA are not dependent on metadata,

while LEfSe identifies differentially abundant species based on prior phenotype

information. Detailed results are shown in **Fig. 4b**. The average consistency with

collected species across the three phenotypes was higher for IDAM (28.35%), than for

QUBIC (15.73%), LEfSe (18.05%), ICA (14.93%), or FABIA (3.08%) (**Fig. 4c**). The

species-level analysis showed that the gene modules inferred by IDAM were biologically

associated with phenotypes.

Fig. 4. Species-level analysis of identified gene modules. a. Phylogenetic tree of the
species associated with gene modules identified by IDAM. Colored dots indicate species
that have been validated by previous studies. **b.** Results of species-level analysis by
different methods. The numbers in the header (34, 32, and 90) represent the number of
collected species associated with the three phenotypes, respectively. Each cell in the
table is represented in the form of X/Y (Z). 1) The denominator Y means the number of
species associated with a particular phenotype identified from the different methods; 2) X
means the number of identified species that are consistent with the collected species;
and 3) Z indicates the consistency index of each method associated with the phenotypes,
which reflects the reliability of identified species of different methods. **c.** Average
consistency for each method. A higher value indicates that inferred species are more
reliable.

**Integration of gene context conservation substantially improved the performance
of gene module inference**

Our hypothesis is that genes with similar expression patterns and high context
conservation during evolution are likely to be functionally related. To evaluate the validity

of this hypothesis, we sought to identify gene modules without using gene context

conservation in IDAM (referred to as IDAM without gene context conservation). As a

result, we found the gene modules identified by this version were less capable of

distinguishing different phenotypes than those from IDAM. IDAM without gene context

conservation detected 37 biclusters from the expression matrix, of which only 64.86%

(24/37) sample subsets were of consistent phenotypes (the percentage of IDAM is

84.62%). Pathway analysis suggested that 78.38% (29/37) gene modules from IDAM

without gene context conservation were significantly associated with MetaCyc functional

categories (with q-values < 0.05 using the hypergeometric test), which is lower than the

94.87% (37/39) of IDAM (**Fig. 5a**). These findings indicated that the integration of

uber-operon structures contributed to the identification of informative gene modules.

We explored the underlying reasons for the superior performance of integrating gene

context conservation. We observed the distribution of the genes in the uber-operons for

each gene module from both IDAM and IDAM without gene context conservation. Overall,

the genes within gene modules that are not significantly associated with any pathways

tended to be spread out among more uber-operons. This finding supports our hypothesis

that functionally related genes are of high context conservation. As an example, we

investigated Bicluster 20 from IDAM without gene context conservation (abbreviated as

'Bicluster 20' hereinafter, **Fig. 5b**). Since the 200 genes within the bicluster had

coordinated expression across the 73 samples, they would be clustered together if we

merely considered co-expression. However, these genes were not significantly

functionally related, and were from eight uber-operons. They cannot reflect the unique

functional characteristics of a particular phenotype. Indeed, the 73 samples within

Bicluster 20 were evenly scattered over different phenotypes: 23 in non-IBD, 11 in UC,

and 39 in CD. When we took gene context conservation into consideration, a new

bicluster occurred (Bicluster 29 from IDAM, abbreviated as 'Bicluster 29' hereinafter).

This bicluster consists of 328 genes and 33 samples, and there was considerable overlap

with Bicluster 20 (**Fig. 5b**). The bicluster was dominated by an uber-operon that is closely

associated with purine metabolism[21], and the enriched pathways are shown in **Fig. 5c**.

Previous studies have shown that the imbalance between the biosynthesis and

degradation of purines can produce excessive uric acid in the gut, leading to the

occurrence of IBD[44]. The facts showed the 33 samples indeed were significantly enriched

in phenotype CD, with a *q*-value < 0.20 using a Z-test. This finding demonstrated the

value of gene context conservation in the clustering of genes with biological relevance,

which improves the ability of IDAM for gene module inference.

Fig. 5. Analysis of the contribution of gene context conservation. a. Comparison

between IDAM and IDAM without gene context conservation. **b.** A specific case of

identified biclusters. Each row represents a gene, and each column represents a sample

(a total of 474 genes and 73 samples). The colored bar at the top represents the

phenotypes of the corresponding samples. **c.** Pathways enriched in Bicluster 20 and

Bicluster 29. The pathways in grey indicate the pathways enriched in Bicluster 20, while

red indicates those enriched in Bicluster 29. The dotted vertical lines indicate the critical

value (a *q*-value of 0.05). The bars located outside the dotted lines correspond to

significantly enriched pathways.

IDAM showed high reproducibility and good generalizability in gene module

inference and provided insights into melanoma-related progression-free survival

after immunotherapy

We used an independent dataset associated with IBD to validate the reproducibility of

inferred gene modules (**Methods**). For each gene module associated with IBD inferred

from IDAM, we estimated the mean difference in centered log-ratio expression between

samples with and without certain phenotypes using the independent dataset. The random

effects, such as age and diet, were balanced by matching the propensity score of the

samples. Results suggested the expression of 79.49% (31/39) gene modules was

significantly different across phenotypes, with a *q*-value < 0.20 (**Supplementary Fig.**

S3a). That means these gene modules were also significantly associated with sample

phenotypes on the independent dataset. When performing the same analysis on single

gene expression (a single gene randomly selected from each gene module), we

observed a much lower rate (33.33%) with significant differences across phenotypes

(with *q*-values < 0.20, **Supplementary Fig. S3b**). This supported the utility of gene

modules for exploring reproducible associations between microbiome and diseases.

To demonstrate the generalizability of IDAM, we applied it to an integrated dataset

containing 60 samples from four different cohorts: cohorts with melanoma, T1DM, without

T1DM (non-T1DM), and IBS (**Supplementary Information and Supplementary Table**

7)[23-25]. The data processing included quality control and the measurement of the

community-level expression of each sample (**Supplementary Information**). A matrix

with the expression of 470,305 genes across 60 samples was produced. Based on this

matrix, IDAM inferred 55 gene modules covering 13,145 genes. Analogous to the

analyses of the IBD-associated dataset, we first evaluated the phenotype consistency of

each gene module-associated sample set using Z-tests and found that 32 of 55 (58.15%)

were consistently enriched for a particular phenotype: seven enriched for T1DM, 18 for

non-T1DM, and seven for melanoma. Comparison with other methods, QUBIC, ICA,

FABIA, and ISA, showed that the gene modules inferred by IDAM could distinguish

between the phenotypes most effectively (**Fig. 6a and Supplementary Table 8**). We

then found that 20.43% of the species from IDAM were consistent with the collected

disease-specific species, and the average consistency across the four phenotypes was

the highest of all the methods (**Fig. 6b and Supplementary Table 9**). Pathway

enrichment for the genes within each gene module showed that IDAM could detect eight

differentially abundant pathways identified by HUMAnN2 (**Supplementary Table 10**) and

many pathways consistent with previous findings[45-47]. For example, the biosynthesis of

low-carbon-number saturated lipid classes, including myristic, stearic, and palmitic acid,

was likely to be higher in those with T1DM than in those without T1DM[46]. Aromatic amino

acids, phenylalanine, and tyrosine were higher in T1DM patients[45], which tends to

increase the risk of T1DM. Collectively, these results demonstrated the high reliability of

the gene modules identified by IDAM.

We also noted that progression-free survival (PFS) after immunotherapy of melanoma

individuals was publicly available. We explored the association between inferred

microbial gene modules of melanoma and PFS (**Methods**). Four out of seven gene

module-related sample groups, with or without the gene module, differed significantly in

their PFS, with a log-rank *p*-value < 0.05 (**Fig. 6c and Supplementary Fig. S3c**). We

assessed the progression risk of patients using Cox proportional hazards models, and

found that the patients with Gene modules 2, 4, and 5 had a lower risk of progression

than those without these gene modules (**Fig. 6d and Supplementary Table 11**), as

assessed during follow-up. Thus, we inferred that the three gene modules were

associated with longer PFS. In contrast, gene module 1 was associated with shorter PFS,

and individuals with this gene module had a high risk of melanoma progression (Hazard

ratio [95% CI] = 9.1195 [3.987,20.615]; **Fig. 6d**). We collected all species and pathways

that have been identified to be associated with shorter or longer PFS from the paper by

Peters *et al.*[23] (**Fig. 6e**). Most of these (6/11 species and 5/8 pathways) can be found in

the gene modules identified by IDAM (shown as red in Fig. 6e). These findings support

the contention that the gut microbiome may have an impact on the efficiency of

immunotherapy on melanoma[23], and provide possible mechanistic insights into means for

improving the efficiency of immunotherapy by manipulating the microbiome.

Fig. 6. Results of IDAM on the integrated dataset. a. Comparison of phenotype

consistency among IDAM, QUBIC, ICA, FABIA, and ISA. The red bars indicate the

percent of gene module-associated sample sets with consistent phenotypes from IDAM,

while the grey bars indicate those from other methods. **b.** Comparison of the average

consistency of collected species across the four phenotypes. **c.** Kaplan-Meier curves of

gene module-associated groups along with corresponding log-rank test *p*-values. **d.**

Results from Cox proportional hazard models. The yellow line indicates a hazard ratio

equal to 1. CI: confidence interval. the *p*-value is from the likelihood-ratio test. **e.** Species

and pathways related to PFS. The red highlights represent those that can be found by

gene modules from IDAM.

Discussion

Although many studies have focused on the microbial characteristic inference of

diseases[9-13], it is still a fundamental challenge to explore critical characteristics in

large-scale, highly heterogeneous microbial data. We developed a new methodology,

IDAM, for matched MG and MT data to identify disease-associated gene modules. The

above results indicate that IDAM is a highly effective approach to the identification of

disease-associated microbial gene modules based on both gene co-expression and

context conservation during evolution. The gene modules of diseases identified can be

useful in understanding disease pathogenesis and providing guidance for the future

development of disease treatment.

IDAM is innovative in that, unlike previous disease-associated microbial characteristics

analysis methods[9-13], it integrates gene context conservation and gene expression

information by leveraging a multi-omic view of microbial communities for the inference of

diseases using functional characteristics. IDAM does not rely on prior knowledge about

the samples. Given that existing metadata is often misannotated and misleading, even

publicly unavailable[18], creating an overly complex environment for sample reanalysis.

This will reduce the application power of previous metadata-based methods. IDAM can

be a good complement to those methods in this case and eliminates bias from misleading

data noise. Additionally, IDAM, as an unsupervised method, is effective to develop new

biological hypotheses, providing novel insights into new disease subtypes or stages.

Previous supervised learning methods, by contrast, cannot provide such information

because they identify key features based on known labels and annotations of samples.

We illustrated and evaluated IDAM using IBD, melanoma, T1DM, and IBS associated

datasets, and found that it outperformed existing methods. We discovered microbial gene

modules significantly associated with specific diseases, some of whose functions are

consistent with what has been reported previously. We also evaluated the gain of using

gene context conservation information in IDAM and found that ignoring this information

can lead to significantly decreased biological relevance between genes within a bicluster.

This research demonstrated that uber-operon structures play a crucial role in functional

indication. Furthermore, IDAM is suitable for the inference of microbial characteristics on

large-scale datasets. One can integrate microbial datasets spanning multiple diseases

and include multiple independent studies for each disease to explore disease-specific

microbial characteristics[48], providing possible insights into disease diagnosis.

IDAM is still not free of limitations. It identifies gene modules at the community level,

which need further taxonomic analysis to improve their clinical utility. This limitation can

be somewhat addressed by tracing genes back to taxa by mapping them to existing

reference genomes. However, there are some genes that cannot be attributed to known

species, which would lead to the biological information in the gene module being

inadequately interpreted. We believe that this issue will be alleviated as more microbial

species and reliable metagenome-assembled genomes are discovered and explored. We

plan to develop a three-dimensional analysis approach in the future[49], which takes both

microbial species and genes and pathways of multiple samples into consideration for

disease-associated microbial characteristic inference. In this way, modules involved in

diseases consisting of genes and pathways with specific functions of specific species will

be detected, providing both functional and taxonomic insights into diseases. Additionally,

IDAM requires relatively large amounts of data. Since IDAM uses expression similarity to

assess the relevance of gene regulation, a large sample size might be needed to

guarantee reliable measurement from data. In practice, new paired MG and MT data from

users can be integrated with the datasets used in this article to increase the sample pool,

thereby improving the practicability of IDAM. Finally, the gene modules identified by IDAM

only provided associations between microbiomes and diseases. Rigorous biological

experiments are required to establish the causality between the gene modules identified

and the disease of interest, and to establish their clinical prevalence and utility.

Methods

Datasets

We downloaded all matched quality-controlled MG and MT data associated with IBD, as

well as the corresponding HUMAnN2 results and metadata of samples at the IBDMDB

website in March 2020 (https://ibdmdb.org)[22]. There were 198 samples from non-IBD

subjects and 615 samples from patients with IBD, consisting of 381 UC samples and 234

CD samples. These samples were collected in two phases: an initial round as a pilot

obtaining 79 samples with matched MG and MT data and the second phase obtaining

734 samples with matched MG and MT[22]. We split the IBD-associated dataset into two

parts based on sample collection phases: one consisting of 79 samples used for

validation and the other consisting of 734 samples used for gene module inference. We

also downloaded matched MG and MT data in .fasta format from the NCBI Sequence

Read Archive with the accession number of SRP197281 for melanoma-associated data[23],

SRP072561 for T1DM- and non-T1DM-associated data[24], and ERP001739 for

IBS-associated data[25]. Uber-operon structures were obtained from the paper by Che *et*

al.[21]. The authors predicted the uber-operons through identifying groups of functionally or

transcriptionally related operons whose gene sets are conserved across multiple

reference genomes[21].

IDAM: a framework for identifying disease-associated microbial gene modules based on metagenomic and metatranscriptomic data

IDAM is a heuristic algorithm for disease-associated microbial gene module inference. It is based on a greedy search to gather genes with similar expression patterns and high context conservation as functional gene modules. IDAM consists of three steps.

Step 1: *Extracting community-level gene expression of samples*

The input data is quality-controlled (quality- and length-filtered and screened for residual host DNA) matched MG and MT sequencing datasets in fasta, fasta.gz, fastq, or fastq.gz format. The datasets of each sample are first processed using HUMAnN2 for gene relative expression within communities[28]. Here, we performed both nucleotide-level search and accelerated translated search using HUMAnN2 for a more comprehensive community-level profiling than only reference genomes based mapping[28]. Given that random sampling may lead to the detection of transcripts for undetected genes for low-abundance species, we used Laplace smooth for RNA and DNA abundance and use smoothing DNA-level features to avoid divide-by-zero errors during normalization. In this study, we directly used the HUMAnN2 results of the datasets from IBDMDB for the IBD-associated datasets. Then, we extract the community-level gene expression of each sample to construct a gene-sample expression matrix $A_{m \times n}$, in which each row represents a gene (total m genes indicating by UniRef90 identifiers[50] and forming a gene set X), and each column represents a sample (total n samples).

Step 2: *Measuring gene context conservation and expression similarity for each gene pair*

We assess context conservation and expression similarity for each gene pair and sort them as a list for module initialization.

2.1 Gene context conservation

Gene context conservation is assessed based on the gene distribution in uber-operons.

Since the genes within X referred to here are indicated as gene families from UniRef90

(a non-redundant protein sequence database)[50], we align the gene family sequences with

genes in uber-operons to determine to which uber-operons these gene families belong.

The sequences of genes in all uber-operons are collected into a custom database. For

each gene family, g_i, within X (1≤ i ≤ m), we extract the corresponding sequence

from the UniProt database[50] and align it with the custom database using BLAST[51]. We

define a gene-to-uber-operon mapping function $\varphi: D \rightarrow H$, in which D consists of all

subsets of X, Y is the set of all uber-operons, and H consists of all subsets of Y. We

assume that a gene family g_i (1≤ i ≤ m) belongs to an uber-operon u if the *E*-value of

its alignment with one of the uber-operon's sequences is less than 0.001. This is denoted

by $\varphi(G_i) = \{u\}$, where G_i is the set consisting of a single gene, g_i. If multiple sequences

from different uber-operons have an *E*-value less than 0.001, we choose the uber-operon

corresponding to the smallest *E*-value. In contrast, if no sequence has an *E*-value less

than 0.001, we set $\varphi(G_i) = \{\emptyset\}$. In this way, we construct a map between an uber-operon

and a gene.

We further define the relationship between uber-operons and gene modules with multiple

genes. Suppose a gene set Q consists of b genes $\{g_1, g_2, \cdots, g_b\}$ (1 < b ≤ m). We

assume $\varphi(Q) = \varphi(\{g_1\} \cup \{g_2\} \cup \cdots \cup \{g_b\}) = \varphi(G_1 \cup G_2 \cup \cdots \cup G_b) = \varphi(G_1) \cup \varphi(G_2) \cup$

$\cdots \cup \varphi(G_b)$, where G_i represents the set consisting of a single gene g_i ($i = 1, 2, \cdots, b$).

For two gene modules, I and J, we set a reward $U(I, J)$ as a gene context conservation

measurement, based on the relative positions of their genes in uber-operons:

$$U(I,J) = \begin{cases} 1, & \varphi(I) \subseteq \varphi(J) \text{ and } \varphi(I) \neq \{\emptyset\}, \\ -1, & \varphi(I) = \{\emptyset\} \text{ and } \varphi(J) \neq \{\emptyset\}, \\ -1, & \varphi(I) \neq \{\emptyset\} \text{ and } \varphi(J) = \{\emptyset\}, \\ -2, & \varphi(I) \neq \varphi(J) \neq \{\emptyset\}, \\ 0, & otherwise. \end{cases} \quad (1)$$

Based on this, we calculate $U(G_i, G_j)$ as the gene context conservation of each pair of

genes g_i and g_j ($1 \leq i \leq j \leq m$).

2.2 Expression similarity of genes

Expression similarity is assessed based on a vector composed of the expression values

of each gene within all samples. We need to measure the expression similarity not only

between two genes, but also across multiple genes, while expanding the modules in *Step*

3. This cannot be achieved using existing similarity measurements, such as Pearson

correlation coefficients, since they are generally used for paired data points. To address

this issue, the continuous expression value of each gene is first discretized to an integer

representation using qualitative representation[34] (**Supplementary Information**). Based

on this representation, we measure the expression similarity $S(G_i, G_j)$ of gene pair

(g_i, g_j) as the number of samples under which the corresponding integers along the

rows of the two genes are identical (or identical integers but with opposite signs).

2.3 Integration of gene context conservation and expression similarity

A combined score, $C(G_i, G_j)$, integrating gene context conservation and the expression

similarity of g_i and g_j is obtained via the following function:

$$C(G_i, G_j) = \left(1 + w \times U(G_i, G_j)\right) \times S(G_i, G_j), \quad (2)$$

where w is a tuning parameter that can be set by users. w ranges between 0 and 1,

and a larger value of w indicates that gene context conservation has a stronger impact

on estimating functional relationships. Here we used 0.3 as the default value

(**Supplementary Information and Supplementary Table 12**). Finally, the gene pairs are

sorted in descending order for the combined scores, generating a seed list, L.

Step 3: Generating biclusters consisting of gene modules and corresponding sample sets

Since the gene set X is a pan-genome of all microorganisms from multiple samples, it

contains a large number of genes. To improve the computational efficiency, we partition

the rows within the matrix equally into t subsets[52,53], where t is determined using a

stochastic model[52,53]. Then, we use gene pairs within each subset instead of all gene

pairs in X, forming the seed list L, which can greatly decrease the number of seeds and

reduce the computational burden.

A submatrix is regarded as biologically relevant if it does not occur randomly in a matrix

$A_{m \times n}$. Since the output submatrices from our algorithm are biologically relevant and

generated by a gene pair (see *Step* 3.1), we need to make sure that at least one gene

pair within each meaningful submatrix can exist as a seed, to keep all biologically relevant

submatrices detectable. This means we can at most divide the rows of matrix *A* into *k* -1

subsets (*i.e.*, $t = k - 1$), where k represents the number of genes of the biologically

relevant submatrix with the fewest genes (referred to with the smallest row size). We

statistically assess the smallest row size of the submatrix that is unlikely to occur by

chance. The submatrices we target are of rank one mathematically, since the

corresponding genes have co-expressed patterns. Suppose a submatrix with rank one is

of size $k \times s$, and its probability of random occurrence in $A_{m \times n}$ is calculated as follows

[52,53]:

644
$$P_{ks} = n \times (n-1) \times \cdots \times (n-s+1) \times \binom{m}{k} \times \left(\frac{2}{s!}\right)^k \times (1 - \frac{2}{s!})^{m-k} \qquad (3)$$

The submatrix is considered to be biologically relevant if $P_{ks} < 0.05$. When users have

prior knowledge of the largest column size, s, we can calculate k based on the formula

above. The smallest row size of the biologically relevant submatrices can be determined.

Users can then set the number of subsets ($t = k - 1$). In this work, the default value of t

was set to 10. That is, the matrix was divided into 10 subsets, which greatly reduced the

computational burden and kept biologically relevant submatrices detectable to a large

extent.

3.1 Initialization

We start with the first gene pair in L that satisfies one of two conditions: i) at least one

gene of these two has not been included in previous biclusters, or ii) these two genes are

included within different previous biclusters, and there is no overlap between the gene

components of the two biclusters. We aggregate samples in which the integers of the two

genes are identical (represented as set P), forming a current bicluster $M = \{Q, P\}$,

where Q represents the gene set consisting of the paired genes. The gene pair is then

removed from L.

3.2 Expansion

We expand M by adding a new gene (if any) that has the highest combined score with M,

giving rise to the updated bicluster $M' = \{Q', P'\}$. The calculation of the combined score

between a gene (represented as g_z ($1 \le z \le m$)) and Q, namely $C(G_z, Q)$, is similar to

that of two genes g_i and g_j. We first assess the context conservation between g_z and

the gene set Q of bicluster M, i.e., $U(G_z, Q)$. Then, we measure the similarity of

expression of g_z and Q, $S(G_z, Q)$, as the number of samples within P under which the

corresponding integers along the rows are identical (or identical integers but with

opposite signs). Based on these, a combined score $C(G_z, Q)$ can be obtained by

replacing $U(G_i, G_j)$ and $S(G_i, G_j)$ in the calculation of $C(G_i, G_j)$ with $U(G_z, Q)$ and

$S(G_z, Q)$, respectively. If $min\{|Q'|, |P'|\} > min\{|Q|, |P|\}$, we set $M = M'$ and repeat Step

3.2. Otherwise, go to *Step* 3.3.

 3.3 Filter and output

676 If the proportion of genes overlapping between Q and the component genes of

677 previously identified biclusters is less than a threshold, f, bicluster M will be provided as

678 the final output, and is ignored otherwise. In this study, the default f was set to 0.1 for

679 more separate biclusters. Go to *Step* 3.1 until the list L is empty.

681 Evaluating phenotype consistency of samples within one bicluster

Z-tests were used to assess whether samples with the same gene module are of highly

consistent phenotypes. Here, the samples in each bicluster were regarded as with the

corresponding gene module. We extracted the phenotypes of all samples from the

metadata as background, and the phenotypes of samples in each bicluster as test sets.

For each phenotype and each test set, we counted the phenotype occurrence in the test

set and background. We evaluated the significance of each phenotype using a right-tailed

Z-test with the null hypothesis that the observed frequency in the test set is less than or

equal to the occurrence probability within the background. The resulting Z-score is

tranfered to *p*-value, and further adjusted using the Benjamini-Hochberg procedure for

false discovery rate. If the *q*-value is less than 0.20, the null hypothesis is rejected,

meaning the phenotype significantly occurred within the test set. In this case, we consider

the samples within the bicluster to be of a highly consistent phenotype.

695 The propensity score matching method

Our aim was to assess whether the association between identified gene modules and

disease phenotypes was reliable. For each gene module, we first identified the underlying

confounding variables that affected its occurrence. We included diet, age, race, and

health status, as well as the other gene modules, and calculated a propensity score for

each sample using the R package MatchIt 4.2.0. We then matched each sample with the

gene module to a sample without the gene module but has a similar propensity score with

the former, obtaining paired samples consisting of samples with and without the gene

module. This was achieved by setting the parameter method = "optimal". Finally, for each

phenotype, we used Wilcoxon signed-rank test to test the null hypothesis that there is no

difference in the phenotype occurrence between sample groups with the gene module

and those without. The *p*-value was adjusted using the R package qvalue 2.18.0. The

q-value less than 0.20 means the gene module is significantly associated with the

phenotype.

Functional analysis of gene modules

We performed an analysis of enriched functions for each gene module. All UniRef90

identifiers were firstly mapped to MetaCyc reactions[28,54]. The reactions were further

associated with MetaCyc pathways[28,54], which enabled a transitive association between

UniRef90 gene families and MetaCyc pathways. We defined a pathway that was satisfied

when the reactions within the pathway significantly occurred via the hypergeometric test,

716
$$P\left(t_{ij}\right) = \frac{\binom{n_j}{t_{ij}}\binom{N-n_j}{m_i-t_{ij}}}{\binom{N}{m_i}} \tag{4}$$

717
$$p - value = \sum_{t_{ij}>T_{ij}}^{n_j} P(t_{ij}) \tag{5}$$

where N is the total number of reactions that were associated with the UniRef90 gene

families and pathways in MetaCyc, m_i is the number of reactions within the ith gene

modules, n_j is the total number of reactions within the jth pathway, and T_{ij} is the

number of observed reactions of pathway j occurring in gene module i. This yielded a

p-value for each gene module and pathway. The resulting *p*-values were adjusted using

the Benjamini-Hochberg procedure for false discovery rates. We selected the pathways

with a *q*-value less than 0.05 as the significantly enriched pathways of each gene module.

We used a widely used tool, LEfSe[11], to infer differentially abundant pathways among

different phenotypes. The pathway abundance information for each sample is from

HUMAnN2. Then, we compared the differentially abundant pathways and significantly

enriched pathways of gene modules from IDAM.

Comparison of IDAM with other methods

The performance of gene module inference of IDAM was compared with the other four

top-performing unsupervised tools for grouping genes into functional modules [55],

including biclustering-based (QUBIC[34], ISA[35], and FABIA[36]) and decomposition-based

(fastICA, an improvement of ICA[33]) methods. All tools were tested and compared based

on the gene-sample matrix constructed. We ran these tools on the Pitzer cluster of the

Ohio Supercomputer Center with memory usage set to 300GB[56] with default parameters.

QUBIC was implemented using the published source code, while the other three

algorithms were implemented using the R packages, isa2 0.3.5, fabia 2.36.0, and fastICA

1.2.2, respectively.

Species-level analysis

For the gene modules from IDAM, ICA[33], QUBIC[34], and FABIA[36], we extracted species

that contribute to the abundance of genes within the identified gene modules, based on

the HUMAnN2 output[28]. In this way, community-level gene expression (abundance) was

decomposed into species-level data. A phylogenetic tree was generated using the

software GraPhlAn[57]. Since the gene modules have been associated with phenotypes

using Z-tests, we assigned the species of each gene module to the corresponding shared

sample phenotype of the samples within the same bicluster. We collected

disease-associated species from the databases Peryton and MicrophenoDB[58], and

previous taxonomic studies. For each phenotype, we counted the total number of species,

as well as the number that matched with the collected species. Then, the consistency

index was calculated as the total number of species in the phenotype dividing by the

number of matched species. We used the average of the consistency indexes across the

three phenotypes as the average consistency index.

LEfSe[11], was used here for comparison. It uses species profiling from MetaPhlAn2 to

identify statistically different species among phenotypes[59,60]. The significantly different

species of each phenotype were aligned to the collected species, from which we

determined the number of matched species and calculated the consistency of each

phenotype.

Gene module validation

We used the 79 samples from an initial round associated with IBD as an independent

dataset to validate the reproducibility of gene modules from IDAM. For each gene module

inferred from IDAM (a total of 39 modules), we test the null hypothesis that the mean

difference in centered log-ratio expression between patients with and without certain

phenotypes was zero using Wilcoxon signed-rank test based on propensity score-based

paired samples. We calculated the q-values using the qvalue R package 2.18.0. The

q-value less than 0.20 means there was a significant difference in the mean expression of

the gene module between patients with and without the phenotype. That was, the gene

module was significantly associated with the phenotype of samples in the independent

dataset.

Survival analysis

We assessed the associations of the seven identified gene modules of melanoma with

PFS, respectively. The samples in each melanoma-associated bicluster were considered

as with the corresponding gene module. In this way, we determined which gene modules

a sample contains. Given that a sample may contain multiple gene modules, it may be

influenced by other gene modules when measuring whether a gene module is associated

with PFS. For each gene module, we first took all the other gene modules as underlying

confounding variables to calculate the propensity score for each sample and constructed

paired sample groups, corresponding to with or without the gene module groups. Then,

the Kaplan-Meier curves of each group were plotted using the "ggsurvplot" function in the

survival R package[61]. The Cox proportional hazards model was used to explore specific

relationships between the gene module and PFS, adjusting for age, antibiotic use in the

last 6 months, sex, BMI, and disease stage. This was achieved using the "coxph" function

in the survival and survminer R packages[62]. Additionally, we identified species and

pathways associated with the gene module with shorter (Gene module 1) or longer PFS

(Gene modules 2, 4, and 5), which is the same as the description in subsections

"Functional analysis of gene modules" and "Species-level analysis".

Data availability

The IBD-associated dataset supporting the conclusions of this article is available in the

IBDMDB database (https://ibdmdb.org). The melanoma, T1DM, and IBS associated

datasets supporting the conclusions of this article are available on NCBI Sequence Read

Archive, with the accession number of SRP197281, SRP072561, and ERP001739,

respectively.

Competing interest statement

The authors declare no competing interests.

Acknowledgements

This work was supported by National Key R&D Program of China (2020YFA0712400),

National Nature Science Foundation of China (NSFC, 61772313 and 11931008), and

Interdisciplinary Science Innovation Group Project of Shandong University (2019).

Author contributions

Q.M. and B.L. conceived the basic idea. Z.L. carried out the computational analysis and data interpretation. L.Z., Q.W., and A.M. designed and drew the figures. D.C. and J.Z. helped polish the manuscript and all the authors wrote the manuscript. All authors read and approved the final manuscript.

References

Gilbert, J. A. *et al.* Current understanding of the human microbiome. *Nature medicine* **24**, 392-400 (2018).

Sepich-Poore, G. D. *et al.* The microbiome and human cancer. *Science* **371**, eabc4552 (2021).

Consortium, T. H. M. P. A framework for human microbiome research. *Nature* **486**, 215-221 (2012).

Proctor, L. M. *et al.* The Integrative Human Microbiome Project. *Nature* **569**, 641-648 (2019).

Qin, J. *et al.* A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* **464**, 59-65 (2010).

Turnbaugh, P. J. *et al.* An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* **444**, 1027-1031 (2006).

Dubinsky, M. & Braun, J. Diagnostic and Prognostic Microbial Biomarkers in Inflammatory Bowel Diseases. *Gastroenterology* **149**, 1265-1274.e3 (2015).

Lecuit, M. & Lortholary, O. Immunoproliferative small intestinal disease associated with Campylobacter jejuni. *Medecine et maladies infectieuses* **35**, S56-58 (2005).

Paulson, J. N., Stine, O. C., Bravo, H. C. & Pop, M. Differential abundance analysis for microbial marker-gene surveys. *Nature methods* **10**, 1200-1202 (2013).

10 Liu, Z. *et al.* Network analyses in microbiome based on high-throughput

multi-omics data. *Briefings in Bioinformatics* **22**, 1639-1655 (2020).

11 Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome*

Biology **12**, R60 (2011).

12 Zhu, Q., Jiang, X., Zhu, Q., Pan, M. & He, T. Graph Embedding Deep Learning

Guides Microbial Biomarkers' Identification. *Frontiers in Genetics* **10** (2019).

13 Wu, Y. *et al.* Identification of microbial markers across populations in early

detection of colorectal cancer. *Nature Communications* **12**, 3063 (2021).

14 Snipen, L., Angell, I.-L., Rognes, T. & Rudi, K. Reduced metagenome sequencing

for strain-resolution taxonomic profiles. *Microbiome* **9**, 79 (2021).

15 Minot, S. S. & Willis, A. D. Clustering co-abundant genes identifies components of

the gut microbiome that are reproducibly associated with colorectal cancer and

inflammatory bowel disease. *Microbiome* **7**, 110 (2019).

16 Zhu, Z., Ren, J., Michail, S. & Sun, F. MicroPro: using metagenomic unmapped

reads to provide insights into human microbiota and disease associations.

Genome Biology **20**, 154 (2019).

17 Franzosa, E. A. *et al.* Relating the metatranscriptome and metagenome of the

human gut. *Proceedings of the National Academy of Sciences* **111**, E2329-E2338

(2014).

18 Kasmanas, J. C. *et al.* HumanMetagenomeDB: a public repository of curated and

standardized metadata for human metagenomes. *Nucleic Acids Research* **49**,

D743-D750 (2020).

19 Ma, A., Sun, M., McDermaid, A., Liu, B. & Ma, Q. MetaQUBIC: a computational

pipeline for gene-level functional profiling of metagenome and metatranscriptome.

Bioinformatics **35**, 4474-4477 (2019).

20 Lathe III, W. C., Snel, B. & Bork, P. Gene context conservation of a higher order

than operons. *Trends in biochemical sciences* **25**, 474-479 (2000).

21   Che, D., Li, G., Mao, F., Wu, H. & Xu, Y. Detecting uber-operons in prokaryotic

genomes. *Nucleic acids research* **34**, 2418-2427 (2006).

22   Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory

bowel diseases. *Nature* **569**, 655-662 (2019).

23   Peters, B. A. *et al.* Relating the gut metagenome and metatranscriptome to

immunotherapy responses in melanoma patients. *Genome medicine* **11**, 61

(2019).

24   Heintz-Buschart, A. *et al.* Integrated multi-omics of the human gut microbiome in a

case study of familial type 1 diabetes. *Nature microbiology* **2**, 16180 (2016).

25   Durbán, A. *et al.* Instability of the faecal microbiota in diarrhoea-predominant

irritable bowel syndrome. *FEMS microbiology ecology* **86**, 581-589 (2013).

26   Niu, S.-Y. *et al.* Bioinformatics tools for quantitative and functional metagenome

and metatranscriptome data analysis in microbes. *Briefings in Bioinformatics* **19**,

1415-1429 (2017).

27   Xie, J. *et al.* QUBIC2: a novel and robust biclustering algorithm for analyses and

interpretation of large-scale RNA-Seq data. *Bioinformatics* **36**, 1143-1149 (2020).

28   Franzosa, E. A. *et al.* Species-level functional profiling of metagenomes and

metatranscriptomes. *Nature Methods* **15**, 962-968 (2018).

29   Morgan, X. C. *et al.* Dysfunction of the intestinal microbiome in inflammatory

bowel disease and treatment. *Genome Biology* **13**, R79 (2012).

30   Imhann, F. *et al.* Interplay of host genetics and gut microbiota underlying the

onset and clinical presentation of inflammatory bowel disease. *Gut* **67**, 108-119

(2018).

31   Klaassen, M. A. Y. *et al.* Anti-inflammatory Gut Microbial Pathways Are

Decreased During Crohn's Disease Exacerbations. *Journal of Crohn's and Colitis*

13, 1439-1449 (2019).

32 Schirmer, M., Garner, A., Vlamakis, H. & Xavier, R. J. Microbial genes and

pathways in inflammatory bowel disease. *Nat Rev Microbiol* **17**, 497-511 (2019).

33 Hyvärinen, A. & Oja, E. Independent component analysis: algorithms and

applications. *Neural Networks* **13**, 411-430 (2000).

34 Li, G., Ma, Q., Tang, H., Paterson, A. H. & Xu, Y. QUBIC: a qualitative biclustering

algorithm for analyses of gene expression data. *Nucleic Acids Research* **37**,

e101-e101 (2009).

35 Bergmann, S., Ihmels, J. & Barkai, N. Iterative signature algorithm for the analysis

of large-scale gene expression data. *Physical review. E, Statistical, nonlinear, and*

soft matter physics **67**, 031902 (2003).

36 Hochreiter, S. *et al.* FABIA: factor analysis for bicluster acquisition. *Bioinformatics*

26, 1520-1527 (2010).

37 O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: current

status, taxonomic expansion, and functional annotation. *Nucleic acids research*

44, D733-745 (2016).

38 Santos-Zavaleta, A. *et al.* RegulonDB v 10.5: tackling challenges to unify classic

and high throughput knowledge of gene regulation in E. coli K-12. *Nucleic acids*

research **47**, D212-D220 (2019).

39 Viladomiu, M. *et al.* Adherent-invasive E. coli metabolism of propanediol in

Crohn's disease regulates phagocytes to drive intestinal inflammation. *Cell Host*

Microbe **29**, 607-619.e8 (2021).

40 Franzosa, E. A. *et al.* Gut microbiome structure and metabolic activity in

inflammatory bowel disease. *Nat Microbiol* **4**, 293-305 (2019).

41 Kuballa, A., Geraci, M., Johnston, M. & Sorrentino, D. The Gut Microbial Profile of

Preclinical Crohn's Disease Is Similar to That of Healthy Controls. *Inflammatory*

bowel diseases **26**, 1682-1690 (2020).

42   Vich Vila, A. *et al.* Gut microbiota composition and functional changes in

inflammatory bowel disease and irritable bowel syndrome. *Sci Transl Med* **10**,

eaap8914 (2018).

43   Skoufos, G. *et al.* Peryton: a manual collection of experimentally supported

microbe-disease associations. *Nucleic Acids Research* **49**, D1328-D1333 (2020).

44   Wu, J. *et al.* Rhein modulates host purine metabolism in intestine through gut

microbiota and ameliorates experimental colitis. *Theranostics* **10**, 10665-10679

(2020).

45   Koo, B. K. *et al.* Identification of novel autoantibodies in type 1 diabetic patients

using a high-density protein microarray. *Diabetes* **63**, 3022-3032 (2014).

46   Wang-Sattler, R. *et al.* Novel biomarkers for pre-diabetes identified by

metabolomics. *Molecular systems biology* **8**, 615 (2012).

47   Mahendran, Y. *et al.* Glycerol and fatty acids in serum predict the development of

hyperglycemia and type 2 diabetes in Finnish men. *Diabetes care* **36**, 3732-3738

(2013).

48   Duvallet, C., Gibbons, S. M., Gurry, T., Irizarry, R. A. & Alm, E. J. Meta-analysis of

gut microbiome studies identifies disease-specific and shared responses. *Nature*

Communications **8**, 1784 (2017).

49   Luo, Y., Ma, Q., Zhang, C. & Zhang, A. R. in *2022 IEEE International Conference*

on Acoustics, Speech and Signal Processing (2022).

50   Suzek, B. E., Huang, H., McGarvey, P., Mazumder, R. & Wu, C. H. UniRef:

comprehensive and non-redundant UniProt reference clusters. *Bioinformatics* **23**,

1282-1288 (2007).

51   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local

alignment search tool. *Journal of molecular biology* **215**, 403-410 (1990).

52   Wang, Z., Li, G., Robinson, R. W. & Huang, X. UniBic: Sequential row-based

biclustering algorithm for analysis of gene expression data. *Scientific reports* **6**,

23466 (2016).

53   Ben-Dor, A., Chor, B., Karp, R. & Yakhini, Z. Discovering local structure in gene

expression data: the order-preserving submatrix problem. *Journal of*

computational biology : a journal of computational molecular cell biology **10**,

373-384 (2003).

54   Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes - a

2019 update. *Nucleic acids research* **48**, D445-d453 (2020).

55   Saelens, W., Cannoodt, R. & Saeys, Y. A comprehensive evaluation of module

detection methods for gene expression data. *Nature Communications* **9**, 1090

(2018).

56   Ohio Supercomputer Center. Pitzer Supercomputer.

http://osc.edu/ark:/19495/hpc56htp (2018).

57   Asnicar, F., Weingart, G., Tickle, T. L., Huttenhower, C. & Segata, N. Compact

graphical representation of phylogenetic data and metadata with GraPhlAn. *PeerJ*

3, e1029 (2015).

58   Yao, G. *et al.* MicroPhenoDB Associates Metagenomic Data with Pathogenic

Microbes, Microbial Core Genes, and Human Disease Phenotypes. *Genomics,*

Proteomics & Bioinformatics **18**, 760-772 (2021).

59   Segata, N. *et al.* Metagenomic microbial community profiling using unique

clade-specific marker genes. *Nature methods* **9**, 811-814 (2012).

60   Truong, D. T. *et al.* MetaPhlAn2 for enhanced metagenomic taxonomic profiling.

Nature Methods **12**, 902-903 (2015).

61   Therneau, T. M. & Lumley, T. Package 'survival'. *R Top Doc* **128**, 28-33 (2015).

62   Andersen, P. K. & Gill, R. D. Cox's regression model for counting processes: a

large sample study. *The annals of statistics* **10**, 1100-1120 (1982).

Supplementary Information

Supplementary information file

This file contains Supplementary Figures S1-3, details in the Problem formulation and

Methods part, and Supplementary references.

Supplementary Tables

This file includes an overview of Supplementary Tables and Supplementary Tables 1-12.