1   A natural genetic variation screen identifies insulin signaling, neuronal communication, and

2   innate immunity as modifiers of hyperglycemia in the absence of *Sirt1*

4   Rebecca A.S. Palu

5   Biology Department, Purdue University-Fort Wayne, Fort Wayne, IN 46818

Running title: Natural modifiers of hyperglycemia

Key words: hyperglycemia, *Drosophila*, genetic variation, modifier genes

Corresponding author:

Rebecca A.S. Palu

Biology Department

Science Building SB330

2101 E. Coliseum Blvd

Fort Wayne IN, 46805

260-481-5704

palur@pfw.edu

ABSTRACT

Variation in the onset, progression, and severity of symptoms associated with metabolic

disorders such as diabetes impairs the diagnosis and treatment of at-risk patients. Diabetes

symptoms, and patient variation in these symptoms, is attributed to a combination of genetic

and environmental factors, but identifying the genes and pathways that modify diabetes in

humans has proven difficult. A greater understanding of genetic modifiers and the ways in which

they interact with metabolic pathways could improve the ability to predict a patient's risk for

severe symptoms, as well as enhance the development of individualized therapeutic

approaches. In this study we use the *Drosophila* Genetic Reference Panel (DGRP) to identify

genetic variation influencing hyperglycemia associated with loss of *Sirt1* function. Through

analysis of individual candidate functions, physical interaction networks, and Gene Set

Enrichment Analysis (GSEA) we identify not only modifiers involved in canonical glucose

metabolism and insulin signaling, but also genes important for neuronal signaling and the innate

immune response. Furthermore, reducing the expression of several of these candidates

suppressed hyperglycemia, making them ideal candidate therapeutic targets. These analyses

showcase the diverse processes contributing to glucose homeostasis and open up several

avenues of future investigation.

INTRODUCTION

Metabolic diseases, and in particular diabetes, are one of the most pressing health crises in the

developed world, with incidences continuing to rise in the last 20 years (CDC 2020). 37.7% of

adults in the US are diagnosed as obese and 10.5% as having some form of diabetes, and it is

estimated that millions more go undiagnosed (Flegal *et al.* 2016; CDC 2020). What is more, the

monetary cost of these disorders to the public has grown to astronomical levels. It is estimated

that $237 billion in direct costs and at least $90 billion in indirect costs were spent on healthcare

related to diabetes and its various complications in 2017 alone, up ~25% from 2012 (Flegal *et*

al. 2016). A focused effort has been made to understand both the genetic and environmental

contributors to metabolic homeostasis, as well as to the disruption of that homeostasis that

leads to disease (Barroso and McCarthy 2019).

Unfortunately, even identifying these contributors has proven difficult. Metabolic diseases are

complex, and the onset, progression, and ultimately the severity of any individual case is

dependent upon a myriad of genetic and environmental variables and the ways in which they

interact with one another (Queitsch *et al.* 2012; Barroso and McCarthy 2019). Even when there

is a strong familial link, phenotypic heterogeneity in disease phenotypes can make it difficult to

identify at-risk patients or make accurate prognostic predictions (Udler *et al.* 2019). This is

particularly true when it comes to predicting complications of diabetes such as neuropathy,

retinopathy, or kidney disease (Barroso and McCarthy 2019; Cabrera *et al.* 2020). Much of this

variation is due to inter-individual differences in genetic background, including silent cryptic

genetic variation that is revealed upon disease or stress (Queitsch *et al.* 2012; Chow 2016;

Barroso and McCarthy 2019).

One example of this kind of symptom heterogeneity can be observed in disease associated with

the deacetylase *SIRT1*. This highly conserved gene was originally identified as a histone

deacetylase important in heterochromatin formation in yeast (Shore *et al.* 1984; Ivy *et al.* 1986;

Rine and Herskowitz 1987). Since then, SIRT1 and its paralogs (the sirtuins) have been found

to have a number of additional targets, many of which are transcription factors and enzymes

with key roles in metabolic homeostasis (Brunet *et al.* 2004; Picard *et al.* 2004; Rodgers and

Puigserver 2007; Li *et al.* 2007; Yang *et al.* 2009; Palu and Thummel 2016). Importantly, as part

of their enzymatic reaction, sirtuins consume the cofactor NAD, which also serves as an

electron carrier in central metabolic pathways such as glycolysis and the TCA cycle. Sirtuin

enzymatic activity, therefore, is directly linked to the availability of this cofactor and thus is

responsive to the energetic state of the cell. This information is then conveyed to the targets,

whose acetylation state alters their activity and stability in the cell (Nogueiras *et al.* 2012). With

this centralized role in regulating the response of metabolic factors to cellular energy availability,

it is unsurprising that variation in *SIRT1* has been linked to, among other things, the

development of diabetes (Zillikens *et al.* 2009; Botden *et al.* 2012; Biason-Lauber *et al.* 2013;

Zhao *et al.* 2017).

Elucidating the mechanism behind this link, however, has proven difficult. Loss-of-function and

gain-of-function studies in model systems have demonstrated a clear role for SIRT1 in

metabolic homeostasis, but the actual impacts on the animals in question have frequently been

contradictory (Boutant and Cantó 2014). It is likely that at least some of these contradictory

results stem from differences in genetic background between the animals used in the various

studies. Understanding the role of this variation and the genes or pathways which modify

metabolic disease will enable the development of improved diagnosis, prediction of prognosis,

and personalized treatment strategies for patients.

Model organism tools, such as the *Drosophila* Genetic Reference Panel (DGRP), provide a way

to study of the impact of natural genetic variation on diseases such as diabetes (Mackay *et al.*

2012; He *et al.* 2014; Ivanov *et al.* 2015; Nelson *et al.* 2016; Jehrke *et al.* 2018; Everman *et al.*

2019). The DGRP is a collection of ~200 isogenic strains derived from a wild population, such

that each strain represents one wild-derived genome (Mackay *et al.* 2012). The variation in the

DGRP is well tolerated under healthy, non-disease conditions and allows for the identification of

genetic polymorphisms that are associated with phenotypic variation in models of human

disease (Chow and Reiter 2017). Importantly, the availability of full-genome sequence for these

strains allows for genome-wide association analyses that link quantitative phenotypes with

genetic variation and modifier genes.

The utility of the DGRP in identifying candidate modifiers of metabolic disease has already been

demonstrated numerous times, with screens associated with misfolded insulin, high sugar and

high fat feeding, and starvation resistance already documented (Mackay *et al.* 2012; He *et al.*

2014; Ivanov *et al.* 2015; Nelson *et al.* 2016; Jehrke *et al.* 2018; Everman *et al.* 2019). While

some of these used biochemical assays to precisely measure metabolite levels in flies as a

quantitative phenotype for the screen (Nelson *et al.* 2016; Jehrke *et al.* 2018; Everman *et al.*

2019), several used more general physiological measurements such as starvation resistance

and lifespan (Mackay *et al.* 2012; Ivanov *et al.* 2015). Although effective, the use of this kind of

general readout reduces the specificity of the modifiers identified. Many genetic factors impact

survival and could lead to a high background signal. The same could be true for otherwise wild-

type flies subjected to different environmental conditions, even when more precise assays are

used as a quantitative phenotype (Nelson *et al.* 2016). A multitude of pathways and processes

impact the response to dietary changes through feeding rate, hormone secretion, anabolism

and catabolism rates, and nutrient absorption. Using a specific genetic model of disease and

then focusing on a specific phenotype, such as hyperglycemia, may reduce some of the noise

and increase specificity of the modifiers identified.

Furthermore, recent studies using the DGRP have demonstrated that the top candidate modifier

genes and pathways differ when different but related models of genetic disease are screened

(Chow *et al.* 2016; Palu *et al.* 2019). These results reinforce the idea that different causative

genes and mutations will interact with different pathways over the course of disease. It also

highlights the importance of exploring multiple disease models in something so diverse as

diabetes, and the utility of the DGRP in precisely distinguishing modifiers of a particular

genotype-phenotype combination.

In this study, we report the results of a natural variation screen in a model of *Sirt1* loss-of-

function. Loss of this gene in *Drosophila* has been shown to lead to progressive metabolic

dysfunction including obesity, hyperglycemia, and ultimately insulin resistance (Palu and

Thummel 2016). Our study design is specifically focused on the phenotype of hyperglycemia

when *Sirt1* expression is disrupted using RNAi in the adipose and liver-like fat body organ. We

observed substantial phenotypic variation across the DGRP for hyperglycemia associated with

loss of *Sirt1*. Using genome-wide association analysis, pathway enrichment, and the generation

of a physical interaction network, we identified a number of modifying pathways and processes,

several of which have known roles in central carbon metabolism, the immune response, and the

kind of neuronal signaling and communication expected to influence the neuroendocrine cells

responsible for insulin secretion in *Drosophila*. Finally, we confirmed that loss of several of the

top candidate modifier genes significantly alters glucose levels in the *Sirt1* RNAi model. Our

findings highlight exciting new areas of study for modifiers of *Sirt1* function, glucose

homeostasis, and insulin sensitivity.

METHODS

Fly stocks and maintenance

Flies were raised at room temperature on a diet based on the Bloomington Stock Center

standard medium with malt. Experimental crosses were maintained on a media containing 6%

yeast, 6% dextrose, 3% sucrose, and 1% agar, with 0.6% propionic acid and 0.1% p-Hydroxy-

benzoic acid methyl ester in 95% ethanol included as antifungal agents. Flies subjected to an

overnight fast were transferred to media containing only 1% agar in water. The *R4>Sirt1i* strain,

which serves as the model of hyperglycemia in this study, is derived from an *R4-GAL4* strain

(BDSC 33832) outcrossed to w^{1118} (Palu and Thummel 2016) and a *Sirt1* RNAi strain from the

Bloomington Stock Center (32481). 185 strains from the DGRP were used for the modifier

screen (Tables S1-S3), wherein virgin females carrying the *R4>Sirt1i* model were crossed to

males of the DGRP strains. Male F1 progeny carrying *R4>Sirt1i* were separated and aged for

one to two weeks. These flies were then either collected under ad libitum fed conditions or

fasted overnight and then collected. The following RNAi and control strains are from the

Bloomington Stock Center: *CG4168* RNAi (28636), *CG5888* RNAi (62175), *uif* RNAi (38354),

CTPSyn RNAi (31924), *smt3* RNAi (36125), *ilp5* RNAi (33683), *Vha55* RNAi (40884), *snRNP-*

U1-70k RNAi (33396), *CG10265* RNAi (43294), *CG15803* RNAi (51449), *Roe* RNAi (57836),

CG34353 RNAi (58291), *CadN2* RNAi (27508), *Ace* RNAi (25958), *CG43897* (31560), *dsxc73A*

RNAi (56987), *bgm* RNAi (56979), *CG3407* RNAi (57762), control *attP40* (36304), and control

attP2 (36303).

Glucose Assay

Samples of five flies each were collected at one or two weeks of age and washed in 1XPBS.

Samples were then either frozen in liquid nitrogen and stored long-term at -80C or immediately

homogenized in 100 uL 1X PBS. Frozen samples were kept frozen until immediately upon

addition of PBS and homogenization. After homogenization samples were subjected to heat

inactivation of enzymes at 70C for approximately 10 minutes. Glucose was measured undiluted

from the lysate using the Sigma HK Glucose Assay kit as described (Tennessen *et al.* 2014).

Protein Assay

Prior to heat inactivation, 10 uL of the fly lysate isolated for glucose measurement was saved

and kept on ice. Protein samples could then be stored long term at -80ºC. Samples were

centrifuged for up to 5 minutes at room temperature and protein measured from the supernatant

after a 1:10 dilution using the Sigma Protein Assay Reagent as described (Tennessen *et al.*

2014).

Phenotypic analysis and genome-wide association

183 For each DGRP line, glucose was measured from 3 samples of 5 flies each. The P-values for

184 association of genetic background and glucose concentration were calculated using one-way

185 ANOVA on R software taking into account all collected data points for each experiment.

186 Average glucose concentration was used for the genome-wide association (GWA). GWA was

187 performed as previously described (Chow *et al.* 2016; Palu *et al.* 2019). DGRP genotypes were

188 downloaded from the website, http://dgrp.gnets.ncsu.edu/. Non-biallelic sites were removed. A

189 total of 3,636,891 variants were included in the analysis. Mean eye glucose concentration for

190 555 samples representing 2775 DGRP/R4>Sirt1i F1 progeny were regressed on each SNP. To

191 account for cryptic relatedness (He *et al.* 2014; Huang *et al.* 2014), GEMMA (v. 0.94) (Zhou and

192 Stephens 2012) was used to both estimate a centered genetic relatedness matrix and perform

193 association tests using the following linear mixed model (LMM):

$$y = \alpha + x\beta + u + \epsilon$$

$$u \sim MVN_n (0, \lambda \tau^{-1} K)$$

$$\epsilon \sim MVN_n (0, \tau^{-1} I_n)$$

197 where, as described and adapted from Zhou and Stephens 2012, y is the n-vector of average

198 glucose concentration for the n lines, α is the intercept, x is the n-vector of marker genotypes, β

199 is the effect size of the marker. u is a n x n matrix of random effects with a multivariate normal

200 distribution (MVN_n) that depends on λ, the ratio between the two variance components, τ^{-1},

201 the variance of residuals errors, and where the covariance matrix is informed by K, the

202 calculated n x n marker-based relatedness matrix. K accounts for all pairwise non-random

203 sharing of genetic material among lines. ϵ, is a n-vector of residual errors, with a multivariate

204 normal distribution that depends on τ^{-1} and I_n, the identity matrix. Quantile-quantile (qq)

205 plots demonstrate an appropriate fit to the LMM at the positive end of the plot, but a greater

206 number of points than expected by chance with an insignificant p-value (Figure S1). Genes

207 were identified from SNP coordinates using the BDGP R54/dm3 genome build. A SNP was

208 assigned to a gene if it was +/- 1 kb from a gene body.

RNAi Validation

Virgin females from the *R4>Sirt1i* model were crossed to males carrying RNAi constructs

targeting candidate modifiers of those models, and the glucose levels of F1 male progeny

expressing both *Sirt1i* and the modifier RNAi construct specifically in the fat body were

measured as described above on 4-5 samples of 5 male flies each. Glucose concentrations

from RNAi-carrying strains are compared directly to genetically matched *attP40* or *attP2*

controls using a Dunnett's multiple comparisons test. Glucose measurements are normalized to

the appropriate genetically matched controls. Normalized controls from individual experiments

are compared in Figure S2. Standard deviation did not significantly vary between controls for

individual experiments.

Bioinformatics Analysis

Genetic polymorphisms were associated with candidate genes within 1 kb of the polymorphism.

Information about candidate genes and their human orthologues was gathered from a number

of databases including Flymine, Flybase, OMIM, and NCBI. Physical interaction maps were

generated using the GeneMANIA plugin on Cytoscape (version 3.8.2) (Shannon *et al.* 2003;

Montojo *et al.* 2010). GSEA was run to generate a rank-list of genes based on their enrichment

for significantly associated polymorphisms. For GSEA analysis, polymorphisms within 1kb of

more than 1 gene were assigned to one gene based on a priority list of exon, UTR, intron, and

upstream or downstream. Genes were assigned to GO categories, and calculation of

enrichment score was performed as described (Subramanian *et al.* 2005). Categories with ES

scores > 0 (enriched for associated genes with low p-values), gene number > 3, and p-values

<0.05 were included in the final output.

RESULTS

Glucose levels in R4>Sirt1i flies vary with genetic background in a consistent pattern across

multiple conditions

Loss of *Sirt1* expression leads to progressive hyperglycemia, obesity, and insulin resistance. To

model the hyperglycemia that is commonly associated with diabetes, we reduced the

expression of the deacetylase *Sirt1* specifically in the fat body of *Drosophila melanogaster*.

RNAi targeting *Sirt1* specifically in the fat body reproduces the hyperglycemia phenotype, with

an approximately 50-60% increase in whole fly glucose levels (P = 0.02, Figure S3) (Palu and

Thummel 2016). The fat body in *Drosophila* performs functions normally undertaken by the

adipose and liver tissues in humans (Géminard *et al.* 2009; DiAngelo and Birnbaum 2009;

Arrese and Soulages 2010). While there are likely roles for *Sirt1* in other metabolic tissues, its

function in the fat body clearly contributes to the maintenance of glucose homeostasis and

insulin sensitivity over time (Palu and Thummel 2016). Modifiers of insulin signaling or glucose

metabolism could alter the degree of hyperglycemia in this model.

The degree of hyperglycemia is determined by a biochemical assay that measures the

concentration of glucose in a whole-fly lysate (Tennessen *et al.* 2014). This is a quantitative

assay where higher concentrations of glucose correlate with more severe hyperglycemia, and

lower concentrations of glucose correlate with milder disease, or potentially hypoglycemia.

Glucose was measured specifically in males, as adult female *Drosophila* devote much of their

physiological output to egg production (Millington and Rideout 2018). Previous studies have

demonstrated that adult males provide a consistent model for metabolic homeostasis in the fly

(Sieber and Thummel 2009; Tennessen *et al.* 2014; Palu and Thummel 2016; Barry and

Thummel 2016; Beebe *et al.* 2020).

Loss of *Sirt1* is induced using the *GAL4/UAS* system, where *R4-GAL4* drives expression of

UAS-Sirt1 RNAi (Figure S4A). *R4-GAL4* is strongly expressed primarily in the fat body of fly,

starting in early development and continuing through adulthood (Lee and Park 2004). The line

containing this model (*R4>Sirt1i*) serves as the donor strain that was crossed to each DGRP

strain. Females from the donor strain were crossed with males of each of 185 DGRP strains to

generate F1 progeny lacking *Sirt1* expression in the fat body. The progeny received 50% of their

genome from the maternal donor strain and 50% from the paternal DGRP strain (Figure S4B).

Therefore, we are measuring the dominant effect of the DGRP background on the *Sirt1* RNAi

hyperglycemia phenotype. This experimental design is similar to a model of *NGLY1* deficiency

using RNAi that was also crossed to the DGRP (Talsness *et al.* 2020).

In the prior study characterizing the impact of the loss of function of *Sirt1* on metabolic

homeostasis, the dysfunction observed was progressive in nature. While larvae and young

adults are immediately obese, hyperglycemia and insulin resistance set in and become worse

with increasing age (Reis *et al.* 2010; Palu and Thummel 2016). To ensure an appropriate set of

conditions with respect to diet and age, we performed a preliminary analysis on 37 DGRP

strains at one or two weeks of age, and under fasted or ad libitum fed conditions (Table S1).

These time points correspond to conditions in the original study where differences in the

metabolic state of the flies corresponded to differences in glycemia (Palu and Thummel 2016).

We wished to select a time point at which hyperglycemia was detectable, but also differed

enough between strains to allow us to identify genetic variation associated with that

heterogeneity.

Three samples were collected for glucose measurements in each strain and condition. Glucose

levels vary across genetic background for each of the conditions being tested (Figure 1A-D).

Average glucose for each strain is significantly correlated between one and two week fasted

flies (R = 0.53, P = 4E-03), between one-week-old fed and fasted flies (R = 0.45, P = 0.020),

and between one-week-old fed and two-week-old fasted flies (R = 0.62, P = 2E-04) (Figure

S5A-C). This supports glucose concentration as a consistent quantitative measurement.

Interestingly, evidence for correlation with two-week-old flies fed ad libitum is not as strong.

While a significant correlation is still detected with one-week-old fasted flies (R = 0.43, P =

0.014), the correlation is not significant with two-week-old fasted flies (R = 0.25, P = 0.219) and

one-week-old flies fed ad libitum (R = 0.34, P = 0.063) (Figure S5D-F). By two weeks of age,

Sirt1 loss-of-function flies are beginning to experience more severe symptoms of disease, and

we expect to see variability in symptoms and behavior in response to those symptoms. Fed flies

at two weeks may have more variable glucose because feeding behavior is a big contributor to

glucose levels in flies that have not been subjected to a fast.

To identify the conditions under which the impact of genetic background was the strongest, we

performed a one-way ANOVA test that included all data points collected. We found that while

there is a significant association between glucose levels and genetic background under all

conditions (p<0.05), this effect is most pronounced in the one-week-old flies fed ad libitum (P =

1.87E-5) and in the two-week-old fasted flies (P = 1.95E-5) (Figure 1A,D). Because fasting

reduces possible intra-strain variation caused by food in the gut, two weeks fasted was selected

for the full screen.

We examined protein concentrations in 90 samples from the first 30 strains collected at 2 weeks

fasted to ensure that any variation we observe in glucose is not due to differences in body size.

Protein levels do not significantly vary across the DGRP (P = 0.63, Figure S6A), nor do protein

levels correlate with glucose levels in individual samples (R = 0.04, P = 0.6577, Figure S6B).

We conclude that the variation we observe in fasting glucose levels are due to differences in

circulating glucose, and not to differences in body size.

Genome-wide association analysis identifies candidate modifiers of Sirt1i-associated

hyperglycemia

Using the conditions determined in the preliminary screen, we proceeded to cross the donor

strain with the remaining 149 DGRP strains (Figure 2, Tables S2,S3). We found a significant

effect of genetic background on glycemia in the R4>Sirt1i flies (P < 2E-16) using one-way

ANOVA including all data points for each strain (N = 555). Individual glucose measurements

ranged from 0.306 µg/fly to 3.416 µg/fly (Table S2), while average concentrations ranged from

0.395 µg/fly (RAL 801) to 2.438 µg/fly (RAL 357) (Figure 2, Table S3).

To identify genetic polymorphisms that may be responsible for this observed variation in

glycemia, we performed a genome-wide association analysis. Average glucose level for each

strain was used as a quantitative phenotype to test for association with polymorphisms in the

DGRP. A total of 3,636,891 variants were tested for the *R4>Sirt1i* model across 186 lines. This

analysis as a result is insufficiently powered for candidates to remain statistically significant after

multiple testing corrections. Instead, the focus is on identification of candidate modifiers and

pathways that can be validated through further study and that will provide the basis for future

projects. This approach has been quite successful in previous studies (Chow *et al.* 2013, 2015,

2016; Palu and Chow 2018; Lavoy *et al.* 2018; Palu *et al.* 2019, 2020; Talsness *et al.* 2020).

Because the analyzed F1 hybrids in this case were male and inherited their X chromosome from

the donor strain and not the DGRP strain, we do not include any X-linked variants in the

resulting candidate modifiers. Using an arbitrary cut-off of $P < 10^{-4}$, we identified 237

polymorphisms on the second and third chromosomes (Table S4). Of these 237, 62 were not

considered further as they were not within +/- 1 kb of a candidate gene. The remaining 175

polymorphisms are associated with a total of 161 candidate genes (Table S5). 100 of these

polymorphisms are intronic, 29 are exonic with 6 producing non-synonymous changes to the

peptide and one a start-gain, 10 are located in the 5' or 3' untranslated regions, and 36 are

within 1 kb up or downstream of the candidate gene (Table S4). Of note in this analysis is that

the results were not filtered for allele frequency > 0.05. This was a concerted choice; several of

the most interesting candidates, including *CG5888* and *ilp5*, would otherwise have been left out

of the analysis. The number of total variants analyzed drops from 237 to 90, and the number of

candidate genes associated at $P < 10^{-4}$ drops from 161 to 75. While this, along with the use of a

low stringency p-value cut-off, increases the probability of false positives, it likewise increases

our power when performing pathway enrichment. Validation of candidate genes with low minor

allele frequencies in later studies will distinguish true positives from false positives.

One concern with using an RNAi model to reduce *Sirt1* expression is that the modifiers

identified might be specific RNAi efficacy, rather than hyperglycemia. The modifiers could be

altering the degree and efficiency of *Sirt1* knockdown, so that hyperglycemia is actually

correlating with the amount of *Sirt1* expression that is achieved. If this were the case, we would

expect to see the top candidate modifiers associated with RNAi machinery or the efficiency of

the GAL4/UAS system. This does not appear to be the case, either from a single-gene function

perspective or when looking at enriched gene categories (Tables S5 and S6). Furthermore, the

GAL4/UAS system is commonly used to model disease in the DGRP, and the candidates

identified have always been unique to the disease and, at times, even the specific model in

question (He *et al.* 2014; Chow *et al.* 2016; Lavoy *et al.* 2018; Palu *et al.* 2019; Talsness *et al.*

2020). All of this suggests that the candidate genes identified through this screen are modifying

Sirt1-associated hyperglycemia directly rather than altering the degree of *Sirt1* knockdown.

Candidate modifiers of Sirt1 are involved in basic metabolic processes, the immune response,

and the regulation of neuronal communication

Because loss of *Sirt1* in the fat body alters glucose metabolism and insulin sensitivity in the

organism, we expected modifiers of hyperglycemia to impact pathways linked to central carbon

metabolism as well as external processes that influence secretion and signaling of hormones

such as insulin. To determine if this is the case, we examined the individual functions of the top

GWA candidates and looked for pathways and processes that are enriched in this list. While we

attempted first to do this through Gene Ontology analysis of our top candidates, we found no

significantly enriched terms. We therefore utilized individual known physical interactions and GO

term enrichment through GSEA to highlight likely candidate pathways.

Analysis of Candidate Modifiers: We expected our top candidates to include genes that

function in pathways or processes related to *Sirt1* regulation or activity. Among the most

interesting candidates are those involved in NAD metabolism. We identified two genes whose

products are part of the NADH dehydrogenase component of Complex I in the electron transport

chain (*NDUFS3* and *ND-PDSW*) (FlyBase Curators 2008; Gaudet *et al.* 2011). There are also

two NADP kinases, enzymes involved in the generation of NADP from NAD (*CG33156* and

CG6145) (Gaudet *et al.* 2011). *DUOX*, an NADPH oxidase, passes electrons from NADPH to

oxygen, generating hydrogen peroxide and altering the redox balance of the cell (Gaudet *et al.*

2011; Anh *et al.* 2011). As Sirt1 utilizes NAD as a cofactor during its enzymatic reaction, altering

the balance of NAD in the cell through differential regulation of these enzymes could further

impact the activity of other pathways that require NAD as an electron carrier, or exacerbate the

phenotypes associated with *Sirt1* loss-of function (Nogueiras *et al.* 2012). We also identified two

genes that have previously been implicated in the regulation and/or extension of lifespan: *sugb*

and *CG42663* (Landis *et al.* 2003; Paik *et al.* 2012). While the role of *Sirt1* in lifespan extension

is still contested, the identification of other genes implicated in this process suggests shared

functions or pathways.

Genes involved in central glucose metabolism as well as insulin signaling are also candidate

modifiers. MFS5 acts as a transporter of both glucose and trehalose for the uptake of these

sugars from circulation (McMullen *et al.* 2021). Glucose-6-phosphatase (*G6P*) is the last rate-

limiting step in both gluconeogenesis and glycogenolysis, which are used to generate glucose

for release into the body during fasting (Gaudet *et al.* 2011; Lizák *et al.* 2019). These two genes

directly regulate circulating glucose levels. Candidates involved in other metabolic pathways

include *CTPSyn*, which encodes the rate limiting step in cytidine synthesis, the very long chain

fatty acid ligase *bgm*, the mannosidase *Edem2*, and the oxoglutarate dehydrogenase complex

subunit *CG33791* (Kang and Ryoo 2009; Gaudet *et al.* 2011; Jang *et al.* 2015; Sivachenko *et al.*

2016; Zhou *et al.* 2019). Partially responsible for regulating general metabolic flux through these

various pathways is insulin. Interestingly, a top candidate is *ilp5*, one of several insulin-like

peptides expressed in the insulin-producing cells (IPCs) in the *Drosophila* brain (Géminard *et al.*

2009). Our analysis also identified *IA-2*, a phosphatase involved in ilp secretion, *CG4168*, an

uncharacterized gene whose closest human orthologue *IGFALS* encodes a protein that binds to

and stabilizes IGF proteins in circulation, and *wrd*, a subunit in the protein phosphatase PP2A

that negatively regulates insulin and TOR signaling (Boisclair *et al.* 1996; Kim *et al.* 2008; Hahn

et al. 2010).

Other potential modifiers of insulin stability and signaling in circulation are *dally*, *cow*, and *Hs3st-*

A. Both *dally* and *cow* encode heparin sulfate proteoglycans, while *Hs3st-A* encodes an O-

sulfotransferase that acts on these proteoglycans (Filmus and Selleck 2001; Gaudet *et al.* 2011;

Chang and Sun 2014). Previous work has demonstrated an impact of the enzyme heparanase,

which cleaves heparan sulfate, on diabetic autoimmunity and complications such as

nephropathy (Rabelink *et al.* 2017). While this is more peripheral to the central insulin signaling

pathway in *Drosophila*, it highlights the utility of such factors in altering disease processes in

subtle ways.

Another interesting group of candidates are those associated with neuronal development and

function. Several members of the defective proboscis extension response (*dpr*) family were

represented in our list (*dpr2*, *dpr6*, and *dpr13*) along with the dpr-interacting protein *DIP-eta*.

The *dpr* gene family is collectively associated with synapse organization and function, as are the

candidate genes *fife*, *CG32373*, and *atilla* (FlyBase Curators *et al.* 2004; Kurusu *et al.* 2008;

Carrillo *et al.* 2015; Bruckner *et al.* 2017). We also noted candidates involved in neuropeptide

signaling (*rk* and *RYa-R*), voltage-gated potassium channels and their regulation (*CG5888* and

CG1688), and axon guidance (*tutl*, *CadN2*, *CG34353*, and *sbb*) (Rao *et al.* 2000; Luo *et al.*

2005; Prakash *et al.* 2005; Al-Anzi and Wyman 2009; Ida *et al.* 2011; Gaudet *et al.* 2011). The

IPCs in *Drosophila* are actually neuroendocrine cells located in the brain, as are the AKH-

producing cells responsible for secreting the glucagon-like hormone AKH (Géminard *et al.*

2006). The secretion of insulin is therefore dependent upon the correct development,

connection, and signaling of neuronal cells.

The immune response is another generally enriched category of modifier genes. Several

members of the nimrod family of immunoglobulins (*NimB2*, *NimC1*, and *NimC3*) were identified

by GWA. All are implicated in the innate immune response, with *NimC1* and *NimC3* in particular

having roles in phagocytosis (Somogyi *et al.* 2010). In response to insecticides, *LRR* regulates

the immune response through NF-kappaB, whose activation is an early protective event in the

progression and pathology of diabetes (Prisco *et al.* 2013; Irvin *et al.* 2018). Two lysozyme

enzymes with links to bacterial defense (*LysX* and *CG7798*) highlight the role of oxidative stress

and redox homeostasis in the innate immune response (FlyBase Curators 2008). As Sirt1 has

roles in regulating the response to oxidative stress, we looked for other genes with similar

functions (Brunet *et al.* 2004). As described above, *DUOX* plays a part in regulating redox

homeostasis through the production of hydrogen peroxide (Anh *et al.* 2011). *CG42331* encodes

a peroxidase that appears to be strongly enriched in the pupal fat body, and *cyp28a5* encodes

an oxidoreductase that, similar to LRR, is involved in the response to insecticides (FlyBase

Curators *et al.* 2004; Graveley *et al.* 2011; Gaudet *et al.* 2011). It is now believed that Type I

and Type II diabetics both suffer at least to some degree from autoimmunity (Candia *et al.*

2019). Exploring the direct and indirect connections of *Sirt1* to the immune response and

oxidative stress directly is an interesting avenue for future direction.

Physical Interaction Network: We generated a network of physical interactions among the 161

candidate genes identified above. These were identified and visualized using Cytoscape

software with the GeneMania plugin (Shannon *et al.* 2003; Montojo *et al.* 2010). The products of

37/161 candidate genes were found to physically interact with at least one other candidate gene

product with no more than one bridging node represented by a non-candidate gene (Figure 3A).

This high degree of interaction suggests that the modifiers identified in this screen are indeed

functioning through shared processes.

Focusing then on the 37 genes involved in physical interactions, we identified several broad

functional categories that could influence glucose homeostasis in the fly. The most obvious

category are enzymes that catalyze steps in basic metabolic pathways (N = 7). This includes

the NAD kinases (*CG6145* and *CG33156*) and one NADH dehydrogenase (*ND-PDSW*)

discussed above (Gaudet *et al.* 2011). Other metabolic candidate modifiers include *MFS3* and

CTPsyn, the rate limiting enzyme in the production of the nucleotide Cytidine triphosphate (Zhou

et al. 2019). Both of these enzymes function in pathways critically dependent upon or feeding

into central carbon metabolism, and their inclusion as candidate modifiers of hyperglycemia

supports a role for those secondary metabolic pathways as a sink for increased circulating

glucose. *Mtmr6* encodes a phosphatidylinositol phosphatase, a key enzyme in several signaling

pathways, including insulin signaling (Gaudet *et al.* 2011). Also interesting is the gene *LRP1*,

which is orthologous to human LDL receptor related protein 1. In additional to its role in lipid

homeostasis, *LRP1* has also been implicated in Alzheimer's disease, for which metabolic

disease and obesity are risk factors (Kang *et al.* 2000; Anstey *et al.* 2011).

Curiously, several of the genes highlighted in this analysis also happen to localize specifically to

the mitochondria (N = 4). *Roe1* and *porin* are both transporters involved in the import of

molecules into the mitochondria (Komarov *et al.* 2004; Jana Alonso *et al.* 2005; FlyBase

Curators 2008; Gaudet *et al.* 2011). The NADH dehydrogenases *ND-PDSW* and *NDUFS3* both

function in the mitochondria as well as part of Complex I in the electron transport chain

(Jana Alonso *et al.* 2005). Closer examination of the top GWA candidates reveals additional

mitochondrial localization candidates including the amino acyl tRNA synthetase *GlyRS*, the

membrane bound regulator of protein kinase A (*pkaap*) and the translation elongation factor

mEFTu1 (Gaudet *et al.* 2011; Lu *et al.* 2016). In adult metabolic homeostasis, central carbon

metabolism is generally used to fuel the electron transport chain in the mitochondria and

generate ATP for the cell (Barry and Thummel 2016). Altering the activity of this essential

downstream pathway could have a clear impact on glucose utilization and disease progression

in diabetes.

Similar to our examination of top candidates, our physical interaction map highlighted the

immune response (*NimC1*, *NimC3*, *NimB2*, and *LRR*) and neuronal function. *Tutl*, *dpr2*, and

DIP-eta, and *wb* are all involved in synapse organization and axon guidance, while *Pax* and

rhea are involved in focal adhesion (Delon and Brown 2009). The identification of genes

important for cellular communication suggests that some of the modifiers identified in this study

have roles in tissues other than the fat body, such as the IPC and APC neurons. This is an

important avenue of future exploration.

Gene Set Enrichment Analysis (GSEA): In the second approach, we performed GSEA

analysis to identify gene ontology terms for which associated variants are enriched. Unlike

traditional GO analysis, which relies upon a set of genes based on a P-value cutoff, GSEA

examines the entire gene set (Dyer *et al.* 2008). For each defined GO category, GSEA

determines whether the members of that category are randomly distributed throughout the

ranked gene list provided or if they are enriched for the lower p-values found at the top of that

list. GO categories enriched at the top of the list describe important functions of the gene set.

GSEA identified 52 significantly associated gene sets (≥ 3 genes) with positive enrichment

scores at a p-value of <0.05 (Table S6, Figure 3B). The top two gene sets implicate neuronal

function and communication in the *Sirt1i*-associated hyperglycemia phenotype: calcium-

activated potassium channel activity (GO:0015269, P = 1.1E-3) and maintenance of presynaptic

active zone structure (GO:0048790, P = 1.2E-3). Similar categories can be found through the

list of significantly associated gene sets, including dendrite morphogenesis (GO:0048813, P =

0.049), which represents the largest group of genes at N = 119 and contains two of the top

GWA candidates (*slit* and *fruitless*). Coupled with the neuronal genes identified in our physical

interaction network, this suggests that function in the neuroendocrine cells could play a big role

in glucose homeostasis in the fat body.

Also enriched are taste receptor activity (GO:0008527, P = 0.013) and sensory reception of

taste (GO:0050909, P = 0.018). These categories highlight a yet unconsidered factor that must

be taken into account with metabolic phenotypes: that of feeding and diet. While all flies were

collected under identical conditions and were maintained on the same diet, it is nonetheless

possible that some may simply be eating more due to differences in the sensing of satiation or

to differences in perception of taste. These differences in perception and consumption can have

detectable impacts on metabolic phenotypes (May *et al.* 2019).

As expected, we also see evidence of general metabolic processes. Some, like alpha,alpha-

trehalase activity (GO:0004555, P = 0.043) and phosphatidylinositol transporter activity

(GO:0008526, P = 0.013) have direct links to glucose metabolism and insulin signaling. Others,

such as oxysterol binding (GO:0008142, P = 0.026), glutamate biosynthetic process

(GO:0006537, P = 0.043), and isoprenoid biosynthetic process (GO:0008299, P = 0.048)

function more peripherally to carbon metabolism and are likely influencing hyperglycemia by

their general contribution to physiological homeostasis.

Another interesting group of GO categories highlighted by GSEA are RNA processing functions.

rRNA (uridine-2'-O-)-methyltransferase activity (GO:0008650, P = 2.4E-3) is the fourth most

associated category as ranked by p-value, and others such as snoRNA binding (GO:0030515, P

= 0.014) reiterating this function. The presence of RNA processing categories is of particular

interest because three of the top candidate genes by GWA are splicing factors (*bru1*, *fand*, and

snRNP-U1-70k) (Park *et al.* 2004; Oas *et al.* 2014; Spletter *et al.* 2015). While it is unclear how

rRNA or mRNA processing may directly or indirectly influence glucose homeostasis in

particular, the identification of this process through several different methods of analysis is

striking and worth further exploration.

Functional analysis of candidate modifier genes

To confirm the roles of our candidate genes in regulating glucose homeostasis, we elected to

test the impact of loss of modifier expression for 16 of the most significant candidates for which

we were able to obtain transgenic RNAi lines (Table 1). We crossed the RNAi strains targeting

each of these modifiers into the *R4>Sirt1i* line, aged the resulting progeny for 2-3 weeks, and

measured glucose in fasted males. We also measured protein levels as a control. Knockdown of

modifier genes did not significantly alter protein levels as compared to a genetically matched

control (Figure S7). Knockdown of the genes *CG4168*, *CG5888*, and *uif* resulted in suppression

of the hyperglycemia phenotype, with a significant decrease in glucose content per fly compared

to controls expressing *R4>Sirt1i* (Figure 4). These results demonstrate that many of the top

GWA candidate modifiers are capable of modifying the hyperglycemia phenotypes associated

with the *R4>Sirt1i* model of diabetes.

550   **DISCUSSION**

Identifying and characterizing the genetic factors influencing the severity of diabetes is critical to

early diagnosis. Prevention is still the best strategy available, and providing patients at high risk

for complications with knowledge of that risk could prevent the worst symptoms from

manifesting. It could even enable intervention before the progression of disease is irreversible.

In this study, we identified and analyzed a number of candidate modifiers of hyperglycemia in a

previously characterized model of diabetes, *Sirt1* loss-of-function. We used the DGRP as an

unbiased source of natural genetic variation for this screen. This is the first time a genetic model

of metabolic dysfunction has been put to this use, as previous screens have either focused on

dietary stress as a source of metabolic disease or on the impact of genotype on metabolic

parameters under non-stressed conditions (Mackay *et al.* 2012; Ivanov *et al.* 2015; Nelson *et al.*

2016; Jehrke *et al.* 2018; Everman *et al.* 2019). We observed very little overlap in modifier

candidates between our observations and these studies. This is consistent with previous work

demonstrating that even when the observed phenotypes are similar or nearly identical, the

overlap in modifiers is often small in different models of disease (Palu *et al.* 2019). One

exception is a screen for the response to starvation resistance performed by Everman *et al.* in

2018. They identified *CG15803*, a transporter of unknown function that was also identified in our

studies (Everman and Morgan 2018). While our preliminary analysis suggests that this gene

does not function in the fat body, it is possible that it could have a role in another physiologically

relevant tissue such as the IPCs or APCs. Indeed, *CG15803* appears to be most highly

expressed in the head and CNS (Graveley *et al.* 2011). This is an interesting avenue of future

exploration.

We did observe overlap in general gene categories with previous studies, even when direct

overlaps were few. This was found to be true for modifiers of neuronal function. In a broad

exploration of genetic variation in the nutrient response that looked at triglycerides, starvation

resistance, mass, and glucose, *NimB3* was identified as a candidate modifier (Nelson *et al.*

2016). While *NimB3* was not identified in this screen, we did find *NimB2*, *NimC1*, and *NimC3*.

Fife, which also has roles in synapse organization, was identified in this study and one for

starvation resistance (Mackay *et al.* 2012). As mentioned above, both the IPCs and APCs are

neuroendocrine cells found in or near the brain (Géminard *et al.* 2006). Maintenance of neuronal

function would be critical to hormonal balance as a result. Furthermore, it is broadly

acknowledged that metabolic homeostasis is also dependent upon feeding rate, over which the

central nervous system has some sway (Owusu-Ansah and Perrimon 2014). The identification

of neuronal genes in each analysis suggests that regulation of particular neuronal pathways and

cells is critical to the maintenance of physiological homeostasis.

Given the prevalence of neuronal and sensory perception genes in the analysis, a concern

could be raised for the role feeding rate could be playing in the variation of glucose across the

DGRP. To assess this, we compared glucose with male feeding rate in a previous analysis

(Table S3) (Garlapow *et al.* 2015). We saw no correlation whatsoever, indicating that while

nutrient sensing may play a role in the response to Sirt1i-induced hyperglycemia, it is not the

driving factor in the variation observed in this screen (Figure S8).

Another interesting category highlighted through several analysis methods is the innate immune

response. While it has long been known that Type I diabetes is an autoimmune disorder, it has

recently been acknowledged that Type II diabetics also display symptoms of autoimmunity

(Candia *et al.* 2019). Furthermore, insulin resistance has frequently been associated with

inflammation, and the presence of macrophages in the adipose tissue is a hallmark of obesity

and diabetes (Wu and Ballantyne 2020). Modifiers associated with innate immunity serve

therefore as validation to the study as a whole, and examination of these genes and their

function in the context of the *Sirt1* loss-of-function model will be an intriguing focus of future

research.

An important component of this study is the validation of top candidate modifiers using RNAi-

mediated knockdown of gene expression. We obtained strains expression transgenic RNAi

constructs targeting 16 of the most significant candidates (Table 1). We found that reduced

expression of three candidates specifically in the fat body resulted in significant suppression of

hyperglycemia: *CG4168*, *CG5888*, and *uif* (Figure 4). We also noted a consistent, though not

significant, decrease in glucose for two independent RNAi constructs targeting *CTPSyn*,

suggesting that this gene warrants further study (Figure 4, data not shown). The remainder of

the genes had no significant or consistent impact on hyperglycemia in the model of *Sirt1* loss-of-

function (Figure 4). There were no enhancing modifiers: loss of modifier expression did not lead

to increased glucose levels for any of the tested genes. This could be because hyperglycemia is

already quite strong in the *Sirt1* loss-of-function model, or because we simply did not hit on any

enhancing modifiers. Of greater concern is the lack of any response for 13 of the 16 tested

candidates. One explanation may be found in the large number of known neuronal genes

identified in this analysis. This modifier RNAi screen specifically focused on the expression of

the RNAi against the candidate genes in the fat body, where expression of *Sirt1* is also reduced.

If, however, the function of a modifier gene is primarily concentrated in the IPCs, as with *ilp5*,

reducing its expression in the fat body would have little effect on the disease phenotypes in

question. We will examine the role of modifier genes not only in the fat body but in the IPCs,

APCs, and other physiologically relevant tissues in future work.

Of immediate interest is of course the mechanism of action for the three suppressor genes that

were confirmed by RNAi in the fat body. Of these, *CG5888* is also the top GWA candidate (P =

2.80E-07). While largely uncharacterized, *CG5888* has been identified as a component or

activator of voltage-gated potassium channels by sequence homology (Gaudet *et al.* 2011).

Intriguingly, it has also been implicated JNK signaling by a previous RNAi screen (Bond and

Foley 2009). JNK signaling is commonly activated by cellular stress and can activate apoptosis.

It also has important roles in the signaling pathways commonly used by the immune system

(Bond and Foley 2009; Shlevkov and Morata 2012). A potential role for *CG5888* in immune

pathways has yet to be explored and could be an exciting area of further discovery.

Uninflatable (*uif*) encodes a single pass transmembrane protein found on the apical membrane

of epithelial cells and has been found to enable Notch signaling (Loubéry *et al.* 2014). It has

also been found to exacerbate disease in a *Drosophila* model of muscular dystrophy

(Kucherenko *et al.* 2011), and its closest human orthologue ELAPOR1 is a regulator of

apoptosis and autophagy (Deng *et al.* 2010). Both of these processes are commonly disrupted

through inappropriate activation in metabolic disease, and might provide some explanation for

the impact of *uif* on *Sirt1i*-associated phenotypes (Bugliani *et al.* 2019).

Perhaps the most intriguing finding is *CG4168* as the modifier with the strongest impact on

Sirt1i-associated hyperglycemia. The protein encoded by *CG4168* is of unknown function, but

its closest human orthologue (*IGFALS*) encodes a serum protein that binds to insulin-like growth

factors (IGF) in circulation (Boisclair *et al.* 1996). In mammals, association with IGFALS

increases the half-life of insulin-like growth factors in the serum as well as their retention in

circulation. While studies of IGFALS in mammals has not shown a role for it in regulating insulin

signaling, the *Drosophila* ilp peptides are used for both IGF and insulin signaling activation

(Géminard *et al.* 2006). It is possible that secretion of the factor encoded by *CG4168* from the

fat body could increase ilp retention in circulation, whereas its loss could result in faster

clearance of ilps from circulation. Under conditions that promote insulin resistance, such as the

loss of *Sirt1*, it is possible that reduced ilp levels in the hemolymph could slow or prevent insulin

resistance and hyperglycemia. Further exploration of the mechanisms behind the action of

CG4168 could reveal important insights into circulating insulin-binding factors and their role in

diabetes.

In conclusion, we have identified a number of pathways and processes involved in the degree of

hyperglycemia in a genetic model of diabetes. Examination of the candidate genes and

pathways described above in this model as well as other models of metabolic dysfunction will

shed new light on the mechanism by which insulin resistance and related complications disease

onset, progression, and severity. Furthermore, the candidates identified as suppressors could

serve as promising targets for therapeutics in diabetes and related metabolic disorders.

665   **DATA AVAILABILITY STATEMENT**

666   Strains and stocks are available upon request, as is code for GSEA. Genomic sequence for the

667   DGRP is available at http://dgrp.gnets.ncsu.edu/. Supplemental material is available at FigShare

668   (https://figshare.com/s/2b0b0237de0f94139a4f).

670   **ACKNOWLEDGEMENTS**

671   We thank Katie G. Owings and Clement Y. Chow for their contributions to the data analysis in

672   this project, as well as reagents and comments. This research was supported by a Faculty

673   Summer Research Grant through the Purdue University-Fort Wayne Office of Sponsored

674   Projects to RASP as well as the Purdue University-Fort Wayne Department of Biology.

REFERENCES

Al-Anzi B., and R. J. Wyman, 2009 The Drosophila immunoglobulin gene turtle encodes

guidance molecules involved in axon pathfinding. Neural Dev. 2009 41 4: 1–15.

https://doi.org/10.1186/1749-8104-4-31

Anh N. T. T., M. Nishitani, S. Harada, M. Yamaguchi, and K. Kamei, 2011 Essential Role of

Duox in Stabilization of Drosophila Wing. J. Biol. Chem. 286: 33244–33251.

https://doi.org/10.1074/JBC.M111.263178

Anstey K. J., N. Cherbuin, M. Budge, and J. Young, 2011 Body mass index in midlife and late-

life as a risk factor for dementia: a meta-analysis of prospective studies. Obes. Rev. 12:

e426–e437. https://doi.org/10.1111/J.1467-789X.2010.00825.X

Arrese E. L., and J. L. Soulages, 2010 Insect Fat Body: Energy, Metabolism, and Regulation.

Annu. Rev. Entomol. 55: 207–225. https://doi.org/10.1146/annurev-ento-112408-085356

Barroso I., and M. I. McCarthy, 2019 The Genetic Basis of Metabolic Disease. Cell 177: 146–

161. https://doi.org/10.1016/j.cell.2019.02.024

Barry W. E., and C. S. Thummel, 2016 The Drosophila HNF4 nuclear receptor promotes

glucose-stimulated insulin secretion and mitochondrial function in adults. Elife 5.

https://doi.org/10.7554/ELIFE.11183

Beebe K., M. M. Robins, E. J. Hernandez, G. Lam, M. A. Horner, *et al.*, 2020 Drosophila

estrogen-related receptor directs a transcriptional switch that supports adult glycolysis and

lipogenesis. Genes Dev. 34: 701. https://doi.org/10.1101/GAD.335281.119

Biason-Lauber A., M. Böni-Schnetzler, B. P. Hubbard, K. Bouzakri, A. Brunner, *et al.*, 2013

Identification of a SIRT1 Mutation in a Family with Type 1 Diabetes. Cell Metab. 17: 448–

455. https://doi.org/10.1016/J.CMET.2013.02.001

Boisclair Y. R., D. Seto, S. Hsieh, K. R. Hurst, and G. T. Ooi, 1996 Organization and

chromosomal localization of the gene encoding the mouse acid labile subunit of the insulin-

like growth factor binding complex. Proc. Natl. Acad. Sci. 93: 10028–10033.

https://doi.org/10.1073/PNAS.93.19.10028

Bond D., and E. Foley, 2009 A quantitative RNAi screen for JNK modifiers identifies Pvr as a

novel regulator of Drosophila immune signaling. PLoS Pathog. 5.

https://doi.org/10.1371/JOURNAL.PPAT.1000655

Botden I. P. G., M. C. Zillikens, S. R. de Rooij, J. G. Langendonk, A. H. J. Danser, et al., 2012

Variants in the SIRT1 Gene May Affect Diabetes Risk in Interaction With Prenatal

Exposure to Famine. Diabetes Care 35: 424–426. https://doi.org/10.2337/DC11-1203

Boutant M., and C. Cantó, 2014 SIRT1 metabolic actions: Integrating recent advances from

mouse models. Mol. Metab. 3: 5–18.

Bruckner J. J., H. Zhan, S. J. Gratz, M. Rao, F. Ukken, et al., 2017 Fife organizes synaptic

vesicles and calcium channels for high-probability neurotransmitter release. J. Cell Biol.

216: 231–246. https://doi.org/10.1083/JCB.201601098

Brunet A., L. Sweeney, J. Sturgill, K. Chua, P. Greer, et al., 2004 Stress-dependent regulation

of FOXO transcription factors by the SIRT1 deacetylase. Science 303: 2011–2015.

https://doi.org/10.1126/SCIENCE.1094637

Bugliani M., S. Mossuto, F. Grano, M. Suleiman, L. Marselli, et al., 2019 Modulation of

Autophagy Influences the Function and Survival of Human Pancreatic Beta Cells Under

Endoplasmic Reticulum Stress Conditions and in Type 2 Diabetes. Front. Endocrinol.

(Lausanne). 10: 52. https://doi.org/10.3389/fendo.2019.00052

Cabrera A. P., R. N. Mankad, L. Marek, R. Das, S. Rangasamy, et al., 2020 Genotypes and

Phenotypes: A Search for Influential Genes in Diabetic Retinopathy. Int. J. Mol. Sci. 2020,

Vol. 21, Page 2712 21: 2712. https://doi.org/10.3390/IJMS21082712

Candia P. de, F. Prattichizzo, S. Garavelli, V. De Rosa, M. Galgani, et al., 2019 Type 2

Diabetes: How Much of an Autoimmune Disease? Front. Endocrinol. (Lausanne). 10: 451.

https://doi.org/10.3389/FENDO.2019.00451

Carrillo R. A., E. Özkan, K. P. Menon, S. Nagarkar-Jaiswal, P.-T. Lee, *et al.*, 2015 Control of

Synaptic Connectivity by a Network of Drosophila IgSF Cell Surface Proteins. Cell 163:

1770–1782. https://doi.org/10.1016/J.CELL.2015.11.022

CDC, 2020 National Diabetes Statistics Report 2020. Estimates of diabetes and its burden in

the United States.

Chang Y.-H., and Y. H. Sun, 2014 Carrier of Wingless (Cow), a Secreted Heparan Sulfate

Proteoglycan, Promotes Extracellular Transport of Wingless. PLoS One 9: e111573.

https://doi.org/10.1371/JOURNAL.PONE.0111573

Chow C. Y., M. F. Wolfner, and A. G. Clark, 2013 Using natural variation in Drosophila to

discover previously unknown endoplasmic reticulum stress genes. Proc Natl Acad Sci U S

A 110: 9013–9018. https://doi.org/10.1073/pnas.1307125110

Chow C. Y., F. W. Avila, A. G. Clark, and M. F. Wolfner, 2015 Induction of excessive

Endoplasmic reticulum stress in the Drosophila male accessory gland results in infertility.

PLoS One 10: 1–12. https://doi.org/10.1371/journal.pone.0119386

Chow C. Y., 2016 Bringing genetic background into focus. Nat. Rev. Genet. 17: 63–64.

https://doi.org/10.1038/nrg.2015.9

Chow C. Y., K. J. P. Kelsey, M. F. Wolfner, and A. G. Clark, 2016 Candidate genetic modifiers

of retinitis pigmentosa identified by exploiting natural variation in Drosophila. Hum. Mol.

Genet. 25: 651–659. https://doi.org/10.1093/HMG/DDV502

Chow C. Y., and L. T. Reiter, 2017 Etiology of Human Genetic Disease on the Fly. Trends

Genet. 33: 391–398. https://doi.org/10.1016/j.tig.2017.03.007

Delon I., and N. H. Brown, 2009 The integrin adhesion complex changes its composition and

function during morphogenesis of an epithelium. J. Cell Sci. 122: 4363–4374.

https://doi.org/10.1242/JCS.055996

Deng L., J. Feng, and R. R. Broaddus, 2010 The novel estrogen-induced gene EIG121

regulates autophagy and promotes cell survival under stress. Cell Death Dis. 1: e32.

https://doi.org/10.1038/CDDIS.2010.9

DiAngelo J. R., and M. J. Birnbaum, 2009 Regulation of Fat Cell Mass by Insulin in Drosophila

melanogaster. Mol. Cell. Biol. 29: 6341–6352. https://doi.org/10.1128/mcb.00675-09

Dyer M. D., T. M. Murali, and B. W. Sobral, 2008 The Landscape of Human Proteins Interacting

with Viruses and Other Pathogens. PLoS Pathog. 4: e32.

https://doi.org/10.1371/journal.ppat.0040032

Everman E. R., and T. J. Morgan, 2018 Antagonistic pleiotropy and mutation accumulation

contribute to age-related decline in stress response. Evolution (N. Y). 72: 303–317.

https://doi.org/10.1111/EVO.13408

Everman E. R., C. L. McNeil, J. L. Hackett, C. L. Bain, and S. J. Macdonald, 2019 Dissection of

Complex, Fitness-Related Traits in Multiple Drosophila Mapping Populations Offers Insight

into the Genetic Control of Stress Resistance. Genetics 211: 1449–1467.

https://doi.org/10.1534/genetics.119.301930

Filmus J., and S. B. Selleck, 2001 Glypicans: proteoglycans with a surprise. J. Clin. Invest. 108:

497–501. https://doi.org/10.1172/JCI13712

Flegal K., D. Kruszon-Moran, M. Carroll, C. Fryar, and C. Ogden, 2016 Trends in Obesity

Among Adults in the United States, 2005 to 2014. JAMA 315: 2284–2291.

https://doi.org/10.1001/JAMA.2016.6458

FlyBase Curators, Swiss-Prot Project Members, and InterPro Project Members, 2004 Gene

Ontology annotation in FlyBase through association of InterPro records with GO terms.

FlyBase Curators, 2008 Assigning Gene Ontology (GO) terms by sequence similarity in

FlyBase.

Garlapow M. E., W. Huang, M. T. Yarboro, K. R. Peterson, and T. F. C. Mackay, 2015

Quantitative Genetics of Food Intake in Drosophila melanogaster. PLoS One 10:

777   e0138129. https://doi.org/10.1371/JOURNAL.PONE.0138129

778 Gaudet P., M. S. Livstone, S. E. Lewis, and P. D. Thomas, 2011 Phylogenetic-based

779   propagation of functional annotations within the Gene Ontology consortium. Brief.

780   Bioinform. 12: 449–462. https://doi.org/10.1093/bib/bbr042

781 Géminard C., N. Arquier, S. Layalle, M. Bourouis, M. Slaidina, *et al.*, 2006 Control of

782   metabolism and growth through insulin-like peptides in Drosophila. Diabetes 55: S5–S8.

783   https://doi.org/10.2337/db06-S001

784 Géminard C., E. J. Rulifson, and P. Léopold, 2009 Remote Control of Insulin Secretion by Fat

785   Cells in Drosophila. Cell Metab. 10: 199–207. https://doi.org/10.1016/j.cmet.2009.08.002

786 Graveley B. R., G. May, A. N. Brooks, J. W. Carlson, L. Cherbas, *et al.*, 2011 The D.

787   melanogaster transcriptome: modENCODE RNA-Seq data for dissected tissues

788 Hahn K., M. Miranda, V. A. Francis, J. Vendrell, A. Zorzano, *et al.*, 2010 PP2A Regulatory

789   Subunit PP2A-B′ Counteracts S6K Phosphorylation. Cell Metab. 11: 438–444.

790   https://doi.org/10.1016/J.CMET.2010.03.015

791 He B. Z., M. Z. Ludwig, D. A. Dickerson, L. Barse, B. Arun, *et al.*, 2014 Effect of Genetic

792   Variation in a Drosophila Model of Diabetes-Associated Misfolded Human Proinsulin.

793   Genetics 196: 557. https://doi.org/10.1534/GENETICS.113.157800

794 Huang W., A. Massouras, Y. Inoue, J. Peiffer, M. Ràmia, *et al.*, 2014 Natural variation in

795   genome architecture among 205 Drosophila melanogaster Genetic Reference Panel lines.

796   Genome Res. 24: 1193–208. https://doi.org/10.1101/gr.171546.113

797 Ida T., T. Takahashi, H. Tominaga, T. Sato, K. Kume, *et al.*, 2011 Identification of the novel

798   bioactive peptides dRYamide-1 and dRYamide-2, ligands for a neuropeptide Y-like

799   receptor in Drosophila. Biochem. Biophys. Res. Commun. 410: 872–877.

800   https://doi.org/10.1016/J.BBRC.2011.06.081

801 Irvin A. E., G. Jhala, Y. Zhao, T. S. Blackwell, B. Krishnamurthy, *et al.*, 2018 NF-κB is weakly

802   activated in the NOD mouse model of type 1 diabetes. Sci. Reports 2018 81 8: 1–7.

https://doi.org/10.1038/s41598-018-22738-3

Ivanov D. K., V. Escott-Price, M. Ziehm, M. M. Magwire, T. F. C. Mackay, *et al.*, 2015 Longevity

GWAS using the drosophila genetic reference panel. Journals Gerontol. - Ser. A Biol. Sci.

Med. Sci. 70: 1470–1478. https://doi.org/10.1093/gerona/glv047

Ivy J., A. Klar, and J. Hicks, 1986 Cloning and characterization of four SIR genes of

Saccharomyces cerevisiae. Mol. Cell. Biol. 6: 688–702.

https://doi.org/10.1128/MCB.6.2.688-702.1986

Jana Alonso, Javier M. Rodriguez, and Luis Alberto Baena-López, and J. F. Santarén*, 2005

Characterization of the Drosophila melanogaster Mitochondrial Proteome. J. Proteome

Res. 4: 1636–1645. https://doi.org/10.1021/PR050130C

Jang B.-Y., H. D. Ryoo, J. Son, K.-C. Choi, D.-M. Shin, *et al.*, 2015 Role of Drosophila EDEMs

in the degradation of the alpha-1-antitrypsin Z variant. Int. J. Mol. Med. 35: 870–876.

https://doi.org/10.3892/IJMM.2015.2109

Jehrke L., F. A. Stewart, A. Droste, and M. Beller, 2018 The impact of genome variation and diet

on the metabolic phenotype and microbiome composition of Drosophila melanogaster. Sci.

Rep. 8. https://doi.org/10.1038/S41598-018-24542-5

Kang D. E., C. U. Pietrzik, L. Baum, N. Chevallier, D. E. Merriam, *et al.*, 2000 Modulation of

amyloid β-protein clearance and Alzheimer's disease susceptibility by the LDL receptor–

related protein pathway. J. Clin. Invest. 106: 1159–1166. https://doi.org/10.1172/JCI11013

Kang M.-J., and H. D. Ryoo, 2009 Suppression of retinal degeneration in Drosophila by

stimulation of ER-associated degradation. Proc. Natl. Acad. Sci. U. S. A. 106: 17043–8.

https://doi.org/10.1073/pnas.0905566106

Kim J., H. Bang, S. Ko, I. Jung, H. Hong, *et al.*, 2008 Drosophila ia2 modulates secretion of

insulin-like peptide. Comp. Biochem. Physiol. Part A Mol. Integr. Physiol. 151: 180–184.

https://doi.org/10.1016/J.CBPA.2008.06.020

Komarov A. G., B. H. Graham, W. J. Craigen, and M. Colombini, 2004 The Physiological

Properties of a Novel Family of VDAC-Like Proteins from Drosophila melanogaster.

Biophys. J. 86: 152–162. https://doi.org/10.1016/S0006-3495(04)74093-X

Kucherenko M. M., A. K. Marrone, V. M. Rishko, H. de F. Magliarelli, and H. R. Shcherbata,

2011 Stress and muscular dystrophy: A genetic screen for Dystroglycan and Dystrophin

interactors in Drosophila identifies cellular stress response components. Dev. Biol. 352:

228–242. https://doi.org/10.1016/J.YDBIO.2011.01.013

Kurusu M., A. Cording, M. Taniguchi, K. Menon, E. Suzuki, *et al.*, 2008 A Screen of Cell-Surface

Molecules Identifies Leucine-Rich Repeat Proteins as Key Mediators of Synaptic Target

Selection. Neuron 59: 972–985. https://doi.org/10.1016/J.NEURON.2008.07.037

Landis G. N., D. Bhole, and J. Tower, 2003 A search for doxycycline-dependent mutations that

increase Drosophila melanogaster life span identifies the VhaSFD, Sugar baby, filamin, fwd

and Cctlgenes. Genome Biol. 2003 42 4: 1–14. https://doi.org/10.1186/GB-2003-4-2-R8

Lavoy S., V. G. Chittoor-Vinod, C. Y. Chow, and I. Martin, 2018 Genetic Modifiers of

Neurodegeneration in a Drosophila Model of Parkinson's Disease. Genetics 209: 1345–

1356. https://doi.org/10.1534/genetics.118.301119

Lee G., and J. H. Park, 2004 Hemolymph Sugar Homeostasis and Starvation-Induced

Hyperactivity Affected by Genetic Manipulations of the Adipokinetic Hormone-Encoding

Gene in Drosophila melanogaster. Genetics 167: 311–323.

https://doi.org/10.1534/GENETICS.167.1.311

Li X., S. Zhang, G. Blander, J. Tse, M. Krieger, *et al.*, 2007 SIRT1 deacetylates and positively

regulates the nuclear receptor LXR. Mol. Cell 28: 91–106.

https://doi.org/10.1016/J.MOLCEL.2007.07.032

Lizák B., A. Szarka, Y. Kim, K. Choi, C. E. Németh, *et al.*, 2019 Glucose Transport and

Transporters in the Endomembranes. Int. J. Mol. Sci. 20.

https://doi.org/10.3390/IJMS20235898

Loubéry S., C. Seum, A. Moraleda, A. Daeden, M. Fürthauer, *et al.*, 2014 Uninflatable and

Notch Control the Targeting of Sara Endosomes during Asymmetric Division. Curr. Biol. 24:

2142–2148. https://doi.org/10.1016/J.CUB.2014.07.054

Lu J., S. J. Marygold, W. H. Gharib, and B. Suter, 2016 The aminoacyl-tRNA synthetases of

Drosophila melanogaster. https://doi.org/10.1080/19336934.2015.1101196 9: 53–61.

https://doi.org/10.1080/19336934.2015.1101196

Luo C.-W., E. M. Dewey, S. Sudo, J. Ewer, S. Y. Hsu, *et al.*, 2005 Bursicon, the insect cuticle-

hardening hormone, is a heterodimeric cystine knot protein that activates G protein-

coupled receptor LGR2. Proc. Natl. Acad. Sci. 102: 2820–2825.

https://doi.org/10.1073/PNAS.0409916102

Mackay T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, *et al.*, 2012 The

Drosophila melanogaster Genetic Reference Panel. Nature 482: 173–178.

https://doi.org/10.1038/nature10811

May C. E., A. Vaziri, Y. Q. Lin, O. Grushko, M. Khabiri, *et al.*, 2019 High Dietary Sugar

Reshapes Sweet Taste to Promote Feeding Behavior in Drosophila melanogaster. Cell

Rep. 27: 1675-1685.e7. https://doi.org/10.1016/J.CELREP.2019.04.027

McMullen E., A. Weiler, H. M. Becker, and S. Schirmeier, 2021 Plasticity of Carbohydrate

Transport at the Blood-Brain Barrier. Front. Behav. Neurosci. 0: 271.

https://doi.org/10.3389/FNBEH.2020.612430

Millington J. W., and E. J. Rideout, 2018 Sex differences in Drosophila development and

physiology. Curr. Opin. Physiol. 6: 46–56. https://doi.org/10.1016/J.COPHYS.2018.04.002

Montojo J., K. Zuberi, H. Rodriguez, F. Kazi, G. Wright, *et al.*, 2010 GeneMANIA Cytoscape

plugin: fast gene function predictions on the desktop. Bioinformatics 26: 2927–8.

https://doi.org/10.1093/bioinformatics/btq562

Nelson C. S., J. N. Beck, K. A. Wilson, E. R. Pilcher, P. Kapahi, *et al.*, 2016 Cross-phenotype

association tests uncover genes mediating nutrient response in Drosophila. BMC

Genomics 17. https://doi.org/10.1186/s12864-016-3137-9

Nogueiras R., K. M. Habegger, N. Chaudhary, B. Finan, A. S. Banks, *et al.*, 2012 Sirtuin 1 and

sirtuin 3: Physiological modulators of metabolism. Physiol. Rev. 92: 1479–1514.

https://doi.org/10.1152/physrev.00022.2011

Oas S. T., A. L. Bryantsev, and R. M. Cripps, 2014 Arrest is a regulator of fiber-specific

alternative splicing in the indirect flight muscles of Drosophila. J. Cell Biol. 206: 895–908.

https://doi.org/10.1083/JCB.201405058

Owusu-Ansah E., and N. Perrimon, 2014 Modeling metabolic homeostasis and nutrient sensing

in Drosophila: implications for aging and metabolic diseases. Dis. Model. Mech. 7: 343–

350. https://doi.org/10.1242/DMM.012989

Paik D., Y. G. Jang, Y. E. Lee, Y. N. Lee, R. Yamamoto, *et al.*, 2012 Misexpression screen

delineates novel genes controlling Drosophila lifespan. Mech. Ageing Dev. 133: 234–245.

https://doi.org/10.1016/J.MAD.2012.02.001

Palu R. A. S., and C. S. Thummel, 2016 Sir2 Acts through Hepatocyte Nuclear Factor 4 to

maintain insulin Signaling and Metabolic Homeostasis in Drosophila. PLoS Genet. 12: 1–

18. https://doi.org/10.1371/journal.pgen.1005978

Palu R. A. S., and C. Y. Chow, 2018 Baldspot/ELOVL6 is a conserved modifier of disease and

the ER stress response. PLoS Genet. 14. https://doi.org/10.1101/261206

Palu R. A. S., E. Ong, K. Stevens, S. Chung, K. G. Owings, *et al.*, 2019 Natural Genetic

Variation Screen in Drosophila Identifies Wnt Signaling, Mitochondrial Metabolism, and

Redox Homeostasis Genes as Modifiers of Apoptosis. G3 Genes|Genomes|Genetics 9:

g3.400722.2019. https://doi.org/10.1534/g3.119.400722

Palu R. A. S., H. M. Dalton, and C. Y. Chow, 2020 Decoupling of Apoptosis from Activation of

the ER Stress Response by the Drosophila Metallopeptidase superdeath. Genetics 214:

913–925. https://doi.org/10.1534/GENETICS.119.303004

Park J. W., K. Parisky, A. M. Celotto, R. A. Reenan, and B. R. Graveley, 2004 Identification of

alternative splicing regulators by RNA interference in Drosophila. Proc. Natl. Acad. Sci.

101: 15974–15979. https://doi.org/10.1073/PNAS.0407004101

Picard F., M. Kurtev, N. Chung, A. Topark-Ngarm, T. Senawong, *et al.*, 2004 Sirt1 promotes fat

mobilization in white adipocytes by repressing PPAR-gamma. Nature 429: 771–776.

https://doi.org/10.1038/NATURE02583

Prakash S., J. C. Caldwell, D. F. Eberl, and T. R. Clandinin, 2005 Drosophila N-cadherin

mediates an attractive interaction between photoreceptor axons and their targets. Nat.

Neurosci. 2005 84 8: 443–450. https://doi.org/10.1038/nn1415

Prisco G. Di, V. Cavaliere, D. Annoscia, P. Varricchio, E. Caprio, *et al.*, 2013 Neonicotinoid

clothianidin adversely affects insect immunity and promotes replication of a viral pathogen

in honey bees. Proc. Natl. Acad. Sci. 110: 18466–18471.

https://doi.org/10.1073/PNAS.1314923110

Queitsch C., K. D. Carlson, and S. Girirajan, 2012 Lessons from model organisms: phenotypic

robustness and missing heritability in complex disease. PLoS Genet 8: e1003041.

https://doi.org/10.1371/journal.pgen.1003041

Rabelink T. J., B. M. van den Berg, M. Garsen, G. Wang, M. Elkin, *et al.*, 2017 Heparanase:

roles in cell survival, extracellular matrix remodelling and the development of kidney

disease. Nat. Rev. Nephrol. 2017 134 13: 201–212. https://doi.org/10.1038/nrneph.2017.6

Rao Y., P. Pang, W. Ruan, D. Gunning, and S. L. Zipursky, 2000 brakeless is required for

photoreceptor growth-cone targeting in Drosophila. Proc. Natl. Acad. Sci. 97: 5966–5971.

https://doi.org/10.1073/PNAS.110135297

Reis T., M. R. Van Gilst, and I. K. Hariharan, 2010 A Buoyancy-Based Screen of Drosophila

Larvae for Fat-Storage Mutants Reveals a Role for Sir2 in Coupling Fat Storage to Nutrient

Availability, (E. Rulifson, Ed.). PLoS Genet. 6: e1001206.

https://doi.org/10.1371/journal.pgen.1001206

Rine J., and I. Herskowitz, 1987 Four Genes Responsible for a Position Effect on Expression

from HML and HMR in Saccharomyces cerevisiae. Genetics 116: 9.

Rodgers J. T., and P. Puigserver, 2007 Fasting-dependent glucose and lipid metabolic

response through hepatic sirtuin 1. Proc. Natl. Acad. Sci. U. S. A. 104: 12861–12866.

https://doi.org/10.1073/pnas.0702509104

Shannon P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, *et al.*, 2003 Cytoscape: a software

environment for integrated models of biomolecular interaction networks. Genome Res. 13:

2498–504. https://doi.org/10.1101/gr.1239303

Shlevkov E., and G. Morata, 2012 A dp53/JNK-dependant feedback amplification loop is

essential for the apoptotic response to stress in Drosophila. Cell Death Differ. 19: 451–460.

https://doi.org/10.1038/cdd.2011.113

Shore D., M. Squire, and K. A. Nasmyth, 1984 Characterization of two genes required for the

position-effect control of yeast mating-type genes. EMBO J. 3: 2817.

Sieber M. H., and C. S. Thummel, 2009 The DHR96 Nuclear Receptor Controls Triacylglycerol

Homeostasis in Drosophila. Cell Metab. 10: 481–490.

https://doi.org/10.1016/J.CMET.2009.10.010

Sivachenko A., H. B. Gordon, S. S. Kimball, E. J. Gavin, J. L. Bonkowsky, *et al.*, 2016

Neurodegeneration in a Drosophila model of adrenoleukodystrophy: the roles of the

Bubblegum and Double bubble acyl-CoA synthetases. Dis. Model. Mech. 9: 377–387.

https://doi.org/10.1242/DMM.022244

Somogyi K., B. Sipos, Z. Pénzes, and I. Andó, 2010 A conserved gene cluster as a putative

functional unit in insect innate immunity. FEBS Lett. 584: 4375–4378.

https://doi.org/10.1016/J.FEBSLET.2010.10.014

Spletter M. L., C. Barz, A. Yeroslaviz, C. Schönbauer, I. R. S. Ferreira, *et al.*, 2015 The RNA-

binding protein Arrest (Bruno) regulates alternative splicing to enable myofibril maturation

in Drosophila flight muscle. EMBO Rep. 16: 178–191.

https://doi.org/10.15252/EMBR.201439791

Subramanian A., P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, *et al.*, 2005 Gene set

enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. Proc. Natl. Acad. Sci. U. S. A. 102: 15545–50.

https://doi.org/10.1073/pnas.0506580102

Talsness D. M., K. G. Owings, E. Coelho, G. Mercenne, J. M. Pleinis, *et al.*, 2020 A drosophila

screen identifies nkcc1 as a modifier of ngly1 deficiency. Elife 9: 1–22.

https://doi.org/10.7554/ELIFE.57831

Tennessen J. M., W. E. Barry, J. Cox, and C. S. Thummel, 2014 Methods for studying

metabolism in Drosophila. Methods 68: 105–115.

https://doi.org/10.1016/j.ymeth.2014.02.034

Udler M. S., M. I. McCarthy, J. C. Florez, and A. Mahajan, 2019 Genetic Risk Scores for

Diabetes Diagnosis and Precision Medicine. Endocr. Rev. 40: 1500–1520.

https://doi.org/10.1210/er.2019-00088

Wu H., and C. M. Ballantyne, 2020 Metabolic Inflammation and Insulin Resistance in Obesity.

Circ. Res. 126: 1549. https://doi.org/10.1161/CIRCRESAHA.119.315896

Yang J., X. Kong, M. Martins-Santos, G. Aleman, E. Chaco, *et al.*, 2009 Activation of SIRT1 by

resveratrol represses transcription of the gene for the cytosolic form of

phosphoenolpyruvate carboxykinase (GTP) by deacetylating hepatic nuclear factor 4alpha.

J. Biol. Chem. 284: 27042–27053. https://doi.org/10.1074/JBC.M109.047340

Zhao Y., J. Wei, X. Hou, H. Liu, F. Guo, *et al.*, 2017 SIRT1 rs10823108 and FOXO1

rs17446614 responsible for genetic susceptibility to diabetic nephropathy. Sci. Reports

2017 71 7: 1–9. https://doi.org/10.1038/s41598-017-10612-7

Zhou X., and M. Stephens, 2012 Genome-wide efficient mixed-model analysis for association

studies. Nat. Genet. 44: 821–4. https://doi.org/10.1038/ng.2310

Zhou X., C. J. Guo, H. H. Hu, J. Zhong, Q. Sun, *et al.*, 2019 Drosophila CTP synthase can form

distinct substrate- and product-bound filaments. J. Genet. Genomics 46: 537–545.

https://doi.org/10.1016/J.JGG.2019.11.006

Zillikens M. C., J. B. J. van Meurs, F. Rivadeneira, N. Amin, A. Hofman, *et al.*, 2009 SIRT1

Genetic Variation Is Related to BMI and Risk of Obesity. Diabetes 58: 2828–2834.

https://doi.org/10.2337/DB09-0536

989   **FIGURE LEGENDS**

Figure 1. Glucose levels vary under a variety of environmental conditions.

Glucose levels were measured in three samples for each of 30-36 strains under one of the

indicated conditions: one week of adult age and fed ad libitum (N = 30 strains, P = 1.87E-05)

(**A**), one week of adult age and fasted for 12-13 hours (N = 30 strains, P = 0.0194) (**B**), two

weeks of adult age and fed ad libitum (N = 30 strains, P = 0.0103) (**C**), and two weeks of adult

age and fasted for 12-13 hours (N = 36 strains, P = 1.95E-05) (**D**). Mean glucose concentrations

are indicated, with error bars indicating standard deviation. DGRP strain or RAL numbers are

indicated along the X-axis. P-values were calculated using one-way ANOVA incorporating all

individual measurements comparing DGRP strain with glucose concentration. Adult flies were

collected within 2-3 days after eclosion from the pupal case and aged to the indicated time

points. * P < 0.05, **** P < 5E-05.

1001

Figure 2. Glucose levels are significantly affected by genetic background.

Glucose levels were measured in three samples for each of 185 strains at two weeks of age

after a 12 hour fast. Adult flies were collected within 2-3 days after eclosion from the pupal case

and aged an additional 9-11 days prior to fasting (11-14 days post-eclosion). Flies were

collected after the overnight fast at 12-15 days post-eclosion. Mean glucose concentrations are

indicated, with error bars indicating standard deviation. P-values were calculated using one-way

ANOVA incorporating all individual measurements comparing DGRP strain with glucose

concentration (P < 2E-16).

1010

Figure 3. Immune responses, neuronal function, and basic metabolic processes are

overrepresented in GWA candidate modifiers of hyperglycemia.

(**A**) *R4>Sirt1i* modifier network, as plotted by the GeneMANIA plugin in Cytoscape (Shannon *et*

al. 2003; Montojo *et al.* 2010). Significant candidate modifiers are indicated in red, with physical

interactions indicated by connecting red lines. Thicker lines indicate stronger evidence for the

interaction. Encircled genes share common pathways or functions. Interacting genes outside of

the candidate modifier list are indicated in gray. (**B**) Top 20 significant ontological categories as

identified by GSEA. Categories are arranged from most significant on top to least significant

along the y-axis. P-values are indicated by red-to-blue gradient, with red the lowest p-values

and blue the highest P-values. Enrichment score (ES) for each category is plotted along the x-

axis. Gene number identified in each category is indicated by dot size.

1022

Figure 4. Loss of candidate gene expression suppresses hyperglycemia in the R4>Sirt1i

model

RNAi against candidate modifiers was expressed under the control of *R4-GAL4* in the *R4>Sirt1i*

model. Glucose level for each sample was normalized to the levels in a genetically matched

control line crossed into the *R4-Sirt1* line. Average *R4>Sirt1i* control glucose levels after

normalization are indicated by a dotted line at 1.0, with standard deviation highlighted by the gray

box. Whole fly glucose concentration was quantified for N = 4-5 samples per strain, each

consisting of 5 flies and individually plotted along the y-axis. Knockdown of *CG4168, CG5888*, or

uif significantly reduces glucose concentrations in the *R4>Sirt1i* model of hyperglycemia

compared to controls (blue). Loss of *CTPSyn* does not significantly alter glucose levels, but a

trending decrease in glucose levels were observed in several independent RNAi strains (light

blue, data not shown). Loss of *smt3*, *ilp5*, *snRNP-U1-70k*, *CG10265*, *CG15803*, *Roe*, *CG34353*,

CadN2, *CG43897*, *CG3407*, *dsxc73A*, or *bgm* do not produce a significant effect (dark gray). P-

values were calculated using one-way ANOVA followed by Dunnett's multiple testing correction.

* P < 0.05, *** P < 0.001.

Figure 1.

Figure 2

Figure 3

Figure 4