Deep learning and CRISPR-Cas13d ortholog discovery for optimized RNA targeting

Jingyi Wei[1,2,3], Peter Lotfy[4], Kian Faizi[4], Sara Baungaard[3], Emily Gibson[3], Eleanor Wang[4,5,6], Hannah Slabodkin[2,3], Emily Kinnaman[2,3], Sita Chandrasekaran[3,5,6], Hugo Kitano[7], Matthew G. Durrant[3,5,6], Connor V. Duffy[3,8], Patrick D. Hsu[3,5,6*], Silvana Konermann[2,3*]

[1]Department of Bioengineering, Stanford University, Stanford, CA

[2]Department of Biochemistry, Stanford University, Stanford, CA

[3]Arc Institute, Palo Alto, CA

[4]Laboratory of Molecular and Cell Biology, Salk Institute for Biological Studies, La Jolla, CA

[5]Department of Bioengineering, University of California, Berkeley, Berkeley, CA

[6]Innovative Genomics Institute, University of California, Berkeley, Berkeley, CA

[7]Department of Computer Science, Stanford University, Stanford, CA

[8]Department of Genetics, Stanford University, Stanford, CA

[*]Corresponding author

Correspondence: silvanak@stanford.edu (S.K.), pdhsu@berkeley.edu (P.D.H.)

Abstract

Transcriptome engineering technologies that can effectively and precisely perturb mammalian RNAs are needed to accelerate biological discovery and RNA therapeutics. However, the broad utility of programmable CRISPR-Cas13 ribonucleases has been hampered by an incomplete understanding of the design rules governing guide RNA activity as well as cellular toxicity resulting from off-target or collateral RNA cleavage. Here, we sought to characterize and develop Cas13d systems for efficient and specific RNA knockdown with low cellular toxicity in human cells. We first quantified the performance of over 127,000 RfxCas13d (CasRx) guide RNAs in the largest-scale screen to date and systematically evaluated three linear, two ensemble, and two deep learning models to build a guide efficiency prediction algorithm validated across multiple human cell types in orthogonal validation experiments (https://www.RNAtargeting.org). Deep learning model interpretation revealed specific sequence motifs at spacer position 15-24 along with favored secondary features for highly efficient guides. We next identified 46 novel Cas13d orthologs through metagenomic mining for activity and cytotoxicity screening, discovering that the metagenome-derived DjCas13d ortholog achieves low cellular toxicity and high transcriptome-wide specificity when deployed against high abundance transcripts or in sensitive cell types, including human embryonic stem cells, neural progenitor cells, and neurons. Finally, our Cas13d guide efficiency model successfully generalized to DjCas13d, highlighting the utility of a comprehensive approach combining machine learning with ortholog discovery to advance RNA targeting in human cells.

* * *

Introduction

The ability to perturb desired RNA molecules with high efficiency and specificity is required for functional elucidation of the transcriptome and its diverse phenotypes. Despite rapid progress in effective technologies for genome engineering, analogous systems for transcriptome engineering lag behind their DNA counterparts. While RNAi has long been used for RNA knockdown, it is challenging to engineer and suffers from widespread off-target effects (Jackson et al., 2003; Sigoillot et al., 2012) due to its important role in endogenous miRNA processing (Doench et al., 2003). The discovery and development of RNA-guided RNA-targeting CRISPR systems, such as Cas13 enzymes, provides an orthogonal and modular approach to overcome these limitations (Abudayyeh et al., 2016; East-Seletsky et al., 2016). Because CRISPR proteins are orthogonal to eukaryotic systems, they can be easily engineered to bind or cleave target RNA molecules. Further, their modular nature enables the facile fusion of effector domains to expand effector functionality. As a result, a broad suite of Cas13-based tools is now able to perturb RNA expression (Abudayyeh et al., 2017; Konermann et al., 2018) or splicing (Konermann et al., 2018), mediate RNA editing (Abudayyeh et al., 2019; Cox et al., 2017; Xu et al., 2021) or methylation (Wilson et al., 2020), as well as profile RNA-protein interactions (Han et al., 2020). These capabilities are now accelerating applications across the study of fundamental RNA biology, RNA-based therapeutics, and molecular diagnostics.

The Cas13 family is unified by the presence of two conserved HEPN ribonuclease motifs, and these enzymes are activated by binding to cognate target RNA as specified by the Cas13 guide RNA (Abudayyeh et al., 2016; East-Seletsky et al., 2016; Slaymaker et al., 2021; Zhang et al., 2018). Several subtypes have been defined on the basis of sequence diversity and domain architecture. Cas13d enzymes – in particular the engineered Cas13d from *R. flavefaciens* strain *XPD3002* (CasRx) (Konermann et al., 2018) – are the smallest and most efficient Cas13 RNA targeting effectors in mammalian and plant cells reported to date (Wessels et al., 2020; Li et al., 2021; Mahas et al., 2019), motivating their further characterization and optimization as RNA targeting tools. In order to successfully apply Cas13d in high-throughput applications, the ability to design highly effective guide RNAs is critical. Recent efforts to understand and predict Cas13d guide activity have taken a first step in this direction, by using a dataset of 2,918 guide RNAs across four transcripts to train a random forest model (Wessels et al., 2020) and by using combined datasets of 10,279 guides to train a deep learning model (Cheng et al. 2023). In addition to the relatively small datasets, the manual selection of guide sequence features (Wessels et al., 2020) or lack of secondary features (Cheng et al. 2023) has limited a broader understanding of Cas13d targeting preferences.

Here, we conducted the largest Cas13d screen to date, quantifying CasRx guide efficiency across >127,000 guide RNAs tiling 55 essential transcripts by measuring their effects on cell proliferation in human cells. We systematically compared a series of computational models on this dataset to predict guide activity. A deep learning convolutional neural network (CNN) model was able to most accurately predict highly effective guides. Model interpretation enabled us to discover a preferred sequence motif at spacer position 15-24 along with a preference for low guide free energy and high target region accessibility for high efficiency guides. We validated

110　the model against orthogonal datasets and confirmed high accuracy across target transcripts
111　and five different cell types.
112
Across the Cas13 subtypes structurally characterized to date, the RNA cleavage site formed by
the two HEPN domains is located distal to the guide binding groove (Liu et al., 2017; Slaymaker
et al., 2021; Zhang et al., 2018), which can result in the cleavage of non-target bystander RNA
molecules (known as 'collateral' cleavage) *in vitro* by the HEPN domains activated upon target
RNA binding. Initial reports for Cas13a, b, and d systems in routinely used mammalian cell lines
reported a low degree of off-target effects in eukaryotic cells (Abudayyeh et al., 2017; Cox et al.,
2017; Konermann et al., 2018). However, more recently, several groups reported cellular toxicity
and more pronounced off-target effects of CasRx, LwaCas13a, and PspCas13b in sensitive cell
types (Ai et al., 2022; Özcan et al., 2021), *in vivo* (Buchman et al., 2020), and when targeting
highly expressed transcripts (Shi et al. 2023).
123
To understand if this cellular toxicity is shared across Cas13 orthologs, we computationally
identified 46 novel Cas13d orthologs from recently reported prokaryotic genomes and
metagenomic contigs and screened them for target transcript knockdown activity and cytotoxic
effects in human cells. We identified DjCas13d, a highly efficient ortholog with minimal
detectable cellular toxicity when targeting highly expressed transcripts across multiple cell
types, including human embryonic stem cells, neural progenitor cells, and neurons.
Furthermore, we show that our CasRx-based guide design model extends to DjCas13d and
accurately selects highly efficient guides, illustrating its generalizability across effectors and cell
types. Overall, we advance the transcriptome engineering toolbox by developing a robust
Cas13d guide design algorithm based on a high-throughput guide screen
(https://www.RNAtargeting.org), and identifying a compact and high-fidelity Cas13d ortholog for
efficient RNA targeting. Finally, we outline a strategy to systematically develop and interpret
robust deep learning models for sequence-based classification.

Results

Deep learning of Cas13d guide RNA efficiency based on large-scale transcript
essentiality screening
In order to systematically understand factors impacting Cas13d guide efficiency, we generated a
library of more than 100,000 RfxCas13d (CasRx) guide RNAs and evaluated their efficiency in a
large-scale pooled screen. Reasoning that CasRx knockdown of essential transcripts would
lead to the depletion of highly effective guides due to reduced cellular proliferation, we selected
a set of 55 essential genes identified in three previously reported survival screens performed
with RNAi and CRISPR interference (CRISPRi) in K562 cells (Hart et al., 2015; Horlbeck et al.,
2016; Luo et al., 2008) for a proliferation-based survival screen. K562 cells were selected due to
their ease of use in pooled screens and our observation of variable CasRx-mediated
endogenous protein knockdown in this cell line **(Figures S1A, B).**
149
To perform the screen, we first generated stable K562 cell lines via transfection of an all-in-one
plasmid encoding the CasRx effector, PiggyBac transposase, and an antibiotic selection

cassette. Next, we designed CasRx guides that tile the 5' UTR, coding sequence (CDS), and 3'
UTR of the 55 essential transcripts with single nucleotide resolution. As controls, we designed
guides tiling 5 non-essential transcripts as well as 3,563 non-targeting guides. The effector cell
line stably expressing CasRx was transduced with a pooled lentiviral library containing all
144,745 guide RNAs. Cells were cultured for 14 days, after which we analyzed guide
abundances by NGS and computed a depletion ratio for each guide compared to its original
abundance in the input library (**Figure 1A**). Analysis of the cumulative distribution of guide
d14/input ratio demonstrated that the top 20th percentile of guides targeting essential transcripts
are clearly separated from guides targeting non-essential transcripts or non-targeting guides
(Figure 1B).
162
Essential transcripts may vary in their magnitude of impact on cell proliferation and survival
upon depletion. A transcript-level analysis of guide depletion confirmed this expectation (**Figure**
S1C). In order to compensate for this in our analysis going forward, we selected the most
effective guides for each individual transcript (see Methods for a full description of selection
parameters) as high efficiency guides. A heat map representation of the positions of these high
efficiency guides within each target transcript revealed a striking degree of clustering, leading to
guide hot spots and deserts along the transcript and clearly deviating from a random distribution
(Figure 1C). Multiple factors could be responsible for the observed clustering of high efficiency
guides, including sequence-, structure-, or position-based effects of the guide RNA or target
transcript.
173
Prediction of CasRx guide activity based on guide RNA sequence alone
We sought to systematically analyze these potential features that could distinguish high
efficiency Cas13d guides and develop computational algorithms to predict guide efficiency.
Initial analysis of the correlation of nucleotide identity with guide efficiency at each position
along the 30 nt spacer showed a preference for G and C at the direct repeat-proximal spacer
positions 15-24 (**Figure S2A**). Therefore, we reasoned that spacer sequence alone might be
predictive of guide efficiency when used as model input. We then developed a series of
computational models for prediction of guide efficiency based on one-hot encoding of the 30 nt
guide spacer sequence without manual sequence feature selection. To understand the impact
of computational model type, we systematically built and assessed the following models: 3
linear models employing logistic regression (Lasso Regression (L1), Ridge regression (L2) or
Elastic Net (EN)), 2 ensemble models (Random forest (RF) and Gradient-boosted tree (GBT))
and 2 deep learning models (convolutional neural network (CNN) and bidirectional long short-
term memory neural network (LSTM)) (**Figure 1D**).
188
All of these models were trained to classify high efficiency guides for target transcripts. Due to
the observed high degree of clustering of effective guides along a transcript (**Figure 1C**),
models that are tested on held-out guides from the same transcripts they were trained on would
potentially be subject to overfitting by learning the targeting hotspots specific to those
transcripts. To alleviate overfitting and ensure model generalizability to other transcripts, we
employed 9-fold cross-validation on the 54 target transcripts (leaving out *RPS19BP1* as it
clustered with non-essential transcripts (**Figure S1C**)), with models being trained and tested on

non-overlapping sets of transcripts. We compared the performance of all 7 models and
observed high model performance for the gradient-boosting tree (GBT) and the two deep
learning models based on Area Under the Receiver Operating Characteristic curve (AUROC),
which evaluates prediction accuracy for both the positive class (high efficiency guides) and the
negative class, and Area under the Precision-Recall Curve (AUPRC) metrics, which focuses
primarily on the prediction accuracy of the positive class (high efficiency guides), across all 9
fold splits (**Figure 1E**).

Overall, the CNN model performed best with a high AUROC of 0.845 (relative to a baseline of
0.5) and a high AUPRC of 0.541 (relative to a baseline of 0.18), so we chose this model for
further refinement and evaluation. The high prediction accuracy of this model based on the
spacer sequence alone indicates that sequence is a primary factor determining guide efficiency.
We further determined that the addition of target flanking sequences of varying length from 1-7
nt to the CNN model did not meaningfully improve model performance (**Figure S2B**), consistent
with our previous biochemical studies suggesting a lack of strong flanking sequence
requirements (Konermann et al., 2018). To understand the minimal spacer length required for
accurate prediction, we computationally truncated the spacer sequence from the 3' end in the
CNN model input, and found only a minor impact on model accuracy until reaching a spacer
length of 24 nt, after which a gradual drop in AUROC and AUPRC was observed (**Figure S2C**).
We validated this experimentally, demonstrating decreasing target knockdown when using
guides shorter than 24 nt in spacer length (**Figure S2D**).

Addition of secondary features improves guide efficiency prediction accuracy
Beyond guide sequence alone, secondary guide attributes such as guide unfolding energy or
target site position (CDS or UTR) may impact guide performance. To understand their potential
contribution, we first evaluated the correlation of such secondary features with guide efficiency
(**Figure 1F schematics, S3A-F**). We found that higher predicted guide and target RNA
unfolding energy, implying more highly structured RNA sequences, were predictive of poor
guide efficiency. We also observed a preference for intermediate spacer GC content (45-55%),
guides targeting the coding region (CDS), as well as guides targeting regions conserved across
transcript isoforms.

As most of the secondary features investigated exhibited a modest correlation with guide
efficiency, we tested whether they would improve model performance when added to the spacer
sequence-only CNN model. When adding these features individually, we found that the guide
target site position had the most prominent effect, followed by target and guide RNA folding
energy (**Figure S3G**). The addition of spacer GC content did not significantly improve model
performance, consistent with our expectation that this feature has been successfully captured
by the spacer sequence-only CNN model. Sequentially including each secondary feature ranked
by their individual contribution into the sequence-only CNN model, we found that AUROC and
AUPRC were improved with each addition, leading to a final model with a very high average
AUROC of 0.875 and a high average AUPRC of 0.638 (**Figure 1F** and **S3H-J** for feature
variations). Adding the same set of secondary features also improved the GBT model (**Figure**

S4), the best performing model not based on deep learning, indicating the contribution of these
secondary features to guide efficiency.
241
One of the key applications of a predictive model like this one would be to accurately predict the
most effective guides in order to aid in guide and library design. The CNN model returns a float
score ranging from 0 to 1 for every guide, and different thresholds can be chosen for high
efficiency guide classification. To evaluate model performance for optimal guide selection, we
set a high model score threshold of 0.8 and plotted the true percentile rank distribution of the
guides above the score threshold. As expected, the guides were heavily skewed towards the
highest efficiency ranks, with a true positive ratio of 0.83 (83% being true high efficiency guides
(top 20th percentile)). Setting an even more stringent model score threshold to 0.9 further
increased the true positive ratio to 93% (**Figure S3K**).
251
Model interpretation reveals favored sequence and secondary features of high efficiency
guides
Having built high performance models that accurately predict efficient guides, we asked whether
these models could help us understand the features contributing to guide efficiency by using
three model interpretation methods. We first used an integrated gradients approach (IG)
(Sundararajan et al., 2017) to provide observability for our CNN model. We began with the
guide sequence preferences learned by the model, and IG analysis on each position in the
guide spacer sequence nominated a core region of position 15-24 as a major contributor to
guide efficiency (**Figure 2A**). Consistent with our original correlation analysis (**Figure S2A**), IG
analysis on each positional nucleotide in the guide sequence revealed a clear preference for an
alternating stretch of guanines, cytosines and guanines ($G_{15-18}C_{19-22}G_{23-24}$) in this core region
(**Figure 2B**).
264
To confirm the favored sequence features across models and model interpretation methods, we
further applied SHapley Additive exPlanations (SHAP), a game theoretic approach (Lundberg et
al., 2020) to our GBT model, and a similar sequence preference in the same core region was
observed (**Figures S5A, B**). In contrast, this unique sequence preference was not found for
Cas13a when we performed a correlational analysis of available datasets (Abudayyeh et al.,
2017; Metsky et al., 2022) (**Figure S6**). Indeed, no consistent sequence preference or core
region emerged across the Cas13a datasets analyzed, which could be due to intrinsic
enzymatic properties of Cas13a or limitations in the size of available datasets.
273
As our IG and SHAP analyses investigated each position in the guide sequence independently,
we further sought to determine the role of specific motifs (nucleotide combinations) in guide
efficiency. We employed Transcription Factor Motif Discovery from Importance Scores (TF-
MoDISco), an algorithm that identifies sequence patterns or motifs incorporated in deep learning
models by clustering important sequence segments based on per-position importance scores
(Shrikumar et al., 2018). We discovered a total of 14 distinct sequence patterns associated with
high efficiency guides from the CNN model, with the top 5 patterns shown in **Figure 2C**. As TF-
MoDISco was initially applied for the identification of transcription factor binding motifs, it is
designed to identify motifs in a position-independent manner. In our analysis, we noticed that all

identified patterns were anchored to a specific position centered around guide spacer
nucleotides 18-20 (**Figure S7A**), consistent with our prior observation of a core region**.**
285
Strikingly, all top 5 sequence patterns contained a cytosine at position 21, with a single guanine
at varying positions in the core region across the different patterns. Taken together, the
identified motifs can be summarized as GN_xC_{21} or $N_xC_{21}G$ within the core region. Generally, the
patterns were sparse and characterized by just two dominant bases (one G and one C), in
contrast to the longer 10-base motif that the individual position-level analysis would have
suggested (**Figures 2B and S5B**). Consistent with our results above, an analysis of enriched
and depleted 3-mers in high efficiency guides across the spacer sequence revealed that
enriched 3-mers were again clustered in the core region (position 15-24) (**Figure S7B**). In
addition to the consistent finding of a prominent enrichment of C at position 21, they revealed a
preference for A or T intercalated with G and C (**Figures S7B, C**), a finding that was obscured
in the per-position analysis. Analysis of enriched and depleted 4-mers in high efficiency guides
also led to a similar finding (**Figure S7D**). A/T substitutions within the 10-base motif ($G_{15-18}C_{19-}$
$_{22}G_{23-24}$) (**Figure 2D**) and analysis of the GC content in the core region (**Figure 2E**) for high
efficiency guides further confirmed a preference for a medium GC content via A/T nucleotides at
the N positions of the key GN_xC_{21} or $N_xC_{21}G$ motif.
301
Next, we used IG and SHAP to investigate the contribution of secondary features in the CNN
and GBT models. IGs revealed that targeting the beginning of the 5′ UTR and the end of the 3′
UTR was the most disfavored, while targeting the coding region (CDS) was generally favored,
with a slight preference for the beginning of the CDS (**Figures 2F, G**). In agreement with our
correlation analysis, guide and target unfolding energy also had a relatively high impact on
guide efficiency, with lower unfolding energy favored by high efficiency guides (**Figures 2H, I**).
SHAP analysis on our GBT model showed a consistent direction of feature contribution to guide
efficiency (**Figure S5C**) and ranked spacer sequence composition as the most important
feature.
311
312   Taken together, our systematic model interpretation was consistent across models and analysis
313   approaches, was able to rank features by their contribution toward guide classification, and
314   significantly expanded our understanding of preferred longer-range sequence motifs that were
315   missed by simpler correlational analyses.
316
Systematic validation of the guide efficiency model across 5 cell types with endogenous
protein knockdown
Next, we sought to experimentally validate our model through CasRx-mediated knockdown of
cell surface markers, reasoning that an orthogonal readout to transcript essentiality and cell
survival would ensure generalizability of our model predictions to multiple readout modalities. To
this end, we performed a screen using a library of 3,218 guides tiling the transcripts of two cell
surface markers, *CD58* and *CD81*, with single-nucleotide resolution. 10 days after lentiviral
transduction of the guide library, cells were FACS sorted into 4 bins on the basis of target
protein expression level (**Figure 3A**) and the enrichment of individual guides in the top and
bottom bins (exhibiting the greatest or least magnitude of knockdown, respectively) was

assessed. We observed clear separation of the most efficient targeting guides from the non-
targeting guides based on the enrichment ratio, with zero non-targeting guides appearing in the
top 20th percentile of guide efficiency (**Figure 3B**).
330
We evaluated our CNN model's performance on this new dataset and found that an ensemble
CNN model comprising all 9 fold splits of the survival screen outperformed each individual split
model (**Figure S8A**) and achieved high prediction accuracy for both *CD58* (AUROC of 0.88 and
AUPRC of 0.66) and *CD81* (AUROC of 0.86 and AUPRC of 0.62) (**Figure 3C**). This
performance is comparable to the model accuracy on held-out essential transcripts from our
initial screen (**Figure 1F**), highlighting its generalizability. Compared with two existing Cas13d
guide design models (Wessels et al., 2020, Cheng et al. 2023), our model showed the highest
AUROC, AUPRC, and Spearman correlation. Importantly, we showed that at a 0.9 score cutoff,
our model exhibited a very high true positive ratio of 0.93 and 0.9 for *CD58* and *CD81*,
respectively, in contrast to the Wessels et al. model (0.52 for both *CD58* and *CD81*) and
DeepCas13 (0.38 for *CD58* and 0.35 for *CD81*) (**Figure 3C**). The far higher true positive ratio at
high score cutoffs underlines the superior utility of our model for key applications such as
predicting the top 3-10 guides per target transcript in individual targeting or library-based
screening approaches. Illustrating this use case, we examined the true percentile rank of the top
10 predicted high efficiency guides for *CD58* and *CD81*, showing that 10/10 guides for *CD58*
and 9/10 for *CD81* were highly effective (**Figure 3D**).
347
To assess generalizability to other cell types, we evaluated our model's performance on a
published CasRx guide tiling dataset (~3000 guides in HEK293FT cells from the Wessels et al.
training dataset). Our model showed high AUROC (0.85, 0.88 and 0.85 for *CD46, CD55 and*
CD71, respectively), AUPRC (0.59, 0.59 and 0.67), Spearman correlation (0.67, 0.69 and 0.66),
and true positive ratio (0.76, 0.9 and 0.94 at a 0.9 score cutoff) (**Figure 3E**). Among the top 10
predicted high efficiency guides, 90% were highly efficient (falling into the top 20% percentile of
efficient guides) (**Figure 3F**). When compared against the Wessels et al. model on opposing
datasets (**Figure S8B**), our model showed significantly higher prediction accuracy using all
evaluation metrics (AUPRC: 0.617 vs 0.379; Spearman correlation r_s: 0.675 vs 0.391; AUROC:
0.873 vs 0.733; true positive ratio (0.9 cutoff): 87% vs 51%), further supporting the
generalizability and high performance of our model.
359
As a final test of the ability of our model to predict efficient guides for knockdown of desired
transcripts in different cell types, we selected 5 top scoring guides and 5 low scoring guides
(excluding the very bottom of our ranking) for two different transcripts (*CD59* and *CD146*), and
tested the knockdown efficiency of each guide in Hela, U2OS, and A375 cells (**Figure 3G**).
Across all three cell lines, the top scoring guides showed very efficient target knockdown (72%-
98% with a median of 90%) while low scoring guides showed variable and significantly lower
levels of knockdown (6%-70% with a median of 35%), confirming the utility and generalizability
of our model across 5 cell types (K562, HEK293FT, Hela, U2OS, and A375).
368
Discovery of DjCas13d, a high-efficiency RNA targeting enzyme with minimal cellular
toxicity in human cells

In genome engineering, two of the most important features are efficiency and specificity. A key
emerging limitation of several Cas13 systems is the induction, in certain contexts, of cellular
toxicity by its RNA trans-cleavage activity (Ai et al., 2022; Buchman et al., 2020; Özcan et al.,
2021), hampering their application as a generalizable transcriptome engineering tool. In the
context of this study, we also observed various degrees of cellular toxicity for CasRx when
paired with highly efficient guides in the A375 cell line (**Figure S9A).**
377
To address this, we reasoned that the evolutionary diversity of Cas13d enzymes may have
already developed solutions to these challenges. To develop a more broadly useful
transcriptome engineering tool, we sought to identify a Cas13d ortholog that combines the key
positive traits of CasRx, like its small size and high targeting efficiency, with low cellular toxicity.
We applied our previously described computational approach for Cas13d discovery (Konermann
et al., 2018) to an expanded database of metagenomic datasets and discovered 46 previously
uncharacterized Cas13d enzymes, expanding the known Cas13d family from 7 to 53 members
(**Figure 4A**, **Table S7**).
386
To evaluate these novel Cas13d enzymes for mammalian transcript knockdown, we
synthesized human codon-optimized constructs of each enzyme with NLS (nuclear localization
sequence) and NES (nuclear export sequence) fusions and measured their ability to knockdown
the mCherry reporter transcript using a matched guide array containing two mCherry targeting
guides. We identified 14 enzymes exhibiting >55% knockdown efficiency (**Figure 4B**) in this
reporter assay. Because reporter knockdown is often weakly predictive of Cas13 performance
on endogenous targets, we further tested the 14 orthologs on our shortlist for their knockdown
efficiency when targeted to the endogenous *CD81* transcript. With this more stringent test, 7
orthologs exhibited >50% knockdown efficiency (**Figure 4C**), and we focused on these for
further characterization.
397
Having identified this shortlist of the most efficient Cas13d enzymes, we next evaluated their
cytotoxic effects in human embryonic stem cells (hESC), since we previously observed issues in
this cell type with CasRx. When targeting the non-essential transcript *CD81* in this highly
sensitive cell type, we were able to observe a significant reduction in viable cells expressing
CasRx and most of the other Cas13d orthologs (**Figure 4D**), consistent with cytotoxic effects on
other sensitive cell types reported in the literature (Özcan et al., 2021). Strikingly, two of the
orthologs we tested (DjCas13d and Ga_0531) led to no detectable reduction of viable cell
counts (**Figure 4D**). Of those two, we chose DjCas13d for additional characterization given its
high knockdown efficiency (>80% in hESCs) (**Figure 4E**) and unusually small size (877aa,
compared to 967aa for CasRx) (**Figure 4C**).
408
In a further evaluation across three guides each for three transcripts in hESCs, DjCas13d
showed no significant effects on viable cell counts in contrast to CasRx, which caused
significantly reduced viable cell counts in eight out of nine guides (**Figure 4F**). In terms of
knockdown efficiency, DjCas13d showed high knockdown efficiency of >70% for most guides
tested (median of 71.5%) – efficiency that was comparable to CasRx (median of 77.4%) (**Figure**
4F).

DjCas13d induces minimal cellular toxicity when targeting highly expressed transcripts

Recent work (Ai et al. 2022; Shi et al. 2023) and our results in stem cells (**Figure 4F**) highlighted high target transcript abundance as a key variable for Cas13-mediated cellular toxicity in addition to the importance of cell type. In our own experiments in hESCs, we also observed the lowest survival rate for CasRx when targeting the most abundant transcript – *CD24* – while no such impact was observed for DjCas13d (**Figure 4F**). In order to further compare CasRx and DjCas13d under conditions known to promote cellular toxicity, we targeted three previously described highly expressed transcripts (*ACTG1*, *HNRNPA2B1*, *FTH1*) (Shi et al. 2023) in HEK293FT cells and confirmed a significant reduction of the number of viable cells when using CasRx but not DjCas13d (**Figure 4G,** all guides significant at P<0.0001). We targeted three medium- and three low expression level transcripts, confirming that lower expression of the target transcript alleviated the toxicity induced by CasRx (**Figure 4G**), consistent with initial reports (Konermann et al., 2018). By contrast, we observed minimal impact on viable cell counts when using DjCas13d to target any of these transcripts (**Figure 4G**), despite comparable knockdown efficiency of DjCas13d (knockdown median of 88%) to CasRx (median of 84%).

In a second head-to-head comparison, we tested DjCas13d against the recently reported Cas7-11 enzyme, which does not belong to the Cas13 family of CRISPR enzymes and was reported to have no impact on cell viability due to its distinct RNA cleavage mechanism (Kato et al., 2022; Özcan et al., 2021). We demonstrate that both DjCas13d and Cas7-11 have a comparably low impact on cell viability and proliferation (90% median cell count for DjCas13d across all targeting conditions, and 73% for Cas7-11) when targeting the same medium to highly expressed transcripts - in stark contrast to CasRx (46% median cell count). However, Cas7-11 suffered from diminished knockdown efficiency (median of 57%) compared to DjCas13d and CasRx (median of 88% and 84%, respectively) (**Figure 4G**).

Overall, we conclude that DjCas13d combines the best features of CasRx and Cas7-11, exhibiting low cellular toxicity and high knockdown efficiency. 84% of guides tested with DjCas13d showed >80% survival rate and >60% knockdown, while only 32% of CasRx guides and no Cas7-11 guides met these cutoffs.

DjCas13d activity can be accurately predicted with our guide efficiency model

Given that DjCas13d belongs to the same subtype of CRISPR effectors as CasRx, we next sought to test whether our Cas13d guide design model could be successfully applied to this new Cas13d ortholog. Encouragingly, our data in **Figure 4F** and **G** demonstrated high efficacy of knockdown with guides recommended by the model when using DjCas13d across 12 transcripts of different expression levels and in different cell types. To further explicitly validate the model performance for DjCas13d, we selected a set of top and bottom scoring guides for a total of eleven transcripts across a range of expression levels in hESCs, HeLa, and U2OS cell lines. Across hESCs (**Figure 4H**) as well as Hela and U2OS cells (**Figure 4I**), the predicted high efficiency guides resulted in a significantly higher degree of protein knockdown (median of 73.9%) compared with low-scoring guides (median of 19.7%) (**Figures 4H, I**). Altogether, these results demonstrate that our model generalizes to the novel DjCas13d ortholog, resulting in

459 reliable knockdown performance and lack of apparent cellular toxicity even in sensitive cell
460 types and for highly abudant transcripts. Given that the sequence divergence between
461 DjCas13d and CasRx (29.9%) is similar to the divergence between other Cas13d orthologs from
462 our new metagenomic mining (~29.4% on average), we expect that our guide design model
463 may generalize to other Cas13d effectors as well.
464
DjCas13d exhibits high transcriptome-wide specificity
The context-dependent cellular toxicity mediated by many Cas13 enzymes is hypothesized to
result from collateral cleavage of bystander transcripts (Buchman et al. 2020; Özcan et al. 2021;
Ai et al. 2022; Shi et al. 2023). This is consistent with the observation that cellular viability and
proliferation are more noticeably impacted when targeting more abundant transcripts – which
would result in a larger number of activated Cas13 enzymes per cell and therefore more
potential collateral RNA cleavage.
472
To investigate this hypothesis and compare the collateral and off-target effects between CasRx
and DjCas13d, we performed RNA-seq two days after CasRx or DjCas13d-mediated
knockdown of *CD81* (307 Transcripts Per Million (TPM)), *FTH1* (1219 TPM) and *ACTG1* (3728
TPM) in HEK293FT cells (**Figure 5A**). Our transcriptome-wide analysis revealed significantly
more non-target transcripts affected by CasRx when targeting more highly expressed transcripts
(*ACTG1>FTH1>CD81*), indicating greater levels of collateral or off-target effects (**Figure 5A**). In
contrast, we observed minimal transcriptome-wide perturbation by DjCas13d apart from
knockdown of the intended target transcript (**Figure 5A**).
481
Next, we extended our RNA-seq analysis to assess consequences of CasRx and DjCas13d in
more sensitive hESC cells when targeting genes with high (*CD24*), medium (*CD81*), or low
(*TFRC*) expression levels. CasRx-mediated knockdown of high and medium expressed genes
resulted in rampant loss of cell viability, making transcriptome analysis impossible in many
samples. Consistent with the high survival of sensitive cell types following DjCas13d treatment
above, this toxicity was not observed for DjCas13d targeting the same transcripts. Similar to the
HEK293FT RNA-seq above, we observed a significant reduction in off-target transcriptome
perturbations when using DjCas13d (0 off-targets for most guides tested, with a modest 7 and
103 off-targets for the two guides targeting *CD24*) compared to CasRx (hundreds of off-targets
even when targeting low- and medium-expression transcripts, and rampant cellular toxicity
when targeting highly expressed transcripts) (**Figure 5B**).
493
Importantly, in order to rule out transcriptome-wide depletion that would be difficult to detect via
differential RNA-seq, we used defined concentrations of exogenous RNA spike-ins to assess
total RNA amount per cell. While CasRx showed a significant decrease in total RNA abundance
across guides targeting *CD71*, DjCas13d did not display significant global RNA depletion with
any guide/target tested, consistent with its low off-targets and low toxicity (**Figure S10A**). As an
additional measure of transcriptome integrity, we visualized total RNA extracted from these
samples and showed that while RNA integrity for DjCas13d was intact, CasRx targeting resulted
in the appearance of a smaller molecular weight band between the 28S and 18S for all targeting
guides (**Figure S10B**), which has also been noted by other groups (Shi et al., 2023).

503
To distinguish between guide-specific off-target effects and universal sequence-indiscriminate
collateral effects in our CasRx datasets, we analyzed the overlap between up- and down-
regulated transcripts among different guides, targets and cell types (**Figures S10C, D, E, F**).
We found a meaningful overlap between the significantly upregulated transcripts across
different CasRx conditions, with enrichment of the unfolded protein response signaling pathway,
suggesting that CasRx mediated non-target-specific collateral activity may stimulate generalized
cellular stress responses.
511
DjCas13d is a effective tool for gene knockdown in many sensitive cell types
Given the promise of DjCas13d as a high-fidelity and low-toxicity RNA targeting tool, we sought
to apply DjCas13d to RNA targeting in sensitive biological processes and therapeutically-
relevant cell types. Our demonstration of CasRx toxicity in hESC cells led us to assess
DjCas13d knockdown in the context of hESC differentiation into neuronal progenitor cells
(NPC), hematopoietic progenitor cells (HPC), and neurons. DjCas13d was delivered via an
inducible Piggybac system at the stem cell stage and induced during differentiation. In NPCs,
we targeted five transcripts including highly-expressed genes like *BSG* and *THY1*, and lower
expressed transcripts such as *CD46* with one or two top-scoring guides per gene. We observed
high cellular survival in all cases with no significant decrease relative to non-targeting
conditions, and effective knockdown efficiencies in most cases, with a median of 63% (**Figure**
6A). In HPCs, we observed 46-69% knockdown of the target proteins CD81 and TRFC in
DjCas13d-expressing cells with no detectable survival defect **(Figure 6B)**. In both of these
cases, we confirmed that the expected markers of differentiation efficiency were not affected by
DjCas13d targeting (SOX1 and PAX6 for NPC, CD43 for HPC) (**Figures 11A,B**). Finally, we
differentiated hESCs to neurons using Neurogenin-2 (Ngn2) directed differentiation and
assessed DjCas13d's ability to knock down two proteins, CD81 and CD24, with 3 top-scoring
guides each. We observed uniform knockdown of approximately 50% in all cases (measured at
the protein level via FACS), coupled with high cell survival near 100% (median of 98%) (**Figure**
6C). Altogether, these data illustrate the broad applicability of DjCas13d across multiple target
genes in sensitive cell types of high biological and therapeutic interest.
533
534 To support easy use of both DjCas13d and CasRx for RNA targeting, we created a freely
535 accessible portal to run our model for Cas13d guide prediction on all human and mouse
536 transcripts and custom target sequences. This community resource is available at
537 http://RNAtargeting.org.

Discussion

In this study, we applied CasRx for large-scale screening across 127,000 guides against 55
target transcripts in human cells, a dataset that is >12 times larger than previous Cas13 guide
design studies (Wessels et al. 2020; Cheng et al. 2023). Using this dataset, we developed a
highly accurate, deep learning-based Cas13d guide efficiency model to nominate highly efficient
guides for transcripts of interest. The model exhibits excellent performance across two screen
modalities, nine cell types, and two diverse Cas13d orthologs, illustrating its generalizability for

predicting highly effective guides across different contexts. The major factors contributing to our
model's generalizability include its primary reliance on the guide RNA spacer sequence - a cell
type-independent feature - as well as the 9-fold cross-validation of the model on non-
overlapping sets of transcripts, which alleviates overfitting to targeting hotspots specific to
certain transcripts.
550
Previous attempts to predict CRISPR guide efficiency have primarily relied on manual selection
of a limited set of guide sequence features combined with simpler machine learning models,
such as elastic nets (Horlbeck et al., 2016), SVM (Doench et al., 2016), or random forest
approaches (Wessels et al., 2020). More recently, deep learning models, which are able to learn
complex, high-order patterns and features automatically from raw data, have been employed to
predict guide efficiency for Cas9 activity (Chuai et al., 2018; Kim et al., 2019; Xue et al., 2019),
Cpf1 (Kim et al., 2018), base editors (Arbab et al., 2020; Koblan et al., 2021), Cas13a (Metsky
et al., 2022) and Cas13d (Cheng et al. 2023).
559
Here, we directly compared two deep learning models with linear and ensemble methods
(elastic nets, random forest, and gradient-boosted trees) for guide efficiency prediction, finding
that the deep learning model (CNN) outperformed the other approaches. This illustrates the
power of deep learning models in sequence-based prediction tasks due to its automatic feature
selection and ability to identify motifs or long-range interactions given a sufficiently large dataset
(>100,000 guides). Furthermore, we show that our model significantly outperforms the current
state-of-the-art models (Wessels et al. 2020; Cheng et al. 2023) (**Figures 3C, S8B**).
567
While deep learning models can extract important higher-order features automatically from raw
inputs, the interpretation of feature contributions is challenging. Prior deep learning models for
Cas9 (Chuai et al., 2018; Xue et al., 2019) and other sequence-based applications (Alipanahi et
al., 2015; Kelley et al., 2016; Lanchantin et al., 2016) mainly employed neuron visualization
methods to unveil important motifs. These approaches are able to successfully identify patterns
recognized by individual filters, but can suffer from redundancy of the identified motifs. Recently
developed interpretation methods such as Integrated Gradients, SHAP, and TF-MoDISco, can
address these limitations and have begun to be applied to identify consolidated and non-
redundant motifs for transcription factor binding (Avsec et al., 2021). In this report, we evaluated
feature importance directly from the deep learning model using these new model interpretation
approaches. This allowed us to discover a core region at guide spacer position 15-24 with a
specific sequence composition predictive of high efficiency guides. Comprehensive motif
analysis revealed a preference for $GW_{1-4}C_{21}$ or $C_{21}W_{0-2}G$ motif. In contrast, analysis of base
preference at individual positions and correlation-based evaluation of feature importance
(Wessels et al., 2020) obscured this motif. This underscores the utility of the combination of
deep learning models that are able to learn higher order sequence features along with
advanced motif-discovery approaches for model interpretation such as TF-MoDISco used here
– the first time, to our knowledge, that such an approach has been applied to CRISPR guide
activity prediction models.
587

In addition to effective guide selection, cellular toxicity has emerged as a significant challenge
for Cas13 applications, effects likely mediated by off-target and/or collateral RNA cleavage
(Buchman et al. 2020; Özcan et al. 2021; Ai et al. 2022; Shi et al. 2023). Initial reports
developing diverse Cas13 effectors for mammalian transcript knockdown demonstrated high
specificity and lack of apparent cellular toxicity in HEK293FT cells, plants, and animal embryos
(Abudayyeh et al., 2017; Cox et al., 2017; Konermann et al., 2018; Kushawah et al., 2020;
Mahas et al., 2019). However, several recent studies have reported marked cellular toxicity of
these effectors in other cell types or target contexts (Buchman et al., 2020; Özcan et al., 2021).
596
Two recent studies aiming to reconcile these reports concluded that collateral RNA cleavage by
Cas13 enzymes is correlated with the expression level of the target transcript, and that the
effect on cellular toxicity is dependent on the cell type (Ai et al. 2022; Shi et al. 2023), indicating
that highly expressed transcripts and sensitive cell types are prone to Cas13-mediated collateral
cleavage and toxicity. Our data comparing CasRx's effect across cell types and endogenous
target RNAs with varying expression levels supports this conclusion. We reasoned that more
robust CasRx RNase activation upon higher target transcript levels would result in a greater
amount of collateral RNA cleavage, which in turn could activate cellular stress pathways and
lead to toxicity.
606
To advance Cas13 applications in sensitive cell types and therapeutic scenarios, our discovery
of the DjCas13d ortholog promises to address current limitations of both CasRx (context-
dependent cellular toxicity) and Cas7-11 (efficiency and size). DjCas13d exhibits minimal
cellular toxicity even in challenging conditions, and achieves high efficiency and transcriptome-
wide targeting specificity against highly expressed transcripts across various cell types. We
further demonstrate efficient and high-viability endogenous RNA targeting with DjCas13d in
hESC-derived neuronal progenitor cells (NPCs), hematopoietic progenitor cells (HPCs), and
neurons. Therefore, DjCas13d is poised to overcome the limitations of previous tools. Future
work characterizing mechanistic distinctions between CasRx and DjCas13d may reveal further
protein engineering opportunities.
617
Taken together, DjCas13d paired with our state-of-the-art Cas13d guide design model provides
a comprehensive solution for 3 key challenges in the RNA targeting toolbox by enabling high
efficiency, cell viability, and specificity. We further envision that the deep learning model
architecture, systematic feature engineering, and model interpretation approach outlined in this
study will be broadly applicable to other sequence-based tasks, such as the prediction of guide
RNA activities for newly discovered CRISPR enzymes, DNA/RNA modifications, and DNA/RNA-
protein interactions.
625
626

Figures

Figure 1: Deep learning of Cas13d guide RNA efficiency based on large-scale transcript essentiality screening

A. Schematic of the pooled CasRx guide tiling screen for essential transcript knockdown as a readout of per-guide knockdown efficiency. Over 127,000 targeting guide RNAs were included.
B. Cumulative distribution of the ratio of relative guide abundance at day 14 compared to the input library across guides targeting essential gene transcripts (blue), non-essential gene transcripts (orange), and non-targeting guides (green). The red dashed line indicates the ratio at the top 20th percentile of essential transcript targeting guides. **C**. Heat map of the positional distribution of high efficiency guides along each transcript. From here forward, high efficiency guides are defined as the top 20% guides within each transcript with a d14/input ratio lower than 0.75 after essential off-target filtering. Heat map color indicates the number of overlapping high efficiency guides at each nucleotide position along the transcript, and the inlaid histogram depicts the observed frequency distribution of these data (blue) as compared to a random distribution of 20% of guides in the library (orange curve). **D.** Schematic of the computational algorithms assessed in this study to predict guide efficiency based on spacer sequence alone.
E. Comparison of prediction accuracy between linear, ensemble and deep learning models across 9-fold splits of held-out transcripts. Averages of Area Under the Receiver Operating Characteristic curve (AUROC) and Area Under the Precision-Recall Curve (AUPRC) across test sets from all 9 folds are shown ± SD. LR - L1, logistic regression with L1 regularization (Lasso Regression); LR - L2, logistic regression with L2 regularization (Ridge regression); LR - EN, logistic regression with elastic net regularization (Elastic Nets) ; GBT, Gradient-Boosted Tree; RF, Random Forest classifier; CNN, Convolutional Neural Network; biLSTM, Bidirectional long short-term memory neural network. Note that the baseline for AUPRC is equal to the fraction of positive class (high efficiency guides), in this case 0.18. **F.** Secondary features were evaluated for their ability to improve sequence-only model performance. Each secondary feature (or feature group) was added to the CNN model sequentially, ordered by its individual contribution to model performance in **Figure S3G**. AUROC and AUPRC (mean ± SD) of all test sets from the 9-fold split of transcripts are shown.

Figure 2: Deep learning model interpretation reveals favored sequence motifs and secondary features of high efficiency guides

A. Evaluation of the importance of each position in the guide spacer sequence in the CNN model using Integrated Gradients (IG). Higher absolute gradient values indicate greater importance for predicting a high efficiency guide. The gray box highlights the identified core region (position 15-24). **B.** Evaluation of the importance of each positional nucleotide in the guide sequence in the CNN model by IG. **C.** Top 5 sequence patterns identified by TF-MoDISco (Transcription Factor Motif Discovery from Importance Scores) in the CNN model. Patterns are aligned to the 30 nt spacer according to the mode position of the seqlets (sequence regions with high importance based on IG scores) in each pattern (**Figure S7A**). **D.** Fraction of high efficiency guides that contain the 10-base motif shown in panel B and A/T substitutions within the 10-base motif. **E.** Fraction of high efficiency guides across different core region GC content. Guides were divided into eleven bins based on the GC content in their core region (position 15-24), and the fraction of high efficiency guides belonging to each bin is plotted. **F.** Contribution of target transcript region (5' UTR, CDS, or 3'UTR) to guide efficiency in the CNN model. The bar plots indicate average IGs of all test samples with different target position flags. **G.** Contribution of position within each transcript target region to guide efficiency in the CNN model. The scatter plots indicate individual IG values against individual input values across all test samples. The reference points are set to 0 for each transcript region. **H.** Contribution of predicted guide unfolding energy to guide efficiency in the CNN model. The reference point is set to 0. **I.** Contribution of predicted target unfolding energy to guide efficiency in the CNN model. The reference point is set to 0.

Figure 3: Systematic validation of the guide efficiency model across 5 cell types with endogenous protein knockdown

A. Schematic of the pooled CasRx guide tiling screen targeting *CD58* or *CD81* transcripts in K562 cells followed by flow cytometry-based readout of cell-surface CD58 or CD81 protein abundance. **B.** Cumulative distribution of guide enrichment ratios for *CD58*, *CD81* and non-targeting guide categories, calculated as the ratio of guide percentage in bin 1 (greatest knockdown) relative to the sum in bin 1 and bin 4 (least knockdown). Red dashed lines indicate the ratio for the top 20th percentile of targeting guides. **C.** Model comparison on *CD58* and *CD81* guides. CNN, the ensemble CNN model built on the survival screen data in this work; Wessels et al. model, a previously published CasRx random forest model (Wessels et al., 2020); DeepCas13, a previously published CasRx deep learning model (Cheng et al. 2023). Model performance is evaluated by AUROC, AUPRC, Spearman's correlation coefficient (r_s) and true positive ratio at 0.8 and 0.9 model score cutoffs across guides targeting *CD58* (left panel) and *CD81* (right panel). **D.** True percentile rank of the top 10 predicted high efficiency guides for *CD58* and *CD81*. The red dashed line indicates the top 20th percentile of *CD58*- or *CD81*-targeting guides. **E.** Performance of the ensemble CNN model on a published CasRx guide tiling dataset of three CD transcripts (*CD46*, *CD55*, and *CD71*) in HEK293FT cells (Wessels et al., 2020). Model AUROC, AUPRC, Spearman's correlation coefficient (r_s), and true positive ratio at 0.8 and 0.9 model score cutoffs are shown for each transcript. **F.** True percentile rank of the top 10 predicted guides by our model for three transcripts in a published CasRx guide tiling dataset in HEK293FT cells (Wessels et al., 2020) predicted by the ensemble CNN model. The red dashed lines indicate the top 20th percentile of targeting guides. **G.** Knockdown efficiency of the predicted 5 top scoring guides and 5 low scoring guides for two transcripts (*CD59* and *CD146*) measured by flow cytometry in Hela, U2OS, and A375 cells. Heat map color indicates the mean knockdown efficiency for each guide across n = 3 biological replicates. The top scoring guides and low scoring guides were significantly different at P<0.0001 for Hela, U2OS and A375 cells based on Welch's t test.

Figure 4: Discovery of DjCas13d, a high-efficiency RNA targeting enzyme with minimal cellular toxicity in human cells

A. Phylogenetic tree of Cas13 enzymes including the expanded Cas13d subtype clade (yellow). 46 additional Cas13d orthologs were identified through mining of recent metagenomic datasets. The 7 previously identified Cas13d orthologs including CasRx (red) are shown in bold text. The newly discovered ortholog DjCas13d is shown in blue. All the ortholog sequences are provided in Table S7. **B.** Evaluation of the knockdown efficiency of all Cas13d orthologs shown in panel A on an mCherry reporter transcript in HEK293FT. Horizontal green dashed line denotes our selected cutoff of >55% knockdown efficiency; the hits are color-coded for further study. **C.** Evaluation of the knockdown efficiency of the selected 14 Cas13d orthologs on an endogenous transcript, *CD81*, as measured by flow cytometry-based readout of protein abundance. The horizontal green dashed line denotes a 50% knockdown efficiency cutoff. Cas13d enzymes are plotted in order of their protein size on the x-axis (small to large). **D-E.** Evaluation of cell viability (panel **D**) and knockdown efficiency (panel **E**) of cells expressing each of the top seven most efficient Cas13d orthologs in H1 hESCs along with a *CD81*-targeting guide. The horizontal green dashed line in panel E denotes an 80% knockdown efficiency cutoff. Orthologs are ordered by their size and color-coded as in panel C. Values are shown as mean ± SEM for n = 3 replicates. **F.** Evaluation of cellular viability (y axis) and knockdown efficiency (x axis) of DjCas13d and CasRx across three transcripts in the hESC line, H1. Three top guides were picked for each transcript based on the CNN model score. Each dot on the scatter plot represents one guide's survival rate and knockdown (mean for n = 3 replicates). The dots are colored by the effector used (CasRx: red, DjCas13d: blue), and the color gradients denote the expression level of the target transcript relative to *GAPDH* (log2 relative expression) in the hESC line H1 based on qPCR. The dashed box denotes guides with >80% survival rate and >60% knockdown. 89% of DjCas13d guides are within the box while only 11% of CasRx guides are within the box. **G.** Evaluation of cellular viability (y axis) and knockdown efficiency (x axis) of DjCas13d, CasRx, and Cas7-11 across nine transcripts of different expression levels in HEK293FT using the same spacer sequences across all three enzymes. Three top guides were picked for each transcript based on the CNN model score. Each dot on the scatter plot represents one guide's survival rate and knockdown (mean for n = 3 replicates). The dots are colored by the effector used (CasRx: red, DjCas13d: blue, Cas7-11: grey), and the color gradients denote the expression level (TPMs (transcript per million), log2(TPM+1)) of the target transcript. As in panel F, the dashed box denotes guides with >80% survival rate and >60% knockdown. 84% of DjCas13d guides are within the box while 32% of CasRx guides and 0 Cas7-11 guides are within the box. **H.** Knockdown efficiency of DjCas13d paired with 3 top scoring guides and 3 low scoring guides from the CNN model prediction on nine transcripts of different expression levels in H1 hESCs. Heat map color indicates the mean extent of knockdown for each guide across n = 3 biological replicates. The top scoring guides and low scoring guides were significantly different at P<0.0001 based on Welch's t test. **I.** Knockdown efficiency of DjCas13d paired with 5 top scoring guides and 5 low scoring guides from the CNN model prediction on two transcripts (*CD59* and *CD146*) in Hela and U2OS cells. Heat map color indicates the mean knockdown efficiency for each guide across n = 3 biological replicates. The sets of top scoring guides and low scoring guides were significantly different at P<0.0001 in Hela and P<0.001 in U2OS based on Welch's t test.

Figure 5: DjCas13d exhibits high transcriptome-wide specificity

A. Volcano plots of differential transcript levels between targeting guide conditions and non-targeting (NT) guide control for CasRx (top) and DjCas13d (bottom) in HEK293FT cells using two top-scoring guides for each target transcript (*CD81* (medium expression level)*, FTH1* (high expression level), and *ACTG1* (high expression level)). Red dots denote significantly affected transcripts with adjusted p value < 0.1 and beta value > |0.5|. Green dots denote target transcript isoforms, with darker green dots denoting the most abundant target transcript isoform, and lighter green dots denoting other significantly changed target transcript isoforms. N=3 biological replicates. **B.** Volcano plots of differential transcript levels between targeting guide conditions and non-targeting (NT) guide control for CasRx (top) and DjCas13d (bottom) in hESC (H1) cells with two top-scoring guides for each target transcript (*TFRC* (low expression), *CD81* (medium expression) and *CD24* (high expression)). Red dots denote significantly affected transcripts with adjusted p value < 0.1 and beta value > |0.5|. Green dots denote target transcript isoforms, with darker green dots denoting the most abundant target transcript isoform, and lighter green dots denoting other significantly changed target transcript isoforms. N=3 biological replicates.

Figure 6: DjCas13d enables toxicity-free RNA perturbation in various sensitive cell types
DjCas13d-mediated RNA targeting in **A:** hESC-derived neuronal progenitor cells (NPCs); **B:**
hESC-derived hematopoietic progenitor cells (HPCs); **C:** hESC-derived neurons. Left panel,
schematic of the experimental workflow. Right panel, scatter plot of cellular viability (y axis) and
knockdown efficiency (x axis) across five transcripts of different expression levels in NPCs and
two transcripts of different expression levels in HPCs and neurons. Each dot on the scatter plot
represents one guide's survival rate and knockdown (mean ± SEM for n = 3 replicates). The
dots are colored by the target transcript listed in the legend. The target transcripts are ranked by
expression levels (high to low). The dots are colored by target transcripts. The black dashed line
indicates a survival rate of 1.0 relative to the average of NT guides, and the shaded box
indicates the SEM of the survival rate for NT guides.

Data and Code Availability

The model is freely accessible at http://RNAtargeting.org. The CasRx screen data and code for this manuscript is available on Github https://github.com/jingyi7777/CasRx_guide_efficiency. The RNAseq data is available at the NCBI Sequence Read Archive (SRA): PRJNA857683.

Acknowledgments

We thank the Konermann laboratory and Hsu laboratory for support and advice; A. Pawluk for help with the manuscript; A. Kundaje and J. Zou for advice on deep learning models; J. Zou for the recommendation of the LinearFold package; W. Zhuk for building the initial deep learning model architecture; HK. Wayment-Steele for advice on RNA structure and energy prediction; A. Shrikumar for advice on the implementation and application of TF-MoDISco to our analysis; The Salk Institute NGS core and the Stanford Shared FACS facility for their support; and B. Hsu for helping build the CasRx guide design website. K.F. was supported by a UCSD Eureka! Scholarship for this work. S.K. is a Hanna Gray Fellow of the Howard Hughes Medical Institute, a Chan Zuckerberg Biohub Investigator, and an Arc Institute Core Investigator. P.D.H. is supported by the NIH (DP5 OD021369, R01 GM131073, R01 GM132465), DARPA, Emergent Ventures, the Shurl and Kay Curci Foundation, and the Rainwater Charitable Foundation and the Arc Institute.

Author Contributions

S.K. and P.D.H. conceived this study and supervised the design and analysis of all experiments. J.W. and H.K. built the computational models, performed feature engineering, and implemented model interpretation. S.K. and J.W. analyzed the NGS data from the screens and calculated secondary features. J.W. performed the validation screen and individual guide testing in cancer cells. J.W. created the Cas13d guide efficiency prediction tool and performed model comparison. S.K. and P.D.H. computationally identified novel Cas13d orthologs. P.L. and E.W. cloned all Cas13d orthologs and tested in HEK293FT. J.W. tested the top Cas13d orthologs in stem cells. J.W., S.B., E.G., H.S., and E.K. cloned individual CasRx and DjCas13d guides. J.W., S.B., E.G., H.S., and E.K. performed individual guide testing in HEK293FT and stem cells. J.W. performed RNA-seq experiments and analyzed the data with C. V. D.. J.W., H.S., E.K., S.B., and E.G. performed RNA knockdown experiments in stem cell-differentiated neuronal progenitor cells, neurons, and hematopoietic progenitor cells. S.C. analyzed Cas13d ortholog sequences. M.D. performed computational mining of additional Cas13 sequences and built the Cas13 phylogenetic tree. P.L. S.K., and P.D.H. adapted CasRx for high-throughput screening. S.K., K.F., P.L., and P.D.H. performed the cell proliferation screen. J.W., S.K., and P.D.H. wrote the manuscript with input from all authors.

Competing Interest Statement

P.D.H. is a cofounder of Spotlight Therapeutics and Moment Biosciences and serves on their boards of directors and scientific advisory boards, and is a scientific advisory board member to

819   Arbor Biotechnologies, Vial Health, and Serotiny. P.D.H. and S.K. are inventors on patents
820   relating to CRISPR technologies, including DjCas13d.

Methods

Plasmid design

For the CasRx expression vector, we designed a piggyBac-based all-in-one plasmid containing
the CasRx effector, piggyBac transposase, and antibiotic selection cassette: PB_EF1a-CasRx-
msfGFP-2A-Blast. The CasRx effector is fused to msfGFP at the C terminus and under the
control of a constitutive EF1a promoter. A nuclear localization signal SV40 NLS was added to
both the N and C terminus of CasRx-msfGFP. The antibiotic selection cassette, blasticidin S
deaminase, is linked with CasRx-msfGFP via a P2A self-cleaving peptide.
829
For the CasRx guide cloning vector, we designed a lentiviral vector: hU6-(CasRx DR)-EF1a-
Puro-WPRE. The CasRx DR is a 36-base direct repeat
(CAAGTAAACCCCTACCAACTGGTCGGGGTTTGAAAC) for CasRx pre-gRNA (Konermann et
al., 2018). The 30 nt guide spacer sequence is cloned into the vector through Gibson cloning
using two BsmBI cleavage sites. For individual guide truncation and individual guide validation
experiments, we designed a piggyBac-based all-in-one plasmid containing the CasRx effector,
guide DR, piggyBac transposase, and antibiotic selection cassette: hU6-(CasRx DR)-TRE-
CasRx-msfGFP-EF1a-rtTA-2A-Puro-CMV-transposase.
838

Guide library design

For the survival screen, we selected 55 essential genes from the intersection of the essential
hits in three previous survival screens performed in K562 cells (Hart et al., 2015; Horlbeck et al.,
2016; Luo et al., 2008). We selected the major transcript isoform of these genes from the
Refseq database and designed guides that tile these transcripts with single nucleotide
resolution. A total of 127,071 targeting guides were generated for the 55 essential transcripts. In
addition, we designed 14111 guides tiling 5 non-essential control transcripts (*CTCFL, SAGE1,*
TLX1, DTX2, OR2C3). Along with 3563 non-targeting guides, we constructed a pooled library of
144745 guides.
848
849   For the validation screen on cell surface markers, 3218 guides were designed that tiled *CD58*
850   transcripts (NM_001779.3, NM_001144822.2) and *CD81* transcripts (NM_004356.4,
851   NM_001297649.2) with single nucleotide resolution. The targeting guides were pooled with
852   1186 non-targeting guides to create the final library.
853

Guide library synthesis, cloning, and library amplification

For each guide spacer sequence in the guide library, we added a constant left overhang
("AACCCCTACCAACTGGTCGGGGTTTGAAAC") and a right overhang
("TTTTTTTTGAATTCAAGCTTGGCGTAACTAGA") to facilitate cloning. The resulting libraries
were synthesized as oligo pools by Twist Biosciences, and then PCR amplified using the primer
pair: Lib_F
("TCTTGTGGAAAGGACGAAACACCGCAAGTAAACCCCTACCAACTGGTCGGGGTTTG") and

Lib_R
("AGAGCTAGCCAGACGTGTGCTCTTCCGATCNNNNNNNNNNTCTAGTTACGCCAAGCTTGA
ATTC") (**Table S1**). The PCR reaction was performed using NEBNext High Fidelity PCR
Master Mix (NEB, catalog no. M0541L) for 20 cycles. The amplified library was gel-purified and
cloned into the BsmBI digested guide cloning vector (hU6-(CasRx DR)-EF1a-Puro-WPRE)
through Gibson assembly. The cloned guide library was then purified and concentrated by
isopropanol precipitation.
868
For guide library amplification, the library plasmid was electroporated to Endura
electrocompetent *E. coli* cells (Lucigen, catalog no. 60242-2) at 50–100 ng/ul. After
electroporation, cells were recovered in LB medium for 1h, and then plated on LB agar plates
with 100 ug/mL carbenicillin at 37°C for 12-14h. The colonies were then harvested at a
coverage of > 500 colonies per guide. The amplified guide library plasmid was extracted using
the Macherey-Nagel NucleoBond Xtra Maxi EF Kit (Macherey-Nagel, catalog no. 740424.10).
To determine guide RNA representation, we PCR amplified the guide region using customized
NGS primers containing Illumina adaptor sequences (**Table S1**). NextSeq sequencing was
performed to determine guide RNA representation in the guide library. We verified that the
library had >87% perfectly matching guides, <0.5% undetected guides, and a skew ratio (90th
percentile:10th percentile read number) of less than 10.
880
Lentivirus production
To produce lentivirus for the guide library, HEK293FT cells, purchased from Thermo Fisher (Cat
R70007) were grown in DMEM supplemented with 10% FBS (D10 media) at 37 °C with 5%
CO2. Cells were passaged at a ratio of 1:2 using TrypLE (Gibco) and seeded 20–24 h before
transfection at 1.8×10^7 cells per T225 flask. For lentiviral plasmid transfection, the guide library
plasmid was mixed with psPAX2 (Addgene, catalog no. 12260) and pMD2.G (Addgene, catalog
no. 12259) in Opti-MEM, and transfected to HEK293FT using Lipofectamine 2000 (Thermo
Fisher, catalog no. 11668027) and PLUS reagent (Thermo Fisher, catalog no. 11514015).
Medium was replaced 4 hours after transfection with fresh, prewarmed D10 medium. Two days
after the start of lentiviral transfection, the supernatant from the HEK293FT cells was harvested
and filtered using a 0.45um Stericup filter. The lentiviral titer was determined through spinfection
on K562 cells prior to the screen.
893
Cell culture and CasRx cell line generation
K562 cells were purchased from ATCC (CCL-243), and cultured in RPMI 1640 medium with
GlutaMAX™ supplement (Thermo Fisher, catalog no.61870036), 10% FBS, and Penicillin-
Streptomycin at 37 °C with 5% CO2. To generate a stable CasRx-expressing K562 cell line, we
transfected K562 cells with the piggyBac-based all-in-one CasRx expression vector (PB_EF1a-
CasRx-msfGFP-2A-Blast) using Lipofectamine 3000 Transfection Reagent (Thermo Fisher,
catalog no. L3000001). Two days after transfection, we selected the cells with 10 µg/ml
blasticidin S (Thermo Fisher, catalog no. A1113903). After selection for 1-2 weeks, we checked
the percentage of CasRx-expressing cells using flow cytometry and confirmed that more than
95% of cells expressed CasRx-GFP.
904

Survival screen

The guide library for the survival screen was lentivirally transduced at MOI=0.2 by spinfection into the stable CasRx-expressing K562 cell line. We ensured the guide library had a coverage of >1000 cells per guide. Two days after transduction, cells were selected with 1 µg/ml puromycin to ensure guide expression and further cultured for 14 days. Cells were harvested at day 14 (end of the screen), and the genomic DNA was extracted using Zymo Research Quick-gDNA MidiPrep (Zymo Research, cat. no. D4075). The guide region was PCR amplified using customized NGS primers containing Illumina adaptor sequences. The resulting PCR products were gel purified and quantified with Nanodrop and Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific, cat. no. Q32851). Pooled guide libraries were sequenced on Illumina NextSeq, with 80 cycles of read 1 (forward) and 8 cycles of index 1. Three biological replicates were performed for the survival screen.

Validation screen on *CD58* and *CD81*

The guide library for the validation screen on *CD81* and *CD58* was lentivirally transduced at MOI=0.1 by spinfection into the stable CasRx-expressing K562 cell line. 45 million cells per biological replicate were transduced to ensure coverage of >1000 cells per guide. After spinfection, cells were selected with 1 µg/ml puromycin and further cultured for 10 days. At the end of the screen, cells were divided into two pools and stained with *CD58* antibody (BD Biosciences, catalog no. 564363) or *CD81* antibody (BD Biosciences, catalog no. 561958) and analyzed using FACSAria II (**Figure S11**). Following calibration with unstained controls, each cell pool was sorted into four bins based on target gene expression level indicated by antibody-conjugated fluorescence intensity. Specifically, cells were first gated by forward and side scatter to select for live, single cells. Next, cells were gated on GFP to select for CasRx-expressing cells. This final population was sorted into four bins based on the intensity of *CD58* or *CD81* antibody-conjugated fluorescence intensity. As high efficiency guides were defined as the top 20% for each gene, we set the bin with the lowest target gene expression (bin 1) at 7-8%, which is equal to the fraction of the target gene's high efficiency guide number in the whole library: 1600*0.2/4401. The rest of the population was equally divided into three bins of the same size (~30%). The genomic DNA for cells in each bin was extracted and sequenced as in the survival screening. Four biological replicates were performed for the validation screen.

Data preprocessing and definition of high efficiency guides

For each guide RNA we calculated the fraction in the day 14 guide pool and the input library pool. Guide efficiency was evaluated by the ratio of guide percentage in the day 14 pool to the input pool (**Table S3**). Guides targeting each transcript were ranked based on their average ratio across the three replicates, and we defined high efficiency guides for each transcript, taking into account three parameters: 1) the top 20% guides per transcript; 2) no essential off-targets predicted by BLAST (see the 'Off-target filtering' section below and **Figure S1D** for details); and 3) with a d14/input ratio lower than 0.75 (the ratio at 5th percentile of the guides for targeting control genes). Guides targeting the transcript *RPS19BP1* were excluded because they clustered with non-essential controls (most guides were not effectively depleted in the screen).

For the validation screen, we first filtered guides with less than 200 counts in all *CD58* bins and
CD81 bins. Less than 1.5% of guides were removed by this filter. We then calculated each
guide's distribution across the 4 bins and used the ratio of guide percentage in bin 1 (greatest
knockdown) to the sum of its percentage in bins 1 and 4 (least knockdown) for evaluation of
guide efficiency. We then ranked guides within each gene based on their average ratio of the
four replicates, and we defined the top 20% guides for each gene as high efficiency guides.
955
Off-target filtering
We performed BLAST to identify potential off target matches for our guides. As the first 24
nucleotides from the 5' end of the CasRx guides were shown to be most indicative of guide
targeting ability (**Figure S2C**), we took the first 24 nucleotides of each guide as BLAST input.
BLAST was performed using a generous E value of 1 (e=1) against the Gencode V33 database.
BLAST results were parsed and off target genes were identified as those with up to three
mismatches to the guide input. To check the essentiality of the off-target genes, we made an
essential gene list by combining the essential gene hits from the three previous survival screens
in K562 cells and we compared the off-target genes with the essential gene list. Guides with
predicted off-targets in essential genes were filtered, as we reasoned they may interfere with the
interpretation of our survival screen readout. For our survival screening, 6790 guides were
filtered and 120281 guides remained for further analysis (**Figure S1D**, Table S4). For the
validation screen, the filtered dataset is provided in Table S5.
969
Analysis of the positional distribution of high efficiency guides
For each transcript, we calculated the number of high efficiency guides overlapping each
position on the transcript, and plotted the results using a heatmap. We further summarized the
distribution of high efficiency guides across all transcripts and positions with a histogram. In
theory, a particular nucleotide position would have at most 30 guides covering it, so the number
of efficient guides ranges from 0 to 30 for each position. We compared the results with a
randomly sampled distribution, which is simulated using 100 random samplings of 20% of the
guides in the library. In theory, the randomly sampled distribution would show a peak at 6
(30*20%), which agrees with our simulation results.
979
Data splits
For model hyperparameter tuning and evaluation, we split our 54 essential transcripts into 9
folds, each containing a unique and non-overlapping set of 6 test transcripts. The 54 transcripts
were distributed evenly across the 9 folds according to their high efficiency guide percent to
make the 9-fold split balanced. Using the predefined transcript splits, we performed 9-fold cross-
validation to tune model hyperparameters and compare prediction accuracy between models.
986
Feature calculation and model inputs
For the sequence input, each 30 nt guide spacer was one-hot encoded into four binary vectors
of length 30 to represent the nucleotide identity at each position.

To predict guide unfolding energy, we used LinearFold, a linear-time RNA secondary structure
prediction algorithm (Huang et al., 2019) on the full-length guide sequence (36nt DR +30nt

spacer). We started with the default parameters and the CONTRAfold v2.0 model (Do et al.,
2006; Lorenz et al., 2011; Wayment-Steele et al., 2020) provided by the LinearFold software at
https://github.com/LinearFold/LinearFold. We subtracted the predicted MFE (minimum free
energy) with the baseline energy (MFE of the unstructured guide with the 30 nt spacer unfolded)
to calculate guide unfolding energy. We also tested the Vienna RNAfold model in LinearFold as
a comparison. To determine whether using the ensemble guide unfolding energy instead of
MFE could improve model prediction, we further tested three RNA structure prediction
algorithms (Contrafold2, Eternafold, Vienna) wrapped by Arnie
(https://github.com/DasLab/arnie) to calculate the ensemble guide unfolding energy with the
partition function (Do et al., 2006; Lorenz et al., 2011; Wayment-Steele et al., 2020). For the
Vienna package, we tested different temperature(T) settings: 37°C , 60 °C, and 70 °C. In our
final model, we used the guide unfolding energy calculated by LinearFold's default CONTRAfold
v2.0 model as it improved model prediction accuracy to the greatest extent.

To calculate target unfolding energy, we first used LinearFold's CONTRAfold v2.0 model to
predict MFE of the native local target region using the local target sequence. We then predicted
MFE of the guide unwound local target region by supplying the algorithm with the constraint that
the 30 nt guide-binding site is unpaired. (This can be achieved by feeding in an additional
constraint structure with the guide-binding site annotated with "."). We then subtracted the
former MFE (MFE of the native target region) by the latter (MFE of the guide unwound target
region) to estimate local target unfolding energy. The local target region was defined as the 30
nt guide-binding site with 15 nt flanking sequence on both sides. Flanking sequences of different
lengths were compared, and the length 15 was chosen for the final model as it improved model
prediction accuracy to the greatest extent.
1015
1016 To calculate the percentage of isoforms targeted by each guide, we obtained all transcript
1017 isoforms for each gene from the Refseq database and evaluated the percentage of isoforms
1018 matched for each 30nt guide target (using perfect matches).
1019
To calculate the three position flags, we obtained Refseq's annotations of the 5′ UTR, CDS, or
3′ UTR region for our target transcripts. Guides that target the 5′ UTR, CDS, or 3′ UTR region
have a flag value of 1 for that correspondent feature, and 0 for the other two flag features. To
calculate the three position floats (5′ UTR position,CDS position,3′ UTR position), we calculated
the relative position of the guide target site in the 5′ UTR, CDS, or 3′ UTR region. Guides
located out of the region have a flag value of 0 for the correspondent feature.
1026
Model architecture
Sequence-only models
For linear models and ensemble models, the one-hot encoded guide sequence was flattened
and converted to 30*4= 120 flag features. The features are then fed into the models to generate
the output. For the CNN model, the one-hot encoded guide was treated as a 4-channel image,
and a few 1D convolutional layers were applied to generate a feature map, which was flattened
and passed to a dense layer to generate the final output. For the biLSTM model, the guide
sequence was treated as a sentence with four characters, and two LSTMs, each processing the

input sequence in one direction (forward or backward), were applied to generate sequence
representations. The resulting vectors were merged, flattened, and passed to a dense layer to
generate the final output.
1038
Full model with secondary features
For the CNN model with secondary features, the one-hot encoded guide was passed to a few
convolutional layers as in the sequence-only model. The output from the CNN layers was
flattened and concatenated with the normalized secondary features. The concatenated feature
vector was sequentially passed to a dense layer, a recurrent dense layer and a final dense layer
of 1 unit to generate the output. All dense layers use leaky ReLU as the activation function. The
CNN layer kernel size, unit number, layer number and the dense layer unit number were defined
after hyperparameter tuning.
For the Gradient-boosted classification tree, the one-hot encoded guide sequence was flattened
and converted to 30*4= 120 flag features. The sequence features are concatenated with the
normalized secondary features, and then fed into the model to generate output.
1050
Model training, hyperparameter tuning and evaluation
All models were trained to solve a binary classification task – predicting high efficiency guides,
and the model output is the probability that a guide is a high efficiency guide.
The linear models and ensemble models were trained in scikit-learn 0.24 and the deep learning
models (LSTM and CNN) were trained in TensorFlow 2.3.1. For the deep learning models, we
used binary cross-entropy as the loss function and applied the Adam optimizer for model
training. Early stopping was used to prevent model overfitting.
For all models, the prediction accuracy is evaluated by AUROC (Area Under the Receiver
Operating Characteristic curve) and AUPRC (The Area Under Precision-Recall Curve).
1060
1061 To tune hyperparameters and evaluate model performance, we used 9-fold cross-validation
1062 over the hyperparameter space. For linear models and ensemble models, we used the
1063 "GridSearchCV" function in scikit-learn to perform a grid search over the hyperparameter set.
1064 For deep learning models, we used the Hyperband tuner in TensorFlow to select top models
1065 quickly by filtering poor models during training.
1066
The hyperparameter sets for all models are listed below:
- logistic regression with L1 regularization: regularization strength - logarithmic in $(10^{-5}, 10^5))$

- logistic regression with L2 regularization: regularization strength - logarithmic in $(10^{-5}, 10^5))$

- logistic regression with elastic net regularization: regularization strength - logarithmic in $(10^{-4},$
$10^4))$, L1 ratio - equally spaced from 0.1 to 1.
- Gradient-boosted classification trees: number of trees –
[100,200,400,800,1000,1200,1500,1800,2000], maximum depth of a tree – [2,4,8], the number of
features to consider when looking for the best split - all, sqrt(n_features), log2(n_features).
- Random forest (RF): number of trees – [100,200,400,800,1000,1200,1500,1800,2000], number of
features to consider when looking for the best split - all, sqrt(n_features), log2(n_features).
- Long short-term memory recurrent neural network (LSTM): LSTM units - [16, 32,64,128], dense
layer units – [8, 16, 32], recurrent dense layer number – [0,1,2,3], dropout rate - [0.0, 0.1, 0.25]

●   Convolutional neural network (CNN): CNN layer kernel size – [3,4,5], CNN units- [8,16,32,64],
CNN layer number – [3,4,5], dense layer units - [8,16,32,64], recurrent dense layer number –
[0,1,2,3]
1082
1083 For all models, we chose the hyperparameter set with the highest average AUROC across all
1084 test sets in the 9-fold splits, and evaluated the final model performance using both the average
1085 AUROC and average AUPRC across test sets.
1086
Secondary feature selection
For the CNN model, we added each secondary feature individually to guide sequence features
and calculated the change in model performance. We selected features that successfully
improved model performance, and added these features sequentially upon guide sequence
features to check feature redundancy. We also tried removing individual features from the final
model to confirm the necessity of the features.
For the Gradient-boosted tree, besides the above methods, we also used Boruta, an all-relevant
feature selection method that aims to find all features useful for prediction (Kursa et al., 2010).
We implemented it using BorutaPy, the Python implementation of Boruta
(https://github.com/scikit-learn-contrib/boruta_py) on our Gradient-boosted tree.
1097
Model interpretation and feature contributions
For the CNN model, we applied "Integrated Gradients" (IG) to investigate feature contributions
in the model. "Integrated Gradients" is an attribution method that evaluates feature importance
by integrating the gradient of output to input features along the straightline path from the
baseline input to the actual input value (Sundararajan et al., 2017). Due to the non-linearity of
the deep learning model, we applied "Integrated Gradients" to the best-performing individual
CNN model on CD genes rather than the ensemble model. To compute integrated gradients, we
first set all-zero baselines for the sequence input, position flags and position floats, and used
average baselines for other features. Next, we generated a linear interpolation between the
baselines and the inputs using 50 steps. We then computed gradients using the
"tf.GradientTape" function in TensorFlow for the interpolated points, and approximated the
gradients integral with the trapezoidal rule. To evaluate the relative importance of each position
on the guide, we averaged the absolute integrated gradient values at each position across all
test sequences. To evaluate the contribution of each nucleotide at each position, we averaged
the integrated gradients for that nucleotide across all test sequences.
For the Gradient-boosted tree, we applied SHAP (SHapley Additive exPlanations) to investigate
feature contributions in the model. SHAP is a game theoretic approach that estimates how each
feature contributes to the model output by providing the SHAP value for each input feature
(Lundberg et al., 2020). We implemented the SHAP package from
https://github.com/slundberg/shap, and applied it to our Gradient-boosted tree. To evaluate the
relative importance of each position on the guide, we averaged the SHAP values at each
position across test sequences. To evaluate the contribution of each nucleotide at each position,
we averaged the SHAP values for that nucleotide across test sequences.
1121
Cas13a guide sequence contribution to guide efficiency

We analyzed three Cas13a guide efficiency datasets: 1) the Luciferase knockdown dataset
containing 186 LwaCas13a guides for *Gaussia* luciferase (Gluc) and 93 guides for *Cypridina*
Luciferase (Cluc) (Abudayyeh et al., 2017); 2) the endogenous gene knockdown dataset
containing 93 LwaCas13a guides for each of *KRAS*, *PPIB* and *MALAT1* (Abudayyeh et al.,
2017); and 3) the ADAPT dataset containing 85 perfect match LwaCas13a guides for virus
detection (Metsky et al., 2022). We calculated the Pearson correlation between each nucleotide
at each position with guide efficiency to evaluate the sequence contribution.
1130

Motif discovery
For motif discovery, we used TF-MoDISco (Transcription Factor Motif Discovery from
Importance Scores), an algorithm that discovers motifs by clustering important regions in
sequences using per-base importance scores (Shrikumar et al., 2018). We implemented TF-
MoDISco from https://github.com/kundajelab/tfmodisco using the integrated gradients of all high
efficiency guides in our training data as input. We ran TF-MoDISco with a sliding window size of
7 and a flank length of 2. For final motif processing, we trimmed the clustered motifs to a
window size of 6, added an initial flank length of 2 and a final flank length of 3 to get the final
motifs. The top 5 active motifs are picked and aligned to the 30 nt spacer according to the mode
position of sequences in each motif.
1141

Nmer analysis
To identify enriched or depleted positional nmers, we divided our survival screen data to 9 folds
as in the model training workflow and calculated the ratio of all possible positional nmers'
percentage in high efficiency guides to non-high efficiency guides in the training set and test set,
respectively, for each fold. We identified enriched (or depleted) nmers based on their ratio in the
training set with a predefined ratio cut-off. We selected the nmers identified as enriched (or
depleted) across all folds, and ranked them by their average percent in high efficiency guides in
the test sets across all folds. The initial ratio cut-off is set as 2 for enriched nmers and 0.5 for
depleted nmers. The cut-off is adjusted during the nmer identification process so that the
percent of guides with enriched nmers are ~20% and the percent of guides with depleted nmers
are ~40%. We mainly focused on 3-mers and 4-mers in this paper.
1153

Final model and model testing on the validation screens
We chose the CNN model as our final model after hyperparameter tuning and model
comparison. We re-trained the model using all of the survival screen data. To prevent
overfitting, we split out a validation set during model training as in the previous 9-fold cross-
validation split. We built 9 individual models using different validation sets from the 9-fold split of
essential transcripts, and we compared their performance on the two cell surface markers,
CD58 and *CD81*. We further built an ensemble model that averaged the prediction of all the
individual models. We found that the ensemble model outperformed all individual models on the
two CD genes, so we set the ensemble CNN model as our final model. As a comparison, we
also retrained the best non-deep learning model, the Gradient-boosted tree (GBT), using all of
the survival screen data. We tested the model on the two CD genes and evaluated model
performance using AUROC and AUPRC.
1166

Model comparison with Wessels et al. model and DeepCas13
We tested the performance of the Random forest model from Wessels et al. on our CD genes
and essential genes using the web server https://cas13design.nygenome.org (Wessels et al.
2020; Guo et al. 2021). We evaluated the model performance using AUROC, AUPRC,
Spearman's correlation coefficient, r_s and true positive ratios at 0.8 and 0.9 model score cutoffs.
As the Random forest model is designed for 23 nt long guides, we extended the guides from
their model output to 30 nt (extends toward the 3′ end) to be in accordance with our screen
data. For comparison, we retrieved the CasRx guide tiling screen dataset on three genes,
CD46, *CD55*, and *CD71*, from Wessels et al. and tested our model's performance. We adjusted
the guide length to 23 nt in our model to be in accordance with their screen data, and we set the
top 20% guides for each gene as "high efficiency guides". The model performance was also
evaluated by AUROC, AUPRC, Spearman's correlation coefficient, r_s and true positive ratios at
0.8 and 0.9 model score cutoffs.
1180
1181 We tested the performance of DeepCas13 (Cheng et al. 2023) on our CD genes using the web
1182 server http://deepcas13.weililab.org. We evaluated the model performance using AUROC,
1183 AUPRC, Spearman's correlation coefficient, r_s and true positive ratios at 0.8 and 0.9 model
1184 score cutoffs.
1185
Cas13d guide efficiency prediction tool and website
A website-based Cas13d guide efficiency prediction tool was developed using our CNN model
for Cas13d guide design across model organism transcriptomes and custom RNA sequences.
For model organism Cas13d guide design, we precomputed the Cas13d guide efficiency for all
coding and non-coding genes of each model organism. Briefly, reference transcriptome
sequences and annotations were obtained from the UCSC Table Browser (Karolchik et al.,
2004) with the NCBI RefSeq track. All possible 30 nt Cas13d guide spacers were extracted from
the transcriptome sequences with single nucleotide resolution. Secondary features were
calculated for each guide as described in the '**Feature calculation and model inputs'** section
above. The final CNN model was applied to all guides for prediction of their efficiency, and the
guides were ranked within each gene based on the model prediction scores.
1197
For custom sequence guide design, all possible 30 nt Cas13d guide spacers are extracted from
the input custom RNA sequences with single nucleotide resolution. Guide unfolding energy and
target unfolding energy are calculated as described in the '**Feature calculation and model**
inputs' section above. A CNN model that uses guide sequence, guide unfolding energy and
target unfolding energy as inputs, trained on the survival screen dataset, is applied to the
custom sequence guides for prediction of their efficiency. Guides are ranked based on the
model prediction scores.
1205
1206 The Cas13d guide efficiency prediction tool is freely available on a public, user-friendly website:
1207 https://www.RNAtargeting.org.
1208
Computational identification of novel Cas13d orthologs through metagenomic database
mining

We applied our previously described pipeline for novel CRISPR effector discovery (Konermann
et al., 2018) to incompletely assembled metagenomic contigs in addition to whole genome,
chromosome, and scaffold-level prokaryotic and metagenomic sample assemblies from the
NCBI Genome database (https://www.ncbi.nlm.nih.gov/), the Gigadb repository
(http://gigadb.org/), as well as the JGI Genome portal (https://genome.jgi.doe.gov/portal/).
Putative effectors encoded near identified CRISPR arrays (<kb distance) were assigned to
previously identified Cas13 families via tBLASTn analysis, where a bit score of at least 60 to any
prior Cas13 subfamily member was required for cluster assignment. As a second round of
discovery independent of CRISPR array identification, tBLASTn was performed on all original
and predicted Cas13d effectors from the first round against all public metagenome whole
genome shotgun sequences without predicted open reading frames (ORFs) from all three
sources listed above. New full-length homologs and homologous fragments were aligned using
Clustal Omega and clustered using PhyML 3.2 (Guindon et al., 2010). All the Cas13d ortholog
sequences are provided in **Table S7**.
1225

Construction of Cas13 phylogenetic tree
A custom sequence database of bacterial isolate and metagenomic sequences was constructed
by aggregating publicly available sequence database, including NCBI, UHGG (Almeida et al.,
2021), JGI IMG (I.-M. A. Chen et al., 2021), the Gut Phage Database (Camarillo-Guerrero et al.,
2021), the Human Gastrointestinal Bacteria Genome Collection (Forster et al., 2019), MGnify
(Mitchell et al., 2020), Youngblut et al animal gut metagenomes (Youngblut et al., 2020),
MGRAST (Meyer et al., 2008), and Tara Oceans samples (Sunagawa et al., 2015). Cas13
sequences from other Cas13 families were identified by searching representative members of
each clade (Cas13a/b/bt/c/x/y) against a collection of protein representatives (clustered at 30%
identity) derived from the custom sequence database using hmmsearch from the hmmer
package (*HMMER*, n.d.). Selected Cas13a, Cas13b, Cas13c, Cas13d representatives were
LbuCas13a, BzoCas13b, AspCas13c, and CasRx respectively. The Cas13bt representative was
collected from (Kannan et al., 2022), and the Cas13X and Cas13Y representatives were
collected from (Xu et al., 2021). All hits that met E < 1e-6 and were 75%-125% the length of the
representative sequence were retained. Sequences were assigned to the best matching
representative. Sequences were then clustered at the 50% identity level along 80% of both
sequences using the mmseqs package (Steinegger & Söding, 2017). Sequences were then
aligned using the MAFFT algorithm mafft-linsi (Katoh et al., 2002). PhyML was used to generate
phylogenetic trees with default parameters (Guindon et al., 2010). Trees were visualized using
the ggtree package in R (Yu, 2020).
1246

Cloning of Cas13d orthologs and Cas7-11
For initial testing and efficiency screening, human codon optimized Cas13d sequences, flanked
by two nuclear localization or export sequences, were cloned into a backbone derived from
pXR001: EF1a-CasRx-2A-EGFP (Addgene #109049) to replace the CasRx coding sequence.
Guide sequences targeting mCherry or *CD81* were cloned into a backbone derived from
pXR003: CasRx gRNA cloning backbone (Addgene #109053) with 5′ full-length direct repeat
(DR) sequences for each Cas13d ortholog. For testing the seven high efficiency Cas13d
orthologs in stem cells, the Cas13d coding sequences and respective mature DR guide scaffold

sites were cloned into the inducible piggyBac-based all-in-one plasmid containing the Cas13d
effector, guide DR, piggyBac transposase, and antibiotic selection cassette: hU6-DR-TRE-
Cas13d-T2A-msfGFP-EF1a-rtTA-T2A-Puro-CMV-transposase. Human codon optimized
DisCas7-11 protein sequence and the mature DR guide scaffold with golden gate sites were
PCR amplified from Addgene plasmids # 172507 and #172508, a gift from Omar Abudayyeh &
Jonathan Gootenberg, and cloned to the constitutive piggyBac-based all-in-one backbone
plasmid as mentioned before. Guide spacers were position matched to CasRx and DjCas13d's
guide spacers and were cloned into the backbone plasmid using Golden Gate cloning. All
individual guide sequences are provided in **Table S6**.
1264
Cell culture for individual guide testing
HEK293FT cells were purchased from Thermo Fisher (Cat # R70007) and grown in DMEM
supplemented with 10% FBS (D10 media) at 37 °C with 5% CO2. Cells were passaged at a
ratio of 1:2 using TrypLE (Gibco). Hela and A375 cells were gifts from the Howard Chang lab
and Scott Dixon lab, respectively. They were both cultured in DMEM supplemented with 10%
FBS (D10 media) at 37 °C with 5% CO2. Cells were passaged at a ratio of 1:2 using TrypLE
(Gibco). U2OS cells were a gift from the Chang lab and grown in McCoy's 5A (modified)
Medium (Thermo Fisher, catalog no. 11668027) supplemented with 10% FBS at 37 °C with 5%
CO2. Cells were passaged at a ratio of 1:2 using TrypLE (Gibco). Stem cell line H1 were
purchased from WiCell (Cat # WA01). Cells were maintained in mTeSR™ Plus media (Catalog
100-0276, STEMCELL Technologies) on Matrigel-coated 6-well plate and passaged 1:12 with
ReLeSR™ (Catalog # 05872, STEMCELL Technologies) every four days.
1277
Transfection of human cell lines
For initial testing and efficiency screening of Cas13d orthologs, HEK293FT cells were plated at
20,000 cells per well in a 96-well plate, then transfected at >80% confluence with 192 ng
Cas13d-2A-EGFP plasmid, 192 ng of crRNA expression plasmid, and 12 ng of mCherry
expression plasmid using Lipofectamine 2000. Cells were harvested 48 hours after transfection
for flow cytometry analysis of mCherry expression. For CD81 knockdown experiments,
HEK293FT cells were transfected with 200 ng Cas13d-2A-EGFP plasmid and 200 ng guide
RNA expression plasmid using Lipofectamine 2000. Cells were harvested 48 hours after
transfection for staining and flow cytometry analysis of CD81 expression.
1287
For experiments comparing CasRx, DjCas13d, and Cas7-11 in HEK293FT cells, cells were
plated at 16,000 cells per well in a 96-well plate and transfected at > 80% confluence with 100
ng of all-in-one PiggyBac plasmids containing CasRx, DjCas13d, or Cas7-11 using
Lipofectamine 2000 (Life Technologies). Cells were selected with 1 μg/ml puromycin 24h after
transfection. 24 hours after selection, cells were harvested for RNA extraction and downstream
processing.
1294
1295 For individual guide testing in Hela cells, low passage cells were plated at a density of 15,000
1296 cells per well in a 96-well plate and transfected at > 80% confluence with all-in-one PiggyBac
1297 plasmids containing CasRx or DjCas13d using FuGENE® HD Transfection Reagent (E2311,
1298 Promega) according to the manufacturer's protocol. Cells were selected with 1 μg/ml puromycin

1299 and induced with Doxycycline (D3072, Sigma) for CasRx or DjCas13d expression 48h after
1300 transfection. Flow analysis was performed seven days after induction.
1301
1302 For individual guide testing in U2OS cells, low passage cells were plated at a density of 15,000
1303 cells per well in a 96-well plate and transfected at > 80% confluence with all-in-one PiggyBac
1304 plasmids containing CasRx or DjCas13d using ViaFect™ Transfection Reagent (E4981,
1305 Promega) according to the manufacturer's protocol. Cells were selected with 0.75 µg/ml
1306 puromycin and induced with Doxycycline (D3072, Sigma) for CasRx or DjCas13d expression
1307 48h after transfection. Flow analysis was performed seven days after induction.
1308
1309 For individual guide testing in A375 cells, low passage cells were plated at a density of 25,000
1310 cells per well in a 96-well plate and transfected at > 80% confluence with all-in-one PiggyBac
1311 plasmids containing CasRx using TransIT-X2 (MIR 6003, Mirus) according to the
1312 manufacturer's protocol. Cells were selected with 0.5 µg/ml puromycin and induced with
1313 Doxycycline (D3072, Sigma) for CasRx expression 48h after transfection. Flow analysis was
1314 performed seven days after induction.
1315
1316 For enzyme comparison and individual guide testing in H1 cells, low passage cells were
1317 passaged with Accutase (Innovative Cell Technologies) and plated into a Matrigel-coated 96-
1318 well plate with mTESR media containing ROCK inhibitor Y-27632 (10 uM, Abcam) at 30,000
1319 cells per well one day before transfection. On day 1, cells were transfected at > 80% confluence
1320 with all-in-one PiggyBac plasmids containing different Cas13d orthologs using FuGENE® HD
1321 Transfection Reagent (E2311, Promega) according to the manufacturer's protocol. Cells were
1322 selected with 0.5 µg/ml puromycin 48h after transfection. 5-7 days after selection, Cas13d
1323 expression was induced with Doxycycline (D3072, Sigma). Flow cytometry analysis was
1324 performed three days after induction.
1325
1326 For RNAseq experiments in H1 cells, low passage cells were passaged with Accutase
1327 (Innovative Cell Technologies) and plated into Cultrex (R&D Systems 343400502)-coated 96-
1328 well plates with mTESR media containing ROCK inhibitor Y-27632 (10 uM, Abcam) at 25,000
1329 cells per well one day before transfection. On day 1, cells were transfected at > 80% confluence
1330 with all-in-one PiggyBac plasmids containing different Cas13d orthologs using FuGENE® HD
1331 Transfection Reagent (E2311, Promega) according to the manufacturer's protocol. Cells were
1332 split and selected with 0.75 µg/ml puromycin 24h after transfection. Puromycin concentration
1333 was increased to 1ug/ml the next day. 72h after transfection, cells were harvested for RNA
1334 extraction and downstream processing.
1335
1336
1337 **Staining and flow cytometry**
1338 For cell surface protein staining, cells were harvested and dissociated with TrypLE, followed by
1339 two washes in cold FACS buffer (DPBS + 2 mM EDTA + 0.02% BSA), and then blocked with
1340 Human TruStain FcX (Biolegend) for 10 minutes. Cells were then stained with target antibodies
1341 for 1 hour at 4°C in the dark, followed by two washes using the FACS buffer, and then analyzed
1342 by flow cytometry.

1343
1344   For intracellular staining, cells were dissociated with Accutase and resuspended in DMEM/F12
1345   with GlutaMAX (ThermoFisher, Cat #10565018) with 20% trypsin inhibitor. Cells were then fixed
1346   with Cytofix/Cytoperm solution (BD) at 4°C for 20 minutes, followed by washes with Perm/Wash
1347   solution (BD). Cells were then stained with target antibodies for 45 minutes at 4°C in the dark,
1348   followed by two washes with the FACS buffer, and then analyzed by flow cytometry.
1349
RT-qPCR
Cells were lysed with BME-supplemented RLT buffer and total RNA was extracted with the
RNeasy Plus 96 Kit (Cat #74192, QIAGEN). The extracted RNA was then reverse transcribed
using RevertAid RT Kit (Thermo Fisher, Cat # K1691) with random hexamer primers at 25°C for
5 min, 42°C for 60 min, and 70°C for 5 min. qPCR was then performed using Taqman Fast
Advanced Master Mix (Thermo Fisher, Cat # 4444965) and Taqman probes for GAPDH control
(Thermo Fisher, Cat # 4326317E) and target genes (IDT, custom gene expression assays).
Custom Taqman probe and primer sets were designed to amplify target regions spanning the
guide target sites. qPCR was performed in 384-well plates using the LightCycler 480 Instrument
II (Roche). Target gene expression change was calculated relative to non-targeting controls
using the ddCt method.
1361
Cell viability assays
For cell viability assays in HEK293FT, cells were plated at 9,000 cells per well in a 96-well plate
the day before transfection. Cells were transfected with 100 ng of all-in-one PiggyBac plasmid
containing constitutive CasRx, DjCas13d, or Cas7-11 using Lipofectamine 2000 (Life
Technologies). 72 hours after transfection, cell viability was measured using WST-1 reagent
(5015944001, Sigma) with an incubation time of 2 hours and measurement of absorbance at
440nm. Cell viability of targeting guide groups for each effector was compared relative to the
corresponding non-targeting guide group. Three biological replicates were performed.
1370
To measure cell viability in stem cells, Hela, U2OS and A375 cells, cells were transfected with
the inducible all-in-one PiggyBac plasmids containing inducible CasRx, DjCas13d, or other
Cas13d orthologs. After selection for plasmid integration with 1 µg/ml puromycin for 5-7 days,
cells were induced for effector (CasRx, DjCas13d or other Cas13d orthologs) expression using
Doxycycline (D3072, Sigma). 3-5 days after induction, flow analysis was performed to quantify
the percent of cells expressing the effector in each experimental group using the GFP reporter.
The GFP+ percentage of cells with targeting guide groups for each effector was normalized to
that of the corresponding non-targeting guide group for evaluation of cell viability upon target
RNA knockdown. Three biological replicates were performed.
1380
To measure cell viability in stem cell derived NPCs, HPCs, or neurons, we transfected stem
cells with the inducible all-in-one PiggyBac plasmids containing inducible DjCas13d and
selection with 1 µg/ml puromycin for 7 days to ensure plasmid integration. Differentiation
procedures were then initiated and cells were induced for DjCas13d expression using
Doxycycline (D3072, Sigma) at the middle time point of differentiation. 5-7 days after induction,
flow analysis was performed to quantify the percent of cells expressing the effector in each

experimental group using the GFP reporter. The GFP+ percentage of cells with targeting guide
groups for each effector was normalized to that of the corresponding non-targeting guide group
for evaluation of cell viability upon target RNA knockdown. Three biological replicates were
performed.
1391
RNA-seq library preparation and sequencing
For HEK293FT cells, total RNA was extracted with the RNeasy Plus 96 Kit (Cat #74192,
QIAGEN) 48h after transfection. For H1 cells, cell numbers were counted and normalized
between different samples (different effectors, guides and replicates) 72h after transfection, and
total RNA was extracted with the RNeasy Plus 96 Kit (Cat #74192, QIAGEN). Stranded mRNA
libraries were prepared using the NEBNext II Ultra Directional RNA Library Prep Kit (NEB, Cat#
E7760L) and NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB, Cat #E7490). The
libraries were sequenced on a partial NovaSeq lane with 150 nt paired end reads. ~20M reads
were demultiplexed per sample.
1401
RNA-seq analysis and pathway analysis of CasRx off targets
Sequencing reads were aligned to the hg38 Ensembl transcriptome using Kallisto (Bray et al.,
2016). Mapping was carried out using default parameters except for a b value (number of
bootstraps) of 100. Differential transcript expression was performed with Sleuth (Pimentel et al.,
2017) using triplicates to compare between targeting and non-targeting conditions. Significantly
differentially expressed transcripts were defined as having an adjusted p value < 0.1 and a beta
value > 0.5. Volcano plots were generated in R using the package EnhancedVolcano (Blighe et
al., 2019). Pathway analysis of CasRx off targets was performed using Enrichr (E. Y. Chen et
al., 2013; Kuleshov et al., 2016; Xie et al., 2021) with the Molecular Signatures Database
(MSigDB).
1412
RNA-seq Spike-In for total RNA quantification
To quantify total RNA amount accurately and determine if uniform transcriptome depletion has
occurred following CasRx- or DjCas13-mediated transcriptome targeting, an equal amount of
ERCC RNA Spike-In Mix (ThermoFisher, Cat #4456740) was added to the total RNA extracted
from cell number-normalized H1 samples using the recommended dilution ratio before library
preparation. After library preparation and NGS sequencing, the ratio of experimental reads to
spike-in reads was calculated for all samples, and then normalized to the ratio of control
samples (non-targeting guides) to get the total RNA amount relative to NT.
1421
1422 **RNA integrity analysis**
1423 To examine RNA integrity, electrophoresis was performed on the extracted RNA and the
1424 electrophoresis graphs were visualized on high sensitivity RNA chips using either Bioanalyzer
1425 (Agilent 2100 Bioanalyzer, G2939BA) (for experiments in HEK293FT) or TapeStation (Agilent
1426 4200 TapeStation system, G2991AA) (for experiments in H1).
1427
1428
Stem cell differentiation to NPC, HPC, neurons and RNA targeting experiments

For RNA targeting experiments in NPC and HPC, human embryonic stem cells (hESCs, H1 line,
WiCell) were first transfected with inducible piggyBac-based all-in-one DjCas13d plasmids
containing a puromycin resistance gene as mentioned above. For RNA targeting experiments in
neurons, H1s were first transfected with inducible piggyBac-based all-in-one DjCas13d plasmids
containing neomycin resistant gene by replacing the puromycin resistance gene in the
piggyBac-based all-in-one DjCas13d plasmid with a neomycin resistance gene. After selection
for plasmid integration with 1 µg/ml puromycin (NPC and HPC) or 100 µg/ml G418 Sulfate
(neurons) for 7 days, differentiation procedures were performed as outlined below.
1438
For differentiation to NPC, stem cells were passaged with Accutase (Innovative Cell
Technologies) and plated at 30,000 cells per well into Matrigel-coated 96-well plates with N2B27
media (DMEM/F12 (Thermo Fisher) + N2 (100x, Thermo Fisher) + B27 without vitamin A (50x,
Thermo Fisher)) containing ROCK inhibitor Y-27632 (10 uM, Abcam) and bFGF (40 ng/mL,
Corning). The following day (day 0), media was replaced with N2B27 media containing AZD-
4547 (50 nM, Abcam, Cat# ab216311), LDN-193189 (250 nM, Sigma, Cat# SML0559), A83-01
(250 nM, Sigma, Cat# SML0788), and XAV-939 (3 uM, Abcam, Cat# ab120897) to achieve dual
SMAD and Wnt inhibition. Media was changed daily. On day 3, AZD-4547 was removed. On
day 4, cells were passaged with Accutase (Innovative Cell Technologies) at 1:3 and plated
again onto Matrigel-coated 96-well plates in N2B27 media containing ROCK inhibitor Y-27632
(10 uM, AbAcam), LDN-193189 (250 nM, Sigma, Cat# SML0559), A83-01 (250 nM, Sigma,
Cat# SML0788), and XAV-939 (3 uM, Abcam, Cat# ab120897). Media was replaced the next
day with N2B27 containing LDN-193189 (250 nM, Sigma, Cat# SML0559), A83-01 (250 nM,
Sigma, Cat# SML0788), and XAV-939 (3 uM, Abcam, Cat# ab120897). Media was changed
daily and cells were induced for DjCas13d expression using Doxycycline (D3072, Sigma) on
day 5. On day 8, all drugs were removed and the media was changed with N2B27 only
(DMEM/F12 + N2 (100x) + B27 without vitamin A (50x)). On day 10, the cells were assayed for
target knockdown and NPC marker expression (Pax6 and Sox1) using flow cytometry.
1457
For differentiation to HPC, stem cells were passaged with ReLeSR (StemCell Technologies)
and plated at ~40 colonies per well into Matrigel-coated 12-well plates with mTesR media
(StemCell Technologies) containing ROCK inhibitor Y-27632 (10 uM, Abcam). The following day
(day 0), media was replaced with 2 mL Hematopoietic Media A (STEMdiff Hematopoietic Basal
Media (StemCell Technologies) with STEMdiff Hematopoietic Supplement A (200x, StemCell
Technologies)). On day 2, a half-media change with Hematopoietic Media A was performed. On
day 3, the media was fully replaced with 2 mL Hematopoietic Media B (STEMdiff Hematopoietic
Basal Media (StemCell Technologies) + STEMdiff Hematopoietic Supplement B (200x,
StemCell Technologies). On day 5, there was a half-media change with Hematopoietic Media B,
and cells were induced for DjCas13d expression using Doxycycline (D3072, Sigma). On day 7
and day 10, 1 mL fresh Hematopoietic B media was added but no media was removed. On day
12, the cells were assayed for target knockdown and HPC marker expression (CD43) using flow
cytometry.
1471
For differentiation to neurons, hESCs (H1) were passaged with Accutase (Innovative Cell
Technologies) and plated at 12,000 cells per well into Cultrex (R&D Systems 343400502)-

coated 96-well plates with mTeSR media (StemCell Technologies) containing ROCK inhibitor Y-
27632 (10 uM, Abcam). The following day cells were infected with lentivirus containing a
doxycycline-inducible Ngn2 cassette in mTeSR media (StemCell Technologies) containing
polybrene (10 mg/mL, Santa Cruz Biotechnology sc-134220). Following infection, media was
changed daily to mTeSR media (StemCell Technologies). When cells reached 70% confluency,
they were passaged with Accutase (Innovative Cell Technologies) and re-plated at 12,000 cells
per well into Cultrex-coated 96-well plates with mTeSR media (StemCell Technologies)
containing ROCK inhibitor Y-27632 (10 uM, Abcam). The day of passage was designated as
day 0 of the differentiation protocol. The following day (day 1), media was replaced with mTeSR
media (StemCell Technologies). On day 2, cells were induced for Ngn2 and DjCas13d
expression using 2 ug/mL Doxycycline (2 ug/mL, Sigma D3072). On day 3, media was replaced
with neural induction media (NIM, DMEM/F12 (Gibco 11330032) + Penicillin-Streptomycin
(Gibco 15140122) + Doxycycline (2 ug/mL, Sigma D3072) + Laminin (1.2 ug/mL, Sigma L4544)
+ Insulin (5 ug/mL, Roche 11376497001) + BSA (10 mg/mL, Sigma A4161) + Apo-transferrin
(10 mg/mL, Sigma T1147) + Putrescine (1.6 mg/mL, Sigma P57800) + Progesterone (0.00625
mg/mL, Sigma P8783) + Sodium selenite (0.00104 mg/mL, S5261) + BDNF (10 ug/mL, Sigma
B3795) + Puromycin (10 ug/mL, Life Technologies A1113803)). Media was changed daily. After
3 days of puromycin selection, cells were passaged with Accumax (Innovative Cell
Technologies) and plated at 87,500 cells per well with neural maturation media (Neurobasal
differentiation media (Neurobasal Media (Gibco 21103049) + DMEM Media (Gibco 10569010) +
HEPES (0.5x, Gibco 15630130) + Penicillin-Streptomycin (Gibco 15140122) + Glutamax (1 mM,
Gibco 35050061)) + Doxycycline (2 ug/mL, Sigma D3072) + Laminin (2.4 ug/mL, Sigma L4544)
+ BDNF (10 ug/mL, Sigma B3795) + dbCAMP (49.14 ug/mL, Sigma Aldrich D0627) + B27 with
vitamin A (1x, Gibco 17504044) + N-acetyl cysteine (5 ug/mL, Sigma A9165) containing ROCK
inhibitor Y-27632 (10 uM, Abcam). Media was changed daily. On day 8, media was replaced
with neural maturation media containing AraC (2.4 ug/mL, Sigma Aldrich C1768) to remove any
post-mitotic neurons from the culture. On day 11, the cells were assayed for target knockdown
using flow cytometry.
1502
1503
1504
1505

References

1. Abudayyeh, O. O., Gootenberg, J. S., Essletzbichler, P., Han, S., Joung, J., Belanto, J. J., Verdine, V., Cox, D. B. T., Kellner, M. J., Regev, A., Lander, E. S., Voytas, D. F., Ting, A. Y., & Zhang, F. (2017). RNA targeting with CRISPR–Cas13. *Nature*, *550*(7675), 280–284.

2. Abudayyeh, O. O., Gootenberg, J. S., Franklin, B., Koob, J., Kellner, M. J., Ladha, A., Joung, J., Kirchgatterer, P., Cox, D. B. T., & Zhang, F. (2019). A cytosine deaminase for programmable single-base RNA editing. *Science*, *365*(6451), 382–386.

3. Abudayyeh, O. O., Gootenberg, J. S., Konermann, S., Joung, J., Slaymaker, I. M., Cox, D. B. T., Shmakov, S., Makarova, K. S., Semenova, E., Minakhin, L., Severinov, K., Regev, A., Lander, E. S., Koonin, E. V., & Zhang, F. (2016). C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. *Science*, *353*(6299). https://doi.org/10.1126/science.aaf5573

4. Ai, Y., Liang, D., & Wilusz, J. E. (2022). CRISPR/Cas13 effectors have differing extents of off-target effects that limit their utility in eukaryotic cells. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkac159

5. Alipanahi, B., Delong, A., Weirauch, M. T., & Frey, B. J. (2015). Predicting the sequence specificities of DNA- and RNA-binding proteins by deep learning. In *Nature Biotechnology* (Vol. 33, Issue 8, pp. 831–838). https://doi.org/10.1038/nbt.3300

6. Almeida, A., Nayfach, S., Boland, M., Strozzi, F., Beracochea, M., Shi, Z. J., Pollard, K. S., Sakharova, E., Parks, D. H., Hugenholtz, P., Segata, N., Kyrpides, N. C., & Finn, R. D. (2021). A unified catalog of 204,938 reference genomes from the human gut microbiome. *Nature Biotechnology*, *39*(1), 105–114.

7. Arbab, M., Shen, M. W., Mok, B., Wilson, C., Matuszek, Ż., Cassa, C. A., & Liu, D. R. (2020). Determinants of Base Editing Outcomes from Target Library Analysis and Machine Learning. *Cell*, *182*(2), 463–480.e30.

8. Avsec, Ž., Weilert, M., Shrikumar, A., Krueger, S., Alexandari, A., Dalal, K., Fropf, R., McAnany, C., Gagneur, J., Kundaje, A., & Zeitlinger, J. (2021). Base-resolution models of transcription-factor binding reveal soft motif syntax. *Nature Genetics*, *53*(3), 354–366.

9. Blighe, K., Rana, S., & Lewis, M. (2019). EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling. *R Package Version*, *1*(0).

10. Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Erratum: Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, *34*(8), 888.

11. Buchman, A. B., Brogan, D. J., Sun, R., Yang, T., Hsu, P. D., & Akbari, O. S. (2020). Programmable RNA Targeting Using CasRx in Flies. *The CRISPR Journal*, *3*(3), 164–176.

12. Camarillo-Guerrero, L. F., Almeida, A., Rangel-Pineros, G., Finn, R. D., & Lawley, T. D. (2021). Massive expansion of human gut bacteriophage diversity. *Cell*, *184*(4), 1098–1109.e9.

13. Chen, E. Y., Tan, C. M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G. V., Clark, N. R., & Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics*, *14*, 128.

14. Chen, I.-M. A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M., Hajek,

P., Ritter, S., Varghese, N., Seshadri, R., Roux, S., Woyke, T., Eloe-Fadrosh, E. A., Ivanova, N. N., & Kyrpides, N. C. (2021). The IMG/M data management and analysis system v.6.0: new tools and advanced capabilities. *Nucleic Acids Research*, *49*(D1), D751–D763.

15. Chuai, G., Ma, H., Yan, J., Chen, M., Hong, N., Xue, D., Zhou, C., Zhu, C., Chen, K., Duan, B., Gu, F., Qu, S., Huang, D., Wei, J., & Liu, Q. (2018). DeepCRISPR: optimized CRISPR guide RNA design by deep learning. *Genome Biology*, *19*(1), 80.

16. Cox, D. B. T., Gootenberg, J. S., Abudayyeh, O. O., Franklin, B., Kellner, M. J., Joung, J., & Zhang, F. (2017). RNA editing with CRISPR-Cas13. *Science*, *358*(6366), 1019–1027.

17. Do, C. B., Woods, D. A., & Batzoglou, S. (2006). CONTRAfold: RNA secondary structure prediction without physics-based models. *Bioinformatics* , *22*(14), e90–e98.

18. Doench, J. G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E. W., Donovan, K. F., Smith, I., Tothova, Z., Wilen, C., Orchard, R., Virgin, H. W., Listgarten, J., & Root, D. E. (2016). Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nature Biotechnology*, *34*(2), 184–191.

19. Doench, J. G., Petersen, C. P., & Sharp, P. A. (2003). siRNAs can function as miRNAs. *Genes & Development*, *17*(4), 438–442.

20. East-Seletsky, A., O'Connell, M. R., Knight, S. C., Burstein, D., Cate, J. H. D., Tjian, R., & Doudna, J. A. (2016). Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection. *Nature*, *538*(7624), 270–273.

21. Forster, S. C., Kumar, N., Anonye, B. O., Almeida, A., Viciani, E., Stares, M. D., Dunn, M., Mkandawire, T. T., Zhu, A., Shao, Y., Pike, L. J., Louie, T., Browne, H. P., Mitchell, A. L., Neville, B. A., Finn, R. D., & Lawley, T. D. (2019). A human gut bacterial genome and culture collection for improved metagenomic analyses. *Nature Biotechnology*, *37*(2), 186–192.

22. Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., & Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology*, *59*(3), 307–321.

23. Han, S., Zhao, B. S., Myers, S. A., Carr, S. A., He, C., & Ting, A. Y. (2020). RNA-protein interaction mapping via MS2- or Cas13-based APEX targeting. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(36), 22068–22079.

24. Hart, T., Chandrashekhar, M., Aregger, M., Steinhart, Z., Brown, K. R., MacLeod, G., Mis, M., Zimmermann, M., Fradet-Turcotte, A., Sun, S., Mero, P., Dirks, P., Sidhu, S., Roth, F. P., Rissland, O. S., Durocher, D., Angers, S., & Moffat, J. (2015). High-Resolution CRISPR Screens Reveal Fitness Genes and Genotype-Specific Cancer Liabilities. *Cell*, *163*(6), 1515–1526.

25. *HMMER*. (n.d.). Retrieved February 25, 2022, from http://hmmer.org

26. Horlbeck, M. A., Gilbert, L. A., Villalta, J. E., Adamson, B., Pak, R. A., Chen, Y., Fields, A. P., Park, C. Y., Corn, J. E., Kampmann, M., & Weissman, J. S. (2016). Compact and highly active next-generation libraries for CRISPR-mediated gene repression and activation. *eLife*, *5*. https://doi.org/10.7554/eLife.19760

27. Huang, L., Zhang, H., Deng, D., Zhao, K., Liu, K., Hendrix, D. A., & Mathews, D. H. (2019). LinearFold: linear-time approximate RNA folding by 5'-to-3' dynamic

1593 programming and beam search. In *Bioinformatics* (Vol. 35, Issue 14, pp. i295–i304).
1594 https://doi.org/10.1093/bioinformatics/btz375

28. Jackson, A. L., Bartz, S. R., Schelter, J., Kobayashi, S. V., Burchard, J., Mao, M., Li, B.,
Cavet, G., & Linsley, P. S. (2003). Expression profiling reveals off-target gene regulation
by RNAi. *Nature Biotechnology*, *21*(6), 635–637.

29. Kannan, S., Altae-Tran, H., Jin, X., Madigan, V. J., Oshiro, R., Makarova, K. S., Koonin,
E. V., & Zhang, F. (2022). Compact RNA editors with small Cas13 proteins. *Nature
Biotechnology*, *40*(2), 194–197.

30. Karolchik, D., Hinrichs, A. S., Furey, T. S., Roskin, K. M., Sugnet, C. W., Haussler, D., &
Kent, W. J. (2004). The UCSC Table Browser data retrieval tool. *Nucleic Acids
Research*, *32*(Database issue), D493–D496.

31. Katoh, K., Misawa, K., Kuma, K.-I., & Miyata, T. (2002). MAFFT: a novel method for
rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids
Research*, *30*(14), 3059–3066.

32. Kato, K., Zhou, W., Okazaki, S., Isayama, Y., Nishizawa, T., Gootenberg, J. S.,
Abudayyeh, O. O., & Nishimasu, H. (2022). Structure and engineering of the type III-E
CRISPR-Cas7-11 effector complex. *Cell*, *185*(13), 2324–2337.e16.

33. Kelley, D. R., Snoek, J., & Rinn, J. L. (2016). Basset: learning the regulatory code of the
accessible genome with deep convolutional neural networks. *Genome Research*, *26*(7),
990–999.

34. Kim, H. K., Kim, Y., Lee, S., Min, S., Bae, J. Y., Choi, J. W., Park, J., Jung, D., Yoon, S.,
& Kim, H. H. (2019). SpCas9 activity prediction by DeepSpCas9, a deep learning–based
model with high generalization performance. In *Science Advances* (Vol. 5, Issue 11, p.
eaax9249). https://doi.org/10.1126/sciadv.aax9249

35. Kim, H. K., Min, S., Song, M., Jung, S., Choi, J. W., Kim, Y., Lee, S., Yoon, S., & Kim, H.
(henry). (2018). Deep learning improves prediction of CRISPR–Cpf1 guide RNA activity.
Nature Biotechnology, *36*(3), 239–241.

36. Koblan, L. W., Arbab, M., Shen, M. W., Hussmann, J. A., Anzalone, A. V., Doman, J. L.,
Newby, G. A., Yang, D., Mok, B., Replogle, J. M., Xu, A., Sisley, T. A., Weissman, J. S.,
Adamson, B., & Liu, D. R. (2021). Efficient C•G-to-G•C base editors developed using
CRISPRi screens, target-library analysis, and machine learning. *Nature Biotechnology*,
39(11), 1414–1425.

37. Konermann, S., Lotfy, P., Brideau, N. J., Oki, J., Shokhirev, M. N., & Hsu, P. D. (2018).
Transcriptome Engineering with RNA-Targeting Type VI-D CRISPR Effectors. *Cell*,
173(3), 665–676.e14.

38. Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z.,
Koplev, S., Jenkins, S. L., Jagodnik, K. M., Lachmann, A., McDermott, M. G., Monteiro,
C. D., Gundersen, G. W., & Ma'ayan, A. (2016). Enrichr: a comprehensive gene set
enrichment analysis web server 2016 update. *Nucleic Acids Research*, *44*(W1), W90–
W97.

39. Kursa, M. B., Rudnicki, W. R., & Others. (2010). Feature selection with the Boruta
package. *Journal of Statistical Software*, *36*(11), 1–13.

40. Kushawah, G., Hernandez-Huertas, L., Abugattas-Nuñez Del Prado, J., Martinez-
Morales, J. R., DeVore, M. L., Hassan, H., Moreno-Sanchez, I., Tomas-Gallardo, L.,

Diaz-Moscoso, A., Monges, D. E., Guelfo, J. R., Theune, W. C., Brannan, E. O., Wang, W., Corbin, T. J., Moran, A. M., Sánchez Alvarado, A., Málaga-Trillo, E., Takacs, C. M., … Moreno-Mateos, M. A. (2020). CRISPR-Cas13d Induces Efficient mRNA Knockdown in Animal Embryos. *Developmental Cell*, *54*(6), 805–817.e7.

41. Lanchantin, J., Singh, R., Lin, Z., & Qi, Y. (2016). Deep Motif: Visualizing Genomic Sequence Classifications. In *arXiv [cs.LG]*. arXiv. http://arxiv.org/abs/1605.01133

42. Li, S., Li, X., Xue, W., Zhang, L., Yang, L.-Z., Cao, S.-M., Lei, Y.-N., Liu, C.-X., Guo, S.-K., Shan, L., Wu, M., Tao, X., Zhang, J.-L., Gao, X., Zhang, J., Wei, J., Li, J., Yang, L., & Chen, L.-L. (2021). Screening for functional circular RNAs using the CRISPR–Cas13 system. In *Nature Methods* (Vol. 18, Issue 1, pp. 51–59). https://doi.org/10.1038/s41592-020-01011-4

43. Liu, L., Li, X., Ma, J., Li, Z., You, L., Wang, J., Wang, M., Zhang, X., & Wang, Y. (2017). The Molecular Architecture for RNA-Guided RNA Cleavage by Cas13a. *Cell*, *170*(4), 714–726.e10.

44. Lorenz, R., Bernhart, S. H., Höner Zu Siederdissen, C., Tafer, H., Flamm, C., Stadler, P. F., & Hofacker, I. L. (2011). ViennaRNA Package 2.0. *Algorithms for Molecular Biology: AMB*, *6*, 26.

45. Lundberg, S. M., Erion, G., Chen, H., DeGrave, A., Prutkin, J. M., Nair, B., Katz, R., Himmelfarb, J., Bansal, N., & Lee, S.-I. (2020). From Local Explanations to Global Understanding with Explainable AI for Trees. *Nature Machine Intelligence*, *2*(1), 56–67.

46. Luo, B., Cheung, H. W., Subramanian, A., Sharifnia, T., Okamoto, M., Yang, X., Hinkle, G., Boehm, J. S., Beroukhim, R., Weir, B. A., Mermel, C., Barbie, D. A., Awad, T., Zhou, X., Nguyen, T., Piqani, B., Li, C., Golub, T. R., Meyerson, M., … Root, D. E. (2008). Highly parallel identification of essential genes in cancer cells. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(51), 20380–20385.

47. Mahas, A., Aman, R., & Mahfouz, M. (2019). CRISPR-Cas13d mediates robust RNA virus interference in plants. *Genome Biology*, *20*(1), 263.

48. Metsky, H. C., Welch, N. L., Haradhvala, N. J., Rumker, L., Zhang, Y. B., Pillai, P. P., Yang, D. K., Ackerman, C. M., Weller, J., Blainey, P. C., Myhrvold, C., Mitzenmacher, M., & Sabeti, P. C. (2021). Designing viral diagnostics with model-based optimization. In *bioRxiv* (p. 2020.11.28.401877). https://doi.org/10.1101/2020.11.28.401877

49. Metsky, H. C., Welch, N. L., Pillai, P. P., Haradhvala, N. J., Rumker, L., Mantena, S., Zhang, Y. B., Yang, D. K., Ackerman, C. M., Weller, J., Blainey, P. C., Myhrvold, C., Mitzenmacher, M., & Sabeti, P. C. (2022). Designing sensitive viral diagnostics with machine learning. *Nature Biotechnology*. https://doi.org/10.1038/s41587-022-01213-5

50. Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E. M., Kubal, M., Paczian, T., Rodriguez, A., Stevens, R., Wilke, A., Wilkening, J., & Edwards, R. A. (2008). The metagenomics RAST server - a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics*, *9*, 386.

51. Mitchell, A. L., Almeida, A., Beracochea, M., Boland, M., Burgin, J., Cochrane, G., Crusoe, M. R., Kale, V., Potter, S. C., Richardson, L. J., Sakharova, E., Scheremetjew, M., Korobeynikov, A., Shlemov, A., Kunyavskaya, O., Lapidus, A., & Finn, R. D. (2020). MGnify: the microbiome analysis resource in 2020. *Nucleic Acids Research*, *48*(D1), D570–D578.

52. Özcan, A., Krajeski, R., Ioannidi, E., Lee, B., Gardner, A., Makarova, K. S., Koonin, E.
V., Abudayyeh, O. O., & Gootenberg, J. S. (2021). Programmable RNA targeting with
the single-protein CRISPR effector Cas7-11. *Nature*, *597*(7878), 720–725.
53. Pimentel, H., Bray, N. L., Puente, S., Melsted, P., & Pachter, L. (2017). Differential
analysis of RNA-seq incorporating quantification uncertainty. *Nature Methods*, *14*(7),
687–690.
54. Shi, P., Murphy, M. R., Aparicio, A. O., Kesner, J. S., Fang, Z., Chen, Z., Trehan, A., &
Wu, X. (2021). RNA-guided cell targeting with CRISPR/RfxCas13d collateral activity in
human cells. In *bioRxiv* (p. 2021.11.30.470032).
https://doi.org/10.1101/2021.11.30.470032
55. Shrikumar, A., Tian, K., Avsec, Ž., Shcherbina, A., Banerjee, A., Sharmin, M., Nair, S., &
Kundaje, A. (2018). Technical Note on Transcription Factor Motif Discovery from
Importance Scores (TF-MoDISco) version 0.5.6.5. In *arXiv [cs.LG]*. arXiv.
http://arxiv.org/abs/1811.00416
56. Sigoillot, F. D., Lyman, S., Huckins, J. F., Adamson, B., Chung, E., Quattrochi, B., &
King, R. W. (2012). A bioinformatics method identifies prominent off-targeted transcripts
in RNAi screens. *Nature Methods*, *9*(4), 363–366.
57. Slaymaker, I. M., Mesa, P., Kellner, M. J., Kannan, S., Brignole, E., Koob, J., Feliciano,
P. R., Stella, S., Abudayyeh, O. O., Gootenberg, J. S., Strecker, J., Montoya, G., &
Zhang, F. (2021). High-resolution structure of cas13b and biochemical characterization
of RNA targeting and cleavage. In *Cell Reports* (Vol. 34, Issue 10, p. 108865).
https://doi.org/10.1016/j.celrep.2021.108865
58. Steinegger, M., & Söding, J. (2017). MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. *Nature Biotechnology*, *35*(11), 1026–
1028.
59. Sunagawa, S., Coelho, L. P., Chaffron, S., Kultima, J. R., Labadie, K., Salazar, G.,
Djahanschiri, B., Zeller, G., Mende, D. R., Alberti, A., Cornejo-Castillo, F. M., Costea, P.
I., Cruaud, C., d'Ovidio, F., Engelen, S., Ferrera, I., Gasol, J. M., Guidi, L., Hildebrand,
F., … Bork, P. (2015). Ocean plankton. Structure and function of the global ocean
microbiome. *Science*, *348*(6237), 1261359.
60. Sundararajan, M., Taly, A., & Yan, Q. (2017). Axiomatic Attribution for Deep Networks.
In D. Precup & Y. W. Teh (Eds.), *Proceedings of the 34th International Conference on
Machine Learning* (Vol. 70, pp. 3319–3328). PMLR.
61. Wayment-Steele, H. K., Kladwang, W., Participants, E., & Das, R. (2020). RNA
secondary structure packages ranked and improved by high-throughput experiments. In
bioRxiv (p. 2020.05.29.124511). https://doi.org/10.1101/2020.05.29.124511
62. Wessels, H.-H., Méndez-Mancilla, A., Guo, X., Legut, M., Daniloski, Z., & Sanjana, N. E.
(2020). Massively parallel Cas13 screens reveal principles for guide RNA design. *Nature
Biotechnology*, *38*(6), 722–727.
63. Wilson, C., Chen, P. J., Miao, Z., & Liu, D. R. (2020). Programmable m6A modification of
cellular RNAs with a Cas13-directed methyltransferase. In *Nature Biotechnology* (Vol.
38, Issue 12, pp. 1431–1440). https://doi.org/10.1038/s41587-020-0572-6
64. Xie, Z., Bailey, A., Kuleshov, M. V., Clarke, D. J. B., Evangelista, J. E., Jenkins, S. L.,
Lachmann, A., Wojciechowicz, M. L., Kropiwnicki, E., Jagodnik, K. M., Jeon, M., &

Ma'ayan, A. (2021). Gene Set Knowledge Discovery with Enrichr. *Current Protocols*,
1(3), e90.

65. Xu, C., Zhou, Y., Xiao, Q., He, B., Geng, G., Wang, Z., Cao, B., Dong, X., Bai, W.,
Wang, Y., Wang, X., Zhou, D., Yuan, T., Huo, X., Lai, J., & Yang, H. (2021).
Programmable RNA editing with compact CRISPR-Cas13 systems from uncultivated
microbes. *Nature Methods*, *18*(5), 499–506.

66. Xue, L., Tang, B., Chen, W., & Luo, J. (2019). Prediction of CRISPR sgRNA Activity
Using a Deep Convolutional Neural Network. *Journal of Chemical Information and*
Modeling, *59*(1), 615–624.

67. Youngblut, N. D., de la Cuesta-Zuluaga, J., Reischer, G. H., Dauser, S., Schuster, N.,
Walzer, C., Stalder, G., Farnleitner, A. H., & Ley, R. E. (2020). Large-Scale Metagenome
Assembly Reveals Novel Animal-Associated Microbial Genomes, Biosynthetic Gene
Clusters, and Other Genetic Diversity. *mSystems*, *5*(6).
https://doi.org/10.1128/mSystems.01045-20

68. Yu, G. (2020). Using ggtree to Visualize Data on Tree-Like Structures. *Current Protocols*
in Bioinformatics / Editoral Board, Andreas D. Baxevanis ... [et Al.], *69*(1), e96.

69. Zhang, C., Konermann, S., Brideau, N. J., Lotfy, P., Wu, X., Novick, S. J., Strutzenberg,
T., Griffin, P. R., Hsu, P. D., & Lyumkis, D. (2018). Structural Basis for the RNA-Guided
Ribonuclease Activity of CRISPR-Cas13d. *Cell*, *175*(1), 212–223.e17.