Exocytosis of the silicified cell wall of diatoms involves extensive membrane disintegration

Diede de Haan[1], Hadas Peled-Zehavi[1], Yoseph Addadi[2], Oz Ben Joseph[1], Lior Aram[1] and Assaf Gal[1*]

[1]Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot, Israel
[2]Life Science Core Facilities, Weizmann Institute of Science, Rehovot, Israel
*email: assaf.gal@weizmann.ac.il

Keywords

Abstract

Diatoms are unicellular algae that are characterized by their silica cell walls. The silica elements form intracellularly in a membrane-bound organelle, and are exocytosed after completion. How diatoms maintain membrane homeostasis during the exocytosis of these large and rigid silica elements is a long-standing enigma. We studied membrane dynamics during cell wall formation and exocytosis in the diatom *Stephanopyxis turris*, using live-cell confocal microscopy and advanced electron microscopy. Our results provide detailed information on the ultrastructure and dynamics of the silicification process, showing that during cell wall formation, the organelle membranes tightly enclose the mineral phase, creating a precise mold of the delicate geometrical patterns. Surprisingly, during exocytosis of the mature silica elements, the proximal organelle membrane becomes the new plasma membrane, and the distal membranes gradually disintegrate into the extracellular space without any noticeable endocytic retrieval or extracellular repurposing. These observations suggest that diatoms evolved an extraordinary exocytosis mechanism in order to secrete their cell wall elements.

Introduction

Diatoms are important members of marine and freshwater phytoplankton communities[1]. This diverse group of unicellular algae is characterized by silica cell walls with intricate species-specific shapes and patterns[2]. Despite immense morphological diversity between species, most diatom cell walls have the same, conserved layout of two similarly shaped silica 'shells' that partially overlap like a petri-dish. Each 'shell' in itself consists of a valve and a series of girdle bands. The valves usually define the shape of the cell, are richly ornamented, and contain hierarchical pore patterns. The girdle bands form partially overlapping rings that surround the sidewalls of the cells.

Diatom cell wall formation is genetically controlled and under strict biological regulation[3–6]. Silicification is usually an intracellular process, taking place inside a membrane-bound organelle, the silica deposition vesicle (SDV)[7–9]. The chemical environment inside the SDV, and the cellular regulation over its contents and shape, exert tight control over the silicification process, resulting in highly specific and reproducible cell wall architectures[10–14]. After intracellular formation, the mature silica elements are exocytosed to cover the cell surface[7,15]. Such exocytosis events are extraordinary in cell biology since the content of the SDV is an enormous solid structure that needs to be secreted without endangering the integrity of the cell.

In classical exocytosis, an intracellular membrane vesicle fuses with the plasma membrane to deliver its content to the extracellular space. Homeostasis of cell surface area and plasma membrane composition can be maintained through a transient fusion that prevents the secreting vesicle from integrating with the plasma membrane, or by offsetting the added membrane with compensatory endocytosis[16]. However, it is difficult to envision how these mechanisms can be applied to the secretion of diatom cell walls due to their large size and rigidity. In some species, the valve covers as much as half of the total cell surface, causing the surface area of the SDV membrane to be similar to that of the entire plasma membrane. The fusion of such an SDV membrane with the plasma membrane would lead to an almost instantaneous doubling of the membrane surface area. Hence, the mechanism that diatoms use to maintain the needed large-scale membrane homeostasis during cell wall exocytosis is a long-standing enigma.

Early ultrastructural studies brought forward several models for silica cell wall exocytosis in diatoms[15,17–24] (Fig. 1). One scenario suggests that the newly formed valve is externalized through

formation of a new plasma membrane underneath it (Fig. 1 A). The SDV membrane and the original

plasma membrane around the new valve remain in place, forming an organic layer around the

silica (Fig. 1 B). An alternative is that the valve is externalized as a result of localized fusion of the

SDV and plasma membrane at the valve edge, such that the proximal part of the SDV membrane

becomes the new plasma membrane (Fig. 1 C). The distal part of the SDV membrane and the

original plasma membrane were suggested to either remain as an extracellular layer protecting

the silica (Fig. 1 D) or shed off to the environment (Fig. 1 E). Another model suggests localized

membrane fusion at the apical part of the valve, followed by pulling of the distal SDV and original

plasma membrane back into the cytoplasm at the valve edge (Fig. 1 F). In this last model, which is

in accordance with classical exocytosis, the composition of the proximal SDV membrane is altered

for its new function as the plasma membrane by incorporation of membranous vesicles (Fig 1 G).

While none of these models have been unequivocally validated, fluorescently-tagged SDV

membrane proteins in *Thalassiosira pseudonana* show rapid diffusion throughout the entire

plasma membrane at the time of valve exocytosis, supporting the notion of (at least partial)

integration of the SDV membrane into the plasma membrane[3,25].

Figure 1. Hypotheses for valve exocytosis in diatoms. The middle row (**A**, **C**, **F**) shows three different scenarios of valve release, and the bottom row (**B**, **D**, **E**, **G**) shows the subsequent fate of the membranes in each of these scenarios. See text for details

Significant experimental challenges have hindered progress in solving the conceptual problems of valve exocytosis, and in elucidating the dynamics and ultrastructure of the SDV. On the one hand, traditional electron microscopy sample preparation for ultrastructural studies causes artefacts like membrane shrinking and structure deformation[26]. On the other hand, the advantages of light microscopy in studying the dynamic aspects of cell wall formation in living cells are limited by the low spatial resolution and scarcity of molecular biology tools[27,28]. Due to these limitations, *in situ* observations of the SDV in the cellular context are sparse, and direct evidence for the nature of the silicification and exocytosis process is missing.

In this study, we investigated membrane dynamics during valve formation and exocytosis in *Stephanopyxis turris* using a combination of transmission electron microscopy (TEM) and live-cell confocal fluorescence microscopy. Compared to commonly studied diatom species, *S. turris* is large and has easily discernible silica structures, allowing detailed observations of individual valves and the dynamic development of their architectural features using fluorescence microscopy[29,30]. In addition, state-of-the-art sample preparation methods for TEM result in excellent preservation of subcellular structures, enabling investigation of the SDV in the near-native state at high resolution without the distorting effects of traditional chemical fixation[31]. Using these methods, we show that the SDV membrane tightly surrounds growing cell walls during the entire silicification process, and that after valve exocytosis, external membranes slowly disintegrate, without signs of endocytic retrieval or extracellular repurposing.

Results

S. turris cells are cylinders of about 50 µm long and 20 µm in diameter (Fig. 2 A). The cells contain a large central vacuole, surrounded by a thin peripheral layer of cytoplasm filled with many small chloroplasts. The cells form chains that are linked through tubular extensions of silica (labiate processes) that extend from the apex of the valves (Fig. 2 B). When the cell walls are examined using a scanning electron microscope (SEM), the two-layered structure of the valve is evident (Fig.

2 B, C). The proximal layer is thin and perforated by nano-sized pores, and is covered on the distal side by a more elevated layer that forms large polygons. The top of the polygonal layer is flattened, forming a 'T'-shape in cross-section.

Timing of cell wall formation is linked to the cell cycle. After cell division, each daughter cell inherits one half of the parental cell wall and forms a new valve (Supplementary Fig. 1). Directly after its exocytosis, the new valve is juxtaposed to the parental valve and covered by the parental girdle bands. The girdle bands are very thin and flexible such that some features of the underlying valve are visible, e.g a 'sheet' of girdle bands draped over the linking processes (Fig. 2 D, arrow). During the interphase, the cell grows in the longitudinal direction, pushing apart the two valves while girdle bands are added to cover the central part of the growing cell (Fig. 2 E, F).

Figure 2. Cell architecture of *S. turris*. (**A**) Light microscope image of a chain of four cells connected through linking extensions (green). Each cell contains many yellow-brown chloroplasts

(blue). The two parental valves (P) are at the extremes of the chain, the two valves formed after the

first cell division (F1) are located in the center, and the four remaining valves (F2) are the youngest,

formed after the second cell division. Girdle bands (pink) cover the youngest valves. The nuclei

(orange) are located at the poles directly under the new valves (F2). Scale bar: 20 µm. (**B**) SEM

image of an apex of a valve, showing the polygonal pattern and linking extensions. Scale bar: 2

µm. (**C**) A fractured valve, revealing the T-shaped cross-section (green) of the polygonal silica layer

that is distal to the thinner perforated layer (purple). Scale bar: 500 nm. (**D-F**) Cells at different

stages of the cell cycle. (**D**) A cell with a new valve (F2, right) that is covered with parental girdle

bands (arrow) (P, left). Scale bar: 10 µm. (**E**) A cell during elongation and girdle band formation. As

valves were pushed apart, the midsection is covered only by girdle bands. Scale bar: 10 µm. (**F**)

Two daughter cells shortly after cell division, located under full-length girdle bands. Scale bar: 20

µm.

To study specific stages along the dynamic process of valve formation, we used previous

reports and established a protocol for *S. turris* cell cycle synchronization that is based on light

starvation[32,33]. For some diatoms, among which *S. turris*, extended dark period causes cell cycle

arrest in interphase, prior to the formation of new girdle bands[32]. When the culture is re-exposed

to light after 20 hours in the dark, the cells resume their cell cycle synchronously. Progression of

the diatom cell cycle can easily be monitored by visualizing the formation of new cell wall elements

using the fluorescent dye PDMPO[34]. PDMPO accumulates in the SDV and stains newly formed silica

(Fig. 3). After the end of light-starvation, the cells will first form their girdle bands and then the

valves (Fig. 3 A). The synchronized growth allowed us to calibrate time points at which the majority

of the cells are at specific stages of cell wall formation (Fig. 3 B, C). To study valve formation, the

cells were harvested 9 to 10 hours after the end of light starvation, when the synchronized culture

contains the highest proportion of cells at that stage.

Figure 3. Light-induced synchronization of *S. turris* **cell cycle.** (**A**) Representative PDMPO fluorescence images of synchronized cultures, at various time points after the end of light starvation and addition of PDMPO. Shortly after the end of light starvation, none of the cells have fluorescently labelled silica, and only PDMPO fluorescence from within the vacuoles is visible. Four hours later, most cells have started forming girdle bands, visible as a fluorescent ring (arrowheads) around the cell. After eight hours, the majority of the culture is forming new, fluorescent valves. After twelve hours, nearly all cells have completed the formation of a new valve. Scale bar: 50 μm. (**B**) High magnification micrographs and schematics of single cells at key stages of cell wall formation. The key stages that are distinguished are: 'start', cells without PDMPO-labelled silica; 'girdle bands', cells during cell elongation and girdle band formation; 'early valve', after cell division, valve formation has started but is prior to growth of linking extensions; 'late valve', cells

at later stages of valve formation when polygons and extensions are visible. Scale bar: 20 μm (**C**)

Percentage of cells at these four key stages during synchronized growth. The lines show the mean

of four independent experiments, indicated with symbols. The highest percentage of cells that are

forming valves is reached around nine hours.

To observe valve formation at a structural level, we used advanced sample preparation

techniques for TEM. In short, synchronized cells were vitrified using high-pressure freezing and

subsequently freeze-substituted and embedded in resin. These methods provide an optimal

combination of near-native state preservation of subcellular structures paired with the ability of

high-throughput room temperature imaging. The TEM images of *S. turris* cells prepared according

to this procedure show exceptional preservation of the cellular environment, and in particular the

SDV membranes and inorganic contents (Fig. 4).

Figure 4. Ultrastructural details of _S. turris_ revealed by TEM imaging. Insets indicate which part of the cell is displayed. (**A**) A cross section through an entire cell, the nucleus (n), and the large central vacuole (v) that is surrounded by darker chloroplasts are dominant. The silica cell wall (dark contrast) is clearly visible at the left side, where the protoplast is not directly adjacent to the cell wall due to slight plasmolysis. Arrowheads indicate girdle bands, and an arrow indicates a linking extension. Scale bar: 10 μm. (**B**) The meeting point of a new valve (nv), and parental valve (pv) with girdle bands (gb) attached to the latter. Scale bar: 1 μm. (**C**) Apex of a cell with the nucleus (n) located directly under the valve. Silica protrusions indicated with arrowheads are cross sections of the polygonal layer. Scale bar: 5 μm. **(D-H)** High magnification images of: (**D**) Golgi-body. Scale bar: 500 nm. (**E**) Chloroplast. Scale bar: 1 μm. (**F**) SDV with growing silica (arrowheads). Bilayers of the proximal SDV membrane (pink), distal SDV membrane (orange), and plasma membrane (yellow) can be distinguished. Scale bar: 100 nm. (**G**) Fully formed polygon (arrow) still inside an SDV. Scale bar: 500 nm. (**H**) Linking extension during its formation in an SDV. Scale bar: 500 nm.

By analysing hundreds of TEM images of synchronized cells at different stages of the cell cycle, and categorizing and sorting these images, we established a timeline of valve formation and exocytosis in _S. turris_ (Fig. 5). Valve formation starts at the cell's apex, where the bilayer membranes of the proximal and distal SDV membranes, as well as the plasma membrane, can be clearly distinguished (Fig. 5 A-A''). With a distance of only 20 nm, the distal SDV membrane is in very close proximity to the plasma membrane. Silica precipitation proceeds radially and the SDV expands with it. After formation of the porous base layer, the polygonal layer is formed distally on top of the base layer (Fig. 5 B-B''). The valve reaches its final morphology while it is still fully enclosed in the SDV (Fig. 5 C-C''). During the entire silicification process, the SDV membrane tightly delineates the growing valve, forming a highly confined space, in which the silica morphology is under the control of the SDV membrane.

The details of the exocytosis process can be observed in the stages following valve maturation. The first indication for exocytosis are isolated discontinuities in the distal membrane coverage of the valve silica, even though the majority of the valve is still tightly surrounded by membranes (Fig. 5 D-D''). From this moment, the membrane ultrastructure changes dramatically. The tight delineation of the distal SDV and plasma membrane around the valve loosens, and we

observe deterioration of the structural integrity of the distal membranes (Fig 5 E-E''). The proximal

SDV membrane is now continuous with the plasma membrane under the parental valve, forming

the outermost surface that separates the cytoplasm from the cell's exterior, and thereby it is the

new plasma membrane under the valve (Fig. 5 E'' arrow). At the end of this exocytosis process, the

new valves can be recognized due to their position under the parental girdle bands. However,

there are no visible traces of the old plasma and distal SDV membrane (Fig. 5 F-F'').

Figure 5. TEM images showing sequential stages of valve formation and exocytosis in *S.*

turris. The top row shows a representative image of each stage, with higher magnification views

of the boxed areas in the rows below. (**A**) Valve formation starts at the apex of the cell. (**A'**) Growing

silica inside the SVD that is located directly under the plasma membrane. (**A''**) The SDV ends at

the growing silica edge. The inset shows the bilayers of the proximal (pink) and distal (orange) SDV

and the plasma membrane (yellow) in higher magnification. (**B**) Later during valve formation, (**B'**)

the polygonal layer (arrow) is formed on top of the base layer, and (**B''**) the growing edge of the

new valve has reached the edge of the parental valve. (**C**) Silicification of the new valve (arrow) is

completed while it is still fully enclosed in the SDV. Mature valves can be recognized by (**C'**) the

fully formed and flattened polygonal layer and (**C''**) pronounced hook shape at the valve edge. (**D**)

At the onset of exocytosis, (**D'**) membranes still surround most parts of the new valve, (**D''**) but

they no longer form a complete enclosure (arrowheads). (**E**) Shortly after exocytosis, (**E'**) distal

membranes lose their structural integrity (arrowheads), and (**E''**) the plasma membrane is now

continuous (arrow) under the new valve. (**F-F''**) After completion of exocytosis, there are no traces

left of the distal membrane remnants. Scale bars represent 5 μm (**A, B, C, D, E, F**), 1 μm (**D', E', F'**),

500 nm (**A', C', C'', D'', E'', F''**), 200 nm (**A'', B', B'', C' inset**) and 50 nm (**A'' inset**).

The high-resolution structural observations using electron microscopy strongly suggest

that valve exocytosis involves degradation of the distal membranes in the extracellular

environment (Fig. 1 E). However, these are 2D frozen snapshots that do not reveal direct and

dynamic information about the exocytosis process. Therefore, we investigated the process

dynamics in living cells using time-lapse confocal microscopy. Synchronized cultures were labelled

with PDMPO to visualize newly formed silica, and with FM4-64 to stain the plasma membrane.

FM4-64 is an amphiphilic, membrane-impermeable dye, that can enter the cell only via endocytic

vesicles[35,36]. We followed doubly-labelled cells during valve formation at 20 to 40 minute intervals.

As the cells are sensitive to these imaging conditions we were able to record cells at only 2-5 time

points. Some cells showed the initial stages of silica deposition (Fig. 6 A-C), whereas other were

captured during valve exocytosis (Fig. 6 D-E).

In accordance with the TEM images, live-cell imaging showed that silica deposition starts

at the cell apex and radiates towards the valve edge (Fig. 6 A-B). The outlines of the growing

polygonal layer can be distinguished in the FM4-64 channel since the plasma membrane is tightly

outlining the growing silica structures within the SDV (Fig. 6 B), and the silicification of the linking

extensions is preceded by membrane extension (Fig. 6 C'', arrowheads). In cells that were imaged

from completion of the valve onwards, recently formed linking extensions still have membranes

tightly surrounding them at the initial time point of imaging (Fig. 6 D). However, the membranes

have largely disintegrated a few hours later, only leaving some FM4-64 stained remnants where

the linking extensions connect to each other (Fig. 6 E).

Figure 6. Coordinated silica and membrane dynamics during valve formation in *S. turris* **revealed by 3D reconstructions of time-lapse confocal fluorescence images.** (**A-E**) Images show an overlay of the fluorescence signals from two cells at different stages of valve formation. (**A-C**) A cell during valve formation. During early valve formation, the emerging silica polygons at the apex can be seen by PDMPO staining (**A**). Later on, membranes can be seen outlining the growing polygons and linking extensions (**B**). The growth of linking extensions is led by a membrane outlining their shape (**C**, arrowheads). (**D-E**) A different cell shortly before (**D**) and after exocytosis (**E**) of a mature valve. **A'-E'** show only the FM4-64 fluorescence highlighted in white. (**A''-E''**) Higher magnification of the boxed area in A, B', C, D', and E'. Scale bars represent 20 μm (**A-E'**) and 5 μm (**A''-E''**).

In order to follow the entire valve formation process in the same cell, some adjustments were made to the imaging protocol. Most importantly, PDMPO imaging was omitted to minimize laser damage, and the concentration of FM4-64 was reduced. This improved protocol enabled us to record the membrane dynamics during the entire process of valve formation and exocytosis at a higher temporal resolution (Fig. 7, Supplementary video 1). At the beginning, formation of the valves is perceptible because the plasma membrane outlines the growing polygons and linking

extensions (Fig. 7 A, t=0 to t=126). Exocytosis of the completed valves is manifested by clear

changes to the membrane arrangement (the valve at the top is exocytosed first, between t=126

and t=138, and the second valve is exocytosed around t=168). First, loosening of the plasma

membrane – SDV – silica complex is evident as the labelled plasma membrane no longer outlines

the polygons. At the same time point, we observe a slight enhancement of the fluorescent signal,

possibly because a fusion between the plasma and SDV membranes allows free FM4-64 from the

medium to infiltrate and stain the SDV. Recently exocytosed valves remain temporarily surrounded

by the distal membranes, which gradually disintegrate. These membrane remnants, which are

visible both in the TEM and live-cell images, gradually lose their tight delineation of the valve

structures. This is most evident around the linking extensions as these are located relatively far

away from the cell body. In some cases, it is clear that the membranes surrounding a linking

extension are no longer connected to the cell body (Fig. 7 B, Supplementary Fig. 2). This labelling

pattern confirms that major fractions of the external membranes are completely detached from

the main cell surface, and that exocytosis of the valve is accompanied by massive membrane

shedding into the extracellular space. Additionally, we can rule out the possibility of significant

retrieval of the external membranes into the cell as such a scenario would result in excess FM4-64

labeling within the cell.

Figure 7. Membrane dynamics in a living *S. turris* **cell going through valve formation and exocytosis.** (**A**) 3D reconstructions of time-lapse confocal fluorescence images of FM4-64 (plasma membrane - red) and chloroplast autofluorescence (magenta). (**A'**) The same images as in (**A**) with only membrane staining (FM4-64) highlighted in white at higher magnification. (**B**) Cropped area around one linking extension (indicated with boxes in **A'**). Onset of exocytosis of the valve on top is between t=126 and 132, after which the membrane remnants around the linking process are no longer connected to the cell and disintegrate gradually. Scale bars: 10 μm (**A**), 5 μm (**B**) and 2 μm (**C**).

Discussion

Extrusion of the intracellularly formed, rigid cell walls of diatoms is an extraordinary exocytosis

event, which is difficult to reconcile with known exocytosis processes. The combination of

ultrastructural studies and live-cell imaging allowed us to investigate the membrane dynamics

during valve formation and exocytosis in *S. turris*. Our observations of the SDV ultrastructure show

that the intricate silica patterns are formed within a tightly confined space, bound by the SDV

membrane, supporting the idea that the SDV membrane is controlling the shape of growing silica

structures[37,38]. Furthermore, we provide a direct evidence from living cells for the actual exocytosis

mechanism. Our data show that the valve is exocytosed through fusion of the plasma membrane

and proximal SDV membrane, and that the distal membranes surrounding recently exocytosed

valves gradually disintegrate after exocytosis (Figs. 5, 7). Altogether, our results indicate that

exocytosis of diatom silica involves two remarkable events. First, the repurposing of an organelle

membrane into a plasma membrane (Fig. 1 C) and second, the shedding of a huge amount of

membranes (Fig. 1 E).

The first event, conversion of the proximal SDV membrane into a new plasma membrane,

requires regulated fusion of the SDV and the old plasma membrane. As the SDV and plasma

membranes carry out specialized, different tasks, it is likely that both require unique lipid and

protein compositions, which need to be adjusted after fusion[39]. The TEM data suggest multiple

fusion events all around the valve edge, resulting in the proximal SDV membrane becoming the

new outermost boundary of the cytoplasm. Subsequently, additional fusion events distally to the

valve might serve to break-up the distal SDV membrane and the old plasma membrane. Fusion of

the SDV and plasma membrane at the time of valve exocytosis was demonstrated previously in

live-cell studies of *T. pseudonana*[3,25], implying a general exocytosis strategy.

Regarding the second point, our results show that the distal membranes disintegrate

gradually while they are no longer connected to the cell membrane (Fig. 7, Supplementary Fig. 2).

Therefore, internalization of these membranes through compensatory endocytosis or

macropinocytotic-like invagination is ruled out. FM4-64 attaches only transiently to membrane

lipids, and can rapidly diffuse out from the degrading debris into the surrounding medium, where

its fluorescence is significantly reduced[35]. Therefore, we cannot identify the fate of these

membrane remnants. Important to note is that the parental girdle bands form a quasi-enclosed

compartment around the newly formed valves, possibly slowing down diffusion away from the cell

and facilitating the uptake of the membrane debris for cellular use after disintegration. On first

sight, this exocytosis process including the extracellular disintegration of large amounts of

membrane appears costly to the cell, but it is conceivable that the costs are minimized by recycling

the building blocks from within the space enclosed by the girdle bands.

Exocytosis is an essential cellular process that has been widely studied in a variety of model

systems. In the classical secretory exocytosis systems, the process proceeds via large amounts of

small vesicles that fuse with the membrane[40]. Nevertheless, there are known alternatives related

to larger exocytosis needs. For example, giant vesicles in the *Drosophila* salivary glands squeeze

out their viscous cargo through a wide pore by crumpling of the vesicular membrane, followed by

membrane retrieval by small endocytic vesicles[41]. In the acrosome reaction, the plasma membrane

and distal acrosomal membrane fuse at several locations forming vesicles that are dispersed in the

environment while the proximal acrosomal membrane remains intact and becomes the new

boundary that separates the spermatozoa from the environment[42,43]. Exocytosis of diatom cell

walls is a parallel strategy to these examples, adapted to account for the need to extrude a rigid,

large vesicle content.

To conclude, this work proposes a remarkable mechanism for the exocytosis of the giant

silica deposition vesicle during silicification in diatoms. The size of the studied unicellular alga,

together with advanced fluorescent and electron microscopy, facilitated the elucidation of the

exocytosis steps. With the toolbox for genetic research in diatom growing, it will soon be possible

to investigate whether similar conserved processes are involved in the regulation of this event as

in classical exocytosis.

Materials and methods

Cell cultures

S. turris was isolated from the North Sea in 2004 and provided by the group of Prof. Eike Brunner, TU Dresden. *S. turris* cultures were maintained in natural Mediterranean seawater that was filtered, its salinity corrected to 3.5% and supplemented with f/2 nutrient recipe. Cultures were maintained at 18°C under 16/8 hours light/dark cycles, by a weekly dilution of 5 ml from a parent culture into 45 ml fresh medium.

Cell cycle synchronization

For synchronization of the cell cycle, 5 ml of a mature cell culture was used to start a new 50 ml culture. After 48 hours of growth under the normal 16L/8D cycle, the culture was placed in darkness for 20 hours. After 20 hours of darkness, the culture was again exposed to light. To track the formation of new silica, PDMPO [2-(4-pyridyl)-5-((4-(2-dimethylaminoethyl-amino-carbamoyl)methoxy)-phenyl)oxazole] (ThermoFisher Scientific, USA) was added to a final concentration of 330 µM. PDMPO fluorescence was monitored by imaging the cultures with an epifluorescence microscope (Nikon Eclipse Ni-U, ex: 365 em: 525). Four cultures were imaged (at least 100 cells per time point) every 2 hours until all of the cells had formed new valves (about 14 hours).

Sample preparation for SEM

Cells were prepared for SEM in two different ways. For the first, quick method, 10 µl of culture was pipetted onto polycarbonate Whatman® Nuclepore™ Track-Etched membranes, which were then dried using a vacuum filter. To prepare cells in which the girdle band region does not collapse, the second method uses critical point drying (CPD). To prepare cells for CPD, cells were fixed in a solution of 2% Glutaraldehyde and 4% Paraformaldehyde in artificial seawater for 1 hour at room temperature while shaking. Then cells were washed 3 times with deionized water (Milli-Q® IQ 7003 Ultrapure Lab Water System, Merck), and dehydrated by washing them in a graded series of ethanol. The final wash was done in 100% anhydrous ethanol overnight. The dehydrated samples

were then dried in a critical point dryer using liquid CO_2 as transitional fluid. The Track-Etched

membranes or dried cells were then placed onto a conductive carbon tape on an aluminium stub.

SEM imaging

The samples were sputter-coated with 4 nm iridium (Safematic). The dried and coated cell wall

samples were imaged with Sigma 500 or Ultra 55 FEG scanning electron microscopes (Zeiss,

Germany), using 3-5 kV, aperture size 20-30 um and a working distance of about 3 mm.

Sample preparation for TEM

To preserve the cells in the native state they were cryo-fixed, using high pressure freezing (HPF),

followed by freeze substitution (FS), according to previously published protocols [31,44]. HPF ensures

rapid vitrification of the cells preventing the formation of ice crystals, which would damage cellular

structures. Through FS, the vitrified ice is replaced with an organic solvent and fixatives.

Synchronized cultures were harvested by collecting the cells on a 5 µm filter membrane. The cells

were collected from the membrane by washing them into an Eppendorf tube using 200 µl of

seawater. After letting the cells sediment for 10 minutes, 2 µl aliquots were pipetted into aluminium

discs (Wohlwend GmbH, Sennwald, Switzerland) and directly loaded into a Leica ICE high pressure

freezing machine (Leica Microsystems GmbH, Wetzlar, Germany). The samples were vitrified in

liquid nitrogen (-192 °C) at 210 MPa (2048 bar). The vitrified samples were stored in liquid nitrogen

until freeze-substitution in an EM ASF2 (Leica Microsystems GmbH, Wetzlar, Germany). The

vitrified water in the sample was substituted with an organic solvent, 100% anhydrous acetone, at

(-90 °C). The acetone was supplemented with chemical fixatives (0.2% uranyl acetate and 0.2%

osmium tetroxide) to enhance cross-linking and contrast of cellular structures. The samples were

immersed in the solution for 48 hours at -90 °C. Then, the samples were allowed to gradually

warm for 24 hours to -20°C, and then gradually warmed to 0°C during one hour. The samples were

washed 3 times with acetone. Then the acetone was replaced with Epon (Agar Scientific Ltd,

Stansted, U.K.) using gradient concentration mixtures (10%, 20%, 30%, 40%, 60%, 80%, 100% Epon

in acetone), twice a day at room temperature. The sample in 100% Epon was hardened at 70 °C

for 72 h. Ultrathin sections of 70 nm were sliced using an ultra-microtome (Ultracut UCT, Leica

Microsystems GmbH, Wetzlar, Germany) equipped with a diamond knife (Ultra 45°, Diatome Ltd,

Nidau, Switzerland). The sections were picked up onto copper TEM grids, coated with carbon film.

The sections were then post-stained by placing them onto a drop of lead citrate solution. After

three minutes of staining, the grids were washed three times in drops of water and then dried by

blotting onto whatman filter paper. In total, we prepared 16 frozen samples that were further

processed using freeze substitution during 3 different cycles, and subsequently imaged hundreds

of cells.

TEM imaging

The TEM samples were imaged with a Tecnai Spirit transmission electron microscope (FEI,

Eindhoven, Netherlands) operated at 120 kV and equipped with Gatan Oneview 4 k × 4 k camera

(Gatan Inc., Pleasanton, U.S.A.).

Live cell imaging

For single-cell time-lapse imaging, a culture was synchronized and stained with PDMPO (330 µM).

About 8 hours after the end of light starvation, when most cells had gone through cytokinesis, an

aliquot of 100 µl was taken and FM4-64 (ThermoFisher Scientific, USA) was added to a final

concentration of 4-8 µM to stain the membranes. After adding the membrane dye, a 20 µl drop

was mounted on a microscope slide and covered with a glass coverslip, using dental wax as spacer.

The samples were visualized using a Leica TCS SP8-STED confocal microscope equipped with a

HCS PL APO 86x/1.20W motCORR objective. FM4-64 and chlorophyll autofluorescence were

acquired by white-light laser using 550 nm laser line (6% laser power) and 650 nm laser line (7%

laser power), respectively. PDMPO fluorescence was acquired using a 405 nm laser (5% laser

power). HyD-SMD detectors were used for PDMPO and FM4-64 with emission collection width set

to 474-530 and 608-640 nm, respectively. Chlorophyll autofluorescence emission was collected

using a HyD detector with 741-779 nm detection width and the transmission channel was detected

with a PMT detector. The cells were imaged for 2 to 4 hours at intervals of 3 to 60 minutes. Images

were analysed using Leica Application Suite X. In total, Time lapses from over 50 cells were

collected over the entire period.

Acknowledgements

We thank Anne Jantschke and Nathalie Pytlik from TU Dresden for providing cultures. We thank Simon Michaeli from the Volcani Institute (ARO) for his contribution in finding the right live cell imaging platform. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 848339). D. dH. was supported by the Sustainability and Energy Research Initiative (SAERI) of Weizmann Institute of Science.

Data availability

All data generated or analysed during this study are included in the manuscript and supporting file; Source Data files for Figures 2-7 have been deposited to Dryad.

Additional information

Supplementary Information accompanies this paper at https://doi.org/XXX.XXX

Competing interests

The authors declare no competing interests.

References

1. Armbrust, E. V. The life of diatoms in the world's oceans. *Nature* **459**, 185–192 (2009).

2. Kroger, N. & Poulsen, N. Diatoms — From Cell Wall Biogenesis to Nanotechnology. *Annu. Rev. Genet.* **42**, 83–107 (2008).

3. Kotzsch, A. *et al.* Silicanin-1 is a conserved diatom membrane protein involved in silica biomineralization. *BMC Biol.* **15**, (2017).

4. Kröger, N., Deutzmann, R. & Sumper, M. Polycationic peptides from diatom biosilica that direct silica nanosphere formation. *Science (80-.).* **286**, 1129–1132 (1999).

5. Sumper, M., Brunner, E. & Lehmann, G. Biomineralization in diatoms: Characterization of novel polyamines associated with silica. *FEBS Lett.* **579**, 3765–3769 (2005).

6. Hildebrand, M. Diatoms, Biomineralization Processes , and Genomics. *Chem. Rev.* **108**, 4855–4874 (2008).

7. Drum, R. W. & Pankratz, H. S. Post mitotic fine structure of Gomphonema parvulum. *J.*

Ultrasructure Res. **10**, 217–223 (1964).

8. Heintze, C. *et al.* An intimate view into the silica deposition vesicles of diatoms. *BMC*

Mater. **2**, 1–15 (2020).

9. Mayzel, B., Wolf, S. G., Varsano, N. & Gal, A. Extracellular silica formation by diatoms.

10. Hildebrand, M., Lerch, S. J. L. & Shrestha, R. P. Understanding Diatom Cell Wall

Silicification — Moving Forward. *Front. Mar. Sci.* **5**, 1–19 (2018).

11. Kröger, N., Lorenz, S., Brunner, E. & Sumper, M. Self-assembly of highly phosphorylated

silaffins and their function in biosilica morphogenesis. *Science (80-.).* **298**, 584–586 (2002).

12. Scheffel, A., Poulsen, N., Shian, S. & Kröger, N. Nanopatterned protein microrings from a

diatom that direct silica morphogenesis. *Proc. Natl. Acad. Sci.* **108**, 3175–3180 (2011).

13. Montagna, M. *et al.* Interactions of long-chain polyamines with silica studied by molecular

dynamics simulations and solid-state NMR spectroscopy. *Langmuir* **36**, 11600–11609

(2020).

14. Jantschke, A. *et al.* Insight into the Supramolecular Architecture of Intact Diatom Biosilica

from DNP-Supported Solid-State NMR Spectroscopy. *Angew. Chemie Int. Ed.* **54**, 15069–

15073 (2015).

15. Stoermer, E. F., Pankratz, H. S. & Bowen, C. C. Fine Structure of the Diatom Amphipleura

pellucida. II. Cytoplasmic Fine Structure and Frustule Formation. **52**, 1067–1078 (1965).

16. Battey, N. H., James, N. C., Greenland, A. J. & Brownlee, C. Exocytosis and Endocytosis.

Plant Cell **11**, 643–659 (1999).

17. Schulz, D. Das Schicksal von Silicalemma und Plasmalemma während der Zellwandbildung

bei Coscinodiscus granii Gough und Thalassiosira eccentrica (Ehrenb.) Cleve. *Ber. Dtsch.*

Bot. Ges. **97**, 411–419 (1984).

18. Volcani, B. E. Cell Wall Formation in Diatoms: Morphogenesis and Biochemistry. in *Silicon*

and Siliceous Structures in Biological Systems 157–200 (Springer, 1981).

19. Bedoshvili, Y. D. *et al.* Frustule morphogenesis of raphid pennate diatom Encyonema

ventricosum (Agardh) Grunow. *Protoplasma* **255**, 911–921 (2018).

20. Chiappino, M. L. & Volcani, B. E. Studies on the Biochemistry and Fine Structure of Silicia

Shell Formation in Diatoms VII. Sequential Cell Wall Development in the Pennate Navicula pelliculosa. *Protoplasma* **93**, 205–221 (1977).

21. Crawford, R. M. Valve Formation in Diatoms and the Fate of the Silicalemma and Plasmalemma. *Protoplasma* **106**, 157–166 (1981).

22. Dawson, P. A. Observations on the structure of some forms of Gomphonema Parvulum Kutz. III. Frustule formation. *J. Phycol.* **9**, 353–365 (1973).

23. Reimann, B. E. F., Lewin, J. C. & Volcani, B. E. Studies on the Biochemistry and fine structure of silica shell formation in diatoms I. The Structure of the Cell Wall of Cylindrotheca fusiformis. *J. Cell Biol.* **24**, 39–55 (1965).

24. Schmid, A.-M. M. Wall Morphogenesis in Coscinodiscus wailesii. *8th Diatom Symp.* 293–314 (1984).

25. Yee, D. P., Hildebrand, M. & Tresguerres, M. Dynamic subcellular translocation of V-type H+-ATPase is essential for biomineralization of the diatom silica cell wall. *New Phytol.* **225**, 2411–2422 (2020).

26. Li, C. W. & Volcani, B. E. Studies on the Biochemistry and Fine Structure of Silica Shell Formation in Diatoms. X Morphogenesis of the Labiate Process in Centric Diatoms. *Protoplasma* **124**, 147–156 (1985).

27. Gröger, P., Poulsen, N., Klemm, J., Kröger, N. & Schlierf, M. Establishing super-resolution imaging for proteins in diatom biosilica. *Sci. Rep.* **6**, 1–8 (2016).

28. Kroth, P. G. *et al.* Genome editing in diatoms: achievements and goals. *Plant Cell Rep.* **37**, 1401–1408 (2018).

29. Chepurnov, V. A. *et al.* In search of new tractable diatoms for experimental biology. *BioEssays* **30**, 692–702 (2008).

30. von Stosch, H. A. & Drebes, G. Entwicklungsgeschichtliche Untersuchungen an zentrischen Diatomeen IV. *Helgoländer Wissenschaftliche Meeresuntersuchungen* **11**, 209–257 (1964).

31. Kadan, Y. *et al.* In situ electron microscopy characterization of intracellular ion pools in mineral forming microalgae. *J. Struct. Biol.* **210**, 107465 (2020).

32. Brzezinski, M. A., Olson, R. J. & Chisholm, S. W. Silicon availability and cell-cycle progression in marine diatoms. *Mar. Ecol. Prog. Ser.* **67**, 83–96 (1990).

503   33.   Vaulot, D., Olson, R. J. & Chisholm, S. W. Light and dark control of the cell cycle in two
504   marine phytoplankton species. *Exp. Cell Res.* **167**, 38–52 (1986).

505   34.   Shimizu, K., Del Amo, Y., Brzezinski, M. A., Stucky, G. D. & Morse, D. E. A novel fluorescent
506   silica tracer for biological silicification studies. *Chem. Biol.* **8**, 1051–1060 (2001).

507   35.   Bolte, S. *et al.* FM-dyes as experimental probes for dissecting vesicle trafficking in living
508   plant cells. *J. Microsc.* **214**, 159–173 (2004).

509   36.   Kühn, S. F. & Brownlee, C. Membrane organisation and dynamics in the marine diatom
510   Coscinodiscus wailesii (Bacillariophyceae). *Bot. Mar.* **48**, 297–305 (2005).

511   37.   Tesson, B. & Hildebrand, M. Extensive and intimate association of the cytoskeleton with
512   forming silica in diatoms: Control over patterning on the meso- and micro-scale. *PLoS*
513   *One* **5**, (2010).

514   38.   Van de Meene, A. M. L. & Pickett-Heaps, J. D. Valve morphogenesis in the centric diatom
515   Proboscia alata Sundstrom. *J. Phycol.* **38**, 351–363 (2002).

516   39.   Holthuis, J. C. M. & Menon, A. K. Lipid landscapes and pipelines in membrane
517   homeostasis. *Nature* **510**, 48–57 (2014).

518   40.   Gundelfinger, E. D., Kessels, M. M. & Qualmann, B. Temporal and spatial coordination of
519   exocytosis and endocytosis. *Nat. Rev. Mol. Cell Biol.* **4**, 127–139 (2003).

520   41.   Kamalesh, K. *et al.* Exocytosis by vesicle crumpling maintains apical membrane
521   homeostasis during exocrine secretion. *Dev. Cell* **56**, 1603-1616.e6 (2021).

522   42.   Okabe, M. The Acrosome Reaction: A Historical Perspective. in *Sperm Acrosome*
523   *Biogenesis and Function During Fertilization* (ed. Buffone, M. G.) 1–13 (Springer
524   International Publishing, 2016). doi:10.1007/978-3-319-30567-7_1

525   43.   Tomes, C. N. Acrosomal Exocytosis. in *Molecular Mechanisms of Exocytosis* 117–147
526   (Springer New York, 2007). doi:10.1007/978-0-387-39961-4_9

527   44.   Weston, A. E., Armer, H. E. J. & Collinson, L. M. Towards native-state imaging in biological
528   context in the electron microscope. *J. Chem. Biol.* **3**, 101–112 (2010).

Supplementary information

Supplementary figure 1. Cell cycle of *S. turris.* New silica elements, formed during the current cell cycle are coloured green. Dashed lines represent girdle bands. (**A**) Start of a cell cycle with a protoplast enclosed by the two parental valves, the valve edges are directly adjacent to each other. The older valve is on the right, and the younger valve on the left. The younger valve is covered under the girdle bands of the older valve. (**B**) Protoplast starts growing in preparation for the next cell division. As the silica valves are rigid and not elastic, growth is only possible in longitudinal direction by pushing apart the two valves. New girdle bands (green) are formed in individual SDVs and added to the edge of the younger valve. (**C**) After the protoplast has reached its final length, the cell undergoes cytokinesis, resulting in two daughter cells enclosed in the parental cell wall.

(**D**) Each daughter cell inherits one half of the parental cell wall and starts forming a new valve,

inside an SDV, shortly after cell division. (**E**) The new valves are completed with attachment of the

linking extensions, after which they are exocytosed. (**F**) Both daughter cells are now ready to

continue to the next cell cycle.

Supplementary figure 2. Distal membranes that undergo degradation are disconnected

from the cellular membrane. Representative (**A**) maximum intensity projections and (**A'**) single

z-slices of time-lapse confocal fluorescence images. Images show membrane staining (FM4-64) in

white and chloroplasts autofluorescence in magenta. The elapsed time since the start of imaging

(in minutes) is shown on the upper right corner. Exocytosis of the valve on the bottom right starts

between t=42 and t=48. The valve on the top left is exocytosed between t=48 and t=54. In t=72

and t=90, distal membranes around the valve are not connected to the cellular membrane. At

t=102 the remnants have largely disintegrated. Scale bar: 10 µm

Supplementary video 1. Time-lapse imaging video of *S. turris* cell, stained with FM4-64, during

valve formation and exocytosis. (MP4 2237 kb)