# A data compendium of *Mycobacterium tuberculosis* antibiotic resistance

Alice Brankin[4,*,**] and Kerri M Malone[19,*,**], Ivan Barilar[29], Simone Battaglia[1], Emanuele Borroni[1], Angela Pires Brandao[2,3], Andrea Maurizio Cabibbe[1], Joshua Carter[5], Daniela Maria Cirillo[1], Pauline Claxton[6], David A Clifton[4], Ted Cohen[7], Jorge Coronel[8], Derrick W Crook[4], Viola Dreyer[29], Sarah G Earle[4], Vincent Escuyer[9], Lucilaine Ferrazoli[3], Philip W Fowler[4], George Fu Gao[10], Jennifer Gardy[11], Saheer Gharbia[12], Kelen Teixeira Ghisi[3], Arash Ghodousi[1,13], Ana Luíza Gibertoni Cruz[4], Louis Grandjean[33], Clara Grazian[14], Ramona Groenheit[44], Jennifer L Guthrie[15,16], Wencong He[10], Harald Hoffmann[17,18], Sarah J Hoosdally[4], Martin Hunt[19,4], Zamin Iqbal[19], Nazir Ahmed Ismail[20], Lisa Jarrett[21], Lavania Joseph[20], Ruwen Jou[22], Priti Kambli[23], Rukhsar Khot[23], Jeff Knaggs[19,4], Anastasia Koch[24], Donna Kohlerschmidt[9], Samaneh Kouchaki[4,25], Alexander S Lachapelle[4], Ajit Lalvani[26], Simon Grandjean Lapierre[27], Ian F Laurenson[6], Brice Letcher[19], Wan-Hsuan Lin[22], Chunfa Liu[10], Dongxin Liu[10], Ayan Mandal[28], Mikael Mansjö[44], Daniela Matias[21], Graeme Meintjes[24], Flávia de Freitas Mendes[3], Matthias Merker[29], Marina Mihalic[18], James Millard[30], Paolo Miotto[1], Nerges Mistry[28], David Moore[31,8], Kimberlee A Musser[9], Dumisani Ngcamu[20], Hoang Ngoc Nhung[32], Stefan Niemann[29,48], Kayzad Soli Nilgiriwala[28], Camus Nimmo[33], Nana Okozi[20], Rosangela Siqueira Oliveira[3], Shaheed Vally Omar[20], Nicholas Paton[34], Timothy EA Peto[4], Juliana Maira Watanabe Pinhata[3], Sara Plesnik[18], Zully M Puyen[35], Marie Sylvianne Rabodoarivelo[36], Niaina Rakotosamimanana[36], Paola MV Rancoita[13], Priti Rathod[21], Esther Robinson[21], Gillian Rodger[4], Camilla Rodrigues[23], Timothy C Rodwell[37,38], Aysha Roohi[4], David Santos-Lazaro[35], Sanchi Shah[28], Thomas Andreas Kohl[29], Grace Smith[21,12], Walter Solano[8], Andrea Spitaleri[1,13], Philip Supply[39], Utkarsha Surve[23], Sabira Tahseen[40], Nguyen Thuy Thuong Thuong[32], Guy Thwaites[32,4], Katharina Todt[18], Alberto

Trovato[1], Christian Utpatel[29], Annelies Van Rie[41], Srinivasan Vijay[42], Timothy M Walker[4,32], A

Sarah Walker[4], Robin Warren[43], Jim Werngren[44], Maria Wijkander[44], Robert J Wilkinson[45,46,26],

Daniel J Wilson[4], Penelope Wintringer[19], Yu-Xin Xiao[22], Yang Yang[4], Zhao Yanlin[10], Shen-Yuan

Yao[20], Baoli Zhu[47]

**Institutions**

[1] IRCCS San Raffaele Scientific Institute, Milan, Italy

[2] Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

[3] Institute Adolfo Lutz, São Paulo, Brazil

[4] University of Oxford, Oxford, UK

[5] Stanford University School of Medicine, Stanford, USA

[6] Scottish Mycobacteria Reference Laboratory, Edinburgh, UK

[7] Yale School of Public Health, Yale, USA

[8] Universidad Peruana Cayetano Heredia, Lima, Perú

[9] Wadsworth Center, New York State Department of Health, Albany, USA

[10] Chinese Center for Disease Control and Prevention, Beijing, China

[11] Bill & Melinda Gates Foundation, Seattle, USA

[12] UK Health Security Agency, London, UK

[13] Vita-Salute San Raffaele University, Milan, Italy

[14] University of New South Wales, Sydney, Australia

[15] The University of British Columbia, Vancouver, Canada

[16] Public Health Ontario, Toronto, Canada

[17] SYNLAB Gauting, Munich, Germany

[18] Institute of Microbiology and Laboratory Medicine, IMLred, WHO-SRL Gauting, Germany

[19] EMBL-EBI, Hinxton, UK

[20] National Institute for Communicable Diseases, Johannesburg, South Africa

[21] Public Health England, Birmingham, UK

[22] Taiwan Centers for Disease Control, Taipei, Taiwan

[23] Hinduja Hospital, Mumbai, India

[24] University of Cape Town, Cape Town, South Africa

[25] University of Surrey, Guildford, UK

[26] Imperial College, London, UK

[27] Université de Montréal, Canada

[28] The Foundation for Medical Research, Mumbai, India

[29] Research Center Borstel, Borstel, Germany

[30] Africa Health Research Institute, Durban, South Africa

[31] London School of Hygiene and Tropical Medicine, London, UK

[32] Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

[33] University College London, London, UK

[34] National University of Singapore, Singapore

[35] Instituto Nacional de Salud, Lima, Perú

[36] Institut Pasteur de Madagascar, Antananarivo, Madagascar

[37] FIND, Geneva, Switzerland

[38] University of California, San Diego, USA

[39] Institut Pasteur de Lille, Lille, France

[40] National TB Reference Laboratory, National TB Control Program, Islamabad, Pakistan

[41] University of Antwerp, Antwerp, Belgium

[42] University of Edinburgh, Edinburgh, UK

[43] Stellenbosch University, Cape Town, South Africa

[44] Public Health Agency of Sweden, Solna, Sweden

[45] Wellcome Centre for Infectious Diseases Research in Africa, Cape Town, South Africa

[46] Francis Crick Institute, London, UK

[47] Institute of Microbiology, Chinese Academy of Sciences, Beijing, China

[48] German Center for Infection Research (DZIF), Hamburg-Lübeck-Borstel-Riems, Germany

*equal contribution authors

**co-corresponding authors

## Abstract

The Comprehensive Resistance Prediction for Tuberculosis: an International

Consortium (CRyPTIC) presents here a global collection of 15,211 *Mycobacterium*

*tuberculosis* clinical isolates, all of which have undergone whole genome sequencing

and have had their minimum inhibitory concentrations to 13 antitubercular drugs

measured. The isolates represent five major *M. tuberculosis* lineages originating from

23 countries across four continents. 6,814 isolates were found resistant to at least one

drug, including 2,129 samples fully satisfy the clinical definitions of RR/MDR, pre-XDR

or XDR.  Resistance status to eight antitubercular drugs can be accurately predicted

using a genetic mutation catalogue for over 90% of the isolates. Furthermore, we show

the presence of suspected resistance conferring mutations for isolates resistant to the

newly introduced drugs bedaquiline, clofazimine, delamanid and linezolid. Finally, a

case study of rifampicin mono-resistance is presented to showcase how this

compendium could be used to advance our genetic understanding of rare resistance

phenotypes and evaluate the likely performance of a widely used molecular diagnostic

tool. It is hoped that this compendium, the largest *M. tuberculosis* matched phenotypic

and genotypic dataset to date, will facilitate and inspire new research projects for years

to come.

**Introduction**

Tuberculosis (TB) is a curable and preventable disease; 85% of those afflicted

can be successfully treated with a six-month regimen. Despite this, TB is the world's

top infectious disease killer with 10 million new cases and 1.2 million deaths estimated

in 2019 alone [1]. Furthermore, drug resistant TB (DR-TB) is a continual threat; almost

half a million people were estimated to have developed resistance to the first-line drug

rifampicin (RR-TB), with three quarters of these cases estimated to be multidrug-

resistant (MDR-TB, resistant to first-line drugs isoniazid and rifampicin) [1]. Worryingly,

only 44% of DR-TB cases were officially notified and just over half of these cases were

successfully treated (57%) [1].

To address these issues, the World Health Organisation (WHO) is encouraging

the development of better, faster and more targeted diagnostic and treatment

strategies through its EndTB campaign. Of particular interest is universal drug

susceptibility testing (DST). Conventionally, DST relies on lengthy (4 weeks minimum)

culture-based methods that require strict biosafety conditions for *Mycobacterium*

*tuberculosis.* The development of rapid genetics-based assays has decreased

diagnostic time to as little as 2 hours through the detection of specific resistance

conferring mutations *e.g.* the Cepheid Xpert® MTB/RIF test [2, 3]. However, assay

bias towards specific genic regions can result in misdiagnosis of resistance, the

prescription of ineffective treatment regimens and subsequent spread of multi-drug

resistant disease, as seen during an outbreak in Eswatini of an MDR strain harbouring

an *rpoB* I491F mutation not detected by the Xpert® MTB/RIF assay [4-6].

Furthermore, detection of rifampicin resistance with the Xpert® MTB/RIF line

probe assay (LPA) assay is used to infer MDR-TB as epidemiologically, rifampicin

resistance tends to coincide with resistance to isoniazid [7]. While this *modus operandi*

is successful at pragmatically identifying potential MDR cases quickly and effectively,

it is not generally true that a single path towards developing MDR or e̲xtensively d̲rug

r̲esistant TB (XDR = MDR/RR + resistance to at least one fluoroquinolone and either

bedaquiline or linezolid) exists. For example, *M. tuberculosis* strains that are sensitive

to rifampicin but resistant to multiple other anti-tubercular drugs would be missed by

current LPA diagnostic testing and remain undetected in the community. A better

understanding of how various combinations of drug resistance in different

backgrounds (nonMDR/MDR) arises is essential to not only prevent the spread of

resistant strains and to promote the use of effective treatment regimens, but also to

facilitate the development of suitable diagnostic assays unbiased in their approaches

to detect *any* and *all* resistance where it occurs.

Whole-genome sequencing (WGS) has the potential to reveal the entirety of

the *M. tuberculosis* genetic resistance landscape for any number of drugs

simultaneously whilst enabling a more rapid turnaround time and affordable cost in

comparison to DST culture methods [8]. The success of WGS as a diagnostic method

wholly depends on a comprehensive and accurate catalogue of resistance-conferring

mutations for each drug. Recent advances have shown that genotypic predictions of

resistance correlate well with DST results for first-line drugs [7]. However, the

mechanisms of resistance to second-line, new and re-purposed drugs (NRDs) are less

well understood despite their increased administration in clinics as MDR cases climb

[9]. To align with the goals of the EndTB campaign, further research is vital to construct

an extensive catalogue of genetic variants associated with resistance, and indeed

susceptibility, if WGS-based diagnostics are to become routine practice for DR-TB

management.

The Comprehensive Resistance Prediction for Tuberculosis: an International

Consortium (CRyPTIC) has collected *M. tuberculosis* clinical isolates worldwide to

survey the genetic variation associated with resistance to 13 antitubercular drugs,

specifically the first-line drugs rifampicin, isoniazid, ethambutol, the second-line drugs

amikacin, kanamycin, rifabutin, levofloxacin, moxifloxacin, ethionamide, and the new

and re-purposed drugs bedaquiline, clofazimine, delamanid and linezolid. Here, we

present a data compendium of 15,211 global clinical isolates, each of which has had

its genomic sequence determined and DST profile measured [10]. This compendium

is the largest drug screening effort to date for *M. tuberculosis* in a 'one isolate – one

microscale assay' format across defined compound concentration ranges. Finally, this

matched genotypic-phenotypic dataset will prove an invaluable resource to accelerate

and improve AMR diagnostic development for TB, both by the enrichment of mutation

catalogues for WGS resistance prediction and the identification of important diagnostic

gaps and drug resistance patterns.

**Methods**

*Ethics*

Approval for CRyPTIC study was obtained by Taiwan Centers for Disease Control IRB

No. 106209, University of KwaZulu Natal Biomedical Research Ethics Committee

(UKZN BREC) (reference BE022/13) and University of Liverpool Central University

Research Ethics Committees (reference 2286), Institutional Research Ethics

Committee (IREC) of The Foundation for Medical Research, Mumbai (Ref nos.

FMR/IEC/TB/01a/2015 and FMR/IEC/TB/01b/2015), Institutional Review Board of

P.D. Hinduja Hospital and Medical Research Centre, Mumbai (Ref no. 915-15-CR

[MRC]), scientific committee of the Adolfo Lutz Institute (CTC-IAL 47-J / 2017) and in

the Ethics Committee (CAAE: 81452517.1.0000.0059) and Ethics Committee review

by Universidad Peruana Cayetano Heredia (Lima, Peru) and LSHTM (London, UK).

***Sample collection***

Participating collection centres varied in their isolate collection approaches, but

the consortium collectively aimed to oversample for *M. tuberculosis* isolates with drug

resistance and multi-drug resistance.

188
***Plate assay***

The CRyPTIC consortium designed two variants of the Sensititre MYCOTB

plate (Thermo Fisher Scientific Inc., USA) named the "UKMYC5" and "UKMYC6"

microtitre plates [10, 11]. These plates contain five to ten doubling dilutions of 13

antibiotics (rifampicin (RIF), rifabutin (RFB), isoniazid (INH), ethambutol (EMB),

levofloxacin (LEV), moxifloxacin (MXF), amikacin (AMI), kanamycin (KAN),

ethionamide (ETH), clofazimine (CFZ), linezolid (LZD), delamanid (DLM), and

bedaquiline (BDQ)). Delamanid and bedaquiline were provided by Otsuka

Pharmaceutical Co., Ltd. and Janssen Pharmaceutica respectively. The UKMYC5

plate also contained para-aminosalicylic acid (PAS), however the MICs were not

reproducible and hence it was excluded from the UKMYC6 plate design and is not

included in any subsequent analysis [10].

A standard operating protocol for sample processing was defined by CRyPTIC

as previously described [10, 11]. Clinical samples were sub-cultured using 7H10 agar

plates, Lowenstein-Jensen tubes or MGIT tubes. Bacterial cell suspensions (0.5

McFarland standard, saline Tween) prepared from (no later than) 14-day old colonies

were diluted 100X in 10 ml enriched 7H9 broth prior to plate inoculation. A semi-

automated Sensititre Autoinoculator (Thermo Fisher, Scientific Inc., USA) was used to

inoculate 100 µl prepared cell suspensions (1.5 x $10^5$ CFU/ml [5 x $10^4$ CFU/ml - 5 x

$10^5$ CFU/ml]) into each well of a UKMYC5/6 microdilution plate. The plate was sealed

and incubated for 14 days at 37°C. Quality control runs were performed periodically

using *M. tuberculosis* H37Rv ATCC 27294, which is sensitive to all drugs on the plates.

***Minimum Inhibitory Concentration (MIC) measurement***

As stated in reference [10], minimum inhibitory concentrations (MICs) for each

drug were read after incubation for 14 days by a laboratory scientist using a Thermo

Fisher Sensititre™ Vizion™ digital MIC viewing system. The Vizion apparatus was

also used to take a high contrast photograph of the plate with a white background,

from which the MIC was measured again using the Automated Mycobacterial Growth

Detection Algorithm (AMyGDA) software [12]. The AMyGDA software detects the

boundaries of each well using a Hough transform for circles and measures growth as

the number of dark pixels within the area contained by this boundary. All images where

the MICs measured by Vizion and AMyGDA were different were uploaded to a citizen

science project, BashTheBug, on the Zooniverse platform [13]. Each image was then classified by ≥11 volunteers and the median classification taken.

MICs were then classified as high (at least two methods concur on the MIC), medium (either there is no plate image, or Vizion and AMyGDA disagree and there is no BashTheBug measurement), or low (all three methods disagree) quality. When considering the pattern of resistance in a sample we took the MIC from the Vizion reading, if it was annotated as having medium or low quality, to ensure adequate data coverage.

### Genomic data processing and variant calling

Sequencing arrangements differed slightly between each CRyPTIC participating laboratory. Short-read, paired end libraries were sequenced on Illumina machines and the resulting FASTQ files were processed using the bespoke pipeline Clockwork (v0.8.3, https://github.com/iqbal-lab-org/clockwork). Briefly, all raw sequencing files are indexed into a relational database with which Clockwork proceeds. Human, nasopharyngeal flora and human immunodeficiency virus related reads are removed and remaining reads are trimmed (adapters and low quality ends) using Trimmomatic and mapped with BWA-MEM to the *M. tuberculosis* H37Rv reference genome (NC000962.3) [14, 15]. Read duplicates are removed. Genetic variants are called independently using Cortex and SAMtools, two variant callers with orthogonal strengths (SAMtools a high sensitivity SNP caller, and Cortex a high specificity SNP and indel caller) [16, 17]. The two call sets are merged to produce a final call set, using the Minos adjudication tool to resolving loci where the two callers disagreed, by remapping reads to an augmented genome containing each alternative allele [18]. Default filters of a minimum depth of 5x and a Fraction of Supporting Reads

of 0.9 (minos) and a Genotype confidence percentile (GCP) filter of 0.5 were applied.

The GCP filter is, roughly speaking, a normalised likelihood ratio test, giving a measure

of confidence in the called allele compared with the other alternatives, and is described

in [18]. This produced one VCF file per sample, each only describing positions where

that sample differed from the reference.

These filtered VCFs were then combined, to produce a single non-redundant

list of all variants seen in the cohort. All samples were then processed a second time

with Minos, remapping reads to a graphical representation of all the segregating

variation within the cohort, generating VCF files which had an entry at all variable

positions (thus for all samples, most positions would be genotyped as having the

reference allele).

In order to remove untrustworthy loci, a genome mask was applied to the

resulting VCF files (regions identified with self-blast matches in [19] comprising of

324,971 bp of the reference genome. Furthermore, positions with less than 90% of

total samples passing default Clockwork/Minos variant call filters (described above)

were filtered out, comprising 95,703 bp of the genome, of which 55,980 bp intersect

with the genome mask.

**_Phylogenetic tree construction_**

A pairwise genetic distance matrix was constructed for 15,211 isolates by

comparing each two variant call format (VCF) files (relative to the reference _M._

_tuberculosis_ H37Rv genome, see [11] and the above methods section for more details)

under consideration position-wise. A neighbourhood-joining tree was constructed from

the distance matrix using *quicktree* [20]. Tree visualisation and annotation was

performed using the R library *ggtree* [21]. *M. tuberculosis* lineages were assigned

using Mykrobe and are represented by the coloured dots at the branch termini of the

tree [18]. For isolates that had 'mixed' lineage classification (*i.e.,* 2 lineages were found

present in the sample by Mykrobe, $n$ = 225, 1.5%), the first of the two lineages was

assigned to the isolate. *ggtree* was also used to construct the trees depicting

bedaquiline, clofazimine and delamanid resistant isolates.

***Data analysis and visualisation***

All data can be found in <ftp address: tbc> and were analysed and visualised

using either R or python3 libraries and packages. See <github: tbc> for codebase.

# 284   Results

### 285   15,211 M. tuberculosis clinical isolates

The CRyPTIC dataset comprises isolates from 27 countries collected by 14

partner sites worldwide. The compendium presented here contains 15,211 isolates for

which both genomic and phenotypic data was collected by 23 of the partner countries

across the continents of Asia, Africa, South America and Europe (Fig. 1). Where the

origin of an isolate was not known, the isolate was assumed to originate in the same

country as the contributing laboratory (269 isolates in Germany, 17 isolates in India, 6

isolates in Peru, 885 isolates in Italy, 510 isolates in South Africa, 357 isolates in

Sweden, 208 isolates in Taiwan, 1 isolate in Brazil and 4 isolates in the UK). The

largest number of isolates were contributed by India ($n$ = 4,004), Peru ($n$ = 2,691),

South Africa ($n$ = 2,155), Vietnam ($n$ = 1,288) and China ($n$ = 1,121).

Lineage assignment revealed that all isolates (except 41, 0.3%) belong to the

four main *M. tuberculosis* lineages (L1-L4). Like previous studies, we see a strong

association between geolocation and lineage (Pearson's chi-squared test, $p < 2.2e-$

$16$, Fig. S3)[22, 23]. The pie-charts in Fig. 1 and supplemental tables show the

breakdown of isolate number -v- lineages (Table S2) and sub-lineages (Table S3) for

each location. The phylogenetic tree in Fig. 2 further highlights the strong population

structure of this collection of isolates, with isolates clustering according to lineage.

Isolates of the ancient Indo-oceanic lineage/L1 ($n = 1,150$) were mostly

contributed by India ($n = 676$ isolates) and Vietnam ($n = 283$ isolates). 85% of the L1

Indian isolates belong to sub-lineages 1.1.2 and 1.2.2 while 66% of the Vietnamese

isolates are sub-lineage 1.1.1.1. No L1 isolates were contributed by 10 of the 23

countries in this study with only 2 isolates collected in South America (one in each of

Brazil and Peru).

There are 5,598 L2 (East Asian) isolates, making it the second largest group in

the dataset. L2 was found most prevalent in Asia and Europe with the largest

proportion found in amongst isolates contributed by China ($n = 722$, 64% of isolates)

and India ($n = 1,573$, 39% of isolates). Sub-lineages 2.2 and 2.2.7 dominate the L2

isolates ($n = 1,421$ and 1,249 respectively); 2.2 was found mostly amongst Peruvian

and South African isolates ($n = 231$ and 161 respectively) apart from those contributed

by the Asian countries of Vietnam ($n = 271$), China ($n = 284$) and India ($n = 272$), while

85% of sub-lineage 2.2.7 isolates were contributed by South Africa ($n = 206$), Vietnam

($n = 164$) and India ($n = 691$).  70% of sub-lineage 2.2.1 originated from South Africa

(10% of isolates found here) and has recently been associated with more favourable

transmission rates [24]. Lastly, 86% and 72% of isolates contributed by Kyrgyzstan

and Turkmenistan respectively belong to L2 with sub-lineage 2.2.10 dominating (16/24

and 75/86 isolates for both countries respectively). 2.2.10 has been previously

described as restricted to Central Asia and this is reflected in the compendium [25].

The majority (1184/1850, 65%) of L3 (East African/Indian) isolates were

contributed by India, followed by 19.6% (363/1850) isolates from Pakistan. L3 is

typically under-sampled and under-studied in current databases and biobanks in

comparison to L2 and L4; the L3 isolates in this study are the largest collected to date

in a single study [26].

L4 (Euro-American) is the largest lineage group in the compendium ($n$ = 6,572).

Isolates donated from Peru dominate; 87% of all L4 isolates are Peruvian with 4.1.2.1

and 4.3.3 being the most prevalent sub-lineages (24% and 22% respectively). There

are 34 different L4 sub-lineages in the dataset, making L4 the most diverse in

comparison to the other lineage groupings.

There were no L5 isolates found in the compendium and only 6 L6 isolates were

identified. Animal-restricted pathogenic mycobacterial isolates are also rare in the

compendium; only 16 cases were identified ($n$ = 15 *M. bovis* and $n$ = 1 *M. caprae*).

**Figure 1:** *Geographical distribution of CRyPTIC M. tuberculosis clinical isolates.*
The total number of isolates contributed by each country is depicted, with pie charts representing
the proportion of *M. tuberculosis* lineages. Where the origin of an isolate was not known, the
collection site identity was assigned (269 isolates in Germany, 17 isolates in India, 6 isolates in
Peru, 885 isolates in Italy, 510 isolates in South Africa, 357 isolates in Sweden, 208 isolates in
Taiwan, 1 isolate in Brazil and 4 isolates in the UK).

350
**Figure 2: *Phylogenetic tree of CRyPTIC M. tuberculosis clinical isolates.*** A
phylogenetic cladogram of 15,211 *M. tuberculosis* clinical isolates. A neighbour-joining tree was
constructed from a pairwise distance matrix using *quicktree* [20]. Coloured dots at the branch
termini represent the lineage assigned to each isolate. "Animal/Other" includes 16 isolates that
were assigned the following lineages: *M. caprae, M. bovis, unknown* along with 17 quality control
isolates representing specific sub-lineages [27].

### *Quality assurance of the drug minimum inhibitory concentrations*

Regular quality assurance checks detected problems with plate inoculation and

reading in two laboratories during collection of the 15,211 isolates. In addition, some

plates were not able to be read, usually because of inadequate growth in the positive

control wells or extensive contamination. These anomalies were investigated, and

samples were removed from the dataset where appropriate, resulting in 2,922 isolates

being held out of the final dataset. Further, due to skip wells and other phenomena

that prevent an MIC being measured, 11.9% of isolates did not return a phenotype for

all 13 drugs on the plate.  The compendium therefore has a total of 157,401 MIC

measurements for 12,289 isolates (Table 1).

|  | MIC MEASUREMENTS | HIGH QUALITY | MEDIUM QUALITY | LOW QUALITY |
|---|---|---|---|---|
| **INH** | 12070 | 9519 | 1351 | 1200 |
| **RIF** | 12099 | 8955 | 1356 | 1788 |
| **EMB** | 12158 | 7506 | 1355 | 3297 |
| **LEV** | 12163 | 7774 | 1354 | 3035 |
| **MXF** | 12194 | 6785 | 1353 | 4056 |
| **AMI** | 12072 | 8973 | 1350 | 1749 |
| **KAN** | 12130 | 9333 | 1355 | 1442 |
| **BDQ** | 12068 | 8536 | 1355 | 2177 |
| **CFZ** | 12049 | 7763 | 1352 | 2934 |
| **DLM** | 11927 | 8095 | 1349 | 2483 |
| **LZD** | 12189 | 7141 | 1355 | 3693 |
| **ETH** | 12132 | 8821 | 1355 | 1956 |
| **RFB** | 12150 | 10042 | 1352 | 756 |
| **TOTAL** | **157401** | **109243** | **17592** | **30566** |

**Table 1: Q***uality metrics for phenotype data* Stated for each drug is the total number of
MIC readings ("PHENOTYPED") stratified into "high" quality (at least two MIC measurement
methods agree), "medium" quality (either Vizion and AMyGDA disagree, or there is no plate
picture) or "low" quality (all three MIC measurements methods disagree) phenotype classifications
as described in Methods. Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol,
LEV = levofloxacin, MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline,
CFZ = clofazimine, DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.

***Resistance classification and distribution***

Binary drug resistant/susceptible phenotypes were calculated from the MICs by

applying an epidemiological cut-off value (ECOFF); samples with MICs at or below the

ECOFF are, by definition, wild-type and hence assumed to be susceptible to the drug

in question [11] (Fig. S1). Samples with MICs above the ECOFF are therefore

assumed to be resistant.

Unsurprisingly given its size, resistance to each of the 13 drugs is represented

within the CRyPTIC compendium (Fig. 3a). The drugs with the highest percentage of

resistance are the first line drugs isoniazid and rifampicin (49.0% and 38.7%

respectively). Of the second line drugs, levofloxacin had the highest proportion of

resistant isolates in the dataset (17.6%) and amikacin the lowest (7.3%). Reassuringly,

a low proportion of isolates were resistant to the NRDs, bedaquiline (0.9%),

clofazimine (4.4%), delamanid (1.6%) and linezolid (1.3%).

Of the 12,289 isolates which returned a binary phenotype for at least one drug,

6,814 (55.4%) were 'resistant' to at least one drug (Fig. 3b).  For the purpose of

describing the broader resistance categories present in the dataset, we assumed that

all MICs that could not be read had susceptible phenotypes in this section.

Consequently, the calculated prevalence of MDR, XDR etc in the dataset (Fig. 3) are

likely under-estimates.

At least 68.8% ($n$ = 4,685) of drug resistant isolates were either MDR or RR

(Fig. 3b) and of these 38.8% ($n$ = 1,819) were pre-XDR (MDR/RR + fluoroquinolone

resistance) and 3.0% were XDR ($n$ = 142). Two of the XDR isolates returned a

resistant phenotype to all 13 of the drugs assayed (Table S1) and therefore could be

reasonably described as totally drug resistant (TDR). One such isolate belonged to L4

and was contributed by South Africa, and the other belonged to L2 with an unknown

country of origin.

Of the 2,129 resistant isolates that did not fit into either of the MDR or RR

categories, more were categorised as isoniazid resistant (73.1%, $n$ = 1,556) than were

categorised as susceptible to both isoniazid and rifampicin and resistant to at least

one other drug (26.9%, $n$ = 573).

Whilst many partner laboratories oversampled for resistance, others used a

less biased collection protocol (Fig. 3c). Therefore, the proportions of resistant

phenotypes are not necessarily representative of the overall distribution of resistance

in the country and are not comparable between countries. Isolates with unknown

country of origin were excluded from this analysis. In all countries that contributed

more than 100 resistant isolates, each of the five phenotypic resistance categories in

Fig. 3b were identified, except for Peru, Vietnam and Nepal which did not contribute

any XDR isolates. Vietnam and Brazil sampled a particularly high proportion of non-

MDR/RR resistant phenotypes; 73.9% and 55.1% of resistant isolates contributed by

these countries, respectively, were neither MDR nor RR.

For Nepal and India, an especially high proportion of the MDR/RR isolates

contributed were fluoroquinolone resistant (92.9% ($n$ = 132/142 isolates) and 69.8%

($n$ = 538/770 isolates) respectively), as has been previously observed for this

geographical region [28, 29].

All five categories of resistance were represented in the four major *M.*

*tuberculosis* lineages. We note that these relative proportions will have been

influenced by the different sampling approaches at each partner laboratory since

lineage distributions are typically geographically distinct (Fig. 1). Bearing this in mind,

we observe that in the compendium, L3 isolates contained the most MDR/RR isolates

as a proportion of resistant isolates (77.6%), followed by L2 (74.3%), L4 (63.4%), and

L1 (50.0%). L2 is known to be associated with MDR/RR and XDR and although it did

not contain the highest proportion of MDR/RR isolates, it did have the most pre-XDR

isolates as a proportion of MDR/RR isolates (54.2%) [30, 31]. L1 and L2 contained the

most XDR isolates as a proportion of MDR/RR isolates (4.7% and 4.2% of isolates
respectively).

**Figure 3: *Drug phenotype data for the CRyPTIC compendium* a)** Prevalence of
resistance to each of 13 drugs in the CRyPTIC dataset. The total number of isolates with a binary
phenotype (of any quality) for the corresponding drug is presented in Table 1. **b)** Phenotypes of
12, 289 CRyPTIC isolates with a binary phenotype for at least one drug. **c)** Geographical
distribution of phenotypes of 12, 289 CRyPTIC isolates. Intensity of blue shows the percentage of
isolates contributed that were categorised as susceptible to all 13 drugs. Donut plots show the

proportions of resistant phenotypes identified in (b) for countries contributing >=100 isolates with drug resistance. **d)** Proportions of resistance phenotypes in the 4 major *M. tuberculosis* lineages. N is the number of isolates of the lineage called resistant to at least one of the 13 drugs. Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, LEV = levofloxacin, MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline, CFZ = clofazimine, DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.

### Co-occurrence of drug resistance amongst the CRyPTIC isolates

Isolates with all possible two-drug resistant combinations were present in this dataset (Fig. 4a). Resistance to any of the remaining 11 drugs was associated with resistance to both isoniazid and rifampicin. Isoniazid resistance was the most strongly associated bar the following two exceptions, both of which arise because drugs from the same class are being compared: a greater proportion of isolates resistant to rifabutin were also resistant to rifampicin (96.8% had rifabutin - rifampicin co-resistance versus 93.3% rifabutin - isoniazid co-resistance) and a greater proportion of isolates resistant to moxifloxacin were associated with levofloxacin resistance (97.6% had moxifloxacin – levofloxacin co-resistance versus 95.0% moxifloxacin – isoniazid co-resistance). Of the second line drugs, levofloxacin and moxifloxacin were more commonly seen as a second resistant phenotype than the injectable drugs kanamycin and amikacin.

Despite this, resistance to both drugs in the aminoglycoside class was common the dataset; 90.4% of amikacin resistant isolates were also resistant to kanamycin although significantly fewer kanamycin resistant isolates were resistant to amikacin (72.0%, *p* <0.00001). In a similar fashion, a smaller proportion of rifampicin resistant isolates were resistant to rifabutin than rifabutin resistant isolates that were resistant to rifampicin (91.3%, 96.8%, *p* <0.00001) while a smaller proportion of levofloxacin

resistant isolates were resistant to moxifloxacin than moxifloxacin resistant isolates

that were resistant to levofloxacin (78.5%, 97.6%, $p$ <0.00001). One possible

explanation is that the cut off, in this case the ECOFFs, have been inconsistently

defined for the different compounds, however here all ECOFFs have been determined

in the same way using a much larger dataset than is typically the case [11]. Differences

in drugs of the same class are also well documented by *in vitro* studies [32-34].

Bedaquiline, clofazimine, delamanid and linezolid were les commonly seen as

a second resistance phenotype and the smallest proportional resistance combinations

involved the NRDs (*e.g.* 1.5% of isoniazid resistant isolates were bedaquiline resistant

and 1.7% were delamanid resistant). Within the NRDs however, co-occurrence of

resistance was proportionally higher; bedaquiline resistance was most seen with

clofazimine resistance (52.4%), clofazimine resistance was most seen with

bedaquiline resistance (10.6%), delamanid resistance was most seen with clofazimine

resistance (26.3%) and linezolid resistance was most seen with clofazimine resistance

(34.2%).

***Additional antibiotic resistance in isolates with MDR or non-MDR phenotypic***

***backgrounds***

We next examined in more detail the correlation structure of phenotypes by

conditioning on different phenotypic "backgrounds". (Fig. 4b-f). We found that a

greater proportion of isolates that were susceptible to isoniazid and rifampicin were

resistant to second-line drugs than the first line drug ethambutol (15.9%) (Fig. 4b). The

proportion of isolates resistant to clofazimine or levofloxacin was particularly high

(32.9% and 24.1%, respectively), and more isolates were resistant to these two drugs

than ethambutol in an isoniazid resistant and rifampicin susceptible background but

not in MDR/RR isolates (Fig. 4c-f).

MDR/RR isolates (asides from rifabutin) were most commonly resistant to the

first line drug ethambutol (46.3%), closely followed by levofloxacin (41.4%), then

moxifloxacin (34.1%). A greater proportion of isolates resistant to levofloxacin than

moxifloxacin was also seen in all other backgrounds and as expected, the proportion

of fluoroquinolone resistance was higher in MDR/RR isolates than non-MDR isolates

[35]. For the aminoglycosides, a greater percentage of MDR/RR isolates were

kanamycin resistant (21.8%) than amikacin resistant (18.1%), a trend seen in all other

backgrounds.

For isolates with an XDR phenotype, a higher proportion were also resistant to

linezolid than bedaquiline (66.7% compared to 44.6%) and 11.3% of XDR isolates

were resistant to both bedaquiline and linezolid (Fig. 4f). XDR isolates were also

resistant to the other NRDs, clofazimine (41.3%) and delamanid (18.8%). For isolates

with an MDR/RR or pre-XDR background, the most common NRD resistance seen

was clofazimine (6.0% and 9.3%, respectively) followed by linezolid (2.3% and 4.7%

respectively), bedaquiline (1.8% and 3.2%, respectively), and delamanid (1.7% and

2.3%, respectively).  In non-MDR/RR backgrounds: clofazimine resistance remained

the most common followed by delamanid resistance, linezolid resistance and

bedaquiline resistance (Fig. 4b-c).

**Figure 4:** *Co-occurrence of antibiotic resistance in CRyPTIC M. tuberculosis isolates* **a)** The heatmap shows the probability of an isolate being resistant to Drug 2 if it is resistant to Drug 1, percentages are given in Table S4. **(b-g)** Percentage of isolates that are resistant to another of the 13 drugs in a background of **(b)** isoniazid susceptible + rifampicin susceptible, **(c)** isoniazid resistant + rifampicin susceptible, **(d)** MDR/RR **(e)** Pre-XDR, **(f)** XDR. Only samples with definite phenotypes for RIF in MDR backgrounds and RIF and INH in non-MDR backgrounds and the additional drug are included. Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, LEV = levofloxacin, MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline, CFZ = clofazimine, DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.

### Genetic determinants of resistance

In order to establish a baseline measure of how well resistance and susceptibility could be predicted based on the state of the art prior to the CRyPTIC project, we constructed a hybrid catalogue of genetic variants associated with resistance to the first-line compounds isoniazid, rifampicin & ethambutol and the

second-line compounds levofloxacin, moxifloxacin, amikacin, kanamycin, ethionamide

based on existing catalogues [7, 36]. (We note in passing that we specifically did not

use the recent WHO catalogue in order to avoid circularity and over-training, as that

catalogue was developed (via prior literature, expert rules and a heuristic algorithm)

based partially on these isolates [37]. We applied our hybrid catalogue to predict the

susceptibility and resistance of all isolates in this compendium to these compounds,

and the resulting genetic-based predictions were then compared with the binary

phenotypes derived from the MICs (Table 2). Since these data were not collected

prospectively or randomly, but indeed are enriched for resistance, the calculated error

rates are not representative of how well such a method would perform in routine clinical

use. We also did not apply the approach used in [7] of refusing to make a prediction if

a novel mutation was detected in a known resistance gene, as we simply wanted to

measure how well a pre-CRyPTIC catalogue could predict resistance in the

compendium. The results were broadly in line with prior measurements on a smaller

(independent) set [18]. The hybrid catalogue does not make predictions for rifabutin,

linezolid, bedaquiline, delamanid or clofazimine; indeed, this is one of the main aims

of the consortium and new catalogues published by CRyPTIC and the WHO will begin

to address this shortcoming (Fig. 5) [19].

563
564
Table 3 shows the top mutations found amongst isolates phenotypically

resistant to first- and second- line drugs. As expected, *rpoB* S450L is the most

prevalent mutation associated with rifampicin resistance, with 62.2% of all

phenotypically rifampicin-resistant isolates carrying it, while 70% of isoniazid resistant

isolates have *katG* S315T mutations. Mutations in *gyrA* dominate amongst

fluoroquinolone resistant isolates; D94G and A90V are the two most frequently

occurring mutations for levofloxacin and moxifloxacin, with just under 77% and just

over 94% of phenotypically resistant isolates carrying at least one mutation in *gyrA*

respectively. Mutations in *rpoC* are amongst those most prevalent for rifabutin

resistant isolates; these mutations are typically described as compensatory, while

G594E was subsequently defined as a lineage-specific marker for L4 LAM and

Haarlem lineages [38].

| | TP | FP | TN | FN | VME | ME | PPV | NPV |
|---|---|---|---|---|---|---|---|---|
| **INH** | 5493 | 142 | 5622 | 224 | 0.039 | 0.025 | 0.961 | 0.975 |
| **RIF** | 4535 | 435 | 6669 | 107 | 0.023 | 0.061 | 0.977 | 0.939 |
| **EMB** | 1919 | 513 | 6702 | 111 | 0.055 | 0.071 | 0.945 | 0.929 |
| **LEV** | 1689 | 255 | 8104 | 184 | 0.098 | 0.031 | 0.902 | 0.969 |
| **MXF** | 1358 | 504 | 9022 | 160 | 0.105 | 0.053 | 0.895 | 0.947 |
| **AMI** | 632 | 84 | 10117 | 163 | 0.205 | 0.008 | 0.795 | 0.992 |
| **KAN** | 735 | 124 | 9043 | 197 | 0.211 | 0.014 | 0.789 | 0.986 |
| **ETH** | 971 | 114 | 9183 | 511 | 0.345 | 0.012 | 0.655 | 0.988 |

**Table 2: *Predicting phenotypic resistance using genetics***
TP: the number of phenotypically resistant samples are that correctly identified as resistant ("true positives"); FP, the number of phenotypically susceptible samples that are falsely identified as resistant ("false positives"); TN, the number of phenotypically susceptible samples that are correctly identified as susceptible ("true negatives"); FN, the number of phenotypically resistant samples that are incorrectly identified as susceptible ("false negative"); VME, very major error rate (false-negative rate), 0-1; ME, major error rate (false-positive rate), 0-1; PPV, positive predictive value, 0-1; NPV, negative predictive value, 0-1.

| | GENE | VARIANT | *n* | % |
|---|---|---|---|---|
| RIF | *rpoB* | S450L | 2914 | 62.2 |
| | *rpoB* | D435V | 506 | 10.8 |
| | *rpoB* | H445D | 202 | 4.3 |
| | *rpoB* | H445Y | 147 | 3.1 |
| | *rpoB* | D435Y | 112 | 2.4 |
| | | | | |
| INH | *katG* | S315T | 3668 | 70. |
| | *fabG1* | c-15t | 829 | 15.8 |
| | *fabG1* | g-17t | 176 | 3.4 |
| | *fabG1* | t-8c | 154 | 2.9 |
| | *inhA* | I194T | 56 | 1.1 |
| | | | | |
| EMB | *embB* | M306V | 1131 | 50 |
| | *embB* | M306I | 1001 | 44.3 |
| | *embB* | Q497R | 449 | 19.9 |
| | *embB* | G406A | 164 | 7.2 |
| | *embB* | G406D | 105 | 4.6 |
| | | | | |
| KAN | *rrs* | a1401g | 660 | 58.9 |
| | *eis* | c-14t | 70 | 6.2 |
| | *eis* | g-10a | 53 | 4.7 |
| | | | | |
| AMI | *rrs* | a1401g | 660 | 74.7 |
| | *rrs* | g1484t | 7 | 0.8 |
| | | | | |
| LEV | *gyrA* | D94G | 783 | 36.5 |
| | *gyrA* | A90V | 487 | 22.7 |
| | *gyrA* | D94N | 157 | 7.3 |
| | *gyrA* | D94A | 133 | 6.2 |
| | *gyrA* | S91P | 92 | 4.3 |
| | | | | |
| MXF | *gyrA* | A90V | 487 | 28.2 |
| | *gyrA* | D94G | 783 | 45.4 |
| | *gyrA* | D94N | 157 | 9.1 |
| | *gyrA* | D94A | 133 | 7.7 |
| | *gyrA* | D94Y | 70 | 4.1 |
| | | | | |
| ETH | *fabG1* | c-15t | 829 | 48 |
| | *fabG1* | L203L | 124 | 7.2 |

**Table 3:** *The top mutations associated with phenotypic drug resistance*
A genetic resistance catalogue was used for genotypic prediction of resistance for each drug in
column 1 [7, 36]. "VARIANT": non-synonymous amino acid mutations are denoted by upper case
letters while nucleotide substitutions for non-coding sequences are denoted by lower case letters.
Negative numbers denote substitutions in promoter regions; "GENE": genic region of interest in
which "Variant" can be found; "*n*": number of phenotypically resistant isolates with "VARIANT"; "%":
percentage of total phenotypically resistant isolates with "VARIANT". Drug acronyms: INH =
isoniazid, RIF = rifampicin, EMB = ethambutol, AMI = amikacin, KAN = kanamycin, LEV =
levofloxacin, MXF = moxifloxacin, ETH = ethionamide.

607
608

### *Resistance to new and re-purposed drugs*

As previously stated, relatively few isolates are resistant to the NRDs, bedaquiline ($n$ = 109), clofazimine ($n$ = 525), delamanid ($n$ = 186) and linezolid ($n$ = 156). South Africa contributed the greatest number of isolates resistant to bedaquiline, clofazimine and linezolid (Fig. 5a), while China and India contributed the most isolates resistant to delamanid. Since the collection protocol differed between laboratories it is not possible to infer any differences in the relative prevalence of resistance to the NRDs in these countries. The results of a survey of all non-synonymous mutations in genes known or suspected to be involved in resistance to these four drugs (*e.g. rv0678, mmpL5, pepQ, ddn, rplC, rrl etc.*) are depicted in Fig. 5b-e [39-43]. Mutations known to be associated with sensitivity were ignored, along with mutations that occurred at a frequency of ≥ 5% amongst all isolates. In contrast to first- and second-line drugs, there are no mutations within a single gene/small group of genes that can fully explain resistance to any NRD. Note that the role of most of these mutations in resistance remains undetermined. Work is ongoing within the consortium to unravel the association of these (and novel) mutations with resistance to the NRDs along with the relevant mechanisms of resistance.

628
**Figure 5:** *Resistance to bedaquiline, clofazimine, delamanid and linezolid*
*amongst M. tuberculosis CRyPTIC isolates* **a)** The prevalence of resistance to
bedaquiline (BDQ), clofazimine (CFZ), delamanid (DLM) and linezolid (LZD) per country or origin
or collection site. Phylotrees are shown for isolates phenotypically resistant to b) BDQ, c)
DLM and e) LZD. Tip point colours denote lineage. Each outer track represents a gene thought to

634 be associated with resistance and coloured blocks denote the presence of a non-synonymous
635 mutation in the relevant gene for a given isolate. Mutations in these genes that are either
636 associated with sensitivity or present in >5% of the collection of isolates as a whole were ignored.
637
638

### *Case study on rifampicin mono-resistance*

Around 1% of TB cases are rifampicin mono-resistant (RMR) and the frequency

is increasing [1, 44]. The WHO does not recommend isoniazid for RMR treatment,

despite it being effective; this is likely due to the reliance on the Xpert® MTB/RIF assay

which cannot distinguish between RMR and MDR. Use of isoniazid could improve

treatment outcomes for RMR patients which are currently similar to that of MDR TB,

including a higher risk of death compared to drug susceptible infections [45, 46]. Due

to its low natural prevalence, RMR has been poorly studied to date but increasingly

large clinical TB datasets, such as the one presented here, make its study now

feasible.

For this case study, we defined RMR as any isolate that was rifampicin resistant

and isoniazid susceptible, discounting isolates with no definite phenotype for either

isoniazid and/or rifampicin. Of the 4,655 rifampicin resistant isolates in the CRyPTIC

database that also had a phenotype for isoniazid, 302 (6.5%) were RMR. These

isolates were contributed by South Africa, Peru, India, Italy, China, Pakistan, Brazil,

Vietnam, Germany, Nigeria, Turkmenistan and Tajikistan (Fig. 6a). South Africa and

Nigeria contributed a significantly higher number of RMR isolates, as a proportion of

rifampicin resistant isolates, than that of the total dataset at 17.5% ($p$ <0.00001) and

27.3% ($p$ = 0.00534) respectively. We note that countries may have oversampled for

solo resistance, but the higher contribution of RMR isolates from South Africa is

consistent with previous studies [44].

RMR has evolved in all four major lineages, with L4 having a significantly higher

RMR prevalence than that of the total rifampicin resistant dataset at 9.4% ($p$ =

0.00031) and L2 significantly lower RMR prevalence at 4.2% ($p$ = 0.00026). Although

these numbers will have been somewhat influenced by differences in sample

collection bias at different geographical sites, we would expect L2 to have a lower

proportion of RMR isolates due to its association with MDR phenotypes [31, 47] (Fig.

6b).

***Rifampicin mono-resistance is incorrectly predicted by current diagnostics***

A widely used, WHO-endorsed diagnostic tool, the Xpert® MTB/RIF assay,

uses a proxy whereby *any* SNP detected in the "rifampicin-resistance determining

region" (RRDR) of *rpoB* results in a prediction of MDR. However, the suitability of the

proxy is dependent upon prevalence of RMR in the population [44]. We tested the

reliability of this on the 4,655 rifampicin resistant isolates in our dataset that had a

phenotype for isoniazid (Fig. 6c).

Of these isolates, 4,353 (93.5%) were MDR and 302 (6.5%) were RMR. 187 of

the MDR isolates had no RRDR mutation and therefore 4.0% of isolates in this study

would be predicted as false negative MDR by the Xpert® MTB/RIF assay. 276 of the

RMR isolates had a mutation in the RRDR of *rpoB* and so the Xpert® MTB/RIF assay

proxy would incorrectly predict 5.9% of the rifampicin resistant isolates as false

positive MDR cases. However, overall, the Xpert® MTB/RIF assay proxy correctly

predicts 89.5% of the rifampicin resistant isolates as MDR and 0.6% of the isolates as

non-MDR in this dataset, which suggests it is a reasonably successful diagnostic tool

with >90% accuracy for MDR classification of rifampicin resistant isolates. As most

sites oversampled for resistance, our dataset likely contains a higher prevalence of

RMR than the global average and hence the Xpert® MTB/RIF assay is likely to perform

better on more representative data. However, the analysis shows how the increasing

global levels of RMR TB cases could increase the number of false positive MDR

diagnoses by the Xpert® MTB/RIF assay, denying isoniazid treatment to a greater

number of patients who would then be moved on to less effective drugs.

***There are genetic differences between rifampicin mono-resistant and***

***multidrug resistant isolates***

We have analysed our matched phenotypic and genotypic data to examine

whether there were any differences in the genetic determinants of rifampicin

resistance between RMR and MDR isolates. The proportion of RMR isolates with no

*rpoB* mutation (5.3%, Fig. 6c) was significantly higher than that of MDR isolates (1.8%,

*p* <0.00001). This suggests that non-target-mediated resistance mechanisms, such as

upregulation of rifampicin specific efflux pumps, could be more influential in providing

protection against rifampicin in RMR isolates than in MDR isolates.

The majority of RMR and MDR isolates contained one or more SNPs in *rpoB*,

with the majority having at least one mutation in the RRDR. To date, several non-

synonymous RRDR mutations have been found in RMR *M. tuberculosis* isolates,

including the resistance conferring mutations S450L, H445D and D435Y, which are

also seen in MDR isolates [48, 49]. For both RMR and MDR isolates in this dataset,

the most common *rpoB* RRDR mutation seen was S450L (63.6% and 41.1% of

isolates respectively, Fig. 6d). Five mutations were present in RMR isolates that were

not seen in MDR isolates: S428G, S441A, S441V, S450M and S450Q, however these

were seen at low prevalence (less than 2%) of RMR isolates. We found more RMR

isolates had His445 mutated than MDR isolates (27.8% of RMR and 9.5% of MDR, *p*

<0.00001), and mutations to amino acids Ser450 and Asp435 were more prevalent in

MDR isolates than RMR isolates (43.7% of RMR and 65.8% of MDR (*p* <0.00001),

and 9.3% of RMR and 15.5% of MDR (*p* = 0.00328) respectively).

Relatively little difference in the mutational profiles for *rpoB* mutations falling

outside the RRDR was found; many RMR mutations were also seen in MDR isolates

and those found exclusively in RMR were at <2% prevalence (Fig. S2). The most

common non-RRDR mutation in both MDR and RMR isolates was a cytosine to

thymine mutation 61 bases upstream of the *rpoB* start codon (10.1% and 8.6% of

isolates respectively). The resistance conferring mutations, V695L and V170F, were

seen at low proportions with no significant difference between MDR and RMR isolates

(V695L was seen in 0.5% MDR isolates and 1.3% of RMR isolates, and V170F was

seen in 0.8% of MDR isolates and 1.7% of RMR isolates).

**Figure 6:** *Rifampicin mono resistance* **a)** Percentage of rifampicin resistant isolates that are rifampicin mono-resistant (RMR) by country of isolate origin. * indicates RMR proportions that were significantly different from that of the total dataset using a two tailed z-test with 95% confidence. **b)** RMR prevalence by lineage. * indicates RMR proportions that were significantly different from that of the total dataset using a two tailed z-test with 95% confidence. **c)** MDR predictions for rifampicin resistant isolates made using the Xpert® MTB/RIF assay proxy. N is the total number of rifampicin resistant isolates. The inner ring shows the proportion of rifampicin resistant isolates that are MDR and RMR. The middle ring represents the proportions of MDR and RMR isolates that have a SNP (synonymous or non-synonymous) in the RRDR of *rpoB* (RRDR), no RRDR SNP but a SNP elsewhere in the *rpoB* gene (rpoB) and no *rpoB* mutations (none). The outer ring shows the expected true positive (TP), true negative (TN), false positive (FP) and false negative (FN) MDR predictions of Xpert® MTB/RIF assay, based on the SNPs present in the rifampicin resistant isolates. **d)** Non-synonymous mutations found in the RRDR of *rpoB* in RMR isolates and MDR isolates. Presence of a coloured spot indicates that the mutation was found in RMR/MDR isolates and spot size corresponds to the proportion of RMR or MDR isolates carrying that mutation.

## Discussion

This compendium of *M. tuberculosis* clinical isolates is the result of an extensive global effort by the CRyPTIC consortium to better map the genetic variation associated with drug resistance. Through its sheer size and by oversampling for resistance, the compendium gives an unparalleled view of resistance patterns in the panel of 13 antitubercular compounds studied, providing more nuance to the traditional classification (nonMDR/MDR/XDR) nomenclature. These insights could be used to inform future treatment guidelines and steer the development of improved diagnostics.

For first-line drugs, molecular based diagnostic assays have vastly improved the detection of and the speed at which we find drug resistant TB cases, resulting in improved quality of care for patients. However, relying solely on these diagnostic methods has several drawbacks. Aside from the Xpert® MTB/RIF assay potentially increasing false positive MDR diagnoses as discussed earlier in the RMR case study, the assay assumes isoniazid resistance upon detection of rifampicin resistance. Thus, less is known about the prevalence of mono isoniazid resistance or 'true' cases of MDR (confirmed rifampicin *and* isoniazid resistance) (REFWHO). A greater amount of isoniazid-resistant and rifampicin-susceptible (Hr-TB) isolates were contributed by CRyPTIC countries than RMR isolates *(n = 1470 versus n = 302),* a pattern also recently observed in a global prevalence study [50]. A modified 6-month treatment regimen is now recommended for Hr-TB (rifampicin, ethambutol, levofloxacin and pyrazinamide), and as a result of inadequate diagnosis many of the 1.4 million global Hr-TB estimated cases would have received inadequate and unnecessarily longer treatment regimens [1, 50]. Encouragingly, CRyPTIC isolates with a Hr-TB background exhibited relatively low levels of resistance to other antitubercular drugs,

including those in the augmented regimen (Fig. 4c). However, without the tools

appropriate to assess and survey this, we will continue to misdiagnose and infectively

treat these clinical cases.

In 2018, CRyPTIC and the 100,000 Genomes project demonstrated using WGS

that genotypic prediction correlates well with culture-based phenotype for first-line

drugs, which is reflected in our summary of the genetic catalogue applied to this

dataset (Table 3) [7]. While predictions can be made to a high level of sensitivity and

specificity, there is still more to learn, as exemplified by the isolates in the compendium

that despite being resistant to rifampicin and isoniazid could not be described

genetically (Table 2). This shortfall, along with the limitations of molecular based

diagnostic assays, highlights the need for continual genetic surveillance as drug

resistant cases rise and shines a favourable light on a WGS-led approach.

One of the particular strengths of this compendium lies with the data collated

for second-line drugs. A greater proportion of drug resistant isolates had additional

resistance to fluoroquinolones than second line injectable drugs (Fig. 4a). This could

be because of more widespread use of fluoroquinolones to treat bacterial infections in

general as well as their ease of administration, hence being recommended over

injectables for longer MDR treatment regimens [1]. Concerningly, we found that

resistance to levofloxacin and moxifloxacin, and kanamycin and amikacin, were more

common than resistance to the mycobacterial specific drug ethambutol in an isoniazid

and rifampicin susceptible background (Fig. 4b) which suggests a level of pre-existing

second line resistance. This concurs with a systematic review that found patients

previously prescribed fluoroquinolones were 3 times more likely to have

fluoroquinolone resistant TB [51]. Careful stewardship of fluoroquinolones both in TB

and other infectious diseases will be paramount for the success of treatment regimens.

Despite the variability in sample collection, the observed high proportions of

fluoroquinolone resistant MDR/RR isolates from some countries also suggests that

MDR treatment regimens could be improved by optimisation on a geographic basis.

Further improvement could also be made by the selection of appropriate drugs

from each class; for example, the WHO recommends switching from kanamycin to

amikacin when treating MDR TB patients [52]. The compendium supports such

recommendations as we saw more resistance to kanamycin than amikacin in not just

MDR/RR but also all other phenotypic backgrounds. For fluoroquinolones, more

isolates were resistant to levofloxacin than moxifloxacin in all phenotypic backgrounds

suggesting moxifloxacin may by the most appropriate fluoroquinolone to recommend.

However, the amenability of drugs to catalogue-based genetic diagnostics is also an

important consideration, and our data suggest levofloxacin resistance could be

predicted more reliably than moxifloxacin with fewer false positives predicted (Table

2).

Testing for fluoroquinolone resistance using molecular diagnostic tests remains

limited. However, the limited global data from the past 15 years suggests that the

proportion of MDR/RR TB cases resistant to fluoroquinolones sits at around 20%, with

these cases primarily found in regions of high MDR-TB burden [1]. While tools such

as the Cepheid Xpert® MTB/XDR cartridge, which has just been approved for use by

the WHO, will permit both isoniazid and fluoroquinolone testing to be increased, the

same pitfalls are to be encountered regarding targeted diagnostic assays [53]. In

contrast, the genetic survey in this study demonstrates the potential of WGS for

genetic prediction of resistance to second-line drugs and studies within the consortium

to investigate this are underway.

The CRyPTIC compendium has facilitated the first global survey of resistance

to NRDs. Reassuringly, prevalence of resistance to the NRDs was lower than for first-

and second-line agents in the dataset as a whole (Fig. 3a), and resistance to the new

drugs bedaquiline and delamanid was less common than the repurposed drugs

clofazimine and linezolid in an MDR/RR background (Fig. 4c). However, the presence

of higher levels of delamanid and clofazimine resistance than ethambutol resistance

in the isoniazid and rifampicin susceptible background does suggest some pre-existing

propensity towards NRD resistance (Fig. 4b).

Co-resistance between all two-drug combinations of NRDs was seen in isolates

in the compendium, the most common being isolates resistant to both bedaquiline and

clofazimine. This link is well documented and has been attributed to shared resistance

mechanisms such as non-synonymous mutations in *rv0678* which were found in both

clofazimine and bedaquiline resistant isolates in the compendium [42] (Fig. 5b,c).

Increased clofazimine use could further increase the prevalence of *M. tuberculosis*

isolates with clofazimine and bedaquiline co-resistance, limiting MDR treatment

options such as using bedaquiline as the backbone of a shorter MDR regimen [54].

Therefore, proposed usage of clofazimine for other infectious diseases, including

COVID-19, should be carefully considered [55].

The WHO recommends not using bedaquiline and delamanid in combination to

prevent the development of co-resistance, which could occur relatively quickly [56];

the rates of spontaneous evolution of delamanid resistance *in vitro* has been shown

to be comparable to that of isoniazid, and likewise bedaquiline resistance arises at a

comparable rate to rifampicin resistance [57]. In this compendium, 12.9% bedaquiline

resistant isolates were resistant to delamanid and 7.1% delamanid resistant isolates

were resistant to bedaquiline. Several scenarios could account for this, including the

presence of shared resistance mechanisms. For example, as bedaquiline targets

energy metabolism within the cell, changes to cope with energy/nutrient imbalances

upon the acquisition of resistance-associated ATPase pump mutations may result in

cross resistance to delamanid in a yet unknown or unexplored mechanism [58].

Additionally, off-target mutations in the *mmpl3* gene (involved in export), are

associated with resistance to delamanid, bedaquiline, clofazimine and linezolid, and

were seen in isolates resistant to each of these drugs, although we note this could be

artefactual (Fig. 5b-e). In any case, it is imperative that genetic determinants of

resistance are fully explored for our last resort NRDs, with special attention given to

those mechanisms that could be shared with other agents. In the meantime, careful

stewardship and phenotypic and genotypic surveillance of the NRDs should be

implemented, including linezolid and clofazimine which are now group A and B drugs

respectively for MDR treatment [1].

Several research avenues are being actively explored by the CRyPTIC

consortium that make use of this compendium, including: *i)* relating genetic mutations

to quantitative changes in the minimum inhibitory concentrations of different drugs

[58]; *ii)* genome-wide association studies [59]; *iii)* training machine learning models

that can predict resistance [60]; *iv*) exploration of the genetic determinants of

resistance to second line and NRDs [61]. Collectively these studies share the same

aim of facilitating the implementation of WGS-directed resistance prediction in the

clinic. Finally, we urge other researchers to explore this large dataset of *M.*

*tuberculosis* clinical isolates and hope it will lead to a wave of new and inciteful studies

that will positively serve the TB community for years to come.

**Author contributions**

D.W.C, T.E.A.P, S.H, A.L.G.C, A.W.S, T.M.W, P.W.F, D.M.C designed the CRyPTIC

study and all contributing laboratories collected samples and provided data. MIC data

and genetic information was retrieved and processed by P.W.F, S.H, A.L.G.C, Z.I, M.H

and J.K. A.B and K.M.M designed and performed all analyses for this manuscript. A.B

and K.M.M wrote the manuscript with feedback from CRyPTIC partners.

893
**Acknowledgements**

We thank Faisal Masood Khanzada and Alamdar Hussain Rizvi (NTRL, Islamabad,

Pakistan), Angela Starks and James Posey (Centers for Disease Control and

Prevention, Atlanta, USA), and Juan Carlos Toro and Solomon Ghebremichael (Public

Health Agency of Sweden, Solna, Sweden), Iñaki Comas and Álvaro Chiner-Oms

(Instituto de Biología Integrativa de Sistemas, Valencia, Spain; CIBER en

Epidemiología y Salud Pública, Valencia, Spain; Instituto de Biomedicina de Valencia,

IBV-CSIC, Valencia, Spain).

**Funding bodies**

This work was supported by Wellcome Trust/Newton Fund-MRC Collaborative Award

(200205/Z/15/Z); and Bill & Melinda Gates Foundation Trust (OPP1133541). Oxford

CRyPTIC consortium members are funded/supported by the National Institute for

Health Research (NIHR) Oxford Biomedical Research Centre (BRC), the views

expressed are those of the authors and not necessarily those of the NHS, the NIHR

or the Department of Health, and the National Institute for Health Research (NIHR)

Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial

Resistance, a partnership between Public Health England and the University of

Oxford, the views expressed are those of the authors and not necessarily those of the

NIHR, Public Health England or the Department of Health and Social Care. J.M. is

supported by the Wellcome Trust (203919/Z/16/Z). Z.Y. is supported by the National

Science and Technology Major Project, China Grant No. 2018ZX10103001. K.M.M. is

supported by EMBL's EIPOD3 programme funded by the European Union's Horizon

2020 research and innovation programme under Marie Skłodowska Curie Actions.

T.C.R. is funded in part by funding from Unitaid Grant No. 2019-32-FIND MDR. R.S.O.

is supported by FAPESP Grant No. 17/16082-7. L.F. received financial support from

FAPESP Grant No. 2012/51756-5. B.Z. is supported by the National Natural Science

Foundation of China (81991534) and the Beijing Municipal Science & Technology

Commission (Z201100005520041). N.T.T.T. is supported by the Wellcome Trust

International Intermediate Fellowship (206724/Z/17/Z). G.T. is funded by the

Wellcome Trust. R.W. is supported by the South African Medical Research Council.

J.C. is supported by the Rhodes Trust and Stanford Medical Scientist Training

Program (T32 GM007365). A.L. is supported by the National Institute for Health

Research (NIHR) Health Protection Research Unit in Respiratory Infections at Imperial

College London. S.G.L. is supported by the Fonds de Recherche en Santé du Québec.

C.N. is funded by Wellcome Trust Grant No. 203583/Z/16/Z. A.V.R. is supported by

Research Foundation Flanders (FWO) under Grant No. G0F8316N (FWO Odysseus).

G.M. was supported by the Wellcome Trust (098316, 214321/Z/18/Z, and

203135/Z/16/Z), and the South African Research Chairs Initiative of the Department

of Science and Technology and National Research Foundation (NRF) of South Africa

(Grant No. 64787). The funders had no role in the study design, data collection, data

analysis, data interpretation, or writing of this report. The opinions, findings and

conclusions expressed in this manuscript reflect those of the authors alone. L.G. was

supported by the Wellcome Trust (201470/Z/16/Z), the National Institute of Allergy and

Infectious Diseases of the National Institutes of Health under award number

1R01AI146338, the GOSH Charity (VC0921) and the GOSH/ICH Biomedical

Research Centre (www.nihr.ac.uk). A.B. is funded by the NDM Prize Studentship from

the Oxford Medical Research Council Doctoral Training Partnership and the Nuffield

Department of Clinical Medicine. D.J.W. is supported by a Sir Henry Dale Fellowship

jointly funded by the Wellcome Trust and the Royal Society (Grant No. 101237/Z/13/B)

and by the Robertson Foundation. A.S.W. is an NIHR Senior Investigator. T.M.W. is a

Wellcome Trust Clinical Career Development Fellow (214560/Z/18/Z). A.S.L. is

supported by the Rhodes Trust. R.J.W. receives funding from the Francis Crick

Institute which is supported by Wellcome Trust, (FC0010218), UKRI (FC0010218),

and CRUK (FC0010218). T.C. has received grant funding and salary support from US

NIH, CDC, USAID and Bill and Melinda Gates Foundation. The computational aspects

of this research were supported by the Wellcome Trust Core Award Grant Number

203141/Z/16/Z and the NIHR Oxford BRC. Parts of the work were funded by the

German Center of Infection Research (DZIF). The Scottish Mycobacteria Reference

Laboratory is funded through National Services Scotland. The Wadsworth Center

contributions were supported in part by Cooperative Agreement No. U60OE000103

funded by the Centers for Disease Control and Prevention through the Association of

Public Health Laboratories and NIH/NIAID grant AI-117312. Additional support for

sequencing and analysis was contributed by the Wadsworth Center Applied Genomic

Technologies Core Facility and the Wadsworth Center Bioinformatics Core. SYNLAB

Holding Germany GmbH for its direct and indirect support of research activities in the

Institute of Microbiology and Laboratory Medicine Gauting. N.R. thanks the

Programme National de Lutte contre la Tuberculose de Madagascar.

**Wellcome Trust Open Access**

This research was funded in part, by the Wellcome Trust/Newton Fund-MRC

Collaborative Award [200205/Z/15/Z]. For the purpose of Open Access, the author has

applied a CC BY public copyright licence to any Author Accepted Manuscript version

arising from this submission.

This research was funded, in part, by the Wellcome Trust [214321/Z/18/Z, and

203135/Z/16/Z]. For the purpose of open access, the author has applied a CC BY

public copyright licence to any Author Accepted Manuscript version arising from this

submission.

973
974
975 **Competing Interest**
976

E.R. is employed by Public Health England and holds an honorary contract with

Imperial College London. I.F.L. is Director of the Scottish Mycobacteria Reference

Laboratory. S.N. receives funding from German Center for Infection Research,

Excellenz Cluster Precision Medicine in Chronic Inflammation, Leibniz Science

Campus Evolutionary Medicine of the LUNG (EvoLUNG)tion EXC 2167. P.S. is a

consultant at Genoscreen. T.R. is funded by NIH and DoD and receives salary support

from the non-profit organization FIND. T.R. is a co-founder, board member and

shareholder of Verus Diagnostics Inc, a company that was founded with the intent of

developing diagnostic assays. Verus Diagnostics was not involved in any way with

data collection, analysis or publication of the results. T.R. has not received any

financial support from Verus Diagnostics. UCSD Conflict of Interest office has

reviewed and approved T.R.'s role in Verus Diagnostics Inc. T.R. is a co-inventor of a

provisional patent for a TB diagnostic assay (provisional patent #: 63/048.989). T.R. is

a co-inventor on a patent associated with the processing of TB sequencing data

(European Patent Application No. 14840432.0 & USSN 14/912,918). T.R. has agreed

to "donate all present and future interest in and rights to royalties from this patent" to

UCSD to ensure that he does not receive any financial benefits from this patent. S.S.

is working and holding ESOPs at HaystackAnalytics Pvt. Ltd. (Product: Using whole

genome sequencing for drug susceptibility testing for *Mycobacterium tuberculosis*).

G.F.G. is listed as an inventor on patent applications for RBD-dimer-based CoV

vaccines. The patents for RBD-dimers as protein subunit vaccines for SARS-CoV-2

have been licensed to Anhui Zhifei Longcom Biopharmaceutical Co. Ltd, China.

1000

1001

**Supplemental**

**Figure S1:** *Per drug MIC distributions of isolates plated on CRyPTIC designed variations on the Thermo Fischer Sensititre MYCOTB MIC plate; UKMYC5 (a) and UKMYC6 (b).* The solid black line represents the ecological resistance cut-off (ECOFF) for each drug as determined by [11]. Isolates with an MIC above the cut-off are considered resistant. N denotes the total number of isolates tested on each plate that returned a phenotype for each drug.

| UNIQUEID | COUNTRY OF ORIGIN | LINEAGE |
|---|---|---|
| site.11.subj.XTB-18-224.lab.XTB-18-224.iso.1 | UNKNOWN | Lineage 2 |
| site.10.subj.YA00026182.lab.YA00026182.iso.1 | ZAF | Lineage 4 |

**Table S1:** *Sample information for isolates classified as resistant to all 13 CRyPTIC drugs tested*. The country of origin is specified using the 3-letter country codes (alpha-3) defined by ISO 3166-1.

**Figure S2:** *Non-synonymous mutations found outside the RRDR of rpoB in RMR isolates and MDR isolates.* Presence of a coloured spot indicates that the mutation was found in RMR/MDR isolates and spot size corresponds to the proportion of RMR or MDR isolates carrying that mutation.

|  | Animal/ other | L1 | L2 | L3 | L4 | L6 | Total |
|---|---|---|---|---|---|---|---|
| **Albania** | 0 | 0 | 0 | 0 | 7 | 0 | 7 |
| **Algeria** | 0 | 0 | 0 | 0 | 25 | 0 | 25 |
| **Brazil** | 1 | 1 | 8 | 0 | 335 | 0 | 345 |
| **Burkina Faso** | 0 | 3 | 1 | 8 | 65 | 1 | 78 |
| **China** | 7 | 2 | 722 | 16 | 374 | 0 | 1121 |
| **Germany** | 1 | 25 | 235 | 61 | 403 | 1 | 726 |
| **India** | 11 | 676 | 1573 | 1184 | 560 | 0 | 4004 |
| **Italy** | 11 | 38 | 192 | 113 | 866 | 4 | 1224 |
| **Japan** | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| **Kyrgyzstan** | 0 | 0 | 25 | 0 | 3 | 0 | 28 |
| **Nepal** | 0 | 5 | 137 | 34 | 21 | 0 | 197 |
| **Nigeria** | 0 | 0 | 0 | 0 | 15 | 0 | 15 |
| **Pakistan** | 1 | 30 | 23 | 363 | 74 | 0 | 491 |
| **Peru** | 2 | 1 | 360 | 0 | 2328 | 0 | 2691 |
| **South Africa** | 1 | 61 | 874 | 57 | 1162 | 0 | 2155 |
| **Sweden** | 0 | 7 | 280 | 6 | 134 | 0 | 427 |
| **Taiwan** | 0 | 18 | 149 | 0 | 44 | 0 | 211 |
| **Tajikistan** | 0 | 0 | 15 | 0 | 4 | 0 | 19 |
| **Tunisia** | 0 | 0 | 0 | 0 | 2 | 0 | 2 |
| **Turkmenistan** | 0 | 0 | 88 | 1 | 30 | 0 | 119 |
| **Ukraine** | 0 | 0 | 21 | 0 | 9 | 0 | 30 |
| **UK** | 0 | 0 | 3 | 3 | 1 | 0 | 7 |
| **Vietnam** | 0 | 283 | 892 | 4 | 109 | 0 | 1288 |
| **Total** | 35 | 1150 | 5598 | 1850 | 6572 | 6 | 15211 |

**Table S2:** *Lineages –v- geographical location of origin/contribution for CRyPTIC isolates*

ALB  ALG  BRZ  BFA  CHN  GER  IND  ITL  JPN  KGZ  NPL  NGA  PAK  PER  ZAF  SWE  TWN  TJK  TUN  TKM  UKR  UK  VNM

| | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1.1 | 0 | 0 | 0 | 2 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 38 |
| 1.1.1.1 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 188 |
| 1.1.2 | 0 | 0 | 0 | 1 | 1 | 12 | 316 | 16 | 0 | 0 | 2 | 0 | 23 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1.3 | 0 | 0 | 1 | 0 | 0 | 1 | 52 | 4 | 0 | 0 | 1 | 0 | 1 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 1.2.1 | 0 | 0 | 0 | 0 | 0 | 8 | 12 | 11 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 3 | 16 | 0 | 0 | 0 | 0 | 0 | 0 | 24 |
| 1.2.2 | 0 | 0 | 0 | 0 | 1 | 2 | 259 | 5 | 0 | 0 | 2 | 0 | 4 | 0 | 54 | 3 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 2 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2.1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 20 | 0 | 0 | 0 | 0 | 0 | 0 | 34 |
| 2.2 | 0 | 0 | 5 | 1 | 284 | 19 | 272 | 14 | 0 | 0 | 68 | 0 | 6 | 231 | 181 | 13 | 56 | 0 | 0 | 0 | 0 | 0 | 0 | 271 |
| 2.2.1 | 0 | 0 | 0 | 0 | 37 | 4 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 3 | 224 | 1 | 21 | 0 | 0 | 0 | 0 | 0 | 0 | 26 |
| 2.2.10 | 0 | 0 | 0 | 0 | 1 | 117 | 1 | 84 | 0 | 16 | 0 | 0 | 3 | 0 | 0 | 70 | 0 | 5 | 0 | 75 | 9 | 1 | 0 | |
| 2.2.2 | 0 | 0 | 0 | 0 | 32 | 8 | 5 | 11 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 1 | 5 | 0 | 0 | 0 | 1 | 0 | 8 | |
| 2.2.3 | 0 | 0 | 2 | 0 | 172 | 3 | 290 | 4 | 0 | 0 | 21 | 0 | 2 | 1 | 3 | 1 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 100 |
| 2.2.4 | 0 | 0 | 0 | 0 | 3 | 3 | 4 | 8 | 0 | 0 | 0 | 0 | 0 | 0 | 37 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 70 |
| 2.2.5 | 0 | 0 | 0 | 0 | 59 | 1 | 12 | 5 | 0 | 0 | 0 | 0 | 2 | 84 | 1 | 1 | 13 | 0 | 0 | 0 | 0 | 0 | 0 | 45 |
| 2.2.6 | 0 | 0 | 1 | 0 | 22 | 8 | 226 | 2 | 0 | 0 | 9 | 0 | 7 | 4 | 204 | 1 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 154 |
| 2.2.7 | 0 | 0 | 0 | 0 | 97 | 4 | 691 | 5 | 0 | 1 | 38 | 0 | 0 | 20 | 206 | 5 | 18 | 0 | 0 | 0 | 0 | 0 | 0 | 164 |
| 2.2.8 | 0 | 0 | 0 | 0 | 4 | 0 | 42 | 0 | 0 | 0 | 1 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 2.2.9 | 0 | 0 | 0 | 0 | 0 | 64 | 0 | 51 | 0 | 6 | 0 | 0 | 0 | 2 | 0 | 174 | 0 | 10 | 0 | 11 | 11 | 2 | 0 | |
| 3 | 0 | 0 | 0 | 8 | 15 | 37 | 828 | 81 | 0 | 0 | 28 | 0 | 346 | 0 | 34 | 5 | 0 | 0 | 0 | 1 | 0 | 2 | 2 | |
| 3.1.1 | 0 | 0 | 0 | 0 | 0 | 17 | 17 | 25 | 0 | 0 | 1 | 0 | 0 | 0 | 19 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 2 | |
| 3.1.2 | 0 | 0 | 0 | 0 | 1 | 5 | 201 | 3 | 0 | 0 | 1 | 0 | 7 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 3.1.2.1 | 0 | 0 | 0 | 0 | 0 | 2 | 119 | 3 | 0 | 0 | 4 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4 | 0 | 0 | 1 | 0 | 6 | 4 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 100 | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | |
| 4.1 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.1.1 | 0 | 0 | 0 | 0 | 0 | 6 | 5 | 10 | 0 | 0 | 1 | 0 | 0 | 154 | 13 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.1.1.1 | 0 | 0 | 5 | 0 | 1 | 3 | 20 | 5 | 0 | 0 | 1 | 0 | 2 | 5 | 54 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 4.1.1.2 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.1.1.3 | 0 | 0 | 1 | 6 | 1 | 0 | 30 | 7 | 1 | 0 | 3 | 0 | 2 | 68 | 138 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 4.1.2 | 0 | 0 | 13 | 0 | 0 | 15 | 3 | 26 | 0 | 1 | 0 | 0 | 0 | 31 | 43 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 |
| 4.1.2.1 | 1 | 8 | 87 | 10 | 0 | 111 | 56 | 183 | 0 | 0 | 3 | 2 | 4 | 653 | 54 | 5 | 0 | 1 | 0 | 0 | 1 | 0 | 17 | |
| 4.1.3 | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 6 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.10 | 4 | 6 | 35 | 5 | 1 | 110 | 166 | 207 | 0 | 1 | 3 | 0 | 28 | 128 | 170 | 4 | 5 | 2 | 1 | 6 | 2 | 0 | 12 | |
| 4.2 | 0 | 0 | 0 | 0 | 3 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.2.1 | 0 | 0 | 0 | 0 | 0 | 9 | 3 | 46 | 0 | 0 | 1 | 0 | 2 | 0 | 2 | 4 | 0 | 0 | 0 | 8 | 3 | 1 | 0 | |
| 4.2.2 | 0 | 1 | 0 | 0 | 53 | 31 | 76 | 51 | 0 | 0 | 5 | 0 | 12 | 2 | 10 | 10 | 1 | 0 | 0 | 0 | 0 | 0 | 11 | |
| 4.2.2.1 | 0 | 0 | 0 | 0 | 0 | 5 | 0 | 9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3 | 0 | 0 | 17 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 96 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.1 | 0 | 3 | 0 | 1 | 0 | 1 | 2 | 21 | 0 | 0 | 0 | 0 | 0 | 8 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 4.3.2 | 0 | 1 | 10 | 0 | 0 | 3 | 1 | 8 | 0 | 0 | 0 | 0 | 0 | 245 | 21 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.2.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 234 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 4.3.3 | 1 | 4 | 34 | 2 | 0 | 33 | 99 | 81 | 0 | 1 | 1 | 0 | 0 | 576 | 120 | 102 | 2 | 1 | 1 | 11 | 2 | 0 | 1 | |
| 4.3.4 | 0 | 0 | 1 | 0 | 0 | 0 | 8 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.4.1 | 0 | 0 | 70 | 0 | 0 | 2 | 6 | 3 | 0 | 0 | 0 | 0 | 0 | 60 | 22 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.4.2 | 0 | 0 | 38 | 0 | 0 | 6 | 7 | 27 | 0 | 0 | 0 | 0 | 0 | 132 | 12 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| 4.3.4.2.1 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 75 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.4 | 0 | 0 | 0 | 0 | 0 | 0 | 13 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.4.1 | 0 | 0 | 10 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 11 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.4.1.1 | 0 | 0 | 4 | 0 | 0 | 16 | 4 | 66 | 0 | 0 | 1 | 0 | 0 | 22 | 180 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4 |
| 4.4.1.2 | 0 | 0 | 0 | 0 | 0 | 1 | 25 | 3 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| 4.4.2 | 0 | 0 | 0 | 0 | 124 | 1 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 0 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 16 |
| 4.5 | 0 | 0 | 0 | 0 | 179 | 28 | 15 | 14 | 0 | 0 | 1 | 0 | 18 | 5 | 1 | 0 | 30 | 0 | 0 | 3 | 0 | 0 | 29 | |
| 4.6 | 0 | 1 | 0 | 1 | 0 | 1 | 1 | 7 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.1.1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.1.2 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.2 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 14 | 0 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.2.1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.2.2 | 1 | 0 | 0 | 34 | 0 | 8 | 1 | 55 | 0 | 0 | 0 | 10 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| 6 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| Bovis | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 10 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| **Caprae** | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| **Mixed** | 0 | 1 | 9 | 0 | 16 | 3 | 89 | 9 | 0 | 1 | 0 | 0 | 6 | 38 | 25 | 12 | 2 | 0 | 0 | 3 | 0 | 0 | 48 |
| **Unknown** | 0 | 0 | 1 | 0 | 7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

1035 **Table S3: *Sub-lineages –v- geographical location of origin/contribution for***
1036 ***CRyPTIC isolates***

1037

1038

1039

1040
1041 **Figure S3**: ***A significant association between country and lineage can be seen***
1042 ***in the CRyPTIC data.*** Pearson's chi-squared test, X-squared = 7935.2, df = 110, *p* < 2.2e-16.
1043 The correlation plot indicates the relative contribution of each row-column pairing to the chi-
1044 square test score (%).
1045
1046

|  | INH | RIF | EMB | LEV | MXF | AMI | KAN | BDQ | CFZ | DLM | LZD | ETH | RFB |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| INH | 100.0 | 74.8 | 38.0 | 34.0 | 27.8 | 14.0 | 17.0 | 1.5 | 5.5 | 1.7 | 2.0 | 28.2 | 70.4 |
| RIF | 93.5 | 100.0 | 46.3 | 41.4 | 34.1 | 17.2 | 20.8 | 1.8 | 6.0 | 1.7 | 2.3 | 29.4 | 91.3 |
| EMB | 98.5 | 95.9 | 100.0 | 53.9 | 47.1 | 23.3 | 26.4 | 2.4 | 8.2 | 2.3 | 3.7 | 35.8 | 87.3 |
| LEV | 93.3 | 90.2 | 56.9 | 100.0 | 78.5 | 27.3 | 31.3 | 3.1 | 9.4 | 2.9 | 4.7 | 39.2 | 87.1 |
| MXF | 95.0 | 92.3 | 61.9 | 97.6 | 100.0 | 29.9 | 34.6 | 3.2 | 9.9 | 2.7 | 5.3 | 41.9 | 88.8 |
| AMI | 93.0 | 90.4 | 58.9 | 65.3 | 57.8 | 100.0 | 90.4 | 2.4 | 13.0 | 5.2 | 8.1 | 40.7 | 82.8 |
| KAN | 89.4 | 86.7 | 53.0 | 59.5 | 53.2 | 72.0 | 100.0 | 1.8 | 10.6 | 3.2 | 5.7 | 40.4 | 80.0 |
| BDQ | 79.4 | 77.8 | 49.5 | 60.7 | 49.5 | 20.0 | 18.9 | 100.0 | 52.4 | 12.9 | 14.0 | 34.9 | 76.9 |
| CFZ | 61.9 | 52.8 | 34.9 | 38.2 | 32.1 | 21.8 | 22.5 | 10.6 | 100.0 | 9.4 | 10.2 | 27.0 | 53.7 |
| DLM | 55.2 | 44.4 | 28.2 | 33.2 | 23.9 | 24.4 | 19.1 | 7.1 | 26.3 | 100.0 | 19.3 | 22.1 | 45.4 |
| LZD | 77.6 | 69.9 | 53.2 | 64.7 | 58.7 | 46.7 | 41.2 | 9.8 | 34.2 | 24.1 | 100.0 | 42.6 | 66.7 |
| ETH | 96.5 | 79.4 | 47.0 | 48.7 | 41.9 | 21.1 | 26.2 | 2.2 | 8.3 | 2.4 | 3.9 | 100.0 | 77.1 |
| RFB | 93.3 | 96.8 | 44.5 | 42.1 | 34.4 | 16.6 | 20.3 | 1.9 | 6.4 | 1.9 | 2.3 | 29.9 | 100.0 |

**Table S4: *Co-occurrence of antibiotic resistance in CRyPTIC M. tuberculosis isolates.*** The probability (%) of an isolate being resistant to Drug 2 (top) if it is resistant to Drug 1 (left). Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, LEV = levofloxacin, MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline, CFZ = clofazimine, DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.

# References

1. WHO, *World Health Organisation: Global Tuberculosis Report 2020.* 2020.
2. Boehme, C.C., et al., *Feasibility, diagnostic accuracy, and effectiveness of decentralised use of the Xpert MTB/RIF test for diagnosis of tuberculosis and multidrug resistance: a multicentre implementation study.* Lancet, 2011. **377**(9776): p. 1495-505.
3. Shinnick, T.M., et al., *Evaluation of the Cepheid Xpert MTB/RIF assay.* Expert Rev Mol Diagn, 2015. **15**(1): p. 9-22.
4. Sanchez-Padilla, E., et al., *Detection of Drug-Resistant Tuberculosis by Xpert MTB/RIF in Swaziland.* New England Journal of Medicine, 2015. **372**(12): p. 1181-1182.
5. Beckert, P., et al., *MDR M. tuberculosis outbreak clone in Eswatini missed by Xpert has elevated bedaquiline resistance dated to the pre-treatment era.* Genome Medicine, 2020. **12**(1): p. 104.
6. Makhado, N.A., et al., *Outbreak of multidrug-resistant tuberculosis in South Africa undetected by WHO-endorsed commercial tests: an observational study.* Lancet Infect Dis, 2018. **18**(12): p. 1350-1359.
7. Consortium, T.C., et al., *Prediction of Susceptibility to First-Line Tuberculosis Drugs by DNA Sequencing.* N Engl J Med, 2018. **379**(15): p. 1403-1415.
8. Pankhurst, L.J., et al., *Rapid, comprehensive, and affordable mycobacterial diagnosis with whole-genome sequencing: a prospective study.* Lancet Respir Med, 2016. **4**(1): p. 49-58.

1078   9.     Kalokhe, A.S., et al., *Multidrug-resistant tuberculosis drug susceptibility and*
1079           *molecular diagnostic testing.* Am J Med Sci, 2013. **345**(2): p. 143-8.
1080   10.    Rancoita, P.M.V., et al., *Validating a 14-Drug Microtiter Plate Containing*
1081           *Bedaquiline and Delamanid for Large-Scale Research Susceptibility Testing of*
1082           *Mycobacterium tuberculosis.* Antimicrob Agents Chemother, 2018. **62**(9).
1083   11.    Consortium, T.C. and P.W. Fowler, *Epidemiological cutoff values for a 96-well broth*
1084           *microdilution plate for high-throughput research antibiotic susceptibility testing of M.*
1085           *tuberculosis.* medRxiv, 2021: p. 2021.02.24.21252386.
1086   12.    Fowler, P.W., et al., *Automated detection of bacterial growth on 96-well plates for*
1087           *high-throughput drug susceptibility testing of Mycobacterium tuberculosis.*
1088           Microbiology (Reading), 2018. **164**(12): p. 1522-1530.
1089   13.    Fowler, P.W., et al., *BashTheBug: a crowd of volunteers reproducibly and accurately*
1090           *measure the minimum inhibitory concentrations of 13 antitubercular drugs from*
1091           *photographs of 96-well broth microdilution plates.* bioRxiv, 2021: p.
1092           2021.07.20.453060.
1093   14.    Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina*
1094           *sequence data.* Bioinformatics, 2014. **30**(15): p. 2114-20.
1095   15.    Li, H. and R. Durbin, *Fast and accurate short read alignment with Burrows-Wheeler*
1096           *transform.* Bioinformatics, 2009. **25**(14): p. 1754-60.
1097   16.    Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics,
1098           2009. **25**(16): p. 2078-9.
1099   17.    Iqbal, Z., et al., *De novo assembly and genotyping of variants using colored de Bruijn*
1100           *graphs.* Nat Genet, 2012. **44**(2): p. 226-32.
1101   18.    Hunt, M., et al., *Antibiotic resistance prediction for Mycobacterium tuberculosis from*
1102           *genome sequence data with Mykrobe.* Wellcome Open Research, 2019. **4**(191).
1103   19.    Walker, T.M., et al., *Assessment of Mycobacterium tuberculosis transmission in*
1104           *Oxfordshire, UK, 2007-12, with whole pathogen genome sequences: an observational*
1105           *study.* Lancet Respir Med, 2014. **2**(4): p. 285-292.
1106   20.    Howe, K., A. Bateman, and R. Durbin, *QuickTree: building huge Neighbour-Joining*
1107           *trees of protein sequences.* Bioinformatics, 2002. **18**(11): p. 1546-7.
1108   21.    Yu, G., *Using ggtree to Visualize Data on Tree-Like Structures.* Curr Protoc
1109           Bioinformatics, 2020. **69**(1): p. e96.
1110   22.    Gagneux, S., *Ecology and evolution of Mycobacterium tuberculosis.* Nat Rev
1111           Microbiol, 2018. **16**(4): p. 202-213.
1112   23.    Coscolla, M. and S. Gagneux, *Consequences of genomic diversity in Mycobacterium*
1113           *tuberculosis.* Semin Immunol, 2014. **26**(6): p. 431-44.
1114   24.    Holt, K.E., et al., *Frequent transmission of the Mycobacterium tuberculosis Beijing*
1115           *lineage and positive selection for the EsxW Beijing variant in Vietnam.* Nat Genet,
1116           2018. **50**(6): p. 849-856.
1117   25.    Chiner-Oms, A. and I. Comas, *Large genomics datasets shed light on the evolution of*
1118           *the Mycobacterium tuberculosis complex.* Infect Genet Evol, 2019. **72**: p. 10-15.
1119   26.    Freschi, L., et al., *Population structure, biogeography and transmissibility of*
1120           *Mycobacterium tuberculosis.* bioRxiv, 2020: p. 2020.09.29.293274.
1121   27.    Borrell, S., et al., *Reference set of Mycobacterium tuberculosis clinical strains: A tool*
1122           *for research and product development.* PLoS One, 2019. **14**(3): p. e0214088.
1123   28.    Sharma, R., et al., *High degree of fluoroquinolone resistance among pulmonary*
1124           *tuberculosis patients in New Delhi, India.* Indian J Med Res, 2019. **149**(1): p. 62-66.
1125   29.    Ghimire, S., et al., *Treatment outcomes of patients with MDR-TB in Nepal on a*
1126           *current programmatic standardised regimen: retrospective single-centre study.* BMJ
1127           Open Respir Res, 2020. **7**(1).

30.    Oppong, Y.E.A., et al., *Genome-wide analysis of Mycobacterium tuberculosis*
*polymorphisms reveals lineage-specific associations with drug resistance.* BMC
Genomics, 2019. **20**(1): p. 252.

31.    Ford, C.B., et al., *Mycobacterium tuberculosis mutation rate estimates from different*
*lineages predict substantial differences in the emergence of drug-resistant*
*tuberculosis.* Nat Genet, 2013. **45**(7): p. 784-90.

32.    Berrada, Z.L., et al., *Rifabutin and rifampin resistance levels and associated rpoB*
*mutations in clinical isolates of Mycobacterium tuberculosis complex.* Diagn
Microbiol Infect Dis, 2016. **85**(2): p. 177-81.

33.    Maitre, T., et al., *Are moxifloxacin and levofloxacin equally effective to treat XDR*
*tuberculosis?* J Antimicrob Chemother, 2017. **72**(8): p. 2326-2333.

34.    Dijkstra, J.A., et al., *In Vitro Susceptibility of Mycobacterium tuberculosis to*
*Amikacin, Kanamycin, and Capreomycin.* Antimicrob Agents Chemother, 2018.
**62**(3).

35.    Ho, J., P. Jelfs, and V. Sintchenko, *Fluoroquinolone resistance in non-multidrug-*
*resistant tuberculosis-a surveillance study in New South Wales, Australia, and a*
*review of global resistance rates.* Int J Infect Dis, 2014. **26**: p. 149-53.

36.    Miotto, P., et al., *A standardised method for interpreting the association between*
*mutations and phenotypic drug resistance in Mycobacterium tuberculosis.* Eur Respir
J, 2017. **50**(6).

37.    Timothy M Walker, P.M., Claudio Köser†, Philip Fowler, Jeff Knaggs, Zamin Iqbal,
Martin Hunt, Leonid Chindelevitch, Maha Farhat, Daniela Cirillo, Inaki Comas,
Jamie Posey, Shaheed Vally Omar, Tim Peto, Anita Suresh, Swapna Uplekar, Sacha
Laurant, Rebecca Colman, Carl-Michael Nathanson, Matteo Zignol, Ann Sarah
Walker, The CRyPTIC Consortium, The Seq&Treat Consortium, Derrick Crook,
Nazir Ismail and Timothy Rodwell, *The 2021 WHO catalogue of Mycobacterium*
*tuberculosis mutations associated with drug resistance: a new global standard for*
*molecular diagnostics (in prep).* 2021.

38.    Comas, I., et al., *Whole-genome sequencing of rifampicin-resistant Mycobacterium*
*tuberculosis strains identifies compensatory mutations in RNA polymerase genes.* Nat
Genet, 2011. **44**(1): p. 106-10.

39.    Almeida, D., et al., *Mutations in pepQ Confer Low-Level Resistance to Bedaquiline*
*and Clofazimine in Mycobacterium tuberculosis.* Antimicrob Agents Chemother,
2016. **60**(8): p. 4590-9.

40.    Degiacomi, G., et al., *In vitro Study of Bedaquiline Resistance in Mycobacterium*
*tuberculosis Multi-Drug Resistant Clinical Isolates.* Front Microbiol, 2020. **11**: p.
559469.

41.    Hartkoorn, R.C., S. Uplekar, and S.T. Cole, *Cross-resistance between clofazimine and*
*bedaquiline through upregulation of MmpL5 in Mycobacterium tuberculosis.*
Antimicrob Agents Chemother, 2014. **58**(5): p. 2979-81.

42.    Nimmo, C., et al., *Population-level emergence of bedaquiline and clofazimine*
*resistance-associated variants among patients with drug-resistant tuberculosis in*
*southern Africa: a phenotypic and phylogenetic analysis.* Lancet Microbe, 2020. **1**(4):
p. e165-e174.

43.    Singh, R., et al., *PA-824 kills nonreplicating Mycobacterium tuberculosis by*
*intracellular NO release.* Science, 2008. **322**(5906): p. 1392-5.

44.    Mvelase, N.R., et al., *Evolving rifampicin and isoniazid mono-resistance in a high*
*multidrug-resistant and extensively drug-resistant tuberculosis region: a retrospective*
*data analysis.* BMJ Open, 2019. **9**(11): p. e031663.

45.  Park, S., et al., *Treatment outcomes of rifampin-sparing treatment in patients with*
*pulmonary tuberculosis with rifampin-mono-resistance or rifampin adverse events: A*
*retrospective cohort analysis.* Respir Med, 2017. **131**: p. 43-48.
46.  Villegas, L., et al., *Prevalence, Risk Factors, and Treatment Outcomes of Isoniazid-*
*and Rifampicin-Mono-Resistant Pulmonary Tuberculosis in Lima, Peru.* PLoS One,
2016. **11**(4): p. e0152933.
47.  Oppong, Y.E.A., et al., *Genome-wide analysis of Mycobacterium tuberculosis*
*polymorphisms reveals lineage-specific associations with drug resistance.* BMC
Genomics, 2019. **20**(1): p. 252.
48.  Pang, Y., et al., *Study of the rifampin monoresistance mechanism in Mycobacterium*
*tuberculosis.* Antimicrob Agents Chemother, 2013. **57**(2): p. 893-900.
49.  Kigozi, E., et al., *Prevalence and patterns of rifampicin and isoniazid resistance*
*conferring mutations in Mycobacterium tuberculosis isolates from Uganda.* PLoS
One, 2018. **13**(5): p. e0198091.
50.  Dean, A.S., et al., *Prevalence and genetic profiles of isoniazid resistance in*
*tuberculosis patients: A multicountry analysis of cross-sectional data.* PLoS Med,
2020. **17**(1): p. e1003008.
51.  Migliori, G.B., et al., *Protecting the tuberculosis drug pipeline: stating the case for*
*the rational use of fluoroquinolones.* Eur Respir J, 2012. **40**(4): p. 814-22.
52.  Heysell, S.K., et al., *Hearing loss with kanamycin treatment for multidrug-resistant*
*tuberculosis in Bangladesh.* Eur Respir J, 2018. **51**(3).
53.  Organisation, W.H., *Update on the use of nucleic acid amplification tests to detect TB*
*and drug-resistant TB: rapid communication.* 2021.
54.  Olaru, I.D., et al., *Bedaquiline-based treatment regimen for multidrug-resistant*
*tuberculosis.* Eur Respir J, 2017. **49**(5).
55.  Yuan, S., et al., *Clofazimine broadly inhibits coronaviruses including SARS-CoV-2.*
Nature, 2021. **593**(7859): p. 418-423.
56.  Falzon, D., et al., *World Health Organization treatment guidelines for drug-resistant*
*tuberculosis, 2016 update.* Eur Respir J, 2017. **49**(3).
57.  Huitric, E., et al., *Rates and mechanisms of resistance development in Mycobacterium*
*tuberculosis to a novel diarylquinoline ATP synthase inhibitor.* Antimicrob Agents
Chemother, 2010. **54**(3): p. 1022-8.
58.  Carter J, The CRyPTIC Consortium, *Quantitative measurement of antibiotic*
*resistance in Mycobacterium tuberculosis reveals genetic determinants of resistance*
*and susceptibility in a target gene approach.* 2021, in preparation.
59.  Earle S and Wilson D, The CRyPTIC Consortium, *Genome-wide association studies*
*of global Mycobacterium tuberculosis resistance to thirteen antimicrobials in 10,228*
*genomes.* 2021, in preparation.
60.  Lachapelle AS, The CRyPTIC Consortium, *Predicting Susceptibility to First- and*
*Second-line Tuberculosis Drugs by DNA sequencing and Machine Learning.* 2021, in
preparation.
61.  Sonnenkalb, L., et al., *Deciphering Bedaquiline and Clofazimine Resistance in*
*Tuberculosis: An Evolutionary Medicine Approach.* bioRxiv, 2021: p.
2021.03.19.436148.
1221
1222