1     # A data compendium of *Mycobacterium tuberculosis*

2     # antibiotic resistance

4     The CRyPTIC Consortium

## Abstract

The Comprehensive Resistance Prediction for Tuberculosis: an International Consortium (CRyPTIC) presents here a compendium of 15,211 *Mycobacterium tuberculosis* global clinical isolates, all of which have undergone whole genome sequencing (WGS) and have had their minimum inhibitory concentrations to 13 antitubercular drugs measured in a single assay. It is the largest matched phenotypic and genotypic dataset for *M. tuberculosis* to date. Here, we provide a summary detailing the breadth of data collected, along with a description of how the isolates were collected and uniformly processed in CRyPTIC partner laboratories across 23 countries. The compendium contains 6,814 isolates resistant to at least one drug, including 2,129 samples that fully satisfy the clinical definitions of rifampicin resistant (RR), multi-drug resistant (MDR), pre-extensively drug resistant (pre-XDR) or extensively drug resistant (XDR). Accurate prediction of resistance status (sensitive/resistant) to eight antitubercular drugs by using a genetic mutation catalogue is presented along with the presence of suspected resistance-conferring mutations for isolates resistant to the newly introduced drugs bedaquiline, clofazimine, delamanid and linezolid. Finally, a case study of rifampicin mono-resistance demonstrates how this compendium could be used to advance our genetic understanding of rare resistance phenotypes. The compendium is fully open-source and it is hoped that the dataset will facilitate and inspire future research for years to come.

## Introduction

Tuberculosis (TB) is a curable and preventable disease; 85% of those afflicted can be successfully treated with a six-month regimen. Despite this, TB is the world's top infectious disease killer (current SARS-CoV-2 pandemic excepted) with 10 million new cases and 1.2 million deaths estimated in 2019 alone (1). Furthermore, drug resistant TB (DR-TB) is a continual threat; almost half a million cases resistant to the first-line drug rifampicin (RR-TB) were estimated, with three quarters of these estimated to be multidrug-resistant (MDR-TB, resistant to first-line drugs isoniazid and rifampicin) (1). Worryingly, only 44% of DR-TB cases were officially notified and just over half of these cases were successfully treated (57%) (1).

To address these issues, the World Health Organisation (WHO) is encouraging the development of better, faster and more targeted diagnostic and treatment strategies through its EndTB campaign (1). Of particular interest is universal drug susceptibility testing (DST). Conventionally, DST relies on lengthy (4 weeks minimum) culture-based methods that require strict biosafety conditions for *Mycobacterium tuberculosis.* The development of rapid genetics-based assays has decreased diagnostic time to as little as 2 hours through the detection of specific resistance conferring mutations *e.g.* the Cepheid Xpert® MTB/RIF test (2,3). However, assay bias towards specific genic regions can result in misdiagnosis of resistance, the prescription of ineffective treatment regimens and subsequent spread of multi-drug resistant disease, as seen during an MDR outbreak in Eswatini (4–6). Furthermore, detection of rifampicin resistance is used to infer MDR-TB epidemiologically as rifampicin resistance tends to coincide with resistance to isoniazid (7). While this *modus operandi* is successful at pragmatically identifying potential MDR cases quickly

and effectively, it is not generally true that a single path exists for developing MDR or

extensively drug resistant TB (XDR = MDR/RR + resistance to at least one

fluoroquinolone and either bedaquiline or linezolid).

Whole-genome sequencing (WGS) has the potential to reveal the entirety of

the *M. tuberculosis* genetic resistance landscape for any number of drugs

simultaneously whilst enabling a more rapid turnaround time and reduction in cost

compared to DST culture-based methods (8). However, the success of WGS as a

diagnostic tool wholly depends on there being a comprehensive and accurate

catalogue of resistance-conferring mutations for each drug. Recent advances have

shown that genotypic predictions of resistance correlate well with DST measurements

for first-line drugs (7). However, the mechanisms of resistance to second-line drugs

along with the new and re-purposed drugs (NRDs) are less well understood despite

their increased administration in clinics as MDR cases climb(1,9).

To address these shortcomings, the Comprehensive Resistance Prediction for

Tuberculosis: an International Consortium (CRyPTIC) has collected *M. tuberculosis*

clinical isolates worldwide to survey the genetic variation associated with resistance

to 13 antitubercular drugs, specifically the first-line drugs rifampicin, isoniazid,

ethambutol, the second-line drugs amikacin, kanamycin, rifabutin, levofloxacin,

moxifloxacin, ethionamide, and the new and re-purposed drugs bedaquiline,

clofazimine, delamanid and linezolid. Here, we introduce and describe these data in

the form of an open-access compendium of 15,211 isolates, each of which has had its

genomic sequence determined and DST profile measured (10). This compendium is

the largest drug screening effort to date for *M. tuberculosis* in a 'one isolate – one

microscale assay' format across defined compound concentration ranges. The

presented dataset forms the backbone for several studies being put forth by the

consortium to achieve its ultimate aim (11–15). By being fully open-access, it is hoped

that this compendium will prove an invaluable resource to accelerate and improve

antimicrobial resistance (AMR) diagnostic development for TB, both by the enrichment

of mutation catalogues for WGS resistance prediction and the identification of

important diagnostic gaps and drug resistance patterns.

**Methods**

***Ethics***

Approval for the CRyPTIC study was obtained by Taiwan Centers for Disease

Control IRB No. 106209, University of KwaZulu Natal Biomedical Research Ethics

Committee (UKZN BREC) (reference BE022/13) and University of Liverpool Central

University Research Ethics Committees (reference 2286), Institutional Research

Ethics Committee (IREC) of The Foundation for Medical Research, Mumbai (Ref nos.

FMR/IEC/TB/01a/2015 and FMR/IEC/TB/01b/2015), Institutional Review Board of

P.D. Hinduja Hospital and Medical Research Centre, Mumbai (Ref no. 915-15-CR

[MRC]), scientific committee of the Adolfo Lutz Institute (CTC-IAL 47-J / 2017) and in

the Ethics Committee (CAAE: 81452517.1.0000.0059) and Ethics Committee review

by Universidad Peruana Cayetano Heredia (Lima, Peru) and LSHTM (London, UK).

***Sample collection***

Participating collection centres varied in their isolate collection approaches and

timescales *(e.g.* longitudinal sampling, rolling patient visits, biobank stocks), but the

consortium collectively aimed to oversample for *M. tuberculosis* isolates with drug

resistance and multi-drug resistance. A standard operating protocol for sample

processing was defined by CRyPTIC as previously described and is discussed in more

detail below in relevant sub-sections (10,11).

***Plate assay***

The CRyPTIC consortium designed two versions of the Sensititre MYCOTB

plate (Thermo Fisher Scientific Inc., USA) named the "UKMYC5" and "UKMYC6"

microtitre plates (10,11). These plates contain five to ten doubling dilutions of 13

antibiotics (rifampicin (RIF), rifabutin (RFB), isoniazid (INH), ethambutol (EMB),

levofloxacin (LEV), moxifloxacin (MXF), amikacin (AMI), kanamycin (KAN),

ethionamide (ETH), clofazimine (CFZ), linezolid (LZD), delamanid (DLM), and

bedaquiline (BDQ)). Delamanid and bedaquiline were provided by Otsuka

Pharmaceutical Co., Ltd. and Janssen Pharmaceutica respectively. The UKMYC5

plate also contained para-aminosalicylic acid (PAS), but the MICs were not

reproducible and hence it was excluded from the UKMYC6 plate design and is not

included in any subsequent analysis (10).

A standard operating protocol for sample processing was defined by CRyPTIC

as previously described (10,11). Clinical samples were sub-cultured using 7H10 agar

plates, Lowenstein-Jensen tubes or MGIT tubes. Bacterial cell suspensions (0.5

McFarland standard, saline Tween) prepared from (no later than) 14-day old colonies

were diluted 100X in 10 ml enriched 7H9 broth prior to plate inoculation. A semi-

automated Sensititre Autoinoculator (Thermo Fisher, Scientific Inc., USA) was used to

inoculate 100 µl prepared cell suspensions ($1.5 \times 10^5$ CFU/ml [$5 \times 10^4$ CFU/ml - $5 \times$

$10^5$ CFU/ml]) into each well of a UKMYC5/6 microdilution plate. The plate was sealed

and incubated for 14 days at 37°C. Quality control runs were performed periodically

using *M. tuberculosis* H37Rv ATCC 27294, which is sensitive to all drugs on the plates.

***Minimum Inhibitory Concentration (MIC) measurements***

Minimum inhibitory concentrations (MICs) for each drug were read after

incubation for 14 days by a laboratory scientist using a Thermo Fisher Sensititre™

Vizion™ digital MIC viewing system (10). The Vizion apparatus was also used to take

a high contrast photograph of the plate with a white background, from which the MIC

was measured again using the Automated Mycobacterial Growth Detection Algorithm

(AMyGDA) software (16). The AMyGDA algorithm was specifically developed to

automate and perform quality control of MIC measurements, and to facilitate machine

learning studies within the consortium. AMyGDA detects the boundaries of each well

using a Hough transform for circles and measures growth as the number of dark pixels

within the area contained by this boundary.

All images where the MICs measured by Vizion and AMyGDA were different

were uploaded to a citizen science project, BashTheBug, on the Zooniverse platform

(17). Each image was then classified by ≥11 volunteers and the median classification

taken. MICs were then classified as high (at least two methods concur on the MIC),

medium (either a scientist recorded a MIC measurement using Vizion but did not store

the plate picture, or Vizion and AMyGDA disagree and there is no BashTheBug

measurement), or low (all three methods disagree) quality.

To ensure adequate data coverage for *this* study, we took the MIC from the

Vizion reading provided by the trained laboratory scientist if it was annotated as having

medium or low quality.

### *Binary phenotype classification*

Binary phenotypes (resistant/susceptible) were assigned from the MICs by applying epidemiological cut-off values (11); samples with MICs at or below the ECOFF are, by definition, wild-type and hence assigned to be susceptible to the drug in question (11). Samples with MICs above the ECOFF are therefore classified as resistant (Fig. S1, Table S1). Please see (11) for the body of work supporting the use of the ECOFF relative to the compendium isolates and supplemental Table S1 for the ECOFFs for each drug tested.

### *Genomic data processing and variant calling*

Short-read, paired end libraries were sequenced on Illumina machines and the resulting FASTQ files were processed using the bespoke pipeline Clockwork (v0.8.3, github.com/iqbal-lab-org/clockwork, (18)). Briefly, all raw sequencing files were indexed into a relational database with which Clockwork proceeds. Human, nasopharyngeal flora and human immunodeficiency virus related reads were removed and remaining reads were trimmed (adapters and low quality ends) using Trimmomatic and mapped with BWA-MEM to the *M. tuberculosis* H37Rv reference genome (NC000962.3) (19,20). Read duplicates were removed. Genetic variants were called independently using Cortex and SAMtools, two variant callers with orthogonal strengths (SAMtools a high sensitivity SNP caller, and Cortex a high specificity SNP and indel caller) (21,22). The two call sets were merged to produce a final call set, using the Minos adjudication tool to resolve loci where the two callers disagreed, by remapping reads to an augmented genome containing each alternative allele (23). Default filters of a minimum depth of 5x, a fraction of supporting reads of 0.9 (minos)

and a genotype confidence percentile (GCP) filter of 0.5 were applied. The GCP filter

is a normalised likelihood ratio test, giving a measure of confidence in the called allele

compared with the other alternatives, and is described in (23). This produced one

variant call format (VCF) file per sample, each only describing positions where that

sample differed from the reference.

These filtered VCFs were then combined, to produce a single non-redundant

list of all variants seen in the cohort. All samples were then processed a second time

with Minos, remapping reads to a graphical representation of all the segregating

variation within the cohort, generating VCF files which had an entry at all variable

positions (thus for all samples, most positions would be genotyped as having the

reference allele). These "regenotyped" VCFs were later used to calculate pairwise

distances (see below).

To remove untrustworthy loci, a genome mask was applied to the resulting VCF

files (regions identified with self-blast matches in (24) comprising of 324,971 bp of the

reference genome). Furthermore, positions with less than 90% of total samples

passing default Clockwork/Minos variant call filters (described above) were filtered out,

comprising 95,703 bp of the genome, of which 55,980 bp intersect with the genome

mask.

***Resistance prediction using a genetic catalogue***

A hybrid catalogue of genetic variants associated with resistance to first- and

second- line drugs based on existing catalogues was created and can be found at

github.com/oxfordmmm/tuberculosis_amr_catalogues/blob/public/catalogues/NC_00

0962.3/NC_000962.3_CRyPTIC_v1.311_GARC1_RUS.csv (7,25). We specifically did not use the recent WHO catalogue to avoid circularity and over-training, as that catalogue was developed (via prior literature, expert rules and a heuristic algorithm) based partially on these isolates (26)).

The resulting VCF file for each isolate (see *"Genomic data processing and variant calling"* section above) was compared to the genetic catalogue to determine the presence or absence of resistance-associated mutations for eight drugs: rifampicin, isoniazid, ethambutol, levofloxacin, moxifloxacin, amikacin, kanamycin and ethionamide. We did not apply the approach used in (7) to make a prediction if a novel mutation was detected in a known resistance gene, as we simply wanted to measure how well a pre-CRyPTIC catalogue could predict resistance in the compendium. These results (found in PREDICTIONS.csv, see "Data availability" section for access) were then compared to the binary phenotypes (see "Binary phenotype classification" section for how these were defined) with the following metrics calculated: TP: the number of phenotypically resistant samples are that correctly identified as resistant ("true positives"); FP, the number of phenotypically susceptible samples that are falsely identified as resistant ("false positives"); TN, the number of phenotypically susceptible samples that are correctly identified as susceptible ("true negatives"); FN, the number of phenotypically resistant samples that are incorrectly identified as susceptible ("false negative"); VME, very major error rate (false-negative rate), 0-1; ME, major error rate (false-positive rate), 0-1; PPV, positive predictive value, 0-1; NPV, negative predictive value.

***Phylogenetic tree construction***

A pairwise genetic distance matrix was constructed for 15,211 isolates by

comparing pairs of regenotyped VCF files (see "*Genomic data processing and variant*

*calling"* section above for more details). A neighbourhood-joining tree was constructed

from the distance matrix using *quicktree* (27). Tree visualisation and annotation was

performed using the R library *ggtree* (28). *M. tuberculosis* lineages were assigned

using Mykrobe and are represented by the coloured dots at the branch termini of the

tree (23). For isolates that had 'mixed' lineage classification (*i.e.,* 2 lineages were found

present in the sample by Mykrobe, $n$ = 225, 1.5%), the first of the two lineages was

assigned to the isolate. *ggtree* was also used to construct the trees depicting

bedaquiline, clofazimine and delamanid resistant isolates.

**The Data**

All data can be found at ftp.ebi.ac.uk/pub/databases/cryptic/reuse/. The FTP

site contains two top level directories: "reuse" and "reproducibility". In total, the

compendium contains data entries for 15,211 isolates. Note that various filtering

criteria have been applied in both this study and other CRyPTIC studies and thus final

isolate numbers presented across the publications will vary.

*"reuse" directory*

We point the reader to this directory to gain access to CRyPTIC project data.

"CRyPTIC_reuse_table_20221019.csv" contains genotypic and phenotypic data

relating to the figures and summaries listed in this manuscript and is what we present

as a general use reference table for most future projects. It includes binary phenotypes

(R/S), MICs, phenotype quality metrics, and ENA sample IDs for 12,288 compendium

isolates (see "Quality assurance of the minimum inhibitory concentrations for 13 drugs"

section below in Results for filters applied to obtain this final set of isolated). It also

includes file paths to each isolate's VCF file and 'regenotyped' VCF file (VCF files

which have an entry at all variable positions, see "Genomic data processing and

variant calling" section above for more).

*"reproducibility" directory*

This directory contains the data used for multiple CRyPTIC project publications

referenced throughout this manuscript. As stated above, each project has taken

slightly different subsets of this data as documented in those papers. For example,

see how tables such as "MUTATIONS.csv" and "GENOTYPES.csv" were used and

filtered, (along with others) in this study to obtain the reuse file

"CRyPTIC_reuse_table_20221019.csv" in Figure 1. Again, for optimal use of

CRyPTIC data in your own project, please refer to

"CRyPTIC_reuse_table_20221019.csv" in the "reuse" directory. All data for this study

were analysed and visualised using either R or python3 libraries and packages. See

github.com/kerrimalone/Brankin_Malone_2022 for codebase.

# Results

## 15,211 M. tuberculosis clinical isolates

The CRyPTIC compendium contains 15,211 isolates for which both genomic and phenotypic data was collected by 23 of the partner countries across the continents of Asia, Africa, South America and Europe. An overview of the processing of the isolates is presented in Figure 1, and for a full description please see Methods.

**Figure 1:** *Processing sequencing and MIC data for 15,211 M. tuberculosis isolates* Briefly: Each isolate was DNA sequenced using an Illumina machine and plated onto 96 well plates (UKMYC5/6) containing 5-10x doubling dilutions of 13 antitubercular drugs for drug susceptibility testing. Associated metadata (including country of origin and processing laboratory) was recorded. DNA variant calling and analysis was performed using Clockwork and minos. After 14 days, MIC measurements were measured by a trained scientist using Vizion, and the plate was

photographed to measure the MIC using the automated AMyGDA software and citizen scientists
from BashTheBug. After quality control procedures, phenotypic MIC data for 2,922 isolates were
removed. The compendium therefore contains 15,211 isolates with whole genome sequencing
data, 12,289 of which have matched phenotypic data, presented in
"CRyPTIC_reuse_table_20211019.csv" via an FTP site (see methods). The data tables
GENOTYPES.csv, VARIANTS.csv, MUTATIONS.csv, SAMPLES.csv and PHENOTYPES.csv
used for the analyses presented in this manuscript are also accessible via the FTP site (see
methods).

The 15, 211 isolates originated from 23 different countries (Fig. 2). The largest

number of isolates were contributed by India (*n* = 4,004), Peru (*n* = 2,691), South

Africa (*n* = 2,155), Vietnam (*n* = 1,288) and China (*n* = 1,121).

**Figure 2: *Geographical distribution of CRyPTIC M. tuberculosis clinical isolates.***
The total number of isolates contributed by each country is depicted. Where labels overlap, labels
are exploded and lines are used to indicate country of origin, from left to right: UK, Albania, India.
Where the origin of an isolate was not known, the collection site identity was assigned (this
occurred for 269 isolates in Germany, 17 isolates in India, 6 isolates in Peru, 885 isolates in Italy,

510 isolates in South Africa, 357 isolates in Sweden, 208 isolates in Taiwan, 1 isolate in Brazil and
4 isolates in the UK).

Lineage assignment revealed that 99.7% of the 15,211 isolates belong to the

four main *M. tuberculosis* lineages (L1-L4). The pie-charts in supplemental Figure S2

show the proportion of isolates amongst the different lineages (Table S2) and sub-

lineages (Table S3) for each location. Like previous studies, we see a strong

association between geolocation and lineage (Pearson's chi-squared test, $p$ < 2.2e-

16, Fig. S3) (29,30). The phylogenetic tree in supplemental figure S4 further highlights

the strong population structure of this collection of isolates, with isolates clustering

according to lineage. Typically under-sampled in current databases and biobanks, the

L3 isolates in this study represent the largest collection to date (31).

Although these 15,211 *M. tuberculosis* isolates were plated to determine their

MICs to 13 antitubercular drugs, regular quality assurance checks detected problems

with plate inoculation and reading in two laboratories.  Therefore, 2,922 isolates were

removed from the dataset, leaving a total of 12,289 isolates with matched phenotypic

and genotypic data for further analysis (Fig.1). Due to wells being skipped and other

phenomena that prevent an MIC being measured, 88.1% of the isolates had a

phenotype for all 13 drugs on the plate. For each drug, the number of isolates with an

MIC measurement, and the associated quality of the reading, is presented in Table 1.

|  | MIC MEASUREMENTS | HIGH QUALITY | MEDIUM QUALITY | LOW QUALITY |
|---|---|---|---|---|
| **INH** | 12070 | 9519 | 1351 | 1200 |
| **RIF** | 12099 | 8955 | 1356 | 1788 |
| **EMB** | 12158 | 7506 | 1355 | 3297 |
| **LEV** | 12163 | 7774 | 1354 | 3035 |
| **MXF** | 12194 | 6785 | 1353 | 4056 |
| **AMI** | 12072 | 8973 | 1350 | 1749 |
| **KAN** | 12130 | 9333 | 1355 | 1442 |
| **BDQ** | 12068 | 8536 | 1355 | 2177 |
| **CFZ** | 12049 | 7763 | 1352 | 2934 |
| **DLM** | 11927 | 8095 | 1349 | 2483 |
| **LZD** | 12189 | 7141 | 1355 | 3693 |
| **ETH** | 12132 | 8821 | 1355 | 1956 |
| **RFB** | 12150 | 10042 | 1352 | 756 |
| **TOTAL** | **157401** | **109243** | **17592** | **30566** |

**Table 1:** *Quality metrics for phenotype data* Stated for each drug is the total number of MIC measurements stratified into "high" quality (at least two MIC measurement methods agree), "medium" quality (either Vizion and AMyGDA disagree, or the scientist recorded a MIC measurement using Vizion but did not store the plate picture) or "low" quality (all three MIC measurements methods disagree) phenotype classifications as described in Methods. Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, LEV = levofloxacin, MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline, CFZ = clofazimine, DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.

***Resistance classification and distribution***

Unsurprisingly, given its size and bias toward the collection of resistant isolates, resistance to each of the 13 drugs is represented within the compendium (Fig. 3a). The drugs with the highest percentage of resistant isolates are the first line drugs isoniazid and rifampicin (49.0% and 38.7% respectively). Of the second line drugs, levofloxacin had the highest proportion of resistant isolates in the dataset (17.6%) and amikacin the lowest (7.3%). A low proportion of isolates were resistant to the NRDs, bedaquiline (0.9%), clofazimine (4.4%), delamanid (1.6%) and linezolid (1.3%).

Of the 12,289 isolates with matched phenotypic and genotypic data, 6,814

(55.4%) were resistant to at least one drug (Fig. 3b).  For the purpose of describing

five broader resistance categories present in the dataset, we assumed that all MICs

that could not be read had susceptible phenotypes. These five resistance categories

comprise: isoniazid and rifampicin susceptible with resistance to another

antitubercular drug, isoniazid resistant but rifampicin susceptible, RR/MDR, pre-XDR

(RR/MDR + fluoroquinolone resistance), and XDR (RR/MDR + fluoroquinolone

resistance + resistance to a group A agent: bedaquiline or linezolid)). Consequently,

the calculated prevalence of MDR, XDR *etc*. in the dataset (Fig. 3) are likely under-

estimates. Of the isolates resistant to one or more drugs, 22.8% were resistant to

isoniazid and not rifampicin, 68.8% were either RR or MDR and 8.4% were resistant

to at least one antitubercular drug, but *not* isoniazid or rifampicin (Fig. 3b). Of the

RR/MDR isolates, 38.8% were pre-XDR and 3.0% were XDR. Two of the XDR isolates

returned a resistant phenotype to all 13 of the drugs assayed (Table S5) and therefore

could be reasonably described as totally drug resistant (TDR). One such isolate

belonged to L4 and was contributed by South Africa, and the other belonged to L2 with

an unknown country of origin contributed by Sweden.

The proportion of drug susceptible isolates collected differed between

countries, as not all laboratories oversampled for resistance (Fig. 3c). In countries that

contributed more than 100 resistant isolates, each of the broad phenotypic resistance

categories in Fig. 3b were seen except for Peru, Vietnam and Nepal which did not

contribute any XDR isolates (Fig. 3c).  Vietnam and Brazil sampled a high proportion

of non-MDR/RR resistant phenotypes; 73.9% and 55.1% of resistant isolates

contributed by these countries, respectively, were neither MDR nor RR. For Nepal and

India, an especially high proportion of the MDR/RR isolates contributed were

fluoroquinolone resistant (92.9% and 69.8% respectively), which has been previously

observed for this geographical region (32,33).

Of the 6814 resistant isolates, 256 were from L1, 2886 were from L2, 625 were

from L3 and 3043 were from L4. All five broader categories of resistance were

represented in the four major *M. tuberculosis* lineages (Fig. 3d). We note that the

relative proportions of resistance categories will have been influenced by the different

local sampling approaches since lineage distributions are typically geographically

distinct (Fig. S2). Bearing this in mind, we observe that in the compendium, L3 isolates

contained the most MDR/RR isolates as a proportion of resistant isolates (77.6%), L2

isolates contained the most pre-XDR isolates as a proportion of MDR/RR isolates

(54.2%) and L2 contained the most XDR isolates as a proportion of MDR/RR isolates

(4.7%).

**Figure 3:** ***Drug phenotype data for the CRyPTIC compendium a)*** Frequency of resistance to each of 13 drugs in the CRyPTIC dataset. The total number of isolates with a binary phenotype (of any quality) for the corresponding drug is presented in Table 1. **b)** Phenotypes of 12,289 CRyPTIC isolates with a binary phenotype for at least one drug. **c)** Geographical distribution of phenotypes of 12,289 CRyPTIC isolates. Intensity of blue shows the percentage of isolates contributed that were categorised as susceptible to all 13 drugs. Donut plots show the proportions of resistant phenotypes identified in (b) for countries contributing >=100 isolates with drug resistance. **d)** Proportions of resistance phenotypes in the 4 major *M. tuberculosis* lineages. N is the number of isolates of the lineage called resistant to at least one of the 13 drugs. Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, LEV = levofloxacin, MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline, CFZ = clofazimine, DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.

***Co-occurrence of drug resistance amongst the CRyPTIC isolates***

As we measured MICs to 13 drugs in parallel, we can ask whether, and how

often, co-occurrence of drug resistance occurs amongst the isolates. We found

isolates with all possible two-drug resistant combinations in this dataset (Fig. 4a, Table

S6).  With the exception of correlations between drugs in the same class (rifabutin vs

rifampicin, moxifloxacin vs levofloxacin), Isoniazid resistance was the most strongly

associated with resistance to each of the other drugs. Resistance to any of the drugs

was also strongly associated with resistance to rifampicin. Of the second line drugs,

levofloxacin and moxifloxacin were more commonly seen as a second resistant

phenotype than the injectable drugs kanamycin and amikacin.

Resistance to both drugs in the aminoglycoside class was common in the

dataset; 90.4% of amikacin resistant isolates were also resistant to kanamycin

although significantly fewer kanamycin resistant isolates were resistant to amikacin

(72.0%, $p$ <0.00001) (Fig. 4a). In a similar fashion, a smaller proportion of rifampicin

resistant isolates were resistant to rifabutin than rifabutin resistant isolates that were

resistant to rifampicin (91.3%, 96.8%, $p$ <0.00001) while a smaller proportion of

levofloxacin resistant isolates were resistant to moxifloxacin than moxifloxacin

resistant isolates that were resistant to levofloxacin (78.5%, 97.6%, $p$ <0.00001).

Differences in drugs of the same class are also well documented by *in vitro*

studies (34–36).

Isolates resistant to the NRDs bedaquiline, clofazimine, delamanid and linezolid

were most likely to also be resistant to isoniazid, followed by rifampicin and rifabutin.

The NRDs were less commonly seen as a second resistance phenotype and the

smallest proportional resistance combinations involved the NRDs (*e.g.* 1.5% of

isoniazid resistant isolates were bedaquiline resistant). Within the NRDs however, co-

occurrence of resistance was proportionally higher; bedaquiline, linezolid and

delamanid resistance was commonly seen with clofazimine resistance (52.4%, 34.2%

and 26.3% of isolates having co-resistance with clofazimine respectively).

***Additional antibiotic resistance in isolates with non-MDR or MDR phenotypic***

***backgrounds***

To further investigate drug resistance patterns amongst the isolates, we

examined in more detail the correlation structure of phenotypes by conditioning on

different resistance backgrounds including isoniazid and rifampicin susceptible,

isoniazid resistant and rifampicin susceptible, rifampicin resistant and isoniazid

susceptible, MDR, pre-XDR and XDR (Fig. 4b-f). We found that a greater proportion

of isolates that were susceptible to isoniazid and rifampicin were resistant to the

second line drugs levofloxacin (24.1%), kanamycin (18.1%), moxifloxacin (13.7%),

and amikacin (8.9%) than the first line drug ethambutol (3.8%) (Fig. 4b). The proportion

of isolates resistant to clofazimine or levofloxacin was particularly high (32.9% and

24.1%, respectively), and more isolates were resistant to these two drugs than

ethambutol in an isoniazid resistant and rifampicin susceptible background but not in

MDR/RR isolates (Fig. 4c-f).

MDR/RR isolates were most commonly resistant (excluding rifabutin) to the first

line drug ethambutol (46.3%), closely followed by levofloxacin (41.4%). As expected,

the proportion of fluoroquinolone resistance was higher in MDR/RR isolates than non-

MDR isolates (37) and we found a greater proportion of isolates were resistant to

levofloxacin than moxifloxacin, a pattern seen in all other backgrounds (Fig. 4c-f). For

the aminoglycosides, a greater percentage of MDR/RR isolates were kanamycin

resistant (21.8%) than amikacin resistant (18.1%), a trend seen in all other

backgrounds.

For isolates with an XDR phenotype, a higher proportion were resistant to

linezolid than bedaquiline (66.7% compared to 44.6%) and 11.3% of XDR isolates

were resistant to both bedaquiline and linezolid (Fig. 4f). XDR isolates were also

resistant to the other NRDs, clofazimine (41.3%) and delamanid (18.8%). In non-XDR

backgrounds the most common NRD resistance seen was clofazimine (Fig. 4b-e).

**Figure 4:** *Co-occurrence of resistance to one drug conditional on resistance to another drug, or to resistance background,* **a)** The heatmap shows the probability of an isolate being resistant to Drug 2 if it is resistant to Drug 1, percentages are given in Table S4. **(b-g)** Percentage of isolates that are resistant to another of the 13 drugs in a background of **(b)** isoniazid susceptible + rifampicin susceptible (but resistant to at least one other antitubercular drug), **(c)** isoniazid resistant + rifampicin susceptible, **(d)** MDR/RR **(e)** Pre-XDR, **(f)** XDR. Only samples with definite phenotypes for RIF in MDR backgrounds and RIF and INH in non-MDR backgrounds and the additional drug are included. Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, LEV = levofloxacin, MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline, CFZ = clofazimine, DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.

### Genetic-based predictions of resistance

To establish a baseline measure of how well resistance and susceptibility could be predicted based on the state of the art prior to the CRyPTIC project, we compared genetic-based predictions of susceptibility and resistance to the binary phenotypes

derived from MICs for eight drugs and for all isolates in this compendium (Table 2).

Since these data were not collected prospectively or randomly, but indeed are

enriched for resistance, the calculated error rates are not representative of how well

such a method would perform in routine clinical use. The results were broadly in line

with prior measurements on a smaller (independent) set (23). The hybrid catalogue

does not make predictions for rifabutin, linezolid, bedaquiline, delamanid or

clofazimine; indeed, this is one of the main aims of the consortium and new catalogues

published by CRyPTIC and the WHO will begin to address this shortcoming (Fig. 5)

(26).

505
506
Table 3 shows the top mutations found amongst isolates phenotypically

resistant to first- and second- line drugs. As expected, *rpoB* S450L was the most

prevalent mutation associated with rifampicin resistance and *katG* S315T was the

most prevalent mutation associated with isoniazid resistance. Mutations in *gyrA*

dominate amongst fluoroquinolone resistant isolates; D94G and A90V are the two

most frequently occurring mutations for levofloxacin and moxifloxacin.

|      | TP   | FP  | TN    | FN  | VME   | ME    | PPV   | NPV   |
|------|------|-----|-------|-----|-------|-------|-------|-------|
| INH  | 5493 | 142 | 5622  | 224 | 0.039 | 0.025 | 0.961 | 0.975 |
| RIF  | 4535 | 435 | 6669  | 107 | 0.023 | 0.061 | 0.977 | 0.939 |
| EMB  | 1919 | 513 | 6702  | 111 | 0.055 | 0.071 | 0.945 | 0.929 |
| LEV  | 1689 | 255 | 8104  | 184 | 0.098 | 0.031 | 0.902 | 0.969 |
| MXF  | 1358 | 504 | 9022  | 160 | 0.105 | 0.053 | 0.895 | 0.947 |
| AMI  | 632  | 84  | 10117 | 163 | 0.205 | 0.008 | 0.795 | 0.992 |
| KAN  | 735  | 124 | 9043  | 197 | 0.211 | 0.014 | 0.789 | 0.986 |
| ETH  | 971  | 114 | 9183  | 511 | 0.345 | 0.012 | 0.655 | 0.988 |

**Table 2: *Predicting phenotypic resistance using genetics***
Statistics on how well phenotypic resistance could be predicted using a standard resistance catalogue that predates the CRyPTIC project. TP: the number of phenotypically resistant samples are that correctly identified as resistant ("true positives"); FP, the number of phenotypically susceptible samples that are falsely identified as resistant ("false positives"); TN, the number of phenotypically susceptible samples that are correctly identified as susceptible ("true negatives"); FN, the number of phenotypically resistant samples that are incorrectly identified as susceptible ("false negative"); VME, very major error rate (false-negative rate), 0-1; ME, major error rate (false-positive rate), 0-1; PPV, positive predictive value, 0-1; NPV, negative predictive value, 0-1. Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, AMI = amikacin, KAN = kanamycin, LEV = levofloxacin, MXF = moxifloxacin, ETH = ethionamide.

| | GENE | VARIANT | *n* | % |
|---|---|---|---|---|
| **Rifampicin** | *rpoB* | S450L | 2914 | 62.2 |
| | *rpoB* | D435V | 506 | 10.8 |
| | *rpoB* | H445D | 202 | 4.3 |
| | *rpoB* | H445Y | 147 | 3.1 |
| | *rpoB* | D435Y | 112 | 2.4 |
| **Isoniazid** | *katG* | S315T | 3668 | 70.0 |
| | *fabG1* | c-15t | 829 | 15.8 |
| | *fabG1* | g-17t | 176 | 3.4 |
| | *fabG1* | t-8c | 154 | 2.9 |
| | *inhA* | I194T | 56 | 1.1 |
| **Ethambutol** | *embB* | M306V | 1131 | 50.0 |
| | *embB* | M306I | 1001 | 44.3 |
| | *embB* | Q497R | 449 | 19.9 |
| | *embB* | G406A | 164 | 7.2 |
| | *embB* | G406D | 105 | 4.6 |
| **Kanamycin** | *rrs* | a1401g | 660 | 58.9 |
| | *eis* | c-14t | 70 | 6.2 |
| | *eis* | g-10a | 53 | 4.7 |
| **Amikacin** | *rrs* | a1401g | 660 | 74.7 |
| | *rrs* | g1484t | 7 | 0.8 |
| **Levofloxacin** | *gyrA* | D94G | 783 | 36.5 |
| | *gyrA* | A90V | 487 | 22.7 |
| | *gyrA* | D94N | 157 | 7.3 |
| | *gyrA* | D94A | 133 | 6.2 |
| | *gyrA* | S91P | 92 | 4.3 |
| **Moxifloxacin** | *gyrA* | D94G | 783 | 45.4 |
| | *gyrA* | A90V | 487 | 28.2 |
| | *gyrA* | D94N | 157 | 9.1 |
| | *gyrA* | D94A | 133 | 7.7 |
| | *gyrA* | D94Y | 70 | 4.1 |
| **Ethionamide** | *fabG1* | c-15t | 829 | 48.0 |
| | *fabG1* | L203L | 124 | 7.2 |

**Table 3: *The top mutations associated with phenotypic drug resistance***
Depicted is a survey of the resistance-associated mutations present in CRyPTIC isolates (7,25).
"VARIANT": non-synonymous amino acid mutations are denoted by upper case letters while
nucleotide substitutions for non-coding sequences are denoted by lower case letters. Negative
numbers denote substitutions in promoter regions; "GENE": genic region of interest in which
"Variant" can be found; "*n*": number of phenotypically resistant isolates with "VARIANT"; "%":
percentage of total phenotypically resistant isolates with "VARIANT".

### *Resistance to new and re-purposed drugs*

As previously stated, relatively few isolates are resistant to the NRDs, bedaquiline ($n$ = 109), clofazimine ($n$ = 525), delamanid ($n$ = 186) and linezolid ($n$ = 156). South Africa contributed the greatest number of isolates resistant to bedaquiline, clofazimine and linezolid (Fig. 5a), while China and India contributed the most isolates resistant to delamanid. Since the collection protocol differed between laboratories it is not possible to infer any differences in the relative prevalence of resistance to the NRDs in these countries. The results of a survey of all non-synonymous mutations in genes known or suspected to be involved in resistance to these four drugs (*e.g. rv0678, mmpL5, pepQ, ddn, rplC, rrl etc.*) are depicted in Fig. 5b-e (38–42). Mutations known to be associated with sensitivity were ignored, along with mutations that occurred at a frequency of ≥ 5% amongst all isolates (as 0.05% of total isolates are resistant to the NRDs). In contrast to first- and second-line drugs, there are no mutations within a single gene/small group of genes that can fully explain resistance to any NRD, and indeed no single gene (if there were, they would be visible as complete rings around the phylotrees in figure 5). Note that the role of most of these mutations in resistance remains undetermined.

**Figure 5:** ***Resistance to bedaquiline, clofazimine, delamanid and linezolid amongst M. tuberculosis CRyPTIC isolates* a)** The prevalence (within these data) of resistance to bedaquiline (BDQ), clofazimine (CFZ), delamanid (DLM) and linezolid (LZD) per country or origin or collection site. Phylotrees are shown for isolates phenotypically resistant to b) BDQ, c) CFZ, d) DLM and e) LZD. Tip point colours denote lineage. Each outer track represents

a gene thought to be associated with resistance and coloured blocks denote the presence of a non-synonymous mutation in the relevant gene for a given isolate. Mutations in these genes that are either associated with sensitivity or present in >5% of the collection of isolates as a whole were ignored.

### *Case study on rifampicin mono-resistance*

Around 1% of TB cases are rifampicin mono-resistant (RMR) and the frequency is increasing (1,43). The WHO does not recommend isoniazid for RMR treatment, despite it being effective; this is likely due to the reliance on assays such as Xpert® MTB/RIF which cannot distinguish between RMR and MDR. Use of isoniazid could improve treatment outcomes for RMR patients which are currently similar to that of MDR TB, including a higher risk of death compared to drug susceptible infections (44,45). Due to its low natural prevalence, RMR has been poorly studied to date but increasingly large clinical TB datasets, such as the one presented here, make its study now feasible.

For this case study, we defined RMR as any isolate that is rifampicin resistant and isoniazid susceptible, and discounted isolates with no definite phenotype for either drug. Of the 4,655 rifampicin resistant isolates in the compendium that also had a phenotype for isoniazid, 302 (6.5%) were RMR. These isolates were contributed by 12 different countries, and we found South African and Nigerian contributions had a significantly higher proportion of RMR isolates than that of the total dataset at 17.5% ($p$ <0.00001) and 27.3% ($p$ = 0.00534) respectively (Fig. 6a) compared with 6.5% for the total dataset. We note that these proportions are influenced by sampling strategies but the higher contribution of RMR isolates from South Africa is consistent with previous studies (43).

***Rifampicin mono-resistance is incorrectly predicted by current diagnostics***

A widely used, WHO-endorsed diagnostic tool, the Xpert® MTB/RIF assay,

uses a proxy whereby *any* SNP detected in the "rifampicin-resistance determining

region" (RRDR) of *rpoB* results in a prediction of MDR. However, the suitability of the

proxy is dependent upon prevalence of RMR in the population (43). We tested the

reliability of this on the 4,655 rifampicin resistant isolates in our dataset that had a

phenotype for isoniazid (Fig. 6b).

Of these isolates, 4,353 (93.5%) were MDR and 302 (6.5%) were RMR. 187 of

the MDR isolates had no RRDR mutation and therefore 4.0% of isolates in this study

would be predicted as false negative MDR by the Xpert® MTB/RIF assay. 276 of the

RMR isolates had a mutation in the RRDR of *rpoB* and so the Xpert® MTB/RIF assay

proxy would incorrectly predict 5.9% of the rifampicin resistant isolates as false

positive MDR cases. However, overall, the Xpert® MTB/RIF assay proxy correctly

predicts 89.5% of the rifampicin resistant isolates as MDR and 0.6% of the isolates as

non-MDR in this dataset, which suggests it is a reasonably successful diagnostic tool

with > 90% accuracy for MDR classification of rifampicin resistant isolates. As our

dataset is oversampled for resistance, it likely contains a higher prevalence of RMR

than the global average and hence the Xpert® MTB/RIF assay is likely to perform

better on more representative data. However, the analysis shows how the increasing

global levels of RMR TB cases could increase the number of false positive MDR

diagnoses by the Xpert® MTB/RIF assay, denying isoniazid treatment to a greater

number of patients who would then be moved on to less effective drugs.

### There are genetic differences between rifampicin mono-resistant and multidrug resistant isolates

We have analysed our matched phenotypic and genotypic data to examine whether there were any differences in the genetic determinants of rifampicin resistance between RMR and MDR isolates as was seen in a recent study of South African isolates (46). The proportion of RMR isolates with no *rpoB* mutation (5.3%, Fig. 6b) was significantly higher than that of MDR isolates (1.8%, $p$ <0.00001). This suggests that non-target-mediated resistance mechanisms, such as upregulation of rifampicin specific efflux pumps, could be more influential in providing protection against rifampicin in RMR isolates than in MDR isolates.

The majority of RMR and MDR isolates contained one or more SNPs in *rpoB*, with the most having at least one mutation in the RRDR. To date, several non-synonymous RRDR mutations have been found in RMR *M. tuberculosis* isolates, including the resistance conferring mutations S450L, H445D and D435Y, which are also seen in MDR isolates (47,48). For both RMR and MDR isolates in this dataset, the most common *rpoB* RRDR mutation seen was S450L (63.6% and 41.1% of isolates respectively, Fig. 6c). Five mutations were present in RMR isolates that were not seen in MDR isolates: S428G, S441A, S441V, S450M and S450Q, however these were seen at low prevalence (< 2%) of RMR isolates. We found more RMR isolates had His445 mutated than MDR isolates (27.8% of RMR and 9.5% of MDR, $p$ <0.00001), and mutations at Ser450 and Asp435 were more prevalent in MDR isolates than RMR isolates (43.7% of RMR and 65.8% of MDR ($p$ <0.00001), and 9.3% of RMR and 15.5% of MDR ($p$ = 0.00328) respectively).

In RMR isolates we observed 27 different *rpoB* mutations that fall outside the

RRDR; 11 were found in RMR but not MDR isolates and all were seen at < 2%

prevalence (Fig. S6). The most common non-RRDR mutation in both MDR and RMR

isolates was a cytosine to thymine mutation 61 bases upstream of the *rpoB* start codon

(10.1% and 8.6% of isolates respectively). The resistance conferring mutations, *rpoB*

I491F, V695L and V170F, were seen at low proportions (< 2% of isolates) with no

significant difference between MDR and RMR isolates.

**Figure 6: *Rifampicin mono resistance* a)** Percentage of rifampicin resistant isolates that
are rifampicin mono-resistant (RMR) by country of isolate origin. * indicates RMR proportions that

were significantly different from that of the total dataset using a two tailed z-test with 95% confidence. **b)** MDR predictions for rifampicin resistant isolates made using the Xpert® MTB/RIF assay proxy. N is the total number of rifampicin resistant isolates. The inner ring shows the proportion of rifampicin resistant isolates that are MDR and RMR. The middle ring represents the proportions of MDR and RMR isolates that have a SNP (synonymous or non-synonymous) in the RRDR of *rpoB* (RRDR), no RRDR SNP but a SNP elsewhere in the *rpoB* gene (rpoB) and no *rpoB* mutations (none). The outer ring shows the expected true positive (TP), true negative (TN), false positive (FP) and false negative (FN) MDR predictions of Xpert® MTB/RIF assay, based on the SNPs present in the rifampicin resistant isolates. **c)** Non-synonymous mutations found in the RRDR of *rpoB* in RMR isolates and MDR isolates. Presence of a coloured spot indicates that the mutation was found in RMR/MDR isolates and spot size corresponds to the proportion of RMR or MDR isolates carrying that mutation.

## Discussion

This compendium of *M. tuberculosis* clinical isolates is the result of an extensive global effort by the CRyPTIC consortium to better map the genetic variation associated with drug resistance. Through its sheer size and by oversampling for resistance, the compendium gives an unparalleled view of resistance and resistance patterns amongst the panel of 13 antitubercular compounds studied. This study serves to summarise the data within the compendium and to highlight the existence of the open-access resource to the wider community to help better inform future treatment guidelines and steer the development of improved diagnostics.

Starting with first-line drugs, molecular based diagnostic assays have vastly improved the detection of and the speed at which we find drug resistant TB cases, resulting in improved quality of care for patients. However, relying solely on these diagnostic methods has several drawbacks. Aside from the Xpert® MTB/RIF assay potentially increasing false positive MDR diagnoses as discussed earlier in the RMR

case study, the assay assumes isoniazid resistance upon detection of rifampicin

resistance. Thus, less is known about the prevalence of mono isoniazid resistance or

'true' cases of MDR (confirmed rifampicin *and* isoniazid resistance) (1) and with large

datasets such as this compendium, we can further investigate these important and

rarer clinical phenotypes (like that of RMR in our case study). Another example of a

rarer phenotype is that of isoniazid-resistant and rifampicin-susceptible (Hr-TB)

isolates; a greater number of these were contributed by CRyPTIC countries than RMR

isolates *(n* = 1470 versus *n* = 302)*,* a pattern also recently observed in a global

prevalence study (49). A modified 6-month treatment regimen is now recommended

for Hr-TB (rifampicin, ethambutol, levofloxacin and pyrazinamide), and as a result of

inadequate diagnosis many of the 1.4 million global Hr-TB estimated cases would

have received inadequate and unnecessarily longer treatment regimens (1,42).

Encouragingly, CRyPTIC isolates with a Hr-TB background exhibited relatively low

levels of resistance to other antitubercular drugs, including those in the augmented

regimen (Fig. 4c). However, without appropriate tools to assess and survey this, we

will continue to misdiagnose and infectively treat these clinical cases. In 2018,

CRyPTIC and the 100,000 Genomes project demonstrated that genotypic prediction

from WGS correlates well with culture-based phenotype for first-line drugs, which is

reflected in our summary of the genetic catalogue applied to this dataset (Table 3) (7).

While predictions can be made to a high level of sensitivity and specificity, there is still

more to learn, as exemplified by the isolates in the compendium that despite being

resistant to rifampicin and isoniazid could not be described genetically (Table 2). This

shortfall, along with the limitations of molecular based diagnostic assays, highlights

the need for continual genetic surveillance and shines a favourable light on a WGS-

led approach.

A strength of this compendium lies with the data collated for second-line drugs.

A greater proportion of drug resistant isolates had additional resistance to

fluoroquinolones than second line injectable drugs (Fig. 4a). This could be due to more

widespread use of fluoroquinolones as well as their ease of administration and hence

them being recommended over injectables for longer MDR treatment regimens (1).

Concerningly, we found that resistance to levofloxacin and moxifloxacin, and

kanamycin and amikacin, were more common than resistance to the mycobacterial

specific drug ethambutol in an isoniazid and rifampicin susceptible background (Fig.

4b), suggesting a level of pre-existing resistance to second-line drugs. This concurs

with a systematic review that found patients previously prescribed fluoroquinolones

were three times more likely to have fluoroquinolone resistant TB (50). Careful

stewardship of fluoroquinolones, both in TB and other infectious diseases, will be

paramount for the success of treatment regimens. Despite variability in sample

collection, we observed high proportions of fluoroquinolone resistant MDR/RR isolates

from some countries and therefore suggest that MDR treatment regimens could be

improved by optimisation on a geographic basis.

Further treatment improvement could also be made by the selection of

appropriate drugs from each class. For example, the WHO recommends switching

from kanamycin to amikacin when treating MDR TB patients (51) and the compendium

supports this recommendation as we saw more resistance to kanamycin than amikacin

in all phenotypic backgrounds. For fluoroquinolones, more isolates were resistant to

levofloxacin than moxifloxacin in all phenotypic backgrounds suggesting moxifloxacin

may by the most appropriate fluoroquinolone to recommend, although we note this

conclusion is critically dependent on the validity of the cutoff, here an ECOFF, used to

infer resistance. However, the amenability of drugs to catalogue-based genetic

diagnostics is also an important consideration, and our data suggest levofloxacin

resistance could be predicted more reliably than moxifloxacin, with fewer false

positives predicted (Table 2). Testing for fluoroquinolone resistance using molecular

diagnostic tests remains limited. Global data from the past 15 years suggests that the

proportion of MDR/RR TB cases resistant to fluoroquinolones sits at around 20%, with

these cases primarily found in regions of high MDR-TB burden (1). While recently

approved tools, such as the Cepheid Xpert® MTB/XDR cartridge, will permit both

isoniazid and fluoroquinolone testing to be increased, the same pitfalls are to be

encountered regarding targeted diagnostic assays (52). In contrast, the genetic survey

in this study demonstrates the potential of WGS for genetic prediction of resistance to

second-line drugs and studies within the consortium to investigate this are underway.

The CRyPTIC compendium has facilitated the first global survey of resistance

to NRDs. Reassuringly, prevalence of resistance to the NRDs was substantially lower

than for first- and second-line agents in the dataset (Fig. 3a), and resistance to the

new drugs bedaquiline and delamanid was less common than the repurposed drugs

clofazimine and linezolid in an MDR/RR background (Fig. 4c). However, the presence

of higher levels of delamanid and clofazimine resistance than ethambutol resistance

in the isoniazid and rifampicin susceptible background does suggest some pre-existing

propensity towards NRD resistance (Fig. 4b).

Co-resistance between NRDs was seen in isolates in the compendium, the

most common being isolates resistant to both bedaquiline and clofazimine. This link is

well documented and has been attributed to shared resistance mechanisms such as

non-synonymous mutations in *Rv0678* which were found in both clofazimine and

bedaquiline resistant isolates in the compendium (42) (Fig. 5b,c). Increased

clofazimine use could further increase the prevalence of *M. tuberculosis* isolates with

clofazimine and bedaquiline co-resistance, limiting MDR treatment options including

using bedaquiline as the backbone of a shorter MDR regimen (53). Therefore,

proposed usage of clofazimine for other infectious diseases should be carefully

considered.

The WHO recommends against the use of bedaquiline and delamanid in

combination to prevent the development of co-resistance, which could occur relatively

quickly (54); the rate of spontaneous evolution of delamanid resistance *in vitro* has

been shown to be comparable to that of isoniazid, and likewise bedaquiline resistance

arises at a comparable rate to rifampicin resistance (55). In this compendium, 12.9%

of bedaquiline resistant isolates were resistant to delamanid and 7.1% of delamanid

resistant isolates were resistant to bedaquiline. Several scenarios could account for

this, including the presence of shared resistance mechanisms. For example, as

bedaquiline targets energy metabolism within the cell, changes to cope with

energy/nutrient imbalances upon the acquisition of resistance-associated ATPase

pump mutations may result in cross resistance to delamanid in a yet unknown or

unexplored mechanism (12).  it is imperative that genetic determinants of resistance

are fully explored for the NRDs, as these are our current treatments of last resort, with

special attention given to those mechanisms that could be shared with other agents.

In the meantime, careful stewardship and phenotypic and genotypic surveillance of

the NRDs should be implemented, including linezolid and clofazimine which are now

group A and B drugs respectively for MDR treatment (1).

Several research avenues are being actively explored by the CRyPTIC

consortium that make further use of this compendium, including: *i)* relating genetic

mutations to quantitative changes in the minimum inhibitory concentrations of different

drugs (12); *ii)* genome-wide association studies (14); *iii)* training machine learning

models that can predict resistance (13); *iv)* exploration of the genetic determinants of

resistance to second line and NRDs (15). Collectively these studies share the same

aim of facilitating the implementation of WGS-directed resistance prediction in the

clinic. Finally, we urge other researchers to explore and analyse this large dataset of

*M. tuberculosis* clinical isolates and hope it will lead to a wave of new and inciteful

studies that will positively serve the TB community for years to come.

**CRyPTIC Consortium Members**

Alice Brankin[5,*,**] and Kerri M Malone[23,*,**], Ivan Barilar[1], Simone Battaglia[2], Emanuele
Borroni[2], Angela Pires Brandao[3,4], Andrea Maurizio Cabibbe[2], Joshua Carter[6], Darren Chetty[7],
Daniela Maria Cirillo[2], Pauline Claxton[8], David A Clifton[5], Ted Cohen[9], Jorge Coronel[10], Derrick
W Crook[5], Viola Dreyer[1], Sarah G Earle[5], Vincent Escuyer[11], Lucilaine Ferrazoli[4], George Fu
Gao[12], Jennifer Gardy[13], Saheer Gharbia[14], Kelen Teixeira Ghisi[4], Arash Ghodousi[2,15], Ana Lúiza
Gibertoni Cruz[5], Louis Grandjean[16], Clara Grazian[17], Ramona Groenheit[18], Jennifer L
Guthrie[19,20], Wencong He[12], Harald Hoffmann[21,22], Sarah J Hoosdally[5], Martin Hunt[23,5], Nazir
Ahmed Ismail[24], Lisa Jarrett[25], Lavania Joseph[24], Ruwen Jou[26], Priti Kambli[27], Rukhsar Khot[27],
Jeff Knaggs[23,5], Anastasia Koch[28], Donna Kohlerschmidt[11], Samaneh Kouchaki[5,29], Alexander S
Lachapelle[5], Ajit Lalvani[30], Simon Grandjean Lapierre[31], Ian F Laurenson[8], Brice Letcher[23],
Wan-Hsuan Lin[26], Chunfa Liu[12], Dongxin Liu[12], Ayan Mandal[32], Mikael Mansjo[18], Daniela
Matias[25], Graeme Meintjes[28], Flávia de Freitas Mendes[4], Matthias Merker[1], Marina Mihalic[22],

James Millard[7], Paolo Miotto[2], Nerges Mistry[32], David Moore[33,10], Kimberlee A Musser[11],

Dumisani Ngcamu[24], Hoang Ngoc Nhung[34], Stefan Niemann[1,35], Kayzad Soli Nilgiriwala[32],

Camus Nimmo[16], Max O'Donnell[36], Nana Okozi[24], Rosangela Siqueira Oliveira[4], Shaheed Vally

Omar[24], Nicholas Paton[37], Timothy EA Peto[5], Juliana Maira Watanabe Pinhata[4], Sara Plesnik[22],

Zully M Puyen[38], Marie Sylvianne Rabodoarivelo[39], Niaina Rakotosamimanana[39], Paola MV

Rancoita[15], Priti Rathod[25], Esther Robinson[25], Gillian Rodger[5], Camilla Rodrigues[27], Timothy C

Rodwell[40,41], Aysha Roohi[5], David Santos-Lazaro[38], Sanchi Shah[32], Thomas Andreas Kohl[1],

Grace Smith[25,14], Walter Solano[10], Andrea Spitaleri[2,15], Philip Supply[42], Adrie JC Steyn[7],

Utkarsha Surve[27], Sabira Tahseen[43], Nguyen Thuy Thuong Thuong[34], Guy Thwaites[34,5],

Katharina Todt[22], Alberto Trovato[2], Christian Utpatel[1], Annelies Van Rie[44], Srinivasan Vijay[45],

Timothy M Walker[5,34], A Sarah Walker[5], Robin Warren[46], Jim Werngren[18], Maria Wijkander[18],

Robert J Wilkinson[47,48,30], Daniel J Wilson[5], Penelope Wintringer[23], Yu-Xin Xiao[26], Yang Yang[5],

Zhao Yanlin[12], Shen-Yuan Yao[24], Baoli Zhu[49], Philip W Fowler[5], Zamin Iqbal[23][**]

[*]equal contribution authors

[**]co-corresponding authors

## Institutions

[1]Research Center Borstel, Borstel, Germany

[2]IRCCS San Raffaele Scientific Institute, Milan, Italy

[3]Oswaldo Cruz Foundation, Rio de Janeiro, Brazil

[4]Institute Adolfo Lutz, Sào Paulo, Brazil

[5]University of Oxford, Oxford, UK

[6]Stanford University School of Medicine, Stanford, USA

[7]Africa Health Research Institute, Durban, South Africa

[8]Scottish Mycobacteria Reference Laboratory, Edinburgh, UK

[9]Yale School of Public Health, Yale, USA

[10]Universidad Peruana Cayetano Heredia, Lima, Perú

[11]Wadsworth Center, New York State Department of Health, Albany, USA

[12]Chinese Center for Disease Control and Prevention, Beijing, China

[13]Bill & Melinda Gates Foundation, Seattle, USA

[14]UK Health Security Agency, London, UK

[15]Vita-Salute San Raffaele University, Milan, Italy

[16]University College London, London, UK

[17]University of New South Wales, Sydney, Australia

[18]Public Health Agency of Sweden, Solna, Sweden

[19]The University of British Columbia, Vancouver, Canada

[20]Public Health Ontario, Toronto, Canada

[21]SYNLAB Gauting, Munich, Germany

[22]Institute of Microbiology and Laboratory Medicine, IMLred, WHO-SRL Gauting, Germany

[23]EMBL-EBI, Hinxton, UK

[24]National Institute for Communicable Diseases, Johannesburg, South Africa

[25]UK Health Security Agency, Birmingham, UK

[26]Taiwan Centers for Disease Control, Taipei, Taiwan

[27]Hinduja Hospital, Mumbai, India

[28]University of Cape Town, Cape Town, South Africa

[29]University of Surrey, Guildford, UK

[30]Imperial College, London, UK

[31]Université de Montréal, Canada

[32]The Foundation for Medical Research, Mumbai, India

[33]London School of Hygiene and Tropical Medicine, London, UK

[34]Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

[35]German Center for Infection Research (DZIF), Hamburg-Lübeck-Borstel-Riems, Germany

[36]Colombia University Irving Medical Center, New York, USA

[37]National University of Singapore, Singapore

[38]Instituto Nacional de Salud, Lima, Perú

[39]Institut Pasteur de Madagascar, Antananarivo, Madagascar

[40]FIND, Geneva, Switzerland

[41]University of California, San Diego, USA

[42]Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL -

Center for Infection and Immunity of Lille, F-59000 Lille, France

[43]National TB Reference Laboratory, National TB Control Program, Islamabad, Pakistan

[44]University of Antwerp, Antwerp, Belgium

[45]University of Edinburgh, Edinburgh, UK

[46]Stellenbosch University, Cape Town, South Africa

[47]Wellcome Centre for Infectious Diseases Research in Africa, Cape Town, South Africa

[48]Francis Crick Institute, London, UK

[49]Institute of Microbiology, Chinese Academy of Sciences, Beijing, China

**Author contributions**

D.W.C, T.E.A.P, S.H, A.L.G.C, A.W.S, T.M.W, P.W.F, D.M.C designed the CRyPTIC

study and all contributing laboratories collected samples and provided data. MIC data

and genetic information was retrieved and processed by P.W.F, S.H, A.L.G.C, Z.I, M.H

and J.K. A.B and K.M.M designed and performed all analyses for this manuscript. A.B

and K.M.M wrote the manuscript with feedback from CRyPTIC partners.

905
**Acknowledgements**

We thank Faisal Masood Khanzada and Alamdar Hussain Rizvi (NTRL, Islamabad,

Pakistan), Angela Starks and James Posey (Centers for Disease Control and

Prevention, Atlanta, USA), and Juan Carlos Toro and Solomon Ghebremichael (Public

Health Agency of Sweden, Solna, Sweden), Iñaki Comas and Álvaro Chiner-Oms

(Instituto de Biología Integrativa de Sistemas, Valencia, Spain; CIBER en

Epidemiología y Salud Pública, Valencia, Spain; Instituto de Biomedicina de Valencia,

IBV-CSIC, Valencia, Spain).

**Funding bodies**

This work was supported by Wellcome Trust/Newton Fund-MRC Collaborative Award

(200205/Z/15/Z); and Bill & Melinda Gates Foundation Trust (OPP1133541). Oxford

CRyPTIC consortium members are funded/supported by the National Institute for

Health Research (NIHR) Oxford Biomedical Research Centre (BRC), the views

expressed are those of the authors and not necessarily those of the NHS, the NIHR

or the Department of Health, and the National Institute for Health Research (NIHR)

Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial

Resistance, a partnership between Public Health England and the University of

Oxford, the views expressed are those of the authors and not necessarily those of the

NIHR, Public Health England or the Department of Health and Social Care. J.M. is

supported by the Wellcome Trust (203919/Z/16/Z). Z.Y. is supported by the National

Science and Technology Major Project, China Grant No. 2018ZX10103001. K.M.M. is

supported by EMBL's EIPOD3 programme funded by the European Union's Horizon

2020 research and innovation programme under Marie Skłodowska Curie Actions.

T.C.R. is funded in part by funding from Unitaid Grant No. 2019-32-FIND MDR. R.S.O.

is supported by FAPESP Grant No. 17/16082-7. L.F. received financial support from

FAPESP Grant No. 2012/51756-5. B.Z. is supported by the National Natural Science

Foundation of China (81991534) and the Beijing Municipal Science & Technology

Commission (Z201100005520041). N.T.T.T. is supported by the Wellcome Trust

International Intermediate Fellowship (206724/Z/17/Z). G.T. is funded by the

Wellcome Trust. R.W. is supported by the South African Medical Research Council.

J.C. is supported by the Rhodes Trust and Stanford Medical Scientist Training

Program (T32 GM007365). A.L. is supported by the National Institute for Health

Research (NIHR) Health Protection Research Unit in Respiratory Infections at Imperial

College London. S.G.L. is supported by the Fonds de Recherche en Santé du Québec.

C.N. is funded by Wellcome Trust Grant No. 203583/Z/16/Z. A.V.R. is supported by

Research Foundation Flanders (FWO) under Grant No. G0F8316N (FWO Odysseus).

G.M. was supported by the Wellcome Trust (098316, 214321/Z/18/Z, and

203135/Z/16/Z), and the South African Research Chairs Initiative of the Department

of Science and Technology and National Research Foundation (NRF) of South Africa

(Grant No. 64787). The funders had no role in the study design, data collection, data

analysis, data interpretation, or writing of this report. The opinions, findings and

conclusions expressed in this manuscript reflect those of the authors alone. L.G. was

supported by the Wellcome Trust (201470/Z/16/Z), the National Institute of Allergy and

Infectious Diseases of the National Institutes of Health under award number

1R01AI146338, the GOSH Charity (VC0921) and the GOSH/ICH Biomedical

Research Centre (www.nihr.ac.uk). A.B. is funded by the NDM Prize Studentship from

the Oxford Medical Research Council Doctoral Training Partnership and the Nuffield

Department of Clinical Medicine. D.J.W. is supported by a Sir Henry Dale Fellowship

jointly funded by the Wellcome Trust and the Royal Society (Grant No. 101237/Z/13/B)

and by the Robertson Foundation. A.S.W. is an NIHR Senior Investigator. T.M.W. is a

Wellcome Trust Clinical Career Development Fellow (214560/Z/18/Z). A.S.L. is

supported by the Rhodes Trust. R.J.W. receives funding from the Francis Crick

Institute which is supported by Wellcome Trust, (FC0010218), UKRI (FC0010218),

and CRUK (FC0010218). T.C. has received grant funding and salary support from US

NIH, CDC, USAID and Bill and Melinda Gates Foundation. The computational aspects

of this research were supported by the Wellcome Trust Core Award Grant Number

203141/Z/16/Z and the NIHR Oxford BRC. Parts of the work were funded by the

German Center of Infection Research (DZIF). The Scottish Mycobacteria Reference

Laboratory is funded through National Services Scotland. The Wadsworth Center

contributions were supported in part by Cooperative Agreement No. U60OE000103

funded by the Centers for Disease Control and Prevention through the Association of

Public Health Laboratories and NIH/NIAID grant AI-117312. Additional support for

sequencing and analysis was contributed by the Wadsworth Center Applied Genomic

Technologies Core Facility and the Wadsworth Center Bioinformatics Core. SYNLAB

Holding Germany GmbH for its direct and indirect support of research activities in the

Institute of Microbiology and Laboratory Medicine Gauting. N.R. thanks the

Programme National de Lutte contre la Tuberculose de Madagascar.

987
988
989 **Competing Interest**
990
E.R. is employed by Public Health England and holds an honorary contract with

Imperial College London. I.F.L. is Director of the Scottish Mycobacteria Reference

Laboratory. S.N. receives funding from German Center for Infection Research,

Excellenz Cluster Precision Medicine in Chronic Inflammation, Leibniz Science

Campus Evolutionary Medicine of the LUNG (EvoLUNG)tion EXC 2167. P.S. is a

consultant at Genoscreen. T.R. is funded by NIH and DoD and receives salary support

from the non-profit organization FIND. T.R. is a co-founder, board member and

shareholder of Verus Diagnostics Inc, a company that was founded with the intent of

developing diagnostic assays. Verus Diagnostics was not involved in any way with

data collection, analysis or publication of the results. T.R. has not received any

financial support from Verus Diagnostics. UCSD Conflict of Interest office has

reviewed and approved T.R.'s role in Verus Diagnostics Inc. T.R. is a co-inventor of a

provisional patent for a TB diagnostic assay (provisional patent #: 63/048.989). T.R. is

**Supplemental Information**

**S1. Lineages of the *M. tuberculosis* isolates of the compendium.**

Isolates of the ancient Indo-oceanic lineage/L1 ($n$ = 1,150) were mostly contributed by India ($n$ = 676 isolates) and Vietnam ($n$ = 283 isolates). 85% of the L1 Indian isolates belong to sub-lineages 1.1.2 and 1.2.2 while 66% of the Vietnamese isolates are sub-lineage 1.1.1.1. No L1 isolates were contributed by 10 of the 23 countries in this study with only 2 isolates collected in South America (one in each of Brazil and Peru).

There are 5,598 L2 (East Asian) isolates, making it the second largest group in the dataset. L2 was found most prevalent in Asia and Europe with the largest

proportion found in amongst isolates contributed by China ($n$ = 722, 64% of isolates)

and India ($n$ = 1,573, 39% of isolates). Sub-lineages 2.2 and 2.2.7 dominate the L2

isolates ($n$ = 1,421 and 1,249 respectively); 2.2 was found mostly amongst Peruvian

and South African isolates ($n$ = 231 and 161 respectively) apart from those contributed

by the Asian countries of Vietnam ($n$ = 271), China ($n$ = 284) and India ($n$ = 272), while

85% of sub-lineage 2.2.7 isolates were contributed by South Africa ($n$ = 206), Vietnam

($n$ = 164) and India ($n$ = 691).  70% of sub-lineage 2.2.1 originated from South Africa

(10% of isolates found here) and has recently been associated with more favourable

transmission rates (56). Lastly, 86% and 72% of isolates contributed by Kyrgyzstan

and Turkmenistan respectively belong to L2 with sub-lineage 2.2.10 dominating (16/24

and 75/86 isolates for both countries respectively). 2.2.10 has been previously

described as restricted to Central Asia and this is reflected in the compendium (57).

1041

1042      The majority (1184/1850, 65%) of L3 (East African/Indian) isolates were

1043      contributed by India, followed by 19.6% (363/1850) isolates from Pakistan. L3 is

1044      typically under-sampled and under-studied in current databases and biobanks in

1045      comparison to L2 and L4; the L3 isolates in this study are the largest collected to date

1046      in a single study (31).

1047

1048      L4 (Euro-American) is the largest lineage group in the compendium ($n$ = 6,572).

1049      Isolates donated from Peru dominate; 87% of all L4 isolates are Peruvian with 4.1.2.1

1050      and 4.3.3 being the most prevalent sub-lineages (24% and 22% respectively). There

1051      are 34 different L4 sub-lineages in the dataset, making L4 the most diverse in

1052      comparison to the other lineage groupings.

1053

1054    There were no L5 isolates found in the compendium and only 6 L6 isolates were

1055    identified. Animal-restricted pathogenic mycobacterial isolates are also rare in the

1056    compendium; only 16 cases were identified (*n* = 15 *M. bovis* and *n* = 1 *M. caprae*).

1057
1058

1059

**Figure S1:** *Per drug MIC distributions of isolates plated on CRyPTIC designed variations on the Thermo Fischer Sensititre MYCOTB MIC plate; UKMYC5 (a) and UKMYC6 (b).* The solid black line represents the epidemiological resistance cut-off (ECOFF) for each drug as determined by (11). Isolates with an MIC above the cut-off are considered resistant. N denotes the total number of isolates tested on each plate that returned a phenotype for each drug.

| DRUG | ECOFF |
|---|---|
| Isoniazid | 0.1 |
| Rifampicin | 0.5 |
| Rifabutin | 0.12 |
| Ethambutol | 4 |
| Ethionamide | 4 |
| Levofloxacin | 1 |
| Moxifloxacin | 1 |
| Amikacin | 1 |
| Kanamycin | 4 |
| Bedaquiline | 0.25 |
| Clofazimine | 0.25 |
| Delamanid | 0.12 |
| Linezolid | 1 |

**Table S1: Epidemiological cut-off values (***ECOFFs) used to binarize MIC measurements into resistant and susceptible.*** Isolates with an MIC above the cut-off are considered resistant and those at or below the cut-off as susceptible.

1074

**Figure S2:** *Geographical distribution of CRyPTIC M. tuberculosis clinical isolates.* The total number of isolates contributed by each country is depicted, with pie charts representing the proportion of *M. tuberculosis* lineages. Where the origin of an isolate was not known, the collection site identity was assigned (269 isolates in Germany, 17 isolates in India, 6 isolates in Peru, 885 isolates in Italy, 510 isolates in South Africa, 357 isolates in Sweden, 208 isolates in Taiwan, 1 isolate in Brazil and 4 isolates in the UK).

1081

1082

1083

1084

1085

1086

1087

1088

1089

|  | Animal/ other | L1 | L2 | L3 | L4 | L6 | Total |
|---|---|---|---|---|---|---|---|
| **Albania** | 0 | 0 | 0 | 0 | 7 | 0 | 7 |
| **Algeria** | 0 | 0 | 0 | 0 | 25 | 0 | 25 |
| **Brazil** | 1 | 1 | 8 | 0 | 335 | 0 | 345 |
| **Burkina Faso** | 0 | 3 | 1 | 8 | 65 | 1 | 78 |
| **China** | 7 | 2 | 722 | 16 | 374 | 0 | 1121 |
| **Germany** | 1 | 25 | 235 | 61 | 403 | 1 | 726 |
| **India** | 11 | 676 | 1573 | 1184 | 560 | 0 | 4004 |
| **Italy** | 11 | 38 | 192 | 113 | 866 | 4 | 1224 |
| **Japan** | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| **Kyrgyzstan** | 0 | 0 | 25 | 0 | 3 | 0 | 28 |
| **Nepal** | 0 | 5 | 137 | 34 | 21 | 0 | 197 |
| **Nigeria** | 0 | 0 | 0 | 0 | 15 | 0 | 15 |
| **Pakistan** | 1 | 30 | 23 | 363 | 74 | 0 | 491 |
| **Peru** | 2 | 1 | 360 | 0 | 2328 | 0 | 2691 |
| **South Africa** | 1 | 61 | 874 | 57 | 1162 | 0 | 2155 |
| **Sweden** | 0 | 7 | 280 | 6 | 134 | 0 | 427 |
| **Taiwan** | 0 | 18 | 149 | 0 | 44 | 0 | 211 |
| **Tajikistan** | 0 | 0 | 15 | 0 | 4 | 0 | 19 |
| **Tunisia** | 0 | 0 | 0 | 0 | 2 | 0 | 2 |
| **Turkmenistan** | 0 | 0 | 88 | 1 | 30 | 0 | 119 |
| **Ukraine** | 0 | 0 | 21 | 0 | 9 | 0 | 30 |
| **UK** | 0 | 0 | 3 | 3 | 1 | 0 | 7 |
| **Vietnam** | 0 | 283 | 892 | 4 | 109 | 0 | 1288 |
| **Total** | 35 | 1150 | 5598 | 1850 | 6572 | 6 | 15211 |

**Table S2:** *Lineages –v- geographical location of origin/contribution for CRyPTIC isolates*

| | ALB | ALG | BRZ | BFA | CHN | GER | IND | ITL | JPN | KGZ | NPL | NGA | PAK | PER | ZAF | SWE | TWN | TJK | TUN | TKM | UKR | UK | VNM |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1.1 | 0 | 0 | 0 | 2 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 38 |
| 1.1.1.1 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 188 |
| 1.1.2 | 0 | 0 | 0 | 1 | 1 | 12 | 316 | 16 | 0 | 0 | 2 | 0 | 23 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1.1.3 | 0 | 0 | 1 | 0 | 0 | 1 | 52 | 4 | 0 | 0 | 1 | 0 | 1 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 1.2.1 | 0 | 0 | 0 | 0 | 8 | 12 | 11 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 3 | 16 | 0 | 0 | 0 | 0 | 0 | 24 |
| 1.2.2 | 0 | 0 | 0 | 0 | 1 | 2 | 259 | 5 | 0 | 0 | 2 | 0 | 4 | 0 | 54 | 3 | 1 | 0 | 0 | 0 | 0 | 0 | 1 |
| 2 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2.1 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 20 | 0 | 0 | 0 | 0 | 0 | 34 |
| 2.2 | 0 | 0 | 5 | 1 | 284 | 19 | 272 | 14 | 0 | 0 | 68 | 0 | 6 | 231 | 181 | 13 | 56 | 0 | 0 | 0 | 0 | 0 | 271 |
| 2.2.1 | 0 | 0 | 0 | 0 | 37 | 4 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 3 | 224 | 1 | 21 | 0 | 0 | 0 | 0 | 0 | 26 |
| 2.2.10 | 0 | 0 | 0 | 0 | 1 | 117 | 1 | 84 | 0 | 16 | 0 | 0 | 3 | 0 | 0 | 70 | 0 | 5 | 0 | 75 | 9 | 1 | 0 |
| 2.2.2 | 0 | 0 | 0 | 0 | 32 | 8 | 5 | 11 | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 1 | 5 | 0 | 0 | 0 | 1 | 0 | 8 |
| 2.2.3 | 0 | 0 | 2 | 0 | 172 | 3 | 290 | 4 | 0 | 0 | 21 | 0 | 2 | 1 | 3 | 1 | 10 | 0 | 0 | 0 | 0 | 0 | 100 |
| 2.2.4 | 0 | 0 | 0 | 0 | 3 | 3 | 4 | 8 | 0 | 0 | 0 | 0 | 0 | 0 | 37 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 70 |
| 2.2.5 | 0 | 0 | 0 | 0 | 59 | 1 | 12 | 5 | 0 | 0 | 0 | 0 | 2 | 84 | 1 | 1 | 13 | 0 | 0 | 0 | 0 | 0 | 45 |
| 2.2.6 | 0 | 0 | 1 | 0 | 22 | 8 | 226 | 2 | 0 | 0 | 9 | 0 | 7 | 4 | 204 | 1 | 4 | 0 | 0 | 0 | 0 | 0 | 154 |
| 2.2.7 | 0 | 0 | 0 | 0 | 97 | 4 | 691 | 5 | 0 | 1 | 38 | 0 | 0 | 20 | 206 | 5 | 18 | 0 | 0 | 0 | 0 | 0 | 164 |
| 2.2.8 | 0 | 0 | 0 | 0 | 4 | 0 | 42 | 0 | 0 | 0 | 1 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 2.2.9 | 0 | 0 | 0 | 0 | 0 | 64 | 0 | 51 | 0 | 6 | 0 | 0 | 0 | 2 | 0 | 174 | 0 | 10 | 0 | 11 | 11 | 2 | 0 |
| 3 | 0 | 0 | 0 | 8 | 15 | 37 | 828 | 81 | 0 | 0 | 28 | 0 | 346 | 0 | 34 | 5 | 0 | 0 | 0 | 1 | 0 | 2 | 2 |
| 3.1.1 | 0 | 0 | 0 | 0 | 0 | 17 | 17 | 25 | 0 | 0 | 1 | 0 | 0 | 0 | 19 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 2 |
| 3.1.2 | 0 | 0 | 0 | 0 | 1 | 5 | 201 | 3 | 0 | 0 | 1 | 0 | 7 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 3.1.2.1 | 0 | 0 | 0 | 0 | 0 | 2 | 119 | 3 | 0 | 0 | 4 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4 | 0 | 0 | 1 | 0 | 6 | 4 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 100 | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| 4.1 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.1.1 | 0 | 0 | 0 | 0 | 0 | 6 | 5 | 10 | 0 | 0 | 1 | 0 | 0 | 154 | 13 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.1.1.1 | 0 | 0 | 5 | 0 | 1 | 3 | 20 | 5 | 0 | 0 | 1 | 0 | 2 | 5 | 54 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 4.1.1.2 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.1.1.3 | 0 | 0 | 1 | 6 | 1 | 0 | 30 | 7 | 1 | 0 | 3 | 0 | 2 | 68 | 138 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 4.1.2 | 0 | 0 | 13 | 0 | 0 | 15 | 3 | 26 | 0 | 1 | 0 | 0 | 0 | 31 | 43 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 |
| 4.1.2.1 | 1 | 8 | 87 | 10 | 0 | 111 | 56 | 183 | 0 | 0 | 3 | 2 | 4 | 653 | 54 | 5 | 0 | 1 | 0 | 0 | 1 | 0 | 17 |
| 4.1.3 | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 6 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.10 | 4 | 6 | 35 | 5 | 1 | 110 | 166 | 207 | 0 | 1 | 3 | 0 | 28 | 128 | 170 | 4 | 5 | 2 | 1 | 6 | 2 | 0 | 12 |
| 4.2 | 0 | 0 | 0 | 0 | 3 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.2.1 | 0 | 0 | 0 | 0 | 0 | 9 | 3 | 46 | 0 | 0 | 1 | 0 | 2 | 0 | 2 | 4 | 0 | 0 | 0 | 8 | 3 | 1 | 0 |
| 4.2.2 | 0 | 1 | 0 | 0 | 53 | 31 | 76 | 51 | 0 | 0 | 5 | 0 | 12 | 2 | 10 | 10 | 1 | 0 | 0 | 0 | 0 | 0 | 11 |
| 4.2.2.1 | 0 | 0 | 0 | 0 | 0 | 5 | 0 | 9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3 | 0 | 0 | 17 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 96 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.1 | 0 | 3 | 0 | 1 | 0 | 1 | 2 | 21 | 0 | 0 | 0 | 0 | 0 | 8 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 4.3.2 | 0 | 1 | 10 | 0 | 0 | 3 | 1 | 8 | 0 | 0 | 0 | 0 | 0 | 245 | 21 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.2.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 234 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.3 | 1 | 4 | 34 | 2 | 0 | 33 | 99 | 81 | 0 | 1 | 1 | 0 | 0 | 576 | 120 | 102 | 2 | 1 | 1 | 11 | 2 | 0 | 1 |
| 4.3.4 | 0 | 0 | 1 | 0 | 0 | 0 | 8 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.4.1 | 0 | 0 | 70 | 0 | 0 | 2 | 6 | 3 | 0 | 0 | 0 | 0 | 0 | 60 | 22 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.3.4.2 | 0 | 0 | 38 | 0 | 0 | 6 | 7 | 27 | 0 | 0 | 0 | 0 | 0 | 132 | 12 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| 4.3.4.2.1 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 75 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.4 | 0 | 0 | 0 | 0 | 0 | 0 | 13 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.4.1 | 0 | 0 | 10 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 11 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.4.1.1 | 0 | 0 | 4 | 0 | 0 | 16 | 4 | 66 | 0 | 0 | 1 | 0 | 0 | 22 | 180 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 4 |
| 4.4.1.2 | 0 | 0 | 0 | 0 | 0 | 1 | 25 | 3 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| 4.4.2 | 0 | 0 | 0 | 0 | 124 | 1 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 0 | 5 | 0 | 0 | 0 | 0 | 0 | 16 |
| 4.5 | 0 | 0 | 0 | 0 | 179 | 28 | 15 | 14 | 0 | 0 | 1 | 0 | 18 | 5 | 1 | 0 | 30 | 0 | 0 | 3 | 0 | 0 | 29 |
| 4.6 | 0 | 1 | 0 | 1 | 0 | 1 | 1 | 7 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.1.1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.1.2 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.2 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 14 | 0 | 0 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.2.1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 4.6.2.2 | 1 | 0 | 0 | 34 | 0 | 8 | 1 | 55 | 0 | 0 | 0 | 10 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |

| 6 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Bovis | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 10 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Caprae | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Mixed | 0 | 1 | 9 | 0 | 16 | 3 | 89 | 9 | 0 | 1 | 0 | 0 | 6 | 38 | 25 | 12 | 2 | 0 | 0 | 3 | 0 | 0 | 48 |
| Unknown | 0 | 0 | 1 | 0 | 7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

**Table S3:** *Sub-lineages –v- geographical location of origin/contribution for CRyPTIC isolates*

**Figure S3**: *A significant association between country and lineage can be seen in the CRyPTIC data.* Pearson's chi-squared test, X-squared = 7935.2, df = 110, *p* < 2.2e-16. The correlation plot indicates the relative contribution of each row-column pairing to the chi-square test score (%).

1111

1112

**Figure S4:** *Phylogenetic tree of CRyPTIC M. tuberculosis clinical isolates.* A
phylogenetic cladogram of 15,211 *M. tuberculosis* clinical isolates. A neighbour-joining tree was
constructed from a pairwise distance matrix using *quicktree* (27). Coloured dots at the branch
termini represent the lineage assigned to each isolate. "Animal/Other" includes 16 isolates that
were assigned the following lineages: *M. caprae* (1)*, M. bovis* (1)*,* along with 17 isolates previously
defined as representative for specific sub-lineages (58).

1119

1120

1121

1122

| UNIQUEID | COUNTRY OF ORIGIN | LINEAGE |
|---|---|---|
| site.11.subj.XTB-18-224.lab.XTB-18-224.iso.1 | UNKNOWN | Lineage 2 |
| site.10.subj.YA00026182.lab.YA00026182.iso.1 | ZAF | Lineage 4 |

1123 **Table S5:** *Sample information for isolates classified as resistant to all 13*
1124 *CRyPTIC drugs tested.* The country of origin is specified using the 3-letter country codes
1125 (alpha-3) defined by ISO 3166-1.

1126

1127

1128
1129

1130

| | INH | RIF | EMB | LEV | MXF | AMI | KAN | BDQ | CFZ | DLM | LZD | ETH | RFB |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| INH | 100.0 | 74.8 | 38.0 | 34.0 | 27.8 | 14.0 | 17.0 | 1.5 | 5.5 | 1.7 | 2.0 | 28.2 | 70.4 |
| RIF | 93.5 | 100.0 | 46.3 | 41.4 | 34.1 | 17.2 | 20.8 | 1.8 | 6.0 | 1.7 | 2.3 | 29.4 | 91.3 |
| EMB | 98.5 | 95.9 | 100.0 | 53.9 | 47.1 | 23.3 | 26.4 | 2.4 | 8.2 | 2.3 | 3.7 | 35.8 | 87.3 |
| LEV | 93.3 | 90.2 | 56.9 | 100.0 | 78.5 | 27.3 | 31.3 | 3.1 | 9.4 | 2.9 | 4.7 | 39.2 | 87.1 |
| MXF | 95.0 | 92.3 | 61.9 | 97.6 | 100.0 | 29.9 | 34.6 | 3.2 | 9.9 | 2.7 | 5.3 | 41.9 | 88.8 |
| AMI | 93.0 | 90.4 | 58.9 | 65.3 | 57.8 | 100.0 | 90.4 | 2.4 | 13.0 | 5.2 | 8.1 | 40.7 | 82.8 |
| KAN | 89.4 | 86.7 | 53.0 | 59.5 | 53.2 | 72.0 | 100.0 | 1.8 | 10.6 | 3.2 | 5.7 | 40.4 | 80.0 |
| BDQ | 79.4 | 77.8 | 49.5 | 60.7 | 49.5 | 20.0 | 18.9 | 100.0 | 52.4 | 12.9 | 14.0 | 34.9 | 76.9 |
| CFZ | 61.9 | 52.8 | 34.9 | 38.2 | 32.1 | 21.8 | 22.5 | 10.6 | 100.0 | 9.4 | 10.2 | 27.0 | 53.7 |
| DLM | 55.2 | 44.4 | 28.2 | 33.2 | 23.9 | 24.4 | 19.1 | 7.1 | 26.3 | 100.0 | 19.3 | 22.1 | 45.4 |
| LZD | 77.6 | 69.9 | 53.2 | 64.7 | 58.7 | 46.7 | 41.2 | 9.8 | 34.2 | 24.1 | 100.0 | 42.6 | 66.7 |
| ETH | 96.5 | 79.4 | 47.0 | 48.7 | 41.9 | 21.1 | 26.2 | 2.2 | 8.3 | 2.4 | 3.9 | 100.0 | 77.1 |
| RFB | 93.3 | 96.8 | 44.5 | 42.1 | 34.4 | 16.6 | 20.3 | 1.9 | 6.4 | 1.9 | 2.3 | 29.9 | 100.0 |

1131 **Table S6:** *Co-occurrence of antibiotic resistance in CRyPTIC M. tuberculosis*
1132 *isolates.* The probability (%) of an isolate being resistant to Drug 2 (top) if it is resistant to Drug
1133 1 (left). Drug acronyms: INH = isoniazid, RIF = rifampicin, EMB = ethambutol, LEV = levofloxacin,
1134 MXF = moxifloxacin, AMI = amikacin, KAN = kanamycin, BDQ = bedaquiline, CFZ = clofazimine,
1135 DLM = delamanid, LZD = linezolid, ETH = ethionamide, RFB = rifabutin.
1136
1137

1138

1139 **Figure S6:** *Non-synonymous mutations found outside the RRDR of rpoB in RMR*
1140 *isolates and MDR isolates.* Presence of a coloured spot indicates that the mutation was
1141 found in RMR/MDR isolates and spot size corresponds to the proportion of RMR or MDR isolates
1142 carrying that mutation.
1143
1144

## References

**References**

1.   WHO. Global Tuberculosis Report 2020. 2021;

2.   Shinnick TM, Starks AM, Alexander HL, Castro KG. Evaluation of the Cepheid Xpert
MTB/RIF assay. Expert review of molecular diagnostics. 2015 Jan;15(1):9–22.

3.   Boehme CC, Nicol MP, Nabeta P, Michael JS, Gotuzzo E, Tahirli R, et al. Feasibility,
diagnostic accuracy, and effectiveness of decentralised use of the Xpert MTB/RIF test
for diagnosis of tuberculosis and multidrug resistance: a multicentre implementation
study. Lancet (London, England). 2011 Apr 30;377(9776):1495–505.

4.   Makhado NA, Matabane E, Faccin M, Pinçon C, Jouet A, Boutachkourt F, et al.
Outbreak of multidrug-resistant tuberculosis in South Africa undetected by WHO-
endorsed commercial tests: an observational study. The Lancet Infectious Diseases.
2018 Dec;18(12):1350–9.

5.   Beckert P, Sanchez-Padilla E, Merker M, Dreyer V, Kohl TA, Utpatel C, et al. MDR
*M. tuberculosis* outbreak clone in Eswatini missed by Xpert has elevated bedaquiline
resistance dated to the pre-treatment era. Genome Medicine. 2020 Dec 25;12(1):104.

6.   Sanchez-Padilla E, Merker M, Beckert P, Jochims F, Dlamini T, Kahn P, et al.
Detection of Drug-Resistant Tuberculosis by Xpert MTB/RIF in Swaziland. New
England Journal of Medicine. 2015 Mar 19;372(12):1181–2.

7.   The CRyPTIC Consortium, 100 000 Genomes ProjecT. Prediction of Susceptibility to
First-Line Tuberculosis Drugs by DNA Sequencing. New England Journal of
Medicine. 2018 Oct 11;379(15):1403–15.

8.   Pankhurst LJ, del Ojo Elias C, Votintseva AA, Walker TM, Cole K, Davies J, et al.
Rapid, comprehensive, and affordable mycobacterial diagnosis with whole-genome
sequencing: a prospective study. The Lancet Respiratory medicine. 2016 Jan;4(1):49–
58.

9.   Kalokhe AS, Shafiq M, Lee JC, Ray SM, Wang YF, Metchock B, et al. Multidrug-
resistant tuberculosis drug susceptibility and molecular diagnostic testing. The
American journal of the medical sciences. 2013 Feb;345(2):143–8.

10.   Rancoita PM v, Cugnata F, Gibertoni Cruz AL, Borroni E, Hoosdally SJ, Walker TM,
et al. Validating a 14-Drug Microtiter Plate Containing Bedaquiline and Delamanid for
Large-Scale Research Susceptibility Testing of *Mycobacterium tuberculosis*.
Antimicrobial agents and chemotherapy. 2018;62(9).

11.   The CRyPTIC Consortium. Epidemiological cutoff values for a 96-well broth
microdilution plate for high-throughput research antibiotic susceptibility testing of *M.
tuberculosis*. European Respiratory Journal. 2022 Mar 17;2200239.

12.   The CRyPTIC Consortium. Quantitative measurement of antibiotic resistance in
*Mycobacterium tuberculosis* reveals genetic determinants of resistance and
susceptibility in a target gene approach. bioRxiv. 2021;

13.   The CRyPTIC Consortium. Predicting Susceptibility to First- and Second-line
Tuberculosis Drugs by DNA sequencing and Machine Learning. In preparation.

14.   The CRyPTIC Consortium. Genome-wide association studies of global
*Mycobacterium tuberculosis* resistance to thirteen antimicrobials in 10,228 genomes.
bioRxiv. 2021;

15.   Sonnenkalb L, Carter J, Spitaleri A, Iqbal Z, Hunt M, Malone K, et al. Deciphering
Bedaquiline and Clofazimine Resistance in Tuberculosis: An Evolutionary Medicine
Approach. bioRxiv. 2021 Jan 1;2021.03.19.436148

16.   Fowler PW, Gibertoni Cruz AL, Hoosdally SJ, Jarrett L, Borroni E, Chiacchiaretta M,
et al. Automated detection of bacterial growth on 96-well plates for high-throughput
drug susceptibility testing of *Mycobacterium tuberculosis*. Microbiology. 2018 Dec
1;164(12):1522–30.

17. Fowler PW, Wright C, Spiers-Bowers H, Zhu T, Baeten EML, Hoosdally SW, et al.
BashTheBug: a crowd of volunteers reproducibly and accurately measure the
minimum inhibitory concentrations of 13 antitubercular drugs from photographs of 96-
well broth microdilution plates. bioRxiv. 2021 Jan 1;2021.07.20.453060

18. Hunt M, Letcher B, Malone KM, Nguyen G, Hall MB, Colquhoun RM, et al. Minos:
variant adjudication and joint genotyping of cohorts of bacterial genomes. bioRxiv.
2021 Jan 1;2021.09.15.460475.

19. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics. 2009 Jul 15;25(14):1754–60.

20. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014 Aug 1;30(15):2114–20.

21. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics (Oxford, England). 2009 Aug
15;25(16):2078–9.

22. Iqbal Z, Caccamo M, Turner I, Flicek P, McVean G. De novo assembly and
genotyping of variants using colored de Bruijn graphs. Nature Genetics. 2012 Feb
8;44(2):226–32.

23. Hunt M, Bradley P, Lapierre SG, Heys S, Thomsit M, Hall MB, et al. Antibiotic
resistance prediction for *Mycobacterium tuberculosis* from genome sequence data with
Mykrobe. Wellcome open research. 2019;4:191.

24. Walker TM, Lalor MK, Broda A, Ortega LS, Morgan M, Parker L, et al. Assessment
of *Mycobacterium tuberculosis* transmission in Oxfordshire, UK, 2007–12, with whole
pathogen genome sequences: an observational study. The Lancet Respiratory
Medicine. 2014 Apr;2(4):285–92.

25. Miotto P, Tessema B, Tagliani E, Chindelevitch L, Starks AM, Emerson C, et al. A
standardised method for interpreting the association between mutations and phenotypic
drug resistance in *Mycobacterium tuberculosis*. European Respiratory Journal. 2017
Dec;50(6):1701354.

26. Walker TM, Miotto P, Köser CU, Fowler PW, Knaggs J, Iqbal Z, et al. The 2021
WHO Catalogue of *Mycobacterium Tuberculosis* Complex Mutations Associated with
Drug Resistance: A New Global Standard for Molecular Diagnostics. SSRN Electronic
Journal. 2021;

27. Howe K, Bateman A, Durbin R. QuickTree: building huge Neighbour-Joining trees of
protein sequences. Bioinformatics. 2002 Nov 1;18(11):1546–7.

28. Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Current Protocols in
Bioinformatics. 2020 Mar 5;69(1).

29. Coscolla M, Gagneux S. Consequences of genomic diversity *in Mycobacterium
tuberculosis*. Seminars in Immunology. 2014 Dec;26(6):431–44.

30. Gagneux S. Ecology and evolution of *Mycobacterium tuberculosis*. Nature Reviews
Microbiology. 2018 Apr 19;16(4):202–13.

31. Freschi L, Vargas R, Husain A, Kamal SMM, Skrahina A, Tahseen S, et al. Population
structure, biogeography and transmissibility of *Mycobacterium tuberculosis*. Nature
Communications. 2021 Dec 20;12(1):6099.

32. Ghimire S, Karki S, Maharjan B, Kosterink JGW, Touw DJ, van der Werf TS, et al.
Treatment outcomes of patients with MDR-TB in Nepal on a current programmatic
standardised regimen: retrospective single-centre study. BMJ Open Respiratory
Research. 2020 Aug 12;7(1):e000606.

33. Sharma R, Sharma SK, Singh BK, Mittal A, Kumar P. High degree of fluoroquinolone
resistance among pulmonary tuberculosis patients in New Delhi, India. The Indian
journal of medical research. 2019 Jan;149(1):62–6.

34. Dijkstra JA, van der Laan T, Akkerman OW, Bolhuis MS, de Lange WCM, Kosterink
JGW, et al. In Vitro Susceptibility of *Mycobacterium tuberculosis* to Amikacin,
Kanamycin, and Capreomycin. Antimicrobial Agents and Chemotherapy. 2018
Mar;62(3).
35. Maitre T, Petitjean G, Chauffour A, Bernard C, el Helali N, Jarlier V, et al. Are
moxifloxacin and levofloxacin equally effective to treat XDR tuberculosis? Journal of
Antimicrobial Chemotherapy. 2017 Aug 1;72(8):2326–33.
36. Berrada ZL, Lin S-YG, Rodwell TC, Nguyen D, Schecter GF, Pham L, et al. Rifabutin
and rifampin resistance levels and associated rpoB mutations in clinical isolates of
*Mycobacterium tuberculosis* complex. Diagnostic Microbiology and Infectious
Disease. 2016 Jun;85(2):177–81.
37. Ho J, Jelfs P, Sintchenko V. Fluoroquinolone resistance in non-multidrug-resistant
tuberculosis—a surveillance study in New South Wales, Australia, and a review of
global resistance rates. International Journal of Infectious Diseases. 2014 Sep;26:149–
53.
38. Singh R, Manjunatha U, Boshoff HIM, Ha YH, Niyomrattanakit P, Ledwidge R, et al.
PA-824 Kills Nonreplicating *Mycobacterium tuberculosis* by Intracellular NO Release.
Science. 2008 Nov 28;322(5906):1392–5.
39. Hartkoorn RC, Uplekar S, Cole ST. Cross-Resistance between Clofazimine and
Bedaquiline through Upregulation of MmpL5 in *Mycobacterium tuberculosis*.
Antimicrobial Agents and Chemotherapy. 2014 May;58(5):2979–81.
40. Degiacomi G, Sammartino JC, Sinigiani V, Marra P, Urbani A, Pasca MR. In vitro
Study of Bedaquiline Resistance in *Mycobacterium tuberculosis* Multi-Drug Resistant
Clinical Isolates. Frontiers in microbiology. 2020;11:559469.
41. Almeida D, Ioerger T, Tyagi S, Li S-Y, Mdluli K, Andries K, et al. Mutations in pepQ
Confer Low-Level Resistance to Bedaquiline and Clofazimine in *Mycobacterium*
*tuberculosis.* Antimicrobial Agents and Chemotherapy. 2016 Aug;60(8):4590–9.
42. Nimmo C, Millard J, van Dorp L, Brien K, Moodley S, Wolf A, et al. Population-level
emergence of bedaquiline and clofazimine resistance-associated variants among
patients with drug-resistant tuberculosis in southern Africa: a phenotypic and
phylogenetic analysis. The Lancet Microbe. 2020 Aug;1(4):e165–74.
43. Mvelase NR, Balakrishna Y, Lutchminarain K, Mlisana K. Evolving rifampicin and
isoniazid mono-resistance in a high multidrug-resistant and extensively drug-resistant
tuberculosis region: a retrospective data analysis. BMJ Open. 2019 Nov
6;9(11):e031663.
44. Villegas L, Otero L, Sterling TR, Huaman MA, van der Stuyft P, Gotuzzo E, et al.
Prevalence, Risk Factors, and Treatment Outcomes of Isoniazid- and Rifampicin-
Mono-Resistant Pulmonary Tuberculosis in Lima, Peru. PLOS ONE. 2016 Apr
5;11(4):e0152933.
45. Park S, Jo K-W, Lee S do, Kim WS, Shim TS. Treatment outcomes of rifampin-
sparing treatment in patients with pulmonary tuberculosis with rifampin-mono-
resistance or rifampin adverse events: A retrospective cohort analysis. Respiratory
Medicine. 2017 Oct;131:43–8.
46. Salaam-Dreyer Z, Streicher EM, Sirgel FA, Menardo F, Borrell S, Reinhard M, et al.
Rifampicin-Monoresistant Tuberculosis Is Not the Same as Multidrug-Resistant
Tuberculosis: a Descriptive Study from Khayelitsha, South Africa. Antimicrobial
Agents and Chemotherapy. 2021 Oct 18;65(11).
47. Pang Y, Lu J, Wang Y, Song Y, Wang S, Zhao Y. Study of the Rifampin
Monoresistance Mechanism in *Mycobacterium tuberculosis*. Antimicrobial Agents and
Chemotherapy. 2013 Feb;57(2):893–900.

48. Kigozi E, Kasule GW, Musisi K, Lukoye D, Kyobe S, Katabazi FA, et al. Prevalence
and patterns of rifampicin and isoniazid resistance conferring mutations in
*Mycobacterium tuberculosis* isolates from Uganda. PLOS ONE. 2018 May
30;13(5):e0198091.
49. Dean AS, Zignol M, Cabibbe AM, Falzon D, Glaziou P, Cirillo DM, et al. Prevalence
and genetic profiles of isoniazid resistance in tuberculosis patients: A multicountry
analysis of cross-sectional data. PLOS Medicine. 2020 Jan 21;17(1):e1003008.
50. Migliori GB, Langendam MW, D'Ambrosio L, Centis R, Blasi F, Huitric E, et al.
Protecting the tuberculosis drug pipeline: stating the case for the rational use of
fluoroquinolones. European Respiratory Journal. 2012 Oct;40(4):814–22.
51. Heysell SK, Ahmed S, Rahman MdT, Akhanda MdW, Gleason AT, Ebers A, et al.
Hearing loss with kanamycin treatment for multidrug-resistant tuberculosis in
Bangladesh. European Respiratory Journal. 2018 Mar;51(3):1701778.
52. WHO. Update on the use of nucleic acid amplification tests to detect TB and drug-
resistant TB: rapid communication. 2021;
53. Olaru ID, Heyckendorf J, Andres S, Kalsdorf B, Lange C. Bedaquiline-based treatment
regimen for multidrug-resistant tuberculosis. European Respiratory Journal. 2017 May
21;49(5):1700742.
54. WHO. World Health Organization treatment guidelines for drug-resistant tuberculosis,
2016 update. 2016;
55. Huitric E, Verhasselt P, Koul A, Andries K, Hoffner S, Andersson DI. Rates and
Mechanisms of Resistance Development in *Mycobacterium tuberculosis* to a Novel
Diarylquinoline ATP Synthase Inhibitor. Antimicrobial Agents and Chemotherapy.
2010 Mar;54(3):1022–8.
56. Holt KE, McAdam P, Thai PVK, Thuong NTT, Ha DTM, Lan NN, et al. Frequent
transmission of the *Mycobacterium tuberculosis* Beijing lineage and positive selection
for the EsxW Beijing variant in Vietnam. Nature Genetics. 2018 Jun 21;50(6):849–56.
57. Chiner-Oms Á, Comas I. Large genomics datasets shed light on the evolution of the
Mycobacterium tuberculosis complex. Infection, Genetics and Evolution. 2019
Aug;72:10–5.
58. Borrell S, Trauner A, Brites D, Rigouts L, Loiseau C, Coscolla M, et al. Reference set
of *Mycobacterium tuberculosis* clinical strains: A tool for research and product
development. PLOS ONE. 2019 Mar 25;14(3):e0214088.
1328