Quantitative measurement of antibiotic resistance in

Mycobacterium tuberculosis reveals genetic determinants of

resistance and susceptibility in a target gene approach

The CRyPTIC Consortium[1]

Abstract

The World Health Organization goal of universal drug susceptibility testing for patients

with tuberculosis is most likely to be achieved through molecular diagnostics; however,

to date these have focused largely on first-line drugs, and always on predicting binary

susceptibilities. Here, we used whole genome sequencing and a quantitative microtiter

plate assay to relate genomic mutations to minimum inhibitory concentration in 15,211

Mycobacterium tuberculosis patient isolates from 27 countries across five continents.

This work identifies 449 unique MIC-elevating genetic determinants across thirteen

drugs, as well as 91 mutations resulting in hypersensitivity for eleven drugs. Our results

provide a guide for further implementation of personalized medicine for the treatment of

tuberculosis using genetics-based diagnostics and can serve as a training set for novel

approaches to predict drug resistance.

[126 words]

---

[1] See the CRyPTIC Author List at the end of manuscript

Introduction (Intro-Discussion 3987/4000 words):

Mycobacterium tuberculosis (Mtb) is a persistent global public health threat that caused

an estimated 10 million new cases of tuberculosis (TB) and 1.4 million deaths in 2019[1].

Of particular concern are the estimated 465,000 rifampicin resistant (RR) cases, 78% of

which were multi-drug resistant (MDR, resistant to both rifampicin and isoniazid)[1]. Drug

resistance poses two major challenges to the successful treatment of TB, as it is both

underdiagnosed (only 38% of RR/MDR cases in 2019)—leading to under-treatment—

and has poor treatment success rates even when identified (57% globally in 2019)[1].

Despite attempts to move to shorter and all-oral MDR TB regimens using new drugs,

most patients are still receiving toxic regimens that decrease patient adherence[1,2].

Collectively, the failure to identify and successfully treat these cases leads to onward

transmission and amplification of drug resistant strains.

The WHO has identified better diagnosis and treatment of drug resistant tuberculosis as

a key part of the global tuberculosis eradication strategy[1]. Rapid genetics-based

diagnostic tools, such as GeneXpert and other line probe assays, have been widely

adopted as they are faster and cheaper than traditional culture-based diagnostic

susceptibility testing (DST). However, outbreaks caused by drug-resistant strains with

mutations not detected by such assays reveal the importance of developing more

comprehensive assays that include a wider range of resistance determinants[3]. Some

approaches incorporate whole-genome sequencing (WGS) or targeted next generation

sequencing to identify all possible resistant variants and recently these methods have

proven to be capable of replacing culture-based DST for the first line drugs; however,

implementation of this technology is not yet feasible globally due to cost and technical

expertise constraints[4–6].

Most current culture and genetics-based DST approaches generate binary results—

'resistant' or 'susceptible'—and thus fail to consistently report elevations in minimum

inhibitory concentration (MIC) below or around the critical concentration[7]. These sub-

threshold elevations in MIC may nevertheless be clinically meaningful, as the

combination of significant interpatient pharmacokinetic variability and elevated MICs

predisposes Mtb strains to development of higher-level resistance, risking treatment

failure and worse patient outcomes[8,9]. A binary system also hampers the wider

implementation of informed high-dose regimens which have been trialed to extend the

clinical utility of relatively less toxic and more widely available drugs such as rifampicin

and isoniazid[10–12]. While some previous efforts have attempted to use quantitative MICs

to identify these lower-level resistance variants, they have been limited by smaller

sample sizes and/or the need to combine heterogenous methods of resistance

determination for meta-analyses of publicly available data[13]. Additionally, relatively few

studies have had adequate sample sizes to systematically investigate new and

repurposed drugs such as bedaquiline, linezolid, clofazimine and delamanid that are

poised to become the new "front-line" drugs for the MDR-TB treatment.

To resolve these issues, we performed WGS and determined the MICs of thirteen drugs

for 15,211 Mtb isolates selected from patient samples gathered from 27 countries over

five continents using a previously validated microtiter plate[14]. This data covers all first-

line drugs (except pyrazinamide), as well as eight drugs from the new MDR-TB treatment guidelines (all Group A, one Group B, and four Group C)[15]. The results can serve as guides for pharmacokinetic and dosing studies to extend the clinical utility of less toxic and more widely available drugs for the treatment of drug-resistant tuberculosis, as well as help to improve the design of genetics-based rapid diagnostics for MDR-TB and the recently published WHO genetic catalogue for tuberculosis[16]. They also provide a large, quality-controlled dataset for development of drug resistance prediction algorithms using machine-learning and other approaches.

Results:

Dataset description

Bacterial isolates were collected from patient samples from 27 different countries and were over-sampled for drug resistance. Of the 15,211 isolates included in the initial CRyPTIC dataset, 5,541 were phenotypically susceptible to all of isoniazid, rifampicin, and ethambutol, 5,602 were isoniazid mono-resistant, 5,261 were rifampicin resistant, and 4,125 were multidrug resistant (MDR, resistant to both rifampicin and isoniazid) based on previously published epidemiological cutoffs (ECOFF, MIC that encompasses 99% of wild type) for the microtiter plates used in this study[17]. Binary phenotypic resistance to the newer drugs was observed at a lower prevalence, with 71 bedaquiline resistant isolates, 106 clofazimine resistant isolates, 76 linezolid resistant isolates, and 85 delamanid resistant isolates (Table S1). Isolate lineages were determined using a published SNP-based protocol from WGS data and the lineage distribution across countries reflects previously described phylogeographic distributions[18–20]. Five out of six major lineages of Mtb were represented in the dataset, with the majority of isolates mapping to L4 (6,066 isolates) and L2 (4,323 isolates), while L3 (1,059), L1 (677), and L6 (6) comprised the remainder. A full description of the CRyPTIC dataset and determination of the ECOFFs has been previously published (also see Methods)[17].

Genetic resistance determinants in Mycobacterium tuberculosis

Previous studies have shown that the vast majority of genetic determinants of resistance to most anti-tuberculosis drugs are related to a relatively small number of genes[6,21]. We thus employed a candidate gene approach and restricted our

investigation of genomic variation to previously identified genes and the 100bp

upstream for each drug (Table 1). All unique genomic variation in the target genes and

upstream regions (SNPs, both synonymous and nonsynonymous, as well as insertions

and deletions <50 base pairs in length) that occurred in isolates with matched high-

quality phenotypic data was included in a separate multivariable linear mixed model

controlling for population structure and technical variation between sites for each drug,

after removing isolates with evidence for mixed allele calls at sites previously identified

as resistant (e.g. rpoB S450X, Methods). Final sample sizes per drug ranged from

6,681 for moxifloxacin to 10,042 for rifabutin (mean sample size 8,353, Figure 1,

Methods). Most isolates had less than five nonsynonymous mutations across all target

genes for each drug (Table S2).

Across thirteen drugs, 540 mutations in 39 genes (out of 4,667 mutations and 49 genes

tested) were found to have independent effects on MIC after correction for multiple

testing (Benjamini-Hochberg correction, false discovery rate<0.05, Figure 2, Table 1,

S3). Ethionamide had the most unique variants associated with reduced susceptibility

(128), while linezolid had the least (8). Effect sizes were measured in log2MIC (where

an increase in 1 log2MIC was equivalent to a doubling of the MIC) and positive effects

for estimates derived from at least three observations ranged from a 0.22 increase in

kanamycin log2(MIC) for *rrs* c492t to a 10.1 increase in isoniazid log2(MIC) for *katG*

W477Stop. Multiple promoter mutations were implicated in resistance to isoniazid,

ethionamide, amikacin, kanamycin, and ethambutol (Figure 2B). The effects of promoter

mutations varied widely, with mutations upstream of *eis* and *embA* being almost

exclusively associated with sub-ECOFF elevations in MIC for amikacin and ethambutol

respectively, while most promoter mutations for the isoniazid and ethionamide-related

fabG1 resulted in MICs above the ECOFF[17]. While a prior study found that common

promoter mutations tended to be associated with lower-level resistance than their

corresponding common gene-body counterparts (e.g. *fabG1* c-15 vs *inhA* I21), we found

that intergenic vs gene body mutation was only associated with significantly different

effects on MIC for isoniazid, ethambutol, and kanamycin (Table S4)[13]. In fact, we found

that the widespread *fabG1* c-15t promoter mutation was associated with higher-level

and equivalent-level resistance to its gene body counterparts *inhA* I21V and I21T

respectively (Figure 2B, Wald test for equality of coefficients p=0.0006, p=0.24

respectively). Resistance-associated promoter mutations were enriched in the region

around each gene's respective -10 element, which is consistent with the essentiality of

the -10 hexamer and increased variability around the -35 position in mycobacterial

promoters (Figure S1, +/- 5 nucleotides, Mantel-Haenszel common OR=4.5,

p=0.0007)[22,23]. Multiple insertion/deletion mutations were associated with resistance to

isoniazid, rifampicin, rifabutin, ethionamide, ethambutol, bedaquiline, clofazimine, and

delamanid (Table S3, Figure 2). Homoplastic mutations (multiple evolutionarily

independent occurrences) were more likely to be associated with resistance for all

drugs except amikacin, clofazimine, linezolid, and delamanid (Woolf test for

homogeneity of ORs p=0.0001, Table S5).

One notable advantage of quantitative MIC measurements is that they also

enable investigation of variants associated with MIC decreases. We identified 63

increased susceptibility-associated mutations (with at least three occurrences) whose

effect sizes ranged from -4.3 rifampicin log2(MIC) for Rv2752c H371Y to -0.23

kanamycin log2(MIC) for *eis* V163I (Figure 2A). Eight of these mutations were

homoplastic with at least three independent occurrences, which raises the question of

what selective pressure may drive increases in drug susceptibility (Methods).

First-line drugs

Rifampicin is a critical first line drug and resistance to it is almost entirely mediated by

mutations within an 81-base pair region of the *rpoB* gene (rifampicin resistance

determining region, RRDR). Most molecular assays target mutations in this region for

rapid prediction of rifampicin resistance, however, mutations outside this region have

been associated with outbreaks[24,25]. We identified 35 mutations in *rpoB* occurring at

least 3 times whose effects collectively ranged from 1.0 to 9.0 increases in log2MIC

(Figure 3A). Notably, seven unique resistance-associated mutations occurred outside

the RRDR, at positions V170, Q172, I491, and L731; however, only V170F was

associated with high level resistance (8.37 increased log2MIC). Although disparate in

primary sequence from the RDRR, positions V170, Q172 and I491 are all near the drug-

binding pocket structurally (Figure 3B). Interestingly, a homoplastic in-frame deletion 12

bp in size in the RRDR was also associated with rifampicin resistance (Figure 3C, Table

S3). Several types of insertion/deletion mutations in the RDRR have previously been

reported, although they are rare, consistent with their greater structural consequences

for the essential RNA polymerase[26].

Prior studies have identified seven "borderline" mutations in *rpoB* (L430P,

D435Y, H445L, H445N, H445S, L452P, and I491F) for rifampicin; resistant isolates with

these mutations are often missed by phenotypic methods such as the Mycobacterial

Growth Indicator Tube (MGIT), possibly due to slower growth rates, which has led to a

reduction in the critical concentration for MGIT in the latest WHO guidelines[27–29]. These

mutations' MICs range on the plate from 5.1 log2MIC for H445L to 2.3 log2MIC for

L430P (rifampicin ECOFF minus baseline MIC=3.3, Table S3). Here, we identify thirteen

additional *rpoB* mutations independently associated with elevated MICs that are less

than 5.1 log2MIC (8/13 located in the RDRR, Table S3). Sixteen *rpoB* mutations in total

were independently associated with elevated MICs at or below the rifampicin ECOFF,

including rpoB L430P, a variant that has been successfully treated with a high dose

rifampicin-containing regimen clinically[30].Several rpoB positions (Q432, D435, H445)

harbored both high and low-level resistance-associated alleles, while others (L430,

L452, I491) were associated exclusively with lower-level resistance regardless of the

amino acid substitution (Figure 3B,C orange and yellow shading respectively). Mapping

these mutations onto the rpoB structure revealed that high--level resistance often

involves disruption of the interactions with the rigid napthol ring while mutations at

positions that contact the ansa bridge had more variable effects, potentially due to

increased structural flexibility in this region of the drug (Figure 3B). Low-level resistance

mutations often co-occurred with other low-level resistance mutations, producing high-

level resistance additively.

Rifabutin is a structural analogue to rifampicin that is sometimes used due to its more

favorable drug interaction profile. It is associated with a lower ECOFF and a unique

resistance profile regarding rpoB mutations (Figure 3A, Table S3). Mutations in *rpoB*

were associated with lower elevations in rifabutin MIC compared to rifampicin MIC

(paired Wilcoxon p=3.7e-9). Interestingly however, all structural features contacted by

these mutations were shared between rifampicin and rifabutin (Figure 3B). A single

mutation, rpoB Q409R (n=24, p=5.0e-3 after Benjamini-Hochberg (BH) correction), was

associated with decreased rifampicin and rifabutin MICs; interestingly, this mutation has

been proposed as a compensatory mutation that may alter the rate of transcription

initiation and resulting transcription efficiency for isolates that harbor other RDRR

mutations[31].

Resistance to isoniazid is mediated primarily through loss-of-function mutations in the

prodrug converting enzyme *katG*, with canonical high-level resistance caused by the

S315T mutation, which was associated with a 6.2 log2 increase in MIC (Figure 4A). Not

all *katG* mutations were associated with high level resistance, nearly half (15/31) being

associated with increases in MIC at or below the ECOFF. No mutations likely to result in

severe loss of function (i.e. nonsense and frameshifting insertion/deletion mutations)

were associated with sub-ECOFF resistance, supporting the consensus of treating

presumptive loss-of-function mutants in *katG* as resistant. The other canonical

isoniazid-related genes, *inhA* and *fabG1*, tended to be associated with lower-level

resistance, with 4/6 and 5/6 mutations associated with sub-ECOFF increases in MIC

respectively (Figure 4A, Table S3). While *fabG1* was previously the only synonymous

mutation known to be associated with resistance to isoniazid, here we identify a

synonymous mutation in the first codon of *katG* that confers high-level resistance to

isoniazid, likely by reducing the rate of translation initiation and subsequent production

of katG enzyme required for activation of isoniazid (4.5 log2MIC, n=3, p=1.4e-8 after

Benjamini-Hochberg (BH) correction, Table S3).

Most isoniazid resistance-associated mutations in katG occurred in the N-terminal lobe

responsible for heme-binding and pro-drug conversion (Figure 4B). The vast majority of

isolates harbored variation at position S315, which is located in the primary isoniazid-

binding pocket on the δ edge of the heme; interestingly however, another cluster of

resistance-associated mutations occurred in the helix made up of residues 138-155. A

recent study found some structural evidence for promiscuous isoniazid binding at this

site and a study of mutations in this region in *Escherichia coli* found they reduced

catalase/peroxidase activity and heme binding; however the precise mechanism of

effect of these mutations in Mtb is unknown[32,33]. Intriguingly, one mutation in this region,

KatG S140N, was associated with decreased isoniazid MIC (n=9, p=5.4e-4 after BH

correction, Figure 4B).

Non-canonical isoniazid resistance-associated variants were identified in *ahpC*, *ndh*,

and *Rv1258c* (*tap*) (Figure 4A). *ahpC* promoter mutations have previously been

identified in resistant isolates and are thought to represent compensatory mutations that

overexpress an alternate peroxidase to compensate for the reduced/loss-of-function of

katG. Mutations in *ahpC* were associated with increased MICs; however, these

mutations almost always co-occurred with mutations in canonical isoniazid genes and

investigation of the interaction between these co-occurring mutation pairs revealed that

ahpC mutations did not result in additive resistance (Figure 4A). Several recent

genome-wide association studies (GWAS) have implicated mutations in the bifunctional

ribonuclease/beta-lactamase *Rv2752c* in resistance and tolerance to both rifampicin

and isoniazid; however, they also identified convergent mutations in drug susceptible

strains[13,34]. While we identified nine nonsynonymous mutations with significant effects

on log2MIC, only one, V218L, was shared between isoniazid and rifampicin, causing a

3.2 elevation in log2MIC for both drugs (Table S3). Only one other *Rv2752c* variant was

associated with elevated rifampicin MICs, while four variants in this gene were

associated with elevated isoniazid MICs (Figure 4A).

Canonical ethambutol resistance is mediated by mutations in *embA* or *embB*. We

identified 45 variants, 12 in the *embC-embA* intergenic region, five in *embA*, and 28 in

embB, that were independently associated with elevated ethambutol MICs (Figure 4C).

Mutations in the *embC-embA* intergenic region have been proposed to upregulate

production of embA and embB by altered promoter structure. Most *embA* variants were

in the upstream region from -16 to -8, however three were located upstream around the

-35 element. Additionally, four synonymous mutations and one missense mutation were

also associated with increased ethambutol MICs, although only two of these were

homoplastic; all *embA* mutations were associated with MIC increases below the ECOFF

(Figure 4C, Table S3). Interestingly, 22/28 mutations in *embB* were also associated

with sub-ECOFF increases in MIC, including the canonical embB M306I. Low-level

resistance mutations often co-occurred, resulting in high-level additive resistance,

consistent with previous studies (Table S6)[35]. Mutations associated with resistance in

embB were clustered around the drug binding pocket (Figure 4D)[36]. We also identified

resistance-associated variants in *embC* and *ubiA*, although these occur less frequently

and require further validation.

Group A and B MDR drugs

The principal mechanism of resistance to fluoroquinolones is mutations in either subunit

of DNA gyrase (*gyrA* or *gyrB*). We identified 22 mutations (12 gyrA, 10 gyrB) that were

independently associated with increased levofloxacin MIC and 19 mutations (10 gyrA, 9

gyrB) independently associated with increased moxifloxacin MIC (Figure 5A).

Resistance-associated mutations in *gyrB* occurred without an accompanying *gyrA*

mutation ~65% of the time (29/44 isolates LEV, 35/54 isolates MXF) but were

associated with lower overall—and in some cases sub-ECOFF—changes in MIC

(Figure 5A, Table S3,S6). Most mutations associated with increased fluoroquinolone

MICs were within 10 Å of the drug binding pocket (Figure 5B). Intriguingly, two

positions—gyrB R446 and gyrB S447—each harbored two unique resistance-

associated missense mutations despite being over 25 Å from the bound

fluoroquinolone. Both residues make contacts with the gyrB protein backbone at

positions 473-475, suggesting they may exert an allosteric effect by either influencing

protein folding and/or the position of residues (notably D461 and R482) that make up

part of the fluoroquinolone binding pocket (Figure 5B). Interestingly, while gyrB E501D

was associated with resistance 1 log2MIC above the moxifloxacin ECOFF, it did not

cause a similar elevation for levofloxacin (0.1 log2MIC above ECOFF), consistent with

previous studies[7,37,38]. We speculate this could be due to alteration in the coordination of

gyrB R482—which must shift to accommodate the bulkier side group of moxifloxacin—

although this remains to be shown experimentally (Figure 5B).

While initial studies on bedaquiline and clofazimine resistance highlighted *atpE*

(bedaquiline), *pepQ*, *Rv0678*, and *Rv1979c* as mediating resistance, surveillance of

clinical samples has revealed the importance of the efflux mechanism mediated by the

mmpL5 membrane transporter, which is controlled by the transcriptional regulator

Rv0678. Consistent with this, we identified sixteen and four mutations in *Rv0678* that

were associated with elevated bedaquiline and clofazimine MICs respectively, of which

four were shared (Figure S2, Table S3). We also identified two *mmpL5* mutations that

were associated with increased MICs for each drug which were not shared between the

two drugs. Finally, we identified both the atpE E61D (n=3) drug binding site mutation

associated with bedaquiline resistance and two mutations in *Rv1979c* associated with

clofazimine resistance. No mutations in *pepQ* were associated with resistance to either

drug. Importantly, 5 unique nonsense and frameshift mutations in *mmpL5* increased

susceptibility to bedaquiline by -1.9 to -4.0 log2MIC, of which one, mmpL5 Y300Stop,

was also shared with clofazimine (Figure 2A). Inactivating mutations in *mmpL5*

abrogated resistance mediated by co-occurring *Rv0678* mutations, consistent with the

proposed mechanism of efflux-mediated resistance and a hypothesis proposed by a

prior study[39].

Resistance to linezolid is mediated by mutations in *rplC* and *rrl*, which tend to cause

higher- and lower-level resistance respectively. We identified the classical rplC C154R

(n=43) mutation and five variants in *rrl* associated with elevated linezolid MICs (Figure

S2, Table S3).

Group C MDR drugs

Aminoglycoside resistance is canonically mediated by mutations in the 16s rRNA

encoded by *rrs*. We identified five and six mutations in *rrs* that were independently

associated with elevated MICs for amikacin and kanamycin respectively (Figure 5C).

Seven promoter mutations in *eis* were associated with elevated MICs to kanamycin,

with three also being associated with elevated amikacin MICs. Interestingly, *eis*

promoter mutations were associated with sub-ECOFF elevations in MIC for amikacin,

while being associated with elevations in MIC comparable to *rrs* mutations for

kanamycin. A deletion in *eis* leading to loss of function was also associated with

increased susceptibility to kanamycin, consistent with an epistatic interaction abrogating

the resistance gained from *eis* overproduction as shown in a prior study[39]. Variants in

aftB, *ccsA*, *whiB6* and *whiB7* were also associated with elevated MICs for at least one

aminoglycoside, however they were infrequent and require further investigation (Figure

5C and Table S3).

Ethionamide is a prodrug that is activated by the monooxygenases *ethA* and *mymA*

(*Rv3083*). More variants (128) were associated with increased ethionamide resistance

than any other drug, with the majority (103) occurring in *ethA*. Notably however, most

(97/103) MIC-elevating *ethA* variants did not raise the ethionamide MIC above the

ECOFF. Variants in *fabG1* and *inhA* were common and strongly associated with

elevated ethionamide MICs (Figure S2). Five resistance-associated variants were

identified in the alternative activating enzyme for ethionamide, *Rv3083*, and three

resistance-associated variants were found in the non-canonical ethionamide gene

mshA. Two mutations in *ethR we*re associated with decreased ethionamide MICs,

consistent with its role as a regulator of the prodrug activating enzyme *ethA*.

Resistance to delamanid is mediated by inactivating mutations in *ddn* or by mutations

that affect the cofactor F_{420} biosynthesis pathway (namely *fgd1* and *fbiA-D*). We

identified eleven mutations in *ddn*, seven in *fbiA*, and one in *fbiC* that were associated

with increases in delamanid MIC (Figure S2, Table S3). Over half (6/11) of the

mutations in *ddn* were nonsense or frameshift mutations suggesting that inactivation of

this gene is a common mechanism of resistance.

Effect of genetic background on MIC

Several studies have noted that the strain genetic background can influence MICs in

addition to primary resistance mutations[35,40,41]. In this study, we found that the effects of

lineage on isolate MIC tended to be small compared to primary resistance allele effects

for most drugs (mean lineage effect 0.41 log2MIC, mean lineage effect to median

primary resistance allele effect ratio 0.15), yet still statistically significant (Figure S3).

Notably however, lineage three was associated with a 1.5 lower moxifloxacin log2MIC

compared to lineage four after controlling for primary resistance alleles in *gyrA* and

gyrB.

Interactions beyond additivity

Given the high rate of co-occurrence of mutations in multiple target genes for a given

drug and additive resistance observed here and elsewhere, we sought to identify

whether there were any effects beyond additivity for co-occurring mutation pairs. Out of

33 pairs tested across 13 drugs, we identified three mutation pairs with greater than

additive effects on ethambutol resistance and one pair (rpoB_L430P:rpoB_D435G) with

greater than additive rifampicin resistance (Figure S4). The interaction of these

mutations resulted in log2MICs increased beyond additivity by 1.4 to 2.4 times, which

resulted in MICs well beyond that of the strongest individual mutations for ethambutol

and even greater than rpoB S450L for rifampicin. The remaining significant mutation-

pairs either consisted of a known resistance mutation with a putative compensatory

mutation (such as rpoB with rpoC) or had additive MICs that were in the tails of the

distribution, suggesting that interactions were reflecting assay thresholds, at least in

part, as opposed to true effects.

Extension beyond the binary 2021 WHO catalogue

In order to assess how measurement of MICs improves our ability to detect meaningful

genetic associations with resistance/susceptibility, we compared our MIC-based

catalogue with the recently published 2021 WHO binary catalogue for tuberculosis[16]

(Table S7). 179 unique mutation-phenotype associations were matched across the two

catalogues, with nearly a third (59/179) classified as "resistant – interim". Our model

finds that 61% (36/59) of these mutations are associated with significant elevations in

MIC in our data, of which 14 were sub-ECOFF and therefore unlikely to be confidently

identified by binary methods. The inability of binary methods to detect these smaller but

significant elevations in MIC is also shown by the lack of associations in *Rv0678* for

bedaquiline and clofazimine.

Discussion

In this study, we used WGS combined with high-throughput quantitative MIC

measurements to develop a quantitative catalogue capable of predicting resistance to

thirteen anti-tuberculosis drugs. This work highlights the use of genetics plus

measurement of a mutation's population mean MIC as a potentially more rapid and

reliable alternative to phenotypic DST for individual isolates, although further work to

verify this clinically is required. A further benefit is that measurement of MICs is robust

to future changes in critical concentration, streamlining and future-proofing use of data

in subsequent meta-analyses. These results can help to improve diagnostics and guide

future study designs trialing high dose therapies of less toxic and more effective drugs

(e.g. rifampicin, isoniazid and moxifloxacin)[10,11,24].

As shown by the MDR outbreak in Eswatini, tuberculosis treatment programs rely on

effective surveillance of drug-resistance and isolates that escape detection through

borderline MICs and/or noncanonical/rarer resistance mechanisms can lead to

outbreaks[24,25]. As well as directly affecting patients, this may also compromise the

efficacy of newer drugs such as bedaquiline that form the core of drug-resistant

tuberculosis treatment regimens. Here, we identify 321 mutations whose effects on MIC

are entirely or partially below their respective ECOFF. Further work is needed to

understand clinically whether these sub-ECOFF mutations lead to increased treatment

failure and/or relapse rates as is the case for the "borderline" mutations in *rpoB* for

rifampicin[27]. If so, rapid molecular assays and other sequencing-based technologies

should be employed to rapidly and accurately detect these variants.

In addition to highlighting variants associated with sub-ECOFF elevations in MICs that

may be missed using traditional binary methods, this work also highlights variants that

result in increased susceptibility to drugs such as bedaquiline, clofazimine, and the

aminoglycosides. Given the relative importance of *Rv0678* mutations in mediating

resistance to bedaquiline and clofazimine in clinical samples, rapid molecular tests

targeting these drugs should also consider incorporating *mmpL5* into their target gene

panels, as here we identify isolates with co-occurring nonsense and frameshift

mutations in *mmpL5* that override mutations in *Rv0678* and result in hypersensitivity to

bedaquiline through epistasis. A recent study showed that mmpS5 assembles with

mmpL5 in the membrane, raising the possibility that mutations in this gene may also

affect bedaquiline/clofazimine susceptibility; however, we failed to identify any

resistance-associated *mmpS5* mutations in this study[42]. The relatively common rate of

inactivating mutations in *mmpL5* we observed raises the intriguing potential of

optimizing therapies to include bedaquiline when the presence of one of these

susceptibility-enhancing mutations is identified. Indeed, previous work has shown that

deletion of the transcriptional regulators *Rv0324* and *Rv0880* also results in increased

susceptibility to bedaquiline, suggesting that other sensitizing mutations may also

occur[43]. Further work to understand the distribution and frequency of these mutations

may help elucidate their clinical relevance globally.

It is notable that despite the high frequency of co-occurring mutation-pairs, we only

identified four mutation pairs that resulted in resistance beyond additivity. One possible

reason is that, in many cases, the additive effects of two resistant mutations were well

beyond the highest MIC measured on the plate. This censoring likely limits our ability to

detect synergistic effects. However, it is also questionable whether accurate

measurement of these effects is even necessary, as in most cases the required drug

concentrations to overcome such synergy would be well beyond what is achievable in a

patient. In this study, we limited our investigation of epistasis to interactions between

significant variants within the target genes for each drug individually, although previous

studies have highlighted that mutations that effect different drugs can have impacts on

the growth rate and development of further resistance[44].

Using our expanded catalog, we were able to explain most binary resistance in

isoniazid, rifampicin, and ethambutol as well as the fluoroquinolones, aminoglycosides,

and rifabutin (Figure S5). This suggests that our target gene panels for these drugs

include almost all major genetic loci involved in resistance. However, ethionamide, as

well as the four new and repurposed drugs, all had comparatively low levels of

heritability in this and other studies, implying that many new variants remain to be

discovered[45]. Ethionamide appears to have a similar heritability profile to pyrazinamide,

in that many mutations in *ethA* have the potential to cause resistance, necessitating

high numbers of resistant isolates to be sequenced to achieve saturation of the

heritability curve. Given the poor heritability of resistance in our target gene panels for

the new and repurposed drugs, it is likely that novel genetic loci mediating resistance to

these drugs remain to be discovered. However, it is notable that high levels of

inactivating variation are observed in both *ddn* (delamanid) and *Rv0678*

(bedaquiline/clofazamine), which may suggest that high numbers of isolates will need to

be sampled to achieve saturation for these drugs as well. Further studies such as

GWAS may identify alternative loci, which may also become more apparent as these

drugs are employed more widely clinically.

Other approaches beyond clinical surveillance have proven advantageous for

understanding resistance determinants in drugs with high levels of variation such as

pyrazinamide, ethionamide, delamanid, and bedaquiline[46]. Random mutagenesis and

directed evolution have the advantage of offering high-throughput methods to screen

large numbers of isolates before they occur in a patient but may not always correlate

with the most common resistance mechanism *in vivo*[47,48]. Another complementary

approach to both *in vitro* and *in vivo* methods is machine learning, which has been

recently been employed to predict resistance to rifampicin, pyrazinamide, and

bedaquiline[49–52]. Importantly, methods that incorporate structure-based features are

capable of making predictions for mutations that have not been previously catalogued,

which is useful for genes with high variation like *pncA*, *ethA*, and *Rv0678*[53]. The

database generated by CRyPTIC can be used as a resource for both approaches, not

only by highlighting which mutations may actually be competitive in a patient, but also by acting as a training set for machine learning algorithms.

There are several limitations to this study. First, while extensive quality control was performed for both measurement of phenotypes and sequencing (Methods), some level of error is unavoidable in a study of this size. Another potential source of error is the exclusion of insertions/deletions greater than 50bp in size, which was required to limit computational complexity of joint variant calling but may result in falsely wild type calls in isolates with large structural variation. Additionally, our target gene approach, designed to limit multiple testing, introduces two problems: one, the inability to detect mutations in genes outside the target panel and two, the potential misattribution of effect to a passenger mutation inside the target genes that co-occurs with a bona fide resistance-conferring mutation elsewhere. We have attempted to limit this through controlling for lineage and population structure in our modelling approach as well as by validating mutations through structural mapping and degree of homoplasy where possible. Another limitation is the use of ECOFFs that have not been extensively validated against other methods, although we have shown good concordance with MGIT results[17]. Our study is also limited by the more limited numbers of isolates resistant to the newer and repurposed drugs, which limits power. Finally, while we investigated genetically-encoded determinants, changes in transcription or translation may also mediate antibiotic tolerance and persistence states, which also impact the efficacy of antibiotics *in vivo*[54].

1   In summary, here we directly quantify the impact of genetic resistance and susceptibility

2   determinants in tuberculosis to advance our understanding of resistance and serve as a

3   guide for further improvement and implementation of genetics-based diagnostic

4   resistance testing for tuberculosis, both through rapid molecular testing and whole-

5   genome sequencing[1].

1 **Figures and Tables**

2 **Table 1. Candidate genes used in this study.**

Drug	Abbrev.	Candidate genes
Isoniazid	INH	*katG, fabG1, inhA, ahpC, ndh, kasA, Rv1258c, Rv2752c*
Ethionamide	ETH	*ethA, ethR, fabG1, inhA, mshA, Rv3083*
Rifampicin	RIF	*rpoA, rpoB, rpoC, rpoZ, Rv2752c*
Rifabutin	RFB	*rpoA, rpoB, rpoC, rpoZ, Rv2752c*
Ethambutol	EMB	*embA, embB, embC, embR, rmlD, iniA, iniC, manB, ubiA*
Amikacin	AMI	*rrs, eis, ccsA, whiB6, whiB7, aftB, fprA*
Kanamycin	KAN	*rrs, eis, ccsA, whiB6, whiB7, aftB, fprA*
Levofloxacin	LEV	*gyrA, gyrB*
Moxifloxacin	MXF	*gyrA, gyrB*
Bedaquiline	BDQ	*atpE, Rv0678, mmpL5, mmpS5, pepQ, Rv3249c*
Clofazimine	CFZ	*Rv1979c, pepQ, Rv0678, mmpL5, mmpS5, Rv3249c*
Linezolid	LZD	*rplC, rrl, Rv3249c*
Delamanid	DLM	*ddn, fgd1, fbiA, fbiB, fbiC, fbiD, Rv3249c*

Figure 1: Data flow and sample sizes for CRyPTIC MIC models. Numbers in

brackets represent the number of variables (mutations plus lineage and site effects)

included in the final model.

1
2 **Figure 2: Variation in effect size by mutation type and drug. A)** Effects on log2MIC

3 for the 540 variants significant after false discovery rate correction using the Benjamini-

1 Hochberg method. Mutation types are delineated by color. Homoplastic mutations are

2 shown as solid circles. ECOFF is shown as tan line. Common resistance mutations are

3 highlighted. **B)** Comparison of effects on log2MIC for promoter and corresponding gene

4 body variants for ethambutol (EMB) and isoniazid (INH). ECOFF (minus baseline MIC)

5 is shown as a tan line.

Figure 3: Heterogenous effects of *rpoB* mutations on rifampicin resistance. A)

Effects of target gene variants on rifampicin (blue) and rifabutin (grey) log2MIC.

ECOFFs are highlighted as lines. **B)** Rifampicin (blue) and rifabutin (grey) bound to

rpoB with resistance-associated variants highlighted (red-high, orange-variable, yellow-

low). **C)** Effects on rifampicin MIC of mutations in rpoB. Text size is weighted by

frequency of occurrence. Colored shading highlights "borderline" variants.

Figure 4: Resistance to isoniazid and ethambutol is a multi-gene phenomenon. A)

Independent effects of variants in target genes on isoniazid log2MIC. ECOFF (minus

baseline MIC) is denoted in red. **B)** KatG dimer with isoniazid (blue) modeled and

resistance-associated positions highlighted in orange. **C)** Independent effects of

variants in target genes on ethambutol log2MIC. ECOFF (minus baseline MIC) is

denoted in red. **D)** EmbA-embB complex bound to ethambutol (blue) with resistant

mutations highlighted in orange.

Figure 5: Resistance to second line drugs. A) Effects of mutations in gyrA and gyrB

on levofloxacin (pink) and moxifloxacin (green) log2MIC. B) Structural mapping of

fluoroquinolone resistance-associated variants reveal that majority lie within 10Å of the

drug binding site. Positions gyrB R446 and S447 are not shown. C) Effects of mutations

in aminoglycoside target genes on amikacin and kanamycin log2MIC.

1
2 **Figure S1: Resistant promoter mutations are enriched near the -10 element.**

3 Promoter mutations are shown by position and colored by the mutant nucleotide.

4 Mutations with significant effects (p<0.05) after Benjamini Hochberg correction are

5 shown in solid circles. The -15 to -5 region is highlighted in tan.

6
7
8

Figure S2: Effects of common (n>10) and/or homoplastic mutations on 13 antituberculosis drugs. Homoplastic mutations are solid circles and ECOFFs are shown as tan lines.

Figure S3: Genetic background has variable influences on strain MIC by drug.
Significant effects of lineage on MIC compared with lineage 4 as reference. No
significant effects were identified for lineage 6.

1
Figure S4: Mutation pairs cause interaction beyond additivity in ethambutol and
rifampicin. Individual, predicted additive (black), and actual combined effects of
mutation pairs on log2MIC. Common single resistance mutations are provided for
reference.

1
2 **Figure S5: Heritability of binary phenotype for 13 antituberculosis drugs.** The
3 proportion of binary resistance explained by each additional mutation is shown by drug.
4 Curves asymptotically approaching horizontal indicate that few major mutations in
5 catalog genes remain to be described.
6

Methods:

Dataset collection

The CRyPTIC dataset collection and processing has been previously described in detail[17]. Briefly, clinical isolates were sub-cultured before inoculation into CRyPTIC-designed 96-well microtiter plates manufactured by ThermoFisher. Plates contained doubling-dilution ranges for 14 different antibiotics (para-aminosalicylic acid was excluded from the study due to poor-quality results on the plate). Isolate MICs were read after 14 days by a laboratory scientist using a Thermo Fisher Sensititre Vizion digital MIC viewing system and an image of the plate was also uploaded to a bespoke web server, allowing for additional MIC measurements by an automated computer vision system (AMYGDA) and by citizen science volunteers (Bash the Bug Zooniverse project) as previously described[55,56]. MIC measurements were classified as high (all three methods agree), medium (only two methods agree), or low (no methods agree). While sequencing processes differed slightly between CRyPTIC laboratories, all sequencing was performed using Illumina. A bespoke processing pipeline took in paired FASTQ files before filtering, mapping, and providing variant calls for each isolate. Isolates that had both phenotypic and whole-genome sequencing data were used as a starting dataset for this study (Clockwork availble from: https://github.com/iqbal-lab-org/clockwork, more detailed description of pipeline available in [16]). ECOFFs were defined elsewhere and are provided in Table S1[17].

Target gene selection

Target genes were selected based on the results of a prior study and through a

literature search for each drug[57].

Statistical modeling

All genetic variation smaller than 50bp occurring in the target genes for each drug

(Table S1) were included as candidates for effects in this study. Large insertions,

deletions, and other structural changes larger than 50bp were not included in this study.

Insertion/deletion mutations that occurred at the same position were pooled as one

candidate effect. Mutations that always co-occurred in the dataset were combined into

one candidate effect with all mutations named. Isolates were excluded from analysis if

they contained evidence for mixed alleles at positions previously associated with

resistance to that drug (i.e. a mixed allele call for position S450 in *rpoB* for rifampicin)[21].

Interval regression was performed in Stata version 16.1 with a genomic cluster variable

(cluster ID where clustering was based on the SNP distance calculated from the whole

genome sequence) as a random effect to control for population structure. A sensitivity

analysis was performed to compare the effects of clustering at 12, 25, 50, and 100

single nucleotide polymorphism (SNP) distances (100 used for all results shown).

Lineage and laboratory performing the MICs (SITEID variable) were included as factor

variables to control for genetic and technical variation in each individual drug model.

MICs were encoded as the interval with upper bound log2(MIC) and lower bound

log2(MIC minus 1 doubling dilution). The bottom and top wells were extended by 3

doubling dilutions to account for censoring.

The Benjamini-Hochberg correction was used to adjust raw p-values and the false

discovery rate was set at 5% for each drug based on the number of variants considered,

including all variants in one mutually adjusted multivariable model. Pairs of mutations

that occurred at least three times with each individual mutation occurring at least 5

times were subsequently tested for interactions in a mixed effect interval regression

model containing all other variants for that drug reaching the significance threshold

(Benjamini-Hochberg adjusted p-value < 0.05).

Data preparation, analysis and figure-making

Data was prepared for analysis using Python, statistical outputs were analyzed using R,

and figures were made using ggPlot2 in R[58]. Homoplasy was calculated by the number

of unique sub-lineages (predicted by SNP-IT and MYKROBE), with a mutation

considered homoplastic if it had evolved in at least 2 independent sublineages[18–20]. A

file that recapitulates all the post-model analysis and figures is available in the

Supplemental material (Supplemental Code). Structural modeling was done using

UCSF Chimera[59].

18
19

Author Contributions:

Members of the CRyPTIC consortium collected, phenotyped, and sequenced all isolates

in the CRyPTIC dataset. JC, ASW, and TMW designed this study, JC performed

statistical analyses and structural mapping, JC wrote the manuscript, JC, PWF, TEAP,

TMW, and ASW revised the manuscript with all partners providing feedback, and PWF,

TMW, ASW, and DWC supervised the work.

Acknowledgements – funders
8
This work was supported by Wellcome Trust/Newton Fund-MRC Collaborative Award
(200205/Z/15/Z); and Bill & Melinda Gates Foundation Trust (OPP1133541). Oxford CRyPTIC
consortium members are funded/supported by the National Institute for Health Research
(NIHR) Oxford Biomedical Research Centre (BRC), the views expressed are those of the authors
and not necessarily those of the NHS, the NIHR or the Department of Health, and the National
Institute for Health Research (NIHR) Health Protection Research Unit in Healthcare Associated
Infections and Antimicrobial Resistance, a partnership between Public Health England and the
University of Oxford, the views expressed are those of the authors and not necessarily those of
the NIHR, Public Health England or the Department of Health and Social Care. J.M. is supported
by the Wellcome Trust (203919/Z/16/Z). Z.Y. is supported by the National Science and
Technology Major Project, China Grant No. 2018ZX10103001. K.M.M. is supported by EMBL's
EIPOD3 programme funded by the European Union's Horizon 2020 research and innovation
programme under Marie Skłodowska Curie Actions. T.C.R. is funded in part by funding from
Unitaid Grant No. 2019-32-FIND MDR. R.S.O. is supported by FAPESP Grant No. 17/16082-7. L.F.
received financial support from FAPESP Grant No. 2012/51756-5. B.Z. is supported by the
National Natural Science Foundation of China (81991534) and the Beijing Municipal Science &
Technology Commission (Z201100005520041). N.T.T.T. is supported by the Wellcome Trust
International Intermediate Fellowship (206724/Z/17/Z). G.T. is funded by the Wellcome Trust.
R.W. is supported by the South African Medical Research Council. J.C. is supported by the
Rhodes Trust and Stanford Medical Scientist Training Program (T32 GM007365). A.L. is
supported by the National Institute for Health Research (NIHR) Health Protection Research Unit
in Respiratory Infections at Imperial College London. S.G.L. is supported by the Fonds de
Recherche en Santé du Québec. C.N. is funded by Wellcome Trust Grant No. 203583/Z/16/Z.
A.V.R. is supported by Research Foundation Flanders (FWO) under Grant No. G0F8316N (FWO
Odysseus). G.M. was supported by the Wellcome Trust (098316, 214321/Z/18/Z, and
203135/Z/16/Z), and the South African Research Chairs Initiative of the Department of Science
and Technology and National Research Foundation (NRF) of South Africa (Grant No. 64787). The
funders had no role in the study design, data collection, data analysis, data interpretation, or
writing of this report. The opinions, findings and conclusions expressed in this manuscript
reflect those of the authors alone. L.G. was supported by the Wellcome Trust (201470/Z/16/Z),

the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award number 1R01AI146338, the GOSH Charity (VC0921) and the GOSH/ICH Biomedical Research Centre (www.nihr.ac.uk). A.B. is funded by the NDM Prize Studentship from the Oxford Medical Research Council Doctoral Training Partnership and the Nuffield Department of Clinical Medicine. D.J.W. is supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (Grant No. 101237/Z/13/B) and by the Robertson Foundation. A.S.W. is an NIHR Senior Investigator. T.M.W. is a Wellcome Trust Clinical Career Development Fellow (214560/Z/18/Z). A.S.L. is supported by the Rhodes Trust. R.J.W. receives funding from the Francis Crick Institute which is supported by Wellcome Trust, (FC0010218), UKRI (FC0010218), and CRUK (FC0010218). T.C. has received grant funding and salary support from US NIH, CDC, USAID and Bill and Melinda Gates Foundation. The computational aspects of this research were supported by the Wellcome Trust Core Award Grant Number 203141/Z/16/Z and the NIHR Oxford BRC. Parts of the work were funded by the German Center of Infection Research (DZIF). The Scottish Mycobacteria Reference Laboratory is funded through National Services Scotland. The Wadsworth Center contributions were supported in part by Cooperative Agreement No. U60OE000103 funded by the Centers for Disease Control and Prevention through the Association of Public Health Laboratories and NIH/NIAID grant AI-117312. Additional support for sequencing and analysis was contributed by the Wadsworth Center Applied Genomic Technologies Core Facility and the Wadsworth Center Bioinformatics Core. SYNLAB Holding Germany GmbH for its direct and indirect support of research activities in the Institute of Microbiology and Laboratory Medicine Gauting. N.R. thanks the Programme National de Lutte contre la Tuberculose de Madagascar.

Competing Interest

E.R. is employed by Public Health England and holds an honorary contract with Imperial College London. I.F.L. is Director of the Scottish Mycobacteria Reference Laboratory. S.N. receives funding from German Center for Infection Research, Excellenz Cluster Precision Medicine in Chronic Inflammation, Leibniz Science Campus Evolutionary Medicine of the LUNG (EvoLUNG)tion EXC 2167. P.S. is a consultant at Genoscreen. T.R. is funded by NIH and DoD and receives salary support from the non-profit organization FIND. T.R. is a co-founder, board member and shareholder of Verus Diagnostics Inc, a company that was founded with the intent of developing diagnostic assays. Verus Diagnostics was not involved in any way with data collection, analysis or publication of the results. T.R. has not received any financial support from Verus Diagnostics. UCSD Conflict of Interest office has reviewed and approved T.R.'s role in Verus Diagnostics Inc. T.R. is a co-inventor of a provisional patent for a TB diagnostic assay (provisional patent #: 63/048.989). T.R. is a co-inventor on a patent associated with the processing of TB sequencing data (European Patent Application No. 14840432.0 & USSN 14/912,918). T.R. has agreed to "donate all present and future interest in and rights to royalties from this patent" to UCSD to ensure that he does not receive any financial benefits from this patent. S.S. is working and holding ESOPs at HaystackAnalytics Pvt. Ltd. (Product: Using whole genome sequencing for drug susceptibility testing for Mycobacterium tuberculosis).

Wellcome Trust Open Access

1
2 *This research was funded in part, by the Wellcome Trust/Newton Fund-MRC Collaborative*
3 *Award [200205/Z/15/Z]. For the purpose of Open Access, the author has applied a CC BY public*
4 *copyright licence to any Author Accepted Manuscript version arising from this submission.*
5
6 This research was funded, in part, by the Wellcome Trust [214321/Z/18/Z, and 203135/Z/16/Z].
7 For the purpose of open access, the author has applied a CC BY public copyright licence to any
8 Author Accepted Manuscript version arising from this submission.
9
10 **Acknowledgements – people**
11
12 JC would like to thank Spencer Dunleavy (Columbia Medical School, New York City, USA). We
13 thank Faisal Masood Khanzada and Alamdar Hussain Rizvi (NTRL, Islamabad, Pakistan), Angela
14 Starks and James Posey (Centers for Disease Control and Prevention, Atlanta, USA), and Juan
15 Carlos Toro and Solomon Ghebremichael (Public Health Agency of Sweden, Solna, Sweden).

16
17 **Ethics Statement**
18
Approval for CRyPTIC study was obtained by Taiwan Centers for Disease Control IRB No.
106209, University of KwaZulu Natal Biomedical Research Ethics Committee (UKZN BREC)
(reference BE022/13) and University of Liverpool Central University Research Ethics
Committees (reference 2286), Institutional Research Ethics Committee (IREC) of The
Foundation for Medical Research, Mumbai (Ref nos. FMR/IEC/TB/01a/2015 and
FMR/IEC/TB/01b/2015), Institutional Review Board of P.D. Hinduja Hospital and Medical
Research Centre, Mumbai (Ref no. 915-15-CR [MRC]), scientific committee of the Adolfo Lutz
Institute (CTC-IAL 47-J / 2017) and in the Ethics Committee (CAAE: 81452517.1.0000.0059) and
Ethics Committee review by Universidad Peruana Cayetano Heredia (Lima, Peru) and LSHTM
(London, UK).
29

Members of the CRyPTIC consortium (in alphabetical order)

Ivan Barilar[29], Simone Battaglia[1], Emanuele Borroni[1], Angela Pires Brandao[2,3], Alice Brankin[4],
Andrea Maurizio Cabibbe[1], Joshua Carter[5], Daniela Maria Cirillo[1], Pauline Claxton[6], David A
Clifton[4], Ted Cohen[7], Jorge Coronel[8], Derrick W Crook[4], Viola Dreyer[29], Sarah G Earle[4], Vincent
Escuyer[9], Lucilaine Ferrazoli[3], Philip W Fowler[4], George Fu Gao[10], Jennifer Gardy[11], Saheer
Gharbia[12], Kelen Teixeira Ghisi[3], Arash Ghodousi[1,13], Ana Luíza Gibertoni Cruz[4], Louis
Grandjean[33], Clara Grazian[14], Ramona Groenheit[44], Jennifer L Guthrie[15,16], Wencong He[10],
Harald Hoffmann[17,18], Sarah J Hoosdally[4], Martin Hunt[19,4], Zamin Iqbal[19], Nazir Ahmed Ismail[20],
Lisa Jarrett[21], Lavania Joseph[20], Ruwen Jou[22], Priti Kambli[23], Rukhsar Khot[23], Jeff Knaggs[19,4],
Anastasia Koch[24], Donna Kohlerschmidt[9], Samaneh Kouchaki[4,25], Alexander S Lachapelle[4], Ajit
Lalvani[26], Simon Grandjean Lapierre[27], Ian F Laurenson[6], Brice Letcher[19], Wan-Hsuan Lin[22],
Chunfa Liu[10], Dongxin Liu[10], Kerri M Malone[19], Ayan Mandal[28], Mikael Mansjö[44], Daniela
Matias[21], Graeme Meintjes[24], Flávia de Freitas Mendes[3], Matthias Merker[29], Marina Mihalic[18],
James Millard[30], Paolo Miotto[1], Nerges Mistry[28], David Moore[31,8], Kimberlee A Musser[9],
Dumisani Ngcamu[20], Hoang Ngoc Nhung[32], Stefan Niemann[29, 48], Kayzad Soli Nilgiriwala[28],

Camus Nimmo[33], Nana Okozi[20], Rosangela Siqueira Oliveira[3], Shaheed Vally Omar[20], Nicholas Paton[34], Timothy EA Peto[4], Juliana Maira Watanabe Pinhata[3], Sara Plesnik[18], Zully M Puyen[35], Marie Sylvianne Rabodoarivelo[36], Niaina Rakotosamimanana[36], Paola MV Rancoita[13], Priti Rathod[21], Esther Robinson[21], Gillian Rodger[4], Camilla Rodrigues[23], Timothy C Rodwell[37,38], Aysha Roohi[4], David Santos-Lazaro[35], Sanchi Shah[28], Thomas Andreas Kohl[29], Grace Smith[21,12], Walter Solano[8], Andrea Spitaleri[1,13], Philip Supply[39], Utkarsha Surve[23], Sabira Tahseen[40], Nguyen Thuy Thuong Thuong[32], Guy Thwaites[32,4], Katharina Todt[18], Alberto Trovato[1], Christian Utpatel[29], Annelies Van Rie[41], Srinivasan Vijay[42], Timothy M Walker[4,32], A Sarah Walker[4], Robin Warren[43], Jim Werngren[44], Maria Wijkander[44], Robert J Wilkinson[45,46,26], Daniel J Wilson[4], Penelope Wintringer[19], Yu-Xin Xiao[22], Yang Yang[4], Zhao Yanlin[10], Shen-Yuan Yao[20], Baoli Zhu[47]

Institutions
IRCCS San Raffaele Scientific Institute, Milan, Italy
Oswaldo Cruz Foundation, Rio de Janeiro, Brazil
Institute Adolfo Lutz, São Paulo, Brazil
University of Oxford, Oxford, UK
Stanford University School of Medicine, Stanford, USA
Scottish Mycobacteria Reference Laboratory, Edinburgh, UK
Yale School of Public Health, Yale, USA
Universidad Peruana Cayetano Heredia, Lima, Perú
Wadsworth Center, New York State Department of Health, Albany, USA
Chinese Center for Disease Control and Prevention, Beijing, China
Bill & Melinda Gates Foundation, Seattle, USA
UK Health Security Agency, London, UK
Vita-Salute San Raffaele University, Milan, Italy
University of New South Wales, Sydney, Australia
The University of British Columbia, Vancouver, Canada
Public Health Ontario, Toronto, Canada
SYNLAB Gauting, Munich, Germany
Institute of Microbiology and Laboratory Medicine, IMLred, WHO-SRL Gauting, Germany
EMBL-EBI, Hinxton, UK
National Institute for Communicable Diseases, Johannesburg, South Africa
Public Health England, Birmingham, UK
Taiwan Centers for Disease Control, Taipei, Taiwan
Hinduja Hospital, Mumbai, India
University of Cape Town, Cape Town, South Africa
University of Surrey, Guildford, UK
Imperial College, London, UK
Université de Montréal, Canada
The Foundation for Medical Research, Mumbai, India
Research Center Borstel, Borstel, Germany
Africa Health Research Institute, Durban, South Africa
London School of Hygiene and Tropical Medicine, London, UK
Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

33 University College London, London, UK
34 National University of Singapore, Singapore
35 Instituto Nacional de Salud, Lima, Perú
36 Institut Pasteur de Madagascar, Antananarivo, Madagascar
37 FIND, Geneva, Switzerland
38 University of California, San Diego, USA
39 Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL -
Center for Infection and Immunity of Lille, F-59000 Lille, France
40 National TB Reference Laboratory, National TB Control Program, Islamabad, Pakistan
41 University of Antwerp, Antwerp, Belgium
42 University of Edinburgh, Edinburgh, UK
43 Stellenbosch University, Cape Town, South Africa
44 Public Health Agency of Sweden, Solna, Sweden
45 Wellcome Centre for Infectious Diseases Research in Africa, Cape Town, South Africa
46 Francis Crick Institute, London, UK
47 Institute of Microbiology, Chinese Academy of Sciences, Beijing, China
48 German Center for Infection Research (DZIF), Hamburg-Lübeck-Borstel-Riems, Germany
18

References:

1   WHO. Global Tuberculosis Report 2020. 2020.

2   Schnippel K, Firnhaber C, Berhanu R, Page-Shipp L, Sinanovic E. Adverse drug
       reactions during drug-resistant TB treatment in high HIV prevalence settings: A
       systematic review and meta-analysis. *J Antimicrob Chemother* 2017; **72**: 1871–9.

3   André E, Goeminne L, Colmant A, Beckert P, Niemann S, Delmee M. Novel rapid
       PCR for the detection of Ile491Phe rpoB mutation of Mycobacterium tuberculosis,
       a rifampicin-resistance-conferring mutation undetected by commercial assays.
       Clin Microbiol Infect 2017; **23**: 267.e5-267.e7.

4   Cox H, Hughes J, Black J, Nicol MP. Precision medicine for drug-resistant
       tuberculosis in high-burden countries: is individualised treatment desirable and
       feasible? *Lancet Infect Dis* 2018; **18**: e282–7.

5   Pankhurst LJ, del Ojo Elias C, Votintseva AA, *et al.* Rapid, comprehensive, and
       affordable mycobacterial diagnosis with whole-genome sequencing: A prospective
       study. *Lancet Respir Med* 2016; **4**: 49–58.

6   Walker TM, Kohl TA, Omar S V., *et al.* Whole-genome sequencing for prediction
       of Mycobacterium tuberculosis drug susceptibility and resistance: A retrospective
       cohort study. *Lancet Infect Dis* 2015; **15**: 1193–202.

7   World Health Organization (WHO). Technical report on critical concentrations for
       TB drug susceptibility testing of medicines used in the treatment of drug-resistant
       TB. *Who* 2018; : 106.

8   Colangeli R, Jedrey H, Kim S, *et al.* Bacterial factors that predict relapse after
       tuberculosis therapy. *N Engl J Med* 2018; **379**: 823–33.

9   McCallum AD, Sloan DJ. The importance of clinical pharmacokinetic–

       pharmacodynamic studies in unraveling the determinants of early and late

       tuberculosis outcomes. *Int J Pharmacokinet* 2017; **2**: 195–212.

10   Grobbelaar M, Louw GE, Sampson SL, van Helden PD, Donald PR, Warren RM.

       Evolution of rifampicin treatment for tuberculosis. *Infect Genet Evol* 2019; **74**:

       103937.

11   Walsh KF, Vilbrun SC, Souroutzidis A, *et al.* Improved Outcomes with High-dose

       Isoniazid in Multidrug-resistant Tuberculosis Treatment in Haiti. *Clin Infect Dis*

       2019; **69**: 717–9.

12   Decroo T, de Jong BC, Piubello A, Souleymane MB, Lynen L, van Deun A. High-

       dose first-line treatment regimen for recurrent rifampicin-susceptible tuberculosis.

       Am J Respir Crit Care Med 2020; **201**: 1578–9.

13   Farhat MR, Freschi L, Calderon R, *et al.* GWAS for quantitative resistance

       phenotypes in Mycobacterium tuberculosis reveals resistance genes and

       regulatory regions. *Nat Commun* 2019; **10**. DOI:10.1038/s41467-019-10110-6.

14   Rancoita PMV, Cugnata F, Gibertoni Cruz AL, *et al.* Validating a 14-drug

       microtiter plate containing bedaquiline and delamanid for large-scale research

       susceptibility testing of mycobacterium tuberculosis. *Antimicrob Agents*

       Chemother 2018. DOI:10.1128/AAC.00344-18.

15   Falzon D, Schünemann HJ, Harausz E, *et al.* World Health Organization

       treatment guidelines for drug-resistant tuberculosis, 2016 update. *Eur Respir J*

       2017; **49**. DOI:10.1183/13993003.02308-2016.

16   WHO. Catalogue of mutations in Mycobacterium tuberculosis complex and their

association with drug resistance. 2021.

The CRyPTIC Consortium. Epidemiological cutoffs for a 96-well broth microtitre plate for high-throughput research antibiotic susceptibility testing of M . tuberculosis. *medRxiv doi:101101/2021022421252386* 2021. DOI:10.1101/2021.02.24.21252386.

Lipworth S, Jajou R, De Neeling A, *et al.* SNP-IT tool for identifying subspecies and associated lineages of Mycobacterium tuberculosis complex. *Emerg Infect Dis* 2019; **25**: 482–8.

Bradley P, Gordon NC, Walker TM, *et al.* Rapid antibiotic-resistance predictions from genome sequence data for Staphylococcus aureus and Mycobacterium tuberculosis. *Nat Commun* 2015; **6**: 1–14.

Hunt M, Bradley P, Lapierre SG, *et al.* Antibiotic resistance prediction for Mycobacterium tuberculosis from genome sequence data with Mykrobe. *Wellcome Open Res* 2019; **4**: 191.

The CRyPTIC Consortium, The 100000 Genomes Project. Prediction of Susceptibility to First-Line Tuberculosis Drugs by DNA Sequencing. *N Engl J Med* 2018; **379**: 1403–15.

Newton-Foot M, Gey Van Pittius NC. The complex architecture of mycobacterial promoters. *Tuberculosis* 2013; **93**: 60–74.

Zhu DX, Garner AL, Galburt EA, Stallings CL. CarD contributes to diverse gene expression outcomes throughout the genome of Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A* 2019; **116**: 13573–81.

Makhado NA, Matabane E, Faccin M, *et al.* Outbreak of multidrug-resistant

tuberculosis in South Africa undetected by WHO-endorsed commercial tests: an observational study. *Lancet Infect Dis* 2018. DOI:10.1016/S1473-3099(18)30496-1.

25 Beckert P, Sanchez-Padilla P, Merker M, Dreyer V, Kohl TA, Utpatel C, Köser CU, Barilar I, Ismail N, Omar SV, Klopper M, Warren RM; Hoffmann H, Maphalala G, Ardizzoni E, de Jong BC, Kerschberger B, Schramm B, Andres S, Kranzer K, Maurer FP, Bonnet M NS. MDR M. tuberculosis outbreak clone in Eswatini missed by Xpert has elevated bedaquiline resistance dated to the pre-treatment era. *Genome Med* 2020; **12**.

26 Brandis G, Hughes D. Mechanisms of fitness cost reduction for rifampicin-resistant strains with deletion or duplication mutations in rpoB. *Sci Rep* 2018; **8**: 1–6.

27 World Health Organization. Technical Report on critical concentrations for drug susceptibility testing of isoniazid and the rifamycins (rifampicin, rifabutin and rifapentine). 2021 https://www.who.int/publications/i/item/technical-report-on-critical-concentrations-for-drugsusceptibility-testing-of-isoniazid-and-therifamycins-(rifampicin-rifabutin-and-rifapentine).

28 Torrea G, Ng KCS, Van Deun A, *et al.* Variable ability of rapid tests to detect Mycobacterium tuberculosis rpoB mutations conferring phenotypically occult rifampicin resistance. *Sci Rep* 2019; **9**: 1–9.

29 Miotto P, Cabibbe AM, Borroni E, Degano M, Cirilloa DM. Role of disputed mutations in the rpoB gene in interpretation of automated liquid MGIT culture results for rifampin susceptibility testing of mycobacterium tuberculosis. *J Clin*

Microbiol 2018; **56**: 1–9.

Jeong DH, Kang YW, Kim JY, Han JS, Jo KW, Shim TS. Successful treatment with a high-dose rifampin-containing regimen for pulmonary tuberculosis with a disputed rpob mutation. *Intern Med* 2018; **57**: 3281–4.

Ma P, Luo T, Ge L, *et al.* Compensatory effects of M. tuberculosis rpoB mutations outside the rifampicin resistance-determining region. *Emerg Microbes Infect* 2021; **10**: 743–52.

Loewen PC, Switala J, Smolenski M, Triggs-Raine BL. Molecular characterization of three mutations in katG affecting the activity of hydroperoxidase I of Escherichia coli. *Biochem Cell Biol* 1990; **68**: 1037–44.

Munir A, Wilson MT, Hardwick SW, *et al.* Using cryo-EM to understand antimycobacterial resistance in the catalase-peroxidase (KatG) from Mycobacterium tuberculosis. *Structure* 2020; **29**: 1–14.

Hicks ND, Yangg J, Zhang X, *et al.* Clinically prevalent mutations in Mycobacterium tuberculosis alter propionate metabolism and mediate multidrug tolerance. *Physiol Behav* 2019; **176**: 139–48.

Safi H, Lingaraju S, Amin A, *et al.* Evolution of high-level ethambutol-resistant tuberculosis through interacting mutations in decaprenylphosphoryl-β-D-Arabinose biosynthetic and utilization pathway genes. *Nat Genet* 2013; **45**: 1190–7.

Zhang L, Zhao Y, Gao Y, *et al.* Structures of cell wall arabinosyltransferases with the anti-tuberculosis drug ethambutol. *Science (80-)* 2020; **9102**: eaba9102.

Disratthakit A, Prammananan T, Tribuddharat C, *et al.* Role of gyrB mutations in

pre-extensively and extensively drug-resistant tuberculosis in Thai clinical

isolates. *Antimicrob Agents Chemother* 2016; **60**: 5189–97.

38 Maruri F, Sterling TR, Kaiga AW, *et al.* A systematic review of gyrase mutations

associated with fluoroquinolone-resistant Mycobacterium tuberculosis and a

proposed gyrase numbering system. *J Antimicrob Chemother* 2012; **67**: 819–31.

39 Vargas Jr R, Freschi L, Spitaleri A, *et al.* The role of epistasis in amikacin,

kanamycin, bedaquiline, and clofazimine resistance in Mycobacterium

tuberculosis complex. *bioRxiv* 2020.

40 Farhat MR, Sixsmith J, Calderon R, Hicks ND, Fortune SM, Murray M. Rifampicin

and rifabutin resistance in 1003 Mycobacterium tuberculosis clinical isolates. *J

Antimicrob Chemother* 2019; **74**: 1477–83.

41 Nebenzahl-Guimaraes H, Jacobson KR, Farhat MR, Murray MB. Systematic

review of allelic exchange experiments aimed at identifying mutations that confer

drug resistance in Mycobacterium tuberculosis. *J Antimicrob Chemother* 2014; **69**:

331–42.

42 Yamamoto K, Nakata N, Mukai T, Kawagishi I, Ato M. Coexpression of MmpS5

and MmpL5 Contributes to Both Ef fl ux Transporter MmpL5 Trimerization and

Drug Resistance in Mycobacterium tuberculosis. 2021; **6**: 1–11.

43 Peterson EJR, Ma S, Sherman DR, Baliga NS. Network analysis identifies

Rv0324 and Rv0880 as regulators of bedaquiline tolerance in Mycobacterium

tuberculosis. *Nat Microbiol* 2016; **1**: 6–12.

44 Borrell S, Teo Y, Giardina F, *et al.* Epistasis between antibiotic resistance

mutations drives the evolution of extensively drug-resistant tuberculosis. *Evol Med*

Public Heal 2013; **2013**: 65–74.

Kadura S, King N, Nakhoul M, *et al.* Systematic review of mutations associated with resistance to the new and repurposed Mycobacterium tuberculosis drugs bedaquiline , clofazimine , linezolid , delamanid and pretomanid. *J Antimicrob Chemother* 2020; : 2031–43.

Lee BM, Harold LK, Almeida D V., *et al.* Predicting nitroimidazole antibiotic resistance mutations in Mycobacterium tuberculosis with protein engineering. *PLoS Pathog* 2020; **16**: 1–27.

Sonnenkalb L, Carter J, Spitaleri A, *et al.* Deciphering Bedaquiline and Clofazimine Resistance in Tuberculosis : An Evolutionary Medicine Approach. *bioRxiv doi:101101/20210319436148* 2021; : 1–38.

Yadon AN, Maharaj K, Adamson JH, *et al.* A comprehensive characterization of PncA polymorphisms that confer resistance to pyrazinamide. *Nat Commun* 2017; **8**: 1–10.

Portelli S, Myung Y, Furnham N, Vedithi SC, Pires DEV, Ascher DB. Prediction of rifampicin resistance beyond the RRDR using structure-based machine learning approaches. *Sci Rep* 2020; **10**: 1–13.

Carter JJ, Walker TM, Walker AS, *et al.* Prediction of pyrazinamide resistance in Mycobacterium tuberculosis using structure-based machine learning approaches. bioRxiv. 2019. DOI:10.1101/518142.

Karmakar M, Rodrigues CHM, Horan K, Denholm JT, Ascher DB. Structure guided prediction of Pyrazinamide resistance mutations in pncA. *Sci Rep* 2020; **10**: 1–10.

Karmakar M, Rodrigues CHM, Holt KE, Dunstan SJ, Denholm J, Ascher DB. Empirical ways to identify novel Bedaquiline resistance mutations in AtpE. *PLoS One* 2019; **14**: 1–14.

Battaglia S, Spitaleri A, Cabibbe AM, *et al.* Characterization of genomic variants associated with resistance to bedaquiline and delamanid in naïve Mycobacterium tuberculosis clinical strains. *J Clin Microbiol* 2020; **58**: 1–16.

Harms A, Maisonneuve E, Gerdes K. Mechanisms of bacterial persistence during stress and antibiotic exposure. *Science (80-)* 2016; **354**. DOI:10.1126/science.aaf4268.

Rancoita PM V., Cugnata F, Gibertoni Cruz AL, *et al.* Validating a 14-drug microtitre plate containing bedaquiline and delamanid for large-scale research susceptibility testing of *Mycobacterium tuberculosis. Antimicrob Agents Chemother* 2018; : AAC.00344-18.

Fowler PW, Gibertoni Cruz AL, Hoosdally SJ, *et al.* Automated detection of bacterial growth on 96-well plates for high-throughput drug susceptibility testing of Mycobacterium tuberculosis. *Microbiology* 2018. DOI:10.1099/mic.0.000733.

Prediction of Susceptibility to First-Line Tuberculosis Drugs by DNA Sequencing. *N Engl J Med* 2018; **379**: 1403–15.

Wickam H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York, 2016 https://ggplot2.tidyverse.org.

Pettersen EF, Goddard TD, Huang CC, *et al.* UCSF Chimera - A visualization system for exploratory research and analysis. *J Comput Chem* 2004; **25**: 1605–12.