Bacterial composition reflects fine-scale salinity changes while phylogenetic diversity

exhibits a strong salt divide

Ariane L. Peralta[1*], Mario E. Muscarella[2], Alexandra Stucy[1], Jo A. Werba[1,3], Michael W.

McCoy[1,4]

[1]Department of Biology, East Carolina University, Howell Science Complex, Greenville, NC

27858, USA; [2]Department of Biology and Wildlife, Institute of Arctic Biology, University of Alaska

Fairbanks, Fairbanks, AK, 99775, USA; [3]Department of Biology, McMaster University, 1280

Main Street West, Hamilton, Ontario, Canada, L8S 4K1; [4]Harbor Branch Oceanographic

Institute, Florida Atlantic University, Fort Pierce, FL, 34946, USA

*corresponding author

ABSTRACT

Climate change induced salinization events are predicted to intensify and lead to increased salt stress in freshwater aquatic ecosystems. As a consequence, formerly distinct abiotic conditions and associated biotic communities merge, and the emergence, loss, and persistence of microbial taxa modify the types and rates of ecosystem processes. This study examined how bacterial taxonomic and phylogenetic diversity and ecosystem function respond to acute salinization events where freshwater and estuarine communities and environments coalesce. We hypothesize that if the salinity change outpaces microbial adaptation or saline microbial populations are not yet established in formerly freshwater conditions, then we predict diminished carbon cycling rates, decreased microbial diversity, and altered the composition of microbial communities compared to historically freshwater communities. We used an experimental mesocosm approach to determine how salinity and the merging of distinct communities influenced resultant bacterial community structure and function. Each mesocosm represented different salinities (0, 5, 9, 13 psu). Two dispersal treatments, representing aquatic communities sourced from brackish 13 psu ponds and a mix of 13 psu and freshwater ponds, were added to all salinity levels and replicated four times. Results revealed that salinity, but not dispersal, decreased bacterial taxonomic and phylogenetic diversity. Carbon mineralization rates were highest in freshwater conditions and associated with low relative abundance indicator taxa. Acute salinity changes, such as localized flooding due to storm surge, will more negatively affect freshwater aquatic communities compared to chronic exposure to salinization where the communities have had time to adapt or turnover resulting in recovered biogeochemical functions.

IMPORTANCE STATEMENT

Climate change induced salinization results in the mixing of formerly distinct environmental

conditions and aquatic communities. This study examined the consequence of short-term, acute

salinity stress on aquatic bacterial taxonomic and phylogenetic diversity and ecosystem function

using an experimental approach. Results revealed that salinity, but not the source of aquatic

communities, decreased bacterial taxonomic and phylogenetic diversity. Carbon mineralization

rates, which represented ecosystem function, were highest in freshwater conditions and also

associated with low relative abundance indicator bacterial taxa. Taken together, acute salinity

changes will more negatively affect freshwater aquatic communities compared to chronic

exposure to salinization where the communities have had time to adapt or turnover resulting in

recovered biogeochemical functions.

INTRODUCTION

Predicting microbial community response to environmental change depends on the tolerance and preference of microbial taxa to current and historical environmental conditions, ongoing changes in environmental conditions, and concurrent community assembly processes (1–3). Environmental changes, such as sea level rise and punctuated salinization events, are predicted to intensify in the coming decades as land development and climate change continue (4). Sea-level rise and saltwater intrusion in previously freshwater wetlands can diminish wetland ecosystem functions by increasing toxic sulfide production, decreasing inorganic nitrogen removal, and decreasing carbon storage. In some cases, diminished microbial functions negatively affect water quality, climate regulation, wetland accretion, and the health of wetland biota (5, 6). However, it is unclear how the composition and function of microbial communities at newly established fresh-salt water interfaces will change as salinization events increase in frequency and duration, especially with regard to subsets of taxa that respond to the coalescence event (i,e,, the merging of distinct microbial communities and environments) (7, 8). As aquatic ecosystems shift from fresh to brackish waters, the concentration of alternative terminal electron acceptors (i.e., Fe(III), Mn(IV)) increases, which prompts CO_2 production (6). As a result, environmental stressors are altering the microbial communities governing these carbon cycling rates.

Salinity tolerance has convergently evolved in unrelated lineages across the tree of life (9, 10). Salinity preference and osmotic tolerance are considered a complex microbial trait (11). Two mechanisms are responsible for osmotic or salinity stress tolerance in microorganisms. First, microorganisms can adapt by excluding harmful solutes (e.g., sodium, chloride) while gathering beneficial solutes necessary for metabolism instead by using active transport systems (12). Second, some microorganisms can create organic compounds to reduce the concentration gradient between the external environment and the cell resulting in the net export of unwanted solutes (13–15). Some studies suggest that long-term salinity changes, like those due to sea level

rise, may occur slowly enough that the organisms can adapt and/or communities reassemble

according to environmental changes. Long-term, incremental salinity increases have promoted

microbial diversity and community establishment of salt-preference in a variety of taxa.

Salinization in tidal and non-tidal systems results in different community responses. For example,

tidal wetland decomposition rates increase as salinity increases due in part to marine/estuarine

subsidies mixing with freshwater ecosystems (16). While experiencing short-term acute

salinization, microbial populations have recovered from previously diminished carbon cycling

functions (carbon dioxide and methane production) (17). Prior studies have revealed both

increases in decomposition rates (15, 18) and decreases in decomposition rates (Neubauer et al.

2013, 2019) as a result of increases in salinity. Punctuated salinity pulses to historically freshwater

ecosystems, such as localized flooding due to storm surge in a climatic event, will more negatively

affect freshwater communities compared to constant exposure to salinization. The challenge to

predicting salinization effects on decomposition rates is due to interactions among the following:

direct effects of redox shifts between oxic and anoxic conditions, direct effects on heterotrophic

communities and local nutrient availability, and the indirect effects due to changes in organic

carbon sources (21).

Salinity tolerance and preference are traits that constrain microbial community response to

salinization (22). As freshwater salinization across inland freshwaters persists, drastic changes to

microbial community structure and function will continue to increase (23). The emergence, loss,

and persistence of microbial taxa can alter the types and rates of ecosystem functions. The

microbial taxa that will survive punctuated salinity changes will have specialized traits. These

'effect' traits can be directly linked to ecosystem functioning (11). Salinity preference is identified

as one of the most complex trait measurements, classifying it based on how many genes are

directly involved in coding the trait and how the trait is integrated with other mechanisms (11).

Because specialized traits are needed for organisms to tolerate increasingly saline environments,

major shifts in microbial composition and diversity are expected (24, 25). When freshwater and

marine microbes are experimentally mixed, results revealed that taxa which were rare at the initial

inoculation became relatively abundant (8). As the community shifts, organisms that are better

adapted to new conditions may become more active and abundant, which can alter the dominant

ecosystem processes (1, 3).

In this study, we address the research question: how does acute salinization affect microbial

taxonomic and phylogenetic diversity and function when freshwater microbial communities mix

with estuarine aquatic communities along freshwater to saltwater gradient? Specialized microbial

traits are necessary to tolerate or thrive in saline environments. We hypothesize that if salinity

change outpaces microbial adaptation or saline microbial populations are not yet established in

formerly freshwater conditions, then we predict diminished carbon cycling rates (i.e., CO_2

respiration) and decreased microbial diversity and altered the composition of microbial

communities when compared to historically freshwater microbial communities. We also expect

that microorganisms that persist in more saline conditions will be more phylogenetically related to

each other (i.e., less genetic variation) compared to historically freshwater microorganisms.

Understanding how salinization alters freshwater wetland bacterial phylogenetic-ecosystem

function relationships can inform the management of carbon storage capacity in coastal wetlands

experiencing increased salinization.

MATERIALS AND METHODS

Experimental Design. Microbial community composition and function were characterized based

on aquatic samples that were collected from a replicated mesocosm experiment conducted from

June-July 2015 (Figure 1) (26). To replicate a freshwater-saltwater aquatic system, peat moss (to

serve as a nutrient pulse), sand (to serve as a benthic substrate), and water were added to 150-

gallon stock watering tanks (filled to 100 gallons). Each mesocosm pond was adjusted to one of

4 different salinities (measured in practical salinity units) using Instant Ocean® Sea Salt (4 levels:

0, 5, 9, 13 psu). Each tank was then seeded with an inoculation consisting of water, zooplankton

and microbes collected from natural ponds in the outer banks of North Carolina (see

Supplemental Table S1 for coastal pond location and information). Twenty 1 L water samples

were collected and filtered through 62.5 μm mesh across a one-hundred meter transect at each

of the 5 coastal ponds and subsequently dispersed into each mesocosm tank. Bacterial

communities represented in the source communities were likely particle- and zooplankton-

associated. Each tank was covered with a shade cloth to reduce the opportunity for other

organisms from colonizing the mesocosms. Populations introduced via the initial inoculation were

allowed to stabilize for 6 weeks before sampling began. After this acclimation period, mesocosms

received additional colonists of zooplankton and microbes via simulated dispersal events from

separate mesocosm ponds established as source tanks for salt (13 psu) and freshwater (0 psu)

zooplankton and microbial communities. The two source tanks served as dispersal treatments,

and consisted of either a saltwater (13 psu; salt dispersal treatment) community or a mix of

communities from freshwater (0 psu) and saltwater (13 psu; fresh+salt dispersal treatment).

Dispersal treatments consisted of 2 L of water from dispersal (source) tanks into the experiment

(treatment) tanks. Sample collection occurred every 9 days (due to average time for completion

of one zooplankton generation cycle) (27) over the 6-week mesocosm experiment for a total of

six time points. The salinity × dispersal setup was replicated 4 times to generate 16 tanks (4 at

each salinity level) with the salt dispersal treatment or with the fresh+salt dispersal treatment (26).

For each tank, the following were measured using a YSI Pro: dissolved oxygen, ammonium

(NH_4^+), temperature, and pH. Community analysis was focused on three of the time points (days

0, 18, and 45; 11 June 2015, 29 June 2015, and 25 July 2015, respectively), representing the

initial, middle, and end of the experiment. Besides salinity levels, measured nitrogen and

phosphorus concentrations were similar among treatment tanks (26).

Microbial Sample Collection and Processing. Sample collection occurred every 9 days over

the time of the 6-week mesocosm experiment for a total of six time points. During each sampling

event, we collected 1 L of water from each tank. Each 1 L bottle of mesocosm water was

homogenized and 200 mL of the water sample were concentrated onto 0.22 µm filters within 24

hours of field sampling (Supor-200; Pall Gelman, East Hills, NY) to collect microbial community

samples**.** Filters were transferred into 2 mL sterile tubes and stored at -80^{o} C until molecular

analyses. We extracted DNA from filters collected at three (of the six) time points representing

the initial, middle, final sampling dates (Days 0, 18, 45) using the PowerWater DNA Isolation Kit

(MO BIO Laboratories).

Microbial Community Sequencing. To examine shifts in microbial community composition and

diversity, aquatic microbial communities in each mesocosm were characterized using paired-end

targeted Illumina sequencing of the 16S rRNA gene (bacteria, archaea) (28). We extracted and

purified the DNA found in the water of each mesocosm using the PowerWater DNA Isolation Kit

(MO BIO Laboratories, Inc., Carlsbad, CA) using the sample collected a 0.22 µm supor filter post-

filtration. We used this DNA as a template in PCR reactions. To characterize bacterial

communities, we used barcoded primers (515FB/806RB) originally developed by the Earth

Microbiome Project (28) to target the V4-V5 region of the bacterial 16S subunit of the ribosomal

RNA gene (29, 30). PCR products were combined in equimolar concentrations and sequenced

using paired-end (2×250 bp) approach using the Illumina MiSeq platform at Indiana University's

Center for Genomics and Bioinformatics.

Raw bacterial sequences were processed using the Mothur pipeline (version 1.41.3) (31).

Briefly, paired-end contigs were assembled and quality trimmed, sequences were then aligned to

the Silva Database (version 132) (32) chimeric sequences were removed using the VSEARCH

algorithm (33), and we removed any sequences classified as Chloroplasts, Mitochondria,

Archaea, or Eukaryotes. Next, we identified operational taxonomic units (OTUs) by separating

sequences based on taxonomic class and then binning sequences using a 97% sequence

similarity cutoff. Representative sequences from each OTU were used to generate a phylogenetic

tree using FastTree v2.1 using the GTR model and CAT rate approximation (Price at al. 2010).

See Supplemental Material for full sequence processing pipeline.

Compositional Analyses. We used a combination of taxonomic and phylogenetic approaches

to characterize the diversity of bacterial communities in each mesocosm. First, we calculated

alpha diversity (within-sample) using either Shannon's diversity (taxonomic) or Faith's Diversity

(phylogenetic). The Faith's Phylogenetic Diversity (PD) metric provides a simple measure of the

phylogenetic relatedness of a community based on the summed branch lengths of its phylogenetic

tree. We expected this measure to capture functional complementarity if more distantly related

species are more functionally unique. We rarefied the OTU table to 20,000 observations before

calculating alpha-diversity. Next, we calculated beta-diversity (among sites) by calculating Bray-

Curtis distance for taxonomic diversity and weighted UniFrac distance for phylogenetic diversity.

Differences in beta-diversity were visualized using Principal Coordinates Analysis.

Carbon Mineralization Assay. On the final mesocosm sampling date, day 45, we measured the

amount of CO_2 respired from the aquatic communities using a laboratory-based bottle assay.

Wheaton bottles (125 mL) fitted with septa were filled with water samples (25 mL) from each

mesocosm tank. The CO_2 concentration readings were determined using an LI-7000 Infrared Gas

Analyzer (IRGA). On the first day (Day 0), bottles were filled with 25 mL of mesocosm tank water,

and the gas samples were collected and analyzed immediately using the IRGA to determine the

baseline CO_2 concentration. A syringe was inserted into the septa and the headspace gas was

mixed 3 times before pulling a sample and beginning analysis using the IRGA. This process was

repeated on days one, three, and seven in order to determine CO_2 respiration rates over time.

Statistical Analyses. We completed statistical calculations in the R environment (R v3.6.3, R

Core Development Team 2020). For bacterial diversity metrics, we conducted a linear mixed

effects model with 'date', 'salinity', and 'dispersal' as fixed effects and 'block' as a random effect using the *lmer()* function in the lmerTest package (Kuznetsova et al. 2017) and the pbkrtest package (Halekoh and Højsgaard 2014). We ran linear mixed models that were fit by REML and produced type II analysis of variances tables (ANOVA) tables based on the Kenward-Roger's denominator degrees of freedom method using the anova() function. Then, we used principal coordinates analysis (PCoA) based on the Bray-Curtis dissimilarity of bacterial community composition and based on the weighted Unifrac distance of phylogenetic diversity along the salinity gradient. We ran a permutational multivariate analysis of variance (PERMANOVA) to examine among-treatment differences in bacterial communities. We identified which bacterial species were most representative of each salinity treatment using an indicator species analysis based on salinity level and also a categorical saline vs non-saline classification, and we included bacterial taxa with a relative abundance greater than 0.05 when summed across all plots. We performed PERMANOVA using the *adonis()* function in the vegan package (Oksanen et al. 2017) and the indval() function in the indicspecies package (Caceres and Jansen 2016). We visualized overlap in indicator taxa across the salinity gradient using an Euler plot. To better understand the phylogenetic underpinnings of saline tolerant and sensitive taxa, we visualized the indicators on a phylogenetic tree and used the ConcenTrait algorithm to determine the phylogenetic depth at which saline tolerance/sensitivity is conserved (34). Briefly, this algorithm finds the nodes in the phylogeny at which 90% of the included tips (taxa) share the same trait. We implemented the ConcenTrait algorithm using a custom R script.

Data Availability

All code and data used in this study are located in a public GitHub repository (https://github.com/PeraltaLab/CSI_Dispersal) and NCBI SRA BioProject ID PRJNA615001.

RESULTS

We examined bacterial community turnover throughout the mesocosm experiment by comparing patterns in bacterial phylogenetic and taxonomic composition across the salinity gradient. Throughout the experiment, freshwater communities were phylogenetically distinct compared to saline communities based on weighted UniFrac distance metric (Fig. 2A, Table S2, PERMANOVA salinity: R^2=0.172, P<0.001). Bacterial communities were phylogenetically similar during day 18, but after 45 days at the experimental salinity level, communities at 9 and 13 psu were distinct from communities at 5 psu (Fig. 1A, Table S2, PERMANOVA salinity x date: R^2=0.048, P<0.001). When bacterial taxonomic composition was compared, salinity was a much stronger environmental filter associated with bacterial community patterns (Werba et a. 2020) (Fig. 2B).

Next, we compared patterns in bacterial phylogenetic diversity to taxonomic composition across the salinity gradient. Communities observed in the freshwater conditions increased in mean PD over the 45 day experiment. The PD of communities at the transition 9 psu tanks declined throughout the experiment duration, while PD of communities in the 13 psu tanks increased in variability from day 0 to 45 (Fig. 2C, Table S3, ANOVA, $F_{2,79}$=5.997, P=0.004). After 45 days, PD was highest in freshwater and lowest in brackish tanks (Fig. 2C).

There are OTUs represented across major phyla that are associated with all salinity levels (Fig. 3A). A striking observation was the smallest number of OTUs represented at 9 psu compared to other treatments, which is represented in the unrooted tree (Fig. 3A). Next, we used a series of Euler plots to reveal shared and unique indicator OTUs. A list of "indicators" that were significantly associated with each salinity treatment is found in Table S3. These indicators, reported at the class-level, are the taxonomic groups best associated with each treatment. Based on OTUs identified as indicator taxa, 254 OTUs are unique to the freshwater treatment, 135 OTUs are unique to the 5 psu treatment, 35 OTUs were unique to the 9 psu treatment, and 124 OTUs are unique to the 13 psu treatment (Fig. 3B). When the OTUs were grouped into salt or no salt,

there are 206 observed indicator OTUs shared across 5, 9, or 13 psu environments, and 257

OTUs unique to the 0 psu treatment (Fig. 3B). Finally, when examining the depth of phylogenetic

relatedness, results from the ConcenTrait analysis revealed that saline vs non-saline microbes

exhibit a similar depth in average phylogenetic relatedness (Fig. 3C).

We examined the relationship between carbon mineralization and phylogenetic diversity and

relative abundance. Dispersal treatment did not affect the relationship between carbon

mineralization rates and taxonomic or phylogenetic diversity (Fig. 4A). At the transition 9 psu

treatments, we measured the lowest measured function compared to other salinity treatments.

Carbon mineralization rates were highest in freshwater conditions and associated relatively low

abundance indicator taxa (Fig. 4B). In contrast, the relative abundance of indicator taxa

represented in the saltwater conditions represented > 25% of the community.

DISCUSSION

This study specifically focused on determining the extent that short-term salinization events

influenced reassembly patterns of aquatic communities and consequent cycling function. Results

revealed that salinity strongly influenced bacterial phylogenetic diversity; bacterial communities

tended to respond to salinity, rather than dispersal, because it is such a strong environmental filter

(8, 10). Past research has shown that strong environmental filters produce phylogenetically

clustered communities (11, 35–37). However, the mixing of two formerly distinct communities and

their environments could affect community structure and function in unpredictable ways.

Outcomes depend on the strength of the environmental change and the capacity for taxa to

successfully establish and reproduce after mixing events occur (2, 3). There are many factors that

determine how bacterial community composition will change in response to an influx of saline-

dominant communities following a salinization event. Both stochastic processes (such as drift)

and deterministic processes (such as the interaction of biotic and abiotic factors) within an

ecosystem can determine patterns of community assembly via four processes: selection,

dispersal, diversification, and drift (35, 38). Abiotic factors in an environment often act as

environmental filters, selecting for organisms that are adapted to establish and persist. In this

study, when fresh and saline communities collided, the environmental conditions strongly dictated

the newly established community assemblages based on taxonomy but not phylogeny.

Bacterial taxa were more related to each other in transition salinity environments (5 and 9

psu) compared to fresh and the most saline endpoint environments. In the case of our study, we

added a mixture of freshwater and mesohaline (13 psu) aquatic communities in equal proportion

as dispersal treatments along an experimental salinity gradient. The dispersal treatments

represented natural dispersal methods of aquatic communities via water; a fresh community being

mixed with a saline community can occur in storm events with coastal flooding, as well as gradual

mixing of these environments via sea level rise. These methods of dispersal are on different time-

scales, but each is potentially important to community assembly at mixing zones. Because

dispersal of microbes is often a passive process, we recognize that the longest transport of

microbial species is via wind, water, and attachment to mobile organisms (Nemergut et al. 2013).

Overall, bacterial communities tended to respond to salinity, rather than dispersal, because it

is such a strong environmental filter and because salinity tolerance is a relatively deeply

conserved trait (10, 11, 39). The depth of phylogenetic relatedness was similar for saline and non-

saline communities. This suggests that there is a deep fresh-salt divide as previously revealed (7,

8, 10). Past studies show that selection may act on traits that are subject to HGT, which can alter

processes by transferring genetic material (i.e., a trait that would allow a previously rare organism

to establish and persist in a previously inhospitable environment) (40, 41). These organisms can

now explore new fitness landscapes and diversify (35). However, gaining a trait is not always a

simple process, as some traits are complex and cannot be easily inherited (10, 11, 39). Ecological

drift (constant changes in relative abundance of organisms) may also potentially affect community

assembly. In particular, low abundance organisms (i.e., majority of microbial species) are more

vulnerable to drift (35). Alternatively, rare taxa that were not detectable based on amplicon

sequencing methods were identified as responding positively to mixing events, making the role of

rare taxa important for driving community composition changes (7, 8).

It is also important to consider the intensity and length of the disturbance affecting a microbial

community (42). The disturbance in this experiment was the addition of salinity and saline-

adapted aquatic species to historically freshwater communities. This disturbance was intended to

represent salinization due to acute salinization events such as storm surge. At the end of the

experiment, phylogenetic diversity (PD) was low for both dispersal treatments at the highest

salinity, indicating that the bacteria persisting in those tanks were more related to each other

(lower PD). This observation supported the hypothesis that PD would be lowest in the most saline

tank, due to salinity tolerance being a complex trait.

After saline and freshwater communities mixed, we observed the highest carbon

mineralization rates in freshwater environments and associated disproportionately lower

abundance taxa. In contrast, microbes carried out lower carbon mineralization rates under saline

conditions, which were associated with indicator taxa at >25% relative abundance. Salinization

influences microbes through osmotic stress and ion-specific toxicity, but salinization can also

provide terminal electron acceptors to drive and change the rates and types of microbial

metabolisms. Some studies suggest that long-term salinity changes, like those due to sea level

rise, may occur slowly enough that the organisms can adapt (e.g., Mansour et al. 2018). Acute

salinity changes caused by "pulses" of salinity influx, such as localized flooding due to storm

surge, will more negatively affect freshwater aquatic communities compared to chronic exposure

to salinization where the communities have had time to adapt or turnover resulting in recovered

biogeochemical functions (e.g., Mansour et al. 2018). Thus, unexpected pulses of salinity are

observed (in some cases) to cause greater damage to ecosystem structure and function than

long-term salinity changes due to the inability of organisms to adapt or change quickly following

one of these events (6, 17, 43).

Global climate change plays a large role in facilitating increased dispersal of microbes; the predicted increases in extreme weather events may potentially promote increased spread of microbes to new, adaptable environments (44, 45). As previously discussed, microbes have several mechanisms for adapting to environmental change; this experiment did not address which mechanisms were used. Future studies focused on bacterial community structure and function under increasing salinity would benefit from examining mechanisms of adaptation that taxa may use when placed under environmental stress.

Acknowledgments

We thank Spencer Wilkinson, Amanda Dunn, and Mary-Grace Lee for help with data collection. We also thank Mike Piehler and Corey Adams for logistical help at the Coastal Studies Institute. This work was supported in part by the East Carolina University Division of Research, Economic Development, and Engagement and the Department of Biology.

References

1. Feckler A, Goedkoop W, Konschak M, Bundschuh R, Kenngott KGJ, Schulz R, Zubrod JP, Bundschuh M. 2018. History matters: heterotrophic microbial community structure and function adapt to multiple stressors. Glob Change Biol 24:e402–e415.

2. Mansour I, Heppell CM, Ryo M, Rillig MC. 2018. Application of the microbial community coalescence concept to riverine networks. Biol Rev 93:1832–1845.

3. Castledine M, Sierocinski P, Padfield D, Buckling A. 2020. Community coalescence: an eco-evolutionary perspective. Philos Trans R Soc B Biol Sci 375:20190252.

4. He Q, Silliman BR. 2019. Climate Change, Human Impacts, and Coastal Ecosystems in the Anthropocene. Curr Biol 29:R1021–R1035.

5. Kirwan ML, Megonigal JP. 2013. Tidal wetland stability in the face of human impacts and sea-level rise. Nature 504:53–60.

6.   Herbert ER, Boon P, Burgin AJ, Neubauer SC, Franklin RB, Ardón M, Hopfensperger KN,

Lamers LPM, Gell P. 2015. A Global Perspective on Wetland Salinization: Ecological

Consequences of a Growing Threat to Freshwater Wetlands. Ecosphere 6:1–43.

7.   Shen D, Langenheder S, Jürgens K. 2018. Dispersal Modifies the Diversity and

Composition of Active Bacterial Communities in Response to a Salinity Disturbance. Front

Microbiol 9.

8.   Rocca JD, Simonin M, Bernhardt ES, Washburne AD, Wright JP. 2020. Rare microbial

taxa emerge when communities collide: freshwater and marine microbiome responses to

experimental mixing. Ecology 101:e02956.

9.   Logares R, Lindström ES, Langenheder S, Logue JB, Paterson H, Laybourn-Parry J,

Rengefors K, Tranvik L, Bertilsson S. 2013. Biogeography of bacterial communities

exposed to progressive long-term environmental change. 5. ISME J 7:937–948.

10.   Paver SF, Muratore D, Newton RJ, Coleman ML. 2018. Reevaluating the Salty Divide:

Phylogenetic Specificity of Transitions between Marine and Freshwater Systems.

mSystems 3.

11.   Martiny JBH, Jones SE, Lennon JT, Martiny AC. 2015. Microbiomes in light of traits: A

phylogenetic perspective. Science 350:aac9323.

12.   Oren A. 1999. Bioenergetic Aspects of Halophilism. Microbiol Mol Biol Rev 63:334–348.

13.   Galinski EA, Trüper HG. 1994. Microbial behaviour in salt-stressed ecosystems. FEMS

Microbiol Rev 15:95–108.

14.   Kempf B, Bremer E. 1998. Uptake and synthesis of compatible solutes as microbial stress

responses to high-osmolality environments. Arch Microbiol 170:319–330.

15.   Weston NB, Vile MA, Neubauer SC, Velinsky DJ. 2011. Accelerated microbial organic

matter mineralization following salt-water intrusion into tidal freshwater marsh soils.

Biogeochemistry 102:135–151.

16.   Morrissey EM, Gillespie JL, Morina JC, Franklin RB. 2014. Salinity affects microbial activity

and soil organic matter content in tidal wetlands. Glob Change Biol 20:1351–1362.

17. Chambers LG, Osborne TZ, Reddy KR. 2013. Effect of salinity-altering pulsing events on

soil organic carbon loss along an intertidal wetland gradient: a laboratory experiment.

Biogeochemistry 115:363–383.

18. Marton JM, Herbert ER, Craft CB. 2012. Effects of Salinity on Denitrification and

Greenhouse Gas Production from Laboratory-incubated Tidal Forest Soils. Wetlands

32:347–357.

19. Neubauer SC, Franklin RB, Berrier DJ. 2013. Saltwater intrusion into tidal freshwater

marshes alters the biogeochemical processing of organic carbon. Biogeosciences

10:8171–8183.

20. Neubauer SC, Piehler MF, Smyth AR, Franklin RB. 2019. Saltwater Intrusion Modifies

Microbial Community Structure and Decreases Denitrification in Tidal Freshwater Marshes.

Ecosystems 22:912–928.

21. Stagg CL, Schoolmaster DR, Krauss KW, Cormier N, Conner WH. 2017. Causal

mechanisms of soil organic matter decomposition: deconstructing salinity and flooding

impacts in coastal wetlands. Ecology 98:2003–2018.

22. Dupont CL, Larsson J, Yooseph S, Ininbergs K, Goll J, Asplund-Samuelsson J, McCrow

JP, Celepli N, Allen LZ, Ekman M, Lucas AJ, Hagström Å, Thiagarajan M, Brindefalk B,

Richter AR, Andersson AF, Tenney A, Lundin D, Tovchigrechko A, Nylander JAA, Brami D,

Badger JH, Allen AE, Rusch DB, Hoffman J, Norrby E, Friedman R, Pinhassi J, Venter JC,

Bergman B. 2014. Functional Tradeoffs Underpin Salinity-Driven Divergence in Microbial

Community Composition. PLOS ONE 9:e89549.

23. Kaushal SS, Likens GE, Pace ML, Utz RM, Haq S, Gorman J, Grese M. 2018. Freshwater

salinization syndrome on a continental scale. Proc Natl Acad Sci 115:E574–E583.

24. Rath KM, Rousk J. 2015. Salt effects on the soil microbial decomposer community and

their role in organic carbon cycling: A review. Soil Biol Biochem 81:108–123.

25.   Trivedi P, Delgado-Baquerizo M, Trivedi C, Hu H, Anderson IC, Jeffries TC, Zhou J, Singh

BK. 2016. Microbial regulation of the soil carbon cycle: evidence from gene–enzyme

relationships. ISME J 10:2593–2604.

26.   Werba JA, Stucy AL, Peralta AL, McCoy MW. 2020. Effects of diversity and coalescence of

species assemblages on ecosystem function at the margins of an environmental shift.

PeerJ 8:e8608.

27.   Thompson PL, Shurin JB. 2012. Regional zooplankton biodiversity provides limited

buffering of pond ecosystems against climate change. J Anim Ecol 81:251–259.

28.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM,

Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012. Ultra-high-

throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms.

ISME J 6:1621–1624.

29.   Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb

Ecol 75:129–137.

30.   Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. Environ Microbiol 18:1403–1414.

31.   Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a

dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence

data on the MiSeq Illumina sequencing platform. Appl Environ Microbiol 79:5112–20.

32.   Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J,

Ludwig W, Glöckner FO. 2014. The SILVA and "All-species Living Tree Project (LTP)"

taxonomic frameworks. Nucleic Acids Res 42:D643–D648.

33.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: a versatile open

source tool for metagenomics. PeerJ 4:e2584.

34.   Martiny AC, Treseder K, Pusch G. 2013. Phylogenetic conservatism of functional traits in

microorganisms. ISME J 7:830.

35.   Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, Knelman JE,

Darcy JL, Lynch RC, Wickey P, Ferrenberg S. 2013. Patterns and Processes of Microbial

Community Assembly. Microbiol Mol Biol Rev 77:342–356.

36.   Graham EB, Crump AR, Resch CT, Fansler S, Arntzen E, Kennedy DW, Fredrickson JK,

Stegen JC. 2016. Coupling Spatiotemporal Community Assembly Processes to Changes

in Microbial Metabolism. Front Microbiol 7.

37.   Reese AT, Lulow K, David LA, Wright JP. 2018. Plant community and soil conditions

individually affect soil microbial community assembly in experimental mesocosms. Ecol

Evol 8:1196–1205.

38.   Vellend M. 2010. Conceptual Synthesis in Community Ecology. Q Rev Biol 85:183–206.

39.   Logares R, Bråte J, Bertilsson S, Clasen JL, Shalchian-Tabrizi K, Rengefors K. 2009.

Infrequent marine–freshwater transitions in the microbial world. Trends Microbiol 17:414–

422.

40.   Soucy SM, Huang J, Gogarten JP. 2015. Horizontal gene transfer: building the web of life.

Nat Rev Genet 16:472–482.

41.   Hall JPJ, Brockhurst MA, Harrison E. 2017. Sampling the mobile gene pool: innovation via

horizontal gene transfer in bacteria. Philos Trans R Soc B Biol Sci 372:20160424.

42.   Graham EB, Averill C, Bond-Lamberty B, Knelman JE, Krause S, Peralta AL, Shade A,

Smith AP, Cheng SJ, Fanin N, Freund C, Garcia PE, Gibbons SM, Van Goethem MW,

Guebila MB, Kemppinen J, Nowicki RJ, Pausas JG, Reed SP, Rocca J, Sengupta A, Sihi

D, Simonin M, Słowiński M, Spawn SA, Sutherland I, Tonkin JD, Wisnoski NI, Zipper SC,

Consortium C, Staal A, Arora B, Oldfield C, Dwivedi D, Larson E, Santillan E, Aaron Hogan

J, Atkins J, Zheng J, Lembrechts J, Patel K, Copes-Gerbitz K, Winker K, Mudge L, Wong

M, Nuñez M, Luoto M, Barnes R. 2021. Toward a Generalizable Framework of Disturbance

Ecology Through Crowdsourced Science. Front Ecol Evol 9.

43. Wilson BJ, Servais S, Mazzei V, Kominoski JS, Hu M, Davis SE, Gaiser E, Sklar F,

Bauman L, Kelly S, Madden C, Richards J, Rudnick D, Stachelek J, Troxler TG. 2018.

Salinity pulses interact with seasonal dry-down to increase ecosystem carbon loss in

marshes of the Florida Everglades. Ecol Appl 28:2092–2108.

44. Litchman E. 2010. Invisible invaders: non-pathogenic invasive microbes in aquatic and

terrestrial ecosystems. Ecol Lett 13:1560–1572.

45. Hutchins DA, Fu F. 2017. Microorganisms and ocean global change. Nat Microbiol 2:1–11.

517 **FIGURES**

518 **Figure 1.** Schematic of experimental mesocosm to test the effect of salinity and dispersal of

519 estuarine communities into freshwater coastal ponds (A) and bacterial sequence processing for

520 statistical analyses (B).

521   (A) Experimental design

522
523   (B) Statistical analyses
524

 Figure 2. Bacterial phylogenetic and taxonomic patterns along a salinity gradient. Ordination

 based on a Principal Coordinates Analysis depicting bacterial phylogenetic distance (A) and

 Bray-Curtis dissimilarity distance (modified from Werba et al. 2020) (B), where symbols are

 colored according to salinity treatment (0, 5, 9, or 13 psu) and symbol shapes represent date of

 sampling (Day 0, 18, 45). Raincloud plots of phylogenetic diversity along a salinity gradient,

 where plots represent raw data depicted as filled circles ('rain'), probability distributions

 ('clouds'), and boxplots displaying mean and 95% confidence intervals (C). Colors represent

 salinity treatment (orange=0 psu, blue=5 psu, green=9 psu, yellow=13 psu), and shapes

 represent date of sampling (square=Day 0, circle=18,triangle=45).

Figure 3. Phylogenetic patterns of indicator bacterial taxa. Phylogenetic tree of bacterial taxa that

were identified to be statistically associated with salinity treatment (0, 5, 9, 13 or >13 psu); each

taxon tip color corresponds to phyla level grouping and is associated with a heat map which

represents OTU relative abundance salinity treatment (A). Euler plots represent bacterial OTU

that are unique or shared across salinity treatments (B). Average cluster depth estimated using

ConcenTrait used to depict phylogenetic distance of bacterial assemblages observed across the

salinity gradient (C).

(3A)

(3B)

(3C)

Figure 4. Relationship between carbon mineralization and bacterial phylogenetic diversity (PD),

and taxonomic-based community metrics (relative abundance, OTU richness, and Shannon

diversity) for Day 45 (last day of experiment).