Bacterial composition reflects fine-scale salinity changes while phylogenetic diversity exhibits a strong salt divide

Ariane L. Peralta[1*], Mario E. Muscarella[2], Alexandra Stucy[1], Jo A. Werba[1,3], Michael W. McCoy[1,4]

[1]Department of Biology, East Carolina University, Howell Science Complex, Greenville, NC 27858, USA; [2]Department of Biology and Wildlife, Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK, 99775, USA; [3]Department of Biology, McMaster University, 1280 Main Street West, Hamilton, Ontario, Canada, L8S 4K1; [4]Harbor Branch Oceanographic Institute, Florida Atlantic University, Fort Pierce, FL, 34946, USA

*corresponding author

ABSTRACT

Climate change induced salinization events are predicted to intensify and lead to increased salt stress in freshwater aquatic ecosystems. As a consequence, formerly distinct abiotic conditions and associated biotic communities merge, and the emergence, loss, and persistence of microbial taxa modify the types and rates of ecosystem processes. This study examined how bacterial taxonomic and phylogenetic diversity and ecosystem function respond to acute salinization events where freshwater and estuarine communities and environments coalesce. We hypothesized that when the salinity change outpaces microbial adaptation or saline microbial populations are not yet established in formerly freshwater conditions, then these aquatic communities will exhibit diminished carbon cycling rates, decreased microbial diversity, and altered composition of microbial communities compared to historically freshwater communities. We used an experimental mesocosm approach to determine how salinity and the merging of distinct communities influenced resultant bacterial community structure and function. Each mesocosm represented different salinities (0, 5, 9, 13 psu). Two dispersal treatments, representing aquatic communities sourced from brackish 13 psu ponds and a mix of 13 psu and freshwater ponds, were added to all salinity levels and replicated four times. Results revealed that salinity, but not dispersal, decreased bacterial taxonomic and phylogenetic diversity. Carbon mineralization rates were highest in freshwater conditions and associated to bacterial taxa represented in low relative abundance. Acute salinity changes, such as localized flooding due to storm surge, will more negatively affect freshwater aquatic communities compared to chronic exposure to salinization where the communities have had time to adapt or turnover.

IMPORTANCE STATEMENT

Climate change induced salinization results in the mixing of formerly distinct environmental conditions and aquatic communities. This study examined the consequence of short-term, acute salinity stress on aquatic bacterial taxonomic and phylogenetic diversity and ecosystem function using an experimental approach. Results revealed that salinity, but not the source of aquatic communities, decreased bacterial taxonomic and phylogenetic diversity. Carbon mineralization rates, which represented ecosystem function, were highest in freshwater conditions and also associated with indicator bacterial taxa in low abundance relative to the total microbial community. Taken together, acute salinity changes will more negatively affect freshwater aquatic communities compared to chronic exposure to salinization where the communities have had time to adapt or turnover resulting in recovered biogeochemical functions.

INTRODUCTION

Predicting microbial community response to environmental change depends on the tolerance and preference of microbial taxa to current and historical environmental conditions, ongoing changes in environmental conditions, and concurrent community assembly processes (1–3). Environmental changes, such as sea level rise and punctuated salinization events, are predicted to intensify in the coming decades as land development and climate change continue (4). Sea-level rise and saltwater intrusion in previously freshwater wetlands can diminish wetland ecosystem functions by increasing toxic sulfide production, decreasing inorganic nitrogen removal, and decreasing carbon storage. In some cases, diminished microbial functions negatively affect water quality, climate regulation, wetland accretion, and the health of wetland biota (5, 6). However, it is unclear how the composition and function of microbial communities at newly established fresh-salt water interfaces will change as salinization events increase in frequency and duration, especially with regard to subsets of taxa that respond to the coalescence event (i.e., the merging of distinct microbial communities and environments) (7, 8). As aquatic ecosystems shift from fresh to brackish waters, the concentration of alternative terminal electron acceptors (i.e., Fe(III), Mn(IV)) increases, which prompts CO_2 production (6). As a result, environmental stressors are altering the microbial communities governing these carbon cycling rates.

Salinity tolerance has convergently evolved in unrelated lineages across the tree of life (9, 10). Salinity preference and osmotic tolerance are considered a complex microbial trait (11). Two mechanisms are responsible for osmotic or salinity stress tolerance in microorganisms. First, microorganisms can adapt by excluding harmful solutes (e.g., sodium, chloride) while gathering beneficial solutes necessary for metabolism by using active transport systems (12). Second, some microorganisms can create organic compounds to reduce the concentration gradient between the external environment and the cell resulting in the net export of unwanted solutes (13–15). Some studies suggest that long-term salinity changes, like those due to sea level rise,

may occur slowly enough that the organisms can adapt and/or communities reassemble according to environmental changes. Long-term, incremental salinity increases have promoted microbial diversity and community establishment of salt-preference in a variety of taxa. Salinization in tidal and non-tidal systems results in different community responses. For example, tidal wetland decomposition rates increase as salinity increases due in part to marine/estuarine subsidies mixing with freshwater ecosystems (16). While experiencing short-term acute salinization, microbial populations have recovered from previously diminished carbon cycling functions (carbon dioxide and methane production) (17). Prior studies have revealed both increases in decomposition rates (15, 18) and decreases in decomposition rates (19, 20) as a result of increases in salinity. Previous studies on salinization effects of freshwaters reveal that punctuated salinity pulses to historically freshwater ecosystems, such as localized flooding due to storm surge in a climatic event, will more negatively affect freshwater communities compared to constant exposure to salinization over several years (2, 17, 21–24). The challenge to predicting salinization effects on decomposition rates is due to interactions among the following: direct effects of redox shifts between oxic and anoxic conditions, direct effects on heterotrophic communities and local nutrient availability, and the indirect effects due to changes in organic carbon sources (25).

Salinity tolerance and preference are traits that constrain microbial community response to salinization (26). As freshwater salinization across inland freshwaters persists, drastic changes to microbial community structure and function will continue to increase (27). The emergence, loss, and persistence of microbial taxa can alter the types and rates of ecosystem functions. The microbial taxa that will survive punctuated salinity changes will have specialized traits. These 'effect' traits can be directly linked to ecosystem functioning (11). Salinity preference is identified as one of the most complex trait measurements, classifying it based on how many genes are directly involved in coding the trait and how the trait is integrated with other mechanisms (11). Because specialized traits are needed for organisms to tolerate increasingly saline environments,

major shifts in microbial composition and diversity are expected (28, 29). When freshwater and marine microbes are experimentally mixed, results showed that taxa which were rare at the initial inoculation became relatively abundant (7). As the community shifts, organisms that are better adapted to new conditions may become more active and abundant, which can alter the dominant ecosystem processes (1–3).

In this study, we address the research question: how does acute salinization affect microbial taxonomic and phylogenetic diversity and function when freshwater microbial communities mix with estuarine aquatic communities along freshwater to saltwater gradient? Specialized microbial traits are necessary to tolerate or thrive in saline environments. We hypothesize that during scenarios when salinity change outpaces microbial adaptation or saline microbial populations are not yet established in formerly freshwater conditions, aquatic communities will exhibit diminished carbon cycling rates (i.e., CO_2 respiration) and decreased microbial diversity and altered the composition of microbial communities when compared to historically freshwater microbial communities. We also expect that microorganisms that persist in more saline conditions will be more phylogenetically related to each other (i.e., less genetic variation) compared to historically freshwater microorganisms. Dispersal of saline communities into freshwater ecosystems have the opportunity to establish and outcompete freshwater communities, but there is likely a period of transition where the mixing of environments and bacterial communities are dynamic and communities are reassembling. Understanding how salinization alters freshwater wetland bacterial phylogenetic-ecosystem function relationships can inform the management of carbon storage capacity in coastal wetlands experiencing increased salinization.

MATERIALS AND METHODS

Experimental Design. Microbial community composition and function were characterized based on aquatic samples that were collected from a replicated mesocosm experiment conducted from June-July 2015 (Figure 1) detailed in Werba et al. (30). To replicate a freshwater-saltwater aquatic

system, peat moss (to serve as a nutrient pulse), sand (to serve as a benthic substrate), and water sourced from the municipal water supply were added to 150-gallon stock watering tanks (filled to 100 gallons). Each mesocosm pond was adjusted to one of 4 different salinities (measured in practical salinity units) using Instant Ocean® Sea Salt (4 levels: 0, 5, 9, 13 psu). Each tank was then seeded with an inoculation from natural ponds in the outer banks of North Carolina that matched the treatment's psu (see **Supplemental Table S1** for coastal pond location and information). The inocula consisted of water, zooplankton and microbes. Twenty 1 L water samples were collected and filtered through 62.5 μm mesh across a one-hundred meter transect at each of the 5 coastal ponds and subsequently dispersed into each mesocosm tank. Bacterial communities represented in the source communities were likely particle- and zooplankton-associated because source communities were collected using a 62.5 μm mesh. Each tank was covered with a shade cloth to reduce the opportunity for other organisms from colonizing the mesocosms. Populations introduced via the initial inoculation were allowed to stabilize for 6 weeks before sampling began. After this acclimation period, mesocosms received additional colonists of zooplankton and microbes via simulated dispersal events from separate mesocosm ponds established as source tanks for salt (13 psu) and freshwater (0 psu) zooplankton and microbial communities. The source tanks served as dispersal treatments and consisted of either a saltwater (13 psu; salt dispersal treatment) community or a mix of communities from freshwater (0 psu) and saltwater (13 psu; fresh+salt dispersal treatment). Dispersal treatments consisted of 2 L of water from dispersal (source) tanks into the experiment (treatment) tanks. Sample collection occurred every 9 days (due to average time for completion of one zooplankton generation cycle) (31) over the 6-week mesocosm experiment for a total of six time points. The salinity × dispersal setup was replicated 4 times to generate 32 tanks (8 at each salinity level divided into 4 in each dispersal treatment) (30). For each tank, the following were measured using a YSI Pro: dissolved oxygen, ammonium (NH_4^+), temperature, and pH. Community analysis was focused on three of the time points (days 0, 18, and 45; 11 June 2015, 29 June 2015, and 25 July 2015, respectively),

representing the initial, middle, and end of the experiment. Besides salinity levels, measured nitrogen and phosphorus concentrations were similar among treatment tanks (30).

Microbial Sample Collection and Processing. Sample collection occurred every 9 days over the time of the 6-week mesocosm experiment for a total of six time points. During each sampling event, we collected 1 L of water from each tank. Each 1 L bottle of mesocosm water was homogenized and 200 mL of the water sample were concentrated onto 0.22 μm filters within 24 hours of field sampling (Supor-200; Pall Gelman, East Hills, NY) to collect microbial community samples. Filters were transferred into 2 mL sterile tubes and stored at -80° C until molecular analyses. We extracted DNA from filters collected at three (of the six) time points representing the initial, middle, final sampling dates (Days 0, 18, 45) using the PowerWater DNA Isolation Kit (MO BIO Laboratories).

Microbial Community Sequencing. To examine shifts in microbial community composition and diversity, aquatic microbial communities in each mesocosm were characterized using paired-end targeted Illumina sequencing of the 16S rRNA gene (bacteria, archaea) (32). We extracted and purified the DNA found in the water of each mesocosm using the PowerWater DNA Isolation Kit (MO BIO Laboratories, Inc., Carlsbad, CA) collected on a 0.22 μm supor filter post-filtration. We used this DNA as a template in PCR reactions. To characterize bacterial communities, we used barcoded primers (515FB/806RB) originally developed by the Earth Microbiome Project (32) to target the V4 region of the bacterial 16S subunit of the ribosomal RNA gene (33, 34). PCR products were combined in equimolar concentrations and sequenced using paired-end (2×250 bp) approach using the Illumina MiSeq platform at Indiana University's Center for Genomics and Bioinformatics.

Raw bacterial sequences were processed using the Mothur pipeline (version 1.41.3) (35). Briefly, paired-end contigs were assembled and quality trimmed, sequences were then aligned to

the Silva Database (version 132) (36) chimeric sequences were removed using the VSEARCH algorithm (37), and we removed any sequences classified as Chloroplasts, Mitochondria, Archaea, or Eukaryotes. Next, we identified operational taxonomic units (OTUs) by separating sequences based on taxonomic class and then binning sequences using a 97% sequence similarity cutoff. Representative sequences from each OTU were used to generate a phylogenetic tree using FastTree v2.1 using the GTR model and CAT rate approximation (38). See Supplemental Material for full sequence processing pipeline.

Compositional Analyses. We used a combination of taxonomic and phylogenetic approaches to characterize the diversity of bacterial communities in each mesocosm. First, we calculated alpha diversity (within-sample) using either Shannon's diversity (taxonomic) or Faith's Diversity (phylogenetic). The Faith's Phylogenetic Diversity (PD) metric provides a simple measure of the phylogenetic relatedness of a community based on the summed branch lengths of its phylogenetic tree. We expected this measure to capture functional complementarity if more distantly related species are more functionally unique. We rarefied the OTU table to 20,000 observations before calculating alpha-diversity. Next, we calculated beta-diversity (among sites) by calculating Bray-Curtis distance for taxonomic diversity and weighted UniFrac distance for phylogenetic diversity. Differences in beta-diversity were visualized using Principal Coordinates Analysis.

Carbon Mineralization Assay. On the final mesocosm sampling date, day 45, we measured the amount of CO_2 respired from the aquatic communities using a laboratory-based bottle assay. Wheaton bottles (125 mL) fitted with septa were filled with water samples (25 mL) from each mesocosm tank. The CO_2 concentration readings were determined using an LI-7000 Infrared Gas Analyzer (IRGA). On the first day (Day 0), bottles were filled with 25 mL of mesocosm tank water, and the gas samples were collected and analyzed immediately using the IRGA to determine the baseline CO_2 concentration. A syringe was inserted into the septa and the headspace gas was

mixed 3 times before pulling a sample and beginning analysis using the IRGA. This process was repeated on days one, three, and seven in order to determine CO_2 respiration rates over time.

Statistical Analyses. We completed statistical calculations in the R environment (R v4.1.2 R Core Development Team 2021). For bacterial diversity metrics, we conducted a linear mixed effects model with 'date', 'salinity', and 'dispersal' as fixed effects and 'block' as a random effect using the *lmer()* function in the lmerTest package (Kuznetsova et al. 2017) and the pbkrtest package (39). We ran linear mixed models that were fit using a restricted maximum likelihood (REML) approach and produced type II analysis of variances tables (ANOVA) tables based on the Kenward-Roger's denominator degrees of freedom method using the *anova()* function. Then, we used principal coordinates analysis (PCoA) based on the Bray-Curtis dissimilarity of bacterial community composition and based on the weighted Unifrac distance of phylogenetic diversity along the salinity gradient. We ran a permutational multivariate analysis of variance (PERMANOVA) to examine among-treatment differences in bacterial communities. We identified which bacterial species were most representative of each salinity treatment using an indicator species analysis based on salinity level and also a categorical saline vs non-saline classification, and we included bacterial taxa with a relative abundance greater than 0.05 when summed across all plots. We performed PERMANOVA using the *adonis()* function in the vegan package (40) and the *indval()* function in the indicspecies package (41). We visualized overlap in indicator taxa across the salinity gradient using an Euler plot. To better understand the phylogenetic underpinnings of saline tolerant and sensitive taxa, we visualized the indicators on a phylogenetic tree and used the ConcenTrait algorithm to determine the phylogenetic depth at which saline tolerance/sensitivity is conserved (42). Briefly, this algorithm finds the nodes in the phylogeny at which 90% of the included tips (taxa) share the same trait. We implemented the ConcenTrait algorithm using a custom R script.

Data Availability

All code and data used in this study are located in a public GitHub repository (https://github.com/PeraltaLab/CSI_Dispersal) and NCBI SRA BioProject ID PRJNA615001.

RESULTS

We examined bacterial community turnover throughout the mesocosm experiment by comparing patterns in bacterial phylogenetic and taxonomic composition across the salinity gradient. Throughout the experiment, freshwater communities were phylogenetically distinct compared to saline communities based on weighted UniFrac distance metric (Fig. 2A, Table 1, PERMANOVA salinity: R^2=0.172, P<0.001). Bacterial communities were phylogenetically similar (based on phylogenetic diversity) during day 18, but after 45 days at the experimental salinity level, communities at 9 and 13 psu were distinct from communities at 5 psu (Fig. 2C, Table 1, PERMANOVA salinity x date: R^2=0.048, P<0.001). When bacterial taxonomic composition was compared, salinity was a much stronger environmental filter associated with bacterial community patterns based on taxonomy (Fig. 2B) compared to phylogeny (Fig. 2A)

Next, we compared patterns in bacterial phylogenetic diversity to taxonomic composition across the salinity gradient. Communities observed in the freshwater conditions increased in mean PD over the 45 day experiment. The PD of communities at the transition 9 psu tanks declined throughout the experiment duration, while PD of communities in the 13 psu tanks increased in variability from day 0 to 45 (Fig. 2C, Table 2, ANOVA, $F_{2,79}$=5.997, P=0.004). After 45 days, PD was highest in freshwater and lowest in brackish tanks (Fig. 2C).

There are OTUs represented across major phyla that are associated with all salinity levels (Fig. 3A). Based on the unrooted tree (Fig. 3A), we observed that the smallest number of OTUs were represented at 9 psu treatment compared to other treatments. Next, we used a series of Euler plots to reveal shared and unique indicator OTUs. A list of "indicators" that were significantly

associated with each salinity treatment is found in Table S2. These indicators, reported at the class-level, are the taxonomic groups best associated with each treatment. Based on OTUs identified as indicator taxa, 254 OTUs are unique to the freshwater treatment, 135 OTUs are unique to the 5 psu treatment, 35 OTUs were unique to the 9 psu treatment, and 124 OTUs are unique to the 13 psu treatment (Fig. 3B). When the OTUs were grouped into salt or no salt, there are 206 observed indicator OTUs shared across 5, 9, or 13 psu environments, and 257 OTUs unique to the 0 psu treatment (Fig. 3B). Finally, when examining the depth of phylogenetic relatedness, results from the ConcenTrait analysis revealed that saline vs non-saline microbes exhibit a similar depth in average phylogenetic relatedness (Fig. 3C).

We examined the relationship between carbon mineralization and phylogenetic diversity and relative abundance (Fig. 4A). At the transition 9 psu treatments, we measured the lowest carbon mineralization values compared to other salinity treatments. Carbon mineralization rates were highest in freshwater conditions and associated with indicator bacterial taxa that were measured at low relative abundance (< 0.01%) (Fig. 4B). In contrast, the relative abundance of indicator taxa represented in the saltwater conditions represented > 25% of the community.

DISCUSSION

This study specifically focused on determining the extent that short-term salinization events influenced reassembly patterns of aquatic communities and carbon mineralization rates. Results revealed that salinity strongly influenced bacterial phylogenetic diversity; bacterial communities tended to respond to salinity, rather than dispersal, because it is such a strong environmental filter (7, 10). Past research has shown that strong environmental filters produce phylogenetically clustered communities (11, 43, 44). However, the mixing of two formerly distinct communities and their environments could affect community structure and function in unpredictable ways. Outcomes depend on the strength of the environmental change and the capacity for taxa to successfully establish and reproduce after mixing events occur (2, 3). There are many factors that

determine how bacterial community composition will change in response to an influx of saline-dominant communities following a salinization event. Both stochastic processes (such as drift) and deterministic processes (such as the interaction of biotic and abiotic factors) within an ecosystem can determine patterns of community assembly via four processes: selection, dispersal, diversification, and drift (43, 45). Abiotic factors in an environment often act as environmental filters, selecting for organisms that are adapted to establish and persist. In this study, once mesocosms were seeded with aquatic inocula, the environmental conditions strongly dictated the newly established community assemblages based on taxonomy (Fig. 2B and (30)) but not phylogeny (Fig. 2A).

Bacterial taxa were more related to each other in transition salinity environments (5 and 9 psu) compared to fresh and the most saline endpoint environments. The composition of the bacterial community (based on phylogeny) was similar in the 5 and 9 psu treatment, but the phylogenetic diversity rebounded for communities in the 5 psu compared to 9 psu treatments. The 9 psu environment could be particularly stressful for the resorted 'transition' community. In the current study, we added a mixture of freshwater and mesohaline (13 psu) aquatic communities in equal proportion as dispersal treatments along an experimental salinity gradient. The dispersal treatments represented natural dispersal methods of aquatic communities via water; a fresh community being mixed with a saline community can occur in storm events with coastal flooding, as well as gradual mixing of these environments via sea level rise. These methods of dispersal are on different time-scales, but each is potentially important to community assembly at mixing zones. Because dispersal of microbes is often a passive process, we recognize that the longest transport of microbial species is via wind, water, and attachment to mobile organisms (43).

Overall, bacterial communities tended to respond to salinity, rather than dispersal. Our results provide further support that salinity is a strong environmental filter especially in light of salinity tolerance being a relatively deeply conserved trait (10, 11, 46). The depth of phylogenetic

relatedness was similar for saline and non-saline communities. This suggests that there is a deep fresh-salt divide as previously revealed (7, 8, 10).

Since salinity tolerance is considered a phylogenetically conserved trait, it is more likely that the reassembly of the mixed community to represent a new state. Past studies show that selection may act on traits that are subject to HGT, which can alter processes by transferring genetic material (i.e., a trait that would allow a previously rare organism to establish and persist in a previously inhospitable environment) (47, 48). These organisms can now explore new fitness landscapes and diversify (43). However, gaining a trait is not always a simple process, as some traits are complex and cannot be easily inherited (10, 11, 46). This study did not support the idea that selection for salinity tolerance was occurring due to the short-term study design. Ecological drift (constant changes in relative abundance of organisms) can potentially affect community assembly. In particular, low abundance organisms (i.e., majority of microbial species) are more vulnerable to drift (43). Alternatively, rare taxa that were not detectable based on amplicon sequencing methods were identified as responding positively to mixing events, making the role of rare taxa important for driving community composition changes (7, 8).

Salinity stressors on freshwater aquatic communities could indirectly influence bacterial communities through influence other members of the food web. For example, microeukaryotes such as protists could shift in composition, changing grazer pressures on bacterial prey (49, 50). While the current study did not examine the other members of the food web, past studies revealed the importance of eco-evolutionary processes of food web interactions that have important implications for climate change effects on carbon cycling (51, 52).

It is also important to consider the intensity and length of the disturbance affecting a microbial community (53). The disturbance in this experiment was the addition of salinity and saline-adapted aquatic species to historically freshwater communities. This disturbance was intended to represent salinization due to acute events such as storm surge. At the end of the experiment, phylogenetic diversity (PD) was low for both dispersal treatments at the highest salinity, indicating

that the bacteria persisting in those tanks were more related to each other (lower PD). This observation supported the hypothesis that PD would be lowest in the most saline tank, due to salinity tolerance being a complex trait.

After saline and freshwater communities mixed, we observed the highest carbon mineralization rates in freshwater environments and associated disproportionately lower abundance taxa. In contrast, microbes carried out lower carbon mineralization rates under saline conditions, which were associated with indicator taxa at >25% relative abundance. Past studies showed that conditionally rare microbial taxa play disproportionate role in community patterns over time and considered keystone taxa important for ecological responses to disturbance (9–11). Salinization influences microbes through osmotic stress and ion-specific toxicity, but salinization can also provide terminal electron acceptors to drive and change the rates and types of microbial metabolisms. Some studies suggest that long-term salinity changes, like those due to sea level rise, may occur slowly enough that the organisms can adapt (e.g., (2)). Acute salinity changes caused by "pulses" of salinity influx, such as localized flooding due to storm surge, will more negatively affect freshwater aquatic communities compared to chronic exposure to salinization where the communities have had time to adapt or turnover resulting in recovered biogeochemical functions (e.g., (2)). Thus, unexpected pulses of salinity are observed (in some cases) to cause greater damage to ecosystem structure and function than long-term salinity changes due to the inability of organisms to adapt or change quickly following one of these events (6, 17, 54). The current study provides insight into salinization effects of saline communities mixing with naïve freshwater communities, an occurrence that is expected at the early onset of salinization events (e.g., sea-level rise, intense storm surges upriver).

Global climate change plays a large role in facilitating increased dispersal of microbes; the predicted increases in extreme weather events may potentially promote increased spread of microbes to new, adaptable environments (55, 56). As previously discussed, microbes have several mechanisms for adapting to environmental change; this experiment did not address which

mechanisms were used. Future studies focused on bacterial community structure and function under increasing salinity would benefit from examining mechanisms of adaptation that taxa may use when placed under environmental stress.

Acknowledgments

We thank Spencer Wilkinson, Amanda Dunn, and Mary-Grace Lee for help with data collection. We also thank Mike Piehler and Corey Adams for logistical help at the Coastal Studies Institute. This work was supported in part by the East Carolina University Division of Research, Economic Development, and Engagement and the Department of Biology.

References

1. Feckler A, Goedkoop W, Konschak M, Bundschuh R, Kenngott KGJ, Schulz R, Zubrod JP, Bundschuh M. 2018. History matters: heterotrophic microbial community structure and function adapt to multiple stressors. Glob Change Biol 24:e402–e415.

2. Mansour I, Heppell CM, Ryo M, Rillig MC. 2018. Application of the microbial community coalescence concept to riverine networks. Biol Rev 93:1832–1845.

3. Castledine M, Sierocinski P, Padfield D, Buckling A. 2020. Community coalescence: an eco-evolutionary perspective. Philos Trans R Soc B Biol Sci 375:20190252.

4. He Q, Silliman BR. 2019. Climate Change, Human Impacts, and Coastal Ecosystems in the Anthropocene. Curr Biol 29:R1021–R1035.

5. Kirwan ML, Megonigal JP. 2013. Tidal wetland stability in the face of human impacts and sea-level rise. Nature 504:53–60.

6.   Herbert ER, Boon P, Burgin AJ, Neubauer SC, Franklin RB, Ardón M, Hopfensperger KN, Lamers LPM, Gell P. 2015. A global perspective on wetland salinization: ecological consequences of a growing threat to freshwater wetlands. Ecosphere 6:art206.

7.   Rocca JD, Simonin M, Bernhardt ES, Washburne AD, Wright JP. 2020. Rare microbial taxa emerge when communities collide: freshwater and marine microbiome responses to experimental mixing. Ecology 101:e02956.

8.   Shen D, Langenheder S, Jürgens K. 2018. Dispersal Modifies the Diversity and Composition of Active Bacterial Communities in Response to a Salinity Disturbance. Front Microbiol 9.

9.   Logares R, Lindström ES, Langenheder S, Logue JB, Paterson H, Laybourn-Parry J, Rengefors K, Tranvik L, Bertilsson S. 2013. Biogeography of bacterial communities exposed to progressive long-term environmental change. 5. ISME J 7:937–948.

10.   Paver SF, Muratore D, Newton RJ, Coleman ML. 2018. Reevaluating the Salty Divide: Phylogenetic Specificity of Transitions between Marine and Freshwater Systems. mSystems 3.

11.   Martiny JBH, Jones SE, Lennon JT, Martiny AC. 2015. Microbiomes in light of traits: A phylogenetic perspective. Science 350:aac9323.

12.   Oren A. 1999. Bioenergetic Aspects of Halophilism. Microbiol Mol Biol Rev 63:334–348.

13.   Galinski EA, Trüper HG. 1994. Microbial behaviour in salt-stressed ecosystems. FEMS Microbiol Rev 15:95–108.

14.   Kempf B, Bremer E. 1998. Uptake and synthesis of compatible solutes as microbial stress responses to high-osmolality environments. Arch Microbiol 170:319–330.

15. Weston NB, Vile MA, Neubauer SC, Velinsky DJ. 2011. Accelerated microbial organic matter mineralization following salt-water intrusion into tidal freshwater marsh soils. Biogeochemistry 102:135–151.

16. Morrissey EM, Gillespie JL, Morina JC, Franklin RB. 2014. Salinity affects microbial activity and soil organic matter content in tidal wetlands. Glob Change Biol 20:1351–1362.

17. Chambers LG, Osborne TZ, Reddy KR. 2013. Effect of salinity-altering pulsing events on soil organic carbon loss along an intertidal wetland gradient: a laboratory experiment. Biogeochemistry 115:363–383.

18. Marton JM, Herbert ER, Craft CB. 2012. Effects of Salinity on Denitrification and Greenhouse Gas Production from Laboratory-incubated Tidal Forest Soils. Wetlands 32:347–357.

19. Neubauer SC, Franklin RB, Berrier DJ. 2013. Saltwater intrusion into tidal freshwater marshes alters the biogeochemical processing of organic carbon. Biogeosciences 10:8171–8183.

20. Neubauer SC, Piehler MF, Smyth AR, Franklin RB. 2019. Saltwater Intrusion Modifies Microbial Community Structure and Decreases Denitrification in Tidal Freshwater Marshes. Ecosystems 22:912–928.

21. Entrekin SA, Clay NA, Mogilevski A, Howard-Parker B, Evans-White MA. 2019. Multiple riparian–stream connections are predicted to change in response to salinization. Philos Trans R Soc B Biol Sci 374:20180042.

22. Hintz WD, Relyea RA. 2019. A review of the species, community, and ecosystem impacts of road salt salinisation in fresh waters. Freshw Biol 64:1081–1097.

23. Hintz WD, Mattes BM, Schuler MS, Jones DK, Stoler AB, Lind L, Relyea RA. 2017. Salinization triggers a trophic cascade in experimental freshwater communities with varying food-chain length. Ecol Appl 27:833–844.

24. Steinmuller HE, Chambers LG. 2018. Can Saltwater Intrusion Accelerate Nutrient Export from Freshwater Wetland Soils? An Experimental Approach. Soil Sci Soc Am J 82:283–292.

25. Stagg CL, Schoolmaster DR, Krauss KW, Cormier N, Conner WH. 2017. Causal mechanisms of soil organic matter decomposition: deconstructing salinity and flooding impacts in coastal wetlands. Ecology 98:2003–2018.

26. Dupont CL, Larsson J, Yooseph S, Ininbergs K, Goll J, Asplund-Samuelsson J, McCrow JP, Celepli N, Allen LZ, Ekman M, Lucas AJ, Hagström Å, Thiagarajan M, Brindefalk B, Richter AR, Andersson AF, Tenney A, Lundin D, Tovchigrechko A, Nylander JAA, Brami D, Badger JH, Allen AE, Rusch DB, Hoffman J, Norrby E, Friedman R, Pinhassi J, Venter JC, Bergman B. 2014. Functional Tradeoffs Underpin Salinity-Driven Divergence in Microbial Community Composition. PLOS ONE 9:e89549.

27. Kaushal SS, Likens GE, Pace ML, Utz RM, Haq S, Gorman J, Grese M. 2018. Freshwater salinization syndrome on a continental scale. Proc Natl Acad Sci 115:E574–E583.

28. Rath KM, Rousk J. 2015. Salt effects on the soil microbial decomposer community and their role in organic carbon cycling: A review. Soil Biol Biochem 81:108–123.

29. Trivedi P, Delgado-Baquerizo M, Trivedi C, Hu H, Anderson IC, Jeffries TC, Zhou J, Singh BK. 2016. Microbial regulation of the soil carbon cycle: evidence from gene–enzyme relationships. 11. ISME J 10:2593–2604.

30. Werba JA, Stucy AL, Peralta AL, McCoy MW. 2020. Effects of diversity and coalescence of species assemblages on ecosystem function at the margins of an environmental shift. PeerJ 8:e8608.

31. Thompson PL, Shurin JB. 2012. Regional zooplankton biodiversity provides limited buffering of pond ecosystems against climate change. J Anim Ecol 81:251–259.

32. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J 6:1621–1624.

33. Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb Ecol 75:129–137.

34. Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples: Primers for marine microbiome studies. Environ Microbiol 18:1403–1414.

35. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. Appl Environ Microbiol 79:5112–20.

36. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig W, Glöckner FO. 2014. The SILVA and "All-species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Res 42:D643–D648.

37. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: a versatile open source tool for metagenomics. PeerJ 4:e2584.

38. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately maximum-likelihood trees for large alignments. PLoS ONE 5:e9490.

39. Halekoh U, Højsgaard S. 2014. A Kenward-Roger Approximation and Parametric Bootstrap Methods for Tests in Linear Mixed Models   The R Package pbkrtest. 1. J Stat Softw 59:1–32.

40. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan: Community Ecology Package (2.5-5).

41. Caceres MD, Jansen F. 2016. indicspecies: Relationship Between Species and Groups of Sites (1.7.6).

42. Martiny AC, Treseder K, Pusch G. 2013. Phylogenetic conservatism of functional traits in microorganisms. 4. ISME J 7:830–838.

43. Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, Knelman JE, Darcy JL, Lynch RC, Wickey P, Ferrenberg S. 2013. Patterns and Processes of Microbial Community Assembly. Microbiol Mol Biol Rev 77:342–356.

44. Graham EB, Crump AR, Resch CT, Fansler S, Arntzen E, Kennedy DW, Fredrickson JK, Stegen JC. 2016. Coupling Spatiotemporal Community Assembly Processes to Changes in Microbial Metabolism. Front Microbiol 7.

45. Vellend M. 2010. Conceptual Synthesis in Community Ecology. Q Rev Biol 85:183–206.

46. Logares R, Bråte J, Bertilsson S, Clasen JL, Shalchian-Tabrizi K, Rengefors K. 2009. Infrequent marine–freshwater transitions in the microbial world. Trends Microbiol 17:414–422.

47. Soucy SM, Huang J, Gogarten JP. 2015. Horizontal gene transfer: building the web of life. Nat Rev Genet 16:472–482.

48. Hall JPJ, Brockhurst MA, Harrison E. 2017. Sampling the mobile gene pool: innovation via horizontal gene transfer in bacteria. Philos Trans R Soc B Biol Sci 372:20160424.

49. Rocca JD, Yammine A, Simonin M, Gibert JP. 2022. Protist Predation Influences the Temperature Response of Bacterial Communities. Front Microbiol 13.

50. Cabrerizo MJ, Marañón E. 2021. Grazing Pressure Is Independent of Prey Size in a Generalist Herbivorous Protist: Insights from Experimental Temperature Gradients. Microb Ecol 81:553–562.

51. Åkesson A, Curtsdotter A, Eklöf A, Ebenman B, Norberg J, Barabás G. 2021. The importance of species interactions in eco-evolutionary community dynamics under climate change. 1. Nat Commun 12:4759.

52. Cavicchioli R, Ripple WJ, Timmis KN, Azam F, Bakken LR, Baylis M, Behrenfeld MJ, Boetius A, Boyd PW, Classen AT, Crowther TW, Danovaro R, Foreman CM, Huisman J, Hutchins DA, Jansson JK, Karl DM, Koskella B, Welch DBM, Martiny JBH, Moran MA, Orphan VJ, Reay DS, Remais JV, Rich VI, Singh BK, Stein LY, Stewart FJ, Sullivan MB, Oppen MJH van, Weaver SC, Webb EA, Webster NS. 2019. Scientists' warning to humanity: microorganisms and climate change. Nat Rev Microbiol 1.

53. Graham EB, Averill C, Bond-Lamberty B, Knelman JE, Krause S, Peralta AL, Shade A, Smith AP, Cheng SJ, Fanin N, Freund C, Garcia PE, Gibbons SM, Van Goethem MW, Guebila MB, Kemppinen J, Nowicki RJ, Pausas JG, Reed SP, Rocca J, Sengupta A, Sihi D, Simonin M, Słowiński M, Spawn SA, Sutherland I, Tonkin JD, Wisnoski NI, Zipper SC, Consortium C, Staal A, Arora B, Oldfield C, Dwivedi D, Larson E, Santillan E, Aaron Hogan J, Atkins J, Zheng J, Lembrechts J, Patel K, Copes-Gerbitz K, Winker K, Mudge L, Wong M, Nuñez M, Luoto M, Barnes R. 2021. Toward a Generalizable Framework of Disturbance Ecology Through Crowdsourced Science. Front Ecol Evol 9.

54. Wilson BJ, Servais S, Mazzei V, Kominoski JS, Hu M, Davis SE, Gaiser E, Sklar F, Bauman L, Kelly S, Madden C, Richards J, Rudnick D, Stachelek J, Troxler TG. 2018. Salinity pulses interact with seasonal dry-down to increase ecosystem carbon loss in marshes of the Florida Everglades. Ecol Appl 28:2092–2108.

55. Litchman E. 2010. Invisible invaders: non-pathogenic invasive microbes in aquatic and terrestrial ecosystems. Ecol Lett 13:1560–1572.

56. Hutchins DA, Fu F. 2017. Microorganisms and ocean global change. Nat Microbiol 2:1–11.

FIGURE LEGENDS

Figure 1. Schematic of experimental mesocosm to test the effect of salinity and dispersal of estuarine communities into freshwater coastal ponds (A) and bacterial sequence processing for statistical analyses (B).

Figure 2. Bacterial phylogenetic and taxonomic patterns along a salinity gradient. Ordination based on a Principal Coordinates Analysis depicting bacterial phylogenetic distance (A) and Bray-Curtis dissimilarity distance (modified from Werba et al. 2020) (B), where symbols are colored according to salinity treatment (0, 5, 9, or 13 psu) and symbol shapes represent date of sampling (Day 0, 18, 45). Raincloud plots of phylogenetic diversity along a salinity gradient, where plots represent raw data depicted as filled circles ('rain'), probability distributions ('clouds'), and boxplots displaying mean and 95% confidence intervals (C). Colors represent salinity treatment (orange=0 psu, blue=5 psu, green=9 psu, yellow=13 psu), and shapes represent date of sampling (square=Day 0, circle=18, triangle=45).

Figure 3. Phylogenetic patterns of indicator bacterial taxa. Phylogenetic tree of bacterial taxa that were identified to be statistically associated with salinity treatment (0, 5, 9, 13 psu or combined salt treatment represented by >0 psu); each taxon tip color corresponds to phyla level grouping and is associated with a heat map which represents OTU relative abundance salinity treatment (A). Euler plots represent bacterial OTU that are unique or shared across salinity treatments. The 0* label refers to combined salt treatments (combining 5, 9, 13 psu into a single 'salt' treatment). (B). Average cluster depth estimated using ConcenTrait used to depict phylogenetic distance of bacterial assemblages observed across the salinity gradient (C).

Figure 4. Relationship between carbon mineralization and bacterial phylogenetic diversity (PD), and taxonomic-based community metrics (relative abundance) for Day 45 (last day of experiment).

Tables

Table 1. Summary PERMANOVA output comparing bacterial communities due to the factors salinity, dispersal, and sampling date.

Factor	Df	Sums Sq	Mean Sq	F.Model	R^2	Pr(>F)
Salinity	1	1.275	1.275	28.576	0.172	**<0.001**
Dispersal	4	0.210	0.052	1.175	0.028	0.200
Date	2	0.515	0.258	5.772	0.069	**<0.001**
Salinity:Dispersal	1	0.041	0.041	0.916	0.006	0.509
Salinity:Date	2	0.353	0.177	3.956	0.048	**<0.001**
Dispersal:Date	8	0.290	0.036	0.811	0.039	0.914
Salinity:Dispersal:Date	2	0.063	0.032	0.707	0.008	0.858
Residuals	105	4.686	0.045	0.6304		
Total	125	7.433	1			

Table 2. Summary of Type II Analysis of Variance Table with Kenward-Roger's method comparing phylogenetic diversity. We conducted a linear mixed effects model with 'date', 'salinity', and 'dispersal' as fixed effects and 'block' as a random effect.

Factor	Df	Sum Sq	Mean Sq	F-value	Pr(>F)
Salinity	1	3505	3505.3	3.703	**0.058**
Dispersal	1	79	79.4	0.084	0.773
Date	2	14396	7197.9	7.604	**<0.001**
Dispersal:Salinity	1	265	264.6	0.280	0.598
Dispersal:Date	2	901	450.4	0.476	0.623
Salinity:Date	2	11353	5676.5	5.997	**0.004**
Dispersal:Salinity:Date	2	136	67.8	0.072	0.931
Residuals	79	74778	946.6		

FIGURES

Figure 1. Schematic of experimental mesocosm to test the effect of salinity and dispersal of estuarine communities into freshwater coastal ponds (A) and bacterial sequence processing for statistical analyses (B).

(A) Experimental design

(B) Statistical analyses

Processing of bacterial sequences

Figure 2. Bacterial phylogenetic and taxonomic patterns along a salinity gradient. Ordination based on a Principal Coordinates Analysis depicting bacterial phylogenetic distance (A) and Bray-Curtis dissimilarity distance (modified from Werba et al. 2020) (B), where symbols are colored according to salinity treatment (0, 5, 9, or 13 psu) and symbol shapes represent date of sampling (Day 0, 18, 45). Raincloud plots of phylogenetic diversity along a salinity gradient, where plots represent raw data depicted as filled circles ('rain'), probability distributions ('clouds'), and boxplots displaying mean and 95% confidence intervals (C). Colors represent salinity treatment (orange=0 psu, blue=5 psu, green=9 psu, yellow=13 psu), and shapes represent date of sampling (square=Day 0, circle=18, triangle=45).

Figure 3. Phylogenetic patterns of indicator bacterial taxa. Phylogenetic tree of bacterial taxa that

were identified to be statistically associated with salinity treatment (0, 5, 9, 13 psu or combined

salt treatment represented by >0 psu); each taxon tip color corresponds to phyla level grouping

and is associated with a heat map which represents OTU relative abundance salinity treatment

(A). Euler plots represent bacterial OTU that are unique or shared across salinity treatments. The

0* label refers to combined salt treatments (combining 5, 9, 13 psu into a single 'salt' treatment).

(B). Average cluster depth estimated using ConcenTrait used to depict phylogenetic distance of

bacterial assemblages observed across the salinity gradient (C).

(3A)

(3B)

(3C)

Figure 4. Relationship between carbon mineralization and bacterial phylogenetic diversity (PD), and taxonomic-based community metrics (relative abundance) for Day 45 (last day of experiment).