Little hope for the polyploid endemic Pyrenean Larkspur (*Delphinium

montanum): evidences from population genomics and Ecological Niche**

Modelling

Pascaline Salvado[1], Pere Aymerich Boixader[2], Josep Parera[3], Albert Vila Bonfill[4], Maria

Martin[3], Céline Quelennec[3], Jean-Marc Lewin[5], Valérie Hinoux[1,5] & Joris A. M. Bertrand[1]

[1] Laboratoire Génome et Développement des Plantes (LGDP, UMR 5096 UPVD/CNRS),

Université de Perpignan Via Domitia, France.

[2] C. Barcelona, 29. 08600 Berga, Spain

[3] Fédération des Réserves Naturelles Catalanes, Prades, France

[4] Parc Natural de Capçaleres del Ter i del Freser, Quelralbs, Spain

[5] Association Charles Flahault, Toulouges, France

Corresponding authors: Joris Bertrand (joris.bertrand@univ-perp.fr) and Valérie Hinoux

(valerie.hinoux@univ-perp.fr)

Word count: 6 466

Abstract

Species endemic to restricted geographical ranges represent a particular conservation issue,

be it for their heritage interest. In a context of global change, this is particularly the case for

plants which belong to high-mountain ecosystems and, because of their ecological

requirements, are doomed to survive or disappear on their 'sky islands'. The Pyrenean

Larkspur (*Delphinium montanum*, Ranunculaceae) is endemic to the Eastern part of the

Pyrenees (France and Spain). It is now only observable at a dozen of localities and some

populations show signs of decline, such as a recurrent lack of flowering. Implementing

population genomic approach (e.g. RAD-seq like) is particularly useful to understand genomic

patterns of diversity and differentiation in order to provide recommendations in term of

conservation. However, it remains challenging for species such as *D. montanum* that are

autotetraploid with a large genome size (1C-value > 10 pg) as most methods currently

available were developed for diploid species. A Bayesian framework able to call genotypes

with uncertainty allowed us to assess genetic diversity and population structure in this system.

Our results show evidence for inbreeding (mean G_{IS} = 0.361) within all the populations and

substantial population structure (mean G_{ST} = 0.403) at the metapopulation level. In addition

to a lack of connectivity between populations, spatial projections of Ecological Niche

Modelling analyses under different climatic scenarios predict a dramatic decrease of suitable

habitat for *D. montanum* in the future. Based on these results, we discuss the relevance and

feasibility of different conservation measures.

Introduction

Extinction risk associated to climate change is expected not only to increase but to accelerate for every degree rise in global temperatures (Urban, 2015). In this context, the number of threatened species by reduced population sizes, habitat degradation and habitat fragmentation is increasing. Small and fragmented populations are particularly prone to be affected by drift and inbreeding depression and generally display decreased levels of genetic diversity (e.g. Bergl et al. 2008; Dixo et al. 2009). The increase of anthropogenic and environmental pressures, both being intimately bound, represents an unprecedented threat for vulnerable species. In particular, populations from species of high-mountain ecosystems might not be able to move upward rapidly enough to keep within their thermal tolerances and might be doomed on their 'sky islands' (see McCormack et al. 2009, e.g. Kidane et al. 2019, De Gabriel Hernando et al. 2021). Thus, species endemic to specific mountain ranges represent a major conservation issue, if only for their heritage interest. It is critical to characterize the distribution of genetic diversity within- and between- their populations to evaluate the inbreeding risk and study population structure to better understand how isolated they really are, in order to undertake effective conservation actions (Calevo et al. 2021).

The endangered Pyrenean larkspur *Delphinium montanum* DC., 1815 perfectly illustrates this situation. This plant (Family Ranunculaceae), endemic to the Eastern part of the Pyrenees (France and Spain), is now only observable at a dozen of localities (two already got extinct over the 20[th] century) and its global population size was estimated to range between 7 800 to 10 300 individuals in total (Aymerich et al. 2020). It occurs between alpine and subalpine levels, between 1 600 and 2 400 m (Lopéz-Pujol et al. 2007), from Serra del Cadí-Pedraforca (Fig. 1, localities 4, 5, 8, and 9) up to the Puigmal massif (localities 1, 3 and 6) via Tosa d'Alp massif (locality 7). There, the distance between the most distant sites is about 60

km, and only one locality is located outside this area: Nohèdes (locality 2), in the Madres

massif. This latter locality is separated from the Puigmal massif populations by the pit of

Cerdagne-Conflent, which could have formed a historical barrier to gene flow. In terms of

population size, Cadí-Pedraforca is the most important with up to 70% of the global population

size (5 450-6 700 individuals). The sites of the Puigmal massif represent 25 to 30% of the global

population (2 200-2 700 individuals). Localities of Madres massif (2) and Tosa d'Alp (7) are

numerically low (less than 300, from which none are actually flowering and 35 individuals,

respectively).

D. montanum is endangered as a result of increasing levels of multiple threats and is

included in the IUCN (International Union for Conservation of Nature) red list of threatened

species, in the Vulnerable (VU) category. Some of its populations are located nearby touristic

sites (*i.e.* mountain resorts or hiking trails) and are negatively impacted by trampling (Simon

et al., 2001). Flowers and fruits predation by herbivores, particularly Pyrenean chamois/isards

(*Rupicapra pyrenaica pyrenaica*) was also shown to be responsible for the seed bank's

decrease in some populations (Simon et al. 2001, Aymerich 2003). Importantly, differences in

population dynamics are observable from one population to another. Some populations are

stable, whereas others show decrease in population size (4-Pedraforca) and/or absence of

flowering individuals since 2011 (2-Nohèdes). Two studies previously investigated population

genetics of *D. montanum* based on allozymes (Simon et al. 2001; Lopéz-Pujol et al. 2007). They

already suggested relatively low effective size as well as, evidence of inbreeding within-

populations and significant population structure.

The features of the genome of the species also make *Delphinium montanum*

particularly challenging from a technical and methodological point of view. *D. montanum* is an

autotetraploid species (Simon et al. 2001; Lopéz-Pujol et al. 2007). Our recent flow cytometry

estimates of 10.32 pg/1C for this species (*i.e.* about twice the 1C-value reported for several

other diploid *Delphinium* species) corroborates previous knowledge and confirms the

important genome size of *D. montanum* (Bertrand et al. unpublished data). Implementing a

population genomic approach on *D. montanum* thus requires genome complexity reduction

protocols such as Reduction site-associated DNA sequencing (or RAD-seq, Baird 2008).

However, the difficulty to infer genotype and allele frequencies on polyploid species with such

techniques as well as the restricted set of methods available to analyze such data (see

Meirmans and van Tienderen 2013; Dufresne et al. 2014; Meirmans et al. 2018) is likely to

explain the relatively low number of population genomics studies dealing with RAD-seq like

techniques on non-model polyploid organisms (Clevenger et al. 2015; van de Peer et al. 2017,

Ahmad 2021).

Out the handful of studies that have addressed population genomic questions on

polyploid organisms without reference genome, some have considered the genotypes as

diploid-like data, an assumption that allows to use classical variant calling tools with default

settings (e.g. Brandrud et al. 2017, see also Brandrud et al. 2019; 2020 and Záveská et al.

2019). Consensus allelic information may also be coded in the form of ambiguous sites at

heterozygous positions (e.g. Wagner et al. 2020). At last, a couple of methods are able to

incorporate ploidy-aware calling algorithms and use genotype likelihoods to infer tetraploid

genotypes (e.g. Záveská et al. 2019; Brandrud et al. 2020; Karbstein et al. 2020; Ahmad et al.

2021). In studies for which both kind of methods were used, the latter one generally outputted

a higher number of variable sites but signals were found to be overall congruent (Záveská et

al. 2019; Brandrud et al. 2020).

The recent studies of Záveská et al. (2019), Brandrud et al. (2020) and Ahmad et al.

(2021) used the RAD-seq loci from the catalog generated in the popular pipeline Stacks

(Catchen et al. 2011; Catchen et al. 2013) to generate a synthetic reference onto which the

raw reads are mapped back before to infer tetraploid genotypes with the approach

implemented in EBG (Empirical Bayes Genotyping in Polyploids, Blischak et al. 2018, see also

Blischak et al. 2016). As a putative alternative, Clark et al. (2019) proposed a package called

PolyRAD to call genotypes with uncertainty from polyploid data in an R environment. PolyRAD

can deal with multiple levels of ploidy and is able to use for example population structure,

linkage disequilibrium and/or self-fertilization as priors to estimate genotype probabilities. To

our best knowledge, this promising and convenient method has been rarely used to

implement a population genomic approach on natural populations of polyploid non-model

organisms.

In this study, we first aimed at reevaluating the conservation status of *Delphinium*

montanum following such population genomics approach. We used a protocol called

normalized Genotyping By Sequencing (nGBS) that provided us with thousands of genetic

markers (SNPs) from which we inferred patterns of diversity and differentiation with an

unprecedented resolution in order to quantify inbreeding and level of connectivity between

populations. We then used an Ecological Niche Modelling (ENM) approach to better

understand the bioclimatic features that constraint the current geographic distribution of *D.*

montanum. Based on several Global Climate Models (CGM), we also predicted the putative

evolution of the geographic distribution of the species in the future, at different time periods

(2011-2040, 2041-2070 and 2071-2100) and under different, more or less pessimistic but still

realistic scenarios of evolution of global warming (or RCPs for Representative Concentration

Pathways).

Materials and Methods

Population sampling

We sampled a total of 106 individuals from 9 localities representative of the geographic

distribution of *Delphinium montanum* (in the Eastern Pyrenees: 3 sites in France and 6 in

Spain) between July and October 2020 (Table 1, Fig. 1, see also Supplementary Appendix S1).

At each locality, we collected about 1-2 cm^2 of leaf tissues that were stored in 90% ethanol at

4°C until DNA extraction. *Delphinium montanum* is not legally protected in France but is

considered as vulnerable (VU) on the Red List of threatened species of IUCN France (UICN

France, FCBN, AFB, MNHN, 2018). It is legally protected in Catalonia (Decret 172/2008. Catàleg

de flora amenaçada de Catalunya) and Spain (Orden AAA/1771/2015. Modificación del

catálogo de especies amenazadas). In France, for all sampling done in natural reserves, special

authorizations were granted by the 'Direction Départementale des Territoires et de la Mer 66'

(DDTM 66). In Spain, we asked for and obtained special permits from the Servei de Fauna i

Flora de la Generalitat de Catalunya and Natural Parks (Cadí-Moixeró and Ter-Freser).

Molecular procedures

Genomic DNA extraction and genotyping were subcontracted to LGC Genomics GmbH (Berlin,

Germany). Genotyping-By-Sequencing (GBS) was performed following a specific protocol

called nGBS for 'normalised GBS'. This double digest Restriction site-Associated DNA seq (or

RAD-seq) like protocol relies on two restriction enzymes (PstI and ApeKI, in our case) to reduce

D. montanum genome complexity and includes a normalization step that aims at avoiding

repetitive regions. The resulting 106 individually barcoded libraries were sequenced in paired-

end mode (2 x 250 bp) on an Illumina NovaSeq 6000 with an expectation of a minimum

number of 1.5 million read pairs per sample.

Genomic data processing

We used Stacks v.2.41 (Catchen et al. 2011; 2013) to build loci from Illumina reads, *de novo* (*i.e.* without aligning reads to a reference genome). Stacks consists of a wrapper of several scripts that are usually run sequentially including: *process_radtags* to demultiplex and clean reads, *denovo_map.pl* to build loci within individuals, create a catalog and match all samples against it and *populations* to further filter the SNPs obtained at the population level and compute basic population genetic statistics. We first optimised several key parameters of the pipeline: *-m* (the minimum number of identical raw reads required to form a putative allele), *-M* (the number of mismatches allowed between alleles to form a locus) and *-n* (the number of mismatches allowed between loci during construction of the catalog) on a subset of 12 individuals representative of the whole data set (*i.e.* geographic origin and coverage). As recommended by several authors, we varied M and n (fixing $M = n$) while keeping $m = 3$ (see Paris et al. 2017; Rochette & Catchen 2017). The combination *-m* 3, *-M* 3 and *-n* 3 was found to be the most suitable to maximize the number of SNPs, assembled and polymorphic loci in our case and was used to run Stacks on the whole data set (see Supplementary Appendix S2). As *D. montanum* is a tetraploid species, we also specified the option '*-X ustacks:--max_locus_stacks* 5' (in the *ustacks* script's) to allow the number of alleles per site to reach the number of 5 instead or 3 by default. However, as the *populations* script is unable to deal with polyploid genotypes, we followed a different approach to call variants and complete genomic data processing.

We then used polyRAD (Clark et al. 2019), a method implemented in an R-package to call genotypes with uncertainty from sequencing data in polyploids. The Bayesian genotype caller implemented in polyRAD is able to import read depth from the catalog and matches files from Stacks through a specific function called polyRAD::read_stacks() to further estimate the

probability of each possible genotype for each individual and each locus, taking into account

features such as possible levels of ploidy and population structure as priors. In order to get

similar filtering parameters than with Stacks and following the polyRAD author's guidelines,

we used the following arguments -*min.ind.with.minor. alleles= 5*, -*min.ind.with.reads= 84* and

a ploidy of 4. We then plotted the distribution of the ratio of the individual over expected

heterozygosity H_{ind}/H_E. The peak of the distribution of H_{ind}/H_E by locus (0.2) was considered to

represent well-behaved markers from which inbreeding (*F*) was estimated at 0.73. From this

estimate of *F*, we then simulated the expected distribution of H_{ind}/H_E, if all markers were

behaving in a Mendelian fashion. Based on the distribution, we defined minimum and

maximum thresholds of 0 and 0.406 comprising 95% of the observations and filtered out

markers for which H_{ind}/H_E were outside this interval. We then tested overdispersion (*i.e.* in our

case, how much does read depth distribution deviate from what would be expected under

binomial distribution) and adjusted its parameter to 6 (from a range of 2 to 14). This

information was used to call genotypes while taking into account population genetic structure.

The most likely genotypes of the remaining 5 095 loci were then exported in several formats

such as genind objects or Structure (.str).

Population genomic analyses

To get an overview of the overall genetic diversity and differentiation among individuals and

populations, we first performed a Principal Component Analysis (PCA) based on a matrix of

106 individuals (as rows) and 5 095 SNPs (as columns) coded as a genind object with the R-

package 'adegenet' (Jombart 2008). We then used Genodive v.3.04 (Meirmans 2020) to

compute expected and observed heterozygosity (H_E and H_O, respectively) as well as the

number of alleles (*A*) and the number of private alleles was computed with the R-package

'poppr' (Kamvar et al. 2014). The deviation from panmixia was evaluated by computing G_{IS}

and the statistical significance of the obtained values was estimated based on 10 000

permutations, also in Genodive. Overall and pairwise genetic differentiation was also assessed

based on *G*-statistics (G_{ST}, Nei 1987) as implemented in Genodive in a similar manner. Pattern

of Isolation By Distance (IBD) was assessed by examining the relationship between linearized

G_{ST}s values (*i.e.* ln(G_{ST}/(1- G_{ST})) and ln-geographical distances (after Rousset 1997) and its

statistical significance was tested with a Mantel test between matrices of pairwise G_{ST} and

geographical distance.

To further investigate population structure and characterize putative

migration/admixture event, we used sNMF (Frichot et al. 2014) as implemented in the R-

package LEA (Frichot & François 2015) to estimate individual ancestry coefficients based on

sparse non-negative matrix factorization algorithms. sNMF is particularly suitable for our

genotype dataset as it can deal with polyploid data and is fast enough to be applied on

hundreds of individuals and thousands of markers with reasonable computation time. The

number of genetic clusters was varied from *K* = 1 to 10, and analyses were run with 10

replicates at each value of *K*.

Ecological Niche Modelling

In order to assess the future trends of spatial dynamics of *D. montanum* distribution in a

context of global change, we conducted Ecological Niche Modelling (ENM) analyses based on

known occurrences of the species. Based on 19 bioclimatic layers we downloaded from

CHELSA v.2.1 (Karger et al. 2017), we aimed to i) model current environmental niche and ii)

predict the spatial evolution of the niche in the future. The bioclimatic layers available in

CHELSA are similar to those available in WORLDCLIM (Fick et al. 2017) in a way that they

consist of downscaled Global Climate Models (GCM) output temperature and precipitation

estimates at a maximal resolution of 30 arc seconds (approximately 1 km). In the current

study, we used the most recent version of CHELSA as the methodology and bias correction it

implements have been shown to outperforms WORLDCLIM in mountainous environments

(Karger et al. 2017; Bobrowski et al. 2021). Current climate data correspond to time averaged

variables over the period 1981-2010.

Future climates correspond to estimates from phase 3b of the Inter-Sectoral Impact

Model Intercomparison Project (ISIMIP3b) based on output of phase 6 of the Coupled Model

Intercomparison Project (CMIP6). They consist of the output of five GCMs (GFDL-ESM4,

UKESM1-0-LL, MPI-ESM1-2-HR, IPSL-CM6A-LR and MRI-ESM2-0) and three climate scenario

specifiers (ssp126, ssp370, ssp585). Climate scenarios specifiers reflect different

Representative Concentration Pathways (RCPs) associated with different radiative forcing

values: 2.6, 7.0 and 8.5 W/m^2 that can be considered as a plausible range of greenhouse gas

concentrations that appear to be optimistic, very likely and pessimistic in term of climate

change, respectively. Annually averaged data were downloaded for three future time periods:

2011-2040 ("2035"), 2041-2070 ("2055") and 2071-2100 ("2085").

The ENM analyses were conducted thanks to the R-package ENMwizard v.0.3.7

(Heming et al. 2018, see also Gutiérrez et al. 2019; Bagley et al. 2020). We first spatially filtered

occurrences to keep only those that were at least 1 km away from each other using the R-

package 'spThin' (Aiello-Lammens et al. 2015). Due to the relatively small spatial scale of the

area under study, this resulted in considering a total of 9 out the 106 occurrences initially

available. We then examined raw values for the 19 bioclimatic variables extracted at each

occurrence point and noticed that the variable bio19 ("precipitation of coldest quarter")

displayed obviously aberrant (*i.e.* negative) values. This variable bio19 was thus discarded

from subsequent analyses. From the 18 remaining variables (bio1 to bio18), we selected the

less correlated ones (Person correlation coefficient < 0.75) thanks to the R-package 'caret'

(Kuhn 2019) and kept five variables: bio5 ("mean daily maximum air temperature of the

coldest month"), bio7 ("annual range of air temperature"), bio14 ("precipitation amount of

the driest month") and bio15 ("precipitation seasonality") for further analyses.

The calibration area for the models was created as a buffer of 1.5° around the

minimum convex polygon encompassing all occurrences. We used the maximum entropy

method (implemented in MaxEnt ver. 3.4.1, Phillips et al. 2006, 2017) to calibrate model and

evaluated model performance thanks to the package ENMeval (Muscarella et al. 2014) as

implemented in ENMwizard. We evaluated models using a geographic partition scheme of

type "block" and optimized two parameters of MaxEnt: the Regularization Multipliers (RM)

and the Feature Classes (FC). RM was varied from 0.5 to 4.5, incremented by 0.5 whereas a

suite of 15 FCs (L, for Linear, P, for Product, Q, for Quadratic and H for Hinge) or combination

of them were evaluated: L, P, Q, H, LP, LQ, LH, PQ, PH, QH, LPQ, LPH, LQH, PQH, LPQH, resulting

in a total of 135 models. Model selection was done by computing the corrected Akaike

Information Criterion ("LowAIC") and the Area Under the receiver operating characteristic

Curve ("AUC") in the function ENMWizard::calib_mdl_b(). The best selected model was then

spatially projected as current niche. For future predictions, a spatial projection of the

consensus of the five CGMs was performed for each time period and each RCPs. Area

calculations were carried out after applying a 10-percentile (x10ptp) and a Maximum training

sensitivity plus specificity (mtss) threshold rules.

Results

Genetic diversity and differentiation

We obtained a total of 159 million read pairs across the 106 individuals. We assembled a total

of 210 756 loci including 22 911 SNPs with Stacks. After filtering the data with polyRAD, we

finally obtained 32 728 loci including 5 095 SNPs.

The two first PCA axes (PC1 and PC2) represented 14.85% and 9.21% of the total

genetic variance, respectively (Fig. 2A). This analysis is consistent with limited amount of

within-population differentiation and a clear pattern of between-population differentiation.

Overall, populations are arranged according to geography, especially along PC1. In term of

genetic diversity, all populations display a significant deviation from panmixia: mean G_{IS} =

0.361, $p < 0.05$ and population G_{IS} values ranging from 0.315 to 0.426 (all $p < 0.05$). However,

G_{IS} values appear as similar from one population to another (this holds true for the number of

total alleles A, too, see Table 1). The number of private alleles Ap is consistent with the relative

level of geographic isolation of the populations.

The degree of genetic differentiation was found to be important overall (mean G_{ST}

value = 0.403, $p < 0.05$) and pairwise (with values ranging from 0.156 to 0.527, Supplementary

Appendix S3). There is a clear and significant pattern of Isolation By Distance across the

dataset ($r = 0.601$, $p = 0.001$, Fig. 3) indicating that gene flow is limited in space at the scale of

the *D. montanum* geographic range. The sNMF analysis is also consistent with a strong

population structure arranged with geography. Cross-entropy was the lowest for K= 6 (Fig. 2B

and 2C). Each cluster corresponds to a sampling site with two exceptions. The individuals from

the locality of Orri de Baix (1) were grouped together with the close locality of Cambre d'Aze

(6), and the Western localities of the Serra del Cadí (Bastanist (8), Serra Pedregosa (5) and to

a lesser extent Pedraforca (4)) formed another group. Only the individuals of Pedraforca (4)

showed mixed ancestry coefficients.

Suitable habitats for *D. montanum* and prediction of its spatial dynamics

ENMeval analyses identified RM = 2 and a combination of Quadratic and Hinge feature (QH)

classes as the best-performing parameters for calibrating the final ENM following the

"LowAIC" and M=2 and a combination of Product and Hinge feature (PH) classes following the

"AUC" optimality criteria. The first model had the best corrected Akaike Information Criterion

(AICc) score (152.25) and an AICc weight of 0.33 whereas the second one had an AICc score of

174.63 and an AICc weight of $4.53.10^{-6}$. These two models had mean omission rates of 0.375

and 0.25 for the 10th percentile and the lowest presence training thresholds, respectively, as

well as a mean test AUC of 0.988. Values of these diagnostic metrics are provided for all

candidate models in Supplementary Appendix S4. Current suitable area for *D. montanum* was

found to be 2 670 and 2 365 km^2 following "LowAIC" and "AUC" selection model criteria,

respectively (the values were identical whatever the threshold method used). All future

consensus spatial projections predict a dramatic decrease of the suitable area (Table 2 and

Fig. 4). For RCPs 7.0 and 8.5, the suitable area would have completely disappeared for the time

period 2071-2100 (but not for RCP 2.6). We also notice some slightly different outcomes

depending on the selection model criteria with RCP 8.5 scenarios giving sometimes higher

values than RCP 7.0, and even RCP 2.6 (e.g. time period 2011-2040, "Low AIC" criterion).

Discussion

Conservation genomics on a polyploid species with important genome size

The use of RAD-seq like (nGBS) data with the method implemented in PolyRAD allowed us to

successfully address the challenge of conducting a population genomic approach on a plant

species that is both polyploid (autotetraploid) and has an important genome size (> 10 Gbp).

It confirms the convenience of the workflow we used, specifically calling genotypes in an R-

environment, as an alternative to the ones followed in the handful of studies recently

published on similar systems (e.g. Záveská et al. 2019; Brandrud et al. 2020; Karbstein et al.

2020; Ahmad et al. 2021). From the > 5 000 SNPs we could rely on, we were able to get an

unprecedented genomic picture of the diversity and the differentiation of the species in order

to propose appropriate conservation recommendations. The results we obtained at the

genome scale are overall consistent with previous knowledge on population genetics of *D.*

montanum (see Simon et al. 2001; López-Pujol 2007).

A set of populations isolated on sky islands

Our results confirm that populations of *D. montanum* can be seen as a set of island-like

dwelling systems separated from each other by a matrix of unsuitable habitats and that gene

flow has little to no chance to be efficiently maintained within this metapopulation. All

populations considered in this study were found to display deficit in heterozygotes ($G_{IS} > 0.315$,

$p < 0.05$). These results confirm previous findings of Simon et al. (2001) and López-Pujol et al.

(2007) and show that the autotetraploid status of the species does not impede relatively low

levels of heterozygosity ($H_O < 0.17$). Polyploidy is expected to be associated with increased

genetic variation, a characteristic that provides more diversity to adapt (especially to extreme

environments such as mountainous ones) and allows buffering effect against deleterious

mutations (see van de Peer et al. 2021 and references therein). From a conservation point of

view, polyploidy does not seem to be associated with increased risk of extinction even though

plant with big genome size may be more prone to be under threat (Vinogradov 2003; Pandit

et al. 2011). As two other closely related congeneric species of *Delphinium montanum*: *D.*

dubium and *D. oxysepalum*, that are endemic to the Alps and the Carpathians, respectively,

are also tetraploid (López-Pujol et al. 2007), we speculate that polyploidy may have allowed

to these species to adapt and survive into these mountainous environments. The open

question is whether current levels of genetic diversity may be sufficient or not to allow *D.*

montanum populations to overcome inbreeding depression and provide sufficient variation

for adaptation in a context of global change. It should be noted that levels of genetic diversity

are overall similar among the populations investigated and we could not associate obvious

signs of decline of some of them (e.g. Nohèdes, Pedraforca) to decreased levels in genetic

diversity or evidence for deviation from panmixia. This suggests that the evidences of possible

decrease in effective size may be too recent to be detectable at the genetic level (see Peery

et al. 2012). Unfortunately, the limited number of available methods for polyploids makes

effective size estimation and test for bottleneck events cumbersome to perform in such

systems.

The PCA, the clustering analyses, the F_{ST} values as well as the pattern of Isolation By

Distance, further confirm limited gene flow between populations and the existence of a

population structure that is consistent with geography. The most likely number of clusters

found in the sNMF analysis (*K*=6) globally distinguishes populations from the Serra del Cadí

(Bastanist (8), Serra Pedregosa (5) and Pedraforca (4)) from those of the Puigmal range (Orri

de Baix (1) and Cambre d'Aze (6)) plus the satellite populations of Vallter (3), Nohèdes (2),

Tosa d'Alp (7) and Torreta de Cadí (9) (which is the westernmost locality, a bit distant from

the others in the Serra del Cadí). Only the locality of Pedraforca (4) shows some ambiguity in

population assignment that could indicate a mixed origin. At *K*=2, the intermediate status of

Tosa d'Alp (7) (and to a lesser extent of Pedraforca (4)) are also in agreement with its

intermediate geographic position and may reflect an ancestral split between Northern and

Southern populations. Altogether, these results support that the insect pollinators (mainly

bumblebees) and the natural dispersal of seeds produced (barochoric) are not able to

maintain gene flow over the geographic distances involved.

Climate change as the main threat for *Delphinium montanum*

The two previous studies of Simon et al. (2001) and López-Pujol et al. (2007) have listed several threats to explain the decline of *D. montanum*, at least in some particular populations. According to Simon et al. (2001), natural disturbance of habitats due to rock falls and avalanches, trampling because of an overall increase in human visits and animal predation (e.g. caterpillar, Pyrenean chamois/isards and perhaps red and roe deers) as well as competition for insect pollinators with simultaneously blooming species (e.g. *Aconitum* spp.) have thus been considered as more or less serious problems for *D. montanum*. The most recent study of López-Pujol et al. (2007) also emphasized on the consequences of the erosion of genetic diversity, especially the loss of rare alleles associated with weak possibility of gene flow between localities that could be however compensated by relatively high effective size in several populations. Without ruling out all these threats on *D. montanum* populations, our results rather support climate change as the main threat for this species. The future predictions of the area of habitat suitability all agree that the spatial extent of the ecological niche of *D. montanum* is very likely to dramatically decrease, and even disappear by the end of the 21th century, with some differences based on the various scenarios we considered. The current suitable habitat for *D. montanum* has an area that was estimated at about 2500 km^2, an area that would decrease to about 500 km^2 (*i.e.* -75 %), by 2040 whatever the scenario envisaged. By 2070, this area would again have decreased to about 150 km^2 based on RCP 2.6 and up to less than 50 km^2 based on the two other scenarios (RCP 7 and RCP 8.5). A Representative Concentration Pathway of 2.6 W/m^2 has to be considered as a rather optimistic pathway (that would keep global temperature rise below 2°C by 2100). It would require that CO_2 emissions start declining by 2020 and go to zero by 2100 (as well as declines

in emission of other greenhouses gases, such as CH_4, SO_2). RCP 7 is usually considered as a

baseline rather than a mitigation target. According to RCP 8.5, emissions would continue to

rise throughout the 21[st] century and can be considered as a pessimistic scenario (with an

increase in average temperature in the order of 4°C by 2100). The various output of these

scenarios likely explains why the bioclimatic niche of *D. montanum* would have completely

disappeared by 2100 for RCP 7 and 8.5 but could have persisted and even started to increase

again to about 10% of its current suitable area following the RCP 2.6 scenario.

Environmental changes as the main threat for *D. montanum* populations is also

supported from field observations. The population of Nohèdes (2) have not flowered since

2011. The individuals there are very small (*i.e.* a few centimetres high) and remain at a

vegetative stage. On this site, we could not notice any evidence of avalanche or landslide that

could have altered the habitat. In addition, other plants species present in the vicinity (e.g.

Orchis spitzelii, Maianthemum bifolium and *Convallaria majalis*) seem to suffer from flowering

default, further supporting that this phenomenon could be non-specific to *D. montanum* and

perhaps caused by bioclimatic or at least environmental modifications. There, output from

Regional Climate Models (RCMs) predict an increase in mean annual temperature that could

reach 3.1 to 4.5°C (+1.5°C already observed since the mean of the period 1961-1990)

associated with a decrease in precipitation of up to 25% and perhaps more compared to

present-day conditions (Lespinas et al. 2014). According to the same study, precipitations

would mostly decrease in summer (-40%) and in spring (-20%) but would remain similar the

rest of the year.

Recommendations for conservation

Ex-situ conservation would be a way to preserve the genetic diversity of *D. montanum*, but

also a bridge for future *in-situ* conservation efforts. A seed collection of different populations

of *D. montanum* could be stored in a bank. However, this measure would be of limited interest

if the decline's cause of *D. montanum* is actually climate change as keeping dormant seeds will

probably not allow this species to adapt to future global changes. Alternatively, conserving

genetic diversity in a botanical garden may be a good option to test whether and if so, leave

time to plants to adapt to rising temperatures and rainfall modifications. Nevertheless, this

approach has potential drawbacks like artificial selection and habitat conversion if *D.*

montanum adapts to *ex-situ* environment, which may differ from their initial natural

environment (Ren et al. 2014). Moreover, if, growing individuals from the all 9 localities

studied in an appropriate botanical garden could be a way to set-up *in-situ* conservation,

preliminary experiments have showed that *cultivating D. montanum* under experimental

conditions in a greenhouse is very difficult because of its environmental requirements

(Aymerich, pers. comm.). This confirms that if any, botanical garden would have to be set up

at a relatively high elevation (*i.e.* > 1500 m), where environmental conditions are similar to

the ecological optimum of the species.

We can identify three main methods of *in-situ* conservation that could be envisaged

for *D. montanum* (see Godefroid et al. 2011; Ren et al. 2014; Mashinski & Albrecht 2017):

reinforcement, translocation and managed relocation (or assisted migration). The first one is

reinforcement, which consists in adding *ex-situ* grown individuals in their natural population

of origin to increase the total number of individuals, increase the spatial occupancy of the

species and thereby reduce extinction risk. On the one hand, the increase in population size

provided by reinforcement could contribute to counteract the effect of genetic drift. On the

other hand, it would have a limited impact to impede inbreeding as all the individuals would

originate from the same genetic pool. The second method consists in translocating individuals

(seeds, seedlings or adult plants) from a population to another one with the same

ecogeography, to increase genetic diversity (the so-called "genetic rescue") and thus increase

the likelihood of population persistence in a changing climate (Mashinski and Albrecht 2017).

Our results show that *D. montanum* populations harbor substantial levels of pairwise genetic

differentiation which suggest that they have exchanged few or no genetic migrants for many

generations and have possibly evolved distinct adaptative differences. As a consequence,

gene pool mixing may not be appropriate, since the risk of outbreeding depression could be

non-negligible between individuals from the different part of the geographic distribution of *D.*

montanum. It could be thus careful to first experiment *ex-situ* crossings to check if hybrids are

viable, and do not suffer from outbreeding depression. However, the genetic benefits of

mixing source populations during genetic rescue have been shown to generally outweigh the

risks of outbreeding depression and loss of local adaptation (see Whiteley et al. 2015;

Frankham 2015; Ralls et al. 2018, see also Liddell 2021). For example, translocations with

mixed source populations of *Arnica montana* showed that genetic variation remained similar

or increased across successive environments and some authors documented a 10-fold

increase in population size following gene flow between wild Trinidadian guppy populations

(Albretch and Edwards 2020; Fitzpatrick et al. 2020).

Finally, the third method consists in introducing a new population in a suitable new

ecosystem, where *D. montanum* has not been observed yet. This method is called managed

relocation or assisted migration (Mashinski and Albrecht 2017). However, it may represent a

risk of modifying the ecosystem, by creating competition for space and pollinators with other

plants naturally present, or by introducing new pathogens. Thus, these three methods of

reinforcement, translocation, and managed relocation can present risks of outbreeding

depression and ecosystem destabilization, and their benefits may not outweigh those risks.

Moreover, since the main threat for *D. montanum* is climate change, on a longer-scale, all

those efforts could be unsuccessful. Keeping individuals in a botanical garden seems to be the

least harmful and risky method for natural populations today and it could be a good start for

crossing experiments between populations.

For the very peculiar situation of Nohèdes's locality (2), which seems to be the most

endangered population at the moment, there is still the question of the lack of flowering. To

test whether this problem may be associated with the absence of a specific environmental

stimulus, we may propose to either i) experimentally modify temperature/water supply, *in*

situ, ii) translocate some individuals to localities where *D. montanum* seem healthy (e.g. Cadí

range) and see if flowering can be restored and/or iii) experimentally modify

temperature/water supply in the lab to better understand the conditions required for

flowering and figure out the limiting factor in the wild. Alternatively, it may be due to the fact

that that this locality is formed by immature individuals and that the local population itself

may be recovering from an important mortality event. At the moment, we have no further

arguments to support this scenario.

Conclusion

Overall, our results confirm that the remaining populations of *D. montanum* show evidence of

decreased genetic diversity and evidence of inbreeding at a genomic scale. The degree of

genetic structure observable is consistent with a set of strongly isolated populations no longer

able to maintain gene flow between their sky islands. In this context, *in-situ* conservation

methods are likely to have an effect on the status of the species. Reinforcement would help

counteracting decrease in local population sizes, and translocation would help increasing

genetic diversity, buffering inbreeding depression and perhaps, bring the variation required

for *D. montanum* to adapt. This being said, *ex-situ* crossings are first recommended to verify

the absence of outbreeding depression as local genetic pools were found to differ

substantially. As our results support that environmental change may represent the main risk

of extinction for *D. montanum* all these measures may be however considered as of limited

interest on the long term, if the ecological niche of the species disappears.

500   Acknowledgements

This study was funded by the interreg-POCTEFA program (FLORALAB Project EFA294/19) and

is set within the framework of the "Laboratoires d'Excellences (LABEX)" TULIP (ANR-10-LBX-

41) as well as within the framework "École Universitaire de Recherche (EUR)" TULIP-GS (ANR-

18-EURE-0019). We also thank Xavier Oliver Martínez-Fornés, Sandra Mendez, Kimberley

Goudedranche, Lucie Schaad, Irene Abad, Queralt Tor and all other contributors from the

FLORALAB network for the help provided in the field.

508   Authors Contributions

509   PS, VH, JAMB analysed the data and wrote the manuscript. PAB, JP, AVB, MM, CC, JML and

510   JAMB contributed to fieldwork. MM, CC and VH coordinated the FLORALAB network and/or

511   project aspects related to this work and JAMB contributed to the conception of the study. All

512   authors have read and approved the manuscript.

514   Conflicts of interest

515   The authors declare no conflict of interest.

Data Archiving

518 Sequencing data have been submitted to the European Nucleotide Archive (ENA;

519 https://www.ebi.ac.uk.ena) under accession n°PRJEB46773.

References

522 Ahmad M, Leroy T, Krigas N, Temsch EM, Weiss-Scheeweiss H, Lexer C, et al. (2021) Spatial

523  and ecological drivers of population structure in *Alkanna tinctiria* (Boraginaceae), a

524  polyploid medicinal herb. Front. Plant. Sci. 12: 706574.

525 Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B, Anderson RP (2015) spThin: an R

526  package for spatial thinning of species occurrence records for use in ecological niche

527  models. Ecography 38: 541-545.

528 Albrecht MA, Edwards CE (2020) Genetic monitoring to assess the success of restoring rare

529  plant populations with mixed gene pools. Mol. Ecol. 29: 4037-4039.

530 Aymerich P (2003) Effects of wild ungulates predation in the conservation of rare plants: two

531  study cases of the Eastern Pyrenees, Acta Bot. Barc. 49: 147-164.

532 Aymerich P, Martínez-Fornés, XO, Mendez S, Mangeot A, Martin M, Tenas B (2019) Seguiment

533  de l'endemisme dels Pireneus orientals *Delphinium montanum* per la xarva

534  transfronterera FloraCat, Actes del XII Colloqui Internacional de Botànica

535  Pirenaica - Cantàbrica, 147-165.

536 Bagley JC, Heming NM, Gutiérrez EE, Devisetty UK, Mock KE, Eckert AJ et al. (2020)

537  Genotyping-by-sequencing and ecological niche modeling illuminate phylogeography,

538  admixture, and Pleistocene range dynamics in quaking aspen (*Populus tremuloides*).

539  Ecol. Evol. 10: 4609-4629.

Baird NA, Etter PD, Atwood TS, Currey MC, Shiver AL, Lewis ZA et al. (2008) Rapid SNP

discovery and genetic mapping using sequenced RAD markers. PloS ONE 3: e3376.

Bergl RA, Bradley BJ, Nsubuga A, Vigilant L. (2008) Effects of habitat fragmentation, population

size and demographic history on genetic diversity: The Cross River Gorilla in a

comparative context. Am. J. Primatol. 70: 848-859.

Blischak PD, Kubatko LS, Wolfe AD (2016) Accounting for genotype uncertainty in the

estimation of allele frequencies in autopolyploids. Mol. Ecol. Resour. 16: 742-754.

Blischak PD, Kubatko LS, Wolfe AD (2018) SNP genotyping and parameter estimation in

polyploids using low-coverage sequencing data. Bioinformatics 34: 407-415.

Bobrowski M, Weidinger J, Schickhoff U (2021) Is new always better? Frontiers in global climate

datasets for modelling treeline species in the Himalayas. Atmosphere 12: 543.

Brandrud MK, Paun O, Lorenzo MT, Nordal I, Brysting AK (2017) RADseq provides evidence for

parallel ecotypic divergence in the autotetraploid *Cochlearia officinalis* in Northern

Norway. Sci. Rep. 7: 5573.

Brandrud MK, Paun O, Lorenz R, Baar J, Hedrén M (2019) Restriction-site associated DNA

sequencing of *Nigritella* and *Gymnadenia* (Orchidaceae). Mol. Phylogenetics Evol. 136:

21-28.

Brandrud MK, Baar J, Lorenzo MT, Athanasiadis A, Bateman RM, Chase MW, Hedrén M et al.

(2020). Phylogenomic relationships of diploids and the origins of allotetraploids in

Dactylorhiza (Orchidaceae). Syst. Biol. 69: 91-109.

Calevo J, Gargiulo R, Bersweden L, Viruel J, González-Montelongo C, Rebbas, K et al. (2021)

Molecular evidence of species- and subspecies-level distinctions in the rare *Orchis*

patens s.l. and implications for conservation. Biodivers. Conserv. 30: 1293-1314.

Catchen J, Amores A, Hohenlohe P, Cresko W, Postlethwait J (2011) Stacks: building and

genotyping loci de novo from short-read sequences. G3 - Genes Genom. Genet. 1: 171-

182.

Catchen J., Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool

set for population genomics. Mol. Ecol. 22: 3124-3140.

Clark LV, Lipka AE, Sacks EJ (2019). polyRAD: Genotype calling with uncertainty from

sequencing data in polyploids and diploids. G3 - Genes Genom. Genet. 9: 663-673.

Clevenger J, Chavarro C, Pearl SA, Ozias-Akins P, Jackson SA (2015) Single Nucleotide

Polymorphism identification in polyploids: a review, example, and recommendations.

Mol. Plant 8: 831-846.

De Gabriel Hernando M, Fernandéz-Gil J, Roa I, Juan J, Ortega F, De la Calzada F, Revilla E

(2021) Warming threatens habitat suitability and breeding occupancy of rear-edge bird

specialists. Ecography 44:1191-1204.

Dixo M, Metzger JP, Morgante JS, Zamudio KR (2009) Habitat fragmentation reduces genetic

diversity and connectivity among toad populations in the Brazilian Atlantic Coastal

Forest. Biol. Conserv. 142: 1560-1569.

Dufresne F, Stift M, Vergilino R, Mable BK (2014) Recent progress and challenges in population

genetics of polyploid organisms: an overview of current state-of-the-art molecular and

statistical tools. Mol. Ecol. 23: 40-69.

Fick SE, Hijmans RJ (2017) New 1 km spatial resolution climate surfaces for global land areas.

Int. J. Climatol. 37: 4302-4315.

Fitzpatrick, SW, BRadburd GS, Kremer CT, Salerno PE, Angeloni LM, Funk WC (2020) Genomic

and fitness consequences of genetic rescue in wild populations. Curr. Biol. 30: 517-522.

Frankham R (2015) Genetic rescue of small inbred populations: meta-analysis reveals large

and consistent benefits of gene flow. Mol. Ecol. 24, 2610-2618.

Frichot E, Mathieu F, Trouillon T, Bouchard G, François O (2014) Fast and efficient estimation

of individual ancestry coefficients. Genetics 4: 973-983.

Frichot E, François O (2015) LEA: An R package for landscape and ecological association

studies. Methods Ecol. Evol. 6: 925-929.

Godefroid S, Piazza C, Graziano R, Buord S, Stevens A-D, Aguraiuja R, et al. (2011) How

successful are plant species reintroductions? Biol. Conserv. 144: 672-682.

Gutiérrez EE, Heming NM, Penido G, Dalponte JC, Lacerda ACR, Moratelli R et al. (2019)

Climate change and its potential impact on the conservation of the Hoary Fox,

Lycalopex vetulus (Mammalia: Canidae). Mammalian Biol. 98: 91-101.

Ren H, Jian S, Liu H, Zhang Q, Lu F (2014) Advances in the reintroduction of rare and

endangered wild plant species. Sci China Life Sci 57, 603–609.

Heming NM, Dambros C, Gutiérrez E (2018) ENMwizard: advanced techniques for Ecological

Niche Modeling made easy. https://github.com/HemingNM/ENMwizard

Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics 24: 1403-1405.

Kamvar ZN, Tabima JF, Grünwald NJ (2014) Poppr: an R package for genetic analysis of

populations with clonal, partially clonal, and/or sexual reproduction. PeerJ 2: e281.

Karbstein K, Tomasello S, Hodač L, Dunkel FG, Daubert M, Hörandl E (2020) Phylogenomics

supported by geometric morphometrics reveals delimitation of sexual species within

the polyploid apomictic *Ranunculus auricomus* complex (Ranunculaceae). Taxon 69:

1191-1220.

Karger DN, Conrad O, Böher J, Kawohl T, Kreft H, Soria-Auza RW et al. (2017). Climatologies at

high resolution for the earth's land surface areas. Scientific Data 4: 170122.

Kidane YO, Steinbauer MJ, Beierkuhnlein C (2019) Dead end for endemic plant species? A

biodiversity hotspot under pressure. Glob. Ecol. Conserv. 19: e00670.

Kuhn M (2019) Caret: classification and regression training. https://CRAN. R-

project.org/package=caret.

Liddell E, Sunnucks P, Cook CN (2021) To mix or not to mix gene pools for threatened species

management? Few studies use genetic data to examine the risks of both actions, but

failing to do so leads disproportionately to recommendations for separate

management. Biol. Conserv. 256: 109072.

Lespinas F, Ludwig W, Heussner S (2014) Hydrological and climatic uncertainties associated

with modeling the impact of climate change on water resources of small

Mediterranean coastal rivers, Journal of Hydrology 511: 403-422.

López-Pujol J, Orellana M, R, Bosch M, Simon J, Blanché C. (2007) Low genetic diversity and

allozymic evidence for autopolyploidy in the tetraploid Pyrenean endemic   larkspur

Delphinium montanum (Ranunculaceae). Biol. J. Linn. Soc. 155: 211-222.

Maschinski J, Albretcht MA (2017) Center for plant conservation's best practice guidelines for

the reintroduction of rare plants. Plant Divers. 39: 390-395.

McCormack JE, Huang H, Knowles LL (2009) Sky Islands. In: Gillespie RG, Clague D (eds)

Encyclopedia of Islands, University of California Press: Berkeley, CA Pp 839-843.

Meirmans PG, Van Tienderen PH (2013) The effects of inheritance in tetraploids on genetic

diversity and population divergence. Heredity 110: 131-137.

Meirmans PG, Liu S, van Tienderen PH (2018) The analysis of polyploid genetic data. J. Hered.

109: 283-296.

Meirmans PG (2020) Genodive version 3.0: Easy-to-use software for the analysis of genetic

data of diploids and polyploids. Mol. Ecol. Res. 20: 1126-1131.

Muscarella R, Galante PJ, Soley-Guardia M, Boria RA, Kass JM, Uriarte M. et al. (2014)

ENMeval: an R package for conducting spatially independent evaluations and

estimating optimal model complexity for Maxent ecological niche models. Methods

Ecol. Evol. 5: 1198-1205.

Nei M (1987) *Molecular Evolutionary Genetics*. Columbia University Press: New York: 1-514.

Pandit MK, Pocock MJO, Kunin WE (2011) Ploidy influences rarity and invasiveness in plants.

J. Ecol. 99: 1108-115.

Paris JR, Stevens JR, Catchen J (2017) Lost in parameter space: a road map for stacks. *Methods*

Ecol. Evol. 8: 1360-1373.

Peery MZ, Kirby R, Reid BN, Stoelting R, Doucet-Bëer E, Robinson S, et al. (2012) Reliability of

genetic bottleneck tests for detecting recent population declines. Mol. Ecol. 21: 3403-

3418.

Phillips SJ, Anderson, RP, Schapire RE (2006) Maximum entropy modelling of species

geographic distributions. Ecological Modelling 190: 231-259.

Philipps SJ, Anderson RP, Dudík M, Shapire RE, Blair ME (2017) Opening the black box: an open-

source release of Maxent. Ecography 40: 887-893.

Ralls K, Ballou JD, Dudash MR, Eldridge MDB, Fenster CB, Lacy RC et al., (2018) Call for a

paradigm shift in the genetic management of fragmented populations. Conserv. Lett.

11: e12412.

Ren H, Jian SG, Liu HW, Zhang QM, Lu HF (2014) Advances in the reintroduction of rare and

endangered wild plant species. Sci. China Life Sci.: 57, 603-609.

Rochette NC, Catchen JM (2017) Deriving genotypes from RAD-seq short read data using

Stacks. Nat. Protoc. 12: 2640-2659.

Rousset F (1997) Genetic differentiation and estimation of gene flow from *F*-statistics under

Isolation by Distance. Genetics 145: 1219-1228.

Simon J, Bosch M, Molero J, Blanché C (2001) Conservation biology of the Pyrenean larkspur

(*Delphinium montanum*): a case of conflict of plant versus animal conservation? Biol.

Conserv. 98: 305-314.

UICN France, FCBN, AFB, MNHN (2018) La liste rouge des espèces menacées en France –

Chapitre Flore vasculaire de France métropolitaine. Paris, France.

Urban MC (2015) Accelerating extinction risk from climate change. Science 348: 571-573.

Van de Peer Y, Mizrachi E, Marchal K (2017) The evolutionary significance of polyploidy. Nature

Rev. Genet. 18: 411-424.

Van de Peer Y, Ahsman T-L, Soltis PS, Soltis DE (2021). Polyploidy: an evolutionary and

ecological force in stressful times. Plant Cell 33: 11-26.

Vinogradov AE (2003) Selfish DNA is maladaptive: evidence from the plant Red List. Trends

Genet 19: 609-614.

Wagner ND, He L, Hörandl E (2021) Phylogenomic relationships and evolution of polyploid

Salix species revealed by RAD sequencing data. Front. Plant Sci. 11: 1077.

Whiteley AR, Fitzpatrick SW, Funk WC, Tallmon DA (2015) Genetic rescue to the rescue. *Trends*

Ecol. Ecol. 30: 42-49.

Záveská E, Maylandt C, Paun O, Bertel C, Božo F, The STEPPE Consortium et al. (2019). Multiple

auto- and alloploidisations marked the Pleistocene history of the widespread Eurasian

steppe plant *Astragalus onobrychis* (Fabaceae). Mol. Phylogenetics Evol. 139: 106572.

Tables

Table 1 Geographic coordinates (in °) and elevation (in m above sea level) of the sampling

localities, sample size (N), observed (H_O) and expected (H_E) levels of heterozygosity, deviation

from panmixia (G_{IS}), total number of alleles (A) and number of private alleles (A_p) per locality.

	Locality	Latitude	Longitude	Elevation	N	H_O	$H_E{}^*$	G_{IS}	A	A_p
1	Orri de Baix	42.444	2.119	2115	12	0.117	0.171	**0.343**	17 549	275
2	Nohèdes	42.615	2.263	1758	12	0.101	0.146	**0.336**	19 570	3,704
3	Vallter	42.426	2.264	2141	12	0.095	0.147	**0.378**	16 862	3,483
4	Pedraforca	42.237	1.704	2366	10	0.144	0.244	**0.426**	22 090	1,069
5	Serra Pedregosa	42.277	1.688	2586	12	0.159	0.237	**0.346**	22 456	524
6	Cambre d'Aze	42.451	2.129	2637	12	0.134	0.209	**0.380**	27 075	668
7	Tosa	42.315	1.886	2153	12	0.125	0.177	**0.315**	21 782	3,857
8	Bastanist	42.289	1.688	1979	12	0.165	0.242	**0.332**	26 822	310
9	Torreta de Cadí	42.283	1,574	2358	12	0.122	0.191	**0.376**	17 876	4,537
					106	**0.129**	**0.196**	**0.361**		

685   Geographic coordinates are in WGS84.

686   Population genetic statistics were computed based on 5 095 SNPs. Statistically significant values are indicated

687   in bold ($p < 0.05$).

688   *H_E: unbiased expected heterozygosity measured following Nei (1987) in Genodive.

Table 2 Current and future *Delphinium montanum* suitable area (in km^2) as modelled and predicted from Ecological Niche Modelling under different model selection criteria (LowAIC and AUC) and thresholding methods (10th percentile of training presence, x10ptp and Maximum training sensitivity plus specificity, mtss. Future predictions are reported for several time periods (2011-2040, 2041-2070 and 2071-2100) and various Representative Concentration Pathways (RCP 2.6, 7.0 and 8.5).

Model selection criterion	Thresholds methods	Current	2011-2040			2041-2070			2071-2100		
			RCP 2.6	RCP 7.0	RCP 8.5	RCP 2.6	RCP 7.0	RCP 8.5	RCP 2.6	RCP 7.0	RCP 8.5
LowAIC	x10ptp mtss	2670	523	492	526	137	17	24	272	0	0
AUC	x10ptp mtss	2365	459	455	446	161	25	43	287	0	0

Figures

Figure 1 Map representing the sampling localities (pink dots) within the extant geographic distribution of *Delphinium montanum*.

Figure 2 A) Principal Component Analysis (PCA) displaying the two first axes (PC1 and PC2) representing 14.85% and 9.21% of the total genetic variance. PCA was computed based on the 106 individuals of *D. montanum* genotyped at 5 095 SNPs whose colours represent sampling localities. **B)** Values of the cross-entropy criterion for a number of clusters ranging from *K* = 1 to 10 (10 sNMF runs each). The optimal number of *K* was found to be 6. **C)** Barplot of ancestry coefficients obtained from sNMF for 106 individuals for *K* = 2 and *K* = 6, based on 5 095 SNPs.

Figure 3 Pattern of Isolation By Distance (IBD) across the data set. Linearized G_{ST}s values (*i.e.* ln(G_{ST}/(1- G_{ST})) are plotted against ln-geographical distances.

Figure 4 Geographic areas holding suitable climatic conditions for *Delphinium montanum* with

projection under present climatic conditions (average for 1981-2010) and consensus

projection (from 5 Global Climate Models (CGMs): GFDL-ESM4, UKESM1-0-LL, MPI-ESM1-2-

HR, IPSL-CM6A-LR and MRI-ESM2-0), for various plausible climatic scenarios for years 2035

(average for 2011-2040), 2055 (average for 2041-2070) and three Representative

Concentration Pathways (RCPs): RCP 26, RCP 70 and RCP 8.5. The time period 2071-2100 is

not represented since the suitable area would have completely disappeared for RCPs 7.0 and

8.5. The Barplot displays current and future area values under various plausible climatic

scenarios. These values correspond to "LowAIC" model ("AUC" model being very similar).