abmR: an R package for agent-based model analysis of large-scale movements across taxa

Benjamin Gochanour[1,2,*], Javier Fernández-López[3], Andrea Contina[4, 5]

[1] Corix Plains Institute, University of Oklahoma, Five Partners Place, 201 Stephenson Parkway, Norman, OK 73019, USA

[2] Oklahoma Biological Survey, University of Oklahoma, 111 East Chesapeake Street, SC Building 134, Norman, OK 73019, USA.

[3] University of Massachusetts Boston, Department of Biology, Integrated Sciences Complex, 100 Morrissey Blvd., Boston, MA 02125

[4] University of Colorado Denver, Department of Integrative Biology, Science Building 2074, Denver, CO 80217, USA

[5] University of Oklahoma, Department of Microbiology and Plant Biology, Center for Earth Observation and Modeling, 101 David L. Boren Blvd, Norman, OK 73019, USA

*Corresponding author

*E-mail: Gochanour.Benjamin@mayo.edu

Benjamin Gochanour ORCID: 0000-0002-0950-1739

Andrea Contina ORCID: 0000-0002-0484-671

Abstract

1. Agent-based modeling (ABM) shows promise for animal movement studies. However, a robust, open-source, and spatially explicit ABM coding platform is currently lacking.

2. We present abmR, an R package for conducting continental-scale ABM simulations across animal taxa. The package features two movement functions, each of which relies on the Ornstein-Uhlenbeck (OU) process.

3. The theoretical background for abmR is discussed and the main functionalities are illustrated using example populations.

4. Potential future additions to this open-source package may include the ability to specify multiple environmental variables or to model interactions between agents. Additionally, updates may offer opportunities for disease ecology and integration with other R movement modeling packages.

Keywords:

Animal migration; R programming; simulations; ecology; open-source

Abbreviated Title: agent-based models in R

1. INTRODUCTION

Animal movement is a complex behavioral trait that affects the survival of populations and species across taxa (Berg, 1983; Dingle, 2014). Long- and short-distance movements can follow predictable environmental constraints, allowing populations to take advantage of seasonal food resources (e.g., migration), or more opportunistic, such as in the case of dispersal behaviors aimed at avoiding predators or finding potential mates (Giuggioli & Bartumeus, 2010). Thus, wild animals make decisions often based on environmental cues that lead to movement patterns characteristic of different populations across the landscape (Nathan et al., 2008; Dodge et al., 2014). However, obtaining a comprehensive understanding of large-scale animal movement behavior and population occurrence under climate change scenarios or habitat loss has proven to be a challenge (Araujo & Guisan, 2006). Moreover, while the research toolbox in movement ecology studies has seen a considerable expansion over the last two decades due to technological advancements of the tracking devices and molecular markers (Cushman and Lewis, 2010; Williams et al., 2020), the limitation of scaling up individual data to population-level inferences is still a substantial obstacle (Hawkes, 2009; but see Holdo & Roach, 2013). A promising research approach that may overcome the limitations of wildlife movement studies hindered by small sample sizes is represented by computer simulations within an Agent-based Modeling (ABM) framework (Tang & Bennett, 2010; Bridge et al., 2017).

The core principle of ABM is to simulate a set of entities, called agents, which are defined by intrinsic properties as well as behavioral rules governing their interactions with the environment (Grimm & Railsback, 2013). That is, agents are described by their inherent attributes while dynamically interacting with external conditions such as the co-occurrence of other agents and/or changing features of their environmental setting. Thus, ABM is used in many areas, including biology, disease risk, social sciences, and economics (Polhill et al.,

2008; Grimm & Railsback, 2013; Kilmek et al., 2015; Willem et al., 2017) with the unifying

goal of investigating and predicting the dynamics of complex systems (Grimm et al., 2005).

In particular, wildlife studies have adopted the ABM approach to simulate population growth,

reproduction, mortality rate, energy budget, and migration ecology, just to cite a few (Brown

& Robinson, 2006; Lustig et al., 2019; Aurbach et al., 2020; Goldstein et al., 2021).

However, we currently lack a robust and spatially explicit ABM coding platform for the

implementation of large-scale animal movement investigations (but see Thiele et al. (2012)

and Chubaty and McIntire (2021)). Here we present a novel ABM framework in R

programming language (R Core Team, 2020) for applications in animal behavior and

movement ecology.

2. PACKAGE OVERVIEW

abmR allows for both computation and visualization of agent movement trajectories through

a set of behavioral rules based on environmental parameters. The two movement functions,

`moveSIM` and `energySIM`, provide the central functionality of the package, allowing the

user to run simulations using an Ornstein-Uhlenbeck movement model (Uhlenbeck &

Ornstein, 1930; hereafter OU). Additional functions provide a suite of visualization and data

summarization tools intended to reduce the effort needed to go from results to presentation-

ready figures and tables (Table 1). The package is currently available on the Comprehensive

R Archive Network (CRAN).

Function	Usage
moveSIM	Runs agent-based model movement simulations based on environmental data
moveVIZ	Creates a plot or table of moveSIM() results
energySIM	Runs agent-based model movement and energy budget simulations based on environmental data
energyVIZ	Creates a plot or table of energySIM() results
tidy_results	Prints results from moveSIM() or energySIM() in an easier-to-read table
get_ex_data	Downloads data that is used in examples in vignette and documentation
as.species	Creates object of class 'species' for input into moveSIM() or energySIM()

Table 1. Functions contained in the abmR package (v. 1.0.6). For more complete function

descriptions, consult the abmR manual.

The abmR package is built to facilitate customization of movement model parameters within

the R environment (Fig. 1). Parameters affecting agent behavior can be obtained from real-

world observations (e.g., GPS data) or approximated from the literature and then manually

entered as arguments into abmR functions. Thus, if the user knows beforehand the behavior

and the ecological constraints of the agents, such as land-cover preferences (e.g., vegetation

composition and structure) or movement direction and average distance traveled per day,

they may transfer this knowledge into abmR to study changes in energy consumption,

mortality rate or alternative routes across different environmental conditions (e.g., raster

layers). Alternatively, for simulations of 'synthetic species' designed to compare relative

mortality rates across habitats, for example, the user may develop an analytical framework

within abmR without exploring the ecology of a particular organism. In this case, the

movement parameters may not be realistic, but selected with the unique goal of comparing a

set of scenarios (e.g., movements in predicted future land-use changes) while studying the

emerging properties of the system (Yin et al. 2022). Finally, abmR functionalities can be

integrated with other approaches, such as a step-selection function (Thurfjell et al. 2014)

and/or Bayesian statistics to estimate movement parameters and explore migration patterns or

habitat occupancy of a particular set of agents (Joo et al. 2020; Cullen et al. 2022).

Workflow 1	Workflow 2	Arguments
Study system (*real species*)	**Synthetic system** (*ideal species*)	replicates, origin x (longitude), and origin y (latitude)
Parameters obtained from the literature	**Virtual parameters for comparative analyses**	days, optimum values, destination (latitude and longitude), motivation, search radius, direction, sigma
Select Environment		environmental raster, single or multiple raster layers
Run abmR		initial energy, energy adjustment (gain/penalty), number of failures and failure threshold
RESULTS		movement visualization, energy visualization, summary results, data output

Figure 1. abmR workflow from data input to result exploration. The black box and black arrows

(Workflow 1, left) show the sequence of the analytical steps, starting with the identification of the

study system (species or population), followed by the acquisition of the movement parameters from

previous observations or the literature. After selecting the environmental conditions (e.g., raster

layers), users can run a suite of abmR functions to study movements, changes in energy consumption

or mortality rate. The blue box and blue arrows (Workflow 2, right) show an alternative analytical

workflow in which users can create synthetic species for comparative analyses across predicted

ecological and environmental scenarios (e.g., loss of migratory behavior and habitat fragmentation).

The panel to the right of both workflows shows a list of the relative arguments that can be entered in

abmR across each step of the analysis.

Both movement functions used by abmR rely on the same OU approach. The OU process is

described by the following stochastic differential equation:

$$dX_t = k(\theta - X_t)\, dt + \alpha dW_t$$

where W_t is a standard Brownian motion on $t \in [0, \infty)$, k is a positive constant that controls

the rate of reversion to the long-term mean, θ, of the OU process (see Blomberg et al 2020),

and $\alpha > 0$ is a constant controlling the volatility of the Brownian motion. In abmR, given the

current agent location (x_t, y_t), agent location at the subsequent timestep (x_{t+1}, y_{t+1}) is modeled

according to the following equations:

130
$$x_{t+1} = x_t + \sigma * Z_x + \phi_x * (\theta_x - x_t) \qquad (1)$$

131
$$y_{t+1} = y_t + \sigma * Z_y + \phi_y * (\theta_y - y_t) \qquad (2)$$

Here, σ is a user-specified multiplier on the random terms Z_x and Z_y, two numbers drawn

from the Normal $(0,1)$ distribution. In addition, ϕ_x and ϕ_y are movement motivation or

attraction strength for the OU process in the longitude and latitude coordinates, respectively,

while θ_x and θ_y are optimal x (longitude) and y (latitude) coordinates, respectively. It is

assumed that the origin point (x_1, y_1) is known. The OU model given in (1) and (2) performs

similarly to a spring-coil (Fig. 2). Greater distance from optimal coordinates θ_x and θ_y acts

like a compressed spring to propel distant agents towards θ_x and θ_y. On the other hand,

agents closer to θ_x and θ_y will travel a shorter distance on that timestep. However, the length

of movement also depends on ϕ_x and ϕ_y, because these motivations serve as a multiplier on

$(\theta_x - x_t)$ and $(\theta_y - y_t)$, respectively (Eqns. 1 and 2).

Figure 2. The Ornstein-Uhlenbeck (OU) model given in (1) and (2) performs like a spring-coil:

agents further from their target location experience higher attraction (and travel further), while agents

closer to their destination experience lesser attraction (and travel less far).

While the two movement functions are distinct (see below), each follows the same

basic two steps. The first, large-scale searching, is illustrated in Fig. 3 and follows an

elliptical pattern, as opposed to a circular distribution, to account for directional migratory

impulses (e.g., genetically controlled migratory behaviors or seasonal Zugunruhe; Merlin &

Liedvogel, 2019). This step finds the 'optimum' location for each agent (θ_x and

θ_y coordinates from Eqns. (1) and (2)). The coordinates θ_x and θ_y are determined according

to an algorithm that selects the location whose observed environmental value has the least

possible difference from the user-specified optimal raster value. For moveSIM, this user-

specified optimal raster value is supplied directly, while for energySIM it is the average of

the lower and upper bounds of a user-specified optimum range. The optimum value or range

of values specified depends on the modeling scenario and the type of environmental raster

that is used (e.g., vegetation, temperature, etc.).

Agents will move toward the selected optimum location. However, if the attraction strength

(ϕ_x and ϕ_y in Eqns. (1) and (2)) is less than 1, agents will have a 'target' location short of the

optimal location. Thus, agents find an 'optimum' location within the semi-circular search

region and then a 'target' location that lies on the line between the 'current location' and the

'optimum' location. Moreover, agents will move towards this target location with some

variance, which is generated by sampling two numbers from the standard normal distribution

and multiplying by σ, as specified by the user (In Fig. 3, σ is 1). Because the support of the

normal distribution consists of all real numbers, large deviations from the 'target' point are

possible. However, because the normal distribution has low density at the extreme tails,

outcomes are most likely to fall within a certain area of the target, as illustrated in Fig. 3. This

first step corresponds to the OU model of Eqns. (1) and (2).

Figure 3. Illustration of large-scale searching specified by the OU model of Eqns. (1) and (2). Agents

find an 'optimum' location within the semi-circular search region and then a 'target' location that lies

on the line between the 'current location' and the 'optimum' location. If there is a tie between

multiple potential 'optimum' cells, one is randomly selected from the list of tied cells to serve as the

optimum. Random variance is added by independently sampling Z_x and Z_y from a standard normal

distribution. Here, $\sigma = 1$ and $\phi_x = \phi_y < 1$, where σ is the multiplier on the random error and ϕ_x and ϕ_y

are the motivations in the x and y directions, respectively. Bounding box represents the most probable

samples from the $N(0,1)$ distribution.

178   The second step is small-scale searching. Here, agents select the 'best' of the 8

neighboring cells (queen's case or Moore neighborhood) after performing step 1, discussed

above and in Fig 3. Again, 'best' here means the cell with the environmental raster value

closest to the agent's user-defined optimum range. These two steps are then repeated for each

timestep until the agent dies or proceeds through all timesteps. Each timestep will use

different environmental raster layers. Users may choose to supply a raster stack to simulate

changes in environment over time (see for example Section 3.1).

185   For the `moveSIM` function, agent death occurs when agents fail to achieve suitable

environmental raster values for more than a user-specified number of consecutive timesteps.

Here, what constitutes a 'suitable' cell is determined by the optimum value and an allowable

deviation proportion, both also specified by the user. For the `energySIM` function, agent

death occurs when energy reaches zero. For both functions, users may choose to disable agent

mortality. In the following subsections, we present the differences between `moveSIM` and

`energySIM` functions and their underlying algorithms.

2.1. Simulation function: moveSIM

The function `moveSIM` runs an OU movement simulation based on environmental conditions

provided by the user (e.g., raster), optionally including agent mortality and adjusted

motivation according to user-specified parameters. The function operates according to the

following algorithm. Here, terms in *italics* are `moveSIM` function arguments (see Table 2).

Argument	Function	Usage
replicates	M, E	# of agents to model
days	M, E	# of timesteps
modeled_species	M, E	Species object from as.species()
env_rast	M, E	Environmental raster
optimum	M	Optimal environmental value
optimum_lo	E	Lowest optimum environmental value
optimum_hi	E	Highest optimum environmental value
dest_x	M,E	Destination Longitude
dest_y	M,E	Destination Latitude
mot_x	M,E	Motivation (x direction)
mot_y	M,E	Motivation (y direction)
search_radius	M,E	Radius of semi-circular search region (km)
direction	M,E	Movement direction: N, S, E, W, or R (Random)
sigma	M,E	Randomness parameter
mortality	M,E	Incorporate agent mortality? T or F
fail_thresh	M	Deviation from optimum constituting failure
n_failures	M	Allowable # of failures before agent death
init_energy	E	Initial energy
energy_adj	E	Energy gain/penalty vector
single_rast	M, E	Using a single-layer raster? T or F
write_results	M, E	Save results as a .csv? T or F
x	S	Origin longitude
y	S	Origin latitude

Table 2. List of arguments used in `moveSIM` (M), `energySIM` (E), or `as.species` (S). T: True,

F: False, SD: standard deviation. In text, these arguments are presented in *italics*. For a more complete

list of argument descriptions, see the abmR documentation.

The following algorithm applies when the argument *direction* is 'N', 'S', 'E', or 'W'.

For random movement (*direction* = 'R') agents simply select a random point from a circle of

radius *search_radius* for each timestep (Step 2). Here, let *env_rast* (x_{t+1}, y_{t+1}) be the value

of *env_rast* at the point (x_{t+1}, y_{t+1}). The core algorithm shown here assumes that *env_rast*

contains no undefined (N/A) grid cells.

1. Specify (x_1, y_1) using *x* and *y* contained in *modeled_species,* set failures = 0.

2. For day t in 1:(*days*-1)

a. Define search area semicircle (radius = *search_radius)* facing *direction*

and centered at (x_t, y_t)

210   b.   Determine (θ_x, θ_y) as location within the search area with *env_rast*

211   value closest to *optimum.*

212   i.   If (*dest_x*, *dest_y*) in search area, set $(\theta_x, \theta_y) = (dest\_x, dest\_y)$

213   c.   Large scale searching: find $(x_{t+1}, y_{t+1})_0$ according to (1) and (2).

214   d.   Small scale searching: set (x_{t+1}, y_{t+1}) as location within eight

215   neighboring cells (queen's case) of $(x_{t+1}, y_{t+1})_0$ with the value closest

216   to *optimum*.

217   Perform (e)-(f) if *mortality* = True.

218   e.   If observed *env_rast*(x_{t+1}, y_{t+1}) - *optimum* > *optimum*fail_thresh,* set

219   failures = failures + 1. If not, set failures = 0.

220   f.   If failures > *n_failures* agent dies. End loop.

221   3.   Return dataframe with *days* rows and 2 columns movement track data.

222   4.   Repeat (1)-(3) *replicates* times.

223   2.2. Simulation function: energySIM

The function `energySIM` builds on `moveSIM` by allowing for dynamic agent energy levels

that are affected by the quality of environmental values achieved. These initial user-defined

energy levels then serve as a driver of mortality and movement distance per timestep. The

energy thresholds range from 0 to 1 and represent proportion deviations from the optimum.

These thresholds are arbitrarily divided into 10 even increments (0.1, 0.2, etc.), but users can

change the energy gain or penalty for attaining each of the thresholds. It operates according to

the following algorithm. Here, terms in *italics* are `energySIM` function arguments (see

Table 2) or calculated variables (e.g., *optimum, energy*).

The following algorithm applies when the argument *direction* is 'N', 'S', 'E', or 'W'.

For random movement (*direction* = '*R*'*)* agents simply select a random point from a circle of

radius *search_radius* for each timestep (Step 3). Here, let *env_rast* (x_{t+1},y_{t+1}) be the value of

env_rast at the point(x_{t+1}, y_{t+1}). The core algorithm shown here assumes that *env_rast*

contains no undefined (N/A) grid cells.

1. Specify (x_1, y_1) using *x* and *y* contained in *modeled_species*.

2. Compute *optimum* as *(optimum_hi - optimum_lo)/2 and* set *energy =*
    init_energy

3. For day t in 1:(*days*-1)

    a. If *mortality* = True, update *search_radius* as *search_radius =*
        *search_radius * (energy/init_energy)*.

    b. Define search area semicircle(radius = *search_radius)* facing *direction*
        and centered at (x_t, y_t).

    c. Determine (θ_x, θ_y) as location within the search area with *env_rast*
        cell value closest to *optimum.*

        i.   If (*dest_x, dest_x*) in search area, set $(\theta_x, \theta_y) = (dest\_x, dest\_y)$

    d. Large scale searching: find$(x_{t+1}, y_{t+1})_0$ according to (1) and (2).

    e. Small scale searching: set (x_{t+1}, y_{t+1}) as location within eight
        neighboring cells (queen's case) of $(x_{t+1}, y_{t+1})_0$ with the cell value
        closest to *optimum*.

    f. Update energy level according to deviation of env_rast (x_{t+1}, y_{t+1})
        from optimum.

    g. If *mortality* = True and *energy* = *0*, the agent dies. End loop.

4. Return dataframe with *days* rows and 2 columns movement track data.

5. Repeat (1)-(4) *replicates* times.

3. EXAMPLE APPLICATIONS

abmR can be used to construct ABM simulations for any desired agent across the globe. In

the following examples, we demonstrate the computation capabilities of the `energySIM`

function, although a similar workflow also applies for `moveSIM`. In the first example, we

show how `energySIM` can be used to compare movements and differential energy

allocations of two synthetic populations of 250 agents each (Box 1). In the second example,

we replicate the movement pattern of the Painted Bunting (*Passerina ciris*), a well-studied

migratory songbird occurring in North America and Mexico.

3.1. Comparisons between populations

In this example, both synthetic populations are characterized by matching number of

replicates and movement timesteps (*days*), equal σ, and the same environmental data

provided by a Normalized Difference Vegetation Index (NDVI) raster stack containing 14

days of data between September 01-14, 2019 (Vermote, 2019). Both populations had an

unspecified destination (indicated with '999' in the arguments *dest_x* and *dest_y*) and were

constrained to move on land. However, Population 1 (P1) agents started their movements

from a different point (105.7° W; 48.2° N) situated about 2,800 km from the origin of

Population 2 (P2) agents (142.7° W; 63.2° N). Additionally, P1 agents had a smaller search

radius (150 km) but higher motivation than P2 agents (P1 motivation = 0.95). P1 agents also

had different optimum ranges (P1 0.2-0.5; P2 0.6-0.8), and different initial energy units (P1

100; P2 70). These differences in simulation parameterization result in clearly dissimilar

movement tracks (Fig. 4).

Population 1	Population 2

``
am.pop.1 = as.species(x = -105.7, y = 48.2)
sim.move <- energySIM(
  replicates = 250,
  days = 14,
  env_rast = as.raster.stack.ndvi.sep,
  search_radius = 150,
  sigma = 0.1,
  dest_x = 999,
  dest_y = 999,
  mot_x = 0.95,
  mot_y = 0.95,
  modeled_species = am.pop.1,
  optimum_lo = 0.2,
  optimum_hi = 0.5,
  init_energy = 100,
  direction = "S",
  mortality = F,
  Energy_adj = c(20,10,8,5,2,0,-2,-5,-8,-10,-20),
  write_results = T,
  single_rast = F)
``
	``
	am.pop.2 = as.species(x = -142.7, y = 63.2)
	simtwo.move <- energySIM(
	  replicates = 250,
	  days = 14,
	  env_rast = as.raster.stack.ndvi.sep,
	  search_radius = 800,
	  sigma = 0.1,
	  dest_x = 999,
	  dest_y = 999,
	  mot_x = 0.8,
	  mot_y = 0.8,
	  modeled_species = am.pop.2,
	  optimum_lo = 0.6,
	  optimum_hi = 0.8,
	  init_energy = 70,
	  direction = "S",
	  mortality = F,
	  energy_adj = c(20,10,8,5,2,0,-2,-5,-8,-10,-20),
	  write_results = T,
	  single_rast = F)
	``

280   **Box 1**. R code used for performing the simulations presented in Fig 4. First, `as.species` is called

281   to initialize two populations with different origin locations. Then, `energySIM` is called to perform a

282   movement simulation for each population; parameters that differ between the two simulations are

283   printed in red, functions in blue, and objects in bold. For argument descriptions, see Table 2 and the

284   package manual.

Figure 4. Movement tracks reveal that Population 1 tended to travel through the central United States, while Population 2 traveled mostly throughout western Canada, United States, and Mexico.

Overall, Population 1 traveled more distance and exhibited more consistent paths near the origin than

did Population 2. The movement tracks are natively produced by abmR. Inset world map provided for

geographic reference.

While we can compare the movement tracks visually, Table 3 provides a numerical

description of the results. In this simulation, P1 traveled a much smaller average distance

(154.6 km) than did P2 (625.5 km). However, P1 traveled more days on average (7.4 days)

before stopping than P2 (3.8 days). Additionally, P2 had higher energy consumption than P1;

its average remaining energy across all timesteps was 61.2 units compared to 99.7 units for

P1. There are several possible reasons for this observed pattern. First, P2 began with a

smaller initial energy (70 units) than P1 (100 units). Additionally, P2 had higher optimum

NDVI values (0.6 - 0.8), which might have been less abundant and generally more difficult to

reach than those of P1 (0.2 - 0.5). Finally, because they began in different places, P1 and P2

agents encountered different raster cells along their journey.

	variable	mean	sd	median	min	max	range
Population 1	day	7.4	4	7	1	14	13
	longitude	-105.5	4.1	-105.2	-121.1	-93.3	27.8
	latitude	41.6	3.4	41.9	30.2	47.5	17.3
	energy	99.7	1.3	100	80	100	20
	delta energy	-0.04	1.5	0	-20	20	40
	distance	154.6	70	154.4	5.5	316.7	311.1
Population 2	day	3.8	3.08	3	1	14	13
	longitude	-126	17.3	-127.1	-166	-80.3	85.7
	latitude	50.2	10.5	52.2	14.4	63.1	48.7
	energy	61.2	17.2	60	0	100	100
	delta energy	-2.3	9.3	-5	-20	20	40
	distance	625.5	294	649.7	5.5	1445	1439.4

302   **Table 3.** A numerical comparison of Populations 1 and 2, created by using values across all timesteps

303   for all agents. 'Day' summarizes the timestep variable of the movement tracks. 'Longitude' and

304   'latitude' summarize the geographical position of agents, while 'energy' summarizes agents'

remaining energy. 'Delta energy' corresponds to the change (gain or loss) of energy between each

timestep, while 'distance' refers to the distance traveled between each timestep. This table was

produced outside of abmR using raw movement data returned by the package.

Fig. 5 visually compares P1 and P2 movement outputs based on longitude and

latitude. This is not a native abmR figure, but rather is produced using the raw data that abmR

generates to show the flexible use of the package. In this figure, P1 movements tended to be

to the east and south of P2. However, P2 trajectory shows a much wider distribution, with

density points extending to the lower values of latitude.

Figure 5. Graphical comparisons of Population 1 and Population 2 movements. Panels A and B show

density plots used to individually compare longitude (Panel A) and latitude (Panel B) coordinates

attained by agents from each population. Panel C compares the distance traveled between each

timestep, while Panel D shows geographical position for all agents in each population across all

timesteps.

Finally, Fig. 6 provides a density surface plot for P2 describing agent energy gains

(blue) and losses (red) across the landscape. This surface was created using the inverse

distance weighted interpolation (IDW) function from the R package 'gstat' (Pebesma, 2004).

IDW interpolates grid cell values across a surface using a linear combination of observed

(sample) points. When interpolating a cell value, the value of the sample points closer to that

cell carry a higher weight, while sample points further from that cell carry smaller weight.

IDW is discussed in more detail in Wong (2017). The results from Fig. 6 match well with

what we observe in Fig. 4. Movement tracks for P2 tend to follow the blue (energy gain)

regions.

Figure 6. Energy gradient plot of Population 2 by timestep. Areas in red reflect energy loss (less

suitable environmental values) while areas in blue reflect energy gain (better environmental values).

This plot is produced directly by energyVIZ. Inset world map added for geographic reference.

3.2. Migration of the Painted Bunting

The Painted Bunting is a migratory songbird that has been intensively studied over the past two decades due to its steeply declining population trend in the United States (U.S.) and its complex molt-migratory behavior (Thompson 1991). This species has been the focus of pioneering light-level geolocator tags that revealed westward post-breeding movements from the southern U.S. throughout Mexico (Contina et al. 2013), a system for genetic analysis of migratory population connectivity (Battey et al. 2018; Contina et al. 2019) and for candidate genes studies related to the migratory behavior (Contina et al. 2018). Moreover, Bridge et al. (2016) modeled the post-breeding movements of a Painted Bunting population departing Oklahoma (U.S.) through an ABM approach and investigated the association between large-scale vegetation productivity changes and the molt-migratory behavior. Therefore, this species provides a well-studied migratory system with ample behavioral and population ecology knowledge against which abmR movement predictions can be tested and compared.

We built a basic ABM movement simulation for the Painted Bunting in abmR by using the same key parameters adopted by Bridge et al. (2016) and predicted that a westward migratory movement would emerge throughout the southern U.S. and the coastal regions of Mexico in late summer. Bridge et al. (2016) found that Painted Bunting agents beginning their migratory journey in Oklahoma in late August and September tend to avoid a direct southern migration and show a clear pattern towards southwestern movements targeting high primary productivity areas in northern Mexico and Sinaloa (northwestern Mexico). In our simulation, we calibrated our model by following the movement parameters adopted by Bridge et al. (2016). We used the identical model start location (-98.8° W; 34.8° N), a similar subset of 14 vegetation index raster files (NDVI; Vermote, 2019) representing primary productivity condition during the first two weeks of September 2011, and no predefined destination coordinates (indicated with '999' in the arguments *dest_x* and *dest_y*). For the full

set of model parameters, see suppl. material S2. While we note that Bridge et al. (2016) used

enhanced vegetation index (EVI) as opposed to NDVI, a multiyear timeframe (2010-2013),

and a complex series of model parametrization implemented in ArcGIS, our simulation offers

a basic but useful illustration of the predictive capabilities of abmR.

The outcome of 12 abmR Painted Bunting simulations revealed a movement pattern

consistent with our predictions based on the results presented by Bridge et al. (2016). Most

agents in our simulation (9 out of 12) showed a southwestern movement towards Sinaloa

(Mexico) in mid-September (Fig. 7), where migratory agents utilize a bloom in primary

productivity due to the Monsoonal precipitations (Rohwer et al. 2005). Three agents showed

a southeastern movement, following vegetation changes along southeastern U.S. and the Gulf

of Mexico. This result is also in line with sporadic but nonetheless documented variations in

migratory strategies in the Painted Bunting where some individuals that breed in south central

U.S. (e.g., western Oklahoma and Arkansas) show southeastern movements in late summer

(Contina et al. 2013).

Figure 7. Outcome of 12 abmR simulations showing a frequent southwestern migration from the breeding ground in Oklahoma (U.S.) towards wintering grounds in Sinaloa (Mexico) and southern Mexico. The NDVI map in the background is an average of the raster stack object used in abmR which contained 14 raster layers ranging from September 1 to September 14, 2011.

4. CONCLUSIONS AND FUTURE WORK

abmR provides a novel and efficient programming platform for simulating large-scale movements of species across taxa. We ran most of the initial test simulations on a local machine equipped with an Intel® Core™ i7-5500U CPU – 2.40GHz and 8 GB of RAM and obtained results for 100-1000 agents within minutes. The novelty of the software includes the capability of concurrently modeling agent movement trajectories and energy budget. This feature enables a broader exploration of the ecological constraints that shape animal dispersal and/or migration. Moreover, abmR built-in arguments, such as *fail_thresh*, *n_failures*, and *energy_adj*, provide additional flexibility when evaluating mortality scenarios that depend on baseline environmental conditions and energy requirement during prolonged movement bouts (see Table 2 for a full list of arguments affecting mortality).

Over the last decades, spatially explicit simulations, pattern-oriented modelling, approximate Bayesian computing, and agent-based models have become more popular in ecological and evolutionary studies (Railsback et al., 2006; DeAngelis & Grimm, 2014; van der Vaart et al. 2015; Gallagher et al. 2021). Analytical platforms, such as InSTREAM, a simulation model approach designed to understand how stream and river salmonid populations respond to habitat alteration (Railsback et al., 2009), or ALMaSS, a predictive modeling tool for answering environmental policy questions regarding the effect of changing landscape structure on threatened animal species (Topping et al., 2003), allow investigation of specific ecological systems using ABM. On the other hand, many programming languages such as Netlogo, R, or Python are widely used to develop custom and more flexible models that can be adapted to address complex ecological or evolutionary research scenarios (Lustig et al., 2019; Chubaty & McIntire, 2021). However, the use of a programming language to develop a flexible ABM from scratch has two important drawbacks. First, it requires

advanced programming skills. Second, its reproducibility can be compromised by the

idiosyncrasies of the simulation algorithm written by the user. These idiosyncrasies,

especially if not well documented, can make it difficult or even impossible for other

researchers to replicate findings or adapt code to suit their modeling scenarios. abmR

provides a novel framework to perform complex movement simulations through standardized

functions and arguments that facilitate model annotation and reproducibility while providing

publication-ready visualizations at the end of each run.

While we developed and tested abmR as a movement and energy budget simulation

tool, its core software functionalities can be adapted to explore other processes such as

disease outbreak scenarios (Dougherty et al., 2018). As an example, pathogen vector

movement can be easily simulated within abmR, allowing the study of areas of confluence

where disease transmission is more probable (Manore et al., 2015). Moreover, potential

future updates will include the ability to specify multiple raster stacks of different movement

predictors and interactions between agents. In abmR, each simulation output can be used as

the input for the next movement model. However, the option of computing agent interactions

affecting movement patterns within the same simulation run is currently missing. This is a

clear area of further package development. Additionally, other code expansions might be

useful to study plant seed dispersal, density-dependent scenarios, and altitudinal movements.

Acknowledgements

We thank three anonymous reviewers who provided invaluable feedback on an earlier draft.

abmR relies heavily on a large set of R packages and we thank the R community for

providing support and open-source code. We thank Jeff Kelly, Xiangming Xiao, and the

Center for Earth Observation and Modeling at the University of Oklahoma for supporting the

research effort.

Author Contributions

Author Contributions

B.G. led the development of the package and wrote the manuscript; J.F.L. contributed to the

package, writing, and critical review of the manuscript; A.C. conceived the manuscript, led

its writing, contributed to the package, and performed package testing.

Conflict of Interest Statement

The authors declare that there are no financial or non-financial conflicts of interest.

Data Availability

abmR is available on CRAN and the simulation results for Populations 1 and 2 are available

for download from Zenodo. The environmental raster data set used in the examples is

available at https://www.ncei.noaa.gov.

Funding

References

Alerstam, T. (1993). *Bird migration*. Cambridge University Press.

Araujo, M. B. & Guisan, A. (2006). Five (or so) challenges for species distribution modelling. J. Biogeogr. 33: 1677–1688.

Aurbach, A., Schmid, B., Liechti, F., Chokani, N., & Abhari, R. (2020). Simulation of broad front bird migration across Western Europe. Ecological Modelling, 415, 108879.

Battey, C. J., Linck, E. B., Epperly, K. L., French, C., Slager, D. L., Sykes Jr, P. W., & Klicka, J. (2018). A migratory divide in the Painted Bunting (*Passerina ciris*). *The American Naturalist*, *191*(2), 259-268.

Berg, H. C. (1983). Random Walks in Biology. Princeton Univ. Press, Princeton, NJ.

Blomberg, S.P., Rathnayake, S.I. and Moreau, C.M., (2020). Beyond Brownian motion and the Ornstein-Uhlenbeck process: stochastic diffusion models for the evolution of quantitative characters. The American Naturalist, 195(2), pp.145-165.

Bridge, E.S., Ross, J.D., Contina, A.J., & Kelly, J.F., (2017). Using Agent-Based Models to Scale from Individuals to Populations. In Aeroecology (pp. 259-275). Springer, Cham.

Brown, D. G., & Robinson, D. T. (2006). Effects of heterogeneity in residential preferences on an agent-based model of urban sprawl. Ecology and society, 11(1).

Contina, A., Bridge, E. S., Seavy, N. E., Duckles, J. M., & Kelly, J. F. (2013). Using geologgers to investigate bimodal isotope patterns in Painted Buntings (*Passerina ciris*). *The Auk*, *130*(2), 265-272.

Contina, A., Bridge, E. S., & Kelly, J. F. (2016). Exploring novel candidate genes from the Mouse Genome Informatics database: Potential implications for avian migration research. *Integrative Zoology*, *11*(4), 240-249.

Contina, A., Alcantara, J. L., Bridge, E. S., Ross, J. D., Oakley, W. F., Kelly, J. F., & Ruegg, K. C. (2019). Genetic structure of the Painted Bunting and its implications for conservation of migratory populations. *Ibis*, *161*(2), 372-386.

Chubaty A. M., & McIntire E.J.B. (2021). SpaDES: Develop and Run Spatially Explicit

Discrete Event Simulation Models. R package version 2.0.7. https://CRAN.R-

project.org/package=SpaDES

Cullen, J.A., Poli, C.L., Fletcher Jr, R.J. and Valle, D. (2022). Identifying latent behavioural

states in animal movement with M4, a nonparametric Bayesian method. Methods in Ecology

and Evolution, 13(2), pp.432-446.

Cushman, S. A., & Lewis, J. S. (2010). Movement behavior explains genetic differentiation

in American black bears. Landscape ecology, 25(10), 1613-1625.

DeAngelis, D. L., & Grimm, V. (2014). Individual-based models in ecology after four

decades. F1000prime reports, 6.

Dingle, H. (2014). Migration: the biology of life on the move. Oxford University Press, USA.

Dodge, S., Bohrer, G., Bildstein, K., Davidson, S.C., Weinzierl, R., Bechard, M.J., Barber,

D., Kays, R., Brandes, D., Han, J., & Wikelski, M. (2014). Environmental drivers of

variability in the movement ecology of turkey vultures (*Cathartes aura*) in North and South

America. Philosophical Transactions of the Royal Society B: Biological Sciences, 369(1643),

p.20130195.

Dougherty, E.R., Seidel, D.P., Carlson, C.J., Spiegel, O., & Getz, W.M., 2018. Going through

the motions: incorporating movement analyses into disease research. Ecology letters, 21(4),

pp.588-604.

Gallagher, C. A., Chudzinska, M., Larsen-Gray, A., Pollock, C. J., Sells, S. N., White, P. J.,

and Berger, U. (2021). From theory to practice in pattern-oriented modelling: identifying and

using empirical patterns in predictive models. Biological Reviews, 96(5), 1868-1888.

Giuggioli, L., & Bartumeus, F. (2010). Animal movement, search strategies and behavioural

ecology: a cross-disciplinary way forward. Journal of Animal Ecology, 79(4), 906-909.

Goldstein, E., Erinjery, J. J., Martin, G., Kasturiratne, A., Ediriweera, D. S., de Silva, H. J., ...

& Iwamura, T. (2021). Integrating human behavior and snake ecology with agent-based

models to predict snakebite in high risk landscapes. PLoS neglected tropical diseases, 15(1),

e0009047.

Grimm, V., Revilla, E., Berger, U., Jeltsch, F., Mooij, W. M., Railsback, S. F., ... & DeAngelis, D. L. (2005). Pattern-oriented modeling of agent-based complex systems: lessons from ecology. science, 310(5750), 987-991.

Grimm, V., & Railsback, S. F. (2013). Individual-based modeling and ecology. Princeton university press.

Hawkes, C. (2009). Linking movement behaviour, dispersal and population processes: is individual variation a key?. Journal of Animal Ecology, 78(5), 894-906.

Hedenström, A. (2008). Adaptations to migration in birds: behavioural strategies, morphology and scaling effects. Philosophical Transactions of the Royal Society B: Biological Sciences, 363(1490), pp.287-299.

Holdo, R. M., & Roach, R. R. (2013). Inferring animal population distributions from individual tracking data: theoretical insights and potential pitfalls. Journal of Animal Ecology, 82(1), 175-181.

Joo, R., Boone, M.E., Clay, T.A., Patrick, S.C., Clusella-Trullas, S. and Basille, M. (2020). Navigating through the R packages for movement. Journal of Animal Ecology, 89(1), pp.248-267.

Klimek, P., Poledna, S., Farmer, J. D., & Thurner, S. (2015). To bail-out or to bail-in? Answers from an agent-based model. Journal of Economic Dynamics and Control, 50, 144-154.

Lustig, A., James, A., Anderson, D., & Plank, M. (2019). Pest control at a regional scale: Identifying key criteria using a spatially explicit, agent-based model. Journal of Applied Ecology, 56(7), 1515-1527.

Manore, C.A., Hickmann, K.S., Hyman, J.M., Foppa, I.M., Davis, J.K., Wesson, D.M., & Mores, C.N. (2015). A network-patch methodology for adapting agent-based models for directly transmitted disease to mosquito-borne disease. Journal of biological dynamics, 9(1), pp.52-72.

Merlin, C., and Liedvogel, M. (2019). The genetics and epigenetics of animal migration and orientation: birds, butterflies and beyond. *Journal of Experimental Biology*, *222*(Suppl_1), jeb191890.

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E.

(2008). A movement ecology paradigm for unifying organismal movement research.

Proceedings of the National Academy of Sciences, 105(49), 19052-19059.

Pebesma, E.J. (2004). Multivariable geostatistics in S: the gstat package. Computers &

Geosciences, 30: 683-691.

Polhill, J. G., Parker, D., Brown, D., & Grimm, V. (2008). Using the ODD protocol for

describing three agent-based social simulation models of land-use change. Journal of

Artificial Societies and Social Simulation, 11(2), 3.

R Core Team. 2020. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria. Available: http://www.r-project.org

Railsback, S. F., Lytinen, S. L., & Jackson, S. K. (2006). Agent-based simulation platforms:

Review and development recommendations. Simulation, 82(9), 609-623.

Railsback, S. F., Harvey, B. C., Jackson, S. K., & Lamberson, R. H. (2009). InSTREAM: the

individual-based stream trout research and environmental assessment model. Gen. Tech. Rep.

PSW-GTR-218. Albany, CA: US Department of Agriculture, Forest Service, Pacific

Southwest Research Station. 254 p, 218.

Rohwer, S., Butler, L. K., Froehlich, D. R., Greenberg, R., & Marra, P. P. (2005). Ecology

and demography of east–west differences in molt scheduling of Neotropical migrant

passerines. *Birds of Two Worlds: The Ecology and Evolution of Migration (R. Greenberg and*

PP Marra, Eds.). Johns Hopkins University Press, Baltimore, Maryland, 87-105.

Tang, W., & Bennett, D.A. (2010). Agent-based modeling of animal movement: a review.

Geography Compass, 4(7), pp.682-700.Thiele, J. C., Kurth, W., & Grimm, V. (2012).

RNetLogo: An R package for running and exploring individual-based models implemented in

NetLogo. Methods in Ecology and Evolution, 3(3), 480-483.

Thompson, C. W. (1991). Is the Painted Bunting actually two species? Problems determining

species limits between allopatric populations. *The Condor*, *93*(4), 987-1000.

Topping, C. J., Hansen, T. S., Jensen, T. S., Jepsen, J. U., Nikolajsen, F., & Odderskær, P.

(2003). ALMaSS, an agent-based model for animals in temperate European landscapes.

Ecological Modelling, 167(1-2), 65-82.

Thurfjell, H., Ciuti, S. and Boyce, M.S. (2014). Applications of step-selection functions in

ecology and conservation. Movement ecology, 2(1), pp.1-12.

Uhlenbeck, G. E., & Ornstein, L. S. (1930). On the theory of the Brownian motion. Physical

review, 36(5), 823

van der Vaart, E., Beaumont, M. A., Johnston, A. S., and Sibly, R. M. (2015). Calibration and

evaluation of individual-based models using Approximate Bayesian Computation. Ecological

Modelling, 312, 182-190.

Vermote, Eric; NOAA CDR Program. (2019): NOAA Climate Data Record (CDR) of

AVHRR Normalized Difference Vegetation Index (NDVI), Version 5. NOAA National

Centers for Environmental Information. https://doi.org/10.7289/V5ZG6QH9.

Willem, L., Verelst, F., Bilcke, J., Hens, N., & Beutels, P. (2017). Lessons from a decade of

individual-based models for infectious disease transmission: a systematic review (2006-

2015). BMC infectious diseases, 17(1), 1-16.

Williams, H. J., Taylor, L. A., Benhamou, S., Bijleveld, A. I., Clay, T. A., de Grissac, S., and

Börger, L. (2020). Optimizing the use of biologgers for movement ecology research. Journal

of Animal Ecology, 89(1), 186-206.

Wong, D. W. (2016). Interpolation: Inverse-Distance Weighting. International Encyclopedia

of Geography: People, the Earth, Environment and Technology: People, the Earth,

Environment and Technology, 1-7.

Yin, S., Xu, Y., de Jong, M.C., Huisman, M.R., Contina, A., Prins, H.H., Huang, Z.Y. and de

Boer, W.F., 2022. Habitat loss exacerbates pathogen spread: An Agent-based model of avian

influenza infection in migratory waterfowl. PLoS Comput Biol 18(8): e1009577.

https://doi.org/10.1371/journal.pcbi.1009577.