Synaptotagmin-1-dependent phasic axonal dopamine release is dispensable for basic

motor behaviors in mice

Benoît Delignat-Lavaud[1,2,3], Jana Kano†[1,2,3], Charles Ducrot†[1,2,3], Ian Massé[4], Sriparna
Mukherjee[1,2,3,5], Nicolas Giguère[1,2,3], Luc Moquin[6], Catherine Lévesque[7], Samuel
Burke[1,2,3], Raphaëlle Denis[1,2,3], Marie-Josée Bourque[1,2,3,5], Alex Tchung[1,2,3], Pedro Rosa-
Neto[6], Daniel Lévesque[7], Louis De Beaumont[4], Louis-Éric Trudeau*[1,2,3,5]

[1]Department of Pharmacology and Physiology, Faculty of Medicine, Université de Montréal,
Montreal, Québec, Canada
[2]Department of Neurosciences, Faculty of Medicine, Université de Montréal, Montreal,
Québec, Canada
[3]SNC and CIRCA Research Groups, Université de Montréal, Montréal, Québec, Canada
[4]Hôpital du Sacré-Cœur-de-Montréal, CIUSSS NIM, Université de Montréal, Montreal,
Québec, Canada
[5]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy
Chase, MD, 20815.
[6]Centre intégré universitaire de santé et de services sociaux (CIUSSS) de l'Ouest-de-l'Île-de-
Montréal; Department of Neurology and Neurosurgery, Psychiatry and Pharmacology and
Therapeutics, McGill University, Montreal, Québec, Canada
[7]Faculty of Pharmacy, Université de Montréal, Montreal, Québec, Canada
22

† Equal contribution

* Corresponding author

Correspondence and material requests should be addressed to Dr. Louis-Eric Trudeau (louis-

eric.trudeau@umontreal.ca).

Abstract

In Parkinson's disease (PD), motor dysfunctions only become apparent after extensive

loss of DA innervation. This resilience has been hypothesized to be due to the ability of many

motor behaviors to be sustained through a diffuse basal tone of DA; but experimental

evidence for this is limited. Here we show that conditional deletion of the calcium sensor

synaptotagmin-1 (Syt1) in DA neurons (Syt1 cKO^{DA} mice) abrogates most activity-

dependent axonal DA release in the striatum and mesencephalon, leaving somatodendritic

(STD) DA release intact. Strikingly, Syt1 cKO^{DA} mice showed intact performance in multiple

unconditioned DA-dependent motor tasks and even in a task evaluating conditioned

motivation for food. Considering that basal extracellular DA levels in the striatum were

unchanged, our findings suggest that activity-dependent DA release is dispensable for such

tasks and that they can be sustained by a basal tone of extracellular DA. Taken together, our

findings reveal the striking resilience of DA-dependent motor functions in the context of a

near-abolition of phasic DA release, shedding new light on why extensive loss of DA

innervation is required to reveal motor dysfunctions in PD.

Keywords: dopamine, exocytosis, synaptotagmin, somatodendritic, movement

Introduction

The neuromodulator dopamine (DA) plays a key role in motor control, motivated behaviors and cognition[1,2]. Blockade of DA receptors severely impairs most motor behaviors[3] and a severe loss of DA in the striatum and other brain regions leads to the characteristic motor dysfunctions of Parkinson's disease (PD). It is striking that in PD, motor dysfunctions only appear when the striatum is severely denervated and that DA replacement therapy with L-DOPA is able to restore motor functions by boosting DA production in the remaining, sparse DA axonal fibers[4]. Such observations are puzzling and ill-understood and suggest the possibility that a minimal basal tone of DA, coupled with adaptations to the DA system to boost its sensitivity, are sufficient to maintain motor functions[5].

Compatible with such observations, DA is thought to act in the brain as a neuromodulator involved in a form of "volume transmission" and not as a point-to-point fast neurotransmitter[6-9]. DA release occurs not only from release sites in the vast axonal arbors of DA neurons, but also from the neurons' somatodendritic (STD) compartment[10]. The molecular machinery underlying exocytotic DA release from terminals[11-14] or dendrites[10,11,15-18] is presently fragmentary. Discoveries on such mechanisms are likely to lead to a better understanding of the functions and connectivity of all classes of modulatory neurons.

Vesicular exocytosis requires the concerted action of SNARE proteins and calcium sensors from the synaptotagmin family (Syt)[19]. Of the 17 Syt isoforms identified so far, only Syt1, 2, 3, 5, 6, 7, 9, and 10 have been reported to bind calcium and drive vesicular fusion[20]. Syt1, 2, and 9 were confirmed as calcium sensors for fast synaptic neurotransmitter release[21]. Syt1 was first shown to play a key role in axonal DA release in primary DA neurons[11] in which it is present at both synaptic and non-synaptic release sites[9], but most likely absent

from dendrites[11]. Recent work has confirmed and extended this finding in the intact brain by

showing that Syt1 is essential for evoked DA release[22]. Syt4 and Syt7 were recently shown to

control STD DA release, suggesting a distinct molecular machinery compared to axonal DA

release[11,23].Very recent functional evidence also supports the possibility that Syt1 is also

involved in DA release occurring in the ventral midbrain[24,25].

Here, we defined the consequence of loss of Syt1 in DA neurons by evaluating the

impact of this deletion on DA-dependent behaviors and by characterizing the selective roles

of Syt1 in axonal and STD DA release. We find that basic unconditioned motor functions,

and conditioned motivation for food are intact in mice lacking Syt1 in DA neurons (Syt1

cKO^{DA}). Combined with our observations that loss of Syt1 leads to extensive loss of axonal

DA release in the striatum, to partial loss of DA release in the mesencephalon and to

unaltered extracellular DA levels in the striatum and mesencephalon, our findings suggest

that basal unconditioned motor functions in rodents only require the maintenance of a basal

tone of extracellular DA largely independent of Syt1-dependent phasic release. We

hypothesize that this situation is very similar to pre-symptomatic stages of PD and sheds light

on the surprising resilience of motor functions after extensive loss of evoked DA.

Results

Syt1 is the main calcium sensor for fast axonal dopamine release.

In line with its high expression and localization in the axonal varicosities of DA neurons, Syt1 is a key regulator of activity-dependent DA release[11,22,9]. We generated conditional deletion of Syt1 in DA neurons (Syt1 cKO^{DA}), by crossing Syt1$^{lox/lox}$ mice with DAT^{IREScre} mice (**Fig. 1A**). Validating earlier results, we found using FSCV *ex vivo* in brain slices of Syt1$^{+/+}$, Syt1$^{+/-}$ and Syt1$^{-/-}$ mice an extensive reduction of phasic DA release induced by single electrical pulses in both the dorsal striatum (**Fig. 1B**: 1.42 ± 0.15 µM in Syt1$^{+/+}$; n = 9 mice (5 males/ 4 females) vs 1.123 ± 0.091 µM in Syt1$^{+/-}$; n = 8 mice (4M/4F) and 0.074 ± 0.012 µM in Syt1$^{-/-}$; n = 8 mice (2M/6F); 1-way ANOVA with Tukey, $F_{(2,22)}$ = 43.02, p = 0,139 for Syt1$^{+/-}$ and p < 0.0001 for Syt1$^{-/-}$) and ventral striatum (**Fig. 1C**: 1.252 ± 0.08 µM in Syt1$^{+/+}$; n = 9 mice (5M/ 4F) vs 0.999 ± 0.03 µM in Syt1$^{+/-}$; n = 8 mice (4M/4F) and 0.144 ± 0.019 µM in Syt1$^{-/-}$; n = 8 mice (2M/6F); 1-way ANOVA with Tukey, $F_{(2,22)}$ = 109, p = 0,0098 for Syt1$^{+/-}$ and p < 0.0001 for Syt1$^{-/-}$) of Syt1$^{-/-}$ mice. In Syt1$^{-/-}$ mice, the residual signal was low in amplitude but still represented DA, as confirmed by the cyclic voltammogram (**Fig. S1A and 1B**). Stimulation with 10 Hz pulse trains, although prolonging DA release kinetics (results not shown), did not enhance peak DA overflow in the dorsal or ventral striatum of Syt1$^{-/-}$ mice (**Fig. S1C**). Mice in which DA neurons expressed a single allele of Syt1 also showed a small decrease in DA release, which reached significance in the ventral striatum (0.99 ± 0.03 µM, n = 8 Syt$^{+/-}$ mice (4M/4F); 1-way ANOVA with Tukey, $F_{(2,22)}$ = 109, p < 0.0001) (**Fig. 1C**). This observation argues for the existence of only a modest safety factor in the amounts of Syt1 required for normal function of DA neuron terminals. We conclude that Syt1 acts as the main calcium sensor for fast activity-dependent DA release in the striatum, but that other calcium sensors are also likely to play a

complementary role, which may be more extensive in the ventral (nucleus accumbens)

compared to dorsal striatum.

Syt1 deletion also reduces DA release in the ventral mesencephalon.

As DA release in cell body region of DA neurons is also calcium-dependent and since

synaptotagmin isoforms and active zone proteins have been suggested to have a role in STD

DA release[11,26,23], we also examined activity-dependent DA overflow in the ventral

mesencephalon (**Fig. 1D**). We first measured DA release in the VTA of Syt1$^{-/-}$ mice using an

optimal FSCV paradigm, with pulse-train stimulation (30 pulses at 10 Hz) and aCSF

containing the DAT blocker nomifensine and the D2 antagonist sulpiride, allowing the

detection of STD DA release without the influence of DA uptake and D2 autoreceptors

activation[23]. We observed a robust ≈68% decrease of activity-dependent DA overflow in

Syt1$^{-/-}$ mice (0.09 ± 0.01 µM, n = 8 (2M/6F), vs 0.28 ± 0.03 µM in n = 9 Syt1$^{+/+}$ mice

(5M/4F); 1-way ANOVA with Tukey, $F_{(2,22)}$ = 20.07, p < 0.0001). This decrease was also

significant in heterozygote mice (0.20 ± 0.02 µM, n = 8 (4M/4F); 1-way ANOVA with

Tukey, $F_{(2,22)}$ = 20.07, p = 0.0310). Similar recordings performed in the SNc also revealed a

robust reduction (≈65%) of evoked DA overflow (**Fig. S1C** - 0.09 ± 0.01 µM, n = 5 Syt1$^{-/-}$

mice (4M/1F) vs 0.25 ± 0.04 µM, n = 6 Syt1$^{+/+}$ male mice; 1-way ANOVA with Tukey, $F_{(2,13)}$

= 8.86, p = 0.0053).

Because there is yet no quantitative evidence for the presence of Syt1 in the STD

compartment of DA neurons[11] or any other neurons, this finding was unexpected.

Interestingly, although the VTA region is well known for containing the cell body and

dendrites of DA neurons, it has also been previously suggested to contain a small contingent

of local dopaminergic axon collaterals with potential axonal release sites[10]. The available

anatomical data is however limited[27,28]. On the other hand, the SNc is believed to be devoid

of axonal DA release sites[29–31], but here again, the supporting data is limited.

Somatodendritic optogenetic stimulation reveal unaltered STD DA release in the

absence of Syt1.

In view of the surprising reduction of evoked DA overflow in the ventral

mesencephalon of Syt1$^{-/-}$ mice, we devised a strategy to trigger STD DA release more

selectively. For this, we combined FSCV with selective optogenetic stimulation of the STD

compartment of neurons by expressing a STD-targeted version of channelrhodopsin (ChR2-

Kv) (**Fig. 2**). Kv2.1 channels were previously reported to be restricted to the somatic and

proximal dendritic membrane and absent from distal dendritic membrane, axons, and nerve

terminals of cortical and hippocampal neurons[32–34]. A 65 amino acid motif of the Kv2.1

voltage-gated potassium channel fused with the carboxy terminus of ChR2-EYFP *(Baker et*

al., 2016) was inserted in a Cre-dependent AAV vector (AAV2/5-hsyn-DIO-ChR2-eYFP-

Kv). A standard cell-wide hChR2 (AAV2/5-hsyn-DIO-ChR2-eYFP), previously used to

trigger DA release in the striatum[36] and mesencephalon[23] was used as a control.

Validation of the construct in primary DA neurons from DAT^{IRES}Cre mice showed the

expected STD expression of the ChR2-Kv construct, with no expression in TH$^+$/MAP2$^-$

processes and varicosities (**Fig. S2**). Expression of the construct *in vivo* in DAT^{IRES}Cre mice

was efficient and restricted to the ventral mesencephalon (**Fig. 2A**). Confocal imaging at 60X

revealed a membrane localization of ChR2-Kv at the soma of DA neurons (**Fig. 2B**). In the

striatum, no eYFP signal was detected in the dorsal striatum but sparse signal was detectable

in the ventral striatum, suggesting limited expression in a subset of VTA DA neuron axons in

addition to the predominant STD expression. In line with this limited axonal expression of

ChR2-Kv, DA release triggered by pulse-train optogenetic stimulation of ChR2-Kv in the

dorsal striatum of Syt1$^{+/+}$ mice was about 10-fold lower compared to release evoked with

conventional ChR2 (0.13 ± 0.02 µM, n = 7 mice (5M/2F) vs 1.22 ± 0.16 µM, n = 5 mice

(3M/2F); 2-way ANOVA with Tukey, $F_{(2, 36)}$ = 62.84, p <0.0001), and 2-fold lower in the

ventral striatum (0.79 ± 0.12 µM, n = 7 mice (5M/2F) vs 1.58 ± 0.29 µM, n = 5 mice

(3M/2F); 2-way ANOVA with Tukey, $F_{(2, 36)}$ = 16.49, p = 0,0003) (**Figs. 2C and 2D**).

Intriguingly, DA overflow evoked with ChR2-Kv in the dorsal striatum of Syt1$^{-/-}$ mice

(0.13 ± 0.05 µM) was not significantly different from the release obtained in Syt1$^{+/+}$ mice

(0.13 ± 0.02 µM) (2-way ANOVA with Tukey, $F_{(2, 36)}$ = 62,84, p >0,9999). In the ventral

striatum, evoked DA overflow in Syt1$^{-/-}$ mice was more than 2-fold lower compared to

Syt1$^{+/+}$ mice (0.33 ± 0.07 µM, n = 9 mice (6M/3F) vs 0.79 ± 0.12 µM, n = 7 mice (5M/2F), 2-

way ANOVA with Tukey, $F_{(2, 36)}$ = 16.49, p = 0,0256) (**Figs. 2C and 2D**). The sodium

channel blocker TTX (1 µM) abolished this small axonal DA release in all mice, thus

demonstrating the requirement of action potentials for this response (**Fig. 2H**).

In the VTA (**Fig. 2E**), the release of DA induced by optogenetic train stimulation with

hChR2 was easily detectable and of an amplitude similar to previously reported (203 ± 13

nM, n = 5 mice (3M/2F)[23]. Release evoked by optogenetic stimulation with ChR2-Kv in

Syt1$^{+/+}$ mice was significantly lower (132 ± 16 nM, n = 7 mice (5M/2F); 2-way ANOVA with

Šidák, $F_{(2, 36)}$ = 30.23, p = 0.0005) compared to hChR2 (203 ± 13 nM, n = 5 mice (3M/2F)

(**Fig. 2E**) and the signal detected in Syt1$^{-/-}$ mice was lower compared to Syt1$^{+/+}$ mice (57 ± 7

nM, n = 9 mice (6M/3F); 2-way ANOVA with Šidák, $F_{(2, 36)}$ = 30.23, p <0.0001). To exclude

any contribution from axonal DA release, additional FSCV recordings were obtained in the

presence of TTX (1 µM). Although STD DA release is normally activity-dependent and

blocked by TTX, we hypothesized that membrane depolarization induced by ChR2-Kv

activation might trigger sufficient calcium influx to trigger DA release from the STD

compartment. Under these conditions, STD DA release was indeed evoked by optogenetic

stimulation of ChR2-Kv in Syt1$^{+/+}$ mice (44 ± 8 nM, n = 7 mice (5M/2F)) and the amount of

release was not significantly reduced in Syt1$^{-/-}$ mice (28 ± 7 nM, n = 9 (6M/3F)). Together

these observations suggest that when STD DA release is triggered using an approach that

minimizes any axonal activation, the signal that is detected is unaltered by loss of Syt1. The

results also suggest that some of the DA release evoked by train stimulation in the ventral

mesencephalon includes an axonal component, with electrical and optical stimulation with

hChR2 therefore triggering a mixture of axonal and STD DA release.

Syt1 cKO^{DA} mice do not exhibit substantial motor defects.

DA is known to play a key role in multiple forms of movement[37,38]. Notably, loss of

striatal DA innervation in PD is responsible for the cardinal motor features observed in the

disease[39]. Blockade of DA receptors induces catalepsy and a range of other motor deficits in

rodents and other species[3,40]. Based on our results showing a dramatic impairment of phasic

axonal DA release in Syt1$^{-/-}$ mice, we hypothesized that this would lead to major motor

deficits in Syt1 cKO^{DA} mice.

First, we evaluated motor coordination using the rotarod (**Fig. 3A**) and the pole test

(**Fig. 3B**). Surprisingly, Syt1$^{-/-}$ and Syt1$^{+/-}$ showed no deficits and even exhibited better

performance on the rotarod task than wild type littermates, with a significantly higher latency

to fall from the rod (124 ± 12s for 8 Syt1$^{+/+}$ mice (5M/3F) vs 197 ± 23s for 8 Syt1$^{+/-}$ mice

(4M/4F); 1-way ANOVA with Dunnett test, $F_{(2, 21)}$ = 4.51, p = 0.0460 and 188 ± 20s for 8

Syt1$^{-/-}$ mice (2M/6F), 1-way ANOVA with Dunnett test, $F_{(2, 21)}$ = 4.51, p = 0.0220). In the

pole test, no statistical difference was observed between genotypes for the time required for

the animals to orient themselves facing in a downward direction (t-turn). For the time

required to climb-down the pole, no difference was detected between Syt1$^{+/+}$ and Syt1$^{-/-}$ mice,

although Syt1$^{+/-}$ mice performed the task faster than Syt1$^{+/+}$ mice (8 ± 0.7s for Syt1$^{+/-}$ vs 11 ±

0.5s for Syt1$^{+/+}$, n = 8 (5M/3F); 1-way ANOVA with Dunnett test, $F_{(2, 21)}$ = 4.706, p = 0.0414). Front paw grip strength was evaluated in the grip test (**Fig. 3C**). No statistical differences were observed regarding the force developed by the mice on the grid.

We next measured spontaneous locomotion and locomotion induced by the psychostimulant drugs cocaine (20 mg/kg) and amphetamine (5 mg/kg). We measured the traveled distances using an open field for 20 min, followed by another 40 min after drug or vehicle injection (0.9% saline). No difference was detected between Syt1$^{+/+}$ (n = 8 (4M/4F)) and Syt1$^{-/-}$ (n = 8 (3M/5F)) mice on basal locomotion after saline injection. A mixed-effects ANOVA model with the Geisser-Greenhouse correction revealed a significant effect of time ($F_{(5,949, 123,8)}$ = 29,21, p < 0,0001), but not of genotype ($F_{(2, 21)}$ = 1,745, p = 0.1991), nor an interaction between time x genotype ($F_{(22, 229)}$ = 1,285p = 0.1825) (**Fig. 3D**), with only a non-significant trend for higher traveled distance within the 5 min post injection in Syt1$^{+/-}$ mice (**Fig. 3L**, 91% of baseline ± 16%, n = 8 (4M/4F) Syt1$^{+/-}$ mice vs 56 ± 13%, n = 8 (4M/4F) Syt1$^{+/+}$ mice; 1-way ANOVA with Dunnett test, $F_{(2, 21)}$ = 3.068, p = 0.1211). The global decrease in locomotion over time after saline injection reflects the normal habituation to the open field.

The locomotor response to the DA transporter blocker cocaine was comparable between genotypes. A mixed-effects ANOVA model with the Geisser-Greenhouse correction revealed only a significant effect of time ($F_{(1,975, 36,63)}$ = 27,28, p < 0,0001), but not of genotype ($F_{(2, 19)}$ = 0,1283, p = 0.8803) nor an interaction between time x genotype ($F_{(22, 204)}$ = 0,6858, p = 0.8509) (**Fig. 3E**). The peak magnitude of the increase in locomotion was also not different between groups (**Fig. 3G**). However, locomotion induced by the DA releaser amphetamine was sharply elevated in Syt1$^{-/-}$ mice. A mixed-effects ANOVA model with the Geisser-Greenhouse correction revealed a significant effect of time ($F_{(2,79, 57,28)}$ = 50,76, p < 0,0001), of genotype $F_{(2, 21)}$ = 6,902, p = 0.005) and an interaction between time x genotype

$F_{(22, 226)} = 3,218$, p < 0,0001). The difference between groups became significant 10 min after

the injection (+714 ± 103% in Syt1$^{-/-}$ mice, n = 8 (2M/6F) vs +252 ± 29% in Syt1$^{+/+}$, n = 8

(4M/4F); Mixed-effect ANOVA model with Dunnett test, $F_{(22, 226)} = 3.218$, p = 0.0046), and

was maximal 20 min post-injection (+1045 ± 126% vs +523 ± 58% ; Mixed-effect ANOVA

model with Dunnett test, $F_{(22, 226)} = 3.218$, p = 0.0069) (**Fig. 3F**). Cumulative distance for the

40 min under treatment confirmed this effect with an average traveled distance for Syt1$^{-/-}$

mice of +777 ± 67% vs +472 ± 53% for Syt1$^{+/+}$ (1-way ANOVA with Tukey, $F_{(2, 20)} = 7.441$,

p = 0.0072) and +481 ± 73% for Syt1$^{+/-}$ mice, n = 8 (4M/4F) (1-way ANOVA with Tukey,

$F_{(2, 20)} = 7.441$, p = 0.0117) (**Fig. 3G**).

The observed increase in amphetamine-induced locomotion could result from

increased amphetamine-induced DA secretion or from striatal DA receptor sensitization. We

next tested this hypothesis by administering the selective D1 receptor agonist SCH23390 (50

µg/kg), the D2 receptor agonist quinpirole (0.2 mg/kg) or the D2 receptor antagonist

raclopride (1 mg/kg). At the selected doses, all drugs caused an abrupt decrease in

locomotion, with no significant differences between genotypes when considering the

complete drug treatment period. A mixed-effects ANOVA with Geisser-Greenhouse

corrections revealed only a significant effect of time (p < 0,0001, with $F_{(6,457, 165,5)} = 80,09$ for

SCH23390, $F_{(4,397, 105,9)} = 272,4$ for quinpirole and $F_{(2,782, 80,69)} = 378,3$ for raclopride), but not

of the genotype (respectively $F_{(2, 26)} = 0,865$, p = 0,433 ; $F_{(2, 25)} = 2,672$, p = 0,089 and $F_{(2, 319)}$

= 2,474, p = 0,086) nor any interaction between time x genotype for SCH23390 and

raclopride (respectively $F_{(22, 282)} = 1,374$, p = 0,126 and $F_{(22, 319)} = 1,411$, p = 0,106), except

for quinpirole treatment $F_{(22, 265)} = 1,753$, p = 0,022)) (**Fig. 3H-K**). However, comparing the

mean traveled distance within the first 5 min following drug injection, a time likely to

correspond to partial receptor occupancy (**Fig. 3L**), Syt1$^{-/-}$ mice showed a lower traveled

distance in response to the D2 agonist quinpirole compared to Syt1$^{+/+}$ mice (respectively 21 ±

3%, n = 10 (3M/7F), vs 47 ± 8%, n = 8 (2M/6F); Brown-Forsythe ANOVA with $F_{(2,13.48)}$ =

5.266, p = 0.0204 and a Dunnett T3 post hoc test, p = 0,035). The opposite pattern was

detected in response to the D2 antagonist raclopride, with a significantly increased inhibition

of locomotion in the Syt1$^{-/-}$ mice compared to Syt1$^{+/+}$ mice (14 ± 3% in Syt1$^{-/-}$, n = 10 males

vs 3 ± 1% in Syt1$^{+/+}$, n = 10 (4M/6F); 1-way ANOVA with Dunnett test, $F_{(2, 27)}$ = 5.343, p =

0.0056). During this period, no significant differences were once again observed among

genotypes in their response to the D1 agonist SCH23390. These results suggest that D2

receptor neuroadaptations occur after genetic deletion of Syt1 in DA neurons, but this has

only a limited impact on DA-mediated motor behaviors.

Phasic DA release is unnecessary for motivation to work for food

Next, we tested if the abrogation of phasic release of DA in Syt1 cKO mice would

impair learning in an operant food-rewarded nose-poking task. We tested Syt1$^{+/+}$ (n = 8,

3M/7F) and Syt1$^{-/-}$ (n = 10, 5M/5F) mice of both sexes with the open-source nose-poking

feeding devices "FED3"[41]. First, the devices were placed in the home cage of Syt1 animals

and a fixed ratio 1 (FR1) paradigm was used, in which mice learned to nose-poke on the

active port for receiving a single food pellet. We quantified how nose-poking for pellets

varied over the circadian cycle, by running mice on the FR1 task for 3 consecutive days. As

expected, pellet consumption and active pokes were higher during the dark cycle, which is

the active period for mice. A mixed-effects ANOVA model with the Geisser-Greenhouse

correction revealed no significant global effect of genotype for pellet consumption ($F_{(1, 16)}$ =

0,1489, p = 0,7047) nor for nose-poking ($F_{(1, 16)}$ = 0,1468, p = 0,7067). However, a significant

effect of time (respectively $F_{(7,988, 126,0)}$ = 17,85 and $F_{(7,985, 125,9)}$ = 17,91, p < 0,0001) and an

interaction between time x genotype was observed (respectively $F_{(35, 552)}$ = 2,611 and $F_{(35, 552)}$

= 2,631, p < 0,0001) (**Fig. 4B, 4C and Fig. S3B, S3C**). Interestingly, we noted a ~4h window

of enhanced nose-poking rates starting before the onset of the dark cycle (**Fig. 4B, 4C**). This

nocturnal peak of activity was found to be significantly increased at ZT16 in Syt1$^{-/-}$ mice,

only during the first night, with 8.5 ± 4 pellets consumed for 8.4 ± 4 active pokes in Syt1$^{+/+}$

mice (n = 8) versus 26.9 ± 3 for both pellets and active pokes in Syt1$^{-/-}$ mice (n = 10) (2-way

ANOVA with Šidák, $F_{(35, 568)} = 2.49$, p <0.0001). The performance of the mice under the FR1

task was evaluated by measuring the poke efficiency (% of total pokes on the active port)

over the 3 days on FR1. We find a significant effect of time (2-way ANOVA, $F_{(35, 502)} =$

7,059, p < 0,0001) but no significant difference between genotypes (2-way ANOVA, $F_{(1, 502)}$

= 0.5806, p = 0.4464) or interaction between time x genotype (2-way ANOVA, $F_{(35, 502)} =$

1,323, p = 0,1061) (**Fig. 4D**), suggesting that overall, all mice learned how to use the active

port to obtain their food at the same rate.

Next, we examined motivation for food. To determine this, we measured their

willingness to work in a closed-economy food-rewarded operant task, which has been used to

quantify the economic demand for food[42,43], as well as changes in economic demand due to

manipulation of the DA system[44,45]. After completion of the FR1 task, the protocol used in

the FED3 device was changed for a 6-day repeating progressive ratio (PR1) task, in which the

nose-poking requirement began on FR1 and increased by 1 poke each time a pellet was

earned. When mice refrained from poking on either active or inactive port for 30 min, the

ratio was reset to FR1. Once again, nose-poking rates of the mice were higher during the dark

phase of their daily cycle. A mixed-effects ANOVA model with the Geisser-Greenhouse

correction revealed no global significant effect of genotype either for pellet consumption ($F_{(1,}$

$_{16)} = 0,639$, p = 0,436) or nose-poking ($F_{(1, 16)} = 0.715$, p = 0,4103). However, a significant

effect of time (respectively $F_{(71, 1134)} = 8,977$ and $F_{(71, 1134)} = 26,00$, p < 0,0001) and an

interaction between time x genotype was detected for poking activity ($F_{(71, 1134)} = 1,589$, p =

0,002) (**Fig. 4E, 4F and Fig. S3D, S3E**) This result shows that motivation for food varies

over the circadian cycle such that mice are willing to work harder during the dark cycle vs.

the light cycle, in line with previous results[41]. Interestingly, although in PR1, the pattern of

response was not significantly difference between $Syt1^{+/+}$ and $Syt1^{-/-}$ animals in term of

pellets earned (**Fig. 4E**), $Syt1^{-/-}$ showed a significant increase of nose-poking activity during

the first night at 4h and 6h after the onset of the dark cycle, with 785 ± 145 pokes vs 412 ± 58

for $Syt1^{+/+}$ mice at 4h (2-way ANOVA with Šidák, $F_{(71, 1134)} = 1.589$, p <0.0001) (**Fig. 4F**)

and 574 ± 118 pokes vs 314 ± 78 at 6h (2-way ANOVA with Šidák, $F_{(71, 1134)} = 1.589$, p

=0.0106), respectively. The performance of the mice under the PR1 task was evaluated by

measuring poke efficiency (% of total pokes on the active port) over the 6 days on PR1. We

find a significant effect of time (2-way ANOVA, $F_{(71, 1123)} = 5,558$, p < 0,0001) but no

significant difference between genotypes (2-way ANOVA, $F_{(1, 1123)} = 2,577$, p = 0,1087), nor

interaction between time x genotype (2-way ANOVA, $F_{(71, 1123)} = 1,095$, p = 0,281), revealing

that all mice show high performance on this task, with about ~95% of correct pokes (**Fig.

4G**). Arguing against changes in motivation for food, daily breaking points were not different

between $Syt1^{+/+}$ and $Syt1^{-/-}$ mice in this experiment, although they were expectedly higher

during the dark cycle (**Fig. 4H**). Overall, these results suggest that loss of Syt1 in DA neurons

and the associated loss of phasic DA release are not necessary for basic motivation to work

for food.

Basal extracellular DA levels and total tissue DA are not altered in Syt1 cKO^{DA} mice

Taken together, our behavioral results suggest that despite a >90% impairment of

phasic DA release in the striatum and a substantial decrease in DA release in the

mesencephalon, Syt1 cKO^{DA} mice do not exhibit any obvious defects in basic unconditioned

DA-dependent motor tasks as well as in a conditioned food motivation task. Our observations

lead us to hypothesise that such behaviors do not require phasic, activity-dependent DA

release and can be maintained by basic resting levels of extracellular DA, previously shown

to be unimpacted in Syt1 cKO^{DA} mice[22]. To validate this, we used microdialysis in

anesthetized mice implanted with two cannulas to measure extracellular DA and metabolites in the dorsal striatum and in the ventral mesencephalon. Following this sampling, the striatum of each animal was also microdissected and the total striatal content of DA and metabolites were quantified by HPLC (**Fig. 5A**). Our results reveal that striatal tissue contents of DA, DOPAC, serotonin (5-HT) and norepinephrine (NE) were not significantly different in Syt1$^{-/-}$ mice (n = 12, 9M/3F)) compared to Syt1$^{+/+}$ mice (n = 10, 4M/6F) (unpaired t-test, respectively $F_{(9,11)}$ = 1.083, p = 0.946 ; $F_{(9,11)}$ = 2.966, p = 0.857 ; $F_{(9,11)}$ = 1.122, p = 0.124 and $F_{(9,11)}$ = 2.049, p = 0.572) (**Fig. 5B**). In addition, the DA/DOPAC ratio was found to be unchanged in Syt1$^{-/-}$ mice, suggesting an intact DA turnover (unpaired t-test, $F_{(11,9)}$ = 1.001, p = 0.7971). Microdialysis similarly revealed that even if activity-dependent DA release was mostly blocked in Syt1$^{-/-}$ mice, basal extracellular DA levels of these mice were not reduced in comparison to Syt1$^{+/+}$ mice in the striatum (unpaired t-test, $F_{(11,9)}$ = 2.326, p = 0.215) and mesencephalon (unpaired t-test, $F_{(11,8)}$ = 2.507, p = 0.157) (**Fig. 5C**). Similarly, extracellular levels of DOPAC (unpaired t-test, $F_{(11,10)}$ = 1.355, p = 0.416 for striatum, $F_{(9,11)}$ = 2.768, p = 0.684 for midbrain), and NE (Welch's t-test, $F_{(11,9)}$ = 4.158, p = 0.268 for striatum, unpaired t-test $F_{(11,9)}$ = 1.223, p = 0.798 for midbrain) were unchanged. However, we found a significant increase of the extracellular levels of 5-HT in the striatum (0.14 ± 0.032 ng/mL in Syt1$^{-/-}$ vs 0.057 ± 0.006 ng/mL in Syt1$^{+/+}$, Welch's t-test, $F_{(11,7)}$ = 34.72, p = 0.025), and although a similar tendency was noted in the mesencephalon, the effect did not reach significance (0.981 ± 0.009 in Syt1$^{-/-}$ vs 0.500 ± 0.031 ng/mL in Syt1$^{+/+}$, $F_{(11,8)}$ = 4.377, p = 0.055 for midbrain).

Together, these results indicate that the pool of releasable DA is not decreased by loss of Syt1 and that these mice can maintain intact and sufficient levels of extracellular DA to maintain a basal tone of D1 and D2 receptor activation.

Increased D2 autoreceptor and DAT function in Syt1 cKO^{DA} mice

Our behavioral results with D2 receptor ligands further suggest that adaptations are likely to occur in the DA system of Syt1 cKO mice, in a manner that is similar to adaptations occurring at early stages of DA neuron degeneration in PD[46,47]. In particular, D2 receptors can be hypothesized to undergo an increased density and/or affinity.

As a first step to examine underlying adaptations to the DA system in Syt1 cKO^{DA} mice, we used FSCV to probe the functionality of D2 autoreceptors and of the membrane DA transporter (DAT) in the dorsal and ventral striatum. We measured the impact on DA release of the D2 receptor agonist quinpirole (1 µM). In line with previous work showing that D2 autoreceptors negatively regulated DA release[48,49], DA overflow evoked by single electrical pulses was decreased by ≈85% in the dorsal striatum (1.38 ± 0.18 µM pre-treatment vs 0.21 ± 0.07 µM after treatment, n = 9 (5M/4F); Welch's t-test, $F_{(8,8)}$ = 7.4, p = 0.0001) and by ≈84% in the ventral striatum (1.22 ± 0.1 µM pre-treatment vs 0.196 ± 0.06 µM after treatment, n = 9 (4M/4F); unpaired t-test, $F_{(8,8)}$ = 2.79, p < 0.0001) after a 15 min perfusion of quinpirole (**Fig. 6A, 6B**). In Syt1$^{-/-}$ mice, quinpirole caused a complete abolition of detectable DA release in both the dorsal striatum (0.06 ± 0.006 µM pre-treatment, vs 0.00 ± 0.00 µM after treatment, n = 7 (1M/6F); Welch's t-test, $F_{(6,6)}$ = 117.3, p < 0.0001) and ventral striatum (0.13 ± 0.02 µM pre-treatment vs 0.00 ± 0.00 µM after treatment, n = 8 (1M/7F); Welch's t-test, $F_{(7,7)}$ = 144.4, p = 0.0014). This observation further confirms that the remaining signal detected by FSCV in Syt1$^{-/-}$ mice is indeed DA. The increased relative effect of D2 autoreceptor stimulation (**Fig. 3**) also supports the hypothesis of an increased sensitivity of D2 receptors in Syt1$^{-/-}$ mice.

We next evaluated the impact of the D2 autoreceptor antagonist sulpiride, previously shown to enhance DA release triggered by pulse trains[50]. DA release was evoked by pulse trains (30 pulses at 10 Hz) to allow a basal level of D2 autoreceptor activation to occur. DA release evoked in this way in the striatum of Syt1$^{+/+}$ mice was robustly increased by sulpiride (5 µM) in both the dorsal striatum (149%) (1.34 ± 0.10 µM pre-treatment vs 2.0 ± 0.19 µM

after treatment, n = 8 (4M/4F); unpaired t-test, $F_{(7,7)}$ = 3.54, p = 0.0079) and ventral striatum

(167%) (1.13 ± 0.08 µM pre-treatment vs 1.89 ± 0.21 µM after treatment, n = 8 (4M/4F);

Welch's t-test, $F_{(7,7)}$ = 6.9, p = 0.0085) (**Fig. 6C, 6D**). Despite a dramatic impairment of DA

release evoked by train stimulation in Syt1$^{-/-}$ mice (**Fig. S1D**), the relative effect of sulpiride

on evoked DA release was similar in magnitude compared to the wild type animals in the

dorsal striatum, with a +140% increase (0.07 ± 0.01 µM pre-treatment vs 0.09 ± 0.01 µM

after treatment, n = 11 (4M/7F); unpaired t-test, $F_{(10,10)}$ = 1.8, p = 0.0466). In the ventral

striatum, the relative effect of sulpiride on train-evoked DA overflow was higher in Syt1$^{-/-}$

compared to Syt1$^{+/+}$ mice, with a +261% increase (0.13 ± 0.02 µM pre-treatment vs 0.34 ±

0.08 µM after treatment, n = 11 (4M/7F); Welch's t-test, $F_{(10,10)}$ = 14.9, p = 0.0318). Together

with our results with quinpirole, these findings again argue that D2 autoreceptor function is

increased in Syt1$^{-/-}$ mice, especially in the ventral striatum region.

Finally, we evaluated the functionality of the DAT by quantifying the impact of the

DAT blocker nomifensine (5 µM) on DA overflow evoked by train stimulation, used here to

have conditions allowing basal DAT function to be optimally revealed. A near saturating

concentration of nomifensine was used to examine the net contribution of DAT activity on

phasic DA release[51] As expected, such treatment greatly enhanced DA overflow in wild type

animals, with a +288% increase in the dorsal striatum (1.79 ± 0.19 µM pre-treatment vs 5.15

± 0.74 µM after treatment, n = 4 (1M/3F); unpaired t-test, $F_{(3,3)}$ = 14.35, p = 0.0045) and a

+221% increase in the ventral striatum (1.51 ± 0.2 µM pre-treatment vs 3.34 ± 0.52 µM after

treatment, n = 4 (1M/3F); unpaired t-test, $F_{(3,3)}$ = 6.62, p = 0.0173) (**Fig. 6E, 6F**). The relative

effect of nomifensine in Syt1$^{-/-}$ mice was significantly higher than in Syt1$^{+/+}$ mice, with a

+685% increase in the dorsal striatum (0.11 ± 0.03 µM pre-treatment vs 0.74 ± 0.22 µM after

treatment, n = 6 (3M/3F); Welch's t-test, $F_{(5,5)}$ = 64.75, p = 0.0367) and a +544% in the

ventral striatum (0.16 ± 0.03 µM pre-treatment vs 0.87 ± 0.13 µM after treatment, n = 6

(3M/3F); Welch's t-test, $F_{(5,5)}$ = 24.1, p = 0.0023) (**Fig. 6E, 6F**). These results demonstrate

that DAT function is intact in Syt1$^{-/-}$ mice. Interestingly, they further suggest that

nomifensine may act in some way to rescue part of the impaired activity-dependent DA

release in these mice. Although speculative, this could be through its electrogenic activity

that directly modulates membrane potential[52].

Higher D2 receptor density and lower D2 affinity in Syt1 cKO^{DA} mice

Because the maintenance of motor behaviors in Syt1$^{-/-}$ mice could be due in part to

adaptations of DA receptors in the striatum, we next used autoradiography to quantify D1 and

D2 binding using the selective D1 and D2 receptor radioligands [^{3}H]-SCH23390 and [^{3}H]-

raclopride, respectively. No change of Bmax and Kd for D1 binding was observed in the

whole striatum (n = 4 females Syt1$^{+/+}$ mice and n = 4 Syt1$^{-/-}$ (3M/1F) mice ; unpaired t-test,

$F_{(3,3)}$ = 3.53, p = 0,9961 for Bmax, and $F_{(3,3)}$ = 2.80, p = 0.466 for Kd) nor any analyzed

striatal region (dorso-lateral, ventro-lateral, dorso-medial and ventro-medial), indicating that

neither D1 receptor density or affinity for [^{3}H]-SCH23390 was changed in Syt1 cKO^{DA} mice

(**Fig. 7A and 7B**). This lack of change in D1 receptor binding in Syt1 cKO^{DA} mice is in line

with the lack of change in the behavioral response to the D1 receptor antagonist SCH23390

in Syt1$^{-/-}$ mice. However, a significant increase in D2 receptor Bmax and Kd (reduced

affinity) was detected in the whole striatum of Syt1$^{-/-}$ mice (n = 5 Syt1$^{+/+}$ (3M/2F) and 4

males Syt1$^{-/-}$; unpaired t-test, $F_{(3,4)}$ = 4.65, p = 0,0002 for Bmax and $F_{(3,4)}$ = 6.65, p = 0,0025

for Kd) (**Fig. 7C and 7D**). Finally, we estimated the binding potential of D1 and D2

receptors, which is defined by the Bmax/Kd ratio. We found that binding potential of both D1

and D2 receptors was not significantly different between Syt1$^{+/+}$ and Syt1$^{-/-}$ mice (unpaired t-

test, $F_{(3,3)}$ = 2.89, p = 0,2065 for D1 and $F_{(4,3)}$ = 1.95, p = 0,3369 for D2) (**Fig. 7E**). This

suggests that D2 receptor overall binding potential was maintained in Syt1$^{-/-}$ mice. This raises

the possibility of a molecular rearrangement, for example implicating desensitization of some

of the receptors, helping to maintain homeostasis[53,54]. These results provide further support

for the hypothesis that in the absence of phasic DA release, adaptations to DA receptors

combined with maintained basal extracellular DA levels are sufficient to support many types

of DA-dependent behaviors.

435 Adaptations of DA synthesis and packaging in Syt1 cKO[DA] mice

Compensatory changes at early stages of DA neuron degeneration in PD include not

only receptor adaptations, but also enhanced DA synthesis[46,47] and increased axonal

sprouting. The later has been linked to reduced activation of D2 autoreceptors *(Parish et al.,*

2001; Tripanichkul et al., 2003; Fasano et al., 2008; Fasano et al., 2010, 2013; Giguère et

al., 2019). We therefore hypothesized that reduced activity-dependent DA release in Syt1[-/-]

mice might lead to an increased density of DA neuron terminal markers in the striatum. We

next used immunohistochemistry and quantitative confocal microscopy to measure striatal

levels of the DA synthesis enzyme TH, the vesicular DA transporter VMAT2 and the DAT.

Serotonin (5-HT) innervation was also examined because previous work showed that this

system undergoes compensatory sprouting in response to reduced DA levels[61].

We measured signal surface and intensity for these markers in a series of 5 different

striatal slices ranging from bregma +0.98 to bregma -1.06 mm, with a total of 22 different

areas for each hemisphere (**Fig. S4A**). We found a global increase of TH signal surface in

Syt1[-/-] mice (n = 10, 3M/7F) compared to Syt1[+/+] animals (n = 10, 4M/6F), both in the dorsal

striatum (135 ± 6% of control n = 20 hemispheres from 10 mice; 2-way ANOVA with Šidák,

$F_{(1, 76)}$ = 51.84, p = 0.0001) and ventral striatum (148 ± 6% of control, n = 20/10 mice; 2-way

ANOVA with Šidák, $F_{(1, 76)}$ = 51.84, p <0.0001) (**Fig. 8A, 8B**). A global increase of TH

signal intensity was also observed, in the dorsal (117 ± 3% of control, n = 20/10 mice; 2-way

ANOVA with Šidák, $F_{(1, 76)}$ = 43.72, p = 0.0005) and ventral striatum (125 ± 4% of control, n

= 20/10 mice; 2-way ANOVA with Šidák, $F_{(1, 76)}$ = 43.72, p <0.0001) (**Fig. 8C**). A similar

increase in the surface of VMAT2 immunoreactivity (145 ± 10% of control, n = 20/10 mice;

2-way ANOVA with Šidák, $F_{(1, 76)}$ = 25.74, p = 0.0043) and in the intensity of VMAT2 signal

(123 ± 5% of control, n = 20/10 mice; 2-way ANOVA with Šidák, $F_{(1, 76)}$ = 35.40, p = 0.0011)

was observed in the dorsal striatum and in the ventral striatum (signal surface; 157 ± 10% of

control, n = 20/10 mice; 2-way ANOVA with Šidák, $_{(1, 76)}$ = 25.74, p = 0.0003, signal

intensity; 130 ± 5% of control, n = 20/10 mice; 2-way ANOVA with Šidák, $F_{(1, 76)}$ = 35.40, p

<0.0001). Overall, these results suggest that in Syt1$^{-/-}$ mice, the total density of dopaminergic

axonal processes is increased, possibly explaining the selective increase in amphetamine-

induced but not cocaine-induced locomotion in these mice.

Opposite changes in striatal DAT immunostaining were observed in Syt1 cKO^{DA}

mice, with a global decrease of signal surface in the dorsal striatum (62 ± 6% of control, n =

20/10 Syt1$^{-/-}$ and 22/11 Syt1$^{+/+}$; 2-way ANOVA with Šidák, $F_{(1, 80)}$ = 16.24, p = 0.0094) and

in the ventral striatum (64 ± 7% of control, n = 20/10 Syt1$^{-/-}$ and 22/11 Syt1$^{+/+}$; 2-way

ANOVA with Šidák, $F_{(1, 80)}$ = 16.24, p = 0.0131) (**Fig. 8A-8C**). The DAT signal intensity

appeared unchanged in the ventral striatum but was significantly decreased in the dorsal

striatum (89 ± 2% of control, n = 20/10 Syt1$^{-/-}$ and 22/11 Syt1$^{+/+}$; 2-way ANOVA with Šidák,

$F_{(1, 80)}$ = 9.764, p = 0.0280).

These results are compatible with the establishment of compensatory mechanisms

acting together to maintain extracellular DA levels in the striatum Syt1 cKO^{DA} mice. These

compensatory changes in TH and VMAT2 occurred in the absence of any change in DA

neuron number in the VTA and SNc of Syt1$^{-/-}$ mice, as confirmed by stereological counting

of TH-positive neurons in these structures (**Fig. 8D**).

Finally, while 5-HT immunoreactive signal intensity in the striatum remained

unchanged after Syt1 deletion, a small decrease of the 5-HT signal surface was visible in the

dorsal striatum (72 ± 8% of control, n = 20/10 Syt1$^{-/-}$ and 22/11 Syt1$^{+/+}$; 2-way ANOVA with

Šidák, $F_{(1, 80)}$ = 13.58, p = 0.0399) and ventral striatum (65 ± 7% of control, n = 20/10 Syt1$^{-/-}$

and 22/11 Syt1$^{+/+}$; 2-way ANOVA with Šidák, $F_{(1, 80)}$ = 13.58, p = 0.0114) (**Fig. 8A-8C**). This

result suggests a lack of serotoninergic compensatory axonal sprouting in the striatum of Syt1

cKO^{DA} mice.

We also evaluated if these adaptations observed for striatal TH, VMAT2 and DAT

also occurred in the SN and VTA of Syt1 cKO^{DA} mice (**Fig. S5**). Using the same analysis in

both midbrain regions, we found no change of DAT signal surface (2-way ANOVA with

Šidák, $F_{(1,44)}$ = 0.7354, p = 0,99 for SNc and p = 0.47 for VTA, n = 12 hemispheres from 6

Syt1$^{-/-}$ mice (3M/3F) and n = 12 hemispheres from 6 Syt1$^{+/+}$ mice (1M/5F)), but a significant

decrease of TH signal surface in the VTA (2-way ANOVA with Šidák, $F_{(1,44)}$ = 12, p =

0,0156) and increase of VMAT2 signal surface in the SNc (2-way ANOVA with Šidák, $F_{(1, 44)}$ = 7,53, p = 0,0111) (**Fig. S5B**). Such changes cannot be attributed to change in DA

neurons number in both regions (**Fig. 8D**). Moreover, no changes of signal intensity (**Fig.**

S5C) were detected for all markers, suggesting that the extent of the adaptations seen in the

axonal compartment was more limited in the mesencephalon compared to the striatum.

Arguing that compensatory adaptations occur gradually in Syt1$^{-/-}$ mice during the

postnatal period as a result of abrogation of activity-dependent DA release, we observed no

significant changes in the general morphological development of primary postnatal

mesencephalic DA neurons isolated from Syt1$^{-/-}$ P0-P3 pups and grown *in vitro* for 2-weeks

(**Fig. S6**). As expected, Syt1 immunoreactivity was absent from the axonal varicosities of

these neurons (**Fig. S6A**) (n = 29 fields for Syt1$^{-/-}$ and 35 for Syt1$^{+/+}$). A subset of these TH-

positive axonal varicosities was also positive for VMAT2, a proportion that was not different

in Syt1$^{-/-}$ mice compared to Syt1$^{+/+}$ mice (**Fig. S6B, S6G,** n = 50 fields for Syt1$^{-/-}$ and 55 for

Syt1$^{+/+}$). Most of these varicosities were non-synaptic in structure, as revealed by the

observation that only a very small subset was in contact with a MAP2-positive dendritic

segment independently of mouse genotype (**Fig. S6C, S6F,** n = 47 fields for Syt1$^{-/-}$ and 64 for

Syt1$^{+/+}$), in line with recent results [9]. The intensity of TH immunoreactivity in these release

sites and of VMAT2 in the cell body of DA neurons was also unchanged in Syt1$^{-/-}$ mice

compared to Syt1$^{+/+}$ mice (**Fig. S6H, S6J)** (TH; n = 138 fields for Syt1$^{-/-}$ and 166 fields for

Syt1$^{+/+}$, VMAT2; n = 17 Syt1$^{-/-}$ fields and 22 Syt1$^{+/+}$ fields). A small increase in VMAT2

signal in the axonal release sites of Syt1$^{-/-}$ mice was however observed, similar to *in vivo*

observations (6002 ± 732 pixels/field, n = 43 for Syt1$^{-/-}$ fields vs 3765 ± 590 pixels/field, n=

45 for Syt1$^{+/+}$ fields; unpaired t-test, $F_{(42,44)}$ = 1.47, p = 0.0191).

Survival of DA neurons in these cultures over 14 days was not significantly different

between genotypes (**Fig. S6K,** n = 10 coverslips Syt1$^{+/+}$ and 12 Syt1$^{-/-}$). Finally, the number

of axonal branches and their length was unchanged (**Fig. S6L-S6O**). These results argue that,

in vitro, the basic development of DA neurons is not impacted by Syt1 deletion.

Finally, we also unveiled compensatory adaptations in adult Syt1 cKO^{DA} mice by

quantification of mRNA levels in total microdissected ventral mesencephalon (**Fig. S7A,** n =

5 Syt1$^{+/+}$ (1M/4F), 5 Syt1$^{+/-}$ (5M) and 5 Syt1$^{-/-}$ (4M/1F) mice). We observed a significant

upregulation of *Slc6a3* (DAT) mRNA (1.31 fold, 1-way ANOVA with Dunnett, $F_{(2, 12)}$ =

3.779, p =0.0324) and of *Syt4* mRNA (1.23 fold, 1-way ANOVA with Dunnett, $F_{(2, 12)}$ = 6.97,

p = 0.0197) in Syt1$^{+/-}$ mice and an upregulation of *Syt4* mRNA (1.26 fold, 1-way ANOVA

with Dunnett, $F_{(2, 12)}$ = 6.97, p = 0,0095) and *th* mRNA (1.39 fold, 1-way ANOVA with

Dunnett, $F_{(2, 12)}$ = 4.771, p = 0,0179) in Syt1$^{-/-}$ mice. It should be noted that these changes of

expression could be underestimated as our analysis was performed on whole microdissected

ventral midbrains, therefore including non-dopaminergic neurons. This dilution effect is

visible for instance in the Syt1 mRNA quantification, which is only reduced by ≈17% in

Syt1$^{+/-}$ mice (p = 0,0603) and by ≈49% in Syt1$^{-/-}$ mice (p <0.0001). Adaptations also occurred

at protein levels, as measured by Western Blot in total microdissected striatum (**Fig. S7B-**

S7F, n = 10 Syt1$^{+/+}$ (4M/6F), 8 Syt1$^{+/-}$ (5M/3F) and 10 Syt1$^{-/-}$ (7M/3F) mice), with a

significant increase of the VMAT2 in Syt1$^{+/-}$ mice (1.2 fold; 1-Way ANOVA with Brown-

Forsythe correction and Dunnett T3 post hoc test, $F_{(2, 20.13)}$ = 13.36, p = 0.0243) and in Syt1$^{-/-}$

mice (1.38, p = 0.0003). No change in relative protein levels was observed for TH (1-Way

ANOVA, $F_{(2, 25)}$ = 0,2591, p = 0.774) and DAT (1-Way ANOVA, $F_{(2, 25)}$ = 2.381, p = 0.1131)

Acute deletion of Syt1 in adult mice impairs DA release but prevents adaptations.

Since Syt1 cKO^{DA} developed strong adaptations, which likely happened during

postnatal development, we knocked out Syt1 in adult mice by injecting 6-7 weeks old

Syt1$^{lox/lox}$ mice with an AAV vector allowing expression of the Cre recombinase in DA

neurons (AAV9-TH-cre-myc-2A-fusion-red). Three weeks after stereotaxic virus injection in

the mesencephalon of these mice, strong expression of the fusion-red reporter was detected in

a majority of DA neurons (**Fig. 9A**). The reporter was also found in a subset of TH-negative

neurons at the sites of infection, revealing the preferential but imperfect selectivity of the

construct used. FSCV recordings in the striatum and mesencephalon of these mice revealed a

robust decrease of DA release evoked by single electrical pulses compared to controls, both

in the dorsal (0.44 ± 0.04 μM, n = 8 (3M/5F) vs 1.12 ±0.07 μM, n = 7 (2M/5F); 2-way

ANOVA with Šidák, $F_{(1, 39)}$ = 120.7, p <0.0001) and ventral striatum (0.37 ± 0.03 μM, n = 8

(3M/5F) vs 0.89 ± 0.1 μM, n = 7 (2M/5F); 2-way ANOVA with Šidák, $F_{(1, 39)}$ = 120.7, p

<0.0001). DA release in the VTA of Cre virus injected mice showed a strong decrease in

comparison to control mice injected with the control AAV (CTRL: 0.24 ± 0.03 μM, n = 7

(2M/5F), Cre AAV: 0.026 ± 0.01 μM, n = 8 (3M/5F); 2-way ANOVA with Šidák, $F_{(1, 39)}$ =

120.7, p = 0.0171). This result validates the key role of Syt1 as the main calcium sensor

controlling phasic DA release in the adult brain.

Immunohistochemical analysis of TH and DAT in these Syt1 adult KO mice (n = 12

hemispheres/6 mice (3M/3F) in CTRL and n = 12/6 (6F) in Cre AAV) showed that both

global TH and DAT surface and signal intensity were unchanged (2-way ANOVA, $F_{(1, 44)}$ =

3,919, p = 0,0540 and $F_{(1, 44)}$ = 0,2478, p = 0,6211 for TH surface and intensity, and $F_{(1, 44)}$ =

1,645, p = 0,2063 and $F_{(1, 44)}$ = 0,6035, p = 0,4414 for DAT surface and intensity), except for

a slight increase of the TH signal surface in the ventral striatum (+139 ± 18% vs control; 2-

way ANOVA with Šidák, $F_{(1, 44)}$ = 3.919, p =0.0348) (**Fig. 9C-9E**). We also evaluated if

adaptations could occur at the level of the injection site in the SNc and VTA, but we failed to

detect any significant change of TH, VMAT2 and DAT signal surface and intensity (**Fig.S8**).

Overall, these results reveal that adult deletion of Syt1 leads to a strong reduction of activity-

dependent DA release, accompanied by blunted neuroanatomical adaptations in comparison

with conditional Syt1 cKO^{DA} mice.

Discussion

In the present study, we find that Syt1 is the main calcium sensor involved in fast

activity-dependent DA release, confirming previous results obtained first *in vitro*[11], and more

recently *in vivo*[22]. Our work confirms, but also extends, these previous observations in three

ways.

First, while previous *in vivo* work using amperometry concluded that release evoked

by single spikes was mostly abrogated in Syt1$^{-/-}$ mice, we find that a small component of

evoked DA release is in fact maintained and that there was less release remaining in the

dorsal (≈5%) compared to the ventral striatum (≈12%) This remaining release was abolished

using the D2 receptor agonist quinpirole. This observation, together with the analysis of

voltammograms, confirms the dopaminergic origin of the signal detected by FSCV in Syt1$^{-/-}$

mice and strongly suggests the involvement of another calcium sensor mediating fast,

activity-dependent release. Syt7 would be a logical candidate because it was previously

demonstrated to be highly expressed in DA neurons and localized in both axonal varicosities

and in the somatodendritic compartment of these neurons[11]. However, Syt7 constitutive KO

mice were recently found to have no impairment of single-pulse evoked DA release in the

striatum, instead playing a role in STD DA release[23]. This absence of effect at the terminals

echoes previous results in GABAergic neurons where Syt7 deletion failed to reduce fast

synchronous release, slow asynchronous release, or short-term synaptic plasticity[62]. However,

a role of Syt7 in asynchronous release of neurotransmitter was demonstrated using double

knock-out of Syt1 and Syt7[63,64]. We can thus speculate that a role of Syt7 in gating evoked

DA release would be unveiled by a double KO of Syt1 and 7 in DA neurons. Other

synaptotagmin isoforms expressed in DA neurons could also be involved in supporting

release in the absence of Syt1. We found that Syt4 mRNA is also upregulated in Syt1$^{-/-}$ mice.

However, this isoform appears to only play a role in STD DA release in DA neurons[11,23].

Syt11, suggested to participate in regulating DA vesicle pool replenishment[65], could also

potentially be involved. Calcium sensors of the Doc2 family are also potential candidates

because they have previously been shown to act as modulators of spontaneous synaptic

transmission by a calcium-independent mechanism in cortical synapses[66]. However,

expression of Syt11 and Doc2b were found to be unchanged in our RT-qPCR analysis of

Syt1 cKO^{DA} mice. Irrespective of the specific calcium sensor mediating residual evoked DA

release in Syt1 cKO^{DA} mice, it is possible that this release could also be mobilized through

activation of nicotinic acetylcholine receptors located on DA neuron terminals[67], something

that should be examined in future work.

Secondly, results from the current study extend previous work by showing that basal

unconditioned motor functions in mice are unimpaired under conditions where most activity-

dependent DA release from VTA and SNc DA neurons is blocked. This observation may be

surprising considering that axonal and STD DA release is known to play a critical role in

motor control[37,38]. Our findings that extracellular levels of DA detected by microdialysis were

unchanged in the Syt1 cKO^{DA} mice is compatible with previous work showing only minimal

changes in similar Syt1 cKO^{DA} mice[22]. Our work suggests that basal levels of DA in the

striatum, deriving either from spontaneous exocytosis and/or from the very small remaining

phasic or desynchronized release, are sufficient to support these basic behaviors and that a

large phasic DA signal is not necessary. Additional measurements of basal extracellular DA

levels using the more precise no-net-flux method would be useful to extend the present

results.

Moreover, we find that Syt1 cKO^{DA} mice, despite a dramatic impairment of evoked

DA release, produce viable, fertile offspring's, with normal body weight (data not shown)

and no gross physical or behavioral abnormalities. However, we observed multiple

behavioral, physiological and anatomical phenotypes pointing towards the establishment of

functional adaptations that tune the DA system to the existing conditions by maximizing DA

production and packaging as well as striatal DA receptor activation.

In open-field experiments, we found that while cocaine-induced locomotion was

unchanged in Syt1$^{-/-}$ animals, amphetamine-induced locomotion was robustly enhanced.

Cocaine is known to increase extracellular DA by blocking reuptake through the DAT, while

amphetamine induces a similar elevation by a more complex mechanism including impaired

VMAT2-dependent vesicular storage and reverse transport[68]. These results, together with our

observation of unchanged tissue DA levels, indicate that while the total pools of DA are not

altered in Syt1 KO animals, there are pre- and postsynaptic adaptations taking place after the

loss of Syt1. The observed increase in VMAT2 expression in Syt1 KO mice might explain in

part the differential impact of loss of Syt1 on amphetamine and cocaine-induced locomotor

activation. Increased mobilization of VMAT2-dependent DA stores might allow a higher rate

of amphetamine-induced DAT reverse transport. Although limited by the use of single doses,

our behavioral results with DA receptor agonists and our autoradiography results suggest that

D2 receptors are most affected, with an increased response to the D2 agonist quinpirole and a

reduced response to the D2 antagonist raclopride, together with increased D2 receptor

binding in the ventro-lateral striatum. The D1 receptor system was unchanged in Syt1$^{-/-}$

animals, as no differences emerged for locomotion induced by the D1 agonist SCH23390, or

for D1 ligand binding. Regarding striatal DA receptors, the absence of effects at D1 receptors

and unchanged binding potential at the D2 receptors suggest that compensatory changes were

able to maintain a normal balance between direct and indirect pathways, which is consistent

with the mild motor phenotype observed in Syt1$^{-/-}$ mice. Further work to distinguish changes

in D2 autoreceptors from changes in striatal D2 receptors would help to clarify the

interpretation. For this, dose-response experiments would be required.

We also find clear presynaptic adaptations illustrated by an increase of TH and

VMAT2 signal surface and intensity and a decrease in DAT immunoreactivity in both the

ventral and dorsal striatum. One interpretation of the increases in TH and VMAT2 signal is

that there may be an increase in the axonal arborization of DA neurons in Syt1$^{-/-}$ mice as well

as a decrease in reuptake capacity. We and others have previously provided strong support

for the hypothesis that DA, through D2 autoreceptors is a key regulator of axon growth by

DA neurons[69,56,60]. In the context of a strong reduction in evoked DA release, it is to be

expected that D2 autoreceptors would be chronically under-stimulated leading to removal of

a brake on axon growth and thus an enhanced density of terminals, in line with our recent

work showing axonal arbor expansion after conditional deletion of the D2 receptor gene in

DA neurons[60]. Further experiments will be required to demonstrate such an axonal arbor

expansion more directly in Syt1$^{-/-}$ mice. The increase in TH immunoreactivity, but without a

similar increase in total TH protein as seen in western blot, suggests the possibility that there

may be more DA neuron release sites, with a lower relative level of TH in each, a model that

would be compatible with the lack of change in total tissue DA levels. The decrease in DAT

immunoreactivity suggests reduced expression of DAT in the KO mice. Our qPCR results

show no significant change in DAT mRNA levels in the KO mice. Western blot also failed to

detect a change in DAT protein, although an increase in VMAT2 levels was confirmed with

this technique. Although speculative, it is possible that the antibody used for

immunohistochemical detection of DAT preferentially recognized cell surface DAT in fixed

tissue and that it is specifically cell surface DAT levels that are decreased. Experiments to

specifically quantify cell surface DAT levels would be useful to extend our findings. A

reduced level of DAT in DA neuron release sites could in theory lead to reduced rates of DA

reuptake and thus contribute to maintaining extracellular DA levels in the Syt1 deficient

mice. Our FSCV experiments were not able to resolve this as the very small responses

remaining in the KO mice precluded a reliable analysis of estimated reuptake kinetics.

Quantifications of DAT function in striatal synaptosome preparations might help to resolve

this.

Compensatory adaptations to the 5-HT system could also potentially be involved in

maintaining performance in some of the behavioral tasks that were examined. While we did

not detect morphological evidence for compensatory axonal sprouting of 5-HT axons in the

striatum, we did observe an increase of extracellular 5-HT levels in the striatum of the Syt1

deficient mice.

Arguing that compensatory changes in the DA system of Syt1$^{-/-}$ mice result from an

early and substantial loss of evoked DA release, we found little if any changes in TH and

DAT immunoreactivity in adult mice in which Syt1 was deleted by injecting a TH-Cre virus

in the ventral mesencephalon. These results indicate that compensatory events are likely to

take place during early pre- and postnatal development of the Syt1 KO mice.

A third notable discovery deriving from the present work is the unexpected

impairment of activity-dependent DA release in the ventral mesencephalon. This observation

is surprising because on the one hand, DA release in this region is thought to derive mostly

from the STD compartment of DA neurons, and on the other hand, Syt1 is typically known as

a release site-specific protein. In line with this, previous work in primary DA neurons failed

to detect Syt1 immunoreactivity in the cell body and dendrites of DA neurons, with the

protein detected in the vast majority of non-synaptic and synaptic terminals along DA neuron

axons[11,9]. One possibility is that part of the DA release detected in the VTA and SNc derives

from local DA neuron axon collaterals, which would contain Syt1. However, very little solid

anatomical data exists currently to support the existence of a large number of such local

collaterals, although minimal quantitative data suggest that the VTA contain axonal

collaterals arising from its own axons and from SNc DA neurons[27,28]. To address this

question more directly, we devised a strategy to selectively activate the STD compartment of

DA neurons using an optogenetic approach. For this, we expressed a Kv2.1-ChR2-YFP

fusion construct in DA neurons that we showed was predominantly expressed in the STD

compartment of DA neurons. While the expression of this ChR2 protein was purely STD in

primary DA neurons, a small contingent of axons in the ventral striatum appeared to contain

the protein. As optically-evoked DA release in the ventral striatum of these mice was blocked

by suppressing firing with TTX, we performed all experiments in ventral mesencephalic

slices in the present of TTX, thus limiting any contribution of axonal processes to evoked DA

release. We found that STD DA release was still detectable under such conditions and this

STD DA release was not reduced by *syt1* gene deletion. Together, these results provide

support for the fact that activity-dependent DA release detected in the VTA, and potentially also in the SNc, implicates a mix of STD and axonal DA release. A dual nature of released DA in the mesencephalon was also previously suggested based on an analysis of the calcium-dependency of DA release in the guinea pig VTA[70], although other work has provided support for a similar calcium-dependency of release in mouse striatum and mesencephalon[71–75,23]. Because the available anatomical data suggests that DA containing axonal varicosities are extremely scarce in the SNc and VTA except in the context of compensatory axonal sprouting associated with partial lesions [76], it would be important to revisit this question in the future with a more thorough anatomical investigation. Such anatomical work would also help to understand the anatomical substrate of the Syt1-dependent DA release detected in the ventral midbrain in the present study and in two recent publications[24,25]. The original optogenetic approach described here in order to trigger STD DA release more selectively should be very useful in the future to further explore the distinct molecular machinery involved in STD and axonal DA release so as to extend recent work showing a role for Syt4 and Syt7 isoforms in this process[11,23].

Considering that electrically evoked DA release in the VTA of Syt1 cKO^{DA} mice was reduced by approximately two-thirds, the near abolition of DA release in the VTA after the adult KO of Syt1 is a puzzling observation. Considering the suboptimal selectivity of the TH-Cre virus used, one obvious possibility is that this could be an artifact due to abrogation of neurotransmitter release from local non-DA neurons that are also stimulated by the extracellular electrodes and that contribute to triggering STD DA release. An approach to resolve this would be to combine this adult KO strategy with the proposed STD optogenetic approach. The use of an inducible DAT-Cre line would also be another possibility to obtain a more selective adult Syt1 KO.

Our results shed new light on the striking resilience of DA-dependent motor functions

at early stages of PD, during which an extensive loss of DA in the striatum gradually occurs

with substantial motor dysfunctions only appearing after more than 70% of terminals in the

striatum are lost[39]. This resilience has been hypothesized to be due to gradually established

compensations in the DA system, including increased DA synthesis in the remaining DA

axons and D2 receptor upregulation, which has been observed in human tissue and in

multiple PD models[77,4,5]. Our work raises the hypothesis that as loss of DA terminals

progresses in PD, even if phasic DA release is greatly impaired, as long as a sufficient tone of

basal extracellular is maintained, many basic motor functions will be preserved. This may

also help explain why even strategies that increase the basal tone of DA without restoring the

normal connectivity of DA neurons, such as the implantation of DA-secreting non-neuronal

cells can help to reduce motor symptoms in PD animal models[78]. In this context, Syt1 cKO^{DA}

mice could represent a model to study compensatory mechanisms and sensitivity in PD.

This model also represents a new tool to study spontaneous release mechanisms and

to tackle the role of phasic DA release in the brain. Previous studies have demonstrated that

although transgenic mice that are hyperdopaminergic or hypodopaminergic can learn various

tasks, their motivation to engage in these tasks is altered. For example, DA-deficient mice

appear to be demotivated and will not engage in some tasks[79], while hyperdopaminergic mice

perform some tasks faster and with less difficulty, suggesting increased motivation[80]. These

results are in line with the well-established role of DA in reward processes, and with its

contribution to motivation for palatable foods ("wanting"), rather than in learning to obtain

this food[81]. In the context of a dramatic reduction in phasic DA release after deletion of Syt1,

we evaluated learning and motivation of Syt1 cKO^{DA} mice in an operant food conditioning

task (**Fig.4**). Although pellets earned from the FR1 and PR1 tasks were their only food

source, body weight remained stable across days (data not shown), demonstrating that mice

satisfied their entire caloric needs with these tasks. Interestingly, we found no global

difference in terms of performance suggesting mice learned both tasks with the same

efficiency and at the same rate (**Fig.4D and G**). In both FR1 and PR1 tasks, we noted a 4-6h

window of enhanced nose-poking rates after the onset of the dark cycle. This period could

coincide with a circadian peak in extracellular DA levels in the striatum[82], and a peak in

activity levels in mice[41]. Interestingly, Ferris et al., by using microdialysis and fast-scan

cyclic voltammetry in rats across the circadian cycle, demonstrated that the amount of

stimulated [DA]o is lowest midway through the dark phase, a time that is characterized by

highest extracellular DA levels. The inverse relationship is exhibited in the light phase, such

that when stimulated [DA]o is highest, extracellular DA is at its lowest point. In the PR1 task,

we also noticed that the nose-poke activity during the first night was significatively higher in

the Syt1 cKO^{DA} mice than in control animals (**Fig. 4F**). Since the number of pellets obtained

was not different between the genotypes, it is possible that the KO mice poke more at the

beginning of the protocol and once they realize the ratio periodically resets, they start acting

thriftier (i.e, fewer pokes to get the same number of pellets). However, poking efficiency and

breaking points were not different between genotypes. Adaptations occurring in Syt1 cKO

mice at the DA transporter and D2 receptor could possibly play a role in this enhancement in

nose poking at the beginning of the dark phases. Ferrer and al. demonstrated that D2

autoreceptor sensitivity is regulated by extracellular DA levels and that the DAT is a critical

governor of diurnal variation in extracellular DA tone. Overall, our results indicate that the

Syt1 conditional deletion and impairment of phasic DA release is dispensable for baseline

motivation to work for food. This result is compatible with observations made in NMDA

cKO^{DA} mice in which phasic DA release was also impaired; in these mice many DA-

dependent behaviors were unaffected[83]. Notably, normal motivation to work for food in a

progressive ratio task was observed[83]. Evaluation of other more complex DA-controlled

functions would be an important next step to document further the roles of phasic DA release.

Our findings are also likely to be relevant to a better understanding of signaling by other

neuromodulators such as serotonin and norepinephrine, which show connectivity patterns that

are similar to that of DA neurons.

Our results are also compatible with previous work obtained in mice in which 90% of

SNc dopaminergic neurons were genetically ablated, leading to a consequent depletion of >

95% of striatal DA [5]. In these mice, motor behaviors of young-adult or aged mice were also

largely unaltered. These mutant mice also exhibited an exaggerated response to L-DOPA

suggesting that preservation of motor functions involves sensitization of striatal DA

receptors. This result is in keeping with our amphetamine-induced locomotion and

autoradiography results, which also point toward the sensitization of DA receptors in $Syt1^{-/-}$

mice. The authors concluded that <5% of striatal dopamine is sufficient to maintain basic

motor functions.

In summary, we conclude that although Syt1 is clearly the main calcium sensor

controlling DA release, a small component of phasic DA release requires another calcium

sensor. We also conclude that Syt1-dependent phasic DA release is dispensable for many

basic unconditioned motor functions that require instead only a basal level of extracellular

DA that could be provided either by (a) spontaneous DA exocytosis, (b) by the small

component of phasic DA release that depends on other calcium sensors, (c) by phasic but

desynchronized exocytosis or even (d) by DAT-mediated reverse transport. Further work will

be required to test these possibilities.

Methods

Detailed protocols for this manuscript are available on the protocols.io repository

(10.17504/protocols.io.n92ldpoj8l5b/v1).

Animal model

The Syt1-floxed mouse line (Syt1$^{lox/lox}$) was obtained from Dr. Schneggenburger who

rederivated the Syt1$^{tm1a(EUCOMM)Wtsi}$ (EMMA, Monterotondo, Italy; EM06829; RRID:

RRID:IMSR_EM:06829) strain with a C57Bl6 background carrying a constitutively

expressing FLP gene in order to remove the Frt-flanked lacZ/neo insert[84]. We then crossed

the Syt1$^{lox/lox}$ mice with B6.SJL-Slc6a3$^{tm1.1(cre)Bkmm}$/J; DATIRES-Cre mice (RRID:IMSR_

JAX:006660), driving expression of the cre recombinase under the control of the DAT

promoter, resulting in the selective deletion of *Syt1* alleles in DA neurons (**Fig. 1A**) (B6.Cg-

Syt1<tm1a(EUCOMM)Wtsi> Slc6a3<tm1.1(cre)Bkmn>/Trudo, RRID:MGI:7488382). Syt1-

floxed/DAT$^{IRES-cre}$ mice were bred from heterozygous crosses to generate Syt1 conditional

knockout in DA neurons (cKO^{DA}), heterozygotes and wild-type animals, referred in the

manuscript as Syt1$^{-/-}$, Syt1$^{+/-}$ and Syt1$^{+/+}$ respectively. Genotyping for Syt1 cKO^{DA} mice was

determined using specific primers to target the *Syt1*-floxed alleles - Syt1^{lox} forward:

GATTCATGATGTCACTGAATCCTATGC   and   Syt1^{lox} reverse

CTGGCAAGTAGCTTAGTGAGTC. Experiments were performed blind with regards to

animal genotype. Male and female were used in this study and their proportion (M/F) is

reported for each experiment. However, the number of mice from each sex did not allow to

separately analyze male and female in statistical analyses. All procedures involving animals

and their care were conducted in accordance with the Guide to care and use of Experimental

Animals of the Canadian Council on Animal Care. The experimental protocols were

approved by the animal ethics committees of the Université de Montréal (protocols 21-113

and 22-108). For microdialysis experiment, the study was approved by the animal care

committee of the research center of the Hôpital du Sacré-Cœur de Montréal (CIUSSS NIM).

Housing was at a constant temperature (21°C) and humidity (60%), under a fixed 12h

light/dark cycle, with food and water available ad libitum.

Virus injections

6-7 week-old cKO^{DA} or Syt1$^{lox/lox}$ mice were anesthetized with isoflurane (Aerrane;

Baxter, Deerfield, IL, USA) and fixed on a stereotaxic frame (Stoelting,Wood Dale, IL,

USA). A small hole was drilled in the exposed skull and a 10 µl Hamilton syringe connected

to a borosilicate injection micropipette pulled using a P-2000 puller (Sutter Instrument)

coupled with a Quintessential Stereotaxic Injector (Stoelting) were used for the injections.

For virally induced Syt1 KO, a AAV9-TH-cre-myc-2A-fusion-red virus (2.1E+13vg/ml) and

AAV9-TH-fusion-red-Sv40 (1.99E+13vg/ml) control virus (kindly provided by Dr. James

Surmeier) were injected bilaterally to Syt1$^{lox/lox}$ mice at the following injection coordinates

[AP (anterior–posterior; ML (medial–lateral); DV (dorsal-ventral), from bregma]: AP -2.8

mm; ML + and – 0.9 mm; DV -4.3 mm and AP -3.2 mm; ML + and – 1.5 mm; DV -4.2 mm

(0.5 µL per sites at a rate of 0.25 µL/min, for a total of 2 µl per mice), in order to infect

neurons in the entire ventral mesencephalon. For optogenetic experiments, a pAAV-syn-

ChR2-eYFP-Kv plasmid was purchased (RRID:Addgene_89256, originally from McLean

Bolton), modified to insert a double-floxed inverse open reading frame (DIO) and packaged

into an AAV2/5 backbone. The AAV2/5-hsyn-DIO-ChR2-eYFP-Kv (6.2 *x10^{12} vg/mL,*

Neurophotonics Core, Québec, CANADA*)* virus or a AAV5-EF1a-DIO-hChR2(H134R)-

eYFP (4.2x10^{12} vg/mL, UNC GTC Vector core, USA, RRID:Addgene_35507) control virus

were injected with the same protocol as above to infect the entire mesencephalon of Syt1$^{+/+}$

and Syt1$^{-/-}$ mice. Each virus was diluted (1:3) with sterile saline solution and kept on ice prior

injections. Animals recovered in their home cage and were closely monitored for 3 days. The

brains were collected for FCSV and immunohistochemistry 3-4 weeks after injections. For

FSCV experiment success of the injections was visually validated each time during the

slicing of the brains by visualizing the presence of the eYFP reporter. Although expression of

the ChR2 constructs was similar in the different mice, small variations in expression level

could have occurred and this represents a limitation of these experiments.

Brain slices preparation

Acute brain slices from 10-12-week-old male or female cKO^{DA} mice were used for all

fast scan cyclic voltammetry (FSCV) recordings. When possible, matched pairs of $Syt1^{+/+}$

and $Syt1^{+/-}/Syt1^{-/-}$ mice were used on each experimental day. The animals were anesthetized

with halothane, decapitated and the brain quickly harvested. Next, the brain was submersed in

ice-cold oxygenated artificial cerebrospinal fluid (aCSF) containing (in mM): NaCl (125),

KCl (2.5), KH_2PO_4 (0.3), $NaHCO_3$ (26), glucose (10), $CaCl_2$ (2.4), $MgSO_4$ (1.3) and coronal

striatal and/or midbrain brain slices of 300 µm thickness were prepared with a Leica

VT1000S vibrating blade microtome. Once sliced, the tissue was transferred to oxygenated

aCSF at room temperature and allowed to recover for at least 1h. For recordings, slices were

placed in a custom-made recording chamber superfused with aCSF at 1 ml/min and

maintained at 32°C with a TC-324B single channel heater controller (Warner Instruments,

USA). All solutions were adjusted at pH 7.35-7.4, 300☐mOsm/kg and saturated with 95%

O_2-5% CO_2 at least 30 min prior to each experiment.

Fast Scan Cyclic Voltammetry

Electrically or optically evoked DA release was measured by FSCV using a 7 µm

diameter carbon-fiber electrode placed into the tissue ∼100 µm below the surface. A bipolar

electrode (Plastics One, Roanoke, VA, USA) or an optical fiber connected to a 470 nm

wavelength LED was placed ∼200 µm away from the recording site. Carbon-fiber electrodes

(Goodfellow Cambridge Limited, UK) of 7☐µm in diameter were aspirated into ethanol-

cleaned glass capillaries (1.2☐mm O.D., 0.68☐mm I.D., 4 inches long; World Precision

Instruments, FL, USA). The glass capillaries were then pulled using a P-2000 micropipette

puller (Sutter Instruments, Novato, USA), dipped into 90°C epoxy for 30s (Epo-Tek 301,

Epoxy Technology, MASS, USA) and cleaned in hot acetone for 3s. The electrodes were

heated at 100°C for 12h and 150°C for 5 days. Electrodes were then polished and filled with

potassium acetate at 4M and potassium chloride at 150 mM. The protruding carbon fibers

were cut using a scalpel blade under direct visualization to a length allowing to obtain

maximal basal currents of 100 to 180 nA. The electrodes were calibrated with 1 µM DA in

aCSF before and after each recorded slice and the mean of the current values obtained were

used to determine the amount of released DA. After use, electrodes were cleaned with

isopropyl alcohol (Bioshop, France). The potential of the carbon fiber electrode was scanned

at a rate of 300 V/s according to a 10 ms triangular voltage wave (–400 to 1000 mV vs

Ag/AgCl) with a 100 ms sampling interval, using a CV 203BU headstage preamplifier

(Molecular Devices) and a Axopatch 200B amplifier (Molecular Devices, USA). Data were

acquired using a Digidata 1440a analog to digital converter board (Molecular Devices, USA)

connected to a computer using Clampex and Clampfit (version 10.7, Molecular Devices,

USA , pClamp RRID:SCR_011323). Slices were left to stabilize for 20 min before any

electrochemical recordings. After positioning of the bipolar stimulation electrode or the

optical probe and carbon fiber electrodes in the tissue, single pulses (400 µA or 30 mW,

1ms,) or pulses-train (30 pulses at 10 Hz) were applied to the tissue to trigger DA release. For

recordings in the VTA and SNc, it is theoretically possible that FSCV signals were partly

contaminated by the presence of 5-HT, as previous studies showed that this biogenic amine is

present in these regions and can be detected together with DA[85,86]. However, the shape of our

cyclic voltammograms suggests that such contamination was minimal or inexistant.

Behavioral analyses

For the behavioral experiments, different cohorts of mice were used for experiments

involving the administration of pharmacological agents. Only mice used for the rotarod and

grip strength tests were from the same cohort.

Grip strength test. A grip strength apparatus (BioSeb instruments, BIO-GS3, France)

was used to evaluate the paw force developed by Syt1 cKO^{DA} mice. 10-12-week-old male or

female mice were held firmly by the tail and slowly lowered until their forepaws grasped the

middle of the grid. Subjects were then lowered to a horizontal position and pulled following

the axis of the sensor until they released their grasp on the grid. The maximum force

exhibited by the subject was recorded. The test was repeated 3 times with 30 min resting

period between each trial. The results are presented as the means of the 3 trials divided by the

body weight of each recorded mouse.

Pole test. The test was conducted with a homemade 48-cm metal rod of 1-cm

diameter covered with adhesive tape to facilitate traction, placed in a cage. 10-12-week-old

male or female Syt1 cKO^{DA} were positioned head-up at the top of the pole and the time

required to turn (t-turn) and climb down completely was recorded. The results are presented

as the mean of 2 sessions, with 3 trial per sessions for each mouse.

Rotarod. Motor coordination was evaluated with a rotarod apparatus (Harvard

Apparatus, LE8205, USA). 10-12-week-old male or female Syt1 cKO^{DA} mice were pre-

trained on the rotarod for two consecutive days to reach a stable performance and then tested

on day three. On the first day, mice were placed on the rotarod, rotating at a constant speed of

4 rpm. They remained on the rotarod until either one min had passed, or up to a maximum of

3 attempts at placing them on the rod. On the second day, mice were trained on the device

with an accelerated rotation of 4 to 40 rpm, over a ten-min period. This training was repeated

3 times with approximately 30 min of rest between each trial. Mice were tested on the

accelerating rod on day 3 and the latency to fall the device of each tested animal was

recorded. The results are presented as the means of the 3 trials on day 3.

Open field. The locomotor behavior of 11-12-week-old male or female Syt1 cKO[DA]

mice was recorded using an infrared actimeter (Superflex sensor version 4.6, Omnitech)

using the Fusion software (v5.6 Superflex Edition, RRID:SCR_017972). A chamber partition

was used to measure two mice at a time. Subjects were not given time to acclimate and spent

a total of 60 min in the chamber with the following protocol: the first 20 min was used to

record basal locomotion, while the following 40 min was used to record locomotion after

saline or drug administration (i.p.). Drug treatments correspond to a single dose of cocaine

hydrochloride at 20 mg/kg (Medisca, cat# 53-21-4, Canada) or D-amphetamine sulfate at 5

mg/kg (Tocris, 2813, UK) at doses known to increase locomotion[87,88]. SCH23390-HCl

(Sigma, D-054, Canada) *at* 50 µg/kg, quinpirole-HCl at 0.2 mg/kg (Sigma, Q-102, Canada)

and raclopride L-tartrate at 1 mg/kg (Sigma, R-121, Canada) were also used at selected high

doses known to reduce locomotion in the open field [89–91]. Each drug was diluted into 0.9%

sodium chloride saline solution (Halyard, #cat 116). For each drug tested, a different cohort

of mice was used. Results are presented as the mean of the traveled distance.

Operant learning. Appetitively motivated operant learning was assessed using the

open source Feeding Experimentation Device 3 (FED)[41]. Individually housed animals were

placed in a 12h/12h light/dark cycle. First, the feeding devices were placed in the home cages

under a free feeding paradigm, which automatically dispenses a new pellet each time the

mouse removes the pellet in the well. This magazine training was performed for 2 days prior

to changing the device mode for a fixed ratio 1 (FR1) paradigm, in which mice learned to

nose-poke on the active port to receive a single pellet. To evaluate the operant learning

capacity of the mice, nose-poking for pellets across the circadian cycle and nose-poking

efficiency (% of total pokes on the active port) were measured by running mice on the FR1

task for 3 consecutive days. After completion of the FR1 task, the device protocol was

changed for a 6-days repeating progressive ratio (PR1) task, in which the nose-poking

requirement began on FR1 and increased by 1 poke each time a pellet was earned. When

mice refrained from poking on either active or inactive port for 30 min, the ratio was reset to

FR1. The nose-poking rate and poke efficiency were once again measured during the PR1

task to evaluate the motivation of mice for food across the circadian cycle.

Dopamine receptor autoradiography (saturation curves)

10-12-week-old male or female Syt1 cKO^{DA} mice were anesthetized using

pentobarbital NaCl solution (7 mg/mL) injected intraperitoneally and then were perfused

intracardially with 20ml of PBS to remove the blood. The brains were extracted and quickly

dipped in isopentane at -30°C for 1 min. 12-microns thick coronal sections of the whole

striatum were then cut using a cryostat (Leica Biosystem, Model 3050) and mounted on

charged microscope slides (X-tra, Leica, Canada). For binding assays, after a pre-wash step

in Tris-HCl buffer (15 min) at room temperature, incubation was performed for 1h at room

temperature in a 50 mM Tris-HCl buffer (pH 7.4), containing 120 mM NaC1, 5 mM KC1, 2

mM CaC1$_2$, 1 mM MgC1$_2$, in the presence of 7 increasing concentrations (0.15 – 10 nM) of

[^{3}H]-raclopride (specific activity 82.8Ci/mmol, Perkin Elmer, #NET975250UC, Canada) for

D2 binding, 7 increasing concentrations (0.15 – 10 nM) of [^{3}H]-SCH23390 (specific activity

83.6Ci/mmol, Perkin Elmer, #NET930025UC, Canada) for D1 binding to generate saturation

curves. To determine the non-specific binding, adjacent sections were incubated respectively

with 5 µM of sulpiride (Sigma, cat#S7771, Canada) for D2 binding and 20 nM mianserin

hydrochloride (Sigma, cat#M2525, Canada) with 1 µM SCH-39166 hydrobromide (Tocris,

cat#2299, Canada) for D1 binding for each radioligand concentrations used. Incubation was

terminated by rinsing sections twice for 5 min in ice-cold Tris-HC1, 50 mM (pH 7.4).

Sections were then dipped briefly in cold distilled water and dried overnight. The microscope

slides with tissue sections were placed in a light-proof X-ray cassette. Autoradiographs were

prepared by placing the slide-mounted tissue sections in contact with a tritium-sensitive film

(Biomax MR, Kodack, cat#8715187, Canada) for 6 weeks at room temperature. The films

were then developed, and autoradiograms analyzed by densitometry. Autoradiograms were

acquired using a grayscale digital camera (Model CFW-1612M, Scion Corporation,

Maryland, USA). The precise topography of D1 and D2 receptors in the striatum was

determined from autoradiograms of 4 serial coronal sections for each radioligand

concentration corresponding to approximately bregma +1.34 mm to 0.38 mm of the Paxinos

and Watson mouse brain atlas[92]. In each section, the caudate-putamen was arbitrarily divided

into 4 quadrants (dorso-lateral, ventro-lateral, dorso-medial and ventro-medial). For each

quadrant, the mean signal intensities were measured on ImageJ software

(ImageJ 1.53v software, Wayne Rasband, NIH, RRID:SCR_003070) and non-specific

binding was subtracted from all density readings. The corrected optical gray densities of the

different brain areas were converted into nCi/mg of tissue equivalent (nCi/mg TE) using a

[^{3}H] radioactive standard disposed on each film (American Radiolabeled Chemicals Inc.,

Cat# ART 0123A). The saturation binding curves were then generated and analyzed on

GraphPad software (GraphPad Prism, V9, RRID:SCR_002798) using a non-linear regression

with the equation $Y = Bmax*X/(Kd + X)$ (one-site binding model), where Y represents the

specific binding values (nCi/mg of tissue equivalent) and X the radioligand concentration to

estimate Bmax, the maximum specific binding and Kd, the dissociation constant at

equilibrium. The two last parameters were extracted and represent respectively the receptor

density and the radioligands affinity for D1 or D2 receptors.

Tissue preparation and immunohistochemistry

10-12-week-old male or female Syt1 cKO^{DA} mice were anesthetized using a

pentobarbital NaCl solution (7 mg/mL) injected intraperitoneally and then were perfused

intracardially with 20 mL of PBS followed by 30 mL of paraformaldehyde (PFA) 4%. The

brains were extracted, placed 48h in PFA followed by 48h in a 30% sucrose solution and

frozen in isopentane at -30°C for 1 min. 40 microns thick coronal sections were then cut

using a cryostat (Leica CM1800) and placed in antifreeze solution at -20°C until used. For

slice immunostaining, after a PBS wash, the tissue was permeabilized, nonspecific binding

sites were blocked and slices were incubated overnight with a rabbit anti-tyrosine

hydroxylase (TH) (1:1000, Millipore Cat# AB152, RRID:AB_390204), a rat anti-DAT

(1:2000, Millipore Cat# MAB369, RRID:AB_2190413), a rabbit anti-VMAT2 (1:2000,

kindly provided by Dr. G.W. Miller, Emory University, USA[93]), a rabbit anti-5-HT (1:2000,

ImmunoStar Cat# 20080, RRID:AB_572263), a chicken anti-GFP (1:1000, Aves Labs Cat#

GFP-1020, RRID:AB_10000240) and/or a rabbit anti-RFP (1:1000, Rockland Cat# 600-401-

379, RRID:AB_2209751) antibody. Primary antibodies were subsequently detected with goat

anti-rabbit, rat or chicken Alexa Fluor-488–conjugated (A-11008 RRID:AB_143165, A-

11006 RRID:AB_2534074, A-11039 RRID:AB_2534096), 546-conjugated (A-11010

RRID:AB_2534077, A-11081 RRID:AB_2534125, A-11040 RRID:AB_2534097) and/or

647-conjugated secondary antibodies (A-21244 RRID: AB_2535812, A-21247

RRID:AB_141778, A-21449 RRID:AB_2535866) (1:500, 2h incubation; Invitrogen,

Canada). Slices were mounted on charged microscope slides (Superfrost/Plus, Fisher

Scientific, Canada) and stored at 4°C prior to image acquisition.

Confocal imaging

Images were acquired using an Olympus Fluoview FV1000 point-scanning confocal

microscope (Olympus FV1000 Confocal Microscope, RRID:SCR_020337) with a 60x oil-

immersion objective (NA 1.42). Images acquired using 488nm, 546 nm and 647 laser

excitations were scanned sequentially to prevent non-specific bleed-through signal. Care was

taken to avoid sensor saturation in all images, thus allowing quantification of signal intensity.

A background correction was first applied at the same level for every image analyzed before

any quantification. This background was determined from the level of signals in striosomes,

regions that contain very low levels of DA neuron markers. An ImageJ (1.53v, National

Institutes of Health, Fiji RRID:SCR_002285) macro developed in-house was used to perform

all quantifications (10.5281/zenodo.8007760). For the immunohistochemical characterization

of brain tissue obtained from Syt1 cKO^{DA} mice or virally induced Syt1 KO mice, grayscale

images were converted to binary (a.k.a. halftone or black & white) by defining a grayscale

cut-off point. Grayscale values below the cut-off became black and those above became

white. The signal contrast/intensity threshold of each image within the same IHC was

adjusted and the relative results (% of WT) were reported to account for signal variability due

to pooling results from different sets of immunostainings. Then the area covered by the signal

and the intensity for each signal were measured with ImageJ in a series of 5 different striatal

sections ranging from bregma +0.98 to bregma -1.06 mm, with a total of 22 different spots

for each hemisphere (**Fig. S4A**).

Stereology

One out of every 6th cryostat section was used for TH immunostaining stereological

counting of DA neurons. After a PBS wash, the tissue was incubated for 10 min with 0.9%

H_2O_2 solution, then washed with PBS again and incubated for 48h with a rabbit anti-TH

antibody (1:1000, Millipore Cat# AB152, RRID:AB_390204) at 4°C, 12h with goat anti-

rabbit biotin-SP-AffiniPure secondary antibody (1:100, Jackson ImmunoResearch Labs Cat#

111-065-003, RRID:AB_2337959) at 4°C and 3h with horseradish peroxidase streptavidin

(1:200, Jackson ImmunoResearch Labs Cat# 016-030-084, RRID:AB_2337238). The

diaminobenzidine (DAB) reaction was carried out for 45s, then stopped by incubation with

0.1M acetate buffer. Slices were mounted on charged microscope slides (Superfrost/Plus,

Fisher Scientific, Canada) and left to dry for 96h after which they were stained with cresyl

violet and went through incubations with increasing concentrations of alcohol. After short

isopropanol and xylene baths, slides were sealed with Permount mounting medium (SP15-

100, Fisher, USA) using glass coverslips. TH-immunoreactive neurons were counted in one

out of every sixth section using a 100x oil-immersion objective on a Leica microscope

equipped with a motorized stage. A 60 x 60 μm^2 counting frame was used in the Stereo

Investigator (MBF Bioscience, RRID:SCR_002526) sampling software with a 12 µm optical

dissector (2 µm guard zones) and counting site intervals of 150 µm after a random start (100

µm intervals for unilateral lesion). Mesencephalic DA nuclei, including the VTA, SNc and

RRF were examined. Stereological estimates of the total number of TH-immunoreactive

neurons within each nucleus were obtained. The number of TH-negative neurons was also

estimated similarly in each region based on cresyl violet staining.

Primary neuronal co-cultures

For all experiments, postnatal day 0-3 (P0-P3) mice were cryoanesthetized, decapited

and used for co-cultures according to a previously described protocol[57]. Primary DA neurons

from VTA or SNc were prepared from $Syt1^{-/-}$ or $Syt1^{+/+}$ pups and co-cultured with ventral

striatum and dorsal striatum neurons from $Syt1^{-/-}$ or $Syt1^{+/+}$ pups, respectively. Neurons were

seeded (60 000 cells/mL) on a monolayer of cortical astrocytes grown on collagen/poly-L-

lysine-coated glass coverslips. All cultures were incubated at 37°C in 5% CO_2 and

maintained in 2/3 of Neurobasal, enriched with 1% penicillin/streptomycin, 1% Glutamax,

2% B-27 supplement and 5% fetal bovine serum (Invitrogen, Canada) plus 1/3 of minimum

essential medium enriched with 1% penicillin/streptomycin, 1% Glutamax, 20mM glucose,

1mM sodium pyruvate and 100 µl of MITO+ serum extender. All primary neuronal co-

cultures were used at 14 days *in vitro* (DIV).

Immunocytochemistry on cell cultures

Cultures were fixed at 14-DIV with 4% paraformaldehyde (PFA; in PBS, pH-7.4),

permeabilized with 0,1% triton X-100 during 20 min, and nonspecific binding sites were

blocked with 10% bovine serum albumin during 10 min. Primary antibodies were: mouse

anti-TH (1:2000, Sigma-Aldrich Cat# MAB318, RRID:AB_2201528), rabbit anti-TH

(1:2000, Millipore Cat# AB152, RRID:AB_390204), rabbit anti-Syt1 (1:1000, Synaptic

Systems Cat# 105 103, RRID:AB_11042457), rabbit anti-VMAT2 (1:2000, gift of Dr. Gary

Miller, Colombia University) and mouse anti-MAP2 (1:2000, Millipore Cat# MAB3418,

RRID:AB_94856). These were subsequently detected using goat anti mouse or rabbit Alexa

Fluor-488-conjugated (A-11001 RRID:AB_2534069, A-11008, RRID:AB_143165), Alexa

Fluor-546-conjugated (A-11003 RRID:AB_2534071, A-11010 RRID:AB_2534077), Alexa

Fluor-568-conjugated (A-11031 RRID:AB_144696, A-21124 RRID:AB_2535766) and

Alexa Fluor-647-conjugated (A-21235 RRID:AB_2535804, A-21244 RRID:AB_2535812)

secondary antibodies (1:500, Invitrogen, Canada).

RT-qPCR

We used RT-qPCR to quantify the amount of mRNA encoding Syt1, 4, 7 and 11,

Doc2b, TH, DAT and VMAT2 in brain tissue from P70 $Syt1^{+/+}$ and $Syt1^{-/-}$ mice. The brains

were quickly harvested and the ventral mesencephalon containing SN/VTA structures were

microdissected and homogenized in 500 µL of trizol. Next, RNA extraction was performed

using RNAeasy Mini Kit (Qiagen, Canada) according to the manufacturer's instructions.

RNA integrity was validated using a Bioanalyzer 2100 (Agilent, RRID:SCR_019715). Total

RNA was treated with DNase and reverse transcribed using the Maxima First Strand cDNA

synthesis kit with ds DNase (Thermo Fisher Scientific). Gene expression was determined

using assays designed with the Universal Probe Library from Roche

(www.universalprobelibrary.com). For each qPCR assay, a standard curve was performed to

ensure that the efficiency of the assay was between 90% and 110%. Assay information is

presented in **supplementary Table 1**. The QuantStudio qPCR instrument (Thermo Fisher

Scientific, RRID:SCR_023003) was used to detect the amplification level. Relative

expression comparison (RQ = 2^-ΔΔCT) was calculated using the Expression Suite software

(Thermo Fisher Scientific), using GAPDH as an endogenous control. Efficiency correction

was performed to mathematically compensate for differences in amplification efficiency of

the targets and endogenous controls when calculating relative quantities.

Western Blot

Striatum samples microdissected from $Sy1^{+/+}$, $Syt1^{+/-}$, $Syt1^{-/-}$ adult mice were lysed in

RIPA buffer (Fisher Scientific, PI89900) containing a protease inhibitor cocktail (Sigma).

Homogenized tissue samples were centrifuged at 12000g for 30 min at 4°C. Supernatant was

collected and protein quantification was done with BCA reagent (Thermo Scientific Pierce

BCA Protein Assay Kit, PI23227). 20 µg of each sample was separated on 8% SDS-PAGE

followed by transfer onto a nitrocellulose membrane. Membrane blocking was done with

10% skimmed milk for 90 min at RT with gentle shaking. The membranes were incubated

overnight at 4°C with gentle shaking with rat anti-DAT (1:1000, Millipore Cat# MAB369,

RRID:AB_2190413), rabbit anti-TH (1:1000, Millipore Cat# AB152, RRID:AB_390204),

rabbit anti-VMAT2 (1:5000, kindly provided by Dr. G.W. Miller, Emory University, USA[93])

and mouse anti-β-actin-HRP (1:5000, Sigma-Aldrich Cat# A3854, RRID:AB_262011)

primary antibodies. Membranes were washed 5 times with TBST buffer for 5 min each time.

After this, appropriate secondary antibodies (1:5000) were added and the incubation was

done at RT for 90 min with gentle shaking. Membranes were washed again with TBST buffer

for 5 times X 5 min and developed using Clarity Western ECL substrate (Bio-Rad,

1004384863). Images were captured on a Luminescent Image Analyzer (GE Healthcare)

using Image quant LAS 4000 software (RRID:SCR_018374). Membranes were stripped and

re-probed for β-actin as a loading control. The unprocessed scans of the blots are provided in

the supplementary information file.

1119 Microdialysis guide cannula implantation

10-12 weeks old male and female Syt1 cKO^{DA} mice were anesthetized with sodium

isoflurane (2.5% isoflurane at 0.5 L/min oxygen flow), coupled with infiltration analgesia of

1.5 mg/kg lidocaine/bupivacaine 10 min prior to incision of the skull and stereotaxic

implantation with a microdialysis guide cannula (CMA 7, Harvard Apparatus) into the left

dorsal subdivision of the striatum (coordinates: 1.0 mm anterior of bregma, 2.0 mm lateral of

bregma, and -3.3 mm below pia) and the right substantia nigra and ventral tegmental area

(coordinates: -3.3 mm anterior of bregma, -0.8 mm lateral of bregma, and -3.4 mm below pia)

following Paxinos coordinates[92]. These guide cannulas were then used to insert the

microdialysis probes (6 kDa MW cut-off, CMA 7, Harvard Apparatus) into the target sites.

Guide cannulas were secured with acrylic dental cement and an anchor screw was threaded

into the cranium. Buprenorphine (0.05 mg/kg, subcutaneously) was used for postoperative

analgesia (once daily for 2 days). Animals were allowed 1-week (housed one per cage) to

recover from cannula implantation before dialysis measurements of extracellular DA, 3,4-

dihydroxyphenylacetic acid (DOPAC), norepinephrine (NE), and serotonin (5-HT) levels by

microdialysis. A removeable obturator was inserted into the cannula to prevent cerebrospinal

fluid seepage and infection.

1136 Microdialysis

Microdialysis probes were calibrated in aCSF containing (in mM): NaCl (126), KCl

(3), NaHCO$_3$ (26), NaH$_2$PO$_4$ (3), MgCl$_2$ (1.3), CaCl$_2$ (2.3), and L-ascorbic acid (0.2). *In vitro*

probe recovery ranged from 4% to 30% at a flow rate of 1 µl/min. A computer-controlled

microinfusion pump *(CMA)* was used to deliver perfusate to the probes, and the dialysate was

collected from the outlet line. Probes were inserted into the indwelling guide cannulas of the

anesthetized animals and perfused with aCSF (flow rate set at 1 µl/min). To minimize the

influence of needle trauma on experimental outcomes, dialysates samples were collected

during a typical 60 min equilibration period but were discarded and not analyzed. Three

samples were then taken at 20 min intervals for 60 min. Each 10 µl dialysate sample was

collected in a fraction vial preloaded with 1 µl of 0.25 mol/l perchloric acid to prevent

analyte degradation and immediately stored at 4 °C for subsequent analysis. Following

microdialysis, mice received an intraperitoneal injection of ketamine/xylazine (120 mg/10

mg/kg) and were intracardially injected with cold saline, decapitated and the brains were

harvested. After the microdialysis experiments, the striatum of each mouse was

microdissected, and flash frozen in isopentane cooled to -35 °C on dry ice and stored at -80

°C prior to HPLC quantification.

High performance liquid chromatography

Extracellular DA, DOPAC, 5-HT and NE concentrations in the left dorsal subdivision

of the striatum and the right substantia nigra (SNc) and ventral tegmental area (VTA) were

determined with high-performance liquid chromatography (HPLC) using a Dionex pump

(Ultimate 3000) coupled with electrochemical detection (EC)[94]. Samples were run through a

Luna C18 (2) 75 mm × 4.6 mm 3 µm analytical column at a flow rate of 1.5 ml/min and

the electrochemical detector (ESA CoulArray, model #5600A) was set at a potential of -250

mV and +300 mV. The mobile phase consisted of 6% methanol, 0.341 mM 1-

octanesulfoic acid sodium salt, 168.2 mM sodium acetate, 66.6 mM citric acid

monohydrate, 0.025 mM ethylenediamine-tetra-acetic acid disodium and 0.71 mM

triethylamine adjusted to pH 4.0–4.1 with acetic acid. Using ESA's CoulArray v3.0 software,

the position of the peaks for each metabolite was compared to an external standard solution

containing 25 ng/ml DA, DOPAC, 5-HT, NE and 50 mM acetic acid prepared fresh daily

from stock solutions and loaded with samples into a refrigerated (10 °C) Dionex RS

autosampler (Ultimate 3000). Under these conditions, the retention time for DOPAC, NE,

DA, and 5-HT was approximately 2.5 min, 1.1 min, 2.8 min, and 3.76 min respectively, with

a total run time of 22 min/sample. Chromatographic peak analysis was accomplished by

identification of unknown peaks in a sample matched according to retention times from

known standards using Chromeleon Chromatography Data System (CDS) Software

(RRID:SCR_016874). The mean standard (n = 5) was used to quantify unknown peaks and to

have the same retention time on each run. The analytical curve, as determined with duplicate

standard solutions of DOPAC, NE, DA and 5-HT in nanopure water, was in the range of 7 to

500 ng/ml (R^2 = 0.9963, 0.9971, 0.9899 or 0.9775, respectively). The analytical curve was

constructed by plotting the area under the curve. For tissular measures, microdissected

striatum were homogenized in a solution which contained 45 µl of 0.25 M perchlorate and

15 µl of 2,3-dihydroxybenzoic acid (100 mg/ml) which served as an internal standard.

Following centrifugation at 10,000 rpm for 15 min at 4 °C, the supernatant was isolated to

detect DA, DOPAC, NE and 5-HT using HPLC-EC as previously described. In parallel,

pellets were reconstituted in 50 µl of 0.1 N NaOH and kept for protein quantification using

a BCA protein assay kit (Thermo Scientific Pierce BCA Protein Assay Kit, PI23227).

Dialysate concentrations are expressed as ng/ml and tissular concentrations as ng/mg of total

proteins.

Statistical analysis

Statistical analyses were performed with Prism 9 software (GraphPad Prism, V9,

RRID:SCR_002798, $p < 0.05$ = *, $p < 0.01$ = **, $p < 0.001$ = ***, $p < 0.0001$ = ****).

Parametric statistical tests were used because samples contained data with normal

distributions. Data are presented as mean ± SEM. The level of statistical significance was

established at $p < 0.05$ in one or two-way ANOVAs with Brown-Forsythe or Geiser-

Greenhouse correction when needed or two-tailed t-tests with Welch's correction when

needed. A ROUT outlier analysis was performed when required (Q = 1%). Post-hoc tests

were only performed when the global ANOVA was significant. The Tukey post-hoc test was

used when all the means were compared to each other and the Sidak post-hoc test was used

when only subsets of means were compared.

1196

Data availability statement

All key data is included in the manuscript and in the source data file. Source data is provided with paper. Access to original raw images and electrochemistry recording traces is available on request from the corresponding author.

References

1. Schultz, W. Multiple Dopamine Functions at Different Time Courses. *Annu. Rev. Neurosci.* **30**, 259–288 (2007).

2. Surmeier, D. J., Graves, S. M. & Shen, W. Dopaminergic modulation of striatal networks in health and Parkinson's disease. *Curr Opin Neurobiol* **29**, 109–117 (2014).

3. Hauber, W. Impairments of movement initiation and execution induced by a blockade of dopamine D1 or D2 receptors are reversed by a blockade of N-methyl-D-aspartate receptors. *Neuroscience* **73**, 121–130 (1996).

4. Mercuri, N. B. & Bernardi, G. The 'magic' of l-dopa: why is it the gold standard Parkinson's disease therapy? *Trends in Pharmacological Sciences* **26**, 341–344 (2005).

5. Golden, J. P. *et al.* Dopamine-dependent compensation maintains motor behavior in mice with developmental ablation of dopaminergic neurons. *J. Neurosci.* **33**, 17095–17107 (2013).

6. Descarries, L., Watkins, K. C., Garcia, S., Bosler, O. & Doucet, G. Dual character, asynaptic and synaptic, of the dopamine innervation in adult rat neostriatum: a quantitative autoradiographic and immunocytochemical analysis. *J. Comp. Neurol.* **375**, 167–186 (1996).

7. Antonopoulos, J., Dori, I., Dinopoulos, A., Chiotelli, M. & Parnavelas, J. G. Postnatal development of the dopaminergic system of the striatum in the rat. *Neuroscience* **110**, 245–256 (2002).

8. Descarries, L. *et al.* Glutamate in dopamine neurons: Synaptic versus diffuse transmission. *Brain Research Reviews* **58**, 290–302 (2008).

9. Ducrot, C. *et al.* Dopaminergic neurons establish a distinctive axonal arbor with a majority of non-synaptic terminals. *The FASEB Journal* **35**, e21791 (2021).

10. Rice, M. E. & Patel, J. C. Somatodendritic dopamine release: recent mechanistic insights. *Philos.*

Trans. R. Soc. Lond., B, Biol. Sci. **370**, (2015).

11. Mendez, J. A., Bourque, M.-J., Fasano, C., Kortleven, C. & Trudeau, L.-E. Somatodendritic

Dopamine Release Requires Synaptotagmin 4 and 7 and the Participation of Voltage-gated

Calcium Channels. *J. Biol. Chem.* **286**, 23928–23937 (2011).

12. Liu, C., Kershberg, L., Wang, J., Schneeberger, S. & Kaeser, P. S. Dopamine Secretion Is

Mediated by Sparse Active Zone-like Release Sites. *Cell* **172**, 706-718.e15 (2018).

13. Banerjee, A. *et al.* Molecular and functional architecture of striatal dopamine release sites.

bioRxiv 2020.11.25.398255 (2020) doi:10.1101/2020.11.25.398255.

14. Liu, C., Goel, P. & Kaeser, P. S. Spatial and temporal scales of dopamine transmission. *Nature*

Reviews Neuroscience 1–14 (2021) doi:10.1038/s41583-021-00455-7.

15. Fortin, G. D., Desrosiers, C. C., Yamaguchi, N. & Trudeau, L. E. Basal somatodendritic

dopamine release requires snare proteins. *J Neurochem* **96**, 1740–1749 (2006).

16. Bergquist, F., Niazi, H. S. & Nissbrandt, H. Evidence for different exocytosis pathways in

dendritic and terminal dopamine release in vivo. *Brain Res.* **950**, 245–253 (2002).

17. Ovsepian, S. V. & Dolly, J. O. Dendritic SNAREs add a new twist to the old neuron theory.

PNAS **108**, 19113–19120 (2011).

18. Hikima, T. *et al.* Activity-dependent somatodendritic dopamine release in the substantia nigra

autoinhibits the releasing neuron. *Cell Rep* **35**, 108951 (2021).

19. Sudhof, T. C. The synaptic vesicle cycle. *Annu. Rev. Neurosci.* **27**, 509–547 (2004).

20. Andrews, N. W. & Chakrabarti, S. There's more to life than neurotransmission: the regulation of

exocytosis by synaptotagmin VII. *Trends Cell Biol* **15**, 626–631 (2005).

21. Xu, J., Mashimo, T. & Südhof, T. C. Synaptotagmin-1, -2, and -9: Ca(2+) sensors for fast release

that specify distinct presynaptic properties in subsets of neurons. *Neuron* **54**, 567–581 (2007).

22. Banerjee, A., Lee, J., Nemcova, P., Liu, C. & Kaeser, P. S. Synaptotagmin-1 is the Ca2+ sensor

for fast striatal dopamine release. *eLife* **9**, e58359 (2020).

23. Delignat-Lavaud, B., Ducrot, C., Kouwenhoven, W., Feller, N. & Trudeau, L.-É. Implication of

synaptotagmins 4 and 7 in activity-dependent somatodendritic dopamine release. *bioRxiv*

2021.01.25.427983 (2021) doi:10.1101/2021.01.25.427983.

24. Hikima, T., Witkovsky, P., Khatri, L., Chao, M. V. & Rice, M. E. Synaptotagmins 1 and 7 Play

Complementary Roles in Somatodendritic Dopamine Release. *J Neurosci* **42**, 3919–3930 (2022).

25. Lebowitz, J. J. *et al.* Synaptotagmin-1 is a Ca2+ sensor for somatodendritic dopamine release.

Cell Rep **42**, 111915 (2023).

26. Robinson, B. G. *et al.* RIM is essential for stimulated but not spontaneous somatodendritic

dopamine release in the midbrain. *eLife* **8**, e47972 (2019).

27. Deutch, A. Y., Goldstein, M., Baldino, F. & Roth, R. H. Telencephalic projections of the A8

dopamine cell group. *Ann. N. Y. Acad. Sci.* **537**, 27–50 (1988).

28. Bayer, V. E. & Pickel, V. M. Ultrastructural localization of tyrosine hydroxylase in the rat ventral

tegmental area: relationship between immunolabeling density and neuronal associations. *J.

Neurosci.* **10**, 2996–3013 (1990).

29. Juraska, J. M., Wilson, C. J. & Groves, P. M. The substantia nigra of the rat: a Golgi study. *J.

Comp. Neurol.* **172**, 585–600 (1977).

30. Wassef, M., Berod, A. & Sotelo, C. Dopaminergic dendrites in the pars reticulata of the rat

substantia nigra and their striatal input. combined immunocytochemical localization of tyrosine

hydroxylase and anterograde degeneration. *Neuroscience* **6**, 2125–2139 (1981).

31. Matsuda, W. *et al.* Single Nigrostriatal Dopaminergic Neurons Form Widely Spread and Highly

Dense Axonal Arborizations in the Neostriatum. *J. Neurosci.* **29**, 444–453 (2009).

32. Lim, S. T., Antonucci, D. E., Scannevin, R. H. & Trimmer, J. S. A novel targeting signal for

proximal clustering of the Kv2.1 K+ channel in hippocampal neurons. *Neuron* **25**, 385–397

(2000).

33. Misonou, H., Mohapatra, D. P. & Trimmer, J. S. Kv2.1: A Voltage-Gated K+ Channel Critical to

Dynamic Control of Neuronal Excitability. *NeuroToxicology* **26**, 743–752 (2005).

34. Jensen, C. S. *et al.* Trafficking of Kv2.1 Channels to the Axon Initial Segment by a Novel

Nonconventional Secretory Pathway. *J Neurosci* **37**, 11523–11536 (2017).

35. Baker, C. A., Elyada, Y. M., Parra, A. & Bolton, M. M. Cellular resolution circuit mapping with

temporal-focused excitation of soma-targeted channelrhodopsin. *eLife* **5**, e14193.

36. O'Neill, B., Patel, J. C. & Rice, M. E. Characterization of Optically and Electrically Evoked

Dopamine Release in Striatal Slices from Digenic Knock-in Mice with DAT-Driven Expression

of Channelrhodopsin. *ACS Chem Neurosci* **8**, 310–319 (2017).

37. Bergquist, F., Shahabi, H. N. & Nissbrandt, H. Somatodendritic dopamine release in rat

substantia nigra influences motor performance on the accelerating rod. *Brain Research* **973**, 81–

91 (2003).

38. Joshua, M., Adler, A. & Bergman, H. The dynamics of dopamine in control of motor behavior.

Current Opinion in Neurobiology **19**, 615–620 (2009).

39. Halliday, G. M. *et al.* Neuropathology of immunohistochemically identified brainstem neurons in

Parkinson's disease. *Annals of Neurology* **27**, 373–385 (1990).

40. Ward, K. M. & Citrome, L. Antipsychotic-Related Movement Disorders: Drug-Induced

Parkinsonism vs. Tardive Dyskinesia—Key Differences in Pathophysiology and Clinical

Management. *Neurol Ther* **7**, 233–248 (2018).

41. Matikainen-Ankney, B. A. *et al.* An open-source device for measuring food intake and operant

behavior in rodent home-cages. *eLife* **10**, e66173 (2021).

42. Bauman, R. An experimental analysis of the cost of food in a closed economy. *J Exp Anal Behav*

56, 33–50 (1991).

43. Chaney, M. A. & Rowland, N. E. Food demand functions in mice. *Appetite* **51**, 669–675 (2008).

44. Beeler, J. A., Daw, N., Frazier, C. R. M. & Zhuang, X. Tonic Dopamine Modulates Exploitation

of Reward Learning. *Front Behav Neurosci* **4**, 170 (2010).

45. Mourra, D., Gnazzo, F., Cobos, S. & Beeler, J. A. Striatal Dopamine D2 Receptors Regulate Cost

Sensitivity and Behavioral Thrift. *Neuroscience* **425**, 134–145 (2020).

46. Donnan, G. A. *et al.* Evidence for plasticity of the dopaminergic system in parkinsonism. *Mol*

Neurobiol **5**, 421–433 (1991).

47. Richard, M. G. & Bennett, J. P. Regulation by D2 dopamine receptors of in vivo dopamine

synthesis in striata of rats and mice with experimental parkinsonism. *Exp Neurol* **129**, 57–63

(1994).

48. Stamford, J. A., Kruk, Z. L. & Millar, J. Actions of dopamine antagonists on stimulated striatal

and limbic dopamine release: an in vivo voltammetric study. *Br J Pharmacol* **94**, 924–932

(1988).

49. Palij, P. *et al.* Presynaptic regulation of dopamine release in corpus striatum monitored in vitro in

real time by fast cyclic voltammetry. *Brain Res* **509**, 172–174 (1990).

50. Fawaz, C. S., Martel, P., Leo, D. & Trudeau, L.-E. Presynaptic action of neurotensin on

dopamine release through inhibition of D2 receptor function. *BMC Neuroscience* **10**, 96 (2009).

51. Hoffman, A. F., Spivak, C. E. & Lupica, C. R. Enhanced Dopamine Release by Dopamine

Transport Inhibitors Described by a Restricted Diffusion Model and Fast-Scan Cyclic

Voltammetry. *ACS Chem Neurosci* **7**, 700–709 (2016).

52. Carvelli, L., McDonald, P. W., Blakely, R. D. & DeFelice, L. J. Dopamine transporters

depolarize neurons by a channel mechanism. *PNAS* **101**, 16046–16051 (2004).

53. Sun, W., Ginovart, N., Ko, F., Seeman, P. & Kapur, S. In Vivo Evidence for Dopamine-Mediated

Internalization of D2-Receptors after Amphetamine: Differential Findings with [3H]Raclopride

versus [3H]Spiperone. *Mol Pharmacol* **63**, 456–462 (2003).

54. Guo, N. *et al.* Impact of D2 receptor internalization on binding affinity of neuroimaging

radiotracers. *Neuropsychopharmacology* **35**, 806–817 (2010).

55. Parish, C. L., Finkelstein, D. I., Drago, J., Borrelli, E. & Horne, M. K. The role of dopamine

receptors in regulating the size of axonal arbors. *J Neurosci* **21**, 5147–5157 (2001).

56. Tripanichkul, W., Stanic, D., Drago, J., Finkelstein, D. I. & Horne, M. K. D2 Dopamine receptor

blockade results in sprouting of DA axons in the intact animal but prevents sprouting following

nigral lesions. *Eur J Neurosci* **17**, 1033–1045 (2003).

57. Fasano, C., Poirier, A., DesGroseillers, L. & Trudeau, L.-E. Chronic activation of the D2

dopamine autoreceptor inhibits synaptogenesis in mesencephalic dopaminergic neurons in vitro.

Eur J Neurosci **28**, 1480–1490 (2008).

58. Fasano, C., Kortleven, C. & Trudeau, L.-E. Chronic activation of the D2 autoreceptor inhibits

both glutamate and dopamine synapse formation and alters the intrinsic properties of

mesencephalic dopamine neurons in vitro. *Eur J Neurosci* **32**, 1433–1441 (2010).

59. Fasano, C. *et al.* Dopamine facilitates dendritic spine formation by cultured striatal medium spiny

neurons through both D1 and D2 dopamine receptors. *Neuropharmacology* **67**, 432–443 (2013).

60. Giguère, N. *et al.* Increased vulnerability of nigral dopamine neurons after expansion of their

axonal arborization size through D2 dopamine receptor conditional knockout. *PLOS Genetics* **15**,

e1008352 (2019).

61. Gagnon, D. *et al.* Evidence for Sprouting of Dopamine and Serotonin Axons in the Pallidum of

Parkinsonian Monkeys. *Front. Neuroanat.* **12**, (2018).

62. Maximov, A. *et al.* Genetic analysis of synaptotagmin-7 function in synaptic vesicle exocytosis.

PNAS **105**, 3986–3991 (2008).

63. Bacaj, T. *et al.* Synaptotagmin-1 and Synaptotagmin-7 Trigger Synchronous and Asynchronous

Phases of Neurotransmitter Release. *Neuron* **80**, 947–959 (2013).

64. Bacaj, T. *et al.* Synaptotagmin-1 and -7 Are Redundantly Essential for Maintaining the Capacity

of the Readily-Releasable Pool of Synaptic Vesicles. *PLOS Biology* **13**, e1002267 (2015).

65. Wang, C. *et al.* Synaptotagmin-11 is a critical mediator of parkin-linked neurotoxicity and

Parkinson's disease-like pathology. *Nat Commun* **9**, 1–14 (2018).

66. Pang, Z. P. *et al.* Doc2 supports spontaneous synaptic transmission by a Ca2+-independent

mechanism. *Neuron* **70**, 244–251 (2011).

67. Sulzer, D., Cragg, S. J. & Rice, M. E. Striatal dopamine neurotransmission: regulation of release

and uptake. *Basal Ganglia* **6**, 123–148 (2016).

68. Sulzer, D., Sonders, M. S., Poulsen, N. W. & Galli, A. Mechanisms of neurotransmitter release

by amphetamines: a review. *Prog. Neurobiol.* **75**, 406–433 (2005).

69. Parish, C. L. *et al.* Effects of long-term treatment with dopamine receptor agonists and

antagonists on terminal arbor size. *European Journal of Neuroscience* **16**, 787–794 (2002).

70. Chen, B. T., Patel, J. C., Moran, K. A. & Rice, M. E. Differential calcium dependence of axonal
versus somatodendritic dopamine release, with characteristics of both in the ventral tegmental
area. *Front Syst Neurosci* **5**, 39 (2011).

71. Beckstead, M. J., Grandy, D. K., Wickman, K. & Williams, J. T. Vesicular dopamine release
elicits an inhibitory postsynaptic current in midbrain dopamine neurons. *Neuron* **42**, 939–946
(2004).

72. Beckstead, M. J., Ford, C. P., Phillips, P. E. M. & Williams, J. T. Presynaptic regulation of
dendrodendritic dopamine transmission. *Eur. J. Neurosci.* **26**, 1479–1488 (2007).

73. Courtney, N. A., Mamaligas, A. A. & Ford, C. P. Species Differences in Somatodendritic
Dopamine Transmission Determine D2-Autoreceptor-Mediated Inhibition of Ventral Tegmental
Area Neuron Firing. *J. Neurosci.* **32**, 13520–13528 (2012).

74. Ford, C. P., Phillips, P. E. M. & Williams, J. T. The Time Course of Dopamine Transmission in
the Ventral Tegmental Area. *J. Neurosci.* **29**, 13344–13352 (2009).

75. Ford, C. P., Gantz, S. C., Phillips, P. E. M. & Williams, J. T. Control of extracellular dopamine at
dendrite and axon terminals. *J. Neurosci.* **30**, 6975–6983 (2010).

76. Fernandes Xavier, F. G., Doucet, G., Geffard, M. & Descarries, L. Dopamine neoinnervation in
the substantia nigra and hyperinnervation in the interpeduncular nucleus of adult rat following
neonatal cerebroventricular administration of 6-hydroxydopamine. *Neuroscience* **59**, 77–87
(1994).

77. Bezard, E., Gross, C. E. & Brotchie, J. M. Presymptomatic compensation in Parkinson's disease
is not dopamine-mediated. *Trends in Neurosciences* **26**, 215–221 (2003).

78. Parmar, M., Torper, O. & Drouin-Ouellet, J. Cell-based therapy for Parkinson's disease: A
journey through decades toward the light side of the Force. *Eur. J. Neurosci.* **49**, 463–471 (2019).

79. Palmiter, R. D. Dopamine signaling in the dorsal striatum is essential for motivated behaviors:
lessons from dopamine-deficient mice. *Ann N Y Acad Sci* **1129**, 35–46 (2008).

80. Cagniard, B. *et al.* Dopamine scales performance in the absence of new learning. *Neuron* **51**,
541–547 (2006).

81. Berridge, K. C. The debate over dopamine's role in reward: the case for incentive salience.

Psychopharmacology (Berl) **191**, 391–431 (2007).

82. Ferris, M. J. *et al.* Dopamine transporters govern diurnal variation in extracellular dopamine tone.

Proc Natl Acad Sci U S A **111**, E2751–E2759 (2014).

83. Zweifel, L. S. *et al.* Disruption of NMDAR-dependent burst firing by dopamine neurons provides

selective assessment of phasic dopamine-dependent behavior. *Proc Natl Acad Sci U S A* **106**,

7281–7288 (2009).

84. Kochubey, O., Babai, N. & Schneggenburger, R. A Synaptotagmin Isoform Switch during the

Development of an Identified CNS Synapse. *Neuron* **90**, 984–999 (2016).

85. Rice, M. E. *et al.* Direct monitoring of dopamine and 5-HT release in substantia nigra and ventral

tegmental area in vitro. *Exp Brain Res* **100**, 395–406 (1994).

86. Cragg, S. J., Hawkey, C. R. & Greenfield, S. A. Comparison of serotonin and dopamine release in

substantia nigra and ventral tegmental area: region and species differences. *J. Neurochem.* **69**,

2378–2386 (1997).

87. Itzhak, Y. & Martin, J. L. Effects of cocaine, nicotine, dizocipline and alcohol on mice locomotor

activity: cocaine–alcohol cross-sensitization involves upregulation of striatal dopamine

transporter binding sites. *Brain Research* **818**, 204–211 (1999).

88. Steinkellner, T. *et al.* In Vivo Amphetamine Action is Contingent on α CaMKII.

Neuropsychopharmacol **39**, 2681–2693 (2014).

89. Centonze, D. *et al.* Distinct Roles of D1 and D5 Dopamine Receptors in Motor Activity and

Striatal Synaptic Plasticity. *J Neurosci* **23**, 8506–8512 (2003).

90. Dourado, M., Cardoso-Cruz, H., Monteiro, C. & Galhardo, V. Effect of Motor Impairment on

Analgesic Efficacy of Dopamine D2/3 Receptors in a Rat Model of Neuropathy. *J Exp Neurosci*

10, JEN.S36492 (2016).

91. Radl, D. *et al.* Differential regulation of striatal motor behavior and related cellular responses by

dopamine D2L and D2S isoforms. *PNAS* **115**, 198–203 (2018).

92. Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates, Compact - 5th Edition.

https://www.elsevier.com/books/paxinos-and-franklins-the-mouse-brain-in-stereotaxic-

coordinates-compact/franklin/978-0-12-816159-3.

93. Cliburn, R. A. *et al.* Immunochemical localization of vesicular monoamine transporter 2

(VMAT2) in mouse brain. *Journal of Chemical Neuroanatomy* **83–84**, 82–90 (2017).

94. Isingrini, E. *et al.* Genetic elimination of dopamine vesicular stocks in the nigrostriatal pathway

replicates Parkinson's disease motor symptoms without neuronal degeneration in adult mice. *Sci*

Rep **7**, (2017).

1418

Acknowledgements

We would like to thank Dr. R. Schneggenburger for kindly providing the Syt1-floxed mouse, Dr. J. Surmeier for kindly provided TH-cre and control viruses, Dr. G. Miller for kindly provided VMAT2 antibody, Dr. A. Gratton for interpretation of HPLC data and M-E. Delignat-Lavaud for illustrations. This work was funded by the National Sciences and Engineering Research Council of Canada (NSERC, grant RGPIN-2020-05279) to L.-E.T. and by the Krembil Foundation (to L.-E.T.). This research was also funded in whole or in part by Aligning Science Across Parkinson's [ASAP 000525] (to L.-E.T.) through the Michael J. Fox Foundation for Parkinson's Research (MJFF). For the purpose of open access, the author has applied a CC BY public copyright license to all author accepted manuscripts arising from this submission. B.D-L received a graduate student award from Parkinson Canada.

Author contributions

B.D-L.: conceptualization and design, acquisition, analysis, validation and interpretation of data (voltammetry, behavior, autoradiography, IHC, viral infections and RT-qPCR), drafting or revising the manuscript. J.K.: acquisition and analysis of data (IHC). C.D. and R.D.: acquisition and analysis of data (ICC). I.M.: acquisition and analysis of data (microdialysis). S.M.: acquisition and analysis of data (Western Blot). N.G.: acquisition and analysis of data (stereology), revising the manuscript. A.T. acquisition and analysis of data (stereology), L.M.: acquisition of data (HPLC). C.L.: acquisition of data (autoradiography). S.B.: acquisition of data (ICC). M-J.B.: acquisition of data (cell culture). P. R-N.: review and editing. D.L.: review, editing and analysis of data (autoradiography). L. D.B.: review and editing. L-E.T.: conceptualization and design, resources, funding acquisition, writing - review and editing.

1443

1444

1445 **Competing interests statement**

1446 The authors declare that they have no conflicts of interest.

 Figures and legends

 Figure 1: Syt1 is the main calcium sensor for fast axonal dopamine release. (A)

 Generation of conditional knock-out of Syt1 in DA neurons by crossing Syt1-floxed mice

 (Syt1$^{lox/lox}$) with DAT^{IREScre} mice. **(B)** Fast scan cyclic recording of Syt1 cKO^{DA} mice in the

 dorsal striatum. Representative traces (top) and quantification of peak amplitude (bottom)

 obtained with single-pulse electrical stimulation (1 ms, 400 µA) in Syt1$^{+/+}$ (n = 18 slices/9

 mice), Syt$^{+/-}$ (n = 16/8) and Syt1$^{-/-}$ mice (n = 16/8). **(C)** Same, but in the ventral striatum

 (NAc core and shell, n =18 slices/9 mice in Syt1$^{+/+}$, n = 16/8 in Syt$^{+/-}$ and n =16/8 in Syt1$^{-/-}$).

 (D) Representative traces (top) and quantification of peak amplitude (bottom) obtained in the

 VTA (n =16 slices/9 mice in Syt1$^{+/+}$, n = 14/8 in Syt$^{+/-}$ and n =16/8 in Syt1$^{-/-}$) with aCSF

 containing nomifensine (DAT blocker) and sulpiride (D2 antagonist) (both at 5 µM), and

 pulse-train stimulation (30 pulses of 1 ms at 10 Hz, 400µA). **(E)** Same for the SNc (n =11

 slices/6 mice in Syt1$^{+/+}$, n = 10/5 in Syt$^{+/-}$ and n =9/5 in Syt1$^{-/-}$). Error bars represent ± S.E.M.

 and the statistical analysis was carried out by 1-way ANOVAs followed by Tukey tests (ns,

 non-significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$). Source data are

 provided as a Source Data file.

 Figure 2: Somatodendritic optogenetics reveals unaltered STD DA release in the

 absence of Syt1. (A) Immunohistochemistry in brain slices of adult Syt1 cKO^{DA} mice

 infected with AAV2/5-hsyn-DIO-ChR2-eYFP-Kv (n = 4 mice, one representative set of

 images is shown), showing expression of ChR2-Kv (eYFP) in the whole ventral

 mesencephalon (scale bar = 500 µm) and striatum (scale bar = 1 mm) using epifluorescence

 microscope at 20x. **(B)** Expression of ChR2-Kv in the same mice evaluated by confocal

microscopy at 60x shows infected DA neurons (TH) in the midbrain (up panel, scale bar = 50

µm). Optical zoom and z-stack on an infected DA neuron showing the membrane distribution

of the eYFP signal (bottom panel, scale bar = 10 µm, insert in the merge image shows the

eYFP signal at a single focal plane). Evaluation of the eYFP signal in the striatum shows no

axonal processes in the dorsal sector but a small contingent of positive fibers in the ventral

sector of the striatum (scale bar = 50 µm). **(C-E)** Fast scan cyclic voltammetry recordings

with average [DA]o peaks obtained in dorsal striatum **(C)**, ventral striatum **(D)** and VTA **(E)**

slices of infected $Syt1^{+/+}$ and $Syt1^{-/-}$ mice with AAV2/5-hsyn-DIO-ChR2-eYFP-Kv (ChR2-

Kv, n = 14 slices/7 mice in $Syt1^{+/+}$, n = 17 slices/9 mice in $Syt1^{-/-}$) or AAV5-EF1a-DIO-

hChR2(H134R)-eYFP (control hChR2, n = 10 slices/ 5 mice). Representative traces of the

striatum (dorsal + ventral) and VTA are shown in **(F)** and **(G)**. DA release was optically

triggered in each region using pulse-train stimulation (30 pulses of 470 nm blue light at 10

Hz) in ACSF containing 5 µM of nomifensine and sulpiride. **(H)** TTX (1 µM) effect on

average [DA]o peaks in the striatum and the VTA (% of inhibition) (n = 14 slices/7 mice in

$Syt1^{+/+}$, n = 17 slices/9 mice in $Syt1^{-/-}$). Error bars represent ± S.E.M. and the statistical

analysis was carried out by 2-way ANOVA followed by Šidák (panel E) and Tuckey (panels

C-D and H) tests (ns, non-significant; *, p < 0.05; **, p < 0.01; ***, p < 0.001; ****, p <

0.0001). Source data are provided as a Source Data file.

1489

Figure 3: Syt1 cKO^{DA} mice do not exhibit any substantial motor defects. (A) Latency to

fall from the device during the rotarod test for $Syt1^{+/+}$, $Syt1^{+/-}$ and $Syt1^{-/-}$ mice (n = 8, mean of

3 attempts) **(B)** Time for $Syt1^{+/+}$, $Syt1^{+/-}$ and $Syt1^{-/-}$ mice (n = 8) to turn (t-turn) and climb

down the vertical pole (mean of 3 attempts/2 days). **(C)** Forelimb grip strength (mean of 3

trials/weight) developed by $Syt1^{+/+}$, $Syt1^{+/-}$ and $Syt1^{-/-}$ mice (n = 8), using a force sensor

connected to a grid. Statistical analysis for A-C were carried out by 1-way ANOVAs

followed by Dunnett tests. **(D-L)** Locomotion of Syt1 cKO^{DA} mice measured as traveled

distance (% of a 20 min baseline) under saline treatment (n = 8 mice) **(D)**, cocaine at 20

mg/kg (n = 8 Syt1$^{+/+}$/Syt1$^{+/-}$ and 6 Syt1$^{-/-}$) **(E)**, amphetamine at 5 mg/kg (n = 8 mice) **(F)**, the

D1 agonist SCH23390 at 50 µg/kg (n = 10 Syt1$^{+/+}$/Syt1$^{-/-}$ and 9 Syt1$^{+/-}$) **(H)**, the D2 agonist

quinpirole at 0.2 mg/kg (n = 10 Syt1$^{-/-}$ / Sy$^{+/-}$ and 8 Syt1$^{+/+}$) **(I)** and the D2 antagonist

raclopride at 1 mg/kg (n = 10 mice) **(J)**. Statistical analyses for D-J were carried out by 2-way

ANOVAs followed by Dunnett tests. **(G&K)** Average drug-induced locomotion (% of

baseline) during a 40 min recording period (mean between 25-60 min) are represented for

cocaine/amphetamine treatment **(G)** and for SCH23390/quinpirole/raclopride treatments **(K)**.

Blue squares indicate the average traveled distance for each mouse, 5 min after receiving an

i.p. injection of each tested drugs **(L)**. Statistical analysis was carried out by 1-way ANOVAs

followed by Dunnett tests (panels A-C and L), mixed-effects ANOVA model with the

Geisser-Greenhouse correction (panels D-F and H-J) and 2-way ANOVAs followed by

Tuckey tests (panels G and K). Error bars represent ± S.E.M. (ns, non-significant; *, p < 0.05;

**, p < 0.01; ***, p < 0.001; ****, p < 0.0001). Source data are provided as a Source Data

file.

1512

Figure 4: Syt1 is dispensable for motivation to work for food. **(A)** Schematic

representation of operant food-rewarded nose-poking protocol in Syt1 cKODA mice. **(B-C)**

time course of the number of pellets earned **(B)** and pokes made **(C)** by Syt1$^{+/+}$ (green lines, n

= 8 mice) and Syt1$^{-/-}$ (black lines, n = 10 mice) mice using "D**C)** Efficiency of the pokes (%

of correct answer) in the FR1 condition. **(E-F)** time course of the number of pellets earned by

Syt1$^{+/+}$ and Syt1$^{-/-}$ mice **(E)** and pokes **(F)** over 6 days in a progressive ratio (PR) 1 paradigm

(the nose-poking requirement began on FR1, increased by 1 poke each time a pellet was

earned and was reset to FR1 if no poking was performed on either the active or inactive port

for 30 min). **(G)** Efficiency of the pokes (% of correct port entries) in the PR1 condition. **(H)**

Maximal breaking points (poke thresholds) achieved by $Syt1^{+/+}$ (green, n = 8) and $Syt1^{-/-}$

(black, n = 10) during each day (D) and night (N) session of the progressive ratio protocol.

For each experiment, mice were placed in a 12h/12h light/dark cycle and the Zeitgeber time

corresponds to the onset of the devices. Statistical analyses were carried out by mixed-effects

ANOVA model with the Geisser-Greenhouse correction (panels A-F) and 2-way ANOVAs

followed by Šidák's test (panel G). Error shading bands in panels A-F (green for $Syt1^{+/+}$, light

grey for $Syt1^{-/-}$) and error bars in panel 4G represent ± S.E.M. (ns, non-significant; *, p <

0.05; #, p < 0.0001). The schematic diagram of the FED3 nose-poking device is modified

from https://github.com/KravitzLabDevices/FED3/blob/main/photos/mouse_feeder.svg.

Source data are provided as a Source Data file.

1532

Figure 5: Basal extracellular DA levels and total tissue DA are not altered in Syt1

cKO^{DA} mice. (A) Schematic representation of the intracerebral microdialysis protocol

performed on anesthetized Syt1 cKO^{DA} and control mice. **(B)** Quantification of total DA,

DOPAC, serotonin and norepinephrine striatal content (ng/mg of total proteins) and

DA/DOPAC ratio (n = 10 $Syt1^{+/+}$ and 12 $Syt1^{-/-}$). Statistical analysis was carried out by two-

tailed unpaired t-test. **(C)** Extracellular quantification (ng/mL) by microdialysis of the same

molecules in the dorsal striatum (n ($Syt1^{+/+}$/ $Syt1^{-/-}$ mice) = 10/12 for DA, 11/12 for DOPAC,

8/12 for 5-HT and 10/12 for NE) and in the mesencephalon (n ($Syt1^{+/+}$/ $Syt1^{-/-}$ mice) = 9/12

for DA, 10/12 for DOPAC, 9/12 for 5-HT and 9/12 for NE) of Syt1 cKO^{DA} mice, with an

average of 3 dialysates for each animal. Statistical analysis was carried out by 2-way

ANOVAs followed by Šidák's tests. Error bars represent ± S.E.M. (ns, non-significant; *, p <

0.05; **, p < 0.01; ***, p < 0.001; ****, p < 0.0001). Source data are provided as a Source

Data file.

Figure 6: Increased D2 autoreceptor function in Syt1 cKO^{DA} mice. (A) FSCV representative traces (top) and quantification of peak amplitude (bottom) obtained in Syt1$^{+/+}$ (n = 9) and Syt1$^{-/-}$ (n = 7) mice with single-pulse electrical stimulation and aCSF containing 1 µM of the D2 agonist quinpirole. **(B)** Same, but in the ventral striatum (n = 9 Syt1$^{+/+}$ and 8 Syt1$^{-/-}$). **(C)** FSCV representative traces (top) and quantification of peak amplitude (bottom) obtained in Syt1$^{+/+}$ (n = 8) and Syt1$^{-/-}$ (n = 11) mice with pulse-train stimulation and aCSF containing 5 µM of the D2 antagonist sulpiride. **(D)** Same, but in the ventral striatum (n = 8 Syt1$^{+/+}$ and 11 Syt1$^{-/-}$). **(E)** FSCV representative traces (top) and quantification of peak amplitude (bottom) obtained in Syt1$^{+/+}$ (n = 4) and Syt1$^{-/-}$ (n = 6) mice with pulse-train stimulation and aCSF containing 5 µM of the DAT blocker nomifensine. **(F)** Same, but in the ventral striatum (n = 4 Syt1$^{+/+}$ and 6 Syt1$^{-/-}$). For each tested drug, the effect on DA release (% of control) for both genotypes is indicated in the right panel. Statistical analysis was carried out by 2-way ANOVAs followed by Šidák's corrections. Error bars represent ± S.E.M. (ns, non-significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$). Source data are provided as a Source Data file.

Figure 7: Higher D2 receptor density and lower D2 affinity in Syt1 cKO^{DA} mice. (A-B) Density of the dopamine D1 receptors (Bmax) **(A)** and affinity (Kd) for the radioligand [^{3}H]-SCH23390 **(B)** in Syt1$^{+/+}$ (n = 4) and Syt1$^{-/-}$ (n = 4) mice. **(C-D)** Density of the dopamine D2 receptors (Bmax) **(C)** and affinity (Kd) for the radioligand [^{3}H]-raclopride **(D)** in Syt1$^{+/+}$ (n = 5) and Syt1$^{-/-}$ mice (n = 4). Each binding was analyzed with a series of 4 serial coronal sections (bregma +1.34 mm to +0.38 mm) and by dividing each section into 4 quadrants (dorso-lateral, ventro-lateral, dorso-medial and ventro-medial). **(E)** Binding potential of D1

1570 and D2 receptors defined as the Bmax/Kd ratio (n = 4 Syt1$^{+/+}$ / 4 Syt1$^{-/-}$ for D1 and 5 Syt1$^{+/+}$ /

1571 4 Syt1$^{-/-}$ for D2). The data followed a log-normal distribution and two-tailed unpaired t-tests

1572 were done after the data were converted to a log scale. Error bars represent ± S.E.M. (ns, non-

1573 significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$). Source data are

1574 provided as a Source Data file.

1575

Figure 8: Adaptations of DA innervation in the striatum of Syt1 cKO^{DA} mice. (A)

Immunochemistry of striatal slices from 10-12-week-old Syt1$^{+/+}$ and Syt1$^{-/-}$ mice (60x

confocal) using (from top to bottom): tyrosine hydroxylase (TH), vesicular monoamine

transporter 2 (VMAT2), dopamine transporter (DAT) and serotonin (5-HT) immunostainings.

Scale bar = 50 µm. **(B)** Quantification of each signal surface (% of WT) in the dorsal and

ventral striatum (n = 20 hemispheres/10 mice for both genotypes). **(C)** Same with signal

intensity (n = 20 hemispheres/10 mice for both genotypes). **(D)** TH immunoreactive cells in

the midbrain of Syt1$^{+/+}$ and Syt1$^{-/-}$ mice, identified by the brown DAB staining. **(E)**

Stereological counts of DA neurons in the SNc (n = 5 Syt1$^{+/+}$ / 5 Syt1$^{-/-}$), VTA (n = 5/5) and

RRF (n = 4/5). Unbiased stereological counting was performed by a blinded observer to

estimate the number of DA neurons. Statistical analysis was carried out by 2-way ANOVAs

followed by Šidák's corrections. Error bars represent ± S.E.M. (ns, non-significant; *, $p <$

0.05; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$). Source data are provided as a Source

Data file.

1590

1591 **Figure 9: Acute deletion of Syt1 in adult mice impairs DA release but prevents**

1592 **adaptations. (A)** Stereotaxic injections of AAV9-TH-Cre and AAV9-TH-FusionRed

1593 (control) viruses in 6-7-week-old Syt1$^{lox/lox}$ mice effectively infects the whole ventral

mesencephalon (epifluorescence imaging of TH (cyan) and FusionRed (RFP) (Cre, magenta) immunostainings were performed in 4 mice per condition and one representative image of the infections is represented). **(B)** FSCV recordings (DA overflow in µM) of dorsal and ventral striatum (single electrical pulse 1 ms, 400 µA, normal aCSF) and VTA (pulse-train 30p, 1ms, 400 µA, 10 Hz in presence of 5 µM of nomifensine/sulpiride) from Syt1$^{lox/lox}$ mice injected with AAV9-TH-Cre (n = 8) and AAV9-TH-Control viruses (n = 7). **(C)** Immunohistochemistry of striatal slices from 9-10-week old Syt1$^{lox/lox}$ injected with AAV9-TH-Cre (black) and AAV9-TH-control (green) illustrating TH and DAT immunostaining (60x confocal). Scale bar = 50 µm. **(D)** Quantification of each signal surface (% of WT) in the dorsal and ventral striatum of injected mice (n = 12 hemispheres/6 mice for each condition). **(E)** Same with signal intensity (n = 12 hemispheres/6 mice for each condition). Statistical analysis was carried out by 2-way ANOVAs followed by Šidák's corrections. Error bars represent ± S.E.M. (ns, non-significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$). Source data are provided as a Source Data file.

A

Syt1-floxed   Ex1   LoxP   Ex2   LoxP   Ex3   Ex4 ...   X   5' — DAT — IRES — NLS-Cre — 3'

Syt1 cKO^DA   Ex1   Ex3   Ex4 ...

B   Dorsal striatum

FSCV   Single pulse

— Syt1 -/-
— Syt1 +/-
— Syt1 +/+

0.4 µM   2s

■ Syt1 +/+
□ Syt1 +/-
□ Syt1 -/-

****
****
ns

Dopamine overflow (µM)

Syt1 +/+   Syt1 +/-   Syt1 -/-

C   Ventral striatum

Single pulse

0.4 µM   2s

****
****
**

Dopamine overflow (µM)

Syt1 +/+   Syt1 +/-   Syt1 -/-

D   VTA

Pulse-train
(30p 10 Hz)

(aCSF + 5 µM nomifensine/sulpiride)

0.1 µM   10s

****
**
*

Dopamine overflow (µM)

Syt1 +/+   Syt1 +/-   Syt1 -/-

E   SNc

Pulse-train
(30p 10 Hz)

(aCSF + 5 µM nomifensine/sulpiride)

0.1 µM   10s

**
*
ns

Dopamine overflow (µM)

Syt1 +/+   Syt1 +/-   Syt1 -/-

A | In vivo (epifluorescence 20x)
B | In vivo (confocal 60x)

Midbrain: eYFP, TH, Merge
Striatum: eYFP, TH, Merge

Midbrain: eYFP, TH, MAP2, Merge
dStr: eYFP, TH, MAP2, Merge
vStr: eYFP, TH, MAP2, Merge

Fast Scan Cyclic Voltammetry

Optical pulse-train 470 nm (30p, 10 Hz)
aCSF + 5 µM nomi./sulpi.

Syt1$^{+/+}$ hChR2 | Syt1$^{+/+}$ ChR2-Kv | Syt1$^{-/-}$ ChR2-Kv

C Dorsal striatum

Dopamine overflow (µM)
ns, ns, ns, ns, ****, ns, ns, ****, ns
ACSF, TTX

D Ventral striatum

vStr
Dopamine overflow (µM)
****, *, ns, ****, ns, ns, ***, ns
ACSF, TTX

E VTA

Dopamine overflow (nM)
ns, ns, ns, ****, ****, ns, ***, ns
ACSF, TTX

F Striatum

hChR2, ChR2-Kv
aCSF, TTX
0.2 µM, 5s
Syt1$^{+/+}$, Syt1$^{-/-}$

G VTA

hChR2, ChR2-Kv
50 nM, 10s
Syt1$^{+/+}$, Syt1$^{-/-}$

H

Dopamine overflow (% of inhibition by TTX)
ns, ns, ns, ns, ****, **, ns, ns
dStr, vStr, VTA

A

Free-feeding (habituation)	Fixed Ratio 1 (learning)	Progressive Ratio 1 (motivation)
2 days	3 days	6 days

Fixed Ratio 1 (FR1) **Progressive Ratio 1 (PR1)**

— Syt1 +/+ (± sem)   — Syt1 -/- (± sem)

B
Pellets vs Zeitgeber time (hours) #

C
Pokes vs Zeitgeber time (hours) #

D
Pokes efficiency (correct %) vs Zeitgeber time (hours)

E
Pellets vs Zeitgeber time (hours)

F
Pokes vs Zeitgeber time (hours) # *

G
Pokes efficiency (correct %) vs Zeitgeber time (hours)

H
Breaking point in PR1 (pokes): D1 ns, N1 ns, D2 ns, N2 ns, D3 ns, N3 ns, D4 ns, N4 ns, D5 ns, N5 ns, D6 ns, N6 ns

A

AAV9-TH-cre-myc-2A-fusion-red (**TH-CRE**)
or AAV9-TH-fusion-red-Sv40 (**TH-CTRL**)

TH | CRE

Merge

B

Single pulse
aCSF

Pulse-train
aCSF + 5 μM nomi./sulpi.

■ Syt1^{lox/lox} AAV-TH-CTRL(FusionRed)
□ Syt1^{lox/lox} AAV-TH-CRE(FusionRed)

Dopamine overflow (μM)

	Dorsal	Ventral	VTA
	****	****	*

C

Syt1^{lox/lox} AAV-CTRL | Syt1^{lox/lox} AAV-CRE

TH

DAT

D Signal surface

TH striatal surface area (% of WT)

ns | *

Dorsal | Ventral

DAT striatal surface area (% of WT)

ns | ns

Dorsal | Ventral

E Signal intensity

TH signal intensity (% of WT)

ns | ns

Dorsal | Ventral

DAT signal intensity (% of WT)

ns | ns

Dorsal | Ventral