Title: Environment-dependence of the expression of mutational load and species' range limits

Running title: Mutational load and environmental stress

Authors: Antoine Perrier[1,2*], Darío Sánchez-Castro[2], Yvonne Willi[2]

[1] Department of Biology, University of Virginia, 485 McCormick Road, Charlottesville, VA 22904, USA

[2] Department of Environmental Sciences, University of Basel, Schönbeinstrasse 6, 4056 Basel, Switzerland.

*Corresponding author

cdt9qe@virginia.edu; https://orcid.org/0000-0002-6447-5692

dario.sancast@gmail.com; https://orcid.org/0000-0002-2552-1692

yvonne.willi@unibas.ch; https://orcid.org/0000-0001-7654-3261

Acknowledgements: This work was supported by the Swiss National Science Foundation (31003 A_166322). We are grateful to Celia Evans (Paul Smith's College, Paul Smith, NY), Joan Edwards (Williams College, Williamstown, MA), Heather Peckham Griscom (James Madison University, Harrisonburg, VA), William K. Smith (Wake Forest University, Winston-Salem, NC) and Rodney Mauricio (University of Georgia, Athens, GA) for logistical support in the US. For field assistance, we thank Mary Anderson, Michael Boyd, Bennet Coe, Scott Cory, Rachel Hillyer, Andrew Jones, Deidre Keating, Larry Kummer, David Lampman, Anastasia Levie-Sprick, Blake Macko, Shannon Malisson, Kathryn McGee, Althea Neighbors, Debra Rogers-Gillig, Caleb Rose, Amber Scarabaggio, Anna Shutley, Caroline Vath and Audrey Werner. For assistance with seed counts, we thank Olivier Bachmann, Markus Funk, and Susanna Riedl. Collection permits were provided by the Clinton

County Conservation Board, Cornell University, Fort Leonard Wood Army Base, Iowa Department of Natural Resources, Missouri Department of Conservation, New York State Office of Parks, Ontario Parks, Palisades Interstate Park Commission, Rock Island Lodge, United States National Park Service, Virginia Department of Conservation and Recreation and the Wisconsin Department of Natural Resources.

Author Contributions: All authors contributed to the study design. YW collected the seeds in the field, AP and DS raised and crossed plants. AP analyzed the data and wrote the manuscript, with support by YW.

Conflict of Interest Statement: The authors declare that there is no conflict of interest.

Data accessibility statement: All data is stored in Dryad (https://doi.org/10.5061/dryad.cc2fqz642).

Title: Environment-dependence of the expression of mutational load and species' range limits

Abstract:

Theoretical and empirical research on the causes of species' range limits suggests the contribution of

several intrinsic and extrinsic factors, with potentially complex interactions among them. An intrinsic

factor proposed by recent theory is mutational load increasing towards range edges because of genetic

drift. Furthermore, environmental quality may decline toward range edges and enhance the expression

of load. Here we tested whether the expression of mutational load associated with range limits in the

North American *Arabidopsis lyrata* is enhanced under stressful conditions, by comparing the

performance of within- *versus* between-population crosses at common garden sites across the species'

distribution and beyond. Heterosis, reflecting the expression of load, increased with heightened

estimates of genomic load and with environmental stress caused by warming, but the interaction was

not significant. We conclude that range-edge populations suffer from a twofold genetic Allee effect

caused by increased mutational load and stress-dependent load linked to general heterozygote

deficiency, but no synergistic effect between them.

Keywords: *Arabidopsis lyrata*, environmental stress, genetic drift, range limits, heterosis, mutational

load.

Introduction

The geographic distribution of species can be constrained by several factors, and they may interact in complex manners (Gaston, 2009; Roy et al., 2009; Sexton et al., 2009; Connallon & Sgrò, 2018; Willi & Van Buskirk, 2019). An important factor put forward by relatively recent theory is the accumulation of mutational load in range-edge populations due to a history of enhanced genetic drift (Peischl et al., 2013; Henry et al., 2015; Willi, 2019). In parallel, environmental conditions may degrade toward the range limits of a species (Brown, 1984; Cahill et al., 2014; Hargreaves et al., 2014; Lee-Yaw et al., 2016). In addition to their respective negative effect on population performance, mutational load and environmental stress may interact synergistically in their effect on performance, as was suggested and found for inbreeding depression (Roff, 1997, pages 285-338; Reed et al., 2012). The contribution of mutational load to range limits could therfore be much stronger than expected by previous theoretical and empirical work, and relevant for shaping the distribution of many species.

Two types of demographic scenarios at range edges are predicted to lead to increased mutational load. First, theoretical models predict that range expansion through serial demographic bottlenecks enhances genetic drift and lowers the efficacy of purifying selection, which leads to the accumulation of deleterious mutations (Peischl et al., 2013; 2015; Peischl & Excoffier, 2015). High load at the expansion front is predicted to slow down the spread of a species, or halt it if genome-wide recombination is low (Peischl et al., 2015). The so-called expansion load is based mainly on recessive deleterious mutations (Peischl & Excoffier, 2015), establishes in the absence of environmental gradients, and can persist in populations for thousands of generations (Peischl et al., 2013). Second, if populations are small and away from the core of distribution, genetic drift is also enhanced and increases mutational load. Simulations showed that when the carrying capacity of suitable habitat declines along a line of habitat patches, the occupied range becomes shorter than the distriubtion of habitat would suggest because of mutation accumulation (Henry et al., 2015). The effect of load is larger if dispersal and population growth rates are low (Henry et al., 2015), and this

scenario of mutational load limiting ranges may apply best to rear edges of species distribution

(Hampre & Petit, 2005). Genomic evidence for the accumulation of mutational load with range

expansion was found for humans (Lohmueller et al., 2008; Peischl et al., 2018), and in plants for

expansion and rear-edge isolation (Zhang et al., 2016; González-Martínez et al., 2017; Willi et al.,

2018; Koski et al., 2019).

At range limits, the environment may also become more stressful. Empirical studies document

a general decline in habitat suitability toward range limits in many taxa (Sexton et al., 2009; Pironon

et al., 2017). A rather consistent and pronounced effect was found in species experimentally

transplanted beyond their range edges as their lifetime performance generally declined compared to

sites within the range (Hargreaves et al., 2014). The decline in performance was found to coincide

with a decline in predicted habitat suitability revealed by niche modelling (Lee-Yaw et al., 2016).

The two meta-level studies suggest that the decay in habitat suitability is rather general. Such

unfavorable conditions may act in isolation and lower population performance, or they may enhance

the expression of mutational load at range edges, further reducing population performance.

Potential mechanisms for a synergistic interaction between mutational load and environmental

stress have rarely been explored *per se*, but have been mainly discussed and tested in the context of

inbreeding depression (Armbruster & Reed, 2005; Willi et al., 2007a; Fox & Reed, 2010). Three

mechanisms have been proposed (Reed et al., 2012). The first is that stress induces the expression of

deleterious mutations that are silent under benign conditions (Kondrashov & Houle, 1994; Elena &

de Visser, 2003). For example in *Drosophila*, recessive alleles were linked to increased mortality

under temperature stress in inbred lines (Vermeulen & Bijlsma, 2004). Extreme stress and inbreeding

impede the function of heat shock proteins such as Hsp90, which are essential in buffering the

expression of deleterious mutations (Rutherford & Lindquist, 1998; Queitsch et al., 2002; Bergman

& Siegal, 2003). A second mechanism is that mutational load may lower stress resistance or tolerance,

by destabilizing cellular homeostasis under stress, or affecting tissue and genome repair after stress

exposure (Agrawal & Whitlock, 2010; Reed et al., 2012). For example, hybrids of *Arabidopsis*

thaliana expressed more metabolites from central pathways linked to higher freezing tolerance or disease resistance compared to inbred lines (Korn et al., 2010; Yang et al., 2015), and the regulation of stress-response pathways led to higher recovery after stress (Miller et al., 2015). The third mechanism is that stress increases phenotypic variance and with it the opportunity for relative fitness measures to differ more among than among outbred individuals (Waller et al., 2008). Reed *et al.* (2012) found that variation in inbreeding depression was predicted by stress itself and – to a weaker extent – also by increased phenotypic variance.

In this study, we tested for the stress dependence of the expression of mutational load at the range limits of North American *Arabidopsis lyrata* subsp. *lyrata* (L.). The species has been strongly impacted by the last glacial maximum (LGM), with more than half of its contemporary distribution being the result of rapid post-glacial range expansion from two separate refugia (Griffin & Willi, 2014; Willi et al., 2018). Both range expansion and rear-edge dynamics were found to be associated with an increase in presumably deleterious mutations assessed on the level of DNA sequences, leading to a current pattern of heightened deleterious mutations in range edge populations (Willi et al., 2018). Another study showed that this heightened genomic load was correlated with expressed load, estimated by heterosis – the fitness increase of between- compared to within-population crosses (Perrier et al., 2020). Here we tested whether the expression of mutational load, again estimated by heterosis, additionally depended on the interaction between genomic load and environmental stressfulness in the direction of a positive synergistic effect. Offsprings of within- (WPC) and between-population crosses (BPC) were raised in a latitudinal transplant experiment with five common garden sites within and beyond the distribution range (plant data as in Perrier *et al.* (2020)). Environmental stressfulness was estimated by (a) the differences in climatic conditions experienced by populations in each common garden relative to their site of origin, assuming local adaptation to climate (Hoffmann & Hercus, 2000), and (b) the relative performance of each population at a garden site compared to the best garden. Finally, we tested whether heterosis in range-edge populations – expressing the highest mutational load compared to populations from elsewhere of the range – was

enhanced when offspring of crosses were transplanted to environments beyond range limits (Lee-

Yaw et al., 2018).

Materials and Methods

Experimental design

Populations. Twenty populations of *A. lyrata* subsp. *lyrata* from across the species distribution were

selected for producing WPC and BPC (Fig. 1; Table S1). The populations represented the two main

ancestral clusters (Griffin & Willi, 2014; Willi et al., 2018), the latitudinal range, and variation in the

genomic signatures of mutational load (Willi et al., 2018). Our sampling also represented the two

mating systems: predominant outcrossing and predominant selfing (MO2, ON1, ON11). In the study

system, selfing populations are mainly restricted to the edges of distriubtion of the two ancestral

clusters (Griffin & Willi, 2014). They have an especially high genomic load, on the one hand because

of their expansion history, and on the other hand because of the mating system shift (Willi et al.,

2018; Perrier et al. 2020). Replicate selfing populations were included to increase the range of

genomic load found in the study system. Eighteen populations (nine per cluster) were considered

target populations in the assessment of expressed mutational load, and two were partner populations

for BPC (NY1, eastern cluster, and IA1, western cluster). Partner populations were selected because

of their core location in each genetic cluster, high genomic diversity and history of little genetic drift.

Seeds of individual plants had been collected in each population over an area of about 450 m^2.

Crossing. One plant of each of 26 randomly selected and presumably unrelated seed families

per population were raised under controlled, indoors conditions. Plants were randomly appointed to

"mothers"/pollen recipients (12), "fathers"/pollen donors (12), and backups (2). Each "mother" was

crossed using pollen from one "father" of the same population (WPC). For target populations, each

"mother" was additionally crossed using pollen from one "father" of the partner population (BPC).

Crosses were non-reciprocal. Pollination was performed at the bud stage to prevent unintended cross-

or self-pollination. We repeated each cross-combination to obtain enough healthy-looking seeds to

perform the transplant experiment (min. 60). In case of systematic cross failure, we replaced the father

plant with a backup plant. In total, 401 family-cross type combinations were used for the transplant

experiment (Table S2).

Transplant experiment. Five common garden sites (CG) were selected along a *ca.* 1400 km

latitudinal gradient crossing the distribution of *A. lyrata* in the eastern USA (Fig. 1; Table S3 with

WorldClim data v 2.0 (Fick & Hijmans, 2017)). The five sites represented the centre of the species'

range (CG3, Harrisonburg, VA), the southern and northern range edges (CG4, Winston-Salem, NC,

and CG2, Williamstown, MA, respectively), and areas beyond the edges (CG5, Athens, GA, and

CG1, in the Adirondacks, NY). In 2017, in each common garden, seeds of all crosses were sown in

pots (2 per pot; pot with opening at the bottom) randomly distributed within each of three spatial

blocks (more details in Perrier *et al.* (2020)). Five data loggers (iButton®, Maxim Integrated Products,

Inc) monitored air temperature every hour (1.5 m above ground, in the shadow) in each site for the

duration of the experiment.

Performance. Germination was recorded for each seed three days a week until peak

germination was over (four to five weeks) and afterwards once a week until thinning to one seedling

per pot. Reproductive output was assessed several weeks after peak flowering in 2018 and 2019, and

calculated for each individual as the sum of fruits, pedicels (flowers that did not develop into fruits),

flowers, and flower buds. Flowers that did not develop into fruits were included because they

contributed to the performance capacity of plants, and their pollen may have fertilized ovules on other

plants. However, seeds per fruit were not estimated as pollen transfer was not controlled for in the

experiment. During the second winter, we also assessed seed survival to estimate how seeds that did

not germinate the first fall could have contributed to population performance. For each population-

cross type combination, we pooled 100 seeds from five to twelve seed families, split them in 10 bags

and placed 2 bags per garden on bare ground next to pots in October 2018. Bags were retrieved in

late spring 2019 and visually screened for germinated seedlings and seeds. Remaining seeds were

tested for germination in the laboratory. Seed survival over the winter was calculated for each bag as

the fraction of germinated seedlings outdoors and in the lab.

Statistical analysis

Initial analyses were based on *Multiplicative performance* (*MP*) calculated on the level of the pot as

the product of the fraction of germinated seeds and the sum of total reproductive output recorded until

year 3 (year 2 for CG 3 that had to be given up early). We tested for a general effect of cross type

(BPC compared to WPC = 0), genomic load, environmental stress and interactions between all fixed

effects. Genomic load was a genomic estimate of mutational load calculated at the population level

as the ratio of non-synonymous polymorphic sites to synonymous polymorphic sites, adjusted for

their mean derived allele frequency (relative to *A. thaliana*), $P_n f_n / P_s f_s$ of Willi *et al.* (2018), calculated

for each population. Between-population crosses were assigned the genomic load of mother

population as all BPC shared the same pollen donor population within genetic cluster. Environmental

stress was depicted by the difference in climate between garden site and site of origin of a population.

Distribution modelling had shown that average minimum temperature in March and April described

best the niche and latitudinal range limits of *A. lyrata* (Lee-Yaw et al., 2018), and therefore,

environmental stress was calculated as the difference in average minimum temperature in early spring

(March and April) between common garden sites and site of origin of populations: $\Delta T_{\min} = T_{\min\,CG}$ -

$T_{\min\,origin}$. $T_{\min\,CG}$ was calculated based on records of temperature loggers (Table S4), and $T_{\min\,origin}$

based on the WorldClim 2.0 database (spatial resolutions of 30 seconds (Fick & Hijmans, 2017);

Table S1) and averaged over target and partner population for BPC. Positive and negative values of

$\Delta T_{\min}$ indicated transplanting toward warmer and colder temperatures, respectively. We assumed that

both positive and negative deviations could be perceived as stressful, resulting in a bell shaped

relationship between $\Delta T_{\min}$ and performance, and therefore included the square term of $\Delta T_{\min}$ in the

model. Evidence for a positive synergistic effect between the expression of genomic load – the

superior performance of between- to within-population crosses depending on genomic load – and the

environmental stressfulness was assumed to be given by positive three-way interaction coefficients.

Analyses were performed in a Bayesian framework on 10 parallel chains (detailed in Method S1),

with the package *MCMCglmm* (Hadfield, 2010; Hadfield, 2019) in *R* (R Core Team, 2021), and

accounted for the zero-inflation of *MP* by modelling a Gaussian process ($MP > 0$) and a logistic

process (probability of 1 assigned to $MP > 0$). Random effects were maternal plant nested within

maternal population and maternal population, for which variance in intercepts and slopes of cross

type were estimated, and block nested within common garden and common garden. Covariates were

mean-centred, and square terms were calculated based on mean-centred values.

Further analyses were – for ease of interpretation – performed on heterosis estimated on the

level of the population and addressed what the best model on the expression of load under

environmental stress was. Heterosis was calculated as the increase in performance of BPC relative to

WPC: $(W_{BPC} - W_{WPC})/W_{WPC}$, where W_{WPC} and W_{BPC} were population averages based on seed family

averages in a common garden. W_{WPC} was further averaged across target and partner population.

Heterosis was also assessed for the finite rate of increase per year, λ, from stage-classified matrices

(Caswell, 2001) based on mean performance of plants of each population-cross type-garden group,

over the three years (detailed in Perrier *et al.* (2020)). These matrices comprised the stages: (1) healthy

seeds in year 1, (2) reproducing individuals of year 2, (3) reproducing individuals of year 3, with a

one-year projection interval between each stage. Heterosis estimates were $\log_{10}$-transformed and

analysed by hierarchical mixed-effects models based on restricted maximum likelihood using the

packages *lme4* (Bates et al., 2015) and *LmerTest* (Kuznetsova et al., 2017) in *R*. Fixed effects were

the genomic estimate of mutational load of the mother population, environmental stress, and

interactions between them. Crossed random effects were population and common garden (detailed in

Method S2A). Here, ΔT_{min} was calculated with $T_{min\ origin}$ again averaged over target and partner

population.

To explore the effect of stress, we depicted it by either signed or absolute ΔT_{min}, included an

orthogonal quadratic term or not, and included interaction terms or not. We performed model

selection based on the corrected Akaike information criterion (AICc (Sugiura, 1978)). For the best

model, we tested the significance of each fixed effect with a likelihood-ratio χ^2 test using the *Anova*

function (type III) of the package *car* (Fox & Weisberg, 2019). We also used the same model to test

for effects on W_{WPC}, as the more interesting component of heterosis. A confounding role of genetic

incompatibility estimated by neutral genetic divergence, and of divergent adaptation between target

and partner populations estimated by environmental divergence could be excluded. Details on these

analyses are presented in Appendix S1.

We repeated the analysis on heterosis on a second estimate of stress, the relative decline in

population-level performance at a site relative to the best site (1 - W_{WPC}/W_{max}) (Reed et al., 2012).

Stress varied between 0 (conditions leading to maximal performance) and 1 (conditions with W_{WPC} =

0). Relative performance was then averaged over both target and partner populations.

Finally, we testeds whether heterosis was higher when range-edge populations (with highest

load) were transplanted beyond the range limits compared to sites at the range limits. We assessed

this effect in two distinc dataset: one considering the six northern edge populations (ON8, ON11,

ON12, NY4, NY5 and NY6; in blue in Fig. 1) raised at the northern edge site (CG2) and beyond the

northern edge (CG1), and one considering the six southern edge populations (MO1, MO2, IN1, NC2,

NC4 and VA1; in red in Fig. 1) raised at the southern edge site (CG4) and beyond the southern edge

(CG5). For both datasets, $\log_{10}$-transformed heterosis was tested in hierarchical mixed-effect models

based on restricted maximum likelihood. Transplant was a fixed effect (edge = 0) and population was

a random effect (detailed in Method S2B).

Results

Analyses on the level of the pot revealed a significant positive effect of cross type on performance in

both the log-normal ($MP > 0$) and a logistic (probability of 1 assigned to $MP > 0$) process (Table 1).

Performance was higher in between- compared to within-population crosses (BPC and WPC = 0),

supporting a general heterosis effect. Performance was also significantly negatively affected by

genomic load and by a positive cross type-by-genomic load interaction in the log-normal process. Performance generally declined with increasing genomic load, but less in BPC than WPC. Performance was further significantly negatively affected by the linear and quadratic term of stress (ΔT_{min} and ΔT_{min}^2) in both the log-normal and logistic process. Performance generally declined when conditions in the common garden diverged from conditions at the site of origin, with a stronger decline under warmer conditions ($\Delta T_{min} > 0$). Furthermore, the interaction cross type-by-ΔT_{min} was significantly positive for the logistic process, indicating that the decline due to stress was stronger for WPC compared to BPC. Finally, the logistic process determining performance was significantly affected by a negative interaction between genomic load and ΔT_{min}. Performance of populations with higher load declined more when exposed to warmer temperatures, but less when exposed to colder temperatures.

Results on population-level heterosis based on MP or λ overlapped with individual-level analysis to the extent that genomic load and the linear term of stress were significant and shared a positive associated with heterosis (Table 1, Fig. 2a). Heterosis was highest in populations with more mutational load, and in populations raised in conditions warmer than at their site of origin ($\Delta T_{min} > 0$). However, neither the quadratic term of stress nor the interactions between stress and genomic load were significant. Heterosis based on MP and λ in the five common gardens ranged from -0.96 to 23.50 (mean: 1.88) and -0.53 to 7.29 (mean: 0.73) respectively (Table S5).

Model selection revealed the best parameters to predict population-level heterosis (Table S6). Among the models compared, the one presented just before – with genomic load, the signed term of stress (ΔT_{min}), its square term, and interactions between genomic load and stress – was the best. The next best model was the one without interactions. Results clearly indicated that heterosis could be predicted better if the climate deviation between garden site and site of origin of populations included information on whether conditions were cooler or warmer, if climate was depicted not only as a linear term, and if interactions between stress and genomic load were included. Similarly to analysis of heterosis, its component of W_{WPC} revealed an effect of genomic load, ΔT_{min} and ΔT_{min}^2 (Table S7),

but no effect of the genomic load-by-ΔT_{min} interaction. The performance of WPC declined with

increasing genomic load, and – in an accelerating manner – with warmer conditions compared to the

site of origin of populations (Fig. S1). The model-predicted maximal decline in WPC performance

for a population with average genomic load was 92% when exposed to warmer temperatures,

(baseline of $\Delta T_{min} = 0$; Table S8).

Analyses on the effect of stress expressed by performance decline on heterosis revealed

similar results as those based on ΔT_{min}. The best model predicting heterosis considered again the

linear and quadratic term for stress and interactions between stress and genomic load (Table S9). The

second model was the one without interactions. Heterosis was significantly positively affected by

genomic load (marginal for λ), and both the linear (except *MP*) and the quadratic term of stress (Table

S10), but again was unaffected by interactions. Heterosis based on *MP* increased almost

symmetrically toward low ($W_{WPC} = W_{max}$) and high stressfulness ($W_{WPC} = 0$; Fig. S2), while heterosis

based on λ increased more strongly toward high stressfulness (Fig. 2b).

Finally, heterosis of range edge populations did not significantly differ between the garden

sites at the range edge and the garden sites beyond the range, neither for the southern nor for the

northern range edge (Table S11, Fig. S3).

Discussion

Recent evolutionary theory proposes that species' range limits may result from high mutational load

of range-edge populations due to prolonged exposure to genetic drift caused by past range expansion

or rear-edge demographics (Willi, 2019). Here we tested whether the expression of such load is

enhanced by environmental stress in a large-scale transplant experiment with *Arabidopsis lyrata*. To

express load, we used heterosis based on multiplicative performance of between- compared to within-

population crosses, assuming that load was mainly due to recessive deleterious mutations. To estimate

stress, we used divergent thermal conditions during a critical part of the year between gardens and

sites of origin of populations, or relative performance decline of each population. Expressed

mutational load was stronger in populations with higher deleterious counts inferred from coding DNA

sequences, as previously found (Willi et al., 2018; Perrier et al., 2020). However, stress did not clearly

synergistically interact with genomic load to enhance heterosis. Nonetheless, we found a general

effect of stress on heterosis linked to a stronger performance decline under warmer conditions than

at sites of origin of populations. Our study therefore suggests that a general heterozygote deficit is

more detrimental under temperature stress, and that interspecific outcrossing can alleviate this effect.

A main result of this study is that expressed mutational load estimated by heterosis was not

significantly enhanced under thermal stress. Indeed, while analyses on pot-level performance were

supportive of an effect of the cross type, genomic load and environmental stress on performance,

these analyses did not support a three-way interaction between effects. This suggests that

environmental stressfulness does not enhance the effect of the higher accumulation of recessive

deleterious mutations on the expression of mutational load. Analyses on population-level heterosis

were not clearly supportive of an interaction either. Even though the model including interactions

between stress and genomic load was better supported than the model without interactions, the

interaction was not significant in the χ^2 test. This suggest that some variation in heterosis may be the

result of an interaction, but is too much variation. Such environment dependence has often been

evoked due to the similarities in the underlying genetics between heterosis (if caused by dominance)

and inbreeding depression (Fenster & Galloway, 2000; Oakley et al., 2015). Indeed, a previous meta-

level study on the stress dependence of heterosis found that the benefit of outcrossing for small and

inbred populations was considerably higher under stress (Frakham, 2015), however the study did not

include large populations for comparison. Some of them too may show a heterotic effect under

outbreeding that is enhanced under stress, as found in our study.

We think that our result for *A. lyrata* is robust despite caveats that were previously mentioned

for the study of environment dependence of load. It was argued that environment dependence of the

expression of load depends on the strength of stress (Fox & Reed, 2010). In our study, exposure to

warmer conditions strongly reduced WPC population performance, by up to 92%. Type and novelty

of stress were also suggested to affect stress dependence of the expression of load (Willi et al., 2007a; Sandner & Matthies, 2016). If *A. lyrata* had regularly experienced conditions we considered stressful here across its evolutionary history, deleterious alleles expressed under these conditions may have already been purged (Hedrick, 1994; Bijlsma et al., 1999; Enders & Nunney, 2016). However, the expression of mutational load was also not enhanced by stress based on relative performance, a broader measure of stress likely depicting also novel stressors. Furthermore, the interaction between genomic load and stress could have been masked by outbreeding depression, previously reported to counteract heterosis under environmental stress such as drought (Prill et al., 2014). Outbreeding depression is expected to be stronger in crosses involving populations of highest divergence (Oakley et al., 2015; Lynch, 1991). In our study, partner populations came from the centers of geographic distribution and were unlikely highly divergent, given the shared postglacial history and moderate climatic differences between populations. Furthermore, the high performance gain especially in populations with high load speaks in favor of an overwhelming positive effect of heterosis and little action of outbreeding depression. For these reasons we conclude that the lack of environment dependence of mutational load seems real.

Heterosis was nonetheless affected by stress in our study, especially warm stress. Heterosis increased under warmer conditions than at the sites of origin of populations (Fig. 2). Perrier *et al.* (2020) previously linked patterns of heightened heterosis with increasing genomic load to a performance decline in WPC, with BPC being fairly constant across that gradient. Here we observed that WPC and BPC performance generally declined with increasing warm stress, but less in BPC, as suggested by the pot-level analys on performance. Lower sensitivity to stress in outbred plants could result from recessive deleterious mutations being frequently heterozygous and their better buffering under stress by *e.g.* heat shock proteins (Rutherford & Lindquist, 1998; Queitsch et al., 2002; Bergman & Siegal, 2003). Higher performance of heterotic hybrids under stress in *A. thaliana* has been linked to disruptions of stress response pathways in inbred parental lines (Korn et al., 2010; Yang et al., 2015; Miller et al., 2015). Previous research in the context of agriculture or genetic rescue

has also provided many examples the effect of heterosis increased under stress (Frakham, 2015; Blum, 2013; Fujimoto et al., 2018). Here, the overall increase in heterosis under warm stress independent of genomic load suggests that most populations of *A. lyrata* suffer from a general heterozygote deficit whose expression is heat-sensitive.

Our results offer some important insights to the understanding of heterosis and its stress-dependence. First, model selection clearly showed that the effect of increasing stressfulness was not linear but better described by a positive accelerating relationship. This was the case both when stress was expressed as the difference in temperature regime during a critical part of the year for *A. lyrata* between common garden and site of origin of populations, or as the performance decline in a garden by populations. Second, when stress was depicted by temperature difference, it was the unsigned temperature difference that depicted stress better. While cooler conditions led to relatively high plant performance, it was the warmer conditions that were stressful and that led to higher heterosis. Third, even though interactions were not significant between genomic load and stressfulness on heterosis, our study suggests that they may still be of some relevance. One indication is that model selection pointed to the inclusion of interactions being a better model predicting heterosis. Another indication is that pot-level performance revealed significant negative interactions between genomic load and the linear term of stress, suggesting that populations with high genomic load performed worse under increasingly warmer conditions than at site of origin, independent of cross type.

Our study has two main implications beyond the study of heterosis. The first is the role of expressed mutational load for range limits. Populations expanding beyond range limits may suffer from a twofold genetic Allee effect (Luque et al., 2016). A first is linked to the process of range expansion itself: in the absence of strong environmental gradients, range expansion may lead to the accumulation of high mutational load in population at range limits (Peischl et al., 2013; Peischl et al., 2015; Gilbert et al., 2017). Further range expansion originating from these range-edge population may therefore lead to an even higher accumulation of mutational load, beyond tolerable levels to maintain population demographic rates (Perrier et al., 2020). A second is due to a general

heterozygote deficit expressed under stressful, particularly warm-stress conditions. However, the result was not confirmed at range limits. Northern and southern range-edge populations did not significantly differ in heterosis when transplanted at their respective range edge or beyond. Nontheless, the stress-dependent genetic Allee effect could be triggered by extreme environmental events more frequent beyond range limits, and by long range dispersal into less suitable habitats than those immediately beyond range limits. Future simulation studies on range dynamic (Gilbert et al., 2017; Polechová, 2018) should therefore consider the consequences of stress-dependent genetic Allee effects on range expansion.

The second implication of our study is that interpopulation outbreeding can have a generally positive effect across populations of the distribution of a species. A great majority of populations indeed benefitted from outbreeding, especially under warmer conditions than what the populations had experienced in the past. Such heterotic effects can often be beneficial beyond the first generation (Willi et al., 2007b; Frakham, 2016). Our results therefore motivate the study of assisted gene flow in the context of climate warming, to preserve species across the entire range. Such management may worth being tested particularly for weak dispersers, potentially combined with the use of climate-adjusted provenances such as explored in restoration ecology (Prober et al., 2015). Populations at the southern range limits, with high genomic load, may also need to be supplemented with material from similar climatic conditions but different demographic history, to maximize heterozygote advantage and ensure the local persistence of species.

References

Agrawal AF, Whitlock MC. 2010. Environmental duress and epistasis: how does stress affect the strength of selection on new mutations. Trends in Ecology and Evolution 25: 450–458.

Armbruster P, Reed DH. 2005. Inbreeding depression in benign and stressful environments. Heredity 95: 235–242.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67: 1–48.

Bergman A, Siegal ML. 2003. Evolutionary capacitance as a general feature of complex gene networks. Nature 424: 549–52.

Bijlsma R, Bundgaard J, Van Putten WF. 1999. Environmental dependence of inbreeding depression and purging in *Drosophila melanogaster*. Journal of Evolutionary Biology 12: 1125–1137.

Blum A. 2013. Heterosis, stress, and the environment: a possible road map towards the general improvement of crop yield. Journal of Experimental Botany 64: 4829–4837.

Brown JH. 1984. On the relationship between abundance and distribution of species. The American Naturalist 124: 255–279.

Cahill AE, Aiello-Lammens ME, Fisher-Reid MC, Hua X, Karanewsky CJ, Ryu HY, Sbeglia CG, Spagnolo F, Waldron JB, Wien JJ. 2014. Causes of warm-edge range limits: systematic review, proximate factors and implications for climate change. Journal of Biogeography 41: 429–442.

Caswell H. 2001. Matrix population models, 2nd edn. Sunderland, MA, USA: Sinauer Associates

Connallon T, Sgrò CM. 2018. In search of a general theory of species' range evolution. PLoS Biology 16: e2006735.

Elena SF, de Visser JAGM. 2003. Environmental stress and the effects of mutation. Journal of Biology 2: 12.

Enders LS, Nunney L. 2016. Reduction in the cumulative effect of stress-induced inbreeding depression due to intragenerational purging in *Drosophila melanogaster*. Heredity 116: 304–313.

Fenster CB, Galloway LF. 2000. Population differentiation in an annual legume: genetic architecture. Evolution 54: 1157–1172.

Fick SE, Hijmans RJ. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology 37: 4302–4315.

Fox CW, Reed DH. 2011. Inbreeding depression increases with environmental stress: an experimental study and meta-analysis. Evolution 65: 246–258.

Fox J, Weisberg S. 2019. An R Companion to Applied Regression, 3rd edn. Thousand Oaks, CA, USA: Sage.

Frankham R. 2015 Genetic rescue of small inbred populations: meta-analysis reveals large and consistent benefits of gene flow. Mol. Ecol. 24, 2610–2618.

Frankham R. 2016 Genetic rescue benefits persist to at least the F3 generation, based on a meta-analysis. Biol. Conserv. 195, 33–36.

Fujimoto R, Uezono K, Ishikura S, Osabe K, Peacock WJ, Dennis ES. 2018. Recent research on the mechanism of heterosis is important for crop and vegetable breeding systems. Breeding Science 68: 145–158.

Gaston K. 2009. Geographic range limits of species. Proceedings of the Royal Society B: Biological Sciences 276: 1391–1393.

Gilbert KJ, Sharp NP, Angert AL, Conte GL, Draghi JA, Guillaime F, Hargreaves AL, Matthey Doret R, Whitlock MC. 2017. Local adaptation interacts with expansion load during range expansion: maladaptation reduces expansion load. The American Naturalist 189: 368–380.

González-Martínez SC, Ridout K, Pannell JR. 2017. Range expansion compromises adaptive evolution in an outcrossing plant. Current Biology 27: 2544–2551.

Griffin PC, Willi Y. 2014. Evolutionary shifts to self-fertilisation restricted to geographic range margins in North American *Arabidopsis lyrata*. Ecology Letters 17: 484–490.

Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. Journal of Statistical Softwares 33: 1–22.

Hadfield JD. 2019. MCMCglmm course notes. [WWW document] URL https://cran.r-
project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf [accessed 24 June 2020].

Hampe A, Petit RJ. 2005. Conserving biodiversity under climate change: the rear edge matters.
Ecology Letters 8: 461–467.

Hargreaves AL, Samis KE, Eckert CG. 2014. Are species' range limits simply niche limits writ large?
A review of transplant experiments beyond the range. The American Naturalist 183: 157–173.

Hedrick PW. 1994. Purging inbreeding depression and the probability of extinction: full-sib mating.
Heredity 73: 363–372.

Henry RC, Barton KA, Travis JMJ. 2015. Mutation accumulation and the formation of range limits.
Biology Letters 11: 20140871–20140871.

Hoffmann AA, Hercus MJ. 2000. Environmental stress as an evolutionary force. BioScience 50: 217–
226.

Kondrashov AS, Houle D. 1994. Genotype-environment interactions and the estimation of the
genomic mutation rate in *Drosophila melanogaster*. Proceedings of the Royal Society B:
Biological Sciences 258: 221–227.

Korn M, Gärtner T, Erban A, Kopka J, Selbig J, Hincha DK. 2010. Predicting *Arabidopsis* freezing
tolerance and heterosis in freezing tolerance from metabolite composition. Molecular Plant 3:
224–235.

Koski MH, Layman NC, Prior CJ, Busch JW, Galloway LF. 2019. Selfing ability and drift load evolve
with range expansion. Evolution Letters 3: 500–512.

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest Package: tests in linear mixed-
effects models. Journal of Statistical Software 82: 1–26.

Lee-Yaw JA, Kharouba HM, Bontrager M, Mahony C, Csergő AM, Noreen AM, Li Q, Schuster R,
Angert AL. 2016. A synthesis of transplant experiments and ecological niche models suggests
that range limits are often niche limits. Ecology Letters 19: 710–722.

Lee-Yaw JA, Fracassetti M, Willi Y. 2018. Environmental marginality and geographic range limits:
a case study with *Arabidopsis lyrata* ssp. *lyrata*. Ecography 41: 622–634.

Lohmueller KE, Indap AR, Schmidt S, Boyko AR, Hernandez RD, Hubisz MJ, Sninsky JJ, White TJ,
Sunyaev SR, Nielsen R, et al. 2008. Proportionally more deleterious genetic variation in
European than in African populations. Nature 451: 994–998.

Luque GM, Vayssade C, Facon B, Guillemaud T, Courchamp F, Fauvergue X. 2016. The genetic
Allee effect: a unified framework for the genetics and demography of small populations.
Ecosphere 7: e01413.

Lynch M. 1991. The genetic interpretation of inbreeding depression and outbreeding depression.
Evolution 45: 622–629.

Miller M, Song Q, Shi X, Juenger TE, Chen ZJ. 2015. Natural variation in timing of stress-responsive
gene expression predicts heterosis in intraspecific hybrids of *Arabidopsis*. Nature
Communications 6: 7453.

Oakley CG, Ågren J, Schemske DW. 2015. Heterosis and outbreeding depression in crosses between
natural populations of *Arabidopsis thaliana*. Heredity 115: 73–82.

Peischl S, Excoffier L. 2015. Expansion load: recessive mutations and the role of standing genetic
variation. Molecular Ecology 24: 2084–2094.

Peischl S, Dupanloup I, Kirkpatrick M, Excoffier L. 2013. On the accumulation of deleterious
mutations during range expansions. Molecular Ecology 22: 5972–5982.

Peischl S, Kirkpatrick M, Excoffier L. 2015. Expansion load and the evolutionary dynamics of a
species range. The American Naturalist 185: E81–E93.

Peischl S, Dupanloup I, Foucal A, Jomphe M, Bruat V, Grenier JC, Gouy A, Gilbert KJ, Gbeha E,
Bosshard L, et al. 2018. Relaxed selection during a recent human expansion. Genetics 208: 763–
777.

Perrier A, Sánchez-Castro D, Willi Y. 2020. Expressed mutational load increases toward the edge of
a species' geographic range. Evolution 74: 1711–1723.

Pironon S, Papuga G, Villellas J, Angert AL, García MB, Thompson JD. 2017. Geographic variation
in genetic and demographic performance: new insights from an old biogeographical paradigm.
Biological Reviews 92: 1877–1909.

Polechová J. 2018. Is the sky the limit? On the expansion threshold of a species range. PLoS Biology
16: e2006735.

Prill N, Bullock JM, van Dam NM, Leimu R. 2014. Loss of heterosis and family-dependent
inbreeding depression in plant performance and resistance against multiple herbivores under
drought stress. Journal of Ecology 102: 1497–1505.

Prober SM, Byrne M, McLean EH, Steane DA, Potts BM, Vaillancourt RE, Stock WD. 2015.
Climate-adjusted provenancing: a strategy for climate-resilient ecological restoration. Frontiers
in Ecology and Evolution 3: 65.

Queitsch C, Sangster TA, Lindquist S. 2002. Hsp90 as a capacitor of phenotypic variation. Nature
417: 618–624.

R Core Team. 2021. R: a language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Reed DH, Fox CW, Enders LS, Kristensen TN. 2012. Inbreeding–stress interactions: evolutionary
and conservation consequences. Annals of the New York Academy of Sciences 1256: 33-48.

Roff DA. 1997. Evolutionary Quantitative Genetics. New York, NY, USA: Chapman and Hall.

Roy K, Hunt G, Jablonski D, Krug AZ, Valentine JW. 2009. A macroevolutionary perspective on
species range limits. Proceedings of the Royal Society B: Biological Sciences 276: 1485–1493.

Rutherford SL, Lindquist S. 1998. Hsp90 as a capacitor for morphological evolution. Nature 396:
336–342.

Sandner TM, Matthies D. 2016. The effects of stress intensity and stress type on inbreeding
depression in *Silene vulgaris*. Evolution 70: 1225–1238.

Sexton JP, McIntyre PJ, Angert AL, Rice KJ. 2009. Evolution and ecology of species range limits.
Annual Review of Ecology, Evolution, and Systematics 40: 415–436.

Sugiura N. 1978. Further analysts of the data by akaike' s information criterion and the finite
corrections. Communications in Statistics - Theory and Methods 7: 13–26.

Vermeulen CJ, Bijlsma R. 2004. Characterization of conditionally expressed mutants affecting age-
specific survival in inbred lines of *Drosophila melanogaster*: lethal conditions and temperature-
sensitive periods. Genetics 167: 1241–1248.

Waller DM, Dole J, Bersch AJ. 2008. Effects of stress and phenotypic variation on inbreeding
depression in *Brassica rapa*. Evolution 62: 917–931.

Willi Y. 2019. The relevance of mutation load for species range limits. American Journal of Botany
106: 757–759.

Willi Y, Van Buskirk J. 2019. A practical guide to the study of distribution limits. The American
Naturalist 193: 773–785.

Willi Y, Dietrich S, Kleunen MV, Fischer M. 2007a. Inter-specific competitive stress does not affect
the magnitude of inbreeding depression. Evolutionary Ecology Research 9: 959–974.

Willi Y, Kleunen MV, Dietrich S, Fischer M. 2007b. Genetic rescue persists beyond first-generation
outbreeding in small populations of a rare plant. Proceedings of the Royal Society B: Biological
Sciences 274: 2357–2364.

Willi Y, Fracassetti M, Zoller S, Van Buskirk J. 2018. Accumulation of mutational load at the edges
of a species range. Molecular Biology and Evolution 35: 781–791.

Yang L, Li B, Zheng XY, Li J, Yang M, Dong X, He G, An C, Deng XW. 2015. Salicylic acid
biosynthesis is enhanced and contributes to increased biotrophic pathogen resistance in
Arabidopsis hybrids. Nature Communications 6: 7309.

Zhang M, Zhou L, Bawa R, Suren H, Holliday JA. 2016. Recombination rate variation, hitchhiking,
and demographic history shape deleterious load in poplar. Molecular Biology and Evolution 33:
2899–2910.

Table 1: Results of model testing for the effect of cross type (between- compared to within-population crosses [0]), genomic estimate of mutational load, environmental stress ($\Delta T_{min} = T_{min\ CG} - T_{min\ origin}$) and their interactions on *multiplicative performance* (*MP*), or on heterosis based on *MP* and the finite rate of increase (λ), at the five common garden sites

Performance	Cross type (CT) Mean	Genomic load (GL) Mean	Stress (S) Mean	S^2 Mean	CT * GL Mean	CT * S Mean	CT * S^2 Mean	GL * S Mean	GL * S^2 Mean	CT * GL * S Mean	CT * GL * S^2 Mean	
Log-normal												
MP	**0.15 ***	**-1.32 ***	**-16.97 ***	**-8.14 ****	**1.50 ***	4.92	-2.27	-69.43	-14.63	24.35	38.02	†, ‡
Logistic												
MP	**0.63 *****	-2.58	**-79.65 ****	**-58.51 *****	2.76	**65.64 *****	15.35	**-385.58 *****	60.68	-161.89	-30.10	‡
Heterosis		β χ²	β χ²	β χ²				β χ²	β χ²			
MP		**1.92 6.63 ****	**1.03 5.77 ***	-0.73 2.98 (*)				-3.21 0.48	3.72 0.70			†, ‡
λ		**0.84 5.36 ***	**0.79 14.78 *****	-0.07 0.15				0.22 0.01	1.68 0.70			†

In the models testing multiplicative performance (MP, $\log_{10}$-transformed if > 0), MP was assumed to follow a Gaussian distribution with 0-inflation. Therefore, the model assessed all fixed and random effects for their importance in both the logistic process (binary variable depicting germination combined with survival and the capacity to initiate flowering) and the Gaussian process (total number of flowers during one or two reproductive seasons). For this model, estimates of coefficients are modes of MCMC samples from the posterior distribution of parameters (*mean*). The logistic part of the model predicts non-zeros in the distribution on the logit scale. In the model testing population heterosis, estimates of population heterosis were $\log_{10}$-transformed prior to analysis. Stress, ΔT_{min}, was an average over target and partner population. Each model testing heterosis was optimized with the *bobyqa* optimizer to improve convergence. Test statistics include regression coefficient (β) and the χ^2-value of each fixed effect. For heterosis based on MP and on λ, the marginal R^2 of the model was of 0.18 and 0.23 respectively, and the conditional R^2 of the model was of 0.37 and 0.46 respectively. Genomic load and stress were standardized prior to analyses (mean = 0). Model fits with significant (positive) intercept are indicated by †. Estimates and regression coefficients with P-values < 0.05 are written in bold; significance is indicated: (*) $P < 0.1$, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Results for random effects are not shown. For one of the five common gardens (CG3), the experiment stopped early and variables consider performance to year 2 only (indicated by ‡).

  Figures:

Figure 1: Distribution of *Arabidopsis lyrata* in eastern North America, with information on habitat suitability, and the location of the 20 populations studied and the 5 common garden sites. The range of *A. lyrata* is represented by the dotted line, habitat suitability by shades of blue, with darker blue indicating higher suitability (Lee-Yaw et al., 2018). Populations are shown by circles, with abbreviations for state (USA) or province (Canada) and a number (Willi et al., 2018). Blue and red circles represent northern- and southern-edge populations in our analysis. Green triangles represent the five common garden (CG) sites; numbers added to labels are in sequence of north to south. State outlines for the USA are shown, and the split between eastern and western genetic cluster is represented by the dashed line. Of the 20 populations, two were used as partner-populations for between-population crosses, NY1 for crosses with eastern populations, and IA1 for crosses with western populations.

Figure 2: Stress dependence of the expression of mutational load estimated by heterosis. Population heterosis was estimated based on the finite rate of increase, λ, of within-population crosses and between-population crosses with a partner population. Stress was depicted as either the difference in minimum temperature in early spring between common garden and the site of origin of a population ($\Delta T_{min} = T_{min\ CG} - T_{min\ origin}$; positive values to the right of the vertical dashed line indicate a warmer environment) **(a)**, or the relative decline in population-level performance W in a garden relative to the garden of highest performance for that population ($1 - W_{WPC}/W_{max}$) **(b)**. For both stress estimates, means of target and partner populations were taken. The horizontal dashed lines indicates when heterosis drops below 0 and outbreeding depression dominates. The black lines represent model-predicted relationships between heterosis and environmental stress (test statistics in Tables 1, Table S10; * $P < 0.05$, *** $P < 0.001$). Genomic mutational load, the ratio of genome-wide non-synonymous to synonymous polymorphic sites adjusted by their mean derived frequency, $P_n f_n/P_s f_s$ (Willi et al., 2018), is represented in shades of yellow (low) to red (high).