Intact reading ability in spite of a spatially distributed visual word form 'area' in an individual born without the left superior temporal lobe

Authors

Jin Li[1], Evelina Fedorenko[2], Zeynep Saygin[1*]

Affiliations

[1]Department of Psychology, The Ohio State University, Columbus, OH, USA.

[2]Department of Brain and Cognitive Sciences / McGovern Institute for Brain Research, MIT, Cambridge, MA, USA

[*]Correspondence to: li.9361@osu.edu and saygin.3@osu.edu

Acknowledgements

We would like to thank our participant – EG – for her time and willingness to participate in brain research. We thank Greta Tuckute, Alexander Paunov, and Benjamin Lipkin for their help in coordinating the collaboration as well as their suggestions for data analysis. We also thank Idan Blank and Zachary Mineroff for help with EG's fMRI and behavioral data collection and members of Saygin Developmental Cognitive Neuroscience Lab for helping with control data collection. We would like to acknowledge the Athinoula A. Martinos Imaging Center at the McGovern Institute for Brain Research, MIT and the support from Center for Cognitive and Behavioral Brain Imaging (CCBBI) and Ohio Supercomputer Center (OSC). E.F. was supported by NIH awards R01 DC016607, R01 DC016950, and research funds from the McGovern Institute for Brain Research and the Department of Brain and Cognitive Sciences at MIT. Z.M.S. was supported by the Alfred P. Sloan Foundation and OSU's Arts & Sciences College and Chronic Brain Injury initiative.

Conflict of interest

The authors declare no competing financial interests.

Abstract

The visual word form area (VWFA) is an experience-dependent brain region in the left ventral temporal cortex of literate adults that responds selectively to visual words. Why does it emerge in this stereotyped location? Past research has shown that the VWFA is preferentially connected to the left-lateralized frontotemporal language network. However, it remains unclear whether the presence of a typical language network and its connections with ventral temporal cortex (VTC) are critical for the VWFA's emergence, and whether alternative functional architectures may support reading ability. We explored these questions in an individual (EG) born without the left superior temporal lobe but exhibiting normal reading ability. Using fMRI, we recorded brain activation to visual words, objects, faces, and scrambled words in EG and neurotypical controls. We did not observe word selectivity either in EG's right homotope of the VWFA (rVWFA)—the most expected location given that EG's language network is right-lateralized—or in her spared left VWFA (lVWFA), in the presence of typical face selectivity in both the right and left fusiform face area (rFFA, lFFA). Interestingly, multivariate pattern analyses revealed voxels in EG's rVWFA and lVWFA that showed 1) higher within- than between-category correlations for words (e.g., Words-Words>Words-Faces), and 2) higher within-category correlations for words than other categories (e.g., Words-Words>Faces-Faces). These results suggest that a typical left-hemisphere language network may be necessary for the emergence of focal word selectivity within ventral temporal cortex, and that orthographic processing may depend on a distributed neural code, which appears capable of supporting reading ability.

Keywords: lesion study; Visual Word Form Area (VWFA); brain plasticity; word selectivity; multivariate pattern analysis (MVPA); category selectivity

Introduction

In the past two decades, numerous regions in the ventral temporal cortex (VTC) have been identified and characterized that respond selectively to different high-level visual categories (e.g., faces: Kanwisher et al., 1997; places: Epstein & Kanwisher, 1998; bodies: Downing et al., 2001; for review, see Kanwisher, 2010). What are the origins of these specialized regions? How do human brains develop this functional topography? Some have hypothesized that the functional organization of the VTC may be innate and related to the evolutionary importance of certain high-level visual categories. Indeed, face perception and recognition abilities appear to be heritable (Wilmer et al., 2010; Zhu et al., 2010), and 4-6 month-old infants already show face- and place-responsive areas within expected locations in the VTC (Deen et al., 2017). A related hypothesis is that pre-existing biases for certain visual attributes (e.g., retinotopy, Hasson et al., 2002; Malach et al., 2002; rectilinearity, Nasr et al., 2014) or perceptual dimensions (e.g., real-world size and animacy, Konkle & Caramazza, 2013) may predispose a brain region to become selective for particular visual categories. However, evolutionary pressures cannot explain the existence of a brain region that specializes for orthography (Hannagan et al., 2015)—the visual word form area (VWFA). The VWFA is a small region in the left lateral VTC that shows strong selectivity for visual words and letter strings in literate individuals (e.g., L. Cohen et al., 2003; Dehaene & Cohen, 2011; Hamamé et al., 2013). Perhaps surprisingly—given its late emergence—the VWFA is located in approximately the same location across individuals and languages (Baker et al., 2007). What sets word forms apart from other visual categories and why does the VWFA land in this stereotyped location?

One compelling possibility is that the specialization of category-selective regions in the VTC is

constrained by their differential connectivity to the rest of the brain (the Connectivity Hypothesis;

Mahon & Caramazza, 2011; Martin, 2006). Indeed, previous work showed that category-

selective responses can be predicted from both structural and functional connectivity (Osher et

al., 2015; Saygin et al., 2012); further, these distinct connectivity patterns may already exist at

birth and may drive future functional specialization (e.g., newborns show functional connectivity

differences between lateral VTC which houses the VWFA vs. medial VTC: Barttfeld et al.,

2018). Thus, written words may be processed in a stereotyped region within the left VTC due to

this region's pre-existing connectivity with the left-lateralized language network (e.g., Behrmann

& Plaut, 2013; Dehaene et al., 2015; Martin, 2006). This network consists of left lateral frontal

and lateral temporal areas and selectively supports language comprehension and production (e.g.,

Fedorenko et al., 2011).

Consistent with this idea, a number of studies have reported both anatomical and functional

connections between the VWFA and the language network in neurotypical adults. For example,

compared to the adjacent FFA, the VWFA shows stronger anatomical connectivity to the left

superior temporal, anterior temporal, and inferior frontal areas (perisylvian putative language

regions) (Bouhali et al., 2014). Several candidate white matter fascicles may serve to connect the

VWFA with frontal and superior temporal language cortex (Wandell et al., 2012; Yeatman &

Feldman, 2013). Similarly, using resting-state functional connectivity, Stevens et al. (2017)

found that the individually defined VWFA connects to the posterior left inferior frontal gyrus as

well as the left planum temporale, both part of the distributed left language network.

Moreover, a longitudinal study in children showed that the location of the VWFA could be

successfully predicted by its connectivity patterns at the pre-reading stage (before any word

selectivity within the VTC is observed), and that the most predictive connections of the relevant

area of the VTC were with the frontal and temporal language areas (Saygin et al., 2016). Even

more strikingly, this pattern of preferential connectivity appears to be already present in neonates

(Li et al., 2020). These connections between the VWFA and the frontotemporal language

network appear to be functionally important such that their damage or stimulation leads to

reading difficulties. For example, a case study of a child with a missing arcuate fasciculus (AF,

presumably connecting the VTC and other parts of the temporal cortex to parietal and frontal

areas; Wandell et al., 2012) found impaired reading ability (Rauschecker et al., 2009). Similarly,

a lesion in the white matter ventral to the angular gyrus resulted in alexia without agraphia,

presumably disrupting the VWFA's connections with the lateral temporal language areas

(Greenblatt, 1976).

A different way to assess the importance of a language network in developing visual word

selectivity is to ask whether language regions and the VWFA occupy the same hemisphere. In

the majority of individuals, the language regions and the VWFA co-lateralize to the left

hemisphere (LH) (e.g., Cai et al., 2010; Gerrits et al., 2019). In rare instances where neurotypical

individuals show right-hemispheric (RH) language dominance, VTC activation during reading

tasks also tends to be right-lateralized (e.g., Cai et al., 2008; Van der Haegen et al., 2012).

Another population where the language network is right-lateralized are individuals with

congenital or early left hemisphere (LH) damage (e.g., Asaridou et al., 2020). In the presence of

early LH damage, linguistic abilities tend to develop normally (e.g., Newport et al., 2017; Staudt,

2010; Staudt et al., 2001; see François et al., 2021 for a review). However, little is known about

the effects of early LH damage on reading ability and on the neural architecture of visual word

processing. In particular, if the left VTC is completely deafferentiated from the downstream LH

language cortex at birth, does the VWFA emerge in the right VTC when the language network is

forcibly remapped to the right hemisphere, or does it still emerge in the left VTC, due to some

pre-existing bias (e.g., innate connectivity with any spared LH cortex)? Indeed, some previous

studies show that language processing and reading may engage opposite hemispheres (e.g., Van

der Haegen et al., 2012). Or—perhaps even more drastically—does the VWFA look atypical

(e.g., smaller or less functionally selective), whatever hemisphere it emerges in? Do these

potential differences affect selectivity for other high-level visual categories in the VTC? And if

the VWFA manifests atypically but reading ability is normal, what does this tell us about the

implementation-level flexibility with respect to orthographic processing?

Here we investigate possible functional reorganization of the visual word selectivity in the

absence of a typical left-lateralized language network. We have a unique opportunity to examine

fMRI responses to stimuli from different visual categories in an individual (EG) born without the

left superior temporal lobe (likely due to pre/perinatal stroke) but with lVTC largely intact. EG's

frontotemporal language network is completely remapped to the right hemisphere; no language-

related responses, as assessed with fMRI, were observed in the remaining parts of EG's left

hemisphere (Tuckute et al., 2021). EG's reading abilities (as well as other linguistic abilities) are

intact. We here investigated a) whether, in the presence of a right-lateralized language network, a

typical VWFA would emerge in the right VTC (in a typical location, or perhaps in other parts of

the right VTC), and b) whether any word-selectivity is observed in the (spared) left VTC. To

foreshadow the results, no word selectivity was observed in EG's right or left VTC, in the

presence of typical selectivity for other visual categories. We then explored the possibility that

orthographic processing is supported by still selective but more distributed neural populations

using multivariate analyses, and indeed observed such multivariate selectivity bilaterally, but

manifesting more strongly in the right VTC.

Methods

Methods

Participants

Critical participant. The participant EG (fake initials; right-handed female with an advanced

professional degree, 54 year-old at the time of testing) contacted Dr. Fedorenko's lab to

participate in brain research studies. Based on her own report, the lack of the left superior

temporal lobe (**Figure 1**) was discovered when she was 25 years old (in her first MRI scan in

1987) and being treated for depression. No known head traumas or injuries were reported as a

child or adult. Several medical MRI scans were performed in subsequent years (1988, 1998, and

2013) and no change was observed compared to the initial scans. Importantly, EG did not report

any difficulties in reading or general language abilities (see details below). She had also acquired

fluency in a second language (Russian). EG was invited to participate in a series of behavioral

and fMRI assessments at MIT in October 2016. Written informed consent was obtained from EG,

and the study was approved by MIT's Committee on the Use of Humans as Experimental

Subjects (COUHES).

To formally evaluate EG's linguistic abilities, five standardized language assessment tasks were

administered: i) an electronic version of the Peabody Picture Vocabulary Test (PPVT-IV) (Dunn

& Dunn, 2007); ii) an electronic version of the Test for Reception of Grammar (TROG-2)

(Bishop, 2003); iii) the Western Aphasia Battery-Revised (WAB-R) (Kertesz, 2006); iv) the

reading and spelling components of PALPA (Kay et al., 1992); and v) an electronic version of

the verbal components of the Kaufman Brief Intelligence Test (KBIT-2) (Kaufman & Kaufman,

2004). PPVT- IV and TROG-2 target receptive vocabulary and grammar, respectively. In these

tasks, the participant is shown sets of four pictures accompanied by a word (PPVT-IV, 72 trials)

or sentence (TROG-2, 80 trials) and has to choose the picture that corresponds to the

word/sentence by clicking on it. WAB-R (Kertesz, 2006) is a more general language assessment

developed for persons with aphasia. It consists of 9 subscales, assessing 1) spontaneous speech, 2)

auditory verbal comprehension, 3) repetition, 4) naming and word finding, 5) reading, 6) writing,

7) apraxia, 8) construction, visuospatial, and calculation tasks, and 9) supplementary writing and

reading tasks. The verbal components of KBIT-2 include 1) the Verbal Knowledge subtest,

which consists of 60 items measuring receptive vocabulary and general information about the

world, and 2) the Riddles subtest consists of 48 items measuring verbal comprehension,

reasoning, and vocabulary knowledge. Most relevant to the current investigation, the reading

component of WAB-R includes comprehension of written sentences and reading commands; the

supplementary reading tasks include reading of irregular words and nonwords. The reading

component of PALPA (Kay et al., 1992) consists of 21 tasks (tasks #18 through 38), which focus

on single letter recognition, visual lexical decision, out-loud reading (of words with diverse

lexical properties and sentences), and homophone definition (see Supplementary Table 2 for

details).

In line with her self-report, EG performed within normal range on all language assessment tasks.

She got 90% correct on PPVT, 99% correct on TROG-2, 97.6, 98.6, and 98.4 on the aphasia,

language, and cortical quotients of the WAR-B (the criterion cut-off score for diagnosis of

aphasia is an aphasia quotient of 93.8), and a score of 130 (98th percentile) on the KBIT-verbal.

EG's performance was therefore not distinguishable from the performance of neurotypical

controls. With respect to the reading assessments, EG made no errors on the main reading

component of WAB-R, no errors in the reading of irregular words, and one error in the reading

of nonwords (reading 'gobter' instead of 'globter'). For the PALPA tasks, she made no errors on

tasks that focus on single letters (tasks #18-23), no errors on the visual decision tasks (tasks #24-

27), no errors on the out-loud reading tasks (tasks #29-37), and no errors on the homophone

definition task (task #38). For the homophone decision task (task #28), EG made three errors

(out of the 60 trials; all were made on nonword pairs: she did not judge the following pairs as

sounding the same: heem-heam, byme-bime, and phex-feks). Based on this comprehensive

assessment, EG appears to have intact visual letter and word recognition ability.

With respect to fMRI tasks that relevant to the current study, EG completed five runs of the

VWFA localizer; see *The VWFA localizer task* section below).

Neurotypical controls. Twenty-five adults (11 female, mean age = 23.6 years old; age range 18-

38 years; standard deviation 5.21 years) from The Ohio State University (OSU) and the

surrounding community were included in the present study. As part of ongoing projects

exploring the relationship between brain function and connectivity, all participants completed a

battery of fMRI tasks, including, critically, the same VWFA localizer task that EG completed

(see *The VWFA localizer task* section below). All participants had normal or corrected-to-normal

vision, and reported no neurological, neuropsychological, or developmental diagnoses. Written

informed consent was obtained from all participants and the study was approved by Institutional

Review Board at OSU. (It is worth noting that although the control group participants were

younger than EG, further examination revealed no significant correlation between age and word-

selectivity (responses to Words minus the average of the other conditions) in our control group

(r=-0.259, p= 0.094; p-value was obtained by a permutation test).

Data acquisition

EG's data were collected on a 3 Tesla Siemens Trio scanner with a 32-channel head coil at the

Athinoula A. Martinos Imaging Center at the McGovern Institute for Brain Research at MIT.

Data of the control group participants were acquired on a 3 Tesla Siemens Prisma scanner with a

32-channel head coil at the Center for Cognitive and Behavioral Brain Imaging at OSU. To

ensure that any differences between EG and our control group were not due to scanner

differences, we compared word-selectivity in the current control group to a smaller group of

adults recruited at MIT and scanned with the same scanner and protocols as EG; no differences

were observed (Supplementary Table 1). For both EG and controls, a whole-head, high

resolution T1-weighted magnetization-prepared rapid acquisition with gradient echo (MPRAGE)

scan was acquired (EG: repetition time (TR) = 2530 ms, echo time (TE) = 3.48 ms; voxel

resolution = 1.0 mm^3; the control group: TR = 1390 ms, TE = 4.62 ms, voxel resolution = 1.0

mm^3). Functional images were acquired with the same echo-planar imaging (EPI) sequence for

both EG and controls: TR=2000ms, TE=30ms, 172 TRs, 100×100 base resolution, voxel

resolution=2.0 mm^3, 25 slices approximately parallel to the base of the temporal lobe to cover

the entire ventral temporal cortex.

The VWFA localizer task

A VWFA localizer was used to define high-level category-selective regions and to measure

category-selective responses (see Saygin et al., 2016, for details). Briefly, black and white line

drawings of words, scrambled words, objects, and faces, along with the fixation condition were

shown in a blocked design. A grid was overlaid on top of the stimuli so that all stimulus types

(not just scrambled words) had edges. Each stimulus was presented for 500ms (ISI=0.193s) and

overlaid on a different single-color background, and 26 stimuli (including 2 repetitions) were

presented in each block. Each run consisted of 19 blocks (4 blocks per stimulus type and 3

fixation blocks), and participants performed a one-back task. The stimuli are available for

download at http://www.zeynepsaygin.com/ZlabResources.html. EG completed 5 runs, and

participants in the control group completed 2 runs. Note that previous studies using the same task

indicated that 2 runs of data are sufficient to successfully identify the VWFA in a neurotypical

population (Saygin et al., 2016); here, we acquired more runs for EG to ensure that we had

sufficient power and that the results obtained for EG were stable across runs (see *Definition of*

functional regions of interest and univariate analyses section for details).

Preprocessing and fMRI analysis

Data were analyzed with Freesurfer v.6.0.0 (http://surfer.nmr.mgh. harvard.edu/), FsFast

(https://surfer.nmr.mgh.harvard.edu/fswiki/FsFast), FSL

(https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FLIRT) and custom MatLab code. All structural MRI data

were processed using a semiautomated processing stream with default parameters (recon-all

function in Freesurfer: https://surfer.nmr.mgh.harvard.edu/fswiki/recon-all/), which includes the

following major steps: intensity correction, skull strip, surface co-registration, spatial smoothing,

white matter and subcortical segmentation, and cortical parcellation. Cortical gray matter and

ventral temporal cortex masks were created based on the Desikan-Killiany (Desikan et al., 2006)

parcellation in native anatomy for each subject. The success of cortical reconstruction and

segmentation was visually inspected for EG.

Functional images were motion-corrected (time points where the difference in total vector

motion from the previous time point exceeded 1mm were excluded), data from each run were

registered to each individual's anatomical brain image using bbregister, and resampled to

$1.0 \times 1.0 \times 1.0 \text{mm}^3$. For EG, instead of registering the functional image of each run to the

anatomical brain separately, we aligned the functional images of the first four runs to the last run

(which had successful functional-anatomical cross-modal registration with bbregister) with linear

affine transformation (FLIRT); then the functional-anatomical transformation for the last run was

applied to all functional runs and was visually inspected (tkregisterfv) for functional to

anatomical alignment.

The preprocessed functional data were then entered into a first-level analysis. Specifically, data

were detrended, smoothed (3mm FWHM kernel), and the regressor for each experimental

condition (Words, Scrambled Words, Objects, and Faces) was defined as a boxcar function (i.e.,

events on/off) convolved with a canonical hemodynamic response function (a standard gamma

function ($d = 2.25$ and $t = 1.25$)). Orthogonalized motion measures from the preprocessing stage

were used as nuisance regressors for the GLM. Resulting beta estimates for each condition and

contrasts of interest between categories (i.e., Words > Others other categories and Faces > Others

other categories) were used in further analyses. For multivariate analyses, no spatial smoothing

was applied.

Definition of functional regions of interest and univariate analyses

The subject-specific functional regions of interests (fROIs) were defined with a group-

constrained subject-specific (GcSS) approach (Fedorenko et al., 2010). In this approach,

individual fROIs are defined by intersecting participant-specific fMRI contrast maps for the

contrast of interest (e.g., Words > Other conditions) with some spatial constraint(s) or 'parcel(s)',

which denote the area(s) in the brain within which most individuals show responses to the

relevant contrast (based on past studies with large numbers of participants). In the present study,

the VWFA and the FFA parcels were derived from previous studies (FFA, Julian et al., 2012;

VWFA, Saygin et al., 2016) (**Figure 1**) and were generated based on probabilistic maps of

functional activation for the relevant contrasts in independent groups of participants. We

registered these parcels to our participants' own anatomy using the combined volume and

surface-based (CVS) non-linear registration method (mri_cvs_register; Postelnicu et al., 2009).

After mapping the functional parcels to each participant's brain, we defined the VWFA fROI by

selecting within the VWFA parcel the top 10% of most active voxels for the contrast Words >

Other conditions (i.e., scrambled words, objects, and faces). Similarly, we defined the FFA by

selecting within the FFA parcel the top 10% of most active voxels for the contrast Faces > Other

conditions. For the control participants, we used run 1 to define the fROIs, and the run 2 to

extract percent signal changes (PSCs; beta estimates divided by baseline activation) for each of

four experimental conditions. For EG, who had 5 runs of data, this procedure was performed

iteratively for every 2-run/3-run split (e.g., defining fROIs with runs 1&2 and extracting PSCs

from runs 3-5, defining with runs 1&3 and extracting PSCs from runs 2, 4, and 5, etc.; 10

combinations in total), and the results were averaged to derive a single estimate per condition per

fROI. Note that although the parcels are relatively large, by design (in order to accommodate

inter-individual variation in the precise location of the functional region), and can overlap, the

resulting fROIs within an individual are small and do not overlap (Saygin et al., 2016).

Multivariate analyses: Split-half correlations

To further examine whether visual words may be represented and processed in a spatially

distributed manner, we performed a multivariate pattern analysis (MVPA) to measure distinctive

activation patterns for different conditions. The analyses were performed with CoSMoMVPA

toolbox (Oosterhof et al., 2016) (https://www.cosmomvpa.org/). In line with the approach

introduced in Haxby et al. (2001), we examined split-half within-category and between-category

correlations. In particular, a searchlight (radius = 3 voxels) was created for each voxel within the

VTC, and response patterns (i.e., beta estimates for each of the four conditions, normalized by

subtracting mean responses across all conditions) were extracted from each searchlight.

Before performing the critical analysis, we asked whether the overall multivariate representation

structure of the VTC is typical in EG. To do so, we constructed a representational similarity

matrix (RSM) from pairwise similarities (i.e., correlations) based on the voxel-wise response

patterns in a given searchlight area (defined above) to different conditions (e.g., the correlation

between activation patterns across voxels to Words and Faces in a given searchlight area).

Correlations of all searchlights within the VTC were then Fisher z-transformed and averaged.

This resulted in one 4x4 RSM from two runs of data for each participant in the control group; for

EG, RSMs from ten run combinations were averaged to get a single RSM. We then calculated

RSM similarity (correlation) between EG and controls, and tested whether this correlation was

different from the correlations between any given control individual and the rest of the control

group (within-controls correlations; similarity of RSM of each subject to the average RSM of the

remaining control subjects).

Then, similarity of response patterns within a category (e.g., Words-Words or Faces-Faces) vs.

between categories (e.g., Words-Faces) was calculated within the VWFA and FFA parcel

boundaries by Pearson's correlation in a pair-wise manner, and then Fisher z-transformed. We

identified voxels of interest that satisfied the following two criteria: 1) voxels whose local

neighborhoods showed higher within- than between-category correlations (and which therefore

represent categories distinctively); and 2) voxels whose local neighborhoods showed higher

within-condition correlations for a particular category (e.g., Words-Words) than within-condition

correlations for other categories (e.g., Faces-Faces). The second criterion identified voxels which

represent a particular category (e.g., visual words) in a more selective fashion. We refer to such

voxels as multivariate-selective voxels.

To examine hemispheric differences, we computed the number of voxels that exhibited

multivariate selectivity for words or faces in the two hemispheres (within the relevant parcels).

To control for the difference in the size of the search spaces (i.e., the parcels), we divided the

number of the multivariate-selective voxels by the total size of the relevant parcel.

Statistical analyses

Paired t-tests were used for comparisons between conditions for EG (across ten run split combinations) and within the control group. For all analyses where we compared EG's response to the control group, we used a Crawford-Howell modified t-test (Crawford & Howell, 1998), which is widely used in single-case studies because it accounts for the variance of the control group, and the percentage of false positives remains the same for various sizes of the control sample (N=5-50) (Crawford et al., 2009). This frequentist approach provided the point estimate (p-value) for the proportion of the control population that will obtain a score lower than the patient's score. One-tailed tests were reported for the critical (directional) hypothesis tested in the current study (unless noted otherwise). In addition, we computed the Bayesian 95% credible interval (a Bayesian alternative to the modified t-test; Crawford & Garthwaite, 2007) to demonstrate the range of p-values based on 10,000 iterations. To assess the significance of RSM correlations between EG and controls, we generated a null distribution of correlation values by shuffling the matrix of EG and controls (i.e., randomizing the labels of values in the RSMs) and then correlating the new shuffled matrices. This procedure was repeated 10,000 times to create the null distribution of the correlation values. The p-value was calculated by counting the number of correlations in the null-distribution that were higher than the correlation value based on the correct category labels, and then divided by 10,000.

Results

Is word selectivity remapped to the right hemisphere when the language network is remapped to the right hemisphere from birth?

We first examined univariate word selectivity in EG's rVWFA. With the VWFA localizer, we

defined the individual-specific rVWFA fROIs, in EG and controls, within a spatial constraint

(rVWFA parcel; see **Figure 1** and Methods) by contrasting Words versus all other categories

(Words > Others). We then extracted the activation to all four conditions from independent data

(see Methods for details). In neurotypical literate individuals, word selectivity is strongly left-

lateralized (McCandliss et al., 2003); selective responses to visual words in the right homotope

of lVWFA are less frequently observed and are less spatially consistent across individuals.

Results from our control group are in line with this picture: we found no selective activation to

Words as compared to other visual categories in the rVWFA (Words vs. Scrambled Words:

t(24)=-0.964, p= 0.345; Words vs. Objects: t(24)=-0.780, p= 0.443; Words vs. Faces: t(24)=-

0.523, p= 0.605; **Figure 2A**).

Figure 1. EG's MRI showing the missing cortex and the parcels for the right and left VWFA and FFA. Top, T1-weighted images in coronal, sagittal, and axial views. Bottom, the VWFA (purple) and FFA (blue) parcels are projected on EG's reconstructed surface. The parcels served as spatial constraints in defining the fROIs (see Methods), but we also explored the entire VTC for category selectivity (outlined with black solid lines). By design, the parcels are relatively large (to accommodate inter-individual variability in the precise locations of these areas) and therefore can overlap, but the individual fROIs are small and do not overlap. Note that even though part of the anterior lVTC is missing in EG, the stereotypical locations for both the VWFA and FFA are spared.

Critically, in EG, whose language network is completely reorganized to the right hemisphere (Tuckute et al., 2021), we asked whether word selectivity was also remapped to the right hemisphere. Surprisingly, no word selectivity was observed in EG's rVWFA: activation to Words in her rVWFA fROI did not significantly differ from those to Scrambled Words and Faces (Words vs. Scrambled Words: t(9)=-0.913, p= 0.385; Words vs. Faces: t(9)=-0.966, p= 0.359) and was significantly *lower* than Objects (Words vs. Objects: t(9)=-2.346, p= 0.044) (**Figure 2B**). To ensure that this result is not due to the choice of a particular threshold (i.e., top 10%) used to define the rVWFA fROI, we performed the same analysis as above at a range of thresholds (5%-50%). The lack of word selectivity was stable across thresholds (**Figure 2C**). We then compared EG to the controls, and found that there was a trend whereby responses to Words (relative to baseline) were slightly higher in EG compared to the control group (t(24)=0.503, p= 0.310, modified t-test; 95% Bayesian CI [0.17, 0.45]), but this difference was not significant.

To test whether selectivity for other visual categories in the right VTC is typical in EG, we examined face selectivity in EG's rFFA, which is spatially proximal to the rVWFA. Similar to the rVWFA analysis above, we defined the rFFA fROI by contrasting Faces versus other categories (Faces > Others), and extracted the activation to the four conditions from independent

data (see Methods for details). EG's face selectivity remained intact (**Figure 2E)** and did not

differ from that of the control group (**Figure 2D**; t(24)=-0.031, p= 0.488, modified t-test,

Bayesian 95% CI of EG's activation [0.35 0.50]). EG's rFFA fROI showed significantly higher

responses to Faces than to other conditions (Faces vs. Words: t(9)=24.813, p= 1.346×10^{-9}; Faces

vs. Scrambled Words: t(9)=16.850, p= 4.092×10^{-8}; Faces vs. Objects: t(9)=20.312, p=7.921×10^{-9};

Figure 2E), just like what we observed in the control group (Faces vs. Words: t(24)=15.299, p=

7.054×10^{-14}; Faces vs. Scrambled Words: t(24)=12.915, p= 2.689×10^{-12}; Faces vs. Objects:

t(24)=10.724, p=1.235×10^{-10}; **Figure 2D**). Moreover, the face selectivity in EG's rFFA was

observed across all thresholds used to define the rFFA (**Figure 2F**).

Figure 2. Responses to four conditions in the rVWFA and rFFA for EG and the control group. (A) Bar
plots show mean PSCs to the four conditions estimated in independent data within individually defined
rVWFA fROIs (i.e., top 10% word-selective voxels within the rVWFA parcel) for the control group. (B)
Mean PSCs to the four conditions estimated in independent data within the individually defined rVWFA
fROI for EG. Here and in E, the results are averaged across run combinations. (C) Parametrically
decreasing the threshold for defining the rVWFA fROI from the top 5% to 50% word-selective voxels
within the rVWFA parcel. Number of voxels in the rVWFA fROIs: 5%=384 voxels, 50%=3007 voxels.
Here and in F, average PSCs across run combinations are shown for each threshold. (D) Mean PSCs to
the four conditions estimated in independent data within individually defined rFFA fROIs for the control
group. (E) Mean PSCs to the four conditions estimated in independent data within the individually
defined rFFA fROI for EG. (F) Parametrically decreasing the threshold for defining the rFFA fROI from
the top 5% to 50% face-selective voxels within the rFFA parcel. Number of voxels in the defined rFFA
fROIs: 5%=90 voxels, 50%=699 voxels. In the bar plots, dots correspond to individual data points for
each condition (controls: n=25 participants; EG: n=10 run combinations, from ten iterations). Horizontal
bars reflect significant paired t-tests $p < 0.05$. Error bars in both the bar and line plots denote standard
errors of the mean by participants (for the control group) and by run combinations (for EG).
449

To ensure that we did not miss any possible word-selective voxels by applying a predefined

spatial constraint (i.e., VWFA parcel) and to account for the possibility that EG's VWFA is

located in a different part of the visual cortex, we searched for word selectivity within the entire

rVTC mask for EG. Specifically, different thresholds from top 1% to top 10% were used to

define the most word-selective voxels (Words > Others). Even within this broad mask, no word-

selective responses were observed in independent data across all thresholds; in fact, the

responses were lowest to words than the other three conditions (**Figure 3A**). In contrast, robust

face-selective responses were observed in independent data across all thresholds when searching

for face-selective voxels (Faces > Others) (**Figure 3B**).

Finally, we supplemented our main analyses, which rely on the Words > Others contrast to

define the VWFA, as is commonly done in the literature (e.g., Dehaene-Lambertz et al., 2018;

Rosenke et al., 2021), with an analysis, which relies on a less stringent localizer contrast (Words

vs. Scrambled Words; Glezer et al., 2009; Lerma-Usabiaga et al., 2018). Unlike the Words >

Others contrast, this contrast does not control for semantics or visual stimulus complexity. Even

with this broader contrast, we found no word selectivity within EG's rVTC across thresholds

(Supplementary Figure 1).

Figure 3. Mean PSCs in the rVTC and lVTC at different thresholds for EG. (A-B) Parametrically
decreasing the threshold for defining word-selective (Words > Others) and face-selective (Faces > Others)
voxels from the top 1% to 10% within the rVTC. Mean PSCs across run combinations (from 10 iterations)
are shown for each threshold. (C-D) Parametrically decreasing the threshold for defining word-selective
and face-selective voxels from the top 1% to 10% within the lVTC. Mean PSCs across run combinations
are shown for each threshold. Number of selected voxels: rVTC: 1%=216 voxels, 10%=2164 voxels;
lVTC: 1%=154 voxels, 10%=1536 voxels.

Is there any word selectivity in the spared left VTC?

Because we did not observe a right-lateralized VWFA and because the left VTC was largely

intact, we also asked whether a canonical lVWFA may have developed in EG (perhaps due to

some specific visual features of word forms that are better represented in the left

hemisphere/right visual field (e.g., Hsiao & Lam, 2013; Seghier & Price, 2011; Tadros et al.,

2013). In the control group, we observed robust word selectivity in the lVWFA (as expected

from previous studies), with Words eliciting greater activation than each of the other conditions:

Words vs. Scrambled Words: $t(24)=5.413$, $p=1.467\times10^{-5}$; Words vs. Objects: $t(24)=4.077$, $p=$

4.333×10^{-4}; Words vs. Faces: $t(24)=5.942$, $p=3.935\times10^{-6}$; **Figure 4A**). In contrast, we found no

word selectivity in EG's lVWFA (**Figure 4B**): activation to Words was around baseline and

significantly lower than those to Scrambled Words and Objects (Words vs. Scrambled Words:

$t(9)=-3.092$, $p=1.289\times10^{-2}$; Words vs. Objects: $t(9)=-2.432$, $p=3.785\times10^{-2}$) and did not

significantly differ from that to Faces (Words vs. Faces: $t(9)=-0.699$, $p=0.503$). Moreover, EG's

activation to Words was significantly lower than the control group's ($t(24)=-1.616$, $p=5.961\times10^{-}$

^{2}, modified t-test; 95% Bayesian credible interval (CI) [0.01, 0.13]). This result was stable

across thresholds that were used to define the lVWFA fROI (**Figure 4C**), and we did not observe

word selectivity in EG when we searched across the entire lVTC with varying thresholds (**Figure**

3C), or when we used a less stringent contrast (Words vs. Scrambled Words; Supplementary

Figure 1).

Figure 4. Responses to four conditions in the lVWFA and lFFA for EG and the controls group. (A), Bar
plots show mean PSCs to four conditions estimated in independent data within individually defined
lVWFA fROIs (i.e., top 10% word-selective voxels within lVWFA parcel) for the control group. (B),
Mean PSCs to the four conditions estimated in independent data within individually defined lVWFA
fROIs for EG. Here and in E, the results are averaged across run combinations.. (C), Parametrically
decreasing the threshold for defining the lVWFA fROIs from the top 5% to 50% word-selective voxels
within the lVWFA parcel. Number of voxels in the defined lVWFA fROIs: 5%=295 voxels, 50%=2402
voxels. Here and in F, average PSCs across run combinations are shown for each threshold. (D), Mean
PSCs across participants to the four conditions estimated in independent data within individually defined
lFFA fROIs for the control group. (E), Mean PSCs to four conditions estimated in independent data
within individually defined lFFA fROIs for EG. (F), Parametrically decreasing the threshold for defining
lFFA fROI from the top 5% to 50% face-selective voxels within the lFFA parcel. Number of voxels in the
defined lFFA fROIs: 5%=18 voxels, 50%=185 voxels. In the bar plots, dots correspond to individual data
points (controls: n=25 subjects; EG: n=10 run combinations, from ten iterations). Horizontal bars reflect
significant paired t-tests p < 0.05. Error bars in both the bar and line plots denote standard errors of the
mean by participants (for the control group) and by run combinations (for EG).

Similar to what we did for the rVTC, to test whether selectivity for other visual categories in the

left VTC is typical in EG, we examined face selectivity in EG's lFFA. EG's face selectivity

remained intact (**Figure 4E)** and did not differ from that of the control group (**Figure 4D**; t(24)=-

0.275, p= 0.393, modified t-test; 95% Bayesian CI, [0.27 0.50]). EG's lFFA fROI showed

significantly higher responses to Faces than to other conditions (Faces vs. Words: t(9)=47.571,

p= 4.015×10^{-12}; Faces vs. Scrambled Words: t(9)=24.681, 1.411×10^{-9}; Faces vs. Objects:

t(9)=34.494, p= 7.141×10^{-11}; **Figure 4E)**, just like what we observed in the control group (Faces

vs. Words: t(24)=7.076, p=1.467×10^{-5}; Faces vs. Scrambled Words: t(24)=8.102, p= 2.518×10^{-8};

Faces vs. Objects: t(24)=5.517, p= 1.130×10^{-5}; **Figure 4D**). As was the case for the rFFA, the

face selectivity in EG's lFFA was observed across all thresholds used to define the lFFA (**Figure

4F**).

Altogether, the examination of EG's right and left VTC suggests that without the typical left-

hemisphere frontotemporal language network from birth—and presumably without the necessary

connections between these areas and parts of the VTC—a canonical VWFA, a word-selective

area, does not develop in either hemisphere.

Distributed neural representation of visual words: multivariate pattern analysis (MVPA)

Our univariate analyses measured the average activation across the voxels most responsive to the

visual category in question. Although this is the classic approach to demonstrate category

selectivity, univariate analyses may be insensitive to potentially meaningful distributed

representation patterns in suprathreshold and/or subthreshold voxels. Previous studies have

found that categorical information can be reliably decoded by comparing within-category versus

between-category correlations in the VTC (e.g., Haxby et al., 2001). Moreover, a recent study

found mature representational similarity structures via multivariate patterns in children with no

univariate selectivity, suggesting that distributed representations may developmentally precede

category selectivity (Cohen et al., 2019). To explore whether words may be represented in a

distributed fashion in EG, we performed a series of multivariate pattern analyses (MVPA).

We first examined the representational similarity matrices (RSMs) in the entire VTC (see

Methods) to investigate whether multivariate representational structure for visual categories was

preserved in EG. Indeed, we found that EG's rVTC RSM was strongly and significantly

correlated with that of the control group (r=0.918, p= 9.100×10^{-3} (permutation test);

Supplementary Figure 3). We tested whether this correlation between EG and the control group

was different from the correlations between any given control individual and the rest of the

control group (see Methods). Single case comparisons showed that the RSM correlation for EG

vs. controls did not significantly differ from the within-controls correlations (t(24)=0.797, p=

0.217, Bayesian 95% CI [0.09 0.351]). Similar results were found for the lVTC: the RSMs of EG

and the control group were strongly and significantly correlated (r=0.888, p= 7.300×10^{-3}), and

the correlation for EG vs. controls did not significantly differ from the within-controls

correlations (t(24)=0.835, p= 0.206, Bayesian 95% CI [0.09 0.34]).

We then asked whether EG's 'VWFA' may contain voxels that show distinct distributed

activation patterns to visual words. Specifically, as described in Methods, we searched for voxels

that satisfied the following criteria: the searchlight around a given voxel should show 1)

distinctive response patterns to Words vs. other categories (e.g., the Words-Words correlation

should be higher than the Words-Faces correlation); and 2) stronger within-category correlations

for the preferred category (e.g., the Words-Words correlation should be higher than the Faces-

Faces, Objects-Objects, and Scrambled Words-Scrambled Words correlations). Previous studies

have shown distributed representations within category-selective regions (e.g., the FFA) of non-

preferred categories (e.g., places), and debate is ongoing over whether this information has

functional relevance (e.g., Kanwisher, 2010). Our second criterion was included to identify

voxels that show more stable multivariate representations for the category of interest (e.g.,

Words) compared to other categories. Indeed, we identified a set of voxels within the rVWFA

and lVWFA parcels that showed a reliable distributed code for words in both controls and EG

(Supplementary Figure 2A, 2C; Supplementary Table 3). In addition, mirroring the univariate

analyses, we also identified a set of voxels that showed a reliable distributed code for faces

within the rFFA and lFFA parcels in both controls and EG (Supplementary Figure 2B, 2D;

Supplementary Table 3).

Critically, to test whether these distributed responses to visual words differ between EG and the

controls, we subtracted the average within-category correlations for all non-word categories from

the within-category correlations for Words. This difference tells us how much stronger the

within-category correlation is for the preferred compared to the non-preferred categories. EG

showed comparable within-category correlation differences to the controls in both the rVWFA

(t(24)=1.081, p= 0.145, modified t-test, 95% Bayesian CI [0.05, 0.26] and lVWFA (t(24)=0.184,

p= 0.428, modified t-test, 95% Bayesian CI [0.30, 0.50]); **Figure 5A**). Similar within-category

correlation differences were also observed between EG and the controls in the rFFA (t(24)=-

0.802, p= 0.215; 95% Bayesian CI [0.09, 0.34]) and lFFA (t(24)=0.488, p= 0.315; 95%

Bayesian CI [0.18, 0.46]); **Figure 5B**).

Figure 5. Results of the multivariate pattern analysis (MVPA). (A) Within-category correlation differences between preferred (i.e., Words-Words) and non-preferred (i.e., average within-category correlation of non-word conditions) conditions for EG and the controls in the rVWFA and lVWFA parcels. Here and in B, correlation values were Fisher's z-transformed. (B) Within-category correlation differences between preferred (i.e., Faces-Faces) and non-preferred (i.e., average within-category correlation of non-face conditions) conditions for EG and the controls in the rFFA and lFFA parcels. (C) Proportion of voxels that show multivariate selectivity for Words in the rVWFA and lVWFA parcels for EG and the controls. (D) Proportion of voxels that show multivariate selectivity for Faces in the rFFA and lFFA parcel for EG and the controls. In the bar plots, dots correspond to individual data points (controls: n=25 subjects; EG: n=10 run combinations from ten iterations). Horizontal bars reflect significant LH-RH differences, p<0.05. Error bars denote standard errors of the mean by participants (for the control group) and by run combinations (for EG).

Finally, we explored potential hemispheric asymmetries with respect to multivariate selectivity

for words and faces. Left-hemisphere dominance for words and right-hemisphere dominance for

faces are well established with univariate measures. Do multivariate patterns also show these

asymmetries? We calculated the number of voxels that show multivariate selectivity for words

and faces within the left and right VWFA and FFA parcels, respectively (controlling for the size

of the search space; see Methods). For Words, mirroring the univariate results in the past

literature, a numerically larger proportion of voxels showing multivariate word selectivity was

found in the left compared to the right VWFA in the controls (although the difference did not

reach significance: $t(24)=1.181$, $p=0.249$). Interestingly, the opposite pattern was observed in EG,

with a larger proportion of voxels showing multivariate word selectivity found in the right

VWFA ($t(9)= -2.469$, $p= 3.564\times10^{-2}$) (**Figure 5C**). For Faces, a numerically larger proportion of

voxels showing multivariate face selectivity was found in the right compared to the left FFA in

the controls (although, like for Words, the difference did not reach significance: $t(24)=0.367$,

$p=0.717$), and EG displayed the opposite pattern: a significantly larger proportion of voxels

showing multivariate face selectivity in the left FFA ($t(9)=8.983$, $p=8.672\times10^{-6}$) (**Figure 5D**).

Discussion

Case studies provide valuable insights for understanding the functional organization of the

human brain and the patterns of reorganization that follow neurological damage (Vaidya et al.,

2019). In the current study, we had a unique opportunity to collect fMRI data from an individual

(EG) born without her left superior temporal lobe in order to ask whether category-selective

responses to visual words within the ventral temporal cortex would be affected. Specifically, we

asked: in the absence of a typical left-hemisphere language network, does word selectivity

emerge, and if so, where? Does the VWFA emerge in the right VTC given that EG's language

network is reorganized to the right hemisphere (Tuckute et al., 2021)? Or do we instead, or

additionally, observe word selectivity in the canonical VWFA within the spared lVTC?

Surprisingly, we found no univariate word selectivity within the rVWFA, the right homotope of

the canonical lVWFA, and even in an expanded search space of the entire rVTC. We also did not

observe word-selective responses in the lVWFA or in any other spared parts of the lVTC.

Moreover, the response magnitude to Words (relative to baseline) was significantly lower in

EG's lVWFA than that of controls. Importantly, this lack of category selectivity was specific to

Words: selective responses to Faces remained intact in EG's rFFA and lFFA.

Absent univariate word selectivity, we then explored multivariate representations of visual words

in EG. We found that EG's VTC showed an overall similar representational structure to that of

the control group, and that, like the control group, EG had a set of voxels whose local

neighborhoods robustly differentiated between Words and other visual categories in their

patterns of response. Critically, these voxels also showed a higher within-category correlation

(across runs) for Words compared to the within-category correlations for other categories, and

the degree of this 'multivariate selectivity' was similar between EG and the controls.

Interestingly, however, in contrast to the typically observed left-hemispheric dominance for

words and right-hemispheric dominance for faces, EG had a larger proportion of voxels that

showed multivariate selectivity for words in her right than her left VWFA, and a larger

proportion of voxels that showed multivariate selectivity for faces in her left FFA than her right

FFA.

Altogether, the current study suggests that in the absence of a typical left-lateralized language

network—at least in cases of early left-hemispheric damage, when language processing is

'forced' to the right hemisphere—neural selectivity for visual words is, or at least can be, highly

atypical. In our participant with extensive left-hemispheric temporal damage and a right-

lateralized language network, we observed no word selectivity in the univariate analyses in either

the right or the spared left VTC. The absence of such selectivity, combined with EG's intact

reading ability, suggests that successful orthographic processing may depend on a more

distributed and more right-lateralized neural representation than that observed in typical literate

adults.

Canonical univariate word selectivity may not emerge when a left-hemisphere language

network is missing from birth

The interesting case of EG allowed us to investigate how visual word processing within the VTC

can be affected by a congenital or early left-hemisphere lesion outside of visual cortex. Our

results provide the first evidence of atypical visual word selectivity in the VTC when the left-

lateralized language cortex is missing, and when the language network consequently develops in

the right hemisphere (Tuckute et al., 2021) during the early stages of language learning and prior

to learning to read. Our study also suggests that even with some remaining anatomical

connections between the spared lVTC and the frontal and temporal areas (presumably via local

U-fibers or remaining long-range fibers), the lVWFA will not emerge at the stereotypical

location when the left hemisphere does not support language processing.

How does a left language network contribute to the development of the VWFA? One possibility

is that in a typical brain, the site of the putative VWFA is predisposed to written scripts via

intrinsic co-activation with speech and/or language/semantic areas in the left temporal (and

maybe frontal) cortex. The processing of speech sounds is tightly linked to reading, and impaired

coding of speech sounds (e.g., phonological awareness) is often considered a precursor of

dyslexia (e.g., Pennington & Bishop, 2008; Shaywitz et al., 2002). In the absence of regions that

typically support speech processing (within the left superior and middle temporal gyri (STG and

MTG); Raschle et al., 2012), EG's left VTC lacked early interactions with these regions. Further,

a typical lVWFA communicates visual orthographic information to higher-level left temporal

(and maybe frontal) regions that integrate visual and auditory information, like the amodal

language regions that process lexical-semantic and combinatorial linguistic information (e.g.,

Fedorenko et al., 2020) and the regions that support abstract conceptual processing (e.g., Binder

et al., 2009; Patterson et al., 2007)—both sets of areas that were also missing in EG's left

hemisphere. Consistent with the idea that connections between the left VTC and the ipsilateral

high-level language areas may be critical for the emergence of the lVWFA, a recent study found

that in newborns, the putative site of the lVWFA already shows preferential functional

connectivity to the areas that will later respond to high-level linguistic information, compared to

adjacent regions (Li et al., 2020); this pre-existing functional coupling may further strengthen

during language and reading acquisition. Because EG was missing both i) speech-responsive

areas, and ii) higher-level language/conceptual areas in her left hemisphere, we cannot evaluate

the relative contributions of these two sets of areas and their connections with the VTC to the

emergence of a canonical VWFA. We speculate that both may be important.

Right-hemispheric neural correlates that support visual word processing

Because EG's language network resides in the right hemisphere (Tuckute et al., 2021), we

expected to find a VWFA in the right VTC. Surprisingly, no univariate word selectivity was

observed in EG's VTC, and this result could not be explained by either a lack of functional

specialization *anywhere* in the rVTC (intact face selectivity was observed), or our particular

procedure for defining the VWFA (the results held across a range of thresholds, when using a

larger search space, and across different contrasts). Interestingly, neurotypical individuals with

right-dominant language activation sometimes also lack a VWFA in the right VTC (e.g., Cai et

al., 2008; Van der Haegen et al., 2012); in these individuals, lVTC appears to be engaged during

word recognition, presumably due to stronger frontotemporal anatomical connections in the LH

than in the RH, and any language activation on the left (even if it's non-dominant) would engage

the lVTC for reading (Powell et al., 2006). In the case of EG, her right-hemisphere language

network may have lacked early interactions with the rVTC, and her left-hemisphere language

network was altogether lacking, resulting in the atypical word selectivity (in both right and left)

that we observed here.

Some have also argued that the development of word-selective cortex directly competes with

face-selective cortex for neural resources, thus contributing to right-hemispheric dominance for

face processing and left-hemispheric dominance for word processing in neurotypical individuals

(e.g., Behrmann & Plaut, 2015; Dehaene & Cohen, 2007). Interestingly, EG showed typical face

selectivity in the right FFA, not different from controls. It is therefore possible that a focal word-

selective area failed to emerge in EG's rVTC because the relevant cortical tissue had already

been 'assigned' to face processing, perhaps due to some specific visual features of faces that are

better represented in the right hemisphere/left visual field (e.g., holistic face processing; Li et al.,

2017; Rossion et al., 2000). Interestingly, although the univariate response magnitude to faces

was higher in EG's right than left FFA (similar to controls), more multivariate-selective face

voxels were present in EG's lFFA, compared to her rFFA, in sharp contrast to controls. The

functional significance of the latter is at present unclear, but can be explored in future work. On

the other hand, EG had more multivariate-selective word voxels in her rVWFA than her lVWFA,

presumably related to the fact that her language network resides in the right hemisphere. These

data suggest that the hemispheric dominance of the language network drives the laterality of

visual processing in the VTC (be it implemented focally, or in a distributed fashion), at least for

words, but perhaps also for faces (Behrmann & Plaut, 2020; Dehaene & Cohen, 2007).

Multivariate responses to words

Despite the absence of typical univariate word selectivity, EG's VTC was similar in its overall

representation similarity structure—across different visual stimulus classes—to that of the

control group. Consistent with our results, Liu et al. (2019) also found typical representational

structure in category-selective regions after resections within or outside VTC; in addition, a

recent study found mature representational similarity structures in children (5-7 year-olds) with

no univariate selectivity, suggesting that distributed representations precede category selectivity

(M. A. Cohen et al., 2019). Our results provide another case where a typical multivariate

representational structure is observed in the absence of univariate selectivity for words.

Moreover, our results showed that EG (like typical adults) displayed distinctive activation

patterns within the general vicinity of both right and left VWFA that could distinguish visual

words from other conditions, with higher within- than between-category correlations (Haxby et

al., 2001) and more robust (consistent over time) representations of Words than other conditions.

Supporting the idea that multivariate selectivity for words may be functionally useful, Stevens et

al. (2017) found the VWFA (defined in individual participants using a standard univariate

contrast) discriminated words from pseudowords, and did so more strongly than other control

regions (e.g., the FFA).

Numerous questions remain about the multivariate word representations observed in EG. We

will highlight two here. First, what is the relationship between univariate and multivariate

representations? Univariate selectivity may partially depend on a connectivity scaffold or other

genetically defined instructions to determine the location of functional specialized areas (Deen et

al., 2017; Kamps et al., 2020; Li et al., 2020). In contrast, distributed representations like those

found in EG may rely less on a connectivity scaffold. Do distributed representations reflect a

fundamentally different neural code from that associated with focal representations? Some have

argued that the answer is no, and that multivariate analyses may simply be more sensitive than

univariate approaches given that they consider the heterogeneity of response across voxels within

a region as well as potential subthreshold voxels (Davis et al., 2014). On the other hand,

multivariate representations may be truly distinct and reflect information derived from bottom-up

visual statistics to a greater extent; and thus perhaps in typical development, representational

similarity structures precede univariate selectivity (M. A. Cohen et al., 2019) and emergence of

univariate selectivity requires experience-dependent interactions with higher-level areas (e.g., the

language cortex in case of the VWFA). Speculating further, perhaps during EG's early stages of

language development (while her language system was maturing in the right hemisphere), the

right visual cortex lacked the critical early interaction with the language cortex, due to the lack of

strong innate connectivity between the two in the right hemisphere (cf. the left hemisphere; e.g.,

Li et al., 2020; Powell et al., 2016), resulting in a more distributed representation for visual

words.

And second, how are multivariate representations read out by downstream brain regions for

further processing (Williams et al., 2007); specifically, whereas the behavioral relevance of

univariate activity has been extensively investigated in both adults and pediatric populations (e.g.,

Furl et al., 2011; Grill-Spector et al., 2004; Huang et al., 2014; Song et al., 2015), the behavioral

relevance of multivariate representations is still under debate. One study found that performance

in a visual search task among different pairs of object categories could be predicted from

underlying neural similarity structures (M. A. Cohen et al., 2016). At this point, it is unclear how

multivariate representations for words contribute to EG's reading ability, and whether the right-

hemisphere representations are more important—questions that will be a focus of future studies.

Limitations and future directions

Overall, our study provides unique evidence that in the absence of a typical left-hemisphere

language network—at least in cases of early left-hemisphere damage—the canonical word-

selective lVWFA may not develop. Some limitations are worth noting. First, our analyses were

restricted to the ventral temporal cortex, which was the focus of our investigation and where we

had coverage. Although we show that normal reading ability is possible in the absence of focal

selectivity for word processing in the VTC, there may be other pathways or neural structures

outside of the VTC that are important for EG's reading ability? For example, Seghier et al. (2012)

reported a patient who acquired dyslexia following extensive left ventral occipitotemporal cortex

(LvOT) resection, but then regained reading ability; they provided evidence to suggest that the

patient's reading ability was now supported via a direct connection between the occipital visual

cortex and the left superior temporal sulcus (STS) without the involvement of the ventral visual

stream. It is therefore possible that the STS in the language-dominant hemisphere (i.e., the right

hemisphere) may support visual word processing through its connections with the occipital

cortex. Future work will explore word processing in EG using whole-brain coverage. Second,

EG's missing left superior temporal cortex led to right-lateralized speech and language

processing. As discussed earlier, it remains unclear whether speech and higher-level language

areas contribute equally to the emergence of the lVWFA, and whether temporal language regions

may be more important than the frontal ones. Finally, here we reported a single case that sheds

new light on the role of speech and language areas in developing a VWFA. Generalization to

other, similar cases, and data from other methods like noninvasive brain stimulation will help us

better understand the causal role of language cortex in developing visual word selectivity.

References

Assem, M., Glasser, M. F., Van Essen, D. C., & Duncan, J. (2020). A Domain-General

Cognitive Core Defined in Multimodally Parcellated Human Cortex. *Cerebral Cortex*,

30(8), 4361–4380. https://doi.org/10.1093/cercor/bhaa023

Baker, C., Liu, J., Wald, L., Kwong, K. K., Benner, T., & Kanwisher, N. (2007). Visual word

processing and experiential origins of functional selectivity in human extrastriate cortex.

Proc Natl Acad Sci USA. https://www.pnas.org/content/104/21/9087

Barttfeld, P., Abboud, S., Lagercrantz, H., Adén, U., Padilla, N., Edwards, A. D., Cohen, L.,

Sigman, M., Dehaene, S., & Dehaene-Lambertz, G. (2018). A lateral-to-mesial

organization of human ventral visual cortex at birth. *Brain Structure and Function*,

223(7), 3107–3119. https://doi.org/10.1007/s00429-018-1676-3

Behrmann, M., & Plaut, D. C. (2013). Distributed circuits, not circumscribed centers, mediate

visual recognition. *Trends in Cognitive Sciences*, *17*(5), 210–219.

https://doi.org/10.1016/j.tics.2013.03.007

Behrmann, M., & Plaut, D. C. (2015). A vision of graded hemispheric specialization: Graded

hemispheric specialization. *Annals of the New York Academy of Sciences*, *1359*(1), 30–46.

https://doi.org/10.1111/nyas.12833

Behrmann, M., & Plaut, D. C. (2020). Hemispheric Organization for Visual Object Recognition:

A Theoretical Account and Empirical Evidence. *Perception*, *49*(4), 373–404.

https://doi.org/10.1177/0301006619899049

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where Is the Semantic

System? A Critical Review and Meta-Analysis of 120 Functional Neuroimaging Studies.

Cerebral Cortex, *19*(12), 2767–2796. https://doi.org/10.1093/cercor/bhp055

Bishop, D. (2003). *Test for Reception of Grammar version 2 (TROG-2).*

Blank, I., Kanwisher, N., & Fedorenko, E. (2014). A functional dissociation between language

834   and multiple-demand systems revealed in patterns of BOLD signal fluctuations. *Journal*

835   *of Neurophysiology*, *112*(5), 1105–1118. https://doi.org/10.1152/jn.00884.2013

836   Bouhali, F., Thiebaut de Schotten, M., Pinel, P., Poupon, C., Mangin, J.-F., Dehaene, S., &

837   Cohen, L. (2014). Anatomical Connections of the Visual Word Form Area. *Journal of*

838   *Neuroscience*, *34*(46), 15402–15414. https://doi.org/10.1523/JNEUROSCI.4918-13.2014

839   Cai, Q., Lavidor, M., Brysbaert, M., Paulignan, Y., & Nazir, T. A. (2008). Cerebral

840   Lateralization of Frontal Lobe Language Processes and Lateralization of the Posterior

841   Visual Word Processing System. *Journal of Cognitive Neuroscience*, *20*(4), 672–681.

842   https://doi.org/10.1162/jocn.2008.20043

843   Cai, Q., Paulignan, Y., Brysbaert, M., Ibarrola, D., & Nazir, T. A. (2010). The Left Ventral

844   Occipito-Temporal Response to Words Depends on Language Lateralization but Not on

845   Visual Familiarity. *Cerebral Cortex*, *20*(5), 1153–1163.

846   https://doi.org/10.1093/cercor/bhp175

847   Cohen, L., Martinaud, O., Lemer, C., Lehéricy, S., Samson, Y., Obadia, M., Slachevsky, A., &

848   Dehaene, S. (2003). Visual Word Recognition in the Left and Right Hemispheres:

849   Anatomical and Functional Correlates of Peripheral Alexias. *Cerebral Cortex*, *13*(12),

850   1313–1333. https://doi.org/10.1093/cercor/bhg079

851   Cohen, M. A., Alvarez, G. A., Nakayama, K., & Konkle, T. (2016). Visual search for object

852   categories is predicted by the representational architecture of high-level visual cortex.

853   *Journal of Neurophysiology*, *117*(1), 388–402. https://doi.org/10.1152/jn.00569.2016

Cohen, M. A., Dilks, D. D., Koldewyn, K., Weigelt, S., Feather, J., Kell, A. JE., Keil, B., Fischl,

B., Zöllei, L., Wald, L., Saxe, R., & Kanwisher, N. (2019). Representational similarity

precedes category selectivity in the developing ventral visual pathway. *NeuroImage*, *197*,

565–574. https://doi.org/10.1016/j.neuroimage.2019.05.010

Crawford, J. R., & Garthwaite, P. H. (2007). Comparison of a single case to a control or

normative sample in neuropsychology: Development of a Bayesian approach. *Cognitive*

Neuropsychology, *24*(4), 343–372. https://doi.org/10.1080/02643290701290146

Crawford, J. R., Garthwaite, P. H., & Howell, D. C. (2009). On comparing a single case with a

control sample: An alternative perspective. *Neuropsychologia*, *47*(13), 2690–2695.

https://doi.org/10.1016/j.neuropsychologia.2009.04.011

Crawford, J. R., & Howell, D. C. (1998). Comparing an Individual's Test Score Against Norms

Derived from Small Samples. *The Clinical Neuropsychologist*, *12*(4), 482–486.

https://doi.org/10.1076/clin.12.4.482.7241

Davis, T., LaRocque, K. F., Mumford, J. A., Norman, K. A., Wagner, A. D., & Poldrack, R. A.

(2014). What do differences between multi-voxel and univariate analysis mean? How

subject-, voxel-, and trial-level variance impact fMRI analysis. *NeuroImage*, *97*, 271–283.

https://doi.org/10.1016/j.neuroimage.2014.04.037

Deen, B., Richardson, H., Dilks, D. D., Takahashi, A., Keil, B., Wald, L. L., Kanwisher, N., &

Saxe, R. (2017). Organization of high-level visual cortex in human infants. *Nature*

Communications, *8*(1), 13995. https://doi.org/10.1038/ncomms13995

Dehaene, S., & Cohen, L. (2007). Cultural Recycling of Cortical Maps. *Neuron*, *56*(2), 384–398.

https://doi.org/10.1016/j.neuron.2007.10.004

Dehaene, S., & Cohen, L. (2011). The unique role of the visual word form area in reading.

Trends in Cognitive Sciences, *15*(6), 254–262. https://doi.org/10.1016/j.tics.2011.04.003

Dehaene, S., Cohen, L., Morais, J., & Kolinsky, R. (2015). Illiterate to literate: Behavioural and

cerebral changes induced by reading acquisition. *Nature Reviews Neuroscience*, *16*(4),

234–244. https://doi.org/10.1038/nrn3924

Dehaene-Lambertz, G., Monzalvo, K., & Dehaene, S. (2018). The emergence of the visual word

form: Longitudinal evolution of category-specific ventral visual areas during reading

acquisition. *PLOS Biology*, *16*(3), e2004103.

https://doi.org/10.1371/journal.pbio.2004103

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R.

L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006). An

automated labeling system for subdividing the human cerebral cortex on MRI scans into

gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.

https://doi.org/10.1016/j.neuroimage.2006.01.021

Downing, P. E., Jiang, Y., Shuman, M., & Kanwisher, N. (2001). A cortical area selective for

visual processing of the human body. *Science (New York, N.Y.)*, *293*(5539), 2470–2473.

https://doi.org/10.1126/science.1063414

Duncan, J., Assem, M., & Shashidhara, S. (2020). Integrated Intelligence from Distributed Brain

Activity. *Trends in Cognitive Sciences*, *24*(10), 838–852.

https://doi.org/10.1016/j.tics.2020.06.012

Dunn, L. M., & Dunn, D. M. (2007). *The Peabody Picture Vocabulary Test, Fourth Edition*.

NCS

Pearson, Inc.

Epstein, R., & Kanwisher, N. (1998). A cortical representation of the local visual environment.

Nature, *392*(6676), 598–601. https://doi.org/10.1038/33402

Fedorenko, E., Behr, M. K., & Kanwisher, N. (2011). Functional specificity for high-level

linguistic processing in the human brain. *Proceedings of the National Academy of*

Sciences, *108*(39), 16428–16433. https://doi.org/10.1073/pnas.1112937108

François, C., Garcia-Alix, A., Bosch, L., & Rodriguez-Fornells, A. (2021). Signatures of brain

plasticity supporting language recovery after perinatal arterial ischemic stroke. *Brain and*

Language, *212*, 104880. https://doi.org/10.1016/j.bandl.2020.104880

Furl, N., Garrido, L., Dolan, R. J., Driver, J., & Duchaine, B. (2011). Fusiform Gyrus Face

Selectivity Relates to Individual Differences in Facial Recognition Ability. *Journal of*

Cognitive Neuroscience, *23*(7), 1723–1740. https://doi.org/10.1162/jocn.2010.21545

Gerrits, R., Van der Haegen, L., Brysbaert, M., & Vingerhoets, G. (2019). Laterality for

recognizing written words and faces in the fusiform gyrus covaries with language

dominance. *Cortex*, *117*, 196–204. https://doi.org/10.1016/j.cortex.2019.03.010

Glezer, L. S., Jiang, X., & Riesenhuber, M. (2009). Evidence for Highly Selective Neuronal

Tuning to Whole Words in the "Visual Word Form Area." *Neuron*, *62*(2), 199–204.

https://doi.org/10.1016/j.neuron.2009.03.017

Greenblatt, S. H. (1976). Subangular alexia without agraphia or hemianopsia. *Brain and*

Language, *3*(2), 229–245. https://doi.org/10.1016/0093-934X(76)90019-5

Grill-Spector, K., Knouf, N., & Kanwisher, N. (2004). The fusiform face area subserves face

perception, not generic within-category identification. *Nature Neuroscience*, *7*(5), 555–

562. https://doi.org/10.1038/nn1224

Hamamé, C. M., Szwed, M., Sharman, M., Vidal, J. R., Perrone-Bertolotti, M., Kahane, P.,

Bertrand, O., & Lachaux, J.-P. (2013). Dejerine's reading area revisited with intracranial

EEG: Selective responses to letter strings. *Neurology*, *80*(6), 602–603.

https://doi.org/10.1212/WNL.0b013e31828154d9

Hannagan, T., Amedi, A., Cohen, L., Dehaene-Lambertz, G., & Dehaene, S. (2015). Origins of

the specialization for letters and numbers in ventral occipitotemporal cortex. *Trends in*

Cognitive Sciences, *19*(7), 374–382. https://doi.org/10.1016/j.tics.2015.05.006

Hasson, U., Levy, I., Behrmann, M., Hendler, T., & Malach, R. (2002). Eccentricity Bias as an

Organizing Principle for Human High-Order Object Areas. *Neuron*, *34*(3), 479–490.

https://doi.org/10.1016/S0896-6273(02)00662-1

Haxby, J. V., Gobbini, M. I., Furey, M. L., Ishai, A., Schouten, J. L., & Pietrini, P. (2001).

Distributed and Overlapping Representations of Faces and Objects in Ventral Temporal

Cortex. *Science*, *293*(5539), 2425–2430. https://doi.org/10.1126/science.1063736

Hsiao, J. H., & Lam, S. M. (2013). The Modulation of Visual and Task Characteristics of a

Writing System on Hemispheric Lateralization in Visual Word Recognition—A

Computational Exploration. *Cognitive Science*, *37*(5), 861–890.

https://doi.org/10.1111/cogs.12033

Huang, L., Song, Y., Li, J., Zhen, Z., Yang, Z., & Liu, J. (2014). Individual differences in

cortical face selectivity predict behavioral performance in face recognition. *Frontiers in*

Human Neuroscience, *8*. https://doi.org/10.3389/fnhum.2014.00483

Julian, J. B., Fedorenko, E., Webster, J., & Kanwisher, N. (2012). An algorithmic method for

functionally defining regions of interest in the ventral visual pathway. *NeuroImage*, *60*(4),

2357–2364. https://doi.org/10.1016/j.neuroimage.2012.02.055

Kamps, F. S., Hendrix, C. L., Brennan, P. A., & Dilks, D. D. (2020). Connectivity at the origins

of domain specificity in the cortical face and place networks. *Proceedings of the National*

Academy of Sciences, *117*(11), 6163–6169. https://doi.org/10.1073/pnas.1911359117

Kanwisher, N. (2010). Functional specificity in the human brain: A window into the functional

architecture of the mind. *Proceedings of the National Academy of Sciences*, *107*(25),

11163–11170. https://doi.org/10.1073/pnas.1005062107

Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The Fusiform Face Area: A Module in

Human Extrastriate Cortex Specialized for Face Perception. *The Journal of Neuroscience*,

17(11), 4302–4311. https://doi.org/10.1523/JNEUROSCI.17-11-04302.1997

Kaufman, A. S., & Kaufman, N. L. (2004). *Kaufman Brief Intelligence Test (2nd ed.)*. American

Guidance Service.

Kay, J., Coltheart, M., & Lesser, R. (1992). *Psycholinguistic assessments of language processing*

in aphasia (PALPA) (1st Edition). Psychology Press.

Konkle, T., & Caramazza, A. (2013). Tripartite Organization of the Ventral Stream by Animacy

and Object Size. *Journal of Neuroscience*, *33*(25), 10235–10242.

Lerma-Usabiaga, G., Carreiras, M., & Paz-Alonso, P. M. (2018). Converging evidence for

functional and structural segregation within the left ventral occipitotemporal cortex in

reading. *Proceedings of the National Academy of Sciences*, *115*(42), E9981–E9990.

https://doi.org/10.1073/pnas.1803003115

Li, J., Huang, L., Song, Y., & Liu, J. (2017). Dissociated neural basis of two behavioral

hallmarks of holistic face processing: The whole-part effect and composite-face effect.

Neuropsychologia, *102*, 52–60. https://doi.org/10.1016/j.neuropsychologia.2017.05.026

Li, J., Osher, D. E., Hansen, H. A., & Saygin, Z. M. (2020). Innate connectivity patterns drive

the development of the visual word form area. *Scientific Reports*, *10*(1), 18039.

https://doi.org/10.1038/s41598-020-75015-7

Liu, T. T., Freud, E., Patterson, C., & Behrmann, M. (2019). Perceptual Function and Category-

Selective Neural Organization in Children with Resections of Visual Cortex. *Journal of*

Neuroscience, *39*(32), 6299–6314. https://doi.org/10.1523/JNEUROSCI.3160-18.2019

Mahon, B. Z., & Caramazza, A. (2011). What drives the organization of object knowledge in the

brain? *Trends in Cognitive Sciences*, *15*(3), 97–103.

https://doi.org/10.1016/j.tics.2011.01.004

Malach, R., Levy, I., & Hasson, U. (2002). The topography of high-order human object areas.

Trends in Cognitive Sciences, *6*(4), 176–184. https://doi.org/10.1016/S1364-

6613(02)01870-3

Martin, A. (2006). Shades of Déjerine—Forging a Causal Link between the Visual Word Form

Area and Reading. *Neuron*, *50*(2), 173–175. https://doi.org/10.1016/j.neuron.2006.04.004

Mbwana, J., Berl, M. M., Ritzl, E. K., Rosenberger, L., Mayo, J., Weinstein, S., Conry, J. A.,

Pearl, P. L., Shamim, S., Moore, E. N., Sato, S., Vezina, L. G., Theodore, W. H., &

Gaillard, W. D. (2009). Limitations to plasticity of language network reorganization in

localization related epilepsy. *Brain*, *132*(2), 347–356.

https://doi.org/10.1093/brain/awn329

McCandliss, B. D., Cohen, L., & Dehaene, S. (2003). The visual word form area: Expertise for

reading in the fusiform gyrus. *Trends in Cognitive Sciences*, *7*(7), 293–299.

https://doi.org/10.1016/S1364-6613(03)00134-7

Morais, J., Cary, L., Alegria, J., & Bertelson, P. (1979). Does awareness of speech as a sequence

of phones arise spontaneously? *Cognition*, *7*(4), 323–331. https://doi.org/10.1016/0010-

0277(79)90020-9

Nasr, S., Echavarria, C. E., & Tootell, R. B. H. (2014). Thinking Outside the Box: Rectilinear

Shapes Selectively Activate Scene-Selective Cortex. *Journal of Neuroscience*, *34*(20),

6721–6735. https://doi.org/10.1523/JNEUROSCI.4802-13.2014

Newport, E. L., Landau, B., Seydell-Greenwald, A., Turkeltaub, P. E., Chambers, C. E.,

Dromerick, A. W., Carpenter, J., Berl, M. M., & Gaillard, W. D. (2017). Revisiting

Lenneberg's Hypotheses About Early Developmental Plasticity: Language Organization

After Left-Hemisphere Perinatal Stroke. *Biolinguistics*, *11*, 407–422.

Oosterhof, N. N., Connolly, A. C., & Haxby, J. V. (2016). CoSMoMVPA: Multi-Modal

Multivariate Pattern Analysis of Neuroimaging Data in Matlab/GNU Octave. *Frontiers in*

Neuroinformatics, *10*. https://doi.org/10.3389/fninf.2016.00027

Osher, D. E., Saxe, R., Koldewyn, K., Gabrieli, J. D. E., Kanwisher, N., & Saygin, Z. M. (2015).

Structural Connectivity Fingerprints Predict Cortical Selectivity for Multiple Visual

Categories across Cortex. *Cerebral Cortex*, *26*(4), 1668–1683.

Patterson, K., Nestor, P. J., & Rogers, T. T. (2007). Where do you know what you know? The

representation of semantic knowledge in the human brain. *Nature Reviews Neuroscience*,

8(12), 976–987. https://doi.org/10.1038/nrn2277

Pennington, B. F., & Bishop, D. V. M. (2008). Relations Among Speech, Language, and

Reading Disorders. *Annual Review of Psychology*, *60*(1), 283–306.

https://doi.org/10.1146/annurev.psych.60.110707.163548

Postelnicu, G., Zollei, L., & Fischl, B. (2009). Combined Volumetric and Surface Registration.

IEEE Transactions on Medical Imaging, *28*(4), 508–522.

https://doi.org/10.1109/TMI.2008.2004426

Powell, H. W. R., Parker, G. J. M., Alexander, D. C., Symms, M. R., Boulby, P. A., Wheeler-

Kingshott, C. A. M., Barker, G. J., Noppeney, U., Koepp, M. J., & Duncan, J. S. (2006).

Hemispheric asymmetries in language-related pathways: A combined functional MRI and

tractography study. *NeuroImage*, *32*(1), 388–399.

https://doi.org/10.1016/j.neuroimage.2006.03.011

Raschle, N. M., Zuk, J., & Gaab, N. (2012). Functional characteristics of developmental dyslexia

in left-hemispheric posterior brain regions predate reading onset. *Proceedings of the*

National Academy of Sciences, *109*(6), 2156–2161.

https://doi.org/10.1073/pnas.1107721109

Rauschecker, A. M., Deutsch, G. K., Ben-Shachar, M., Schwartzman, A., Perry, L. M., &

Dougherty, R. F. (2009). Reading impairment in a patient with missing arcuate fasciculus.

Neuropsychologia, *47*(1), 180–194.

https://doi.org/10.1016/j.neuropsychologia.2008.08.011

Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K., & Goebel, R. (2021). A

Probabilistic Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cerebral*

Cortex, *31*(1), 603–619. https://doi.org/10.1093/cercor/bhaa246

Rossion, B., Dricot, L., Devolder, A., Bodart, J.-M., Crommelinck, M., Gelder, B. de, &

Zoontjes, R. (2000). Hemispheric Asymmetries for Whole-Based and Part-Based Face

Processing in the Human Fusiform Gyrus. *Journal of Cognitive Neuroscience*, *12*(5),

793–802. https://doi.org/10.1162/089892900562606

Saygin, Z. M., Osher, D. E., Koldewyn, K., Reynolds, G., Gabrieli, J. D. E., & Saxe, R. (2012).

Anatomical connectivity patterns predict face selectivity in the fusiform gyrus. *Nature*

Neuroscience, *15*, 321–327.

Saygin, Z. M., Osher, D. E., Norton, E. S., Youssoufian, D. A., Beach, S. D., Feather, J., Gaab,

N., Gabrieli, J. D. E., & Kanwisher, N. (2016). Connectivity precedes function in the

development of the visual word form area. *Nature Neuroscience*, *19*(9), 1250–1255.

https://doi.org/10.1038/nn.4354

Seghier, M. L., Neufeld, N. H., Zeidman, P., Leff, A. P., Mechelli, A., Nagendran, A., Riddoch, J.

M., Humphreys, G. W., & Price, C. J. (2012). Reading without the left ventral occipito-

temporal cortex. *Neuropsychologia*, *50*(14), 3621–3635.

https://doi.org/10.1016/j.neuropsychologia.2012.09.030

Seghier, M. L., & Price, C. J. (2011). Explaining Left Lateralization for Words in the Ventral

Occipitotemporal Cortex. *Journal of Neuroscience*, *31*(41), 14745–14753.

https://doi.org/10.1523/JNEUROSCI.2238-11.2011

Shaywitz, B. A., Shaywitz, S. E., Pugh, K. R., Mencl, W. E., Fulbright, R. K., Skudlarski, P.,

Constable, R. T., Marchione, K. E., Fletcher, J. M., Lyon, G. R., & Gore, J. C. (2002).

Disruption of posterior brain systems for reading in children with developmental dyslexia.

Biological Psychiatry, *52*(2), 101–110. https://doi.org/10.1016/S0006-3223(02)01365-3

Song, Y., Zhu, Q., Li, J., Wang, X., & Liu, J. (2015). Typical and Atypical Development of

Functional Connectivity in the Face Network. *Journal of Neuroscience*, *35*(43), 14624–

14635. https://doi.org/10.1523/JNEUROSCI.0969-15.2015

Staudt, M. (2010). Reorganization after pre- and perinatal brain lesions*. *Journal of Anatomy*,

217(4), 469–474. https://doi.org/10.1111/j.1469-7580.2010.01262.x

Staudt, M., Grodd, W., Niemann, G., Wildgruber, D., Erb, M., & Krägeloh–Mann, I. (2001). Early left periventricular brain lesions induce right hemispheric organization of speech. *Neurology*, *57*(1), 122–125. https://doi.org/10.1212/WNL.57.1.122

Stevens, W. D., Kravitz, D. J., Peng, C. S., Tessler, M. H., & Martin, A. (2017). Privileged Functional Connectivity between the Visual Word Form Area and the Language System. *The Journal of Neuroscience*, *37*(21), 5288–5297. https://doi.org/10.1523/JNEUROSCI.0138-17.2017

Tadros, K., Dupuis-Roy, N., Fiset, D., Arguin, M., & Gosselin, F. (2013). Reading laterally: The cerebral hemispheric use of spatial frequencies in visual word recognition. *Journal of Vision*, *13*(1), 4–4. https://doi.org/10.1167/13.1.4

Tuckute, G., Paunov, A., Kean, H., Small, H., Mineroff, Z., Blank, I., & Fedorenko, E. (2021). Frontal language areas do not emerge in the absence of temporal language areas: A case study of an individual born without a left temporal lobe. *BioRxiv*, 2021.05.28.446230. https://doi.org/10.1101/2021.05.28.446230

Vaidya, A. R., Pujara, M. S., Petrides, M., Murray, E. A., & Fellows, L. K. (2019). Lesion Studies in Contemporary Neuroscience. *Trends in Cognitive Sciences*, *23*(8), 653–671. https://doi.org/10.1016/j.tics.2019.05.009

Van der Haegen, L., Cai, Q., & Brysbaert, M. (2012). Colateralization of Broca's area and the visual word form area in left-handers: FMRI evidence. *Brain and Language*, *122*(3), 171–178. https://doi.org/10.1016/j.bandl.2011.11.004

Wandell, B. A., Rauschecker, A. M., & Yeatman, J. D. (2012). Learning to See Words. *Annual Review of Psychology*, *63*(1), 31–53. https://doi.org/10.1146/annurev-psych-120710-100434

Williams, M. A., Dang, S., & Kanwisher, N. G. (2007). Only some spatial patterns of fMRI

response are read out in task performance. *Nature Neuroscience*, *10*(6), 685–686.

https://doi.org/10.1038/nn1900

Wilmer, J. B., Germine, L., Chabris, C. F., Chatterjee, G., Williams, M., Loken, E., Nakayama,

K., & Duchaine, B. (2010). Human face recognition ability is specific and highly

heritable. *Proceedings of the National Academy of Sciences*, *107*(11), 5238–5241.

https://doi.org/10.1073/pnas.0913053107

Yeatman, J. D., & Feldman, H. M. (2013). Neural plasticity after pre-linguistic injury to the

arcuate and superior longitudinal fasciculi. *Cortex*, *49*(1), 301–311.

https://doi.org/10.1016/j.cortex.2011.08.006

Zhu, Q., Song, Y., Hu, S., Li, X., Tian, M., Zhen, Z., Dong, Q., Kanwisher, N., & Liu, J. (2010).

Heritability of the Specific Cognitive Ability of Face Perception. *Current Biology*, *20*(2),

137–142. https://doi.org/10.1016/j.cub.2009.11.067

1092