Title: Green spotted puffer can detect an almost nontoxic TTX analog odor

using crypt olfactory sensory neurons

Running title： Green spotted puffer smell TTX analog

Authors: Takehisa Suzuki[1], Ryota Nakahigashi[2], Masaatsu Adachi[2], Toshio Nishikawa[2],

and Hideki Abe[1] *

Affiliations:

1. Laboratory of Fish Biology, Graduate School of Bioagricultural Sciences,

Nagoya University, Furo-Cho, Chikusa-Ku, Nagoya, Aichi 464-8601, Japan

2. Laboratory of Organic Chemistry, Graduate School of Bioagricultural Sciences,

Nagoya University, Furo-Cho, Chiku-Ku, Nagoya, Aichi 464-8601, Japan

† Present address: Laboratory of Heterocyclic Chemistry, Graduate School of

Pharmaceutical Sciences, Tohoku University, 6-3 Aoba, Aramaki, Aoba-ku, Sendai 980-

8578, Japan

Corresponding author's email address

habe@agr.nagoya-u.ac.jp (Hideki Abe)

Keywords

Tetrodotoxin (TTX), TTX analog, crypt olfactory sensory neurons, chemoattractant, toxic

puffer

Summary statement:

Behavioral and immunohistochemical experiments suggest that an almost nontoxic TTX

analog, 5,6,11-trideoxyTTX, acts as an olfactory chemoattractant for green spotted puffers, and

crypt-type olfactory sensory neurons detect that.

Abstract

Tetrodotoxin (TTX) is a well-known neurotoxin that functions as a defense substance for

toxic puffers. Several behavioral studies reported that TTX attracts toxic puffers belonging to

the genus *Takifugu*. Although our electrophysiological and behavioral studies showed that a

TTX analog, 5,6,11-trideoxyTTX, acts as an olfactory chemoattractant for grass puffers

(*Takifugu alboplumbeus*), it is unclear whether toxic puffers are commonly attracted to

5,6,11-trideoxyTTX, and which types of olfactory sensory neurons (OSNs) detect 5,6,11-

trideoxyTTX.

Here we investigated whether green spotted puffer (*Dichotomyctere nigroviridis*), a

phylogenetically different species from the grass puffer, is attracted to 5,6,11-trideoxyTTX.

Administration of 5,6,11-trideoxyTTX attracted green spotted puffers, but TTX or Vehicle did

not. Furthermore, immunohistochemistry of the olfactory epithelium exposed to 5,6,11-

trideoxyTTX with an antibody against phosphorylated ribosomal protein S6 (pS6), a neuronal

activity marker, labeled oval cells with apical invagination. Such oval cells were also labeled

by the antibody against S100, a specific marker of crypt OSNs. Thus, our results suggest that

5,6,11-trideoxyTTX acts as an olfactory chemoattractant that is detected by crypt-type OSNs

in the olfactory epithelium of green spotted puffers. Toxic puffers may use 5,6,11-

trideoxyTTX as an olfactory chemoattractant involved in reproduction and parental care or as

an olfactory cue of TTX- bearing organisms for effective toxification.

Introductions

Tetrodotoxin (TTX) is a well-known neurotoxin that functions as a defense substance for toxic puffers. TTX has been believed to be accumulated in toxic puffer's bodies through the food chain, originally produced by bacteria such as *Pseudoalteromonas, Pseudomonas*, and *Vibrio* (Noguchi et al., 2006). However, TTX has also been detected in various animals, such as flatworms, ribbon worms, crabs, and gobies (Noguchi and Arakwa,2008). Recently, TTX-bearing animals and their DNA were detected in the intestinal contents of toxic puffers, and toxic puffers had been eating egg plates of toxic flatworm containing highly concentrated TTX (Itoi et al., 2015, 2018; Okabe et al., 2019).

Several studies have reported that TTX attracts toxic puffers. For example, Matsumura (1995) reported that mature male grass puffers (*Takifugu alboplumbeus*) are attracted to TTX during the spawning season. Saito et al. (2000) also reported that juvenile tiger puffers (*T. rubripes*) were attracted to TTX, and Okita et al. (2013) reported that nose ablation eliminated the attractions of tiger puffers to TTX. Our electrophysiological and behavioral study has recently shown that an almost nontoxic TTX analog, 5,6,11-trideoxyTTX, acts as an olfactory chemoattractant for grass puffers (Noguchi et al., 2021). However, these studies have only been conducted on toxic puffers of the genus *Takifugu*. Thus, it is unclear whether toxic puffers are commonly attracted to 5,6,11-trideoxyTTX and which types of olfactory sensory neuron (OSN)s respond to the 5,6,11-trideoxyTTX.

In this study, we demonstrated that green spotted puffers are attracted to the 5,6,11-trideoxyTTX, and crypt-type OSNs are activated specifically by 5,6,11-trideoxyTTX. These findings provide a possibility that attraction to the 5,6,11-trideoxyTTX is a common feature of toxic puffers that may be involved in reproduction, parental care, or effective toxification of these toxic puffers.

Materials and Methods

Animals

Green spotted puffers (*Dichotomyctere nigroviridis*; standard body length 1.5–3.0 cm, bodyweight 0.5–1.5 g) were purchased from a local-dealer and maintained in 60 L tanks (28ºC; 20–30 fish per each tank) filled with artificial brackish water (ABW; Instant Oceans® Sea-Salt, Cl^- 543 mM, Na^+ 471 mM, Mg^{2+} 54.3 mM, K^+ 10.7 mM, Ca^{2+} 10 mM Br^- 0.7 mM, Sr^{2+} 0.1 mM, F^- 0.05 mM, I^- 0.002 mM, CO_3^{2-}/HCO_3^- 3.3 mM, pH 8.2 ; Instant Ocean, Blacksburg, VA)

with a specific gravity of 1.010 under 12-h light/12-h dark photoperiod. Fish were fed dried

shrimp once a day. In this study, both males and females were used without distinction. All fish

were maintained and used in accordance with the Nagoya University regulations on animal

care and use in research.

<u>Odorant solutions</u>

L-Arginine (L-Arg, Sigma-Aldrich, St. Louis, MO, USA) and TTX (for biochemistry, citrate

buffered, FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan) were dissolved in

ultrapure water and preserved as a stock solution at -30ºC. 5,6,11-trideoxyTTX was synthesized

and purified according to the methods of Adachi et al. (2014). 5,6,11-trideoxyTTX was

dissolved in a 0.025 N acetic acid solution (for behavioral assay) or ultrapure water (for

immunohistochemistry) at 10^{-3} M and stored as a stock solution at -30ºC until use.

For the behavioral experiment, stock odorant solutions were diluted with ABW

containing 0.8% methylcellulose. For immunohistochemistry, all stock solutions were diluted

with Artificial Cerebro-Spinal Fluid for teleost (ACSF; NaCl 140 mM, KCl 5 mM, $MgCl_2$ 1.3

mM, $CaCl_2$ 2.4 mM, HEPES 10 mM, glucose 10 mM; pH 7.4).

<u>Behavioral experiments</u>

An overview of the system used in the experiment is shown in Fig. 1. All behavioral

experiments were performed using three green spotted puffers at a time. First, three fish were

transferred to a circular test aquarium (radius = 12.5 cm, depth = 8 cm) filled with ABW (2 L)

prepared from deionized water and Instant Ocean® Sea-Salt and allowed to acclimate to the

experimental environment for 1–2 h. The test aquarium was surrounded by a cardboard

partition (25-cm height) so that the experimenter was invisible to the fish.

A video CCD camera (XC-ST50, Sony, Tokyo, Japan), equipped with a low distortion

C-mount lens (f = 4.4 mm; LM4NC1M, Kowa, Nagoya, Japan), was mounted vertically above

the aquarium, and the camera system was adjusted so that the aquarium filled the camera frame.

Time-lapse images (2 frames/s, 640 × 480 pixels) from the top of the test aquarium were

acquired to a laptop computer (ThinkPad T410; Lenovo, Beijing, China) up to 10 min using

image acquisition software (Micro-Manager 1.4.22; https://micro-manager.org) and a USB-

video capture unit (GV-USB2, I-O DATA DEVICE, INC., Ishikawa, Japan). Five minutes after

the start of time-lapse image acquisition, 1 mL of 2×10^{-4} M odorant solution (Vehicle [ABW

or 2.5×10^{-4} N acetic acid in ABW], L-Arg, TTX, or 5,6,11-trideoxyTTX; diluted with ABW

containing 0.8% methylcellulose) was gently administered to the center of the test aquarium

through an injection needle (21G, TERUMO Corporation, Tokyo, Japan) connected to a 1-mL

syringe via polyvinyl chloride tube (The estimated final odorant concentration would be 10^{-7}

M if the administered solution were uniformly diffused in the test aquarium).

The numeric x-y coordinates representing the movement of the fish over time (one

coordinate in a frame) were extracted from each video using a UMA tracker software

(http://ymnk13.github.io/UMATracker/; Yamanaka and Takeuchi, 2018). We first binarized the

grayscale images using appropriate filters that the UMATracker produced. We then tracked the

fish's position using the "Group tracker GMM" algorithm and exported the coordinates to a

text file. To quantify the degree of attraction, we calculated the time spent on the central (a 4

cm radius) area from the center of the test aquarium. This criterion was based on the preliminary

diffusion experiment that 0.8% methylcellulose-ABW colored with methylene blue remained

in that area even after 10 min of administration (Movie S1). We excluded the data that fish

stopped swimming longer than 5 seconds during image acquisition.

Statistics

128   Data are expressed as mean ± standard error of the mean (SEM) and represented as a dot plot

129   and a box–whisker plot (box: median and quartiles; whisker: $1.5 \times$ interquartile range).

130   Parametric data were analyzed using Welch's *t*-test. Outliers were detected and excluded using

131   Grubb's test. Statistical analysis and graph preparations in this study were performed using R

132   (version 4.0.4; https://www.r-project.org).

Immunohistochemistry

Fish were anesthetized by tricaine methanesulfonate (MS–222; 0.02%, Sigma-Aldrich) and

excised bilateral olfactory organs. The excised olfactory organs were halved by a fine blade.

The olfactory epithelium （OE） was distributed on one surface of the halved olfactory organs

and washed three times for 5 min in ACSF. Each OE was exposed to an odorant solution

(Vehicle, L-Arg, TTX, or 5,6,11-trideoxyTTX; 10^{-6} M, 1 mL, diluted with ACSF) for 5 min in

a 1.5 mL test tube. After odorant stimulation, OE was fixed for 1 h in 4% paraformaldehyde at

4ºC, rinsed in phosphate-buffered saline (PBS), and cryoprotected in 20% sucrose in PBS

overnight at 4ºC. OE was embedded in the Tissue-Tek® O.C.T. compound (Sakura Finetek

Japan Co. Ltd., Tokyo, Japan). Cross cryosection (4–6 μm) of OE mounted onto gelatin-coated

slides and dried. After rehydration in PBS, the sections were blocked in 2% normal goat serum

（NGS） in PBS with 0.05% Tween-20 (PBST) for 3h at 28ºC.

For single immunohistochemistry, sections were incubated successively in PBST with

1) an anti-phosphorylated ribosomal protein S6 (pS6) rabbit polyclonal antibody (Ser244,

Ser247 anti-pS6, 44-923G, Thermo Fisher Scientific, MA, USA; 1:5000), that was used in

mouse OSNs as validated in Sharma et al. (2017), as a neural activity marker for 2 days at 4 °C;

2) an Alexa Fluor® 488-conjugated secondary antibody (goat anti-rabbit IgG (H+L), A-11008,

Thermo Fisher Scientific, 1:500) for 1 day at 4°C; 3) 4',6-diamidino-2-phenylindole (DAPI;

1:1000 dilution) for 5 min at room temperature.

For double immunohistochemistry, sections were successively incubated with 1) a

mixture of primary antibodies, anti-pS6 mouse monoclonal antibody (Ser235, Ser236 anti-pS6,

62016S, Cell Signaling Technology, Danvers, MA; 1:5000) diluted with anti-S100 rabbit serum

solution (Envision FLEX-S-100, Agilent Technologies, Santa Clara, CA), that was used in

zebrafish crypt OSNs as validated in Germanà et al. (2004), for 2 days at 4°C; 2) a mixture of

secondary antibodies, Alexa Fluor® 555-conjugated donkey anti-mouse IgG (A-31570,

Thermo Fisher Scientific; 1:500) and biotinylated goat anti-rabbit IgG (BA1000, VECTOR

Laboratories, Burlingame, CA, USA; 1:500) for 1 day at 4°C; 3) an Alexa Fluor® 488-

conjugated streptavidin (S11223, Thermo Fisher Scientific; 1:500) for 2 h at 28 °C. The

sections were rinsed triples for 5 min in PBST after each step of incubation, mounted in a

mounting medium (10% Mowiol 4-88, 25% glycerol, 0.2 M Tris-HCl (pH 9.0), and 2.5%–3.0%

1,4-diazabicyclo [2,2,2] octane; diluted with ultrapure water), and coverslipped.

<u>Confocal microscopy</u>

Optical sections of the immunohistochemistry and propidium iodide-stained olfactory organs

were acquired using a FV1000D IX81 confocal laser-scanning microscope (Olympus, Tokyo,

Japan). All microscopic images used in this study were processed to pseudocolor stacks and

were adjusted for brightness and contrast using Fiji (https://fiji.sc/). Some confocal microscopy

images were analyzed using FluoRender (http://www.sci.utah.edu/software/fluorender.html;

Wan et al., 2012) for three-dimensional (3D) reconstructions and visualization.

Results

5,6,11-trideoxyTTX, an almost nontoxic TTX analog, attracts green spotted puffers.

To investigate the behavioral response of green spotted puffers, we first created an odorant-

induced behavior assay system indexed by the attraction to odorant stimuli administered to the

center of a circular test aquarium. Three individuals were put in the test aquarium, and

following acclimation, L-Arg (a food odor for teleost) was administered to the center of the test

aquarium (final concentration: 10^{-7} M). Before L-Arg administration, fish swam around

peripherally. However, after L-Arg administration, they frequently swam into the center of the

test aquarium (Fig. 2B). Vehicle administration did not change the swimming pattern of the

fish (Fig. 2A). Those changes in swimming patterns were shown in Movie S2. Thus, in addition

to our preliminary diffusion test using the colored odorant solution, we considered the invasion

to the test aquarium's central area (within 4 cm radius: gray circles in Fig. 2) as an indicator of

attraction toward the odorant. To quantify the attractant activity, we measured the time spent

on the central area every minute and compared the cumulative spending time at 5 min after L-

Arg administration with Vehicle control (ABW). The mean cumulative spending time in the L-

Arg group (23.7 $\pm$ 11.8 s, n = 6) was longer than that in the Vehicle group (7.8 ± 6.5 s, n =

8) at the end of the experiment (p = 0.029; Fig. 3A).

Then, we next examined the changes in swimming behavior when 5,6,11-trideoxyTTX

(final concentration: 10^{-7} M) or its Vehicle (acetic acid in ABW; final concentration: 1.25×10^{-7} N) was administered. Before Vehicle or 5,6,11-trideoxyTTX administration (-4 min, -2 min),

green spotted puffers swam around the peripherally and did not invade the central area of the

test aquarium (Fig. 2C and D, respectively). However, after the 5,6,11-trideoxyTTX

administrations (+2 min, +4 min), the fish actively swam in the central area (Fig. 2D, Movie

S3). In contrast, the fish remained swimming peripherally after the Vehicle administrations and

rarely invaded the central area (Fig. 2C). The cumulative spending time in the central area (1

min period) were calculated, and their time changes are plotted in Fig. 3B. Cumulative

spending time increased after the 5,6,11-trideoxyTTX administration. The mean cumulative

spending time of the 5,6,11-trideoxyTTX group (25.8 ± 13.0 s, n = 11) was longer than that of

the Vehicle group (9.4 ± 7.8 s, n = 12) at the end of the experimental period (p = 0.003; Fig.3B).

By contrast, when TTX was administered, the fish continued to swim around

peripherally and rarely invaded the central area (Fig. 2E, Movie S4). The cumulative spending

time in the central area was calculated, and their time changes are plotted in Fig. 3C. The mean

cumulative spending time in the TTX group (13.3 ± 8.7 s, n = 8) was longer than that in the

Vehicle group (9.4 ± 7.8 s, n = 12), but the Welch's *t*-test showed no significant difference

between the two groups (p = 0.345). These results suggested that 5,6,11-trideoxyTTX acts as

an olfactory chemoattractant in green spotted puffers as well as in grass puffers.

5,6,11-trideoxyTTX activated oval-shaped OSNs.

The olfactory organs of teleost have nasal cavities under the body surface, and their OE is

located on the floor of the nasal cavity (Olivares & Schmachtenberg, 2019). Green spotted

puffer also has olfactory organs, but their olfactory organs protrude from the dorsal head

surface with stalks and are exposed to the external environment (Fig. 4A, arrows). The stalk is

split into two fleshy flaps (Fig. 4B). The sensory epithelium (SE) is located on the inside (arrow

in Fig. 4B), and the non-sensory epithelium (NSE) is located on the outside of the flap

(arrowhead in Fig. 4B). Fig. 4C is a 3D reconstructed confocal microscopy image of the whole

mount OE stained with propidium iodide. The inside of the flap was shallowly curved, and SE

was surrounded and separated by the NSE at the edge of the flap. OSNs are densely distributed

throughout SE. The x-z confocal section shows the cross-sectional structure of SE (Fig. 4D).

Most OSNs in SE had spindle-shaped soma (Fig.4 D, arrow), although some large oval-shaped

OSNs (minor axis diameter = c.a. 10 μm) were scattered in SE (Fig. 4C and D, arrowheads).

To identify the 5,6,11-trideoxyTTX-sensitive OSN, we immunohistochemically labeled

odorant-activated OSNs using a pS6 antibody. Excised OE from green spotted puffer was

exposed to Vehicle (ACSF), L-Arg, 5,6,11-trideoxyTTX, or TTX. L-Arg activated spindle-

shaped OSNs (Fig. 5B, arrow). The cilia-like structure was frequently observed on the L-Arg

activated OSNs (Fig. 5B inset, arrowhead). 5,6,11-trideoxyTTX activated large oval-shaped

OSNs without cilia (Fig. 5C, arrow). The apical ends of 5,6,11-trideoxyTTX-sensitive OSNs

were invaginated, and their cell bodies were located in the superficial layer of OE. TTX

administration weakly labeled tiny numbers of oval-shaped OSNs (Fig. 5D, arrow). However,

Vehicle-treatment did not activate any OSNs (Fig. 5A). Besides, a small number of 5,6,11-

trideoxyTTX -sensitive OSNs were scattered over the OE, some of which were present as cell

clusters (Fig. 5E, arrows).

<u>Crypt OSNs were activated by 5,6,11-trideoxyTTX.</u>

5,6,11-trideoxyTTX-sensitive OSNs were large oval-shaped OSNs and scattered over the OE.

To identify the cellular subtype of 5,6,11-trideoxyTTX-sensitive OSNs, we performed double

immunohistochemistry of 5,6,11-trideoxyTTX-treated OE with a pS6 antibody and an S100

antibody. S100 protein is a specific marker for crypt OSNs (Germanà et al., 2004). The results

showed that both pS6 and S100 antibodies labeled oval-shaped OSNs with apical invagination.

(Fig. 6A and B). The nuclei of oval-shaped OSNs were large and located in the superficial layer

of OE (Fig. 6C, arrows). Merged confocal microscopy image indicated that 5,6,11-

trideoxyTTX-activated pS6 immunopositive OSNs are overlapped with S100 immunopositive

crypt OSNs (Fig. 6D, arrow). However, Vehicle-treatment did not activate any OSNs (Fig. 6E).

Discussion

We demonstrated that green spotted puffers are attracted to 5,6,11-trideoxyTTX, and crypt-type

OSNs detect 5,6,11-trideoxyTTX.

Previous studies reported that TTX attracts tiger puffer (*T. rubripes*, Okita et al., 2013;

Saito et al., 2000) and grass puffer (*T. alboplumbeus*, Matsumura,1995; Noguchi et al., 2021).

These puffers live in the seawater, but the green spotted puffer lives in brackish water. Many

brackish and marine Tetraodontidae species have TTX and its analogs in their body. The whole

mitogenome sequence shows that Tetraodontidae can be phylogenetically divided into four

clades (clade A~D; Yamanoue et al., 2011). All these four clades contain both toxic and

nontoxic puffers, and the genus *Takifugu* and the genus *Dichotomyctere* belong to the different

clades (clade C and D, respectively). Although further studies using pufferfish belonging to the

other clades are needed, these phylogenic data suggest that the 5,6,11-trideoxyTTX-attracting

nature appears to be a synapomorphy for marine or brackish toxic puffers belonging to these

clades. The olfactory sensing ability of 5,6,11-trideoxyTTX is probably a common feature

among toxic puffers.

In teleosts, there are four types of OSNs intermingled in OE: Ciliated, microvillous,

crypt, and Kappe OSNs; the latter has hitherto not been described in teleost other than zebrafish

(Olivares and Schmachtenberg, 2019). These OSNs are different in morphology, location of

the cell bodies in OE, and responsiveness to the odorants (Hamdani and Døving, 2007). Ciliated

OSNs are bipolar cells with a long and fine apical dendrite with a few cilia, and the cell bodies

are in the deeper layer of OE. Microvillous OSNs are also spindle-shaped bipolar cells with a

short and thick apical dendrite with microvilli, and their cell bodies are in the middle layer of

OE. Crypt OSNs are oval-shaped with few cilia and microvilli, and their cell bodies are in the

superficial layer of OE. The most striking feature of crypt OSNs is apical invaginations

(Hansen and Finger, 2000). Although ciliated OSNs and microvillous OSNs are numerously

distributed, crypt OSNs are few and scattered in OE.

5,6,11-trideoxyTTX-activated pS6-immunopositive OSNs were oval-shaped with

apical invagination, but L-Arg-activated pS6-immunopositive OSNs were bipolar cells with

cilia; thus, the shapes of each odorant-activated OSNs are different. Additionally, a small

number of 5,6,11-trideoxyTTX-activated pS6-immunopositive OSNs are scattered in OE.

These morphological and locational features suggest that 5,6,11-trideoxyTTX-activated pS6-

immunopositive OSNs are crypt OSNs, so we performed double immunohistochemistry for

pS6 and S100 on OE. Double immunohistochemistry demonstrated that 5,6,11-trideoxyTTX-

activated pS6-immunopositive OSNs were also S100-immunopositive and had an oval-shape

with apical invagination, which is a characteristic of crypt OSNs. These findings suggest that

5,6,11-trideoxyTTX-sensitive OSNs are crypt OSNs.

The functional role of crypt OSN remains unclear. Crypt OSNs respond to the major

histocompatibility complex (MHC) peptides in zebrafish (*Danio rerio*; Bienchl et al., 2016)

and amino acids in mackerel (*Trachurus symmetricus*; Vielma et al., 2008). Crypt OSNs of the

juvenile rainbow trout respond either to amino acids, bile salts, and reproductive pheromones,

whereas those of the mature rainbow trout respond to gonadal extracts and hormones from the

opposite sex (Bazaes & Schmachtenberg, 2012). The number and the density of crypt OSNs

are sex-specific; moreover, the number and location of crypt OSNs in OE varies throughout

the year (Bettini et al., 2012; Hamdani et al., 2008). The studies in crucian carp showed that

the axons of crypt OSNs extend to the ventral part of the olfactory bulb that extends their axons

to the lateral part of the medial olfactory tract (lMOT) (Hamdani & Døving, 2006), and

disconnection of lMOT reduces male reproductive behavior (Weltzien et al., 2003). Axons in

the lMOT projected to the ventromedial telencephalon and the preoptic area in goldfish (Oka

et al., 1982; von Bartheld et al., 1984). Moreover, electrical stimulations of the ventral

telencephalon and preoptic area elicit spawning acts in male landlocked sockeye salmon, and

lesions of the same areas impair courtship behavior in male goldfish (Koyama et al., 1984;

Satou et al., 1984). These findings suggest that 5,6,11-trideoxyTTX-evoked signals from crypt

OSNs may be involved in the reproduction of puffers.

5,6,11-trideoxyTTX has almost no toxicity and does not act as a defense substance

against predators because the 5,6,11-trideoxyTTX has almost no affinity to voltage-gated sodium channels (Yotsu-Yamashita et al., 1999). Toxic puffers accumulate TTX and its analogs in their bodies, but their amounts and proportions vary by species and organs. 5,6,11-trideoxyTTX is especially accumulated in the ovary, but the ratio of the amount of TTX and 5,6,11-trideoxyTTX in the ovary is double in grass puffer (Jang et al., 2010). Green spotted puffers have TTX and its analogs in their liver and gonads (Chulanetra et al., 2011), and the amount of 5,6,11-trideoxyTTX in the whole body is half the amount of TTX (Jang et al., 2010). Why are toxic puffers having such a harmless toxin?

The first hypothesis is that 5,6,11-trideoxyTTX acts as an olfactory chemoattractant for conspecific puffers. Our study suggests that 5,6,11-trideoxyTTX is detected by crypt OSNs, and crypt OSNs are thought to detect olfactory information involved in reproduction. Matsumura (1995) reported that mature male grass puffers are attracted to the extremely low concentration of TTX (1.5×10^{-11} M) during the spawning season. They gather and spawn in early summer (Motohashi et al., 2010), and TTX levels in the ovaries increase during the spawning periods (Itoi et al., 2016). In addition to grass puffers, male tiger puffers chase females before spawning (Shibata et al., 2006). Whereas little is known about the reproductive behavior of green spotted puffers, but it is observed that their eggs are attached to hard and flat submerged surfaces in shallow water and are guarded by males until hatching; moreover, adults protect fry schools by directing them to the roots of mangroves (Pethiyagoda 1991; Rüdiger and Baensch, 1991). It has also been reported that males of some freshwater puffers protect their eggs and use their fins to provide fresh water until they hatch. (Doi et al., 2014). These studies suggest that male green spotted puffers may use 5,6,11-trideoxyTTX to find a female for mating or locate their eggs for parental behavior.

The second hypothesis is that toxic puffers use 5,6,11-trideoxyTTX as an olfactory cue to find TTX-bearing organisms for feeding. Recently, toxic puffer (*Takifugu pardalis*) eggs and toxic flatworms (*Planocera multitentaculata*) DNA had been detected in the intestinal contents of grass puffers, and grass puffers eat egg plates of toxic flatworm containing highly concentrated TTX (Itoi et al., 2015; Okabe et al., 2019). In another toxic puffer (*Chelonodon patoca*), DNAs from the toxic flatworm and goby (*Yongeichthys criniger*) were detected in intestinal contents (Itoi et al., 2020). Furthermore, some toxic puffers secreted TTX from the glands in their skin when they were stimulated (Kodama et al., 1986), and predators ingested toxic puffer larva or their eggs but spat them out promptly (Itoi et al., 2014). These studies suggested that toxic puffers accumulate TTX and use it as a defense substance against predators.

Because it has been reported that TTX-bearing prey organisms, such as bivalves and gastropods (Biessy et al., 2019), ribbon worms (*Cephalothrix* cf. *simula*; Vlasenko and Magarlamov, 2020), and flatworm (*Prosthiostomum trilineatum*; Suo et al.,2021) have 5,6,11-trideoxyTTX as well as TTX. Green spotted puffer feed on mollusks, crustaceans, and other invertebrates (Rainboth, 1996), and these phyla appeared to contain various TTX-bearing organisms (Noguchi and Arakawa, 2008). Thus, green spotted puffers may use 5,6,11-trideoxyTTX as an olfactory cue to selectively feed TTX-bearing prey organisms and effectively accumulate TTX and its analogs in their body.

This study demonstrated that the green spotted puffers are attracted to 5,6,11-trideoxyTTX, and 5,6,11-trideoxyTTX activates crypt OSNs, which has been reported to sense odor information about reproduction. These results raise the possibility that the attraction to 5,6,11-trideoxyTTX is a common characteristic among toxic puffers. Additionally, recent studies on feeding behavior suggest that 5,6,11-trideoxyTTX, as an olfactory chemoattractant, may be involved in the reproduction, parental care, or effective toxification of these toxic puffers.

Acknowledgment

We would like to express our gratitude to Prof. Naoyuki Yamamoto and Drs. Maki Goto and Hanako Hagio (Nagoya University) and Dr. Kazuya Fukuda (Currently: Hiroshima Univ.) for their suggestions during experimentation.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: T.S., H.A.; Methodology: T.S., H.A., R.N., M.A., T.N.; Validation: T.S., H.A.; Formal analysis: T.S.; Investigation: T.S., H.A.; Resources: H.A.; Data curation: T.S., H.A.; Writing - original draft: T.S., H.A.; Writing - review & editing: TS, T.N., H.A.; Visualization: T.S.; Supervision: T.N., H.A.; Project administration: H.A.; Funding acquisition: H.A., T.N.

Funding

This work was supported by Grant-in-Aids for Scientific Research (19K06762 to H.A., and

17K19195 to T.N. and H.A., and 17H06406 to T.N.).

References

Adachi, M., Sakakibara, R., Satake, Y., Isobe, M. and Nishikawa, T. (2014). Synthesis of

5,6,11-trideoxytetrodotoxin. *Chem. Lett.* **43**, 1719-1721. doi:10.1246/cl.140684

Bazáes, A. and Schmachtenberg, O. (2012). Odorant tuning of olfactory crypt cells from

juvenile and adult rainbow trout. *J. Exp. Biol.* **215**, 1740–1748. doi:10.1242/jeb.067264

Bettini, S., Lazzari, M. and Franceschini, V. (2012). Quantitative analysis of crypt cell

population during postnatal development of the olfactory organ of the guppy, *Poecilia*

reticulata (Teleostei, Poecilidae), from birth to sexual maturity. *J. Exp. Biol.* **215**, 2711–

2715. doi:10.1242/jeb.069039

Biechl, D., Tietje, K., Gerlach, G. and Wullimann, M. F. (2016). Crypt cells are involved

in kin recognition in larval zebrafish. *Sci. Rep.* **6**, 24590. doi:10.1038/srep24590

Biessy, L., Boundy, M. J., Smith, K. F., Harwood, D. T., Hawes, I., Wood, S. A. (2019)

Tetrodotoxin in marine bivalves and edible gastropods: A mini-review. *Chemosphere*. **236**,

124404. doi: 10.1016/j.chemosphere.2019.124404

Chulanetra, M., Sookrung, N., Srimanote, P., Indrawattana, N., Thanongsaksrikul, J.,

Sakolvaree, Y., Chongsa-Nguan, M., Kurazono, H. and Chaicumpa, W. (2011). Toxic

marine puffer fish in Thailand Seas and tetrodotoxin they contained. *Toxins* **3**, 1249–1262.

doi:10.3390/toxins3101249

Doi, H., Yamanoue, Y., Sonoyama, T. and Ishibashi, T. (2014). Spawning of eight

Southeast Asian brackish and freshwater puffers of the genera *Tetraodon* and

Carinotetraodon in captivity. *Fish. Sci.* **81**, 291–299. doi:10.1007/s12562-014-0842-7

Germanà, A., G. Montalbano, R. Laurà, E. Ciriaco, M. E. del Valle, and José A. Vega.

(2004). S100 protein-like immunoreactivity in the crypt olfactory neurons of the adult

zebrafish. *Neurosci. Lett.* **371**, 196–198. doi:10.1016/j.neulet.2004.08.077

Hamdani, E. H. and Døving, K. B. (2006). Specific projection of the sensory crypt cells in

the olfactory system in crucian carp, *Carassius carassius*. *Chem. Senses* **31**, 63–67.

doi:10.1093/chemse/bjj006

Hamdani, E. H. and Døving, K. B. (2007). The functional organization of the fish olfactory

system. *Prog. Neurobiol.* **82**, 80–86. doi:10.1016/j.pneurobio.2007.02.007

Hamdani, E. H., Lastein, S., Gregersen, F. and Døving, K. B. (2008). Seasonal variations

in olfactory sensory neurons — fish sensitivity to sex pheromones explained? *Chem. Senses* **33**, 119–123. doi:10.1093/chemse/bjm072

Hansen, A. and Finger, T. E. (2000). Phyletic distribution of crypt-type olfactory receptor neurons in fishes. *Brain Behav. Evol.* **55**, 100–110. doi:10.1159/000006645

Itoi, S., Ishizuka, K., Mitsuoka, R., Takimoto, N., Yokoyama, N., Detake, A., Takayanagi, C., Yoshikawa, S. and Sugita, H. (2016). Seasonal changes in the tetrodotoxin content of the pufferfish *Takifugu niphobles. Toxicon* **114**, 53–58. doi:10.1016/j.toxicon.2016.02.020

Itoi, S., Kozaki, A., Komori, K., Tsunashima, T., Noguchi, S., Kawane, M. and Sugita, H. (2015). Toxic *Takifugu pardalis* eggs found in *Takifugu niphobles* gut: Implications for TTX accumulation in the pufferfish. *Toxicon* **108**, 141–146. doi:10.1016/j.toxicon.2015.10.009

Itoi, S., Sato, T., Takei, M., Yamada, R., Ogata, R., Oyama, H., Teranishi, S., Kishiki, A., Wada, T., Noguchi, K. et al. (2020). The planocerid flatworm is a main supplier of toxin to tetrodotoxin-bearing fish juveniles. *Chemosphere* **249**, 126217. doi:10.1016/j.chemosphere.2020.126217

Itoi, S., Ueda, H., Yamada, R., Takei, M., Sato, T., Oshikiri, S., Wajima, Y., Ogata, R., Oyama, H., Shitto, T. et al. (2018). Including planocerid flatworms in the diet effectively toxifies the pufferfish, *Takifugu niphobles. Sci. Rep.* **8**, 12302. doi:10.1038/s41598-018-30696-z

Itoi, S., Yoshikawa, S., Asahina, K., Suzuki, M., Ishizuka, K., Takimoto, N., Mitsuoka, R., Yokoyama, N., Detake, A., Takayanagi, C. et al. (2014). Larval pufferfish protected by maternal tetrodotoxin. *Toxicon* **78**, 35–40. doi:10.1016/j.toxicon.2013.11.003

Jang, J. H., Lee, J. S. and Yotsu-Yamashita, M. (2010). LC/MS analysis of tetrodotoxin and its deoxy analogs in the marine puffer fish *Fugu niphobles* from the southern coast of Korea, and in the brackish water puffer fishes *Tetraodon nigroviridis* and *Tetraodon biocellatus* from Southeast Asia. *Mar. Drugs* **8**, 1049–1058. doi:10.3390/md8041049

Kodama, M., Sato, S., Ogata, T., Suzuki, Y., Kaneko, T. and Aida, K. (1986). Tetrodotoxin secreting glands in the skin of puffer fishes. *Toxicon* **24**, 819–829. doi:10.1016/0041-0101(86)90107-8

Koyama, Y., Satou, M., Oka, Y. and Ueda, K. (1984). Involvement of the telencephalic hemispheres and the preoptic area in sexual behavior of the male goldfish, *Carassius*

auratus: A brain-lesion study. *Behav. Neural Biol.* **40**, 70–86. doi:10.1016/s0163-
1047(84)90182-1

Matsumura, K. (1995). Tetrodotoxin as a pheromone. *Nature* **378**, 563–564.
doi:10.1017/CBO9781107415324.004

Motohashi, E., Yoshihara, T., Doi, H. and Ando, H. (2010). Aggregating behavior of the
grass puffer, *Takifugu niphobles*, observed in aquarium during the spawning period. *Zool.*
Sci. **27**, 559–564. doi:10.2108/zsj.27.559

Noguchi, T. and Arakawa, O. (2008). Tetrodotoxin – distribution and accumulation in
aquatic organisms, and cases of human intoxication. *Mar. Drugs* **6**, 220–242.
doi:10.3390/md20080011

Noguchi, T., Arakawa, O. and Takatani, T. (2006). TTX accumulation in pufferfish. *Comp.*
Biochem. Physiol. Part D: Genomics Proteomics **1**, 145–152.
doi:10.1016/j.cbd.2005.10.006

Noguchi, Y., Suzuki, T., Matsutani, K., Nakahigashi, R., Satake, Y., Adachi, M.,
Nishikawa, T., and Abe, H. (2021). Almost nontoxic tetrodotoxin analog, 5,6,11-
trideoxytetrodotoxin, as an olfactory chemoattractant for the grass puffers.
bioRxiv doi:10.1101/2021.09.12.459881

Oka, Y., Ichikawa, M. and Ueda, K. (1982). Synaptic organization of the olfactory bulb and
central projection of the olfactory tract. In *Chemoreception in Fishes*. (ed. T. J. Hara)*, pp.
61–75. Amsterdam: Elsevier

Okabe, T., Oyama, H., Kashitani, M., Ishimaru, Y., Suo, R., Sugita, H. and Itoi, S.
(2019). Toxic flatworm egg plates serve as a possible source of tetrodotoxin for pufferfish.
Toxins **11**, 402. doi:10.3390/toxins11070402

Okita, K., Yamazaki, H., Sakiyama, K., Yamane, H., Niina, S., Takatani, T., Arakawa, O.
and Sakakura, Y. (2013). Puffer smells tetrodotoxin. *Ichthyol. Res.* **60**, 386–389.
doi:10.1007/s10228-013-0353-z

Olivares, J. and Schmachtenberg, O. (2019). An update on anatomy and function of the
teleost olfactory system. *PeerJ* **7**, e7808. doi:10.7717/peerj.7808

Pethiyagoda, R. (1991). *Freshwater fishes of Sri Lanka.* Colombo, Sri Lanka: Wildlife
Heritage Trust of Sri Lanka. pp.294-295.

Rainboth, W. (1996). *Fishes of The Cambodian Mekong. FAO species identification field*

guide for fishery purposes, Rome, Italy: FAO. pp.225-227.

Rüdiger, R. and Baensch, H. A. (1991). *Aquarium Atlas* (vol. 1). Mergus-Verlag GmbH,

Osnabrück, Germany, pp.866-867

Saito, T., Kageyu, K., Goto, H., Murakami, K. and Noguchi,T. (2000). Tetrodotoxin

attracts pufferfish ("torafugu" *Takifugu rubripes*). *Bull. Inst. Ocean. Res. Dev. Tokai Univ.*

21, 93–96.

Satou, M., Oka, Y., Kusunoki, M., Matsushima, T., Kato, M., Fujita, I. and Ueda, K.

(1984). Telencephalic and preoptic areas integrate sexual behavior in hime salmon

(landlocked red Salmon, *Oncorhynchus nerka*): Results of electrical brain stimulation

experiments. *Physiol. Behav.* **33**, 441–447. doi:10.1016/0031-9384(84)90167-7

Schmachtenberg, O. (2006). Histological and electrophysiological properties of crypt cells

from the olfactory epithelium of the marine teleost *Trachurus symmetricus. J. Comp.*

Neurol. **121**, 113–121. doi:10.1002/cne.20847

Sharma, R., Ishimaru, Y., Davison, I., Ikegami, K., Chien, M. S., You, H., Chi, Q.,

Kubota, M., Yohda, M., Ehlers, M. et al. (2017). Olfactory receptor accessory proteins

play crucial roles in receptor function and gene choice. *eLife*, **6**.

https://doi.org/10.7554/ELIFE.21895

Shibata, R., Aono, H. and Machida, M. (2006). Spawning ecology of tiger puffer, *Takifugu*

rubripes (in Japanese with English abstract). Bull. Fish. Res. Agen. (Japan) suppl. **4**,

131–135. https://www.fra.affrc.go.jp/bulletin/bull/bull-b4/18.pdf

Suo, R., Kashitani, M., Oyama, H., Adachi, M., Nakahigashi, R., Sakakibara, R.,

Nishikawa, T., Sugita, H. and Itoi, S. (2021). First detection of tetrodotoxins in the

Cotylean flatworm *Prosthiostomum trilineatum. Mar. Drugs* **19**, 40.

doi:10.3390/md19010040

Vielma, A., Ardiles, A., Delgado, L. and Schmachtenberg, O. (2008). The elusive crypt

olfactory receptor neuron: Evidence for its stimulation by amino acids and cAMP pathway

agonists. *J. Exp. Biol.* **211**, 2417–2422. doi:10.1242/jeb.018796

Vlasenko, A. E. and Magalamov, T. Y. (2020). Tetrodotoxin and its analogues in

Cephalothrix cf. simula (Nemertea: Palaeonemertea) from the sea of Japan (Peter the

Great Gulf): Intrabody distribution and secretions. *Toxins* **12**, 745.

doi:10.3390/toxins12120745

von Bartheld, C. S., Meyer, D. L., Fiebig, E. and Ebbesson, S. O. E. (1984). Central connections of the olfactory bulb in the goldfish, *Carassius auratus. Cell Tissue Res.* **238**, 475–487. doi:10.1007/BF00219862

Wan, Y., Otsuna, H., Chien, C. B. and Hansen, C. (2012). FluoRender: An Application of 2D Image Space Methods for 3D and 4D Confocal Microscopy Data Visualization in Neurobiology Research. In *2012 IEEE Pacific Visualization Symposium,* pp. 201–208. doi:10.1109/pacificvis.2012.6183592

Weltzien, F., Höglund, E., Hamdani, E. H. and Døving, K. B. (2003). Does the lateral bundle of the medial olfactory tract mediate reproductive behavior in male crucian carp? *Chem. Senses* **28**, 293–300. doi: 10.1093/chemse/28.4.293

Yamanaka, O. and Takeuchi, R. (2018). UMATracker: an intuitive image-based tracking platform, *J. Exp. Biol.* **221**, jeb182469. doi:10.1242/jeb.182469

Yamanoue, Y., Miya, M., Doi, H., Mabuchi, K., Sakai, H. and Nishida, M. (2011). Multiple invasions into freshwater by pufferfishes (Teleostei: Tetraodontidae): A mitogenomic perspective. *PLoS ONE* **6**, e17410. doi:10.1371/journal.pone.0017410

Yotsu-Yamashita, M., Sugimoto, A., Takai, A. and Yasumoto, T. (1999). Effects of specific modifications of several hydroxyls of tetrodotoxin on its affinity to rat brain membrane. *J. Pharmacol. Exp. Ther.* **289**, 1688–1696.

Figure Legends

Figure 1: **Schematic drawing of the test aquarium used in this study.**

Side view (left) and Top view (right) of the test aquarium with odorant administration syringe. Three green spotted puffers were acclimatized in the aquarium until the odorant administration from the syringe. See the Materials and Methods for details.

Figure 2: **Swimming trajectories of green spotted puffers in response to an odorant administration.**

Swimming trajectories of green spotted puffers before and after the administration of Vehicle (ABW; A), L-Arg (10^{-7} M; B), Vehicle (1.25×10^{-7} N acetic acids in ABW; C), 5,6,11-trideoxyTTX (10^{-7} M; D), and TTX (10^{-7} M; E). Each panel represents the swimming trajectories of three fish in 2 min under each condition, and the time scale added on the top is the elapsed time with the start of odorant administration as time 0. The outer circle represents the outer edge of the test aquarium, and the inner circle represents the central area within a 4 cm radius from the center of the aquarium. The swimming trajectories in the central area are shown as the bold line, and the trajectories in the other area are shown as gray lines.

Figure 3: **Green spotted puffers were attracted to 5,6,11-trideoxyTTX.**

A: Cumulative spending time of individual puffers in the central area of the test aquarium (1 min period) before and after the administration of L-Arg (10^{-7} M; gray box; n=6) or its Vehicle (ABW; open box; n=8).

B: Cumulative spending time of puffers before and after the administration of 5,6,11-trideoxyTTX (10^{-7} M; gray box; n = 11) or its vehicle (1.25×10^{-7} N acetic acid; open box; n = 12).

C: Cumulative spending time of puffers before and after the administration of TTX (10^{-7} M; gray box; n = 8) or its Vehicle (1.25×10^{-7} N acetic acid; open box; n = 12).

All the X-axis represent the elapsed time with the start of the odor administration as 0.

Box plots show the median, quartiles (boxes) and $1.5 \times$ interquartile range (whiskers). Welch's t-test. *: $p < 0.05$, **: $p < 0.01$.

Figure 4: **Olfactory epithelium of the green spotted puffer.**

A: Dorsal view of the head region of green spotted puffer. Arrows: Olfactory organ. Scale

546 bar: 2 mm.

547 B: Magnified view of the olfactory organ. Arrow: Sensory epithelium, Arrowhead: Non-

548  sensory epithelium. Scale bar: 0.5 mm.

549 C: Reconstructed confocal microscopy image of the whole mount olfactory organ stained

550  by propidium iodide. SE: sensory epithelium, NSE: Non-sensory epithelium. Scale bar:

551  20 μm.

552 D: x-z confocal section of the dashed rectangular region in (C). Arrowhead: Large oval-

553  shaped OSNs. Arrow: spindle-shaped OSNs. Scale bar: 10 μm.

555 Figure 5: **5,6,11-trideoxyTTX activates oval-shaped OSNs.**

556 A~D: pS6 immunohistochemistry of OE cross-sections exposed to Vehicle (ACSF; A), L-

557  Arg (10^{-6} M; B), 5,6,11-trideoxyTTX (10^{-6} M; C), or TTX (10^{-6} M; D). Arrow:

558  activated OSNs. Scale bar: 10 μm. Insets in B and C: Magnified view of the pS6-

559  immunopositive OSNs. Arrowhead: Cilia of activated OSN. Scale bar: 2.5 μm.

560 E: Lower magnification confocal microscopy image of pS6-immunopositive OSNs in

561  OE cross-section exposed to 5,6,11-trideoxyTTX. Arrow: 5,6,11-trideoxyTTX-

562  activated OSNs. Scale bar: 50 μm.

564 Figure 6: **5,6,11-trideoxyTTX activates crypt OSNs.**

565 A~C: Reconstructed confocal microscopy images of 5,6,11-trideoxyTTX-administered OE

566  displaying pS6-immunopositive OSNs (A) or S100-immunopositive OSNs (B) or

567  stained with DAPI (C).

568 D: A merged image of A ~ C (pS6: magenta, S100: green, DAPI: blue). Each

569  immunopositive OSNs had apical invagination, and their cell bodies were located

570  superficial layer in OE. Arrow: Nucleus of 5,6,11-trideoxyTTX-activated OSNs. Scale

571  bar: 10μm.

572 E: No pS6-immunopositive OSN was seen in Vehicle (ACSF)-treated OE. Scale bar:

573  10μm.

575 Movie S1: **Diffusion test of odorant solution.**

576 0.8% methylcellulose-ABW solution (1 mL) colored with methylene blue, which was used

577 to dissolve the odorant solution, was administered from the center of the test aquarium filled

578 with 2L ABW. The diffusion area of methylene blue remained within a 4 cm radius from the

center of the aquarium (purple circle) even after 10 min.

Movie S2: **Attractive swimming behavior of green spotted puffers toward L-Arg**.

Vehicle or L-Arg solutions were administered from the center of the test aquarium at time 0
and remained in the aquarium. As a result, the fish frequently swam into the central area
(purple circle) after L-Arg administration.

Movie S3: **Attractive swimming behavior of green spotted puffers**
toward 5,6,11-trideoxyTTX.

The 5,6,11-trideoxyTTX solution was added from the center of the test aquarium at time 0
and remained in the aquarium. As a result, the fish frequently swam into the central area
(purple circle).

Movie S4: **No attractive swimming behavior of green spotted puffers toward TTX.**

TTX solution was added from the center of the test aquarium at time 0 and remained in the
aquarium. The fish continued to swim around peripherally and rarely invaded the central
area (purple circle).

CCD camera

Cardboard partition (height: 25 cm)

Circular aquarium (r =12.5 cm)

ABW (2 L)

Central area
(r = 4 cm)

Odorant solution (1 mL)