1 Photoinduced Isomerization Sampling of Retinal in
2 Bacteriorhodopsin

Zhong Ren

Department of Chemistry, University of Illinois at Chicago, Chicago, IL 60607, USA
Renz Research, Inc., Westmont, IL 60559, USA

zren@uic.edu
ORCID 0000-0001-7098-3127

Keywords
charge separation, proton pump, serial crystallography, singular value decomposition,
X-ray free electron laser

Abstract
Photoisomerization of retinoids inside a confined protein pocket represents a critical
chemical event in many important biological processes from animal vision, non-visual
light effects, to bacterial light sensing and harvesting. Light driven proton pumping in
bacteriorhodopsin entails exquisite electronic and conformational reconfigurations
during its photocycle. However, it has been a major challenge to delineate transient
molecular events preceding and following the photoisomerization of the retinal from
noisy electron density maps when varying populations of intermediates coexist and
evolve as a function of time. Here I report several distinct early photoproducts
deconvoluted from the recently observed mixtures in time-resolved serial
crystallography. This deconvolution substantially improves the quality of the electron
density maps hence demonstrates that the all-*trans* retinal undergoes extensive
isomerization sampling before it proceeds to the productive 13-*cis* configuration. Upon
light absorption, the chromophore attempts to perform *trans*-to-*cis* isomerization at
every double bond together with the stalled *anti*-to-*syn* rotations at multiple single
bonds along its polyene chain. Such isomerization sampling pushes all seven
transmembrane helices to bend outward, resulting in a transient expansion of the
retinal binding pocket, and later, a contraction due to recoiling. These ultrafast

responses observed at the atomic resolution support that the productive photoreaction
in bacteriorhodopsin is initiated by light-induced charge separation in the prosthetic
chromophore yet governed by stereoselectivity of its protein pocket. The method of a
numerical resolution of concurrent events from mixed observations is also generally
applicable.
39

Significance Statement

Photoisomerization of retinal is a critical rearrangement reaction in many important
biological processes from animal vision, non-visual light effects, to bacterial light
sensing and harvesting. It has been a major challenge to visualize rapid molecular
events preceding and following photoisomerization so that many protein functions
depending on such reaction remain vaguely understood. Here I report a direct
observation of the stereoselectivity of bacteriorhodopsin hence delineate the structural
mechanism of isomerization. Upon a light-induced charge separation, the retinal in a
flat conformation attempts to perform double bond isomerization and single bond
rotation everywhere along its polyene chain before it proceeds to the specific,
productive configuration. This observation improves our understanding on how a non-
specific attraction force could drive a specific isomerization.
52

Introduction

Bacteriorhodopsin (bR) pumps protons outward from the cytoplasm against the
concentration gradient via photoisomerization of its retinal chromophore. The trimeric
bR on the native purple membrane shares the seven transmembrane helical fold and the
same prosthetic group (Fig. S1) with large families of microbial and animal rhodopsins
(Ernst et al., 2014; Kandori, 2015). An all-*trans* retinal in the resting state is covalently
linked to Lys216 of helix G through a protonated Schiff base (SB; Fig. S1e), of which the
double bond $C_{15}=N_\zeta$ is also in *trans* (traditionally also noted as *anti* (McCarty, 1970)).
Upon absorption of a visible photon, the all-*trans* retinal in bR isomerizes efficiently and
selectively to adopt the 13-*cis* configuration (Govindjee et al., 1990; Tittor and
Oesterhelt, 1990). In contrast, an all-*trans* free retinal in organic solvents could
isomerize about various double bonds with a preference to form 11-*cis*, but much
slower (Hamm et al., 1996; Logunov et al., 1996) and with poor quantum yields
(Freedman and Becker, 1986; Koyama et al., 1991).
67

A broad consensus is that the isomerization event takes place around 450-500 fs

during the transition from a blue-shifted species I to form a red-shifted intermediate J

(Herbst, 2002; Mathies et al., 1988). This ultrafast event is explained in excited state

dynamics as a barrierless reaction, or at least with a very small activation barrier, from

the Frank-Condon point to the conical intersection between the potential energy

surfaces of the ground and excited states (Birge, 1990; Gozem et al., 2017). Various

molecular events prior to the isomerization have also been detected. Vibrational

spectroscopy showed a variety of possible motions, such as torsions about $C_{13}=C_{14}$ and

$C_{15}=N_\zeta$, H-out-of-plane wagging at C_{14}, and even protein responses (Diller et al., 1995;

Kobayashi et al., 2001). Nevertheless, the species I or a collection of species detected

before 30 fs remain in a good *trans* configuration about $C_{13}=C_{14}$ instead of a near 90°

configuration (Zhong et al., 1996). Recently, deep-UV stimulated Raman spectroscopy

revealed strong signals of Trp and Tyr motions in the protein throughout the I and J

intermediates (Tahara et al., 2019). Despite extensive studies, fundamental questions on

the photoisomerization of retinal remain unanswered at the atomic resolution. It is

assumed that the specific isomerization at $C_{13}=C_{14}$ in bR occurs faster with a high

quantum yield compared to the slower, nonspecific isomerization at multiple sites for

free retinal in solution and gas phase (Freedman and Becker, 1986; Hamm et al., 1996;

Kiefer et al., 2019; Koyama et al., 1991; Logunov et al., 1996) because the interactions

between the retinal chromophore and its protein environment somehow catalyze the

photoreaction by tuning the potential energy surfaces of the ground and/or excited

states. What is the structural basis of such tuning or catalysis? What is the quantum

mechanical force that causes the all-*trans* retinal to isomerize specifically to 13-*cis* in bR?

Why not isomerize elsewhere in bR? How is the quantum yield of this specific

isomerization enhanced by the protein compared to those of free retinal in solution?

Does any isomerization sampling occur, that is, is isomerization even attempted at other

double bonds? This work addresses these questions by solving a series of structures of

the early intermediates based on the electron density maps unscrambled from the

published serial crystallography datasets using singular value decomposition (SVD).

These structures of "pure" photoproducts at atomic resolution reveal widespread

conformational changes in all seven helices prior to the all-*trans* to 13-*cis* isomerization

and after its completion, suggesting that isomerization sampling takes place in bR,

where rapid photoisomerizations and single bond rotations are attempted everywhere

along the polyene chain of the retinal before the only successful one flips the SB at ~500

fs. The implication of these findings to the proton pumping and directional

conductance is presented in a companion paper (Ren, 2021).

Several international consortia carried out large operations of serial crystallography

at X-ray free electron lasers (XFELs). It is now possible to capture transient structural

species at room temperature in the bR photocycle as short-lived as fs (Bränden and

Neutze, 2021). Compared to cryo-trapping, authentic structural signals from these

XFEL data are expected to be greater in both amplitude and scope. However, the

signals reported so far do not appear to surpass those obtained by cryo-trapping

methods, suggesting much needed improvements in experimental protocols and data

analysis methods. Two major sources of data are used in this study (Table S1). Nogly

et al. captured retinal isomerization to 13-*cis* by the time of 10 ps and attributed its

specificity to the H-bond breaking between the SB and a water (Nogly et al., 2018).

Kovacs et al. deposited datasets at many short time delays (Kovacs et al., 2019). Those

sub-ps datasets demonstrate oscillatory signals at frequencies around 100 cm^{-1} (see

below). But no signal of conformational change, other than the extensive oscillations, is

found from these data in the transmembrane helices or the chromophore depicting the

retinal isomerization despite the original report. The essence of this work is a numerical

resolution of structural heterogeneity, a common difficulty often encountered in cryo

trapping and time-resolved crystallography. To what extend a specific structural

species can be enriched in crystals depends on the reaction kinetics governed by many

experimental parameters including but not limited to the fluence, wavelength, and

temperature of the light illumination. While it is possible to reach higher fractional

concentrations at specific time points for more stable species such as K and M states of

bR due to the ratio between the rates going into and exiting from that species, transient

species such as I and J are often poorly populated. If such structural heterogeneity is

not resolved, it is very difficult, if not impossible, to interpret the electron density maps

and to refine the intermediate structures (Ren et al., 2013). An assumption in nearly all

previous studies has been that each dataset, at a cryo temperature or at a time delay, is

derived from a mixture of a single conformer of a photoinduced species and the ground

state. Therefore, the difference map reveals a pure intermediate structure. This

assumption is far from the reality thus often leads to misinterpretation of the observed

electron density maps. This work is yet another case study to demonstrate the

application of our analytical protocol based on SVD (Methods) that makes no

assumption on how many excited intermediates that contribute to the captured signals

at each time point (Ren, 2019; Ren et al., 2013; Yang et al., 2011). More importantly, this

work showcases that our resolution of structural heterogeneity enables new mechanistic

insights into the highly dynamic chemical or biochemical processes.

Results and Discussion

A total of 22 datasets and 18 time points up to 10 ps are analyzed in this study (Table

S1). Difference Fourier maps at different time points and with respect to their

corresponding dark datasets are calculated according to the protocols previously

described (Methods). A collection of 126 difference maps at short delays ≤ 10 ps are

subjected to singular value decomposition (SVD; Methods) followed by a numerical

deconvolution using the previously established Ren rotation in a multi-dimensional

Euclidean space (Ren, 2016, 2019). Such resolution of electron density changes from

mixed photoexcited species in the time-resolved datasets results in four distinct

intermediate structures in the early photocycle (Fig. 1), which are then refined against

the reconstituted structure factor amplitudes (Table S2; Methods).

Low frequency oscillations upon photoexcitation

Ten out of 17 major components derived from the sub-ps delays of Kovacs et al. (Fig.

S2c) describe five two-dimensional oscillatory behaviors at frequencies ranging from 60

to 400 cm^{-1} (Fig. S3). Compared to a bond stretching frequency commonly observed in

vibrational spectroscopy, these oscillations are at much lower frequencies. The lowest

frequency is 61±2 cm^{-1}, that is, a period of 550±20 fs (Fig. S3a), observed for more than

an entire period, which matches exactly the oscillation detected in transient absorption

changes in visual rhodopsin (Wang et al., 1994). The second lowest frequency at 150±3

cm^{-1} is observed for more than one and a half periods, and the highest frequency

oscillation is observed for nearly six periods at 396±3 cm^{-1} (Fig. S3). Although these ten

components follow the oscillatory time dependencies nicely, they do not show any

association with the chromophore or any secondary structure of the protein (Fig. S4).

Similar oscillatory components were also extracted from the time-resolved XFEL

datasets of carbonmonoxy myoglobin (Ren, 2019). Therefore, the same conclusion

stands, that is, these low frequency vibrations induced by short laser pulses often

detected by ultrafast spectroscopy are the intrinsic property of a solvated protein

molecule (Johnson et al., 2014; Liebel et al., 2014). However, the functional relevance of

these oscillations is unclear. Interestingly, the isomerization sampling and productive

photoisomerization observed in this study occur within the first oscillatory period at the

lowest frequency. While such coincidence begs the question whether the protein

oscillation is required for isomerization (see below), direct evidence is lacking in these

XFEL data to support any functional relevance of these oscillatory signals.

These oscillations originate solely from the sub-ps datasets (Fig. S2ab yellow, green,

blue, and purple traces) of Kovacs et al. derived from excessive peak power density of

their pump laser (Kovacs et al., 2019; Miller et al., 2020). The datasets of Nogly et al. in

the same delay range carry coefficients ≈ 0 corresponding to these oscillatory

components, that is, free of oscillation (Fig. S2ab red and brown traces near the origin).

On the other hand, no dataset of Kovacs et al. features significantly nonzero coefficient

other than those corresponding the oscillatory components (Fig. 1a yellow trace near

the origin). Therefore, these datasets containing oscillations do not carry any

interpretable signal in the protein or on the chromophore, at least not in the outstanding

components identified by SVD (Fig. S2c). As Miller et al. pointed out, it is entirely

possible that the specimens were pumped onto higher potential energy surfaces due to

multiphoton absorption and the subsequent decays follow some biologically irrelevant

reaction pathways (Miller et al., 2020).

Intermediates I', I, and expansion of retinal binding pocket

In contrast to the oscillating signals, three components U_{10}, U_{14}, and U_{17} reveal strong

light-induced structural signals in terms of both extensiveness and quality (Figs. 2ab

and S5). These signals originate exclusively from a few time points of Nogly et al., too

few to fit the time dependency with exponentials. Instead, a spline fitting through these

time points gives rise to the estimated coefficients c_{10}, c_{14}, and c_{17} in the linear

combination of $c_{10}U_{10} + c_{14}U_{14} + c_{17}U_{17}$ for reconstructing the electron density maps of the

states I, J, and their respective precursors I', J' (Fig. 1a). A reconstituted difference map

of I' − bR = 2000U_{14} + 3000U_{17} (Fig. 2c) is located on the spline trajectory from the origin,

that is, bR at the time point of 0-, to the first time point of 49-406 fs (PDB entry 6g7i).

This state is denoted I' as a precursor leading to the I state judged by the time point at

~30 fs. However, this is not to say that a single species I' exists around 30 fs. Quite the

opposite, the population of the time-independent conformational species I' rises and

falls and peaks around 30 fs, while many other species during isomerization sampling

coexist with I′ at the same time (see below). The reconstituted difference map is used to

calculate a set of structure factor amplitudes that would produce this difference map of

I′ – bR (Methods). And the structure of I′ is refined against this reconstituted dataset

(beige; Figs. 2cd and S6). The same protocol is used to refine the structure of I state

(purple; Fig. S7) with a reconstituted difference map I – bR = $3300U_{10} + 2700U_{14}$ (Fig. 3d).

This SVD-dependent refinement strategy extends the commonly used method based on

an extrapolated map to another level. This newly developed method is able to refine a

structure against any linear combination of signal components while eliminating noise

and systematic error components, and components identified as other intermediate

species mixed in the data. Therefore, this method enables the refinement of an

unscrambled, hence pure, structural species (Methods).

Five geometric parameters are calculated along the long-chain chromophore in the

refined intermediate species to depict the transient conformation (Fig. 1c): 1) Atomic

displacements along the polyene chain and the lysyl side chain of Lys216 with respect

to the ground state structure demonstrate the overall amplitude of conformational

change of the chromophore (Fig. 1c top panel). 2) The retinal in the ground state is

largely flat except C_{15} and the methyl groups of C_{16} and C_{17} (Fig. S1e). Five consecutive

double bonds from C_5 to C_{14} of the retinal are largely coplanar. Atoms along the

polyene chain and the lysyl side chain of Lys216 are located on both sides of this retinal

plane. The side toward the three-fold axis of the bR trimer is called inboard; the

opposite side is outboard (see Fig. S1 for orientations in bR molecule). Distances from

the atoms along the long-chain chromophore to the retinal plane in the ground state are

calculated for all intermediates (Fig. 1c 2^{nd} panel). This parameter shows the curvature

of the refined conformation. 3) Changes in the distances from the atoms along the

chromophore to C_4 show extension or contraction of the chromophore in various

intermediates (Fig. 1c 3^{rd} panel). 4) Torsion angles of the single and double bonds along

the chromophore clearly demonstrate *syn*/*anti* and *cis*/*trans* conformation as well as

possible deviation from the ideal bond geometry in the refined intermediates (Fig. 1c 4^{th}

panel). 5) Six double bonds of the chromophore are refined to either *cis* or *trans*

configurations in the ground state and all intermediates with small deviations from the

ideal geometry. Therefore, each double bond can be fitted with a plane. The fitted

plane of each double bond tilts from the ground state as the conformation evolves

through the intermediates (Fig. 1c bottom panel). These conformational parameters

provide a good metric to evaluate and compare the intermediate species.

In contrast to the flat all-*trans* retinal chromophore in the ground state (Fig. 1c 2nd

panel), the side chain of Lys216 is highly twisted forming two near-90° single bonds

(Fig. 1c 4th panel), which results in a corner at C_ε that deviates dramatically from the

plane of the all-*trans* retinal (Fig. 1c 2nd panel). This corner and its interaction with helix

C is important to the function of this proton pump (Ren, 2021), which is a topic outside

the scope of isomerization sampling here. The refined geometry of the retinal in I'

retains a near perfect all-*trans* configuration, including the Schiff base (SB) double bond

$C_{15}=N_\zeta$, while various single bonds along the polyene chain deviate from the standard

anti conformation significantly (Fig. 1c 4th panel). The torsional deviations from *anti* are

in a descending order from the β-ionone ring to the SB. However, the largest torsional

twist to 90° is found at the single bond C_δ-C_γ in the Lys216 anchor (Fig. 1c 4th panel).

These torsional changes at single bonds are among the earliest conformational

responses to photon absorption due to the low cost in energy. On the contrary, no

immediate change is visible at the double bonds (see below). The torsional changes of

the single bonds result in an S-shaped retinal shortened by ~4% (Fig. 1c 3rd panel). The

distal segment C_6-C_{12} moves inboard up to 0.9 Å and the proximal segment C_{13}-C_ε,

including the SB, moves outboard up to 1.6 Å (Fig. 1c 1st and 2nd panels). This creased

retinal observed here at around 30 fs (Fig. 2d) is attributed to the direct consequence of

a compression under an attraction force between the β-ionone ring and the SB (see

below).

The refined structure of the I state (Fig. S7) shows that the retinal remains in near

perfect all-*trans*, including the SB, and as creased as its precursor I' (Fig. 1b). The

torsional deviations from *anti* single bonds become even more severe compared to the I'

state and remain in a descending order from the β-ionone ring to the SB (Fig. 1c 4th

panel). The major difference from its precursor is that the single bond N_ζ-C_ε now

adopts a perfect *syn* conformation, and C_δ-C_γ is restored to perfect *anti* (Figs. 1c 4th panel

and 3c). As a result, the anchor Lys216 has largely returned to its resting conformation.

This prompt restoration of the resting conformation is found important to the pumping

mechanism of bR presented in the companion paper (Ren, 2021). Such a lack of

substantial change between the ground state and the intermediate I was previously

noted by a comparison of the native retinal to a chemically locked $C_{13}=C_{14}$ double bond

(Zhong et al., 1996).

Remarkably, the major component U_{10} reconstituted into the difference map of I – bR

contains widespread signal associated with all seven helices (Fig. 3c). The reconstituted

map clearly shows collective outward motions from the center (Fig. 3d) suggesting an

expansion of the retinal binding pocket at hundreds of fs, which is confirmed by the

refined structure of the I state (Fig. 3e top panel). For example, the distances between

the C_α atoms increase by 0.8 Å between Arg82 and Phe208 and by 0.7 Å between Tyr83

and Trp182. It is noteworthy that similar signals in the protein are present in the raw

difference map calculated from the time point of 457-646 fs from Nogly et al. (6g7j) prior

to an SVD analysis (Fig. S8). The refined structure also shows the expansion to a lesser

extent (Fig. S9 top panel). But this expansion of the retinal binding pocket was not

reported (Nogly et al., 2018).

Transient bleaching at near UV of 265-280 nm was observed before 200 fs and

attributed to structural changes in the retinal skeleton and the surrounding Trp residues

(Schenkl et al., 2005). Recent deep-UV stimulated Raman spectroscopy also

demonstrated that motions of Trp and Tyr residues start to emerge at 200 fs and remain

steady until the isomerization is over at 30 ps (Tahara et al., 2019). Here the refined

structure of the I state with displaced helices and an expanded retinal binding pocket

offers an explanation for the stimulated Raman gain change at hundreds of fs.

However, it is unclear why and how such extensive protein responses take place even

before the retinal isomerization. According to the broadly accepted concept of

proteinquake, initial motions are generated at the epicenter where the chromophore

absorbs a photon and then propagated throughout the protein matrix (Ansari et al.,

1985). It is plausible that these ultrafast protein responses are the direct consequence of

isomerization sampling in a confined protein pocket. It was observed in organic

solvents using high-pressure liquid chromatography (HPLC) that all-*trans* retinal could

isomerize at various double bonds along the polyene chain to adopt 9-, 11-, and 13-*cis*

configurations, but with rather poor quantum yields (Freedman and Becker, 1986;

Koyama et al., 1991). This intrinsic property of the all-*trans* retinal would behave the

same even when it is incorporated in the protein except that the protein matrix herds

the chromophores on the right track of the productive photocycle and keeps the

concentrations of the attempted byproducts low. These byproduct conformations of the

retinal during isomerization sampling are too numerous and too minor to be observed

experimentally. Nevertheless, they cause a common effect, an expansion of its binding

pocket, since the all-*trans* retinal in the resting state is tightly boxed by massive side

chains all around (Figs. 3a and S10). Any attempt to isomerize would push against this

box one way or another. For instance, triple attempts to isomerize simultaneously at 11,

13, and 15 positions were suggested by a quantum mechanics/molecular mechanics

simulation (Altoè et al., 2010). A molecular dynamics simulation also concluded that

the protein arrests many different photoisomerization paths and funnels them into a

single, productive path (Hayashi et al., 2003). When the retinal binding pocket is

altered in mutants, the quantum yield of each isomerization byproduct is expected to

increase resulting in an impaired productive pathway (see below).

Intermediates J′, J and productive isomerization of retinal

The time point of 10 ps of Nogly et al. (6g7k) differs from the previous time point of

457-646 fs (6g7j) by negating the component of U_{10} (Fig. 1a), which leads to a restoration

of the normal retinal binding pocket in J′ from an expanded one in the I state followed

by a contraction in J (Fig. 3e bottom panel). Two time-independent structures of J′

(green; Fig. S11) and J (gray; Fig. S12) are refined with the same protocol based on the

respective reconstituted difference maps J′ – bR = $2700U_{14}$ - $1300U_{17}$ and J – bR = $-4200U_{10}$

+ $2000U_{14}$ - $300U_{17}$ (Methods). Their populations peak at the approximate time of ~700 fs

and ~20 ps, respectively. The observed contraction of the retinal binding pocket is likely

due to an elastic recoiling of the seven helical bundle following its transient expansion

caused by the isomerization sampling. A slight contraction is also present in the refined

structure of 6g7k at 10 ps (Fig. S9 bottom panel).

The creased retinal persists in J′ but starts to relax back to a flat conformation in J

(Fig. 1b and 1c 2[nd] panel). The difference map of J′ – bR clearly shows the 13-*cis*

configuration (Fig. 4a). Indeed, near perfect 13-*cis* is successfully refined in both J′ and J

structures (Fig. 1c 4[th] panel). While the SB double bond $C_{15}=N_\zeta$ is momentarily distorted

from the *trans* configuration in J′ with a torsion angle of 133°, a perfect *trans*

configuration at $C_{15}=N_\zeta$ is promptly restored in J (Fig. 1c 4[th] panel). The refined

structures of this series of early intermediates show that the SB N_ζ is rotating clockwise

in the entire process of the isomerization of I′ → I → J′ → J, if the retinal is viewed from

the proximal to distal direction. It seems that the isomerization starts in an expanded

retinal binding pocket and finishes in a tighter one. Whether the pocket expansion and

contraction are required for the productive isomerization and what role the low

frequency oscillations play in isomerization will need more time points at short delays

to further isolate the molecular events temporally.

The previous interpretations of the mixed signals of multiple intermediate species

presented a rotation of the C_{14} atom, instead of N_ζ, in the counterclockwise direction if

viewed from the proximal end of the retinal (Fig. 4b) (Kovacs et al., 2019; Nogly et al.,

2018). This rotation starts at the time point of 49-406 fs (6g7i) and proceeds

continuously through hundreds of fs to ps. At 10 ps time point (6g7k), this rotation still

does not complete and results in the remaining twist of 37° in the torsion angle of

$C_{13}=C_{14}$ (Fig. 4c bottom panel). An experimentally determined double bond with a

torsion angle of 82°, for example 6g7j, indicates that this twisted double bond

conformation is more populated than any other conformation at hundreds of fs. Such

interpretation is in direct contradiction of the current view in molecular dynamics. In

theory, a double bond conformation spends most time in either energy well of *trans* or

cis. The transition time between these energy wells is relatively short. A twisted double

bond conformation cannot be significantly populated compared to the discrete

populations in *trans* and *cis*, that is, cannot be observed, unless an additional energy

well emerges near the torsion angle of 90°, which no evidence supports. The gradual

rotation of $C_{13}=C_{14}$ double bond presented in the refined structures of Nogly et al. and

Kovacs et al. was a misinterpretation of the mixed signals of multiple intermediate

species as a single conformer. The gradual rotation does not occur in isomerization,

rather it merely reflects the gradual population shift from discrete states of *trans* to *cis*.

In stark contrast, this work presents that perfect *trans* to perfect *cis* isomerization occurs

in a single move during the transition of I → J′ (Fig. 4c bottom panel). Are these two

different interpretations of the same data equivalent and inconsequential to mechanistic

understanding? The discrete states in *trans* and *cis* presented here result in two sharp

U-turns of the SB N_ζ not observed in the previous interpretations, the first one at I′

around 30 fs and the second one during I → J′ isomerization at ~500 fs (Fig. 1b inset).

The functional implication of these ultrafast SB U-turns is presented in the companion

paper (Ren, 2021).

Coulombic attraction as driving force of isomerization sampling

The fundamental questions remain: What is the driving force that causes the all-*trans*

retinal to isomerize after a photon absorption, at several double bonds with a preference

at $C_{11}=C_{12}$ if not restrained but exclusively at $C_{13}=C_{14}$ in bR? Which structural feature(s)

of the chromophore and/or its protein environment differentiate the excited state from

the ground state? How does the protein environment enhance the quantum yield and

speed up the reaction rate of the isomerization to 13-*cis* by tuning the shape of the

potential energy surfaces? When does the crossing of the conical intersection occur

along the photocycle trajectory? Here I hypothesize that a Coulombic attraction

between the β-ionone ring and the SB at the Frank-Condon point, 0+ time point,

provides the initial driving force upon a photon absorption. The electric field spectral

measurements (Mathies and Stryer, 1976) and the quantum mechanics simulation

(Nogly et al., 2018) suggested that a charge separation occurs along the polyene chain at

the excited state of bR. Such a dipole moment was also detected through a transient

bleaching signal at near UV region (Schenkl et al., 2005). It can be shown that a

plausible charge separation of ±0.1e between the β-ionone ring and the SB would cause

an attraction force > 1 pN. If calibrated with the measured range of dipole moment of

10-16 D (Mathies and Stryer, 1976), the charge separation could reach the level of ±0.16e

to ±0.26e, giving rise to an attraction force of 3.5-9 pN between the β-ionone ring and

the SB. This attraction force is evidently sufficient to crease the flat all-*trans* retinal into

an S-shape and to compress it slightly within tens of fs as observed in I′ and I states

(Figs. 1b, 1c 2nd and 3rd panels, and 2d). In the meanwhile, this very attraction force also

triggers simultaneous attempts of double bond isomerizations and single bond

rotations along the polyene chain that cause the expansion of the retinal binding pocket

as observed at hundreds of fs. Despite such observable expansion, the elastic strength

of the seven helical bundle blocks all reaction pathways of the isomerization sampling

except the productive one. Following the only successful isomerization at $C_{13}=C_{14}$, the

chromophore segment from C_{15} to C_δ is attracted to the β-ionone ring; and these two

parts become significantly closer (Fig. 1c 3rd panel). None of the single bond rotations

can complete under the restraints of the protein (Figs. 3a and S10). Especially, the

segment closer to the midpoint of the retinal is more confined due to the steric

hinderance of Thr90 and Tyr185 from the inboard and outboard sides, respectively (Fig.

3a). Therefore, the single bonds deviate from *anti* less and less towards the midpoint

(Fig. 1c 4th panel).

I could further speculate that the charge separation along the polyene chain

differentiates the excited state from the ground state. Therefore, the disappearance of

any structural effects caused by the charge separation marks the crossing of the conical

intersection. The light-induced charge separation seems to subside gradually as the

reaction proceeds beyond the K state as indicated by the slow restoration of the perfect

anti conformation and the reluctant flattening of the retinal plane (Fig. 1c 2nd and 4th

panel), where the refined structures of K and L are taken from the companion paper

(Ren, 2021). If it is safe to assume that the single bond distortion away from a perfect

anti and the extent of the crease in the retinal plane are direct measures of the charge

separation with a delay no longer than ns time scale, the observed crossing of the

conical intersection is surprisingly late during the K $\rightarrow$ L transition at a few µs. The

structural basis of tuning potential energy surface arises from the stereochemical

hinderance of the retinal binding pocket that makes all isomerization attempts and

single bond rotations energetically costly except one.

Apparently, the same charge separation and the attraction force upon photon

absorption also take place in a solution sample of free retinal. Compared to the retinal

embedded in protein, photoisomerization in solution is nonspecific, resulting in a range

of byproducts, since an isomerization at any position would bring the SB significantly

closer to the β-ionone ring. The major photoproduct is 11-*cis* among several possibilities

in solution simply because $C_{11}=C_{12}$ double bond is located at the midpoint from the β-

ionone ring to the SB thus most vulnerable to a mechanical twist. It is also

understandable that each of the byproducts could only achieve a poor quantum yield

(Freedman and Becker, 1986; Koyama et al., 1991) as rotations at multiple single bonds

driven by the same attraction force and achieving a similar kink in the polyene chain

would further sidetrack the double bond isomerizations thus diminishing their

quantum yields. Since the total quantum yield of isomerization at various double

bonds is ~1/4, it can be predicted that the total quantum yield of single bond rotations is

~3/4 in solution. Because the deeper valleys on the potential energy surface leading to

single bonds in *syn* are more energetically favorable routes than the shallower valleys

leading to *cis* double bonds. However, these byproducts due to single bond rotations

are short-lived beyond detection by HPLC as they spontaneously revert back in

solution. The protein environment in bR plays a major role in enhancing the quantum

yield and accelerating the reaction rate of the isomerization to 13-*cis* by shutting down

all other reaction pathways triggered by the same charge separation. This is further

elucidated by the mutant functions below.

Isomerization byproducts permitted by mutant protein environments

The structure of a double mutant T90A/D115A (3cod) showed little difference from the

wildtype (Joh et al., 2008) while the single mutants T90V and T90A retain < 70% and <

20% of the proton pumping activity, respectively (Marti et al., 1991; Perálvarez et al.,

2001). These observations illustrate that some nonproductive pathways of the

isomerization sampling succeed more in the altered retinal binding pocket. In the

wildtype structure, Thr90 in helix C points towards the $C_{11}=C_{12}-C_{13}-C_{20}$ segment of the

retinal from the inboard with its C_γ atom 3.7 Å from the retinal plane (Fig. 3a). Given

the van der Waals radius r_C of 1.7 Å, only 0.3 Å is spared for the H atoms of the C_γ

methyl group thereby effectively shutting down the nonproductive pathways of the

isomerization sampling. In addition, the C_ε methyl group of Met118 and the C_δ ethyl

group of Ile119 in helix D are 3.7-4.6 Å away from the retinal segment of $C_7=C_8-C_9=C_{10}$

and C_{19} from the inboard (Fig. S10). Any motion of the retinal would have to push

helices C and D toward inboard causing an expansion of its binding pocket. Missing

the close contact in T90A increases the room to 1.9 Å for isomerization byproducts,

which would greatly reduce the quantum yield of the 13-*cis* productive isomerization

thus retain < 20% of the activity.

In addition to 13-*cis*, the retinal in the light adapted T90V mutant showed 9- and 11-

cis configurations at the occupancies of 3% and 18%, respectively, while these

configurations were not detected in light adapted wildtype (Marti et al., 1991). Then

why would a Val residue at this position with an equivalent C_γ methyl group permit the

formation of some isomerization byproducts? In wildtype bR, the side chain of Thr90

engages two strong H-bonds Trp86O-Thr90O$_\gamma$-D115O$_\delta$ so that its C_γ methyl group is

aligned toward the retinal (Fig. 3b). Without these H-bonds in T90V, the isopropyl

group of Val90 is free to adopt other rotameric positions so that neither of the C_γ methyl

groups has to point directly to the retinal, which increases the available room for the

formation of some isomerization byproducts. The conformational change caused by the

Val substitution at this position was detected by spectroscopy (Kandori et al., 1999).

Compared to the light adapted state with 3% 9-*cis* and 18% 11-*cis*, these isomerization

byproducts could reach even higher percentages during active photocycles thus reduce

the proton pumping activity below 70%.

It is important that the C_γ methyl group of Thr90 in the wildtype is pointing to

$C_{11}=C_{12}$ double bond located at the midpoint of the retinal as 11-*cis* is the most probable

byproduct (Hamm et al., 1996; Logunov et al., 1996). The potential energy surface of the

excited state in solution must feature a major branch of energetic valley leading to 11-

cis. But this valley is blocked by the stereo hinderance of Thr90 braced by the H-bonds.

From the outboard, the side chain of Tyr185 in helix F is nearly parallel to the mid

segment of the retinal plane with a distance of 3.5 Å (Fig. 3a), and Pro186 is also parallel

to the distal segment with 3.7-4.4 Å distance (Fig. S10). These close contacts of flat areas

from C_5 to C_{14} of the retinal prevents any significant motion of the retinal toward the

outboard. Even slight motions would push helix F away as observed here in the

expansion of the retinal binding pocket. The mutant Y185F largely retains the flat

contact so that its proton pumping activity does not reduce much (Hackett et al., 1987;

Mogi et al., 1987). However, it is predictable that various single mutants at this position

with smaller and smaller side chains would promote more and more isomerization

byproducts and eventually shut down proton pumping.

Two massive side chains of Trp86 and 182 from the extracellular and cytoplasmic

sides respectively do not seem to play a significant role in suppressing byproduct

formation as shown by the mutant W182F that retains the most of the wildtype activity

(Hackett et al., 1987), since the motions involved in isomerization sampling are oriented

more laterally. The transient expansion and contraction of the retinal binding pocket

(Fig. 3e) indicate that the tight box surrounds the mid-segment of the retinal (Fig. 3a) is

not completely rigid. Rather, its plasticity must carry sufficient strength to prevent

isomerization byproducts. Presumably, this strength originates from the mechanical

property of the helical bundle.

Concluding remarks

In summary, this work employs a numerical resolution from concurrent structural

events, hence substantially improves the quality of electron density maps of

unscrambled intermediate species that are otherwise difficult to observe. This approach

reveals the transient structural responses to many unsuccessful attempts of double

bond isomerization and single bond rotation. These findings underscore an important

implication, that is, a nonspecific Coulombic attraction provides the same driving force

for the isomerization sampling with and without a protein matrix. A productive

isomerization at a specific double bond is guided by the incorporation of the

chromophore in a specific protein environment. The productive pathway is selected

from numerous possibilities via stereochemical hinderance. This case study offers a

detailed example elucidating how a protein structure achieves regioselectivity in a

biochemical reaction that catalyzes a specific double bond isomerization by accelerating

its reaction rate and shutting down all other byproduct pathways. Nevertheless, this

nonspecific Coulombic attraction force may not be directly applicable to the

photoisomerization of retinal from 11-*cis* to all-*trans* in the activation of visual

rhodopsins. The commonality among microbial and animal rhodopsins perhaps exists

only at a high level in a rather abstract sense, if any. These photoreactions to and from

the all-*trans* retinal not necessarily share any common trajectory on the potential energy

surfaces. The key difference is bR as an energy convertor versus a visual rhodopsin as a

quantum detector (Lewis, 1978).

Acknowledgements

This work is supported in part by the grant R01EY024363 from National Institutes of

Health. The following database and software are used in this work: CCP4 (ccp4.ac.uk),

Coot (www2.mrc-lmb.cam.ac.uk/Personal/pemsley/coot), dynamiX™ (Renz Research,

Inc.), gnuplot (gnuplot.info), PDB (rcsb.org), PHENIX (phenix-online.org), PyMOL

(pymol.org), Python (python.org), and SciPy (scipy.org).

Competing Interests

ZR is the founder of Renz Research, Inc. that currently holds the copyright of the

computer software dynamiX™.

Data Availability

Depositions of the refined structures of I′, I, J′, and J to PDB-Dev (pdb-dev.wwpdb.org)

are underway.

542 References

Adams, P.D., Afonine, P.V., Bunkóczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.-W.,
Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive Python-based system for
macromolecular structure solution. Acta Crystallogr. D Biol. Crystallogr. *D66*, 213–221.
https://doi.org/10.1107/S0907444909052925.

Altoè, P., Cembran, A., Olivucci, M., and Garavelli, M. (2010). Aborted double bicycle-pedal
isomerization with hydrogen bond breaking is the primary event of bacteriorhodopsin proton pumping.
Proc. Natl. Acad. Sci. *107*, 20172–20177. https://doi.org/10.1073/pnas.1007000107.

Ansari, A., Berendzen, J., Bowne, S.F., Frauenfelder, H., Iben, I.E., Sauke, T.B., Shyamsunder, E., and
Young, R.D. (1985). Protein states and proteinquakes. Proc. Natl. Acad. Sci. *82*, 5000–5004.
https://doi.org/10.1073/pnas.82.15.5000.

Berman, H.M., Kleywegt, G.J., Nakamura, H., and Markley, J.L. (2012). The Protein Data Bank at 40:
Reflecting on the Past to Prepare for the Future. Structure *20*, 391–396.
https://doi.org/10.1016/j.str.2012.01.010.

Birge, R.R. (1990). Nature of the primary photochemical events in rhodopsin and bacteriorhodopsin.
Biochim. Biophys. Acta BBA - Bioenerg. *1016*, 293–327. https://doi.org/10.1016/0005-2728(90)90163-X.

Bonvin, A.M.J.J. (2021). 50 years of PDB: a catalyst in structural biology. Nat. Methods *18*, 448–449.
https://doi.org/10.1038/s41592-021-01138-y.

Brändén, G., and Neutze, R. (2021). Advances and challenges in time-resolved macromolecular
crystallography. Science *373*, eaba0954. https://doi.org/10.1126/science.aba0954.

Chandonia, J.-M., and Brenner, S.E. (2006). The impact of structural genomics: expectations and
outcomes. Science *311*, 347–351. https://doi.org/10.1126/science.1121018.

Diller, R., Maiti, S., Walker, G.C., Cowen, B.R., Pippenger, R., Bogomolni, R.A., and Hochstrasser, R.M.
(1995). Femtosecond time-resolved infrared laser study of the J–K transition of bacteriorhodopsin. Chem.
Phys. Lett. *241*, 109–115. https://doi.org/10.1016/0009-2614(95)00598-X.

Ernst, O.P., Lodowski, D.T., Elstner, M., Hegemann, P., Brown, L.S., and Kandori, H. (2014). Microbial
and animal rhodopsins: Structures, functions, and molecular mechanisms. Chem. Rev. *114*, 126–163.
https://doi.org/10.1021/cr4003769.

Freedman, K.A., and Becker, R.S. (1986). Comparative investigation of the photoisomerization of the
protonated and unprotonated n-butylamine Schiff bases of 9-cis-, 11-cis-, 13-cis-, and all-trans-retinals. J.
Am. Chem. Soc. *108*, 1245–1251. https://doi.org/10.1021/ja00266a020.

Glynn, C., and Rodriguez, J.A. (2019). Data-driven challenges and opportunities in crystallography.
Emerg. Top. Life Sci. ETLS20180177. https://doi.org/10.1042/ETLS20180177.

Govindjee, R., Balashov, S.P., and Ebrey, T.G. (1990). Quantum efficiency of the photochemical cycle of
bacteriorhodopsin. Biophys. J. *58*, 597–608. https://doi.org/10.1016/S0006-3495(90)82403-6.

Gozem, S., Luk, H.L., Schapiro, I., and Olivucci, M. (2017). Theory and simulation of the ultrafast double-
bond isomerization of biological chromophores. Chem. Rev. *117*, 13502–13565.
https://doi.org/10.1021/acs.chemrev.7b00177.

Hackett, N.R., Stern, L.J., Chao, B.H., Kronis, K.A., and Khorana, H.G. (1987). Structure-function studies
on bacteriorhodopsin. V. Effects of amino acid substitutions in the putative helix F. J. Biol. Chem. *262*,
9277–9284. https://doi.org/10.1016/S0021-9258(18)48077-5.

Hamm, P., Zurek, M., Röschinger, T., Patzelt, H., Oesterhelt, D., and Zinth, W. (1996). Femtosecond
spectroscopy of the photoisomerisation of the protonated Schiff base of all-trans retinal. Chem. Phys. Lett.
263, 613–621. https://doi.org/10.1016/S0009-2614(96)01269-9.

Hayashi, S., Tajkhorshid, E., and Schulten, K. (2003). Molecular dynamics simulation of
bacteriorhodopsin's photoisomerization using ab initio forces for the excited chromophore. Biophys. J. *85*,
1440–1449. https://doi.org/10.1016/S0006-3495(03)74576-7.

Henry, E.R., and Hofrichter, J. (1992). Singular value decomposition: Application to analysis of
experimental data. In Numerical Computer Methods, (Academic Press), pp. 129–192.

Herbst, J. (2002). Femtosecond infrared spectroscopy of bacteriorhodopsin chromophore isomerization.
Science *297*, 822–825. https://doi.org/10.1126/science.1072144.

Joh, N.H., Min, A., Faham, S., Whitelegge, J.P., Yang, D., Woods, V.L., and Bowie, J.U. (2008). Modest
stabilization by most hydrogen-bonded side-chain interactions in membrane proteins. Nature *453*, 1266–
1270. https://doi.org/10.1038/nature06977.

Johnson, P.J.M., Halpin, A., Morizumi, T., S. Brown, L., I. Prokhorenko, V., P. Ernst, O., and Miller, R.J.D.
(2014). The photocycle and ultrafast vibrational dynamics of bacteriorhodopsin in lipid nanodiscs. Phys.
Chem. Chem. Phys. *16*, 21310–21320. https://doi.org/10.1039/C4CP01826E.

Jung, Y.O., Lee, J.H., Kim, J., Schmidt, M., Moffat, K., Šrajer, V., and Ihee, H. (2013). Volume-conserving
trans–cis isomerization pathways in photoactive yellow protein visualized by picosecond X-ray
crystallography. Nat. Chem. *5*, 212–220. https://doi.org/10.1038/nchem.1565.

Kandori, H. (2015). Ion-pumping microbial rhodopsins. Front. Mol. Biosci. *2*.
https://doi.org/10.3389/fmolb.2015.00052.

Kandori, H., Kinoshita, N., Yamazaki, Y., Maeda, A., Shichida, Y., Needleman, R., Lanyi, J.K., Bizounok,
M., Herzfeld, J., Raap, J., et al. (1999). Structural change of threonine 89 upon photoisomerization in
bacteriorhodopsin as revealed by polarized FTIR spectroscopy. Biochemistry *38*, 9676–9683.
https://doi.org/10.1021/bi990713y.

Kiefer, H.V., Gruber, E., Langeland, J., Kusochek, P.A., Bochenkova, A.V., and Andersen, L.H. (2019).
Intrinsic photoisomerization dynamics of protonated Schiff-base retinal. Nat. Commun. *10*, 1210.
https://doi.org/10.1038/s41467-019-09225-7.

Kobayashi, T., Saito, T., and Ohtani, H. (2001). Real-time spectroscopy of transition states in
bacteriorhodopsin during retinal isomerization. Nature *414*, 531–534. https://doi.org/10.1038/35107042.

Kovacs, G.N., Colletier, J.-P., Grünbein, M.L., Yang, Y., Stensitzki, T., Batyuk, A., Carbajo, S., Doak, R.B.,
Ehrenberg, D., Foucar, L., et al. (2019). Three-dimensional view of ultrafast dynamics in photoexcited
bacteriorhodopsin. Nat. Commun. *10*, 3177. https://doi.org/10.1038/s41467-019-10758-0.

Koyama, Y., Kubo, K., Komori, M., Yasuda, H., and Mukai, Y. (1991). Effect of protonation on the
isomerization properties of n-butylamine Schiff base of isomeric retinal as revealed by direct HPLC
analyses: Selection of isomerization pathways by retinal proteins. Photochem. Photobiol. *54*, 433–443.
https://doi.org/10.1111/j.1751-1097.1991.tb02038.x.

Lanyi, J.K., and Schobert, B. (2007). Structural changes in the L photointermediate of bacteriorhodopsin. J.
Mol. Biol. *365*, 1379–1392. https://doi.org/10.1016/j.jmb.2006.11.016.

Lewis, A. (1978). The molecular mechanism of excitation in visual transduction and bacteriorhodopsin.
Proc. Natl. Acad. Sci. *75*, 549–553. https://doi.org/10.1073/pnas.75.2.549.

Liebel, M., Schnedermann, C., Bassolino, G., Taylor, G., Watts, A., and Kukura, P. (2014). Direct
observation of the coherent nuclear response after the absorption of a photon. Phys. Rev. Lett. *112*,
238301. https://doi.org/10.1103/PhysRevLett.112.238301.

Liebschner, D., Afonine, P.V., Baker, M.L., Bunkóczi, G., Chen, V.B., Croll, T.I., Hintze, B., Hung, L.-W.,
Jain, S., McCoy, A.J., et al. (2019). Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix. Acta Crystallogr. Sect. Struct. Biol. *75*, 861–877.
https://doi.org/10.1107/S2059798319011471.

Logunov, S.L., Song, L., and El-Sayed, M.A. (1996). Excited-state dynamics of a protonated retinal Schiff
base in solution. J. Phys. Chem. *100*, 18586–18591. https://doi.org/10.1021/jp962046d.

Marti, T., Otto, H., Mogi, T., Rösselet, S.J., Heyn, M.P., and Khorana, H.G. (1991). Bacteriorhodopsin
mutants containing single substitutions of serine or threonine residues are all active in proton
translocation. J. Biol. Chem. *266*, 6919–6927. https://doi.org/10.1016/S0021-9258(20)89590-8.

Mathies, R., and Stryer, L. (1976). Retinal has a highly dipolar vertically excited singlet state: implications
for vision. Proc. Natl. Acad. Sci. *73*, 2169–2173. https://doi.org/10.1073/pnas.73.7.2169.

Mathies, R., Brito Cruz, C., Pollard, W., and Shank, C. (1988). Direct observation of the femtosecond
excited-state cis-trans isomerization in bacteriorhodopsin. Science *240*, 777–779.
https://doi.org/10.1126/science.3363359.

McCarty, C.G. (1970). Chapter 9 syn-anti isomerizations and rearrangements. In The Chemistry of the
Carbon-Nitrogen Double Bond, (John Wiley & Sons, Ltd), p. 363.

Miller, R.J.D., Paré-Labrosse, O., Sarracini, A., and Besaw, J.E. (2020). Three-dimensional view of ultrafast
dynamics in photoexcited bacteriorhodopsin in the multiphoton regime and biological relevance. Nat.
Commun. *11*, 1240. https://doi.org/10.1038/s41467-020-14971-0.

Mogi, T., Stern, L.J., Hackett, N.R., and Khorana, H.G. (1987). Bacteriorhodopsin mutants containing
single tyrosine to phenylalanine substitutions are all active in proton translocation. Proc. Natl. Acad. Sci.
84, 5595–5599. https://doi.org/10.1073/pnas.84.16.5595.

Nogly, P., Weinert, T., James, D., Carbajo, S., Ozerov, D., Furrer, A., Gashi, D., Borin, V., Skopintsev, P.,
Jaeger, K., et al. (2018). Retinal isomerization in bacteriorhodopsin captured by a femtosecond x-ray laser.
Science *361*, eaat0094. https://doi.org/10.1126/science.aat0094.

Perálvarez, A., Barnadas, R., Sabés, M., Querol, E., and Padrós, E. (2001). Thr90 is a key residue of the
bacteriorhodopsin proton pumping mechanism. FEBS Lett. *508*, 399–402. https://doi.org/10.1016/S0014-
5793(01)03080-0.

Ren, Z. (2013a). Reaction trajectory revealed by a joint analysis of Protein Data Bank. PLoS ONE *8*, e77141.
https://doi.org/10.1371/journal.pone.0077141.

Ren, Z. (2013b). Reverse engineering the cooperative machinery of human hemoglobin. PLoS ONE *8*,
e77363. https://doi.org/10.1371/journal.pone.0077363.

Ren, Z. (2016). Molecular events during translocation and proofreading extracted from 200 static
structures of DNA polymerase. Nucleic Acids Res. *6*, 1–13. https://doi.org/10.1093/nar/gkw555.

Ren, Z. (2019). Ultrafast structural changes decomposed from serial crystallographic data. J. Phys. Chem.
Lett. *10*, 7148–7163. https://doi.org/10.1021/acs.jpclett.9b02375.

Ren, Z. (2021). Directional proton conductance in bacteriorhodopsin is driven by concentration gradient,
not affinity gradient. BioRxiv doi:10.1101/2021.10.04.463074. https://doi.org/10.1101/2021.10.04.463074.

Ren, Z., Perman, B., Srajer, V., Teng, T.-Y., Pradervand, C., Bourgeois, D., Schotte, F., Ursby, T., Kort, R.,
Wulff, M., et al. (2001). A molecular movie at 1.8 Å resolution displays the photocycle of photoactive
yellow protein, a eubacterial blue-light receptor, from nanoseconds to seconds. Biochemistry *40*, 13788–
13801. https://doi.org/10.1021/bi0107142.

Ren, Z., Chan, P.W.Y., Moffat, K., Pai, E.F., Royer, W.E., Šrajer, V., and Yang, X. (2013). Resolution of
structural heterogeneity in dynamic crystallography. Acta Cryst *D69*, 946–959.
https://doi.org/10.1107/S0907444913003454.

Schaffer, J.E., Kukshal, V., Miller, J.J., Kitainda, V., and Jez, J.M. (2021). Beyond X-rays: an overview of
emerging structural biology methods. Emerg. Top. Life Sci. ETLS20200272.
https://doi.org/10.1042/ETLS20200272.

Schenkl, S., Mourik, F. van, Zwan, G. van der, Haacke, S., and Chergui, M. (2005). Probing the ultrafast
charge translocation of photoexcited retinal in bacteriorhodopsin. Science *309*, 917–920.
https://doi.org/10.1126/science.1111482.

Schmidt, M., Rajagopal, S., Ren, Z., and Moffat, K. (2003). Application of singular value decomposition to
the analysis of time-resolved macromolecular X-ray data. Biophys. J. *84*, 2112–2129.
https://doi.org/10.1016/S0006-3495(03)75018-8.

Schmidt, M., Graber, T., Henning, R., and Srajer, V. (2010). Five-dimensional crystallography. Acta
Crystallogr. A *66*, 198–206. https://doi.org/10.1107/S0108767309054166.

Šrajer, V., Ren, Z., Teng, T.-Y., Schmidt, M., Ursby, T., Bourgeois, D., Pradervand, C., Schildkamp, W.,
Wulff, M., and Moffat, K. (2001). Protein conformational relaxation and ligand migration in myoglobin: A
nanosecond to millisecond molecular movie from time-resolved Laue X-ray diffraction. Biochemistry *40*,
13802–13815. https://doi.org/10.1021/bi010715u.

Tahara, S., Kuramochi, H., Takeuchi, S., and Tahara, T. (2019). Protein dynamics preceding
photoisomerization of the retinal chromophore in bacteriorhodopsin revealed by deep-UV femtosecond
stimulated Raman spectroscopy. J. Phys. Chem. Lett. *10*, 5422–5427.
https://doi.org/10.1021/acs.jpclett.9b02283.

Tittor, J., and Oesterhelt, D. (1990). The quantum yield of bacteriorhodopsin. FEBS Lett. *263*, 269–273.
https://doi.org/10.1016/0014-5793(90)81390-A.

Ursby, T., and Bourgeois, D. (1997). Improved estimation of structure-factor difference amplitudes from
poorly accurate data. Acta Crystallogr. A *53*, 564–575. https://doi.org/10.1107/S0108767397004522.

Wang, Q., Schoenlein, R.W., Peteanu, L.A., Mathies, R.A., and Shank, C.V. (1994). Vibrationally coherent
photochemistry in the femtosecond primary event of vision. Science *266*, 422–424.
https://doi.org/10.1126/science.7939680.

Yang, X., Ren, Z., Kuk, J., and Moffat, K. (2011). Temperature-scan cryocrystallography reveals reaction
intermediates in bacteriophytochrome. Nature *479*, 428–432. https://doi.org/10.1038/nature10506.

Zhong, Q., Ruhman, S., Ottolenghi, M., Sheves, M., Friedman, N., Atkinson, G.H., and Delaney, J.K.
(1996). Reexamining the primary light-induced events in bacteriorhodopsin using a synthetic C13=C14-
locked chromophore. J Am Chem Soc *118*, 12828–12829. https://doi.org/10.1021/ja961058+.

703
704

705 **Figures and Legends**

Figure 1. Intermediate species in early photocycle identified in SVD space. (a) Multi-

dimensional spaces of SVD. The SVD analysis of difference Fourier maps at short

delays ≤ 10 ps results in time-dependent coefficients $c_k(t)$, where k = 1, 2, …, each

corresponding to a time-independent components U_k. Each raw difference map at a

time delay t can be closely represented by a linear combination of these components,

$c_1(t)U_1 + c_2(t)U_2 + …$, that is called a reconstituted difference map. Each of these

components U_k and the reconstituted difference maps can be rendered in the same way

as an observed difference map. The coefficient set $c_k(t)$ is therefore a trace of the

photocycle trajectory, when these time-dependent functions are plotted in a multi-

dimensional space or plotted together against the common variable of time t.

Coefficients corresponding to components U_{10}, U_{14}, and U_{17} are plotted in three

orthographical views. Three time points from Nogly et al. in small red dots contain U_{14}

equally. These time points vary in U_{10} and U_{17}. Datasets from Kovacs et al. in small

yellow dots do not carry any of these signals, therefore cluster near the origin. The

component map of U_{10} is displayed in Fig. 3c. U_{14} is displayed in Figs. 2a and S5. U_{17} is

displayed in Fig. 2b. Several apices of the spline fitting are chosen as the potential pure

states of I′, I, J′, and J marked by large dots. This choice is only an approximate due to

the insufficient number of time points observed. (b) The refined retinal conformations

compared with the resting state in white. I′, I, J′, and J are in beige, purple, green, and

bluish gray, respectively. The creased S-shape of the retinal is established in I′ (Fig. 2)

and easing gradually (Fig. 1bc). 13-*trans* in I′ and I, 13-*cis* in J′ and J are two discrete

conformations without any other conformations in the middle (c 4th panel). The inset is

a zoom-in view of the SB double bond $C_{15}=N_\zeta$. The SB N_ζ atom performs two rapid U-

turns at ~30 fs and ~500 fs. (c) Conformational parameters calculated from the refined

chromophore. The chemical structure of the chromophore on top is aligned to the

horizontal axis. Double bonds are shaded in gray. Intermediates K and L are taken

from the companion paper (Ren, 2021). 1) Atomic displacements of each intermediate

from the resting state show greater changes in the proximal segment around the SB (top

panel). See Fig. S1 for definitions of proximal, distal, inboard, outboard, and other

orientations in bR molecule. 2) A plane is least-squares fitted to C_4 through C_{14} of the

resting state as boxed on the chemical structure. The distances of all atoms to this plane

in the inboard and outboard directions show the curvature of the chromophore. The

creased retinal in early intermediates and the inboard protruding corner at C_ε in the

resting state are clearly shown (2nd panel). 3) Distances to atom C_4 are calculated for all

refined chromophores. Changes in these distances with respect to the resting state

show the shortened chromophore in I′ and I. Once isomerization to 13-*cis* occurs, the

segment from C_{15} to C_δ around the SB becomes significantly closer to the β-ionone ring

due to the Coulombic attraction force, while the distal segment of the retinal from C_{14}

and beyond stretches (3rd panel). 4) The torsion angles of single and double bonds

quantify *anti*/*syn* or *trans*/*cis* for the ground state and all intermediates (4th panel).

5) Only a single bond can be twisted with its torsion angle near 90°. A twisted double

bond would be energetically costly. Each double bond is least-squares fitted with a

plane. The interplanar angle between a double bond and the corresponding one in the

ground state measures the local tilting of the retinal (bottom panel).

Figure 2. Shortened retinal in S-shape since earliest intermediate I'. (a) Cross sections
of component map U_{14}. The middle cross section is an integration ±0.2 Å around the
curved surface through the retinal. The top cross section is an integration 1.2-1.8 Å
outboard from the retinal surface and the bottom one is an integration 0.8-1.2 Å
inboard. See Fig. S1 for definitions of inboard, outboard, proximal, distal, and other
orientations in bR molecule. Green and red indicate electron density gain and loss,
respectively. Nearly the entire retinal is in negative densities. The proximal segment
and three waters are in intense negative densities. On the other hand, strong positive
densities flank the proximal and distal segments from the outboard and inboard,

respectively. Such signal distribution results in the S-shaped retinal by the refinement

shown in (d). (b) Cross sections of component map U_{17}. The middle cross section is an

integration ±0.2 Å around the surface through the retinal. The top panel is an

integration 0.5-0.9 Å outboard and the bottom is an integration 0.8-1.2 Å inboard.

Negative and positive densities flank the retinal from the outboard and inboard,

respectively. (c) Difference map of I′ – bR reconstituted from U_{14} and U_{17} (a and b). The

map is contoured at ±3σ in green and red mesh, respectively. The opposite

displacements of the distal and proximal segments of the retinal are obvious. Extensive

signals indicate changes in the water network and Asp85 and 212. (d) Refined retinal

conformation in beige overlaid on the resting state in white. This view is orthographical

to (c). The marked distances from C_3 to N_ζ show a shortened retinal creased into an S-

shape. C_{20} methyl group is tilted 33° toward outboard from its resting state bR (Fig. 1c

bottom panel). Wat402 remains in H-bonds with both Asp85 and 212.

$I - bR = 3300\boldsymbol{U}_{10} + 2700\boldsymbol{U}_{14}$

Figure 3. Retinal binding pocket expansion and contraction. (a) Two orthographical views of the retinal tightly boxed at its middle segment. The closest contact is Thr90 and Tyr185 on the inboard and outboard sides of the retinal plane, respectively. The minimum distance between them is 7.0 Å = $4r_C$ + 0.2 Å, where r_C = 1.7 Å is the van der Waals radius of C atom. The C_γ methyl group is 3.7 Å = $2r_C$ + 0.3 Å away from the retinal plane. See also (Kandori, 2015). (b) H-bond network involving Thr90. The O_γ hydroxyl group of Thr90 is strongly H-bonded with the main chain carbonyl of Trp86 in helix C and the carboxyl group of Asp115, simultaneously. These H-bonds fixate the

side chain of Thr90 so that the C_γ methyl group is pointing to the retinal plane directly.

(c) Component map $\boldsymbol{U}_{10}$. The main chain and side chains of the protein are rendered

with ribbons and/or sticks. The retinal and Lys216 are in purple sticks. Several key

waters are in blue spheres. Parts of the structure are omitted to reveal more of the

interior. The map is contoured at $\pm 3\sigma$ in green and red, respectively. Three

orthographical views of $\boldsymbol{U}_{10}$ clearly show that the signals are distributed around the

middle segment of the molecule and taper off to both cytoplasmic and extracellular

surfaces. The signals also concentrate on all seven helices. (d) Reconstituted difference

map I – bR from $\boldsymbol{U}_{10}$ (c) and $\boldsymbol{U}_{14}$ (Figs. 2a and S5). The map is contoured at $\pm 2.5\sigma$ in

green and red mesh, respectively. The difference map at three middle segments of

helices B, C, and G show main chain displacements toward inboard or outboard as

indicated by the arrows marking the negative and positive pairs of densities. These

difference densities are the direct evidence of the expansion of the retinal binding

pocket. The refined structure of I is in purple, and the resting state is in white. (e) The

refined structures of I and J compared with the resting state viewed along the trimer

three-fold axis from the extracellular side. Atomic displacements in the main chain

from bR to I and J are color coded and marked by arrows with lengths 20× of the actual

displacements. All seven helices in I move away from the center except a small segment

in helix C showing an expansion of the retinal binding pocket (top panel). However, all

seven helices in J move closer to one another showing a contraction with respect to the

resting state bR (bottom panel). This contraction is much more significant if compared

directly with the expanded I state.

Figure 4. Photoisomerization at $C_{13}=C_{14}$ of retinal in bR. (a) Reconstituted difference

map J' – bR from $\boldsymbol{U}_{14}$ and $\boldsymbol{U}_{17}$ (Figs. 2ab and S5). The map is contoured at $\pm 3.5\sigma$ in green

and red mesh, respectively. These difference densities are the direct evidence of

isomerization at hundreds of fs. The refined structure of J' in 13-*cis* is in green. $C_{13}=C_{14}$

in perfect *cis* is shown in the circled green positive density. J' is the earliest 13-*cis*

species at hundreds of fs. (b) Published structures of Nogly et al. The chromophore

conformations in darker and darker blues are overlayed with the resting state in white.

Photoisomerization is interpreted as a gradual change in conformation from tens to

hundreds of fs to ps. Even at 10 ps, $C_{13}=C_{14}$ is not yet in perfect *cis* (Nogly et al., 2018).

(c) Conformational parameters calculated from the published structures of Nogly et al.

The chemical structure of the chromophore on top is aligned to the horizontal axis.

Double bonds are shaded in gray. 1) A plane is least-squares fitted to C_4 through C_{14} of

the resting state as boxed on the chemical structure. The distances of all atoms to this

plane in the inboard and outboard directions show slight crease of the chromophore

captured by the structures of Nogly et al. (top panel). 2) The torsion angles indicate no

significant rotation of single bonds captured by the structures of Nogly et al. (bottom

panel). However, $C_{13}=C_{14}$ is transitioning gradually from *trans* to *cis*. In contrast, the

refined structures of I and J' show a complete photoisomerization from perfect *trans* to

perfect *cis* directly.

Photoinduced Isomerization Sampling of Retinal in

Bacteriorhodopsin

Zhong Ren

Department of Chemistry, University of Illinois at Chicago, Chicago, IL 60607, USA

Renz Research, Inc., Westmont, IL 60559, USA

zren@uic.edu

ORCID 0000-0001-7098-3127

Methods

From the outset, the key presumption is that every crystallographic dataset, at a given

temperature and a given time delay after the triggering of a photochemical reaction,

captures a mixture of unknown number of intermediate species at unknown fractions.

Needless to say, all structures of the intermediates are also unknown except the

structure at the ground state that has been determined and well refined by static

crystallography. A simultaneous solution of all these unknowns requires multiple

datasets that are collected at various temperatures or time delays so that a common set

of intermediate structures are present in these datasets with variable ratios. If the

number of available datasets is far greater than the number of unknowns, a linear

system can be established to overdetermine the unknowns with the necessary

stereochemical restraints (Ren et al., 2013). The analytical methods used in this work to

achieve such overdetermination have been incrementally developed in the past years

and recently applied to another joint analysis of the datasets of carbonmonoxy

myoglobin (Ren, 2019). Time-resolved datasets collected with ultrashort pulses from an

X-ray free electron laser were successfully analyzed by these methods to visualize

electron density components that reveal transient heating, 3d electrons of the heme iron,

and global vibrational motions. This analytical strategy is recapped below.

The methodological advance in this work is the refinement of each pure

intermediate structure that has been deconvoluted from multiple mixtures. Structure

factor amplitudes of a single conformation free of heterogeneity are overdetermined.

Given the deconvoluted structure factor amplitude set of a pure state, the standard

structural refinement software with the built-in stereochemical constraints is taken full

advantage of, e.g. PHENIX (Adams et al., 2010; Liebschner et al., 2019). In case that the

computed deconvolution has not achieved a single pure structural species, the

structural refinement is expected to make such indication.

Difference Fourier maps

A difference Fourier map is synthesized from a Fourier coefficient set of F_{light}-$F_{reference}$

with the best available phase set, often from the ground state structure. Before Fourier

synthesis, F_{light} and $F_{reference}$ must be properly scaled to the same level so that the

distribution of difference values is centered at zero and not skewed either way. A

weighting scheme proven effective assumes that a greater amplitude of a difference

Fourier coefficient F_{light}-$F_{reference}$ is more likely caused by noise than by signal (Ren et al.,

2001, 2013; Šrajer et al., 2001; Ursby and Bourgeois, 1997). Both the dark and light

datasets can serve as a reference in difference maps. If a light dataset at a certain delay

is chosen as a reference, the difference map shows the changes since that delay time but

not the changes prior to that delay. However, both the dark and light datasets must be

collected in the same experiment. A cross reference from a different experimental

setting usually causes large systematic errors in the difference map that would swamp

the desired signals. Each difference map is masked 3.5 Å around the entire molecule of

bacteriorhodopsin (bR). No lipid density is analyzed.

Meta-analysis of protein structures

Structural meta-analysis based on singular value decomposition (SVD) has been

conducted in two forms. In one of them, an interatomic distance matrix is calculated

from each protein structure in a related collection. SVD of a data matrix consists of

these distance matrices enables a large-scale joint structural comparison but requires no

structural alignment (Ren, 2013a, 2013b, 2016). In the second form, SVD is performed

on a data matrix of electron densities of related protein structures (Ren, 2019; Ren et al.,

2013; Schmidt et al., 2003, 2010). Both difference electron density maps that require a

reference dataset from an isomorphous crystal form and simulated annealing omit

maps that do not require the same unit cell and space group of the crystals are possible

choices in a structural meta-analysis (Ren, 2019; Ren et al., 2013). The interatomic

distances or the electron densities that SVD is performed on are called core data. Each

distance matrix or electron density map is associated with some metadata that describe

the experimental conditions under which the core data are obtained, such as

temperature, pH, light illumination, time delay, mutation, etc. These metadata do not

enter the SVD procedure. However, they play important role in the subsequent

interpretation of the SVD result. This computational method of structural analysis takes

advantage of a mathematical, yet practical, definition of conformational space with

limited dimensionality (Ren, 2013a). Each experimentally determined structure is a

snapshot of the protein structure. A large number of such snapshots taken under a

variety of experimental conditions, the metadata, would collectively provide a survey of

the accessible conformational space of the protein structure and reveal its rection

trajectory. Such joint analytical strategy would not be effective in early years when far

fewer protein structures were determined to atomic resolution. Recent rapid growth in

protein crystallography, such as in structural genomics (Berman et al., 2012; Bonvin,

2021; Chandonia and Brenner, 2006) and in serial crystallography (Glynn and

Rodriguez, 2019; Schaffer et al., 2021), has supplied the necessarily wide sampling of

protein structures for a joint analytical strategy to come of age. The vacancies or gaps in

a conformational space between well-populated conformational clusters often

correspond to less stable transient states whose conformations are difficult to capture, if

not impossible. These conformations are often key to mechanistic understanding and

could be explored by a back calculation based on molecular distance geometry (Ren,

2013a, 2016), the chief computational algorithm in nucleic magnetic resonance

spectroscopy (NMR), and by a structure refinement based on reconstituted dataset, a

major methodological advance in this work (see below). These structures refined to

atomic resolution against reconstituted datasets may reveal short-lived intermediate

conformation hard to be captured experimentally. Unfortunately, a protein structure

refined against a reconstituted dataset currently cannot be recognized by the Protein

Data Bank (PDB). Because crystallographic refinement of a macromolecular structure is

narrowly defined as a correspondence from one dataset to one structure. A never-

observed dataset reconstituted from a collection of experimental datasets does not

match the well-established crystallographic template of PDB; let alone a refinement of

crystal structure with the NMR algorithm.

A distance matrix contains M pairwise interatomic distances of a structure in the

form of Cartesian coordinates of all observed atoms. An everyday example of distance

matrix is an intercity mileage chart appended to the road atlas. Differences in the

molecular orientation, choice of origin, and crystal lattice among all experimentally

determined structures have no contribution to the distance matrices. Due to its

symmetry, only the lower triangle is necessary. A far more intimate examination of

protein structures in PDB is a direct analysis of their electron density maps instead of

the atomic coordinates. M such (difference) electron densities, often called voxels in

computer graphics, are selected by a mask of interest. In the case of difference maps,

only the best refined protein structure in the entire collection supplies a phase set for

Fourier synthesis of electron density maps. This best structure is often the ground state

structure determined by static crystallography. Other refined atomic coordinates from

the PDB entries are not considered in the meta-analysis. That is to say, a meta-analysis

of difference electron density maps starts from the X-ray diffraction data archived in

PDB rather than the atomic coordinates interpreted from the diffraction data, which

removes any potential model bias.

Singular value decomposition of (difference) electron density maps

An electron density map, particularly a difference map as emphasized here, consists of

density values on an array of grid points within a mask of interest. All M grid points in

a three-dimensional map can be serialized into a one-dimensional sequence of density

values according to a specific protocol. It is not important what the protocol is as long

as a consistent protocol is used to serialize all maps of the same grid setting and size,

and a reverse protocol is available to erect a three-dimensional map from a sequence of

M densities. Therefore, a set of N serialized maps, also known as vectors in linear

algebra, can fill the columns of a data matrix **A** with no specific order, so that the width

of **A** is N columns, and the length is M rows. Often, $M \gg N$, thus **A** is an elongated

matrix. If a consistent protocol of serialization is used, the corresponding voxel in all N

maps occupies a single row of matrix **A**. This strict correspondence in a row of matrix

A is important. Changes of the density values in a row from one structure to another

are due to either signals, systematic errors, or noises. Although the order of columns in

matrix **A** is unimportant, needless to say, the metadata associated with each column

must remain in good bookkeeping.

SVD of the data matrix **A** results in $\mathbf{A} = \mathbf{UWV}^{\mathrm{T}}$, also known as matrix factorization.

Matrix **U** has the same shape as **A**, that is, N columns and M rows. The N columns

contain decomposed basis components U_k, known as left singular vectors of M items,

where $k = 1, 2, …, N$. Therefore, each component U_k can be erected using the reverse

protocol to form a three-dimensional map. This decomposed elemental map can be

presented in the same way as the original maps, for example, rendered in molecular

graphics software such as Coot and PyMol. It is worth noting that these decomposed

elemental maps or map components U_k are independent of any metadata. That is to

say, these components remain constant when the metadata vary. Since each left

singular vector U_k has a unit length due to the orthonormal property of SVD (see

below), that is, $|U_k| = 1$, the root mean squares (rms) of the items in a left singular

vector is $1/\sqrt{M}$ that measures the quadratic mean of the items.

The second matrix $\mathbf{W}$ is a square matrix that contains all zeros except for N positive

values on its major diagonal, known as singular values w_k. The magnitude of w_k is

considered as a weight or significance of its corresponding component U_k. The third

matrix $\mathbf{V}$ is also a square matrix of $N \times N$. Each column of $\mathbf{V}$ or row of its transpose $\mathbf{V}^{\mathrm{T}}$,

known as a right singular vector V_k, contains the relative compositions of U_k in each of

the N original maps. Therefore, each right singular vector V_k can be considered as a

function of the metadata. Right singular vectors also have the same unit length, that is,

$|V_k| = 1$. Effectively, SVD separates the constant components independent of the

metadata from the compositions that depend on the metadata.

A singular triplet denotes 1) a decomposed component U_k, 2) its singular value w_k,

and 3) the composition function V_k. Singular triplets are often sorted in a descending

order of their singular values w_k. Only a small number of n significant singular triplets

identified by the greatest singular values w_1 through w_n can be used in a linear

combination to reconstitute a set of composite maps that closely resemble the original

ones in matrix $\mathbf{A}$, where $n < N$. For example, the original map in the ith column of

matrix $\mathbf{A}$ under a certain experimental condition can be closely represented by the ith

composite map $w_1 v_{1i} U_1 + w_2 v_{2i} U_2 + … + w_n v_{ni} U_n$, where $(v_{1i}, v_{2i}, …)$ is from the ith row of

matrix $\mathbf{V}$. The coefficient set for the linear combination is redefined here as $c_{ki} = $

$w_k v_{ki}/\sqrt{M}$. The rms of the values in a map component, or the average magnitude

measured by the quadratic mean, acts as a constant scale factor that resets the modified

coefficients c_{ki} back to the original scale of the core data, such as Å for distance matrices

and $e^-/Å^3$ for electron density maps if these units are used in the original matrix $\mathbf{A}$.

Practically, an electron density value usually carries an arbitrary unit without a

calibration, which makes this scale factor unnecessary. In the linear combination $c_{1i}U_1$ +

$c_{2i}U_2$ + … + $c_{ni}U_n$, each component U_k is independent of the metadata while how much of

each component is required for the approximation, that is, c_{ki}, depends on the metadata.

1008

Excluding the components after U_n in this approximation is based on an assumption

that the singular values after w_n are very small relative to those from w_1 through w_n. As

a result, the structural information evenly distributed in all N original maps is

effectively concentrated into a far fewer number of n significant components, known as

information concentration or dimension reduction. On the other hand, the trailing

components in matrix $\mathbf{U}$ contain inconsistent fluctuations and random noises.

Excluding these components effectively rejects noises (Schmidt et al., 2003). The least-

squares property of SVD guarantees that the rejected trailing components sums up to

the least squares of the discrepancies between the original core data and the

approximation using the accepted components.

1019

1020   However, no clear boundary is guaranteed between signals, systematic errors, and

1021   noises. Systematic errors could be more significant than the desired signals. Therefore,

1022   excluding some components from 1 through n is also possible. If systematic errors are

1023   correctly identified, the reconstituted map without these significant components would

1024   no longer carry the systematic errors.

1025

The orthonormal property of SVD

The solution set of SVD must guarantee that the columns in $\mathbf{U}$ and $\mathbf{V}$, the left and right

singular vectors U_k and V_k, are orthonormal, that is, $U_h \bullet U_k = V_h \bullet V_k = 0$ (ortho) and $U_k \bullet U_k$

$= V_k \bullet V_k = 1$ (normal), where $h \neq k$ but both are from 1 to N. The orthonormal property

also holds for the row vectors. As a result, each component U_k is independent of the

other components. In other words, a component cannot be represented by a linear

combination of any other components. However, two physical or chemical parameters

in the metadata, such as temperature and pH, may cause different changes to a

structure. These changes are not necessarily orthogonal. They could exhibit some

correlation. Therefore, the decomposed components U_k not necessarily represent any

physically or chemically meaningful changes (see below).

1037

Due to the orthonormal property of SVD, an N-dimensional Euclidean space is

established, and the first n dimensions define its most significant subspace. Each

coefficient set $c_i = (c_{1i}, c_{2i}, \ldots, c_{ni})$ of the ith composite map is located in this n-dimensional

subspace. All coefficient sets for $i = 1, 2, \ldots, N$ in different linear combinations to

approximate the N original maps in a least-squares sense can be represented by N

points or vectors $c_1, c_2, \ldots, c_N$ in the Euclidean subspace. This n-dimensional subspace is

essentially the conformational space as surveyed by the jointly analyzed core data. The

conformational space is presented as scatter plots with each captured structure

represented as a dot located at a position determined by the coefficient set c_i of the ith

observed map. When the subspace has greater dimensionality than two, multiple two-

dimensional orthographical projections of the subspace are presented, such as Fig. 1a.

These scatter plots are highly informative to reveal the relationship between the

(difference) electron density maps and their metadata.

1051

If two coefficient sets $c_i \approx c_j$, they are located close to each other in the

conformational space. Therefore, these two structures i and j share two similar

conformations. Two structures located far apart from each other in the conformational

space are dissimilar in their conformations, and distinct in the compositions of the map

components. A reaction trajectory emerges in this conformational space if the temporal

order of the core data is experimentally determined (Fig. 1a). Otherwise, an order could

be assigned to these structures based on an assumed smoothness of conformational

changes along a reaction trajectory (Ren, 2013a, 2013b, 2016). Causation and

consequence of structural motions could be revealed from the order of the structures in

a series, which may further lead to structural mechanism. In addition, an off-trajectory

location in the conformational space or a location between two clusters of observed

structures represents a structure in a unique conformation that has never been

experimentally captured. Such a hypothetical structure can be refined against a

reconstituted distance matrix using molecular distance geometry (Ren, 2013a, 2013b,

2016) or a reconstituted electron density map with the method proposed below.

1067

Rotation in SVD space

Dimension reduction is indeed effective in meta-analysis of protein structures when

many datasets are evaluated at the same time. However, the default solution set of SVD

carries complicated physical and chemical meanings that are not immediately obvious.

The interpretation of a basis component U_k, that is, "what-does-it-mean", requires a

clear demonstration of the relationship between the core data and their metadata. The

outcome of SVD does not guarantee any physical meaning in a basis component.

Therefore, SVD alone provides no direct answer to "what-does-it-mean", thus its

usefulness is very limited to merely a mathematical construction. However, the

factorized set of matrices $\mathbf{U}$, $\mathbf{W}$, and $\mathbf{V}$ from SVD is not a unique solution. That is to say,

they are not the only solution to factorize matrix $\mathbf{A}$. Therefore, it is very important to

find one or more alternative solution sets that are physically meaningful to elucidate a

structural interpretation. The concept of a rotation after SVD was introduced by Henry

& Hofrichter (Henry and Hofrichter, 1992). But they suggested a protocol that fails to

preserve the orthonormal and least-squares properties of SVD. The rotation protocol

suggested by Ren incorporates the metadata into the analysis and combines with SVD

of the core data. This rotation achieves a numerical deconvolution of multiple physical

and chemical factors after a pure mathematical decomposition, and therefore, provides

a route to answer the question of "what-does-it-mean" (Ren, 2019). This rotation shall

not be confused with a rotation in the three-dimensional real space, in which a

molecular structure resides.

1089

1090   A rotation in the n-dimensional Euclidean subspace is necessary to change the

1091   perspective before a clear relationship emerges to elucidate scientific findings. It is

1092   shown below that two linear combinations are identical before and after a rotation

1093   applied to both the basis components and their coefficients in a two-dimensional

1094   subspace of h and k. That is,

1095

$$c_h U_h + c_k U_k = f_h R_h + f_k R_k, \tag{1}$$

1096

1097   where c_h and c_k are the coefficients of the basis components U_h and U_k before the

1098   rotation; and f_h and f_k are the coefficients of the rotated basis components R_h and R_k,

1099   respectively. The same Givens rotation of an angle θ is applied to both the components

1100   and their coefficients:

1101

$$\begin{cases} R_h = U_h \cos\theta - U_k \sin\theta; \\ R_k = U_h \sin\theta + U_k \cos\theta. \end{cases} \tag{2}$$

1102

1103 Obviously, the rotated components R_h and R_k remain mutually orthonormal and

1104 orthonormal to other components. And

1105

$$\begin{cases} f_h = s_h t_h = c_h \cos\theta - c_k \sin\theta; \\ f_k = s_k t_k = c_h \sin\theta + c_k \cos\theta. \end{cases} \tag{3}$$

1106

Here $s_{h|k} = \sqrt{\sum f_{h|k}^2}$ are the singular values that replace w_h and w_k, respectively, after the

rotation. They may increase or decrease compared to the original singular values so

that the descending order of the singular values no longer holds. $T_{h|k} = (t_{h|k1}, t_{h|k2}, \ldots,$

$t_{h|kN}) = (f_{h|k1}, f_{h|k2}, \ldots, f_{h|kN})/s_{h|k}$ are the right singular vectors that replace V_h and V_k,

respectively. T_h and T_k remain mutually orthonormal after the rotation and

orthonormal to other right singular vectors that are not involved in the rotation.

1113

1114   Eq. 1 holds because the dot product of two vectors does not change after both

1115 vectors rotate the same angle. To prove Eq. 1 in more detail, Eqs. 2 and 3 are combined

1116 and expanded. All cross terms of sine and cosine are self-canceled:

1117

1118 $f_h R_h + f_k R_k = (c_h \cos\theta - c_k \sin\theta)(U_h \cos\theta - U_k \sin\theta) + (c_h \sin\theta + c_k \cos\theta)(U_h \sin\theta + U_k \cos\theta)$

1119 $\qquad = c_h U_h \cos^2\theta + c_k U_k \sin^2\theta + c_h U_h \sin^2\theta + c_k U_k \cos^2\theta \pm c_h U_k \sin\theta\cos\theta \pm c_k U_h \sin\theta\cos\theta$

1120 $\qquad = c_h U_h (\cos^2\theta + \sin^2\theta) + c_k U_k (\sin^2\theta + \cos^2\theta)$

1121 $\qquad = c_h U_h + c_k U_k$

1122

A rotation in two-dimensional subspace of h and k has no effect in other dimensions,

as the orthonormal property of SVD guarantees. Multiple steps of rotations can be

carried out in many two-dimensional subspaces consecutively to achieve a multi-

dimensional rotation. A new solution set derived from a rotation retains the

orthonormal property of SVD. The rotation in the Euclidean subspace established by

SVD does not change the comparison among the core data of protein structures. Rather

it converts one solution set $\mathbf{A} = \mathbf{UWV}^T$ to other alternative solutions $\mathbf{A} = \mathbf{RST}^T$ so that

an appropriate perspective can be found to elucidate the relationship between the core

data and metadata clearly and concisely.

1132

For example, if one physical parameter could be reoriented along a single dimension

k but not involving other dimensions by a rotation, it can be convincingly shown that

the left singular vector U_k of this dimension illustrates the structural impact by this

physical parameter. Before this rotation, the same physical parameter may appear to

cause structural variations along several dimensions, which leads to a difficult

interpretation. Would a proper rotation establish a one-on-one correspondence from all

physical or chemical parameters to all the dimensions? It depends on whether each

parameter induces an orthogonal structural change, that is, whether structural

responses to different parameters are independent or correlated among one another. If

structural changes are indeed orthogonal, it should be possible to find a proper rotation

to cleanly separate them in different dimensions. Otherwise, two different rotations are

necessary to isolate two correlated responses, but one at a time.

1145

For another example, if the observed core datasets form two clusters in the

conformational space, a rotation would be desirable to separate these clusters along a

single dimension k but to align these clusters along other dimensions. Therefore, the

component U_k is clearly due to the structural transition from one cluster to the other.

Without a proper rotation, the difference between these clusters could be complicated

with multiple dimensions involved. A deterministic solution depends on whether a

clear correlation exists between the core data and metadata. A proper rotation may

require a user decision. A wrong choice of rotation may select a viewpoint that hinders

a concise conclusion. However, it would not alter the shape of the reaction trajectory,

nor create or eliminate an intrinsic structural feature. A wrong choice of rotation cannot

eliminate the fact that a large gap exists between two clusters of observed core datasets

except that these clusters are not obvious from that viewpoint. A different rotation may

reorient the perspective along another direction. But the structural conclusion would be

equivalent. See example of before and after a rotation in (Ren, 2016).

1160

This rotation procedure finally connects the core crystallographic datasets to the

metadata of experimental conditions and accomplishes the deconvolution of physical or

chemical factors that are not always orthogonal to one another after a mathematical

decomposition. SVD analysis presented in this paper employs rotations extensively

except that no distinction is made in the symbols of components and coefficients before

and after a rotation except in this section. This method is widely applicable in large-

scale structural comparisons. Furthermore, Ren rotation after SVD is not limited to

crystallography and may impact other fields wherever SVD is used. For example, SVD

is frequently applied to spectroscopic data, images, and genetic sequence data.

1170

Structural refinement against reconstituted dataset

The linear combination $\Delta\rho(t) = f_1(t)R_1 + f_2(t)R_2 + \ldots + f_n(t)R_n$ after a rotation reconstitutes

one of the observed difference maps at a specific time point t. This time-dependent

difference map depicts an ever-evolving mixture of many excited species. A

reconstituted difference map $\Delta\rho(E)$ for a time-independent, pure, excited species E =

intermediate I', I, J', and J deconvoluted from many mixtures would take the same form

except that only one or very few coefficients remain nonzero if a proper rotation has

been found (Table S2). In order to take advantage of the mature refinement software for

macromolecular structures with extensive stereochemical restraints, a set of structure

factor amplitudes is needed. Therefore, it is necessary to reconstitute a set of structure

factor amplitudes that would produce the target difference map $\Delta\rho(E)$ based on a

known structure at the ground state. First, an electron density map of the structure at

the ground state is calculated. This calculated map is used as a base map. Second, this

base map of the ground state is combined with the positive and negative densities in the

target difference map $\Delta\rho(E)$ so that the electron densities at the ground state are skewed

toward the intermediate state. Third, structure factors are calculated from the

combined map. Finally, the phase set of the calculated structure factors is discarded,

and the amplitudes are used to refine a single conformation of the intermediate species

E that $\Delta\rho(E)$ represents.

1190

This protocol following the SVD and Ren rotation of components achieves a

refinement of a pure structural species without the need of alternative conformations.

Several points are noteworthy. First, the minimization protocol in this refinement is

performed against a numerically reconstituted amplitude set that has never been

directly measured from a crystal. This reconstituted dataset could be considered as an

extrapolated dataset "on steroids" if compared to the traditional extrapolation of small

differences, such as, the Fourier coefficient set to calculate a 3Fo-2Fc map, a technique

often used to overcome a partial occupancy of an intermediate structure. An

extrapolation of small differences is not directly observed either but computed by an

exaggeration of the observed difference based on an assumption that the intermediate

state is partially occupied, such as the doubling of the observed difference in 3Fo-2Fc =

Fo + 2(Fo-Fc). In contrast to the conventional technique of extrapolation, the

deconvolution method applied here is an interpolation among many experimental

datasets rather than an extrapolation. Secondly, the deconvolution is a simultaneous

solution of multiple intermediate states mixed together instead of solving a single

excited state.

1207

Second, a map calculated from the ground state structure is chosen as the base map

instead of an experimental map such as Fo or 2Fo-Fc map. If the second step of the

protocol is skipped, that is, no difference map is combined with the ground state map,

the refinement would result in an R factor of nearly zero, since the refinement is

essentially against the calculated structure factors (bR in Table S2). This is to say, the

residuals of the refinement are solely due to the difference component instead of the

base map. This is desirable since errors in the static structure of the ground state are

gauged during its own refinement. On the other hand, if an experimental map is

chosen as a base map, the refinement R factors would reflect errors in both the base map

and the difference map, which leads to a difficulty in an objective evaluation of this

refinement protocol.

1219

Third, the combination of the base map and a difference map is intended to

represent a pure intermediate species. Therefore, alternative conformations in

structural refinement that model a mixture of species would defeat this purpose.

However, this combined map could be very noisy and may not represent a single

species without a proper rotation. This is particular the case, if the target difference

map $\Delta\rho$ is not derived from an SVD analysis and Ren rotation. The SVD analysis

identifies many density components that are inconsistent among all observed difference

maps and excludes them, which greatly reduces the noise content. Therefore, this

refinement protocol may not be very successful without an SVD analysis. Another

source of noise originates from the phase set of the structure factors. Prior to the

refinement of the intermediate structure, the phase set remains identical to that of the

ground state. This is far from the reality when an intermediate structure involves

widespread changes, such as those refined in this study. If the rotation after SVD is not

properly selected, the target difference map would remain as a mixture minus the

ground state. Therefore, the refinement of a single conformation would encounter

difficulty or significant residuals, as judged by the R factors, the residual map, and the

refined structure. A proper solution to this problem is a better SVD solution by Ren

rotation rather than alternative conformations. A successful refinement of near perfect

trans or *cis* double bonds is a good sign to indicate that the reconstituted amplitude set

after a rotation reflects a relatively homogeneous structure. If a double bond could not

be refined well to near perfect *trans* or *cis* configuration, the dataset of structure factor

amplitudes is likely from a mixture of heterogeneous configurations, which occurred

frequently in previous studies of bR and photoactive yellow protein (Jung et al., 2013;

Lanyi and Schobert, 2007; Nogly et al., 2018). It has been a great difficulty in

crystallographic refinement in general that a heterogeneous mixture of conformations

cannot be unambiguously refined even with alternative conformations. This difficulty

becomes more severe when a mixture involves more than two conformations or when

some conformations are very minor.

1248

Lastly, the refinement protocol proposed here could be carried out in the original

unit cell and space group of the crystal at the ground state. However, this is not always

applicable as the original goal of the meta-analysis is a joint examination of all available

structures from a variety of crystal forms. It would be highly desirable to evaluate

difference maps of the same or similar proteins from non-isomorphous crystals together

by SVD. Alternatively, the refinement protocol could also be performed in the space

group of P1 with a virtual unit cell large enough to hold the structure, which is the

option in this study (Table S2). This is to say, the entire analysis of SVD-rotation-

refinement presented here could be extracted and isolated from the original crystal

lattices, which paves the way to future applications to structural data acquired by

experimental techniques beyond crystallography, most attractively, to single particle

reconstruction in cryo electron microscopy.

1261

1262

1263 **Supplementary Tables**

1264

Table S1. Datasets analyzed in this work

Publication	PDB	Label	Resolution	Main conclusions	New findings in this work
Nogly et al. Science 361, eaat0094, 2018	6g7h	dark6	1.5 Å	Retinal fully isomerizes by 10 ps. But the SB water dissociates earlier.	The short-delay datasets contribute to the structures of I' → I → J' → J. Photoisomerization in J'; retinal binding pocket expansion before 1 ps in I and contraction at ~20 ps in J
	6g7i	49-406fs	1.9 Å		
	6g7j	457-646fs	1.9 Å		
	6g7k	10ps	1.9 Å		
Kovacs et al. Nat. Commun. 10, 3177, 2019	6ga1	dark1	1.7 Å	The exceedingly high power density of the pump laser causes two-photon absorption. Vibrational motions were observed.	The sub-ps datasets exhibit extensive vibrations at various frequencies. The vibrational signals are widespread over the entire bR molecule and not associated with any structural elements. Therefore, it is concluded that these global vibrations are intrinsic properties of bR induced by short laser pulses. The vibrational signals are more prominent under higher power density of the laser pulses. However, these vibrations are irrelevant to the light-driven proton pumping function of bR.
	6ga2	dark2	1.8 Å		
	6rmk	dark3	1.8 Å		
	6ga7	240fs	1.8 Å		
	6ga8	330fs	1.8 Å		
	6ga9	390fs	1.8 Å		
	6gaa	430fs	1.8 Å		
	6gab	460fs	1.8 Å		
	6gac	490fs	1.8 Å		
	6gad	530fs	1.8 Å		
	6gae	560fs	1.8 Å		
	6gaf	590fs	1.8 Å		
	6gag	630fs	1.8 Å		
	6gah	680fs	1.8 Å		
	6gai	740fs	1.8 Å		
	6ga4	1ps	1.8 Å		
	6ga5	3ps	1.9 Å		
	6ga6	10ps	1.8 Å		

1265

1266

Table S2. Refinement statistics

Intermediate	bR	I′	I	J′	J
Time period	0-	< 50 fs	40-700 fs	0.5-2 ps	1-30 ps
Coefficient c_{10}	0	0	3,300	0	-4,200
Coefficient c_{14}	0	2,000	2,700	2,700	2,000
Coefficient c_{17}	0	3,000	0	-1,300	-300
Starting model	PDB 6g7h				
Resolution range	50-2.1 Å				
Space group	P1				
Unit cell	$a = b = 62.32$ Å; $c = 111.10$ Å; $\alpha = \beta = 90°$; and $\gamma = 120°$				
Unique reflections	80,354 in working set + 4,236 in test set = 84,590 total				
Completeness	95% in working set + 5% in test set = 100% reconstituted				
R (%)	1.8	29.4	31.0	29.1	30.0
R_{free} (%)	1.9	31.1	32.4	30.4	30.7
Refined content	230 protein residues + retinal + water molecules				
Number of atoms	1,798	1,795	1,798	1,796	1,795
Water molecules	8	5	8	6	5
RMSD bonds (Å)	0.005	0.009	0.009	0.009	0.009
RMSD angles (°)	0.793	1.206	1.105	1.085	1.068
Rama. favored (%)	98.7	96.5	95.6	96.1	96.5
Rama. outliers (%)	0.0	0.0	0.4	0.4	0.4
Clash score	4	9	5	4	6

1267

1268

1269 **Supplementary Figures and Legends**

1270

Figure S1. Orientations in bacteriorhodopsin. (a) Bacteriorhodopsin (bR) trimer viewed

from the extracellular (EC) side along the three-fold axis. (b) An orthographical view to

(a) looking from the outside of the trimer. (c and d) Two orthographical views of the

retinal chromophore looking along the three-fold and normal to the three-fold axis. The

plane of retinal is largely parallel to the three-fold axis. Therefore, two sides of the

plane are called inboard and outboard with respect to the three-fold axis. The direction

toward the anchor Lys216 is called proximal. The β-ionone ring direction is therefore

distal. (e) Chemical structure of retinal incorporated to its anchor Lys216. The atom

numbers and various segment names are marked.

1280

1281

1282

Figure S2. SVD applied to difference Fourier maps. Difference Fourier maps at the

short delays $t \leq 10$ ps are decomposed into component maps. Each difference map at a

time delay t can be represented by a linear combination of these components, $c_1(t)\boldsymbol{U}_1 +$

$c_2(t)\boldsymbol{U}_2 + \ldots$, where $\boldsymbol{U}_k$ are the time-independent components and $c_k(t)$ are their

corresponding time-dependent coefficients (Methods). (a and b) Two example plots

show circular correlations between c_1 and c_2, c_9 and c_{11}. These circular correlations

indicate two-dimensional oscillations. Each colored trace represents difference maps in

a time series calculated with a common reference. Those time series with a dark

reference are plotted with thick lines. Other series are in thin lines. (c) Singular values

before and after Ren rotation (Ren, 2016, 2019) (Methods). Singular values derived from

SVD indicate the significance of the components. 17 of them stand out.

1294

Ren: Isomerization sampling

Figure S3. Oscillations of SVD components. The coefficients of ten components c_1, c_2; c_4,

c_7; c_{15}, c_{16}; c_9, c_{11}; and c_{12}, c_{13} are found oscillating at frequencies ranging from 60 to 400

cm^{-1}. Each pair of the coefficients oscillate at a common frequency. These frequencies

are 61±2, 150±3, 224±7, 334±8, and 396±3 cm^{-1}, respectively. These coefficients are

plotted against the time delay t (left) and against each other in a pair (right). Each

coefficient is fitted with a sine function around a straight baseline $c_k = a \sin\left(\frac{2\pi t}{T} + \varphi\right) +$

$b + ct$. Both the fitted function and the baseline are plotted. The amplitude a for the

last pair of coefficients c_{12} and c_{13} are replaced with a Gaussian function $a -$

$\exp\left(-\frac{(t-t_0)^2}{\tau^2}\right)$ to implement a node at $t_0 = 509\pm5$ fs (e).

1305

1306

a. U_1 b. U_2 c. U_4 d. U_7 e. U_9

f. U_{11} g. U_{12} h. U_{13} i. U_{15} j. U_{16}

Figure S4. No structural signal in oscillating components. Ten oscillating components are contoured at $\pm 3\sigma$ in green and red, respectively. The main chain and side chains of the protein are rendered with ribbon and sticks, respectively. The retinal and Lys216 are in purple sticks. Several key waters are in blue spheres. Parts of the structure are omitted to reveal more of the interior. Despite that the time-dependent coefficients to these components contain strong oscillatory signals (Figs. S2 and S3), these components themselves display no obvious association with any structural features such as the retinal or secondary structures. They are in stark contrast to the signal distributions of the non-oscillating components (Figs. 2ab, 3c, and S5).

Figure S5. Two orthographical views of component map U_{14}. The main chain and side chains of the protein are rendered with ribbons and sticks, respectively. The retinal and Lys216 are in purple sticks. Several key waters are in blue spheres. Parts of the structure are omitted to reveal more of the interior. The map is contoured at $\pm 3\sigma$ in green and red, respectively. The signals are largely associated with the chromophore and its immediate vicinity.

1328
1329
1330 Figure S6. Two orthographical views of the 2Fo-Fc map of I′ contoured at 3.5σ. Here Fo
1331 is the reconstituted structure factor amplitudes rather than observed amplitudes (Table
1332 S2). Fc is the structure factor amplitudes calculated from the refine structure (Methods).

1333   The same protocol applies to the Fourier synthesis of 2Fo-Fc maps of other

1334   intermediates (Figs. S7, S11, and S12).

1335

1336

Figure S7. Two orthographical views of the 2Fo-Fc map of I contoured at 3σ. Here Fo is the reconstituted structure factor amplitudes rather than observed amplitudes (Table S2). Fc is the structure factor amplitudes calculated from the refine structure (Methods).

1344 Figure S8. Raw difference Fourier map at 457-646 fs. This difference Fourier map is

1345 calculated from the dataset 6g7j at the time point of 457-646 fs by subtracting the dark

1346 dataset 6g7h. The map is contoured at $\pm 3\sigma$ in green and red, respectively. This map is

1347 prior to SVD analysis. Compared with $\boldsymbol{U}_{10}$ (Fig. 3c) and the reconstituted map (Fig. 3d),

1348 it is clear that this is the original source of the widespread signals except that the σ

1349 value of this map is higher than those after SVD. (a) The raw difference map contoured

1350 in the entire molecule shows the association of the signals with the structural elements

1351 at an excellent signal-to-noise ratio. (b) Details of the raw difference map show

1352 displacements of helices. The raw difference map is largely the same as the

1353 reconstituted map (Fig. 3d).

1354

1355

1356

Figure S9. Refined structures at 457-646 fs and 10 ps compared to the ground state. The refined structures of 6g7j and 6g7k and compared with the resting state 6g7h viewed along the trimer three-fold axis from the extracellular side. Atomic displacements in the main chain from bR to 6g7j and 6g7k are color coded and marked by arrows with lengths 20× of the actual displacements. Compared to the structures of I and J species (Fig. 3e), smaller expansion and slight contraction were also captured.

1363

1364

1365

1366 Figure S10. Retinal binding pocket at the distal end. Two orthographical views of the

1367 retinal at its distal end. The closest contacts are 3.7 Å to Ile119 and Pro186.

1368

1369
1370
1371   Figure S11. Two orthographical views of the 2Fo-Fc map of J′ contoured at 4σ. Here Fo
1372   is the reconstituted structure factor amplitudes rather than observed amplitudes (Table
1373   S2). Fc is the structure factor amplitudes calculated from the refine structure (Methods).
1374

1375
1376
1377 Figure S12. Two orthographical views of the 2Fo-Fc map of J contoured at 5σ. Here Fo
1378 is the reconstituted structure factor amplitudes rather than observed amplitudes (Table
1379 S2). Fc is the structure factor amplitudes calculated from the refine structure (Methods).

1380 Photoinduced Isomerization Sampling of Retinal in

1381 Bacteriorhodopsin

1382

1383 Zhong Ren

1384

1385 Department of Chemistry, University of Illinois at Chicago, Chicago, IL 60607, USA

1386 Renz Research, Inc., Westmont, IL 60559, USA

1387

1388 zren@uic.edu

1389 ORCID 0000-0001-7098-3127

1390

1391 Point-by-point response to peer review

1392 I am grateful to the editor and reviewers for their constructive comments and

1393 suggestions. I have clarified a few points and expanded discussion throughout. I have

1394 also added several figures for structural comparison as requested. Here I respond to

1395 the reviewers' comments point by point. These comments are copied below in *italics*

1396 followed by my response. The topics are arranged by relevance to one another.

1397

1398 *Editor: All three reviewers find the work to be interesting and note that, as written, will require*

1399 *a major revision of our understanding of the biophysics of light induced isomerization of retinal*

1400 *in bacteriorhodopsin.*

1401

Reviewer 1: The manuscript reports SDV analysis of time-resolved x-ray data directed at the I to

J intermediates of the bacteriorhodopsin mechanism. The SDV approach that has been developed

over the past decade by the author helps to eliminate noise and systematic error components in

the data sets to generate better models of the intermediates. A key new aspect of the study from

prior directed at bacteriorhodopsin is the inclusion of structural heterogeneity, providing a truer

description of proteins than by modeling based on pure species. The work reveals that excitation-

induced charge separation causes retinol to initially form an S shaped conformation with

isomerization sampled at many carbons before being fixed at 13C; rapid protein expansion

followed by contraction during the retinol motions are uncovered. While bacteriorhodopsin has

been extensively studied as a classical model protein, this work more rigorously elucidates

structural evolution that provides a more complete explanation for the selectivity of

isomerization at the 13C of retinol. The demonstrated functional role of the mechanical

flexibility involving such rapid (fs) protein motion is not well appreciated, and I consider this a

particularly novel contribution to our general understanding of protein biophysics.

Reviewer 3: The paper addresses the mechanism of light induced isomerization of retinal in

microbial bacteriorhodopsin in light of recent time-resolved serial femtosecond X-ray

crystallography (TR-SFX) experiments. The significance of the work is that photoisomerization

of retinal occurs in the case of microbial rhodopsins as well as light reception and signaling in

animal rhodopsins. However, at the atomistic level much remains to be learned about the

regiospecificity and the stereospecificity and how it is controlled by the protein environment.

That knowledge in turn can build a better understanding of light energy conservation, as well as

applications to optogenetics as well as understanding visual signaling. The work is potentially

appropriate for publication; however, further justification and clarification of the major findings

is needed. The paper has some nice insights and commentary which might form the basis for a

contribution after it is revised and reviewed again. The research involves a refinement of the

previously published time-resolved X-ray structures of bacteriorhodopsin in the range from zero-

10 ps reported by Nogly, Neutze, Scherlter et al. in two earlier papers published in Science. As

well there is as a more comprehensive and definitive study by G. Nass Kovacs, I. Schlichting et

al. published in Nature Communications (2019). In those papers the results of time-resolved

serial crystallography (TR-SFX) were interpreted in terms of photointermediates evolving with

time. However, the electron density maps correspond to a mixture of photoexcited species

present in the early photocycle. The current paper is a refinement of the previous time-resolved

data sets involving four intermediate structures which are refined against the predicted structure

factor amplitudes. The method involves singular value decomposition which is clearly and nicely

explained in the supplementary material.

I appreciate the encouraging words of the editor and reviewers as well as their

constructive comments.

Reviewer 3: General Comments: The paper is very well-written with informative and well-

prepared figures. The results of the structural refinement involve deconvolution of several

distinct early photoproducts which improves the quality of the electron density maps. It is

shown that the all-trans retinal undergoes isomerization sampling at the various double bond

positions, which are however constrained by the protein binding cavity leading to conformational

1447 *selection of the 13-cis photoproduct. Because of the isomerization sampling, it is proposed that*

1448 *there is a transient expansion of the retinal binding pocket followed by a contraction or a recoil as*

1449 *the 13-cis photoproduct is formed. At this stage, the work is basically a refinement of what is*

1450 *already known.*

1451

The transient, sub-picosecond expansion of the retinal binding pocket and the later

recoil that results in a contraction of the binding pocket were not known previously.

Such overall response of the protein framework reflects that numerous reaction dead

ends fail in simultaneous attempts to isomerize or rotate along the polyene chain of the

retinal. The raw difference Fourier map at the key time point of 457-646 fs (Nogly et al.,

2018) without the treatment of SVD already shows excellent signal of the pocket

expansion (Fig. S8). The refined structures of 6g7j and 6g7k also show the expansion,

although to a lesser extent, and a slight contraction (Fig. S9). The previous failure to

discern the pocket expansion could be caused by the simplest reason – didn't look. An

alternative explanation could be a suboptimal numerical algorithm or software

implementation in calculation or rendering of difference Fourier maps. I have long

demonstrated that using difference Fourier maps to reveal weak signal of structural

changes is highly nontrivial (Ren et al., 2001). However, these early work of mine

received little attention.

1466

1467 *Reviewer 3: The general reader will want to know whether the work is a refinement of the earlier*

1468 *papers, or whether it is transformative with new biophysical insights. My recommendation is*

1469 *that the paper is potential suitable for publication, but that it needs to more fully address all of*

1470 *the points above and below to justify the significance. Are the structures a refinement of the*

1471 *earlier structures or do they lead to transformative insights?*

1472

1473 Whether this work is transformative or merely an incremental improvement from the

1474 published literature is up to the readers' judgement. There is no need to make any

1475 claim. However, I have made changes and additions throughout to address every point

1476 raised by the reviewers. A major addition is the structural comparison with the

1477 previous structures as requested (see below).

1478

Reviewer 3: However, there is also a proposed mechanism involving a nonspecific Coulombic

attraction of the all-trans polyene whose projected length is less in 13-cis conformer. The author

needs to bring in more persuasively what is known of the excited state photochemistry and

molecular orbitals of bacteriorhodopsin, as extensively investigated in earlier work by Birge, El

Sayed, Hudson, and others in the earlier literature. The proposal that a greater Coulomb

electrostatic force is consistent with the reduced projected length in the 13-cis photoproduct as a

general mechanism ignores other aspects of the excited state photochemistry. The excited state of

a polyene is expected to be more polar consistent with the experimentally measured dipole

moment of Mathies et al.

1488

I appreciate this constructive suggestion, which would no doubt elevate this manuscript

to a new level. A brief mention of the barrierless excited state potential energy surface

is added (Line 70-73). A number of quantum mechanics and molecular dynamics

simulations are discussed and compared to the findings in this study (Altoè et al., 2010;

Hayashi et al., 2003). Questions regarding the correspondence between isomerization

theory and experimental observation may not have concrete answers yet. At least, I

attempt to pose the questions and speculate the correspondence (Line 373-380, 407-420,

433-442, and 476-480).

1497

Reviewer 3: In addition, one needs to consider the greater anti-bonding character of the excited

state molecular orbitals, which is expected to favor an opposite retinal elongation as in the all-

trans configuration. There may be other contributions besides the electrostatic and steric

contributions described by the author. The resultant energies may not be simply attributable to

the light-induced charge separation known from previous experimental work. Probably the

stereoselectivity of the protein pocket and conformational selection also come into play, in

addition to the Coulombic attractive forces in the isomerization sampling.

Reviewer 3: Please clarify whether the comments pertain to the excited singlet state or the

ground state potential energy surface. The Coulombic attraction explanation would not work for

11-ciss to all-trans isomerization in visual rhodopsin. Typically, electronic excited states are

more polar so one would want to know whether the proposal it unique to rhodopsins or more

general. The Coulombic attraction is counteracted by greater anti-bonding character of the

excited state molecular orbitals so please explain further. What is the dominant effect, is it the

greater dipole moment or the greater anti-bonding character?

1512

I have been speculating the correspondence between isomerization theories and my

structural observation (see above). However, this study does not provide any evidence

to support anti-bonding character of the excited state. Quite the contrary, what we can

learn here is how a non-specific force causes a highly selected result. Nevertheless, I do

not believe that I have intentionally excluded any contribution from other factors. I

would be happy to reword if pointed out.

1519

Reviewer 3: The idea about nonspecific electrostatic forces leading to collapsing of the all-trans

retinal by double bond isomerizations along the polyene with conformational selection by the

protein is belied by the 11-cis to all-trans retinal isomerization in visual rhodopsin. That would

need to be clarified.

Reviewer 3: With regard to animal rhodopsins there is ample evidence for expansion of the

protein binding cavity upon 11-cis to all-trans isomerization. However, the retinal elongation is

opposite to that proposed for bacteriorhodopsin and other microbial rhodopsins.

1527

It has been reasoned repeatedly that studies on all-*trans* → 13-*cis* isomerization in

bacteriorhodopsin would shed light into the mechanism of 11-*cis* → all-*trans*

isomerization in visual rhodopsin. The commonality perhaps exists only at a high level

in a rather abstract sense, if any. The reactions from all-*trans* and to all-*trans* not

necessarily share any common trajectory. This is noted in the Concluding remarks

(Line 516-523).

1534

Reviewer 3: Are the refined structures due to multiphoton excitation coming from hot

vibrational levels back to the ground state potential surface? Or do they represent alterative

non-biological pathways besides the biologically relevant process?

Reviewer 3: Very importantly, the role of multiphoton excitation as discussed by Nass Kovacs

and also by R.J.D. Miller et al. Nature Communications (2020) needs to be brought more

explicitly into the picture, to fully justify the novelty and significance of the present

contribution.

Reviewer 3: And importantly, whether the intermediate structures correspond to multiphoton

excitations, or involve just the first excited singlet state to ground state potentials.

Reviewer 3: References - It is essential to mention the paper of R.J.D. Miller et al. Nature

Communications 11, 1240 (2020) and comment on the possibilities of multiphoton excitation as

an explanation for the coexisting refined structural intermediates (!).

1547

These comments and suggestions are important and constructive. The refinement of

the intermediate structures in this study is largely driven by the captured signals in

datasets of Nogly et al. Datasets from Kovacs et al. contribute zero signal. But these

datasets contain significant oscillatory signals. I kept the section on low frequency

oscillations as short as possible without discussion on multiphoton absorption because

it deviates from the main subject of isomerization sampling. The experimental data of

Kovacs and Schlichting et al. contain no biologically relevant signal but only

oscillations. The reason was pointed out by Miller et al. – multiphoton absorption

(Miller et al., 2020). I agree with Reviewer 3 that it would be responsible to discuss in

greater detail on the issue of laser peak power and multiphoton absorption rather than

leaving it for the readers to conclude. The section on low frequency oscillations has

been expanded (Line 157-162 and 176-188).

1560

1561   *Reviewer 1: The main paper also states that the results were based on only the few data sets of*

1562   *Nogly, et al. This point could be clarified, as the data appear to have 17 significant components*

1563   *which would require more than the few data sets. Do the oscillations dominate the sets of*

1564   *Kovacs, et al., such that they contribute little support for U10?*

1565

Yes. The section on low frequency oscillations has been expanded (see above).

Discussion on multiphoton excitation is added (Line 117-119 and 176-188). The datasets

of Kovacs et al. require a lot of singular values, that is, components. Unfortunately,

they do not contribute any useful signal. Their datasets contain near perfect oscillations

from 60 to 400 cm^{-1}. These oscillations must occur on the potential energy surfaces of

multiphoton excitation.

1572

Reviewer 1: It's also surprising that a single protein-based component U10 could sufficiently

describe both intermediates I and J; does this reflect the crudeness of the current description of

the protein response, as one would expect the protein structural changes involved in the

expansion around a cis-retinol and contraction for the trans-retinol would differ somewhat in

their nature due to the distinct chromophore conformations. Expanded presentation of the

specific analysis of bacteriorhopsin in the SI would help to clarify the limitations of the model

generated from the study.

1580

1581   I and J species contain positive and negative U_{10}, respectively. But they also require U_{14}.

1582   Therefore, strictly speaking, a single component is insufficient to describe both

1583   intermediates I and J. The coefficients are listed in Table S2. I have also explicitly stated

1584   these coefficients in the main text so that some confusion may be avoided (Line 196-197,

1585   208, and 322-325).

1586

1587   *Reviewer 3: To justify the significance of the work, I would expect to see a much stronger*

1588   *comparison to the results of the Nass Kovacs and Nogly papers, with more definitive statements*

1589   *about what is learned of the photointermediate structures versus the simpler intermediate*

1590   *analysis. The paper needs to present a more direct comparison to the earlier structures to more*

1591   *directly substantiate the isomerization sampling and its role in the binding pocket expansion.*

1592

This is a very constructive suggestion. I left these direct comparisons out in the hope

that the readers shall conclude by themselves. I added the direct comparison in text

and figures and expanded the discussion (Line 282-284, 328-329, 345-371; Figs. 4bc and

S9). The key difference is that Nogly et al. and Kovacs et al. presented a gradual

rotation of the double bond $C_{13}=C_{14}$ instead of discrete states of *trans* and *cis* concluded

in this study. The gradual rotation is a misinterpretation of the gradual shift of two or

more discrete populations directly caused by the inability to deconvolute mixed signals.

This difference is highly consequential to the mechanistic understanding presented in

the companion paper (Ren, 2021).

1602

Reviewer 1: Maybe a matter of taste, but placement of figures could be reconsidered, as in some

ways it didn't follow the sequence of my mind's digestion of the results. The collection of panels

for Figure 2 is primarily the source of this comment. Figure 2a summarizes the structural

evolution, so perhaps placing it before all data would be appropriate. I would find it more useful

to have Figure 2C focused on the retinol motions placed with information about retinol-based

components U14 and U17 in Figure 1. It also seems that Figure 2b which shows the U10

component would be better associated with the panels in Figure 3.

1610

1611   This is an excellent suggestion. I reorganized the main figures accordingly.

1612

1613   *Reviewer 3: Figures - The figures are high-quality and quite beautiful; however, it is difficult to*

1614   *make cross comparisons due to the small structural differences both in the main text as well as in*

1615 *the supplemental information. It will be more helpful to the reader if an expanded overlay of the*

1616 *refined intermediate structures is provided, together with comparison in particular to the Nass*

1617 *Kovacs structures.*

1618

1619 I added Fig. 4 mainly for structural comparison (Fig. 4bc). Fig. S9 is also added to show

1620 the expansion and contraction captured by Nogly et al. but to a lesser extent. These

1621 changes in the retinal binding pocket must have been overlooked by the original

1622 authors.

1623

1624 *Reviewer 3: Figures - Please provide a figure showing the overlay of the Nogly and Nass Kovacs*

1625 *photointermediate structures so the reader can directly see the extent of the structural differences*

1626 *brought about by the new refinements. This could be combined with overlay of the*

1627 *photointermediate structures, which are lost in Figure (c).*

1628

1629 Overlay of too many structures could be messy. However, the point is well taken. I

1630 have managed to present structural comparison without too many figures and

1631 extensive overlay of structures.

1632

1633 *Reviewer 3: With regard to figure 2(c) it is necessary to more clearly describe what is actually*

1634 *presented, both in the figure caption and the main text.*

1635 *Reviewer 3: Figure 2 - In part (c) please state more clearly what exactly is plotted in the various*

1636 *panels, both in the caption as well as in the main text. The color coding is not enough.*

1637

1638 An entire paragraph is inserted to describe the conformational parameters (Line 216-

1639 238).

1640

1641 *Reviewer 3: Specific Comments: Page 2 bottom - Please clarify the protonation state of the*

1642 *retinylidene Schiff base (PSB or SB) for the general reader, which is important with regards to*

1643 *the proposed non-specific Columbia attractive force.*

1644

1645 Done.

1646

1647 *Reviewer 1: Figure 3(a) caption: "2ab" should be "2b".*

1648

1649   Corrected.

1650

Reviewer 1: The one minor recommendation for improvement I have regards the detail of the

presentation of the analysis results. The description of the SVD methodology applied is very

thorough explained in a general manner in the SI; however, some details of the analysis in this

specific study could be more clearly presented. One example, the main paper states that 24 data

sets and 18 time points up to 10 ps were analyzed and referred to Table SI1, while Table SI1 lists

only 22 data sets under 10 ps.

1657

1658   The errors in Table S1 are corrected.

1659

References

Altoè, P., Cembran, A., Olivucci, M., and Garavelli, M. (2010). Aborted double bicycle-pedal
isomerization with hydrogen bond breaking is the primary event of bacteriorhodopsin proton pumping.
Proc. Natl. Acad. Sci. *107*, 20172–20177. https://doi.org/10.1073/pnas.1007000107.

Hayashi, S., Tajkhorshid, E., and Schulten, K. (2003). Molecular dynamics simulation of
bacteriorhodopsin's photoisomerization using ab initio forces for the excited chromophore. Biophys. J. *85*,
1440–1449. https://doi.org/10.1016/S0006-3495(03)74576-7.

Miller, R.J.D., Paré-Labrosse, O., Sarracini, A., and Besaw, J.E. (2020). Three-dimensional view of ultrafast
dynamics in photoexcited bacteriorhodopsin in the multiphoton regime and biological relevance. Nat.
Commun. *11*, 1240. https://doi.org/10.1038/s41467-020-14971-0.

Nogly, P., Weinert, T., James, D., Carbajo, S., Ozerov, D., Furrer, A., Gashi, D., Borin, V., Skopintsev, P.,
Jaeger, K., et al. (2018). Retinal isomerization in bacteriorhodopsin captured by a femtosecond x-ray laser.
Science *361*, eaat0094. https://doi.org/10.1126/science.aat0094.

Ren, Z. (2021). Directional proton conductance in bacteriorhodopsin is driven by concentration gradient,
not affinity gradient. BioRxiv doi:10.1101/2021.10.04.463074. https://doi.org/10.1101/2021.10.04.463074.

Ren, Z., Perman, B., Srajer, V., Teng, T.-Y., Pradervand, C., Bourgeois, D., Schotte, F., Ursby, T., Kort, R.,
Wulff, M., et al. (2001). A molecular movie at 1.8 Å resolution displays the photocycle of photoactive
yellow protein, a eubacterial blue-light receptor, from nanoseconds to seconds. Biochemistry *40*, 13788–
13801. https://doi.org/10.1021/bi0107142.

1679