Nonsense-mediated decay is not a major mechanism for regulating the uORF-containing

mRNAs in Arabidopsis

Author Affiliations: Hsin-Yen Larry Wu[1] and Polly Yingshan Hsu[1, #]

[1] Department of Biochemistry & Molecular Biology, Michigan State University, East Lansing, MI

48824 USA

[#] Corresponding author: pollyhsu@msu.edu, phone: 517-353-5284

ORCIDs:

0000-0001-9407-338X (H.L.W.)

0000-0001-7071-5798 (P.Y.H.)

Keywords: translation, upstream open reading frames, nonsense-mediated decay, mRNA half-

life, ribosome profiling, Arabidopsis

Figures 1-6

Supplementary Figures 1-10

Supplementary Tables 1-4

ABSTRACT

Background:

Upstream ORFs (uORFs) are widespread cis-regulatory elements in the 5' untranslated regions

of eukaryotic genes. In addition to repressing main ORF translation, uORF translation in animals

also reduces mRNA stability through nonsense-mediated decay (NMD). In contrast, the roles of

uORFs in plants are less understood. One limitation of most prior studies is the inclusion of both

translated and untranslated uORFs for analysis.

Results:

Here, we identified actively translated uORFs (TuORFs) in Arabidopsis through ribosome

profiling and examined their roles in gene expression by integrating multiple genome-wide

datasets. Similar to animal systems, Arabidopsis TuORFs are associated with 38%, 14%, and

43% reductions in translation efficiency, mRNA half-lives, and protein levels, respectively.

Surprisingly, we found TuORF-containing mRNAs in plants have 51% higher transcript levels,

which is opposite to the observations in animals. The higher transcript levels of TuORF-

containing mRNAs are persistent in diverse tissues and developmental stages across different

plants. We present multiple lines of evidence supporting the hypothesis that uORF translation

generally does not trigger NMD in Arabidopsis. First, TuORF-containing mRNAs are not

increased in various NMD mutants. Second, TuORF-containing mRNAs and known NMD

targets have distinct expression patterns, and they are translationally repressed via different

mechanisms. Finally, TuORF-containing mRNAs and NMD targets are degraded through

separate pathways.

Conclusions:

In summary, our work reveals that Arabidopsis TuORFs reduce mRNA stability and translation

through mechanisms different from NMD. Our results highlight a fundamental difference in gene

regulation mediated by TuORFs in plants and animals.

INTRODUCTION

Upstream open reading frames (uORFs) are cis-acting translational repressors that regulate many eukaryotic protein-coding transcripts. Mutations or modifications of uORFs are associated with human diseases and plant traits [1–4]. Considering AUG as a translation start codon, 30%–70% of genes contain at least one potential uORF in moss, Arabidopsis, fly, mouse, and human [5–7]. Genes with predicted uORFs are enriched for signal transduction proteins, transcription factors, and membrane proteins in animals [7] and for transcription factors and protein kinases in plants [5], underlining the potential importance of uORFs in modulating the expression of vital cellular regulators.

Many studies in eukaryotes have reported that uORF translation reduces main ORF (mORF) translation [6, 8–11], and it is estimated to reduce protein production by 30%–80% [1, 8]. According to the scanning model of mRNA translation, the 43S pre-initiation complex, which contains the small ribosomal subunit, several translation initiation factors, and initiator tRNA, binds the 5' cap and scans the 5' untranslated region (UTR) to search for an optimal initiation sequence [12]. When a uORF is translated, the ribosome may stall during elongation or at termination, or it may dissociate from the mRNA after termination, thus reducing the translation of the downstream mORF [13]. The mORFs downstream of uORFs may get translated through leaky scanning or reinitiation [14].

uORF translation is also expected to trigger mRNA degradation, presumably through nonsense-mediated decay (NMD). NMD is an evolutionarily conserved regulatory mechanism to eliminate aberrant mRNAs, such as those that contain premature termination codons and unusually long 3' UTRs [15, 16]. Two pathways have been established in mammals: the exon junction complex (EJC)-dependent and long 3' UTR pathways. Normally, EJC and two NMD factors (UP-FRAMESHIFT SUPPRESSOR 2 and 3; UPF2 and UPF3) deposited during splicing of mRNAs will be evicted by the ribosome in the first round of translation. In the presence of a premature termination codon, EJC-UPF2-UPF3 complexes downstream of the termination

codon remain attached to the mRNA. This allows UPF1, which is part of the surveillance

complex recruited at translation termination, to interact with UPF2 and UPF3 in proximity. The

interaction of UPF1, UPF2, and UPF3 activates the SUPPRESSOR WITH MORPHOLOGICAL

EFFECT ON GENITALIA1 (SMG1) kinase, which phosphorylates UPF1. This establishes the

first phase of NMD, substrate recognition, and activates downstream signaling and eventually

the second phase of NMD, substrate degradation, which involves SMG5, SMG6, and SMG7.

Phosphorylated UPF1 also inhibits translation initiation [17]. Alternatively, unusually long 3'

UTRs, which result in poor translation termination, are believed to promote NMD through

phosphorylated UPF1 independent of EJC [18].

The plant NMD pathway is best characterized in *Arabidopsis thaliana*. Arabidopsis

possesses homologous NMD factors involved in the first phase, including UPF1, UPF2, and

UPF3, and in the second phase, SMG7 [19]. However, Arabidopsis is missing the homologs of

animal SMG1, SMG5, and SMG6 [20]. The absence of key NMD components in plants

suggests significant variations in the NMD mechanisms between plants and animals. It remains

an open question how the differences in NMD components affect substrate recognition and

degradation in plants.

In Arabidopsis, putative NMD targets have been identified based on transcripts that

overly accumulate in NMD-defective mutants [21–24]. However, there is a concern that the

upregulation of some transcripts in NMD mutants might be caused by secondary effects of NMD

[25]. For example, NMD mutants affect numerous aspects of plant physiology, and some alleles

cause severe growth retardation and lethality due to constitutive immune responses [26–32].

Moreover, UPF1 has been reported to have multiple functions in plants, such as alternative

splicing regulation and translational control, in addition to NMD [25]. To identify *bona fide* NMD

targets, *upf1* and *smg7*, which are defective in the first and second phases of NMD,

respectively, were introduced into *pad4*, in which the immune response is inactivated [25, 28].

The 333 transcripts upregulated in both *upf1 pad4* and *smg7 pad4* were defined as high-

confidence NMD targets. It was shown these NMD targets are translationally regulated by UPF1

and that their turnover requires decapping and 5' to 3' decay [25].

As uORF translation could result in a transcript with an unusually long 3' UTR, uORF-

containing mRNAs are predicted to be targeted by NMD [16, 33]. Consistent with this

expectation, genome-wide studies in zebrafish, mouse, and human have found that transcripts

with translated uORFs have lower mRNA levels globally [6, 8]. Moreover, transcripts containing

translated uORFs are upregulated in knockdown of NMD factors in mouse embryonic stem cells

and human cell lines [34, 35], supporting the idea that these transcripts are NMD targets.

In plants, whether uORF translation triggers NMD is still under debate. Two genome-

wide studies suggest that uORF-containing mRNAs are targets of NMD, as transcripts

possessing predicted uORFs are enriched in the genes upregulated in Arabidopsis *upf1-1* and

upf3-1 mutants [21, 23]. In contrast, three studies suggested that NMD only controls a small

fraction of uORF-containing mRNAs in plants, such as those containing uORFs with > 35

codons (~2% of uORF-containing genes), or those responsive to certain environmental stimuli,

such as polyamine or blue light [36–38]. Moreover, evidence from several gene-specific or

genome-wide studies suggests that uORFs do not trigger NMD. First, the mRNA levels of

several uORF-containing transcripts, including *NIP5;1* and *AtMHX*, are not increased in NMD

mutants [39, 40]. Second, a genome-wide study reported that genes with increasing numbers of

uORFs did not show lower transcript abundances in Arabidopsis and tomato [41]. Third, genes

with predicted uORFs are not over-represented in the consensus NMD target list identified from

three NMD mutants, including *upf1-5*, *upf3-1*, and *smg7-1* [27]. It was also reported that 47

mRNAs that contain Conserved Peptide uORFs (CPuORFs) behave differently from NMD

targets prior to their degradation mediated by XRN4 [42]. It is unknown whether some uORF-

containing mRNAs escape NMD or whether uORF-containing mRNAs are generally not targets

of NMD in plants. It has been concluded that the mRNA features studied so far are insufficient

to infer NMD targetability in plants [43]. Overall, it remains unclear how translated uORFs affect

mRNA stability and what the role of NMD in this process is in plants.

One limitation of the previous studies is that predicted uORFs, which include both

translated and untranslated uORFs, were commonly included in the analyses. Although many

uORFs have been predicted based on nucleotide sequences [7], determining which uORFs are

actively translated still requires experimental evidence. Recently, deep sequencing of ribosome

footprints – known as ribosome profiling or Ribo-seq – has been exploited to study genome-

wide mRNA translation [44, 45]. Several Ribo-seq studies in Arabidopsis have deployed

different strategies to identify translated uORFs (TuORFs) [9, 46–50]. Among the parameters

used, 3-nucleotide (3-nt) periodicity, which corresponds to translating ribosomes deciphering 3

nts per codon, is considered the most stringent criterion to identify high-confidence translation

events [44, 51–53]. Our previous work and a recent study using RiboTaper [51], which

computes the statistical significance of 3-nt periodicity, uncovered 187–1378 AUG-initiated

TuORFs in Arabidopsis [47, 50].

Here we further improved the Ribo-seq methodology and identified 2093 AUG-initiated

TuORFs in Arabidopsis seedlings. We systematically examined the assumed roles of TuORFs

in mORF translation and mRNA decay by integrating multiple genome-wide datasets. As

predicted, we found that TuORF mRNAs are associated with lower translation efficiency, shorter

mRNA half-lives, and lower protein abundance. Surprisingly, we found that, in contrast to the

animal paradigm, plant TuORF mRNAs have higher steady-state mRNA levels, and they are not

increased in the NMD mutants. Examinations of numerous parameters revealed that uORFs

and NMD regulate gene expression through separate mechanisms. Taken together, our results

reveal unexpected features and regulation of uORFs in Arabidopsis and suggest that plants and

animals have evolved different uORF-dependent mechanisms for gene regulation.

RESULTS

Translated uORFs identified via an improved Ribo-seq method

Previously we optimized Ribo-seq to reach high 3-nt periodicity, and we exploited this

unique feature of mRNA translation to identify translated ORFs in Arabidopsis [47]. Although the

data revealed many unannotated translation events, the number of TuORFs uncovered was

low, presumably due to insufficient Ribo-seq coverage within uORFs. We proposed that

increasing Ribo-seq coverage could improve the efficiency of translated ORF identification,

especially for relatively short ORFs, such as uORFs. Here we optimized the coverage of Ribo-

seq in 7-day-old Arabidopsis seedlings by minimizing the potential loss of ribosome footprints

during the purification process (see METHODS for details).

In addition to high correlations among biological replicates (**Figure S1A**), our new Ribo-

seq data displayed characteristics expected for high-quality data, including strong 3-nt

periodicity (92% in-frame reads), high enrichment for coding sequences (CDSs), and

characteristic ribosome footprint lengths (**Figure 1A-C**). The improved coverage is evident when

examining the number of unique ribosome footprints detected globally. At a sequencing depth of

20 million reads from randomly selected reads, the number of unique ribosome footprints in our

new data is 4.01-fold higher than that in our previous dataset (**Figure 1D**). The coverage

improvement is also evident in individual transcript profiles within both uORFs and mORFs. For

example, a TuORF in *REPRESSOR OF GA1* (*RGA1*) was identified in our current data but

missed in our previous study despite the mRNA levels being similar (**Figure S1B**).

The high-coverage data increased the number of TuORFs uncovered. In total, 2093

TuORFs were identified in 1768 genes based on significant 3-nt periodicity using RiboTaper and

Araport11 annotation (**Figure 1E-G** and **Table S1A**). This result is in strong contrast to the

relatively few TuORFs identified in Arabidopsis seedling shoots (93 TuORFs) and roots (138

TuORFs) from our previous data obtained using plants grown under similar conditions and using

the same statistical analysis and annotations (**Figure 1E** and **Table S1B-C**). The new data also

revealed translation from a higher number of Conserved Peptide uORFs (CPuORFs) and

annotated coding sequences (mORFs of protein-coding genes) (**Figure 1E** and **Table S1D-F**).

We manually plotted all TuORF genes to confirm strong 3-nt periodicity present in the expected

uORF regions using RiboPlotR [54] (examples shown in **Figure 1F-G, Figure S2**). Taken

together, these results show that our enhanced-coverage Ribo-seq data dramatically improved

the identification of TuORFs.

Previous analyses of predicted uORFs found that transcription factors and protein

kinases are overrepresented among uORF-containing genes [5, 55]. Consistent with these

analyses, functions related to transcription factors and protein kinases/phosphatases were

identified as being enriched in our Gene Ontology (GO) analysis of TuORF-containing genes

(**Table S2**). We found that many TuORF-containing genes encode extensively studied

regulators of plant growth and environmental responses. For example, this list included well-

known transcription factors, such as those involved in hormone signaling (RGA1 and AUXIN

RESPONSE FACTOR2/3/8), root development (SCARECROW), the circadian clock (TIMING

OF CAB EXPRESSION1), and light signaling (PHYTOCHROME INTERACTING FACTOR3/4/7)

[56–63]. The list also included well-characterized protein kinases, such as Phytochromes B and

E, ribosomal-protein S6 kinases (S6K1/2), and SNF1-related protein kinase 2 (SNRK2.1/2.5)

(**Figure S1B, S2A-N**) [64–69]. These results support the notion that TuORFs control important

cellular regulators in plants.

TuORF-containing genes have higher mRNA abundance but are associated with lower

translation efficiency, shorter mRNA half-lives, and lower protein abundance

To understand how uORF translation affects gene expression, we compared mRNA

abundance and translation efficiency of TuORF-containing genes (hereafter, TuORF genes),

genes containing sequence-predicted uORFs but undetected by RiboTaper (hereafter, UuORF

genes), and genes containing an annotated 5' UTR but no potential uORFs (hereafter, no-uORF

genes). Note that some UuORFs may be translated but missed in our analysis because their 3-

nt periodicity is insufficient to pass the statistical test: these include extremely short uORFs,

those translated at a low frequency, and those overlapping with other uORFs, thus disrupting

the 3-nt periodicity. To study the global effects of uORFs on mRNA stability and protein

abundance, we also compared genes with TuORFs, UuORFs, or no-uORFs, using available

mRNA half-life data determined by metabolic labeling in Arabidopsis seedlings and two

quantitative proteomics datasets from shoots and roots of Arabidopsis seedlings [70, 71]. The

growth conditions and source of the data are summarized in **Table S3**. We presented the data

in box plots to show the statistical results and in cumulative plots to show the trends of different

gene groups (**Figure 2A-D, Figure S3, S4**).

Studies in vertebrates have shown that uORF translation is associated with lower

steady-state mRNA abundance, consistent with the expectation that TuORF mRNAs are

regulated by NMD [6, 8]. Surprisingly, we found that TuORF genes in Arabidopsis have higher

steady-state mRNA levels than those of no-uORF genes (**Figure 2A, S3A**), despite having

shorter mRNA half-lives as expected (**Figure 2C, S3C**). The median mRNA level of TuORF

genes is 51% higher than that of no-uORF genes. The higher steady-state mRNA levels and

shorter mRNA half-lives together suggest that TuORF genes are highly transcribed. To examine

whether the high transcript levels of TuORF genes are conserved in other plant species, we

reanalyzed our previous tomato Ribo-seq data from seedling roots [72] and found the TuORF

genes in tomato also have higher steady-state mRNA levels relative to no-uORF genes (**Figure**

2E). Given that Arabidopsis and tomato diverged more than 100 million years ago [73], our

results suggest that TuORF-associated higher mRNA abundance could be prevalent in plants.

To investigate whether our observation is specific to seedlings or common to different

developmental stages/tissues/genetic backgrounds in Arabidopsis and tomato, we examined

the mRNA levels of the uORF-containing genes in 128 independent RNA-seq studies collected

by the EMBO-EBI expression atlas [74]. Consistent with our Arabidopsis seedlings and Tomato

seedling root data, we found the TuORF genes are expressed at higher levels in diverse

tissues, developmental stages, and ecotypes in Arabidopsis and tomato (**Figure 2G and S5A-**

C), supporting the hypothesis that TuORF genes are preferentially expressed at high levels in

these plants under a wide range of sampling conditions.

241   CPuORFs are uORFs that encode evolutionarily conserved peptides, and they were

identified independently from our approach by evolutionary conservation [75–79]. Many of the

CPuORFs were also identified as TuORFs in our study (**Figure 1E, 1G**). We found that the

median mRNA abundance of CPuORF genes is 2.2-fold higher than that of no-uORF genes

(**Figure S6**), consistent with our observations that genes with TuORFs are expressed at higher

levels. Our observation of the high mRNA abundance of plant TuORF genes reveals that plants

and animals implement TuORFs differently in gene regulation.

248   We cannot rule out the possibility that TuORFs are more likely to be identified in highly

expressed transcripts, as TuORF translation levels are positively correlated with transcript

abundance (r = 0.38, **Figure S7A**). However, high mRNA abundance alone is insufficient to

determine whether a potential uORF is translated, as many UuORF genes are highly expressed

as well and the distributions of TuORF and UuORF mRNA levels largely overlap (**Figure S7B**).

The fact that the TuORFs identified are functionally associated with lower translation efficiency,

shorter mRNA half-lives, and less protein accumulation than UuORF genes (**Figure 2B-D, F,**

S4) support the idea that TuORFs are selectively translated to repress the translation and

mRNA stability.

257   Interestingly, even though globally TuORF genes in Arabidopsis start with 51% more

mRNAs than no-uORF genes, they produce 41%–43% less protein, on average (**Figure 2A, 2D,**

S3, S4), likely due to the combined effects of a 38% reduction in translation efficiency and a

14% reduction in mRNA half-lives (**Figure 2B, 2C**). These observations suggest that the global

repression of TuORF gene protein synthesis is determined by multilayer post-transcriptional

regulation. Notably, for the four parameters examined here, UuORF genes consistently have

levels between those of TuORF genes and no-uORF genes: they show a 4% increase in mRNA

levels and 11%, 6%, and 21% reductions in translation efficiency, mRNA half-lives, and protein

abundance, respectively (**Figure 2A-D, S3A-D**). These results indicate that UuORFs have

smaller effects on gene expression than TuORFs and highlight the importance of distinguishing

TuORFs and UuORFs in genome-wide analysis.

TuORF genes are larger and likely evolutionarily older than other genes

We next investigated whether there are general structural differences among TuORF,

UuORF, and no-uORF mRNAs. Considering the directionality of ribosome scanning, we

hypothesized TuORFs might be associated with longer 5' UTRs. Indeed, we found that the

median 5' UTR length of TuORF mRNAs is 2.5-fold of that of no-uORF mRNAs (**Figure 3A**),

which is consistent with a previous report that uORFs in longer 5' UTRs are more likely to be

translated [9]. However, unexpectedly, we found that TuORF mRNAs also have longer CDSs

and 3'UTRs, as well as longer transcript lengths, than UuORF mRNAs (**Figure 3B-D**). No-uORF

mRNAs are the shortest in all categories examined (**Figure 3A-D**). These results support the

hypothesis that TuORFs are more likely to be found in longer 5' UTRs and reveal that TuORF

mRNAs tend to encode larger proteins (**Figure 3B**). The enrichment of TuORF genes for

transcription factors, protein kinases, and related functions in GO term analysis (**Table S2**) also

implies TuORF genes tend to encode proteins with conserved domains. It is known that

evolutionarily older genes are generally longer, expressed at higher levels, and have protein

sequences that are under stronger purifying selection [80, 81]. Our observations that TuORF

genes possess many features of evolutionarily older genes imply that older genes are more

likely to be regulated through mRNA translation and degradation mediated by TuORFs.

TuORF mRNAs are not upregulated in the NMD mutants

The established paradigm in vertebrates is that uORF translation triggers NMD and thus

results in low steady-state mRNA levels [6, 8, 34, 35]. Despite having higher steady-state

mRNA levels (**Figure 2A**), TuORF genes in Arabidopsis also have shorter mRNA half-lives

(**Figure 2C**). Therefore, we investigated whether TuORF-containing transcripts in Arabidopsis

are subject to NMD.

If TuORFs trigger NMD, one would expect TuORF mRNAs to be among the known NMD

targets and their mRNA levels to be increased in NMD-deficient mutants. We first examined

whether TuORF mRNAs were reported in the high-confidence NMD target list [25]. Among the

333 NMD targets that were upregulated in both *upf1 pad4* and *smg7 pad4* adult leaves [25], 248

genes with an annotated 5' UTR were expressed in our data (**Table S4A**). However, only 49 of

1748 TuORF genes were identified in the high-confidence NMD target list (**Table S4B**). The 49

genes represent 2.8% of TuORF-containing genes and 20% of the NMD targets. These results

suggest that TuORFs generally do not trigger NMD; rather, NMD targets are more likely to be

regulated by TuORFs than other genes (Chi-square test with Yates' correction: p-value =

7.555e-10). One caveat is that the poor overlap between the two gene lists could be due to

differences in growth conditions, developmental stages, and sample handling in different labs

(**Table S3**).

To further investigate if TuORF mRNAs are NMD targets, we examined whether they are

upregulated in NMD-deficient mutants by analyzing four published RNA-seq datasets for NMD

mutants [23, 25, 30, 82]. Although these datasets were collected from different NMD mutants

and at various developmental stages (**Table S3**), the high-confidence NMD targets are

consistently and significantly upregulated in all the NMD mutants examined compared with the

control plants (i.e., 3.05-, 1.54-, 4.98-, and 2.65-fold for NMD targets without TuORFs; 3.04-,

2.23-, 5.04-, and 2.66-fold for NMD targets with TuORFs)(**Figure 4A-D**). In contrast, TuORF

mRNA levels show no or little change in the NMD mutants (i.e., 1.01-, 0.98-, 1.10- and 1.10-

fold). Thus, only the NMD targets, but not TuORF mRNAs, are dramatically increased in the

NMD mutants, supporting the idea that uORF translation in Arabidopsis generally does not

trigger NMD.

Given different uORFs are translated at various levels (See **Figure 1F-G, S1B, S2A-N**

for examples), it is possible only the highly repressive TuORFs efficiently trigger NMD. To test

this hypothesis, we split the TuORF genes into two groups based on their uORF relative

translation efficiency (RTE) (i.e., TE of an uORF is normalized to TE of its mORF). However,

even the higher TuORF RTE group shows no or little effects on the mRNA accumulation in the

NMD mutants (i.e., 1.13-, 0.99-, 1.15- and 1.11-fold)(**Figure 4E-H**). We also tested whether

having more TuORFs in a transcript efficiently triggers NMD. We split the TuORF genes into two

groups based on the number of TuORFs detected. We found that transcripts with 2~5 TuORFs

do not lead to significant mRNA increase in the NMD mutants, either (i.e., 0.93-, 0.98-, 0.99-

and 1.09-fold)(**Figure 4I-L**). Together, these results further support the idea that uORF

translation in Arabidopsis generally does not induce NMD.

Lengths of the TuORF and 3' UTR alone are insufficient to predict NMD targetability

A previous study suggested that uORFs that encode a peptide longer than 35 amino

acids are likely to trigger NMD in Arabidopsis [36]. To test whether TuORF length is associated

with NMD targetability, we analyzed TuORF lengths in the NMD targets and non-NMD targets.

We found that the NMD target TuORF genes tend to have longer uORF peptides than non-NMD

target TuORF genes (median length 28 amino acids vs. 20.5 amino acids, p = 9.874e-07,

Wilcoxon signed-rank test). However, the distributions of the uORF peptide lengths in NMD

targets and non-NMD targets largely overlap (**Figure S8**), indicating that TuORF length alone is

insufficient to predict whether a TuORF triggers NMD of a transcript.

We also examined whether the high-confidence NMD targets have longer 3' UTRs, a

feature that potentially triggers NMD. Interestingly, the NMD targets do not have significantly

longer 3' UTRs, regardless of whether they contain TuORFs (**Figure S9**). Taken together, our

results are consistent with a recent suggestion that no definite RNA features studied to date are

sufficient to predict NMD targets in plants [43].

TuORF mRNAs and NMD targets have distinct expression patterns

To further characterize the roles of NMD and TuORFs in gene expression regulation, we

compared the expression patterns of the NMD targets and TuORF mRNAs (**Figure 5**).

Consistent with the expectation that NMD targets are being actively degraded, we found that

NMD targets have lower mRNA abundance than other genes; this is in strong contrast to

TuORF mRNAs, which have higher mRNA abundance than other genes (**Figure 5A, 5E**). While

both NMD targets and TuORF genes are associated with lower translation efficiency, shorter

mRNA half-lives, and lower protein abundance than other genes (**Figure 5B-D, 5F-H**), the NMD

targets are associated with even lower protein levels than TuORF genes (**Figure 5D, 5H**).

If TuORFs effectively trigger NMD, one would expect TuORF-containing genes to have

translation efficiency similar to TuORF-containing NMD targets. While both NMD targets and

TuORF mRNAs have lower translation efficiency (**Figure 5B, 5F**), TuORF-containing NMD

targets have significantly lower translation efficiency than either NMD targets without TuORFs or

TuORF-containing non-NMD target genes (**Figure 5B, 5F**). This additive repression of

translation efficiency in TuORF-containing NMD targets suggests that NMD and TuORFs

suppress translation through separate mechanisms and supports that TuORFs do not effectively

trigger NMD. Our observation is consistent with the expectation that TuORFs prevent the

scanning pre-initiation complex from reaching the mORF while NMD suppresses translation

through UPF1 and/or other translational repressors [17, 25, 83].

Together, the differences in the expression patterns between TuORF mRNAs and NMD

targets further support the idea that TuORFs are not an mRNA feature that efficiently triggers

NMD in Arabidopsis.

TuORF mRNAs and NMD targets are degraded through different mRNA decay

mechanisms

mRNA decay can occur in the 5' to 3' or 3' to 5' direction, which is carried out by different

machinery [84]. We next investigated the mechanisms underlying the decay of TuORF mRNAs

and NMD targets. A genome-wide survey of mRNA decay rates in wild type and mutants

defective in either 5' to 3' decay or 3' to 5' decay, or both, has been conducted in Arabidopsis

[85]. In that survey, the *vcs-7* mutant, a null mutant of *VARICOSE*, which encodes a scaffold

protein for the mRNA decapping complex, was used to examine the 5' to 3' mRNA decay; the

sov mutant, defective in *SUPPRESSOR OF VARICOSE*, which encodes a ribonuclease II, was

used to examine 3' to 5' decay; and the *vcs/sov* double mutant was used to examine decay

affected in both directions [86–89]. We exploited this published dataset [85] to study the genetic

requirements for the mRNA decay of TuORF mRNAs, NMD targets, TuORF-containing NMD

targets, and other mRNAs.

We first compared the decay rates of each gene group across the four genetic

backgrounds (**Figure 6**). Similar to mRNAs that are non-NMD targets and lack TuORFs (a.k.a.

other mRNAs, **Figure 6D, 6H**), TuORF mRNAs have dramatically reduced decay rates in the

vcs mutant compared to the wild type (**Figure 6A, 6E**), indicating that decapping and 5' to 3'

decay are critical for degrading these transcripts. In the *vcs/sov* double mutant, TuORF mRNAs

have slightly but significantly lower decay rates than in the *vcs* mutant (**Figure 6A and 6E**),

suggesting *SOV* also plays a minor role in degrading TuORF mRNAs.

In contrast, the mRNA decay rates of NMD targets without TuORFs are only significantly

lower in the *vcs/sov* double mutant, suggesting that NMD targets are efficiently degraded by

both the 5' to 3' and 3' to 5' pathways (**Figure 6B, 6F**). In other words, in the absence of either

SOV or *VCS*, the remaining pathway could completely take over the decay of NMD targets

(**Figure 6B, 6F**). Although there are fewer NMD targets containing TuORFs, which resulted in

small sample size and statistical insignificance (**Figure 6C, 6G**), the decay rates in this group of

genes in the four genetic backgrounds show trends similar to those of NMD targets without

TuORFs (**Figure 6B, 6F**).

Next, we compared the decay rates of the four gene groups within each genetic

background (**Figure S10**). In all four genetic backgrounds, NMD targets have higher decay

rates than TuORF mRNAs and other genes (**Figure S10A-H**). This pattern is still persistent in

the *vcs/sov* double mutant **(Figure S10D, S10H)**, suggesting additional decay pathway(s) is

involved in degrading the NMD targets in the absence of both VCS and SOV.

Taken together, our results suggest that TuORF mRNAs and NMD targets are

preferentially degraded through different decay pathways. NMD targets consistently have higher

decay rates than TuORF mRNAs and appear to be regulated by additional pathway(s) other

than *VCS* and *SOV* (see the model in **Figure 6I**).

DISCUSSION

In this study, we identified Arabidopsis TuORFs using improved Ribo-seq data and

systematically examined their presumed roles in gene expression. While our results support the

expected roles of uORFs in repressing mORF translation and mRNA stability, our study reveals

distinct features and regulation of uORFs in plants compared with animals.

Our observations that plant TuORF genes have higher expression levels (**Figure 2**),

encode larger proteins (**Figure 3**), and are enriched for evolutionarily conserved domains

(**Table S2**) raised an interesting question as to whether older genes are more likely to be

controlled at the protein levels through mRNA degradation and translation via uORFs. Higher

mRNA abundance may allow TuORF mRNAs to be highly translated under certain conditions

that require a high level of specific proteins. For example, it has been suggested that

derepression of downstream translation is a general mechanism of uORF-mediated expression

under stress [90]. Thus, TuORFs could enable fast and dynamic regulation of mORF translation.

Future work monitoring the translation of TuORFs and their mORFs under different conditions

will help decipher the regulation and physiological roles of specific TuORFs.

If TuORFs trigger NMD in Arabidopsis, we would expect that TuORF mRNAs behave

similarly to NMD targets. In this report, we presented six observations suggesting that

Arabidopsis TuORF mRNAs and NMD targets are two distinct groups of transcripts: (1) NMD

targets, but not TuORF mRNAs, are upregulated in various NMD mutants (**Figure 4**). (2)

TuORF mRNAs have significantly higher transcript levels, while the NMD targets have lower

transcript levels than other genes (**Figure 5A**). (3) NMD and TuORFs additively repress

translation, suggesting they repress translation through separate mechanisms (**Figure 5B, 5F**).

(4) While NMD targets are efficiently degraded through both 5' to 3' and 3' to 5' decay, TuORF

mRNAs, like other genes, mainly rely on 5' to 3' decay (**Figure 6**). (5) The decay rates of NMD

targets are consistently higher than those of TuORF mRNAs (**Figure S10**). (6) NMD targets also

appear to be degraded by additional pathway(s) in the absence of both *VCS* and *SOV* (**Figure**

S10H). These differences between TuORF mRNAs and NMD targets suggest that NMD is not a

major mechanism regulating TuORF mRNAs.

The fact that Arabidopsis is missing key NMD factors found in animals, including SMG1,

SMG5, and SMG6, suggests significant variations in the NMD mechanisms between plants and

animals. In particular, SMG1 is the kinase responsible for UPF1 phosphorylation required for

substrate recognition in animal NMD [15, 16]. The lack of SMG1 and the use of an alternative

kinase for Arabidopsis UPF1 phosphorylation [91] may lead to a difference in substrate

recognition: while TuORF mRNAs are recognized as NMD targets in animals, plant TuORF

mRNAs are not.

Despite the difference in NMD components in plants and animals, our findings that

Arabidopsis NMD targets are degraded through both 5' to 3' decay and 3' to 5' decay is

consistent with the two-directional decay of NMD targets in animals. Note that plant SOV is the

homolog of DIS3L2, the exonuclease responsible for 3' to 5' decay of NMD targets in animals

[92, 93]. Two-directional decay could be advantageous to quickly remove aberrant mRNAs from

the cytoplasm to prevent further translation and facilitate the recycling of NMD machinery.

Genes controlled by both NMD and TuORFs might encode critical regulators, and their

protein levels require precise regulation. These genes include key enzymes in gibberellin

hormone and polyamine biosynthesis and many transcription factors (**Figure 1F-G, and Table**

S4B). For example, one of the transcription factors, *SUPPRESSOR OF ACAULIS 51* (*SAC51*),

was discovered in a genetic screen for suppressors of the dwarf phenotype of the *acaulis5*

mutant [94]. *SAC51* contains a CPuORF that is controlled by the thermospermine [95]. When

the CPuORF was mutated, it increased the transcript level of *SAC51* [94]. *SAC51* encodes a

bHLH transcription factor that has been proposed to activate the genes required for stem

elongation.

It is well established that there is an important interplay between translation efficiency

and mRNA stability. Two mechanisms have been proposed to explain this phenomenon: the

"stalled ribosome-triggered decay model," which predicts that ribosome stalling or slowing

promotes mRNA degradation, and the "translation factor-protected model," which predicts that

active translation promotes mRNA stability because the translation initiation factors protect the

mRNA from the decapping complex [96–101]. Consistent with the first model, several CPuORF-

containing mRNAs in Arabidopsis were reported to have strong ribosome stalling patterns, and

the endonucleolytic cleavages upstream of the stalled ribosomes were observed [102, 103].

While most of the TuORFs we identified did not display clear stalling patterns (e.g., **Figure 1F-**

G, S1B, S2A-N), it remains possible that ribosomes have lower translation speeds, on average,

during uORF translation. Because ribosomes will quickly approach termination after initiation

due to the short length of uORFs, and because both initiation and termination are major time-

consuming steps in translation [104, 105], the average translation speed per codon for uORFs is

likely to be slower than that for longer ORFs. Alternatively, if cells mainly monitor the translation

efficiency of the mORF, then the poorer mORF translation caused by uORFs could explain the

faster degradation of TuORF mRNAs predicted by this model. In contrast, since ribosomes

actively engage with TuORFs, the "translation factor-protected" model does not fit TuORF

mRNAs, especially because their degradation relies on decapping and 5' to 3' decay in plants.

Further investigation into TuORF-mediated mRNA degradation will help elucidate the role of

uORFs in regulating mRNA stability.

CONCLUSION

Consistent with the observations in animals, we found that TuORFs in Arabidopsis reduce

mRNA stability and translation. However, unlike in animals, uORF translation generally does not

activate NMD in Arabidopsis. Our results reveal major differences in TuORF-mediated gene

expression between plants and animals.

MATERIALS AND METHODS

Plant growth conditions and lysate preparation

Arabidopsis Col-0 seeds were surface sterilized with 70% ethanol for 5 min, followed by 33%

bleach and 0.03% Tween 20 for 10 min, then rinsed with sterile water 5 times. The seeds were

imbibed at 4°C in the dark for 2 days, then grown hydroponically in sterile liquid media (2.15 g/L

Murashige and Skoog salt, 1% sucrose, 0.5 g/L MES, pH 5.7) while shaking at 85 rpm under a

16-h light (75–80 µmol m^{-2}·s^{-1} from cool white fluorescent bulbs) and 8-h dark cycle at 22°C for

7 days. At Zeitgeber time 4 (4 h after lights on), DMSO corresponding to 0.1% of the media

volume was added to the media (these were mock samples of our large-scale experiment). After

20 and 60 min, three biological replicates (~300 plants per sample) were harvested at each time

point and immediately flash frozen in liquid nitrogen.

Plant lysates were prepared as previously described [47]. Briefly, per 0.1 g of ground tissue

powder was resuspended in 400 µL of lysis buffer (100 mM Tris-HCl [pH 8], 40 mM KCl, 20 mM

MgCl$_2$, 2% [v/v] polyoxyethylene [10] tridecyl ether [Sigma, P2393], 1% [w/v] sodium

deoxycholate [Sigma, D6750], 1 mM dithiothreitol, 100 µg/mL cycloheximide [Sigma, C4859],

100 µg/mL chloramphenicol [Sigma R4408], and 10 units/mL DNase I [Epicenter, D9905K]).

The lysates were spun at 3,000 g for 3 min, and the supernatant was transferred to a new tube

and subsequently centrifuged at 20,000 g for 10 min. The supernatant was transferred to a new

tube and the RNA concentration was determined with 10x dilutions using the Qubit RNA HS

assay (Thermo Fisher Scientific; Q32852). Aliquots of 100 µL and 200 µL of the lysates were

made, and they were flash frozen in liquid nitrogen and stored at -80°C until further processing.

Ribo-seq library construction

Ribosome footprints were obtained using 200 µL of the lysates described above, and

sequencing libraries were constructed according to our previous method [47] with the following

modifications: briefly, after RNase I digestion (50 units nuclease per 40 µg of RNA; the nuclease

was included in TruSeq Mammalian Ribo Profile Kit, Illumina, RPHMR12126), adding 15 µl of

SUPERase-IN (Invitrogen, AM2696), and passing through a size exclusion column (Illustra

MicroSpin S-400 HR Columns; GE Healthcare; 27-5140-01), RNA > 17 nt was isolated with the

RNA Clean & Concentrator-25 kit (Zymo Research, R1017) and separated on 15% urea-TBE

gels (Invitrogen, EC68852BOX). Gel slices between 28 and 30 nt were isolated, and the RNAs

were purified as previously described. Next, rRNA depletion was performed using RiboZero

Plant Leaf kit (Illumina, MRZPL1224) in one quarter of the recommended reaction volume. The

major change we made here was to gel-purify the ribosome footprints prior to the rRNA

depletion. In our previous protocol, we purified the RNA twice (first >17 nt and then <200 nt) to

enrich ribosome footprints before the rRNA depletion and gel purification. We reasoned that

some footprints might be lost during the two RNA purification steps. In addition, the rRNA

depletion step requires a limited amount of input RNA according to the manufacturer's

recommendations. Therefore, we expected modifying the order of these steps could maximize

the input and minimize the footprint loss during the processes. Ribo-seq libraries were then

constructed using the TruSeq Mammalian Ribo Profile Kit (illumina, RPHMR12126) as

previously described with nine cycles of PCR amplification. Libraries with equal molarity were

pooled and sequenced on a Hi-Seq 4000 sequencer using single-end 50-bp sequencing.

RNA-seq library construction

Total RNA greater than 200 nt was purified from 100 µL of the lysates described above using

the RNA Clean & Concentrator-25 kit (Zymo Research, R1017) as previously described [47].

RNA integrity was evaluated using a Bioanalyzer (Agilent) RNA pico chip, and RNA integrity

numbers (RINs) ranging from 7.2 to 7.7 were observed among the samples. A total of 4 µg of

RNA per sample was subjected to rRNA depletion using the RiboZero Plant Leaf kit (Illumina,

MRZPL1224) following the manufacturer's recommendations. Then, 100 ng of rRNA-depleted

RNA was fragmented to around 200-nt long based on the RIN reported by the Bioanalyzer, and

strand-specific sequencing libraries were made using the NEBNext Ultra II Directional RNA

Library Prep Kit (New England Biolabs, E7760S) with eight cycles of amplification. Libraries of

equal molarity were pooled and sequenced on a Hi-Seq 4000 using paired-end 100-bp

sequencing.

Data pre-processing and analysis

Data pre-processing and analysis were performed similarly to that previously described [47],

except that the Araport11 annotation was used in this study. Briefly, for Ribo-seq libraries, the

adaptor (AGATCGGAAGAGCACACGTCT) was clipped with fastx_clipper (FASTX toolkit

v0.0.14) (http://hannonlab.cshl.edu/fastx_tool- kit/). For both RNA-seq and Ribo-seq, we used

Bowtie2 (v2.3.4.1) [106] to remove rRNA/tRNA/snRNA/snoRNA sequences. Both the RNA-seq

and Ribo-seq reads were mapped to the transcriptome with STAR aligner [107] (RNA-seq

parameters: --outFilterMismatchNmax 2 --outFilterMultimapNmax 20 --outFilterType BySJout --

alignSJoverhangMin 8 --alignSJDBoverhangMin 2; all parameters used for Ribo-seq were the

same except --outFilterMismatchNmax 1). We then used the bam files from both RNA-seq and

Ribo-seq for RiboTaper (v1.3.1a) [51] to identify translated ORFs. The distribution of Ribo-seq

reads in different genome features was calculated using Ribo-seQC. The Ribo-seq metaplot

was created using the create_metaplots.bash function in RiboTaper. The Ribo-seq read lengths

and offsets for RiboTaper were 24, 25, 26, 27, 28 and 8, 9, 10, 11, 12, respectively. Combining

both 20-min and 60-min samples, the total reads mapped to the genome for Ribo-seq and RNA-

seq were 298.01 million and 180.67 million, respectively. The discovered uORFs and mORFs

were extracted from the RiboTaper output ORF_max_filt file. To calculate translation efficiency,

we first used STAR to map the RNA-seq and Ribo-seq reads to the CDS of annotated coding

genes. The resulting bam files were used to quantify the transcripts per million (TPM) of each

gene with RSEM (v1.3.1) [108]. Then, translation efficiency was calculated by dividing the Ribo-

seq TPM by the RNA-seq TPM. The root and shoot data from Hsu et al., 2016 [47] were

reanalyzed using the Araport11 annotation. The P_sites_all files from RiboTaper were

processed using the following code: cut -f 1,3,6 P_sites_all | sort | uniq -c | sed -r 's/^(*[^]+)

+/\1\t/' > output.txt to combine the read counts at each P-site. Data visualization and statistical

analysis were performed in R (v4.0.3) [109]. In particular, the gene levels from the Ribo-seq and

RNA-seq reads were visualized with RiboPlotR [54].

The mRNA half-life and mRNA decay rate data were downloaded from Szabo et al. and

Sorenson et al., respectively [70, 85]. The quantitative proteomics data for Arabidopsis shoots

and roots were downloaded from Song et al. [71]. The expression levels (TPM) used for

investigating mRNA levels of uORF genes in different tissues, developmental stages, and

ecotypes in Arabidopsis and in tomato were directly downloaded from the EMBO-EBI

expression atlas (E-MTAB-7978: Arabidopsis tissue atlas, E-GEOD-53197: 17 Arabidopsis

thaliana accessions, E-MTAB-4812: tomato root, leaf, flower [two stages], and fruit [six stages],

and E-MTAB-4813: three longitudinal sections of six stages during tomato fruit development)

[74]. Our previous Ribo-seq and RNA-seq data for Arabidopsis roots and shoots were described

in [47]; our previous Ribo-seq and RNA-seq data for tomato roots were described in [72]. The

published RNA-seq data for the NMD mutants *upf1* (*lba1*) *upf3-1* and its wild-type control (GEO:

GSE41432), *smg7-1 pad4* and its *pad4* control (GEO: GSE55884), *upf1-3 pad4* and its *pad4*

control (GEO: GSE146340), and *upf2-10* and its wild-type control (SRA: SRP056795) were

downloaded, and the mRNA levels were quantified by Kalisto [110] and the TPM values were

further analyzed to compare the steady-state mRNA levels in different categories of genes.

AVAILABILITY OF DATA AND MATERIALS

All raw and processed sequencing data generated in this study have been submitted to the

NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession

number GSE183264. The genomic locations of TuORFs are available on TAIR genome browser

(https://www.arabidopsis.org). Codes for this study are available at

https://github.com/hsinyenwu/Wu-and-Hsu-2022-TuORF-nonNMD

COMPETING INTEREST STATEMENT

The authors declare that they have no competing interests.

FUNDING

This work was supported by a National Science Foundation grant (2051885) and a Michigan

State University startup grant to PYH.

ACKNOWLEDGEMENTS

We thank Robert Last and Melissa Lehti-Shiu for critical reading and helpful comments on the

manuscript. This work used the Vincent J. Coates Genomics Sequencing Laboratory at UC

Berkeley, supported by an NIH S10 OD018174 Instrumentation Grant.

AUTHOR CONTRIBUTIONS

HLW and PYH designed the research, PYH performed the sequencing experiments, HLW

analyzed the sequencing data, HLW and PYH interpreted the data, and HLW and PYH wrote

the paper.

REFERENCES

1. Calvo SE, Pagliarini DJ, Mootha VK. Upstream open reading frames cause widespread

reduction of protein expression and are polymorphic among humans. Proceedings of the

National Academy of Sciences of the United States of America. 2009;106:7507–12.

2. Xu G, Yuan M, Ai C, Liu L, Zhuang E, Karapetyan S, et al. uORF-mediated translation allows

engineered plant disease resistance without fitness costs. Nature. 2017;545:491–4.

3. Zhang H, Si X, Ji X, Fan R, Liu J, Chen K, et al. Genome editing of upstream open reading

frames enables translational control in plants. Nature Biotechnology. 2018;36:894–8.

4. Lee DSM, Park J, Kromer A, Baras A, Rader DJ, Ritchie MD, et al. Disrupting upstream

translation in mRNAs is associated with human disease. Nature Communications 2021 12:1.

2021;12:1–14.

5. Kim BH, Cai X, Vaughn JN, Von Arnim AG. On the functions of the h subunit of eukaryotic

initiation factor 3 in late stages of translation initiation. Genome Biology. 2007;8:R60.

6. Johnstone TG, Bazzini AA, Giraldez AJ. Upstream ORFs are prevalent translational

repressors in vertebrates. The EMBO Journal. 2016;35:706–23.

7. Zhang H, Wang Y, Wu X, Tang X, Wu C, Lu J. Determinants of genome-wide distribution and

evolution of uORFs in eukaryotes. Nature Communications. 2021;12:1–17.

8. Chew GL, Pauli A, Schier AF. Conservation of uORF repressiveness and sequence features

in mouse, human and zebrafish. Nature Communications. 2016;7:11663.

9. Liu MJ, Wu SH, Wu JF, Lin WD, Wu YC, Tsai TY, et al. Translational landscape of

photomorphogenic Arabidopsis. Plant Cell. 2013;25:3699–710.

10. Wu HYL, Song G, Walley JW, Hsu PY. The tomato translational landscape revealed by

transcriptome assembly and ribosome profiling. Plant Physiology. 2019;181:367–80.

11. Lin Y, May GE, Kready H, Nazzaro L, Mao M, Spealman P, et al. Impacts of uORF codon

identity and position on translation regulation. Nucleic Acids Research. 2019;47:9358–67.

12. Hinnebusch AG, Ivanov IP, Sonenberg N. Translational control by 5′-untranslated regions of

eukaryotic mRNAs. Science. 2016;352:1413–6.

13. Dever TE, Ivanov IP, Sachs MS. Conserved Upstream Open Reading Frame Nascent

Peptides That Control Translation. Annual Review of Genetics. 2020;54:237–64.

14. Kozak M. Initiation of translation in prokaryotes and eukaryotes. Gene. 1999;234:187–208.

15. He F, Jacobson A. Nonsense-Mediated mRNA Decay: Degradation of Defective Transcripts

Is only Part of the Story. Annual Review of Genetics. 2015;49:339–66.

16. Kurosaki T, Popp MW, Maquat LE. Quality and quantity control of gene expression by

nonsense-mediated mRNA decay. Nature Reviews Molecular Cell Biology 2019 20:7.

2019;20:406–20.

17. Isken O, Kim YK, Hosoda N, Mayeur GL, Hershey JWB, Maquat LE. Upf1 Phosphorylation

Triggers Translational Repression during Nonsense-Mediated mRNA Decay. Cell.

2008;133:314–27.

18. Kurosaki T, Li W, Hoque M, Popp MWL, Ermolenko DN, Tian B, et al. A Post-Translational

regulatory switch on UPF1 controls targeted mRNA degradation. Genes and Development.

2014;28:1900–16.

19. Raxwal VK, Riha K. Nonsense mediated RNA decay and evolutionary capacitance.

Biochimica et Biophysica Acta. 2016;1859:1538–43.

20. Lloyd JPB, Davies B. SMG1 is an ancient nonsense-mediated mRNA decay effector. The

Plant journal : for cell and molecular biology. 2013;76:800–10.

21. Kurihara Y, Matsui A, Hanada K, Kawashima M, Ishida J, Morosawa T, et al. Genome-wide

suppression of aberrant mRNA-like noncoding RNAs by NMD in Arabidopsis. Proceedings of

the National Academy of Sciences of the United States of America. 2009;106:2453–8.

22. Kalyna M, Simpson CG, Syed NH, Lewandowska D, Marquez Y, Kusenda B, et al.

Alternative splicing and nonsense-mediated decay modulate expression of important regulatory

genes in Arabidopsis. Nucleic Acids Research. 2012;40:2454–69.

23. Drechsel G, Kahles A, Kesarwani AK, Stauffer E, Behr J, Drewe P, et al. Nonsense-

mediated decay of alternative precursor mRNA splicing variants is a major determinant of the

Arabidopsis steady state transcriptome. The Plant Cell. 2013;25:3726–42.

24. Degtiar E, Fridman A, Gottlieb D, Vexler K, Berezin I, Farhi R, et al. The feedback control of

UPF3 is crucial for RNA surveillance in plants. Nucleic Acids Research. 2015;43:4219–35.

25. Raxwal VK, Simpson CG, Gloggnitzer J, Entinze JC, Guo W, Zhang R, et al. Nonsense-

Mediated RNA Decay Factor UPF1 Is Critical for Posttranscriptional and Translational Gene

Regulation in Arabidopsis. The Plant Cell. 2020;32:2725–41.

26. Jeong HJ, Kim YJ, Kim SH, Kim YH, Lee IJ, Kim YK, et al. Nonsense-mediated mRNA

decay factors, UPF1 and UPF3, contribute to plant defense. Plant and Cell Physiology.

2011;52:2147–56.

27. Rayson S, Arciga-Reyes L, Wootton L, de Torres Zabala M, Truman W, Graham N, et al. A

role for nonsense-mediated mrna decay in plants: Pathogen responses are induced in

arabidopsis thaliana nmd mutants. PLoS ONE. 2012;7:e31917.

28. Riehs-Kearnan N, Gloggnitzer J, Dekrout B, Jonak C, Riha K. Aberrant growth and lethality

of Arabidopsis deficient in nonsense-mediated RNA decay factors is caused by autoimmune-like

response. Nucleic Acids Research. 2012;40:5615–24.

29. Filichkin SA, Cumbie JS, Dharmawardhana P, Jaiswal P, Chang JH, Palusa SG, et al.

Environmental stresses modulate abundance and timing of alternatively spliced circadian

transcripts in Arabidopsis. Molecular Plant. 2015;8:207–27.

30. Merchante C, Brumos J, Yun J, Hu Q, Spencer KR, Enríquez P, et al. Gene-specific

translation regulation mediated by the hormone-signaling molecule EIN2. Cell. 2015;163:684–

97.

31. Chiam N-C, Fujimura T, Sano R, Akiyoshi N, Hiroyama R, Watanabe Y, et al. Nonsense-

Mediated mRNA Decay Deficiency Affects the Auxin Response and Shoot Regeneration in

Arabidopsis. Plant and Cell Physiology. 2019;60:2000–14.

32. Yoine M, Ohto MA, Onai K, Mita S, Nakamura K. The lba1 mutation of UPF1 RNA helicase

involved in nonsense-mediated mRNA decay causes pleiotropic phenotypic changes and

altered sugar signalling in Arabidopsis. The Plant Journal. 2006;47:49–62.

33. Kertész S, Kerényi Z, Mérai Z, Bartos I, Pálfy T, Barta E, et al. Both introns and long 3′-

UTRs operate as cis-acting elements to trigger nonsense-mediated decay in plants. Nucleic

Acids Research. 2006;34:6147–57.

34. Hurt JA, Robertson AD, Burge CB. Global analyses of UPF1 binding and function reveal

expanded scope of nonsense-mediated mRNA decay. Genome Research. 2013;23:1636–50.

35. Baird TD, Cheng KCC, Chen YC, Buehler E, Martin SE, Inglese J, et al. ICE1 promotes the

link between splicing and nonsense-mediated mRNA decay. eLife. 2018;7.

36. Nyikó T, Sonkoly B, Mérai Z, Benkovics AH, Silhavy D. Plant upstream ORFs can trigger

nonsense-mediated mRNA decay in a size-dependent manner. Plant Molecular Biology.

2009;71:367–78.

37. Uchiyama-Kadokura N, Murakami K, Takemoto M, Koyanagi N, Murota K, Naito S, et al.

Polyamine-responsive ribosomal arrest at the stop codon of an upstream open reading frame of

the AdoMetDC1 gene triggers nonsense-mediated mRNA decay in Arabidopsis thaliana. Plant

and Cell Physiology. 2014;55:1556–67.

38. Kurihara Y, Makita Y, Kawashima M, Fujita T, Iwasaki S, Matsui M. Transcripts from

downstream alternative transcription start sites evade uORF-mediated inhibition of gene

expression in Arabidopsis. Proceedings of the National Academy of Sciences of the United

States of America. 2018;115:7831–6.

39. Saul H, Elharrar E, Gaash R, Eliaz D, Valenci M, Akua T, et al. The upstream open reading

frame of the Arabidopsis AtMHX gene has a strong impact on transcript accumulation through

the nonsense-mediated mRNA decay pathway. The Plant Journal. 2009;60:1031–42.

40. Tanaka M, Sotta N, Yamazumi Y, Yamashita Y, Miwa K, Murota K, et al. The minimum open

reading frame, AUG-Stop, induces boron-dependent ribosome stalling and mRNA degradation.

The Plant Cell. 2016;28:2830–49.

41. Li YR, Liu MJ. Prevalence of alternative AUG and non-AUG translation initiators and their

regulatory effects across plants. Genome Research. 2020;30:1418–33.

42. Nagarajan VK, Kukulich PM, Von Hagel B, Green PJ. RNA degradomes reveal substrates

and importance for dark and nitrogen stress responses of Arabidopsis XRN4. Nucleic Acids

Research. 2019;47:9216–30.

43. Ohtani M, Wachter A. NMD-Based Gene Regulation - A Strategy for Fitness Enhancement

in Plants? Plant and Cell Physiology. 2019;60:1953–60.

44. Ingolia NT, Ghaemmaghami S, Newman JRS, Weissman JS. Genome-wide analysis in vivo

of translation with nucleotide resolution using ribosome profiling. Science. 2009;324:218–23.

45. Brar GA, Weissman JS. Ribosome profiling reveals the what, when, where and how of

protein synthesis. Nature Reviews Molecular Cell Biology. 2015;16:651–64.

46. Juntawong P, Girke T, Bazin J, Bailey-Serres J. Translational dynamics revealed by

genome-wide profiling of ribosome footprints in Arabidopsis. Proceedings of the National

Academy of Sciences of the United States of America. 2014;111:E203–12.

47. Hsu PY, Calviello L, Wu H-YL, Li F-W, Rothfels CJ, Ohler U, et al. Super-resolution

ribosome profiling reveals unannotated translation events in Arabidopsis. Proceedings of the

National Academy of Sciences of the United States of America. 2016;113:E7126–35.

48. Hu Q, Merchante C, Stepanova AN, Alonso JM, Heber S. Genome-Wide Search for

Translated Upstream Open Reading Frames in Arabidopsis Thaliana. IEEE Transactions on

Nanobioscience. 2016;15:150–9.

49. Bazin J, Baerenfaller K, Gosai SJ, Gregory BD, Crespi M, Bailey-Serres J. Global analysis

of ribosome-associated noncoding RNAs unveils new modes of translational regulation.

Proceedings of the National Academy of Sciences of the United States of America.

2017;114:E10018–27.

50. Kurihara Y, Makita Y, Shimohira H, Fujita T, Iwasaki S, Matsui M. Translational landscape of

protein-coding and non-protein-coding RNAs upon light exposure in Arabidopsis. Plant and Cell

Physiology. 2020;61:536–45.

51. Calviello L, Mukherjee N, Wyler E, Zauber H, Hirsekorn A, Selbach M, et al. Detecting

actively translated open reading frames in ribosome profiling data. Nature Methods.

2016;13:165–70.

52. Xu Z, Hu L, Shi B, Geng S, Xu L, Wang D, et al. Ribosome elongating footprints denoised

by wavelet transform comprehensively characterize dynamic cellular translation events. Nucleic

Acids Research. 2018;46:e109.

53. Harnett D, Meerdink E, Calviello L, Sydow D, Ohler U. Genome-Wide Analysis of Actively

Translated Open Reading Frames Using RiboTaper/ORFquant. Methods in Molecular Biology.

2021;2252:331–46.

54. Wu H-YL, Hsu PY. RiboPlotR: a visualization tool for periodic Ribo-seq reads. Plant

Methods. 2021;17:1–7.

55. Von Arnim AG, Jia Q, Vaughn JN. Regulation of plant translation by upstream open reading

frames. Plant Science. 2014;214:1–12.

56. Silverstone AL, Ciampaglio CN, Sun TP. The Arabidopsis RGA gene encodes a

transcriptional regulator repressing the gibberellin signal transduction pathway. The Plant Cell.

1998;10:155–69.

57. Strayer C, Oyama T, Schultz TF, Raman R, Somers DE, Mas P, et al. Cloning of the

Arabidopsis clock gene TOC1, an autoregulatory response regulator homolog. Science.

2000;289:768–71.

58. Di Laurenzio L, Wysocka-Diller J, Malamy JE, Pysh L, Helariutta Y, Freshour G, et al. The

SCARECROW gene regulates an asymmetric cell division that is essential for generating the

radial organization of the Arabidopsis root. Cell. 1996;86:423–33.

59. Ellis CM, Nagpal P, Young JC, Hagen G, Guilfoyle TJ, Reed JW. AUXIN RESPONSE

FACTOR1 and AUXIN RESPONSE FACTOR2 regulate senescence and floral organ abscission

in Arabidopsis thaliana. Development. 2005;132:4563–74.

60. Nagpal P, Ellis CM, Weber H, Ploense SE, Barkawi LS, Guilfoyle TJ, et al. Auxin response

factors ARF6 and ARF8 promote jasmonic acid production and flower maturation. Development.

2005;132:4107–18.

61. Ni M, Tepperman JM, Quail PH. PIF3, a phytochrome-interacting factor necessary for

normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. Cell.

1998;95:657–67.

62. Huq E, Quail PH. PIF4, a phytochrome-interacting bHLH factor, functions as a negative

regulator of phytochrome B signaling in Arabidopsis. EMBO Journal. 2002;21:2441–50.

63. Leivar P, Monte E, Al-Sady B, Carle C, Storer A, Alonso JM, et al. The Arabidopsis

phytochrome-interacting factor PIF7, together with PIF3 and PIF4, regulates responses to

prolonged red light by modulating phyB levels. The Plant Cell. 2008;20:337–52.

64. Sharrock RA, Quail PH. Novel phytochrome sequences in Arabidopsis thaliana: structure,

evolution, and differential expression of a plant regulatory photoreceptor family. Genes &

Development. 1989;3:1745–57.

65. Devlin PF, Patel SR, Whitelam GC. Phytochrome E Influences Internode Elongation and

Flowering Time in Arabidopsis. The Plant Cell. 1998;10:1479–87.

66. Zhang SH, Lawton MA, Hunter T, Lamb CJ. atpk1, a novel ribosomal protein kinase gene

from Arabidopsis. I. Isolation, characterization, and expression. Journal of Biological Chemistry.

1994;269:17586–92.

67. Mizoguchi T, Hayashida N, Yamaguchi-Shinozaki K, Kamada H, Shinozaki K. Two genes

that encode ribosomal-protein S6 kinase homologs are induced by cold or salinity stress in

Arabidopsis thaliana. FEBS Letters. 1995;358:199–204.

68. Park YS, Hong SW, Oh SA, Kwak JM, Lee HH, Nam HG. Two putative protein kinases from

Arabidopsis thaliana contain highly acidic domains. Plant Molecular Biology. 1993;22:615–24.

69. Boudsocq M, Barbier-Brygoo H, Laurière C. Identification of nine sucrose nonfermenting 1-

related protein kinases 2 activated by hyperosmotic and saline stresses in Arabidopsis thaliana.

Journal of Biological Chemistry. 2004;279:41758–66.

70. Szabo EX, Reichert P, Lehniger MK, Ohmer M, de Francisco Amorim M, Gowik U, et al.

Metabolic labeling of RNAs uncovers hidden features and dynamics of the Arabidopsis

transcriptome. The Plant Cell. 2020;32:871–87.

71. Song G, Hsu PY, Walley JW. Assessment and Refinement of Sample Preparation Methods

for Deep and Quantitative Plant Proteome Profiling. Proteomics. 2018;18:e1800220.

72. Wu H-YL, Song G, Walley JW, Hsu PY. Translational landscape in tomato revealed by

transcriptome assembly and ribosome profiling. bioRxiv. 2019;181:367–80.

73. Clarke JT, Warnock RCM, Donoghue PCJ. Establishing a time-scale for plant evolution. The

New Phytologist. 2011;192:266–301.

74. Papatheodorou I, Moreno P, Manning J, Fuentes AMP, George N, Fexova S, et al.

Expression Atlas update: From tissues to single cells. Nucleic Acids Research. 2020;48:D77–

83.

75. Hayden CA, Jorgensen RA. Identification of novel conserved peptide uORF homology

groups in Arabidopsis and rice reveals ancient eukaryotic origin of select groups and

preferential association with transcription factor-encoding genes. BMC Biology. 2007;5:32.

76. Jorgensen RA, Dorantes-Acosta AE. Conserved peptide upstream open reading frames are

associated with regulatory genes in angiosperms. Frontiers in Plant Science. 2012;3 AUG:191.

77. Takahashi H, Takahashi A, Naito S, Onouchi H. BAIUCAS: A novel BLAST-based algorithm

for the identification of upstream open reading frames with conserved amino acid sequences

and its application to the Arabidopsis thaliana genome. Bioinformatics. 2012;28:2231–41.

78. Vaughn JN, Ellingson SR, Mignone F, Arnim A von. Known and novel post-transcriptional

regulatory sequences are conserved across plant families. RNA. 2012;18:368–84.

79. Van Der Horst S, Snel B, Hanson J, Smeekens S. Novel pipeline identifies new upstream

ORFs and non-AUG initiating main ORFs with conserved amino acid sequences in the 5′ leader

of mRNAs in Arabidopsis thaliana. RNA. 2019;25:292–304.

80. Wolfa YI, Novichkovb PS, Kareva GP, Koonina E V., Lipmana DJ. The universal distribution

of evolutionary rates of genes and distinct characteristics of eukaryotic genes of different

apparent ages. Proceedings of the National Academy of Sciences of the United States of

America. 2009;106:7273–80.

81. Cui X, Lv Y, Chen M, Nikoloski Z, Twell D, Zhang D. Young genes out of the male: An

insight from evolutionary age analysis of the pollen transcriptome. Molecular Plant. 2015;8:935–

45.

82. Gloggnitzer J, Akimcheva S, Srinivasan A, Kusenda B, Riehs N, Stampfl H, et al. Nonsense-

mediated mRNA decay modulates immune receptor levels to regulate plant antibacterial

defense. Cell Host and Microbe. 2014;16:376–90.

83. Zinshteyn B, Sinha NK, Enam SU, Koleske B, Green R. Translational repression of NMD

targets by GIGYF2 and EIF4E2. PLoS Genetics. 2021;17 10 October:e1009813.

84. Sieburth LE, Vincent JN. Beyond transcription factors: Roles of mrna decay in regulating

gene expression in plants [version 1; referees: 3 approved]. F1000Research. 2018;7:1940.

85. Sorenson RS, Deshotel MJ, Johnson K, Adler FR, Sieburth LE. Arabidopsis mRNA decay

landscape arises from specialized RNA decay substrates, decapping-mediated feedback, and

redundancy. Proceedings of the National Academy of Sciences of the United States of America.

2018;115:E1485–94.

86. Deyholos MK, Cavaness GF, Hall B, King E, Punwani J, Van Norman J, et al. Varicose, a

WD-domain protein, is required for leaf blade. Development. 2003;130:6577–88.

87. Goeres DC, Van Norman JM, Zhang W, Fauver NA, Spencer M Lou, Sieburth LE.

Components of the Arabidopsis mRNA decapping complex are required for early seedling

development. The Plant Cell. 2007;19:1549–64.

88. Xu J, Yang JY, Niu QW, Chua NH. Arabidopsis DCP2, DCP1, and VARICOSE form a

decapping complex required for postembryonic development. The Plant Cell. 2006;18:3386–98.

89. Zhang W, Murphy C, Sieburth LE. Conserved RNaseII domain protein functions in

cytoplasmic mRNA decay and suppresses Arabidopsis decapping mutant phenotypes.

Proceedings of the National Academy of Sciences of the United States of America.

2010;107:15981–5.

90. Andreev DE, Arnold M, Kiniry SJ, Loughran G, Michel AM, Rachinskii D, et al. TASEP

modelling provides a parsimonious explanation for the ability of a single uORF to derepress

translation during the integrated stress response. eLife. 2018;7:e32563.

91. Kerényi F, Wawer I, Sikorski PJ, Kufel J, Silhavy D. Phosphorylation of the N- and C-

terminal UPF1 domains plays a critical role in plant nonsense-mediated mRNA decay. Plant

Journal. 2013;76:836–48.

92. Chang HM, Triboulet R, Thornton JE, Gregory RI. A role for the Perlman syndrome

exonuclease Dis3l2 in the Lin28-let-7 pathway. Nature. 2013;497:244–8.

93. Lubas M, Damgaard CK, Tomecki R, Cysewski D, Jensen TH, Dziembowski A.

Exonuclease hDIS3L2 specifies an exosome-independent 3'-5' degradation pathway of human

cytoplasmic mRNA. The EMBO journal. 2013;32:1855–68.

94. Imai A, Hanzawa Y, Komura M, Yamamoto KT, Komeda Y, Takahashi T. The dwarf

phenotype of the Arabidopsis acl5 mutant is suppressed by a mutation in an upstream ORF of a

bHLH gene. Development. 2006;133:3575–85.

95. Cai Q, Fukushima H, Yamamoto M, Ishii N, Sakamoto T, Kurata T, et al. The SAC51 family

plays a central role in thermospermine responses in arabidopsis. Plant and Cell Physiology.

2016;57:1583–92.

96. Presnyak V, Alhusaini N, Chen YH, Martin S, Morris N, Kline N, et al. Codon optimality is a

major determinant of mRNA stability. Cell. 2015;160:1111–24.

97. Radhakrishnan A, Chen YH, Martin S, Alhusaini N, Green R, Coller J. The DEAD-Box

Protein Dhh1p Couples mRNA Decay and Translation by Monitoring Codon Optimality. Cell.

2016;167:122-132.e9.

98. Beelman CA, Parker R. Differential effects of translational inhibition in cis and in trans on the

decay of the unstable yeast MFA2 mRNA. Journal of Biological Chemistry. 1994;269:9687–92.

99. LaGrandeur T, Parker R. The cis acting sequences responsible for the differential decay of

the unstable MFA2 and stable PGK1 transcipts in yeast include the context of the translational

start codon. RNA. 1999;5:420–33.

100. Schwartz DC, Parker R. mRNA Decapping in Yeast Requires Dissociation of the Cap

Binding Protein, Eukaryotic Translation Initiation Factor 4E. Molecular and Cellular Biology.

2000;20:7933–42.

101. Chan LY, Mugler CF, Heinrich S, Vallotton P, Weis K. Non-invasive measurement of

mRNA decay reveals translation initiation as the major determinant of mRNA stability. eLife.

2018;7:e32536.

102. Hou C-Y, Lee W-C, Chou H-C, Chen A-P, Chou S-J, Chen H-M. Global analysis of

truncated RNA ends reveals new insights into ribosome stalling in plants. The Plant Cell.

2016;28:2398–416.

103. Yu X, Willmann MR, Anderson SJ, Gregory BD. Genome-wide mapping of uncapped and

cleaved transcripts reveals a role for the nuclear mrna cap-binding complex in cotranslational

rna decay in Arabidopsis. The Plant Cell. 2016;28:2385–97.

104. Shah P, Ding Y, Niemczyk M, Kudla G, Plotkin JB. Rate-Limiting Steps in Yeast Protein

Translation. Cell. 2013;153:1589–601.

105. Lawson MR, Lessen LN, Wang J, Prabhakar A, Corsepius NC, Green R, et al.

Mechanisms that ensure speed and fidelity in eukaryotic translation termination. Science.

2021;373:876–82.

106. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods.

2012;9:357–9.

107. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29:15–21.

108. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. BMC Bioinformatics. 2011;12:323.

109. R Core Team (2013). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. 2017.

110. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq

quantification. Nature biotechnology. 2016;34:525–7.

FIGURE and FIGURE LEGENDS

Figure 1

Figure 1. Enhanced Ribo-seq coverage improves uORF identification

(A) Metagene analysis of 28-nt ribosome footprints mapped to regions near the start and

stop codons of annotated ORFs. The reads are presented with their first nt at the P-site,

which is the 13th nt for 28-nt footprints. The reads are colored in red, blue, and green to

indicate they are in the first (expected), second, and third reading frames, respectively.

The majority of footprints were mapped to the CDS in the expected reading frame

(92.4% in frame).

(B) Genomic features mapped by ribosome footprints. Reads that mapped to nuclear (Nuc)-,

mitochondrial (Mt)-, and plastid (Pt)-encoded genes are shown.

(C) Length distribution of ribosome footprints. Reads that mapped to nuclear-encoded genes

are presented.

(D) Distinct P-sites detected in 1 to 20 million randomly selected ribosome footprints from

our current and previous datasets.

(E) Numbers of TuORFs, CPuORFs, and mORFs identified in our current and previous

datasets.

(F-G) RNA-seq and Ribo-seq profiles of *ATKAO1 and SAMDC1* from our current data. RNA-

seq coverage is shown with a light-yellow background. Ribo-seq reads are presented

with their first nt at the P-site, and they are colored in red, blue, and green to indicate

they are in the first (expected), second, and third reading frames, respectively. Within the

gene models, yellow boxes represent the TuORFs, black boxes represent the annotated

mORFs, and gray and white regions indicate 5' UTRs and 3' UTRs, respectively. The

specific isoform being considered is indicated to the left of the gene model. Within the

profiles, green and orange vertical dashed lines represent translation start and stop,

respectively, for the TuORF. Black and gray vertical dashed lines represent translation

start and stop, respectively, for the annotated mORF.

924
925
926

Figure 2

Arabidopsis   Tomato

A ... **B** ... **C** ... **D**   **E** ... **F**

(A-D figures showing mRNA level (TPM), Translation efficiency, mRNA half-life, Protein level (MS count); E-F figures showing mRNA level (TPM), Translation efficiency)

A mRNA level (TPM)
1752 6442 13256
Ratio: 1.51 1.04 1

B Translation efficiency
1748 6344 13060
0.62 0.89 1

C mRNA half-life
1449 4053 8015
0.86 0.94 1

D Protein level (MS count)
1091 3334 7231
0.57 0.79 1

E mRNA level (TPM)
1155 6654 7608
1.46 0.63 1

F Translation efficiency
1155 6654 7608
0.59 0.82 1

G mRNA level (TPM)
TuORF UuORF no-uORF

Figure 2. TuORF-containing genes have higher mRNA abundance but are associated with

lower translation efficiency, shorter mRNA half-lives, and lower protein abundance

(A-D) The mRNA levels, translation efficiency, mRNA half-lives, and protein levels of TuORF,

UuORF, and no-uORF genes in Arabidopsis seedlings. The mRNA half-life data, which

were measured with metabolic labeling in Arabidopsis seedlings, were extracted from [70],

and the protein abundance as determined by quantitative proteomics in Arabidopsis

seedling shoots were extracted from [71]. Within the boxplots, the dashed lines mark the

median level of the no-uORF genes; the number of genes in each category is listed below

the lower whisker. The ratios underneath the plots indicate the median of each group

normalized to that of the no-uORF genes. The statistical significance was determined by

Wilcoxon signed-rank test. The adjusted p-value was determined using the Benjamini and

Yekutieli procedure to control for the false discovery rate in multiple testing (* 0.05 > p >

0.01, ** 0.01 > p > 0.001, *** 0.001 > p > 1e-4, **** 1e-4 > p > 0). These statistical

analyses and graphic layouts are the same for all boxplots in the study.

(E-F) The mRNA levels and translation efficiency of TuORF, UuORF, and no-uORF genes from

a reanalysis of our previous tomato root data [72]. The dashed lines mark the median level

of the no-uORF genes.

(G) The mRNA levels of TuORF, UuORF, and no-uORF genes in 46 Arabidopsis tissues and

developmental stages from independent studies. The TPM values from RNA-seq

quantification were downloaded from the EMBO-EBI expression atlas [74]. Additional

Arabidopsis and tomato RNA-seq datasets were presented in Figure S5.

953
954

Figure 3

Figure 3. TuORF genes are larger than other genes

Lengths of the 5' UTRs (A), CDSs (B), 3' UTRs (C), and entire transcripts (D) of TuORF,

UuORF, and no-uORF genes. Within the boxplots, the dashed lines mark the median level of

the no-uORF genes; the number of genes in each category is listed below the lower whisker.

The ratios underneath the plots indicate the median of each group normalized to that of the no-

uORF genes. The statistical significance was determined as described in the Figure 2 legend.

Figure 4: TuORF mRNAs are not upregulated in the NMD mutants

The ratios of mRNA levels between NMD mutants and their control plants from four RNA-seq

datasets [23, 25, 30, 82] were used to examine the effects of the NMD pathway on NMD targets

(NMDt) and TuORF genes. (A) *upf1* (*lba1* allele) *upf3-1* double mutant compared with wild type

(Col-0). (B) *upf2-10* compared with wild type (Col-0). (C) *upf1-3 pad4* compared with *pad4*. (D)

smg7-1 pad4 compared to *pad4*. See Table S3 for specific growth conditions for each dataset.

(E-H) Same datasets as 4A-D, but the TuORF genes were grouped based on low or high

relative TuORF translation efficiency (RTE) (i.e., TE $_{TuORF}$ / TE $_{mORF}$). (I-L) Same datasets as 4A-

D, but the TuORF genes were grouped based on the TuORF number(s). Within the boxplots,

the dashed lines mark the median level of other mRNAs; the number of genes in each category

is listed below the lower whisker. The ratios underneath the plots indicate the median of each

group normalized to that of other mRNAs. The statistical significance was determined as

described in the Figure 2 legend.

Figure 5. Expression patterns of NMD targets and TuORF mRNAs.

Cumulative plots (A-D) and boxplots (E-H) for the mRNA levels (A, E), translation efficiency (B, F), mRNA half-lives (C, G), and protein levels (D, H) of NMD targets (NMDt) without or with TuORFs compared with those of TuORF mRNAs and other mRNAs. Within the boxplots, the dashed lines mark the median level of other mRNAs; the number of genes in each category is listed below the lower whisker. The ratios underneath the plots indicate the median of each group normalized to that of other mRNAs. The statistical significance was determined as described in the Figure 2 legend.

Figure 6

Figure 6. TuORF mRNAs and NMD targets are degraded through different mRNA decay

mechanisms

Cumulative plots (A-D) and boxplots (E-H) showing the mRNA decay rates for NMD targets and

TuORF mRNAs in wild type and mutants defective in either 5' to 3' decay (*vcs-7*), 3' to 5' decay

(*sov*), or both (*vcs sov*). The mRNA decay rates were extracted from [85]. Because Col-0 is

naturally an *sov* mutant, the wild type here is Col-0 carrying the functional *SOV* allele from

Landsberg *erecta* driven by its native promoter [87, 89]. The *sov* mutant here is Col-0, and *vcs*

is Col-0 with the *vsc-7* allele and the L*er SOV* transgene. The *vcs sov* double mutant is Col-0

carrying the *vcs-7* mutation [85]. Within the boxplots, the dashed lines mark the median level of

998   wild type; the number of genes in each category is listed below the lower whisker. The ratios

999   underneath the plots indicate the median of each group normalized to that of wild type. The

1000   statistical significance was determined as described in the Figure 2 legend. (I) Model of mRNA

1001   decay for NMD targets, TuORF mRNAs, and other mRNAs. The orange packman indicates 5'-

1002   to-3' decay machinery, and the hot pink packman indicates 3'-to-5' decay machinery.

1003

SUPPLEMENTARY DATA

Figure S1. Correlations among samples and comparisons between our previous and current datasets

Figure S2. Examples of TuORFs in important regulatory genes.

Figure S3. Cumulative plots for the mRNA levels, translation efficiency, mRNA half-lives and protein abundance of uORF-containing genes

Figure S4. Protein abundance of the uORF-containing genes in Arabidopsis root

Figure S5. mRNA levels of uORF-containing genes in different ecotypes, growth stages, and tissues in Arabidopsis and tomato

Figure S6. mRNA levels of CPuORF-containing genes compared to those of no-uORF genes

Figure S7. (A) TuORF translation levels are positively correlated with the mRNA levels. (B) The distributions of the mRNA levels of TuORF and UuORF genes in Arabidopsis seedlings. Vertical dashed lines indicate the median value of each group.

Figure S8. Lengths of the uORF peptides for NMD target and non-NMD target TuORFs

Figure S9. Lengths of UTRs and CDS for NMD targets and TuORF genes

Figure S10. The mRNA decay rates of NMD targets and TuORF-containing mRNAs in different genetic backgrounds

Table S1. Translated ORFs identified by RiboTaper

- **Table S1A-C.** Translated uORFs identified in the current dataset and our previous Arabidopsis shoot and root datasets

- **Table S1D-F.** Translated mORFs identified in the current dataset and our previous Arabidopsis shoot and root datasets

Table S2. GO-term analysis of TuORF genes

Table S3. Summary of the large-scale datasets used in this study

Table S4. NMD target gene list

- **Table S4A.** High-confidence NMD targets expressed in our data

1030   • **Table S4B.** 49 genes identified as high-confidence NMD target TuORF genes

SUPPLEMENTARY FIGURES

A

B

Figure S1. Correlations among samples and comparisons between previous and current RNA-seq and Ribo-seq datasets. (A) Pearson correlations among three biological replicates of our current RNA-seq and Ribo-seq analyses of samples harvested after 20 or 60 min of DMSO treatment (these are control samples of our large-scale experiment). TPM values in the CDSs of annotated protein-coding genes were used. (B) Higher Ribo-seq read coverage was obtained in

both the uORF and mORF regions in our current dataset; *RGA1* is shown as an example. Note

that the mRNA levels are similar between the previous data (Hsu et al. 2016 [1], shoot) and the

current data. Data representations are the same as those described in the Figure 1F legend.

Figure S2. TuORFs in known important regulatory genes

A. *TOC1*

C. *ARF2*

E. *ARF8* (continued)

uORF4

AT5G37020

F. *Phytochrome B*

AT2G18790

G. *Phytochrome E*

AT4G18130

H. *PIF3*

AT1G09530

Figure S2. Examples of TuORFs in important regulatory genes. Data are presented as

described in the Figure 1F legend. Isoform numbers are shown to the left of the gene models,

and <u>the isoform being considered is bolded</u>. For genes that have multiple TuORFs identified (B,

E, and I), the 3-nt periodicity of each TuORF is presented in a separate panel.

Figure S3. Cumulative plots for the mRNA levels (A), translation efficiency (B), mRNA half-lives (C) and protein abundance (D) of TuORF, UuORF and no-uORF genes.

Figure S4. Protein abundance of the uORF-containing genes in Arabidopsis roots. The

protein quantification data from Arabidopsis seedling roots were extracted from [2]. Within the

boxplot, the dashed line marks the median level of no-uORF genes; the number of genes in

each category is listed below the lower whisker. The ratios underneath the plots indicate the

median of each group normalized to that of no-uORF genes. The statistical significance was

determined as described in the Figure 2 legend.

A

B

C

Figure S5. mRNA levels of uORF-containing genes in different tissues, developmental stages and ecotypes in Arabidopsis and tomato.

TuORF, UuORF, and no-uORF genes were compared.

(A) Arabidopsis roots and aerial and floral parts from different ecotypes.

(B) Various developmental stages of tomato fruit (Heinz 1706 cultivar).

(C) Roots, shoots, flowers, and fruits of tomato (Heinz 1706 cultivar).

The TPM values from RNA-seq quantification were extracted from the EMBO-EBI expression atlas [3] (see METHODS).

Figure S6

Figure S6. mRNA levels of CPuORF-containing genes compared with no-uORF genes.

Within the boxplot, the dashed line marks the median level of no-uORF genes; the number of genes in each category is listed below the lower whisker. The ratios underneath the plots indicate the median of each group normalized to that of no-uORF genes. The statistical significance was determined as described in the Figure 2 legend.

Figure S7. (A) TuORF translation levels are positively correlated with the mRNA levels. TuORF

normalized read count is TuORF Ribo-seq count divided by TuORF length. (B) The distributions

of the mRNA levels of TuORF and UuORF genes. Vertical dashed lines indicate the median

value of each group.

Figure S8. Distributions of uORF peptide lengths in NMD target and non-NMD target TuORF

genes. Vertical dashed lines indicate the median value of each group.

Figure S9. Lengths of UTRs and CDSs for NMD targets and TuORF mRNAs.

Lengths of the 5' UTRs (A), CDSs (B), 3' UTRs (C), and entire transcripts (D) of NMD targets

(NMDt) without a TuORF, NMD targets with a TuORF, non-NMD TuORF mRNAs, and other

mRNAs. Within the boxplots, the dashed lines mark the median level of other mRNAs; the

number of genes in each category is listed below the lower whisker. The ratios underneath the

plots indicate the median of each group normalized to that of other mRNAs. The statistical

significance was determined as described in the Figure 2 legend.

Figure S10. The mRNA decay rates of NMD targets and TuORF-containing mRNAs in different genetic backgrounds. Cumulative plots (A-D) and boxplots (E-H) showing the mRNA decay rates for NMD targets and TuORF mRNAs in wild type and mutants defective in either 5' to 3' decay (*vcs-7*), 3' to 5' decay (*sov*), or both (*vcs sov).* The mRNA decay rates were extracted from [4]. Within the boxplots, the dashed lines mark the median level of other mRNAs; the number of genes in each category is listed below the lower whisker. The ratios underneath the plots indicate the median of each group normalized to that of other mRNAs. The statistical significance was determined as described in the Figure 2 legend.

Reference:

1. Hsu PY, Calviello L, Wu H-YL, Li F-W, Rothfels CJ, Ohler U, et al. Super-resolution ribosome profiling reveals unannotated translation events in Arabidopsis. Proceedings of the National Academy of Sciences of the United States of America. 2016;113:E7126–35.

2. Song G, Hsu PY, Walley JW. Assessment and Refinement of Sample Preparation Methods for Deep and Quantitative Plant Proteome Profiling. Proteomics. 2018;18:e1800220.

3. Papatheodorou I, Moreno P, Manning J, Fuentes AMP, George N, Fexova S, et al. Expression Atlas update: From tissues to single cells. Nucleic Acids Research. 2020;48:D77–83.

4. Sorenson RS, Deshotel MJ, Johnson K, Adler FR, Sieburth LE. Arabidopsis mRNA decay landscape arises from specialized RNA decay substrates, decapping-mediated feedback, and redundancy. Proceedings of the National Academy of Sciences of the United States of America. 2018;115:E1485–94.