Context, competition, and symbiont-induced bet-hedging between *Dictyostelium*

discoideum* and *Paraburkholderia

Trey J. Scott*

David C. Queller

Joan E. Strassmann

Department of Biology, Washington University in St. Louis

*Corresponding Author

E-mail: tjscott@wustl.edu

21 **Keywords: symbiosis, context-dependence, competition, bet-hedging, *D. discoideum*,**

22 ***Paraburkholderia***

Abstract

Symbiotic interactions change with environmental context. We investigated

context-dependence and bet-hedging in the symbiosis between social amoeba hosts and

Paraburkholderia bacteria, where the context is the abundance of host food bacteria.

Paraburkholderia have been shown to harm hosts dispersed to food-rich environments,

but aid hosts dispersed to food-poor environments by allowing hosts to carry food

bacteria. Through measuring symbiont density and host spore production, we show that

this food context matters in three other ways. First, it matters for symbionts, who suffer a

greater cost from competition with food bacteria in the food-rich context. Second, it

matters for host-symbiont conflict, changing how symbiont density negatively impacts

host spore production. Third, data-based simulations show in some cases this context-

dependence can lead to a symbiont-induced bet-hedging advantage for hosts. These

results show how food context can have many consequences for the *Dictyostelium*-

Paraburkholderia symbiosis and suggest that symbionts can induce bet-hedging in hosts.

Introduction

Context-dependence, where the environment can change the sign or magnitude of one partner's effect on the other, is common in symbioses (Bronstein 1994; Thompson 1994; Chamberlain *et al.* 2014). These context-dependent effects on partners can be crucial to understanding the nature of symbiotic interactions (Keeling & McCutcheon 2017; Iwai 2019). For example, in the symbioses between *Paramecium bursaria* hosts and their *Chlorella* endosymbionts, hosts benefitted from symbiosis in light environments, but were harmed in the dark. For *Chlorella*, the effects of symbiosis were negative in co-culture, indicating that hosts exploited their endosymbionts for the benefits hosts receive in light conditions (Lowe *et al.* 2016). However, in the context of an environment with a *Chlorella* competitor, hosts benefited their symbionts by eating these competitors (Iwai 2019). This example illustrates that understanding how partners affect each other across multiple contexts can change our view of the interaction, sometimes from one of exploitation to one of mutual benefit.

Measures of interaction effects in individual contexts may miss beneficial interactions in symbioses if benefits accrue across contexts at the expense of benefits in any individual context. These benefits that occur across contexts are an example of bet-hedging, where organisms adapt to uncertain environments by avoiding the worst effects of harsh contexts while being suboptimal in more favorable contexts (Slatkin 1974; Philippi & Seger 1989; Starrfelt & Kokko 2012). This reduces the variance in fitness across contexts (increasing geometric mean fitness) at the expense of the conventional, arithmetic mean fitness (Seger & Brockmann 1987). For example, in *Lobelia inflata,* limiting the timing of reproduction may reduce the variance in fitness from unpredictable

season lengths while being suboptimal in any individual year (Simons & Johnston 2003).

Bet-hedging is thus defined as an adaptation that involves a trade-off between geometric

mean fitness and arithmetic mean fitness.

Bet-hedging is commonly non-social, but joining together with others may be a

powerful way to hedge one's bets against harsh conditions. Cooperative breeding in birds

(Rubenstein 2011) and plant-fungus mutualisms (Lekberg & Koide 2014) have now been

suggested as bet-hedging phenotypes. Symbiont-induced bet-hedging may be common

because symbionts often give hosts novel traits that allow hosts to gain access to new,

sometimes harsh, environments (Moran 2007; Oliver *et al.* 2010; DiSalvo *et al.* 2015).

For example, endosymbiotic bacteria in aphids play important roles in whether hosts can

withstand heat (Montllor *et al.* 2002) or survive on different host plants (Tsuchida 2004).

These novel traits may be more likely to involve bet-hedging if symbionts induce a cost

in other contexts so geometric mean fitness is increased at the expense of arithmetic mean

fitness. However, symbiont-induced bet-hedging in context-dependent symbioses

remains relatively unexplored.

Context-dependence is important in the lifecycle of the social amoeba

Dictyostelium discoideum. Amoebae need edible bacteria to grow and proliferate (Raper

1937), but the abundance (Young 2004; Vos *et al.* 2013) and quality (Kuserk 1980;

Brock *et al.* 2018) of food bacteria in the soil is known to vary at smaller scales. This

results in a patchy environment where some patches are food-rich and other patches are

food-poor. In response to starvation, amoebae aggregate and form a multicellular fruiting

body to disperse resistant spores to new environments (smith *et al.* 2014). The patchy soil

environment is considered an important selection pressure for this fruiting body structure

(Bonner 1982; Kessin 2001).

D. discoideum interacts with three species of mostly inedible *Paraburkholderia*

bacterial symbionts — *P. agricolaris*, *P. hayleyella*, and *P. bonniae* (Brock *et al.* 2020).

Throughout this paper, we will use "*Paraburkholderia*" or "symbionts" as shorthand for

the three symbiotic *Paraburkholderia* species. Hosts infected with *Paraburkholderia*

have been isolated from multiple locations in the United States with around 25% of

screened hosts being infected by at least one species (Haselkorn *et al.* 2019).

Paraburkholderia are able to enter and live inside *D. discoideum* cells and spores, but can

also proliferate, albeit sometimes only slowly, without their hosts (DiSalvo *et al.* 2015;

Shu *et al.* 2018a; Brock *et al.* 2020) unlike the obligate endosymbionts that are also found

in *D. discoideum* (Haselkorn *et al.* 2021). There is some evidence consistent with

coevolution between hosts and symbionts (Brock *et al.* 2016; Shu *et al.* 2018a; Garcia *et*

al. 2019; Brock *et al.* 2020). For example, host clones naturally infected with *P.*

hayleyella are harmed less by infection with this symbiont than host clones that were not

infected in the wild (Shu *et al.* 2018a) indicating that *P. hayleyella* hosts have adaptations

favoring symbiosis. Symbionts also have the ability to move towards hosts (Shu *et al.*

2018b), suggesting that being able to find hosts is beneficial.

The symbiosis with *Paraburkholderia* bacteria impacts the growth and

proliferation of *D. discoideum*. Having symbionts allows hosts to carry food bacteria

inside the spore-containing part of fruiting bodies called the sorus (DiSalvo *et al.* 2015).

Whether this novel trait is advantageous or not depends on the presence of food bacteria

after dispersal. When food is abundant, having symbionts can be costly, as shown by

infected amoebae producing fewer spores than uninfected amoebae (Brock *et al.* 2011;

DiSalvo *et al.* 2015). In food-poor environments, the cost of having *Paraburkholderia* is

compensated by hosts gaining the ability to carry food bacteria (in addition to inedible

Paraburkholderia) in dispersing spores. This allows amoebae to disperse and grow where

they ordinarily could not (Brock *et al.* 2011; DiSalvo *et al.* 2015). These context-

dependent effects on the host could be extremely important in the natural soil

environment, where food-poor patches arise frequently (Kessin 2001).

Less is known about how symbionts are affected across food contexts. Gaining

the ability to disperse to new locations may be a major reason for symbionts to seek out

social amoeba hosts (Garcia & Gerardo 2014), but could also make the context of host

food bacteria in the new environment important for symbionts. A benefit from being

dispersed to patches with few bacteria could be that symbionts face reduced competition.

If few bacteria are present, symbionts will mostly compete with food bacteria that were

also carried in the sorus. This should be a relatively low competition situation because

symbionts outnumber food bacteria in sori (Khojandi *et al.* 2019). Having few

competitors should advantage symbionts while environments with plentiful bacteria

could strongly limit symbiont growth because of their relatively slow growth rates, at

least as measured in the lab (Brock *et al.* 2020). We will use "food-rich" and "food-poor"

to describe newly colonized patches with many and few bacteria, respectively, of the sort

edible by *D. discoideum*. These categories reflect the relationship to *D. discoideum* and

could be called high and low competition in terms of their effect for *Paraburkholderia*.

It is also unclear how the number of extracellular *Paraburkholderia* in the

environment impacts hosts since previous studies have focused on intracellular

Paraburkholderia (Shu *et al.* 2018b; Miller *et al.* 2020). When they are outside the

amoebae, *Paraburkholderia* could affect *D. discoideum* fitness through interactions with

food bacteria perhaps by reducing the amount of food for hosts through competition or by

releasing diffusible toxins that affect amoebae. Thus host food context could also affect

the relationship between symbiont density and host spore production.

It is also not known whether symbionts can induce a bet-hedging phenotype in

hosts. This phenotype would give hosts the ability to evade the harsh effects of food-poor

contexts at the expense of reduced fitness in food-rich contexts. This could increase host

success across contexts (measured by geometric mean fitness) by reducing the variance in

fitness across contexts, but will appear maladaptive according to measures of arithmetic

mean fitness (Seger & Brockmann 1987; Starrfelt & Kokko 2012). Infected hosts could

alternatively gain an outright advantage from carrying symbionts without a trade-off

between arithmetic and geometric mean fitness.

To understand context-dependence in the symbiosis between amoebae and

Paraburkholderia, we used *D. discoideum* infected with either *P. agricolaris* or *P.*

hayleyella — the two most common and best-studied species of *D. discoideum*

symbionts. We investigate whether *Paraburkholderia* benefit from reduced interspecific

competition when dispersed to food-poor contexts, how symbiont density and food

context impact host spore production, and whether symbionts induce a bet-hedging

phenotype in their hosts when food conditions vary across soil patches.

Methods

To understand the effects of symbiosis across food contexts, we used 4 naturally

uninfected *D. discoideum* clones, 4 clones naturally infected with *P. agricolaris*, and 4

clones naturally infected with *P. hayleyella* (Figure 1). We cured the infected clones and

re-infected them with their native symbionts to equalize infection density. Uninfected

clones were left uninfected, but were otherwise treated the same as infected clones. This

resulted in three host infection conditions: uninfected, infected with *P. agricolaris*, and

infected with *P. hayleyella*. To mimic natural dispersal, we collected sori and transferred

them to food-rich (with additional *K. pneumoniae* bacteria) or food-poor (KK2 buffer

with no *K. pneumoniae*) nutrient plates. Bacteria appear on food-poor plates only if

transferred sori contain bacteria, as expected for infected samples. We grew replicate

experimental sets involving all conditions beginning on two separate dates, July 13 and

22, 2020, and followed up with additional experiments (see results) beginning on January

26 and April 23, 2021.

Paraburkholderia isolation

To isolate *Paraburkholderia* from their hosts, we grew wild collected *D.*

discoideum clones on SM/5 plates (2 g glucose (Fisher Scientific), 2 g Bacto Peptone

(Oxoid), 2 g yeast extract (Oxoid), 0.2 g $MgSO_4 * 7H_2O$ (Fisher Scientific), 1.9 g

KH_2PO_4 (Sigma-Aldrich), 1 g K_2HPO_4 (Fisher Scientific), and 15 g agar (Fisher

Scientific) per liter). Wild *D. discoideum* clones were grown with *K. pneumoniae* food

bacteria that were suspended in KK2 buffer (2.25 g KH_2PO_4 (Sigma-Aldrich) and 0.67 g

K_2HPO_4 (Fisher Scientific) per liter). After wild clones completed the social cycle

(feeding, starvation, and fruiting body formation), we collected sori with pipette tips and

placed them on SM/5 plates. We allowed the bacteria and amoebae contained within to

proliferate and then streaked out the resulting bacteria to get single colonies.

Paraburkholderia removal

To generate uninfected clones, we treated infected *D. discoideum* clones with

antibiotics by plating on 30 μg/mL tetracycline SM/5 plates with 200 μL of 1.5 optical

density (OD_{600}) tetracycline-resistant *K. pneumoniae* suspended in KK2 buffer. After

passage on SM/5 plates without tetracycline to let the amoebae recover from any effects

of the antibiotic, we collected single sori with a pipette tip and placed ten of them in

different locations on SM/5 plates to confirm that we had successfully removed the

bacteria. If bacteria are present, these spot tests will show bacterial growth and

Dictyostelium proliferation as the spores hatch and eat the bacteria (Brock *et al.* 2011).

Without bacteria, amoebae cannot proliferate and the spot will stay blank. We considered

a clone to be cured if no bacteria showed up in spot tests. We similarly treated naturally

uninfected hosts with tetracycline to control for any effect of curing on our results.

Paraburkholderia re-infection

We re-infected cured *D. discoideum* clones with their native *Paraburkholderia*

isolates by plating 200 μL $2x10^5$ spores with 200 μL of 0.1% *Paraburkholderia* solution.

This solution consisted of 1.5 OD_{600} *Paraburkholderia* and 1.5 OD_{600} *K. pneumoniae* in a

1:1000 ratio. To confirm re-infection (and also successful isolation), we performed spot

tests as above, where successful re-infection was inferred when bacteria grew on 8 or

more spots out of the 10 we put down.

Artificial dispersal to food-rich and food-poor plates

To obtain sori to transfer to food-rich and food-poor plates, we started by growing

D. discoideum clones from frozen stock, as described above, on 200 μL 1.5 OD *K.*

pneumoniae expressing green fluorescent protein (GFP). We obtained GFP-expressing *K.*

pneumoniae (strain ID DBS0349837) from the Dicty Stock Center at dictyBase (Fey *et*

al. 2013). This initial growth period is to remove freezer effects and ensure that food

bacteria that are carried to new plates are GFP-expressing since stocks were fed non-GFP

bacteria before freezing. After six days of growth, we used pipette tips to collect sori

from mature fruiting bodies. We counted spores using a hemocytometer and diluted

spores to a concentration of $2x10^5$ per mL, and then plated them on plates with (food-

rich) or without (food-poor) an additional 200 μL of the GFP-expressing food bacterium

K. pneumoniae (Figure 1). In order to survive on food-poor plates, the host must carry

food bacteria from the previous plate. We grew food-rich and food-poor plates for six

days unless otherwise stated, enough time for mature fruiting bodies to form.

Measurement of bacteria density

To measure *Paraburkholderia* density, we measured the quantity of bacteria left

on plates after *D. discoideum* formed fruiting bodies. We first collected plate contents by

washing plates with 15 mL of KK2 buffer. To remove fruiting bodies and bacteria

associated with fruiting bodies, we centrifuged wash solutions for three minutes at 13000

rpm. We measured bacteria using optical density measured at 600 nm (OD_{600}), a

frequency at which bacteria commonly scatter light. Because the OD_{600} is due to both

Paraburkholderia and *K. pneumoniae*, we used GFP fluorescence measurements (with an

excitation wavelength of 485 and emission wavelength of 515 nm) and a standard curve

relating *K. pneumoniae* fluorescence to its OD_{600} to subtract out the component due to

GFP-expressing *K. pneumoniae*. Both OD_{600} and fluorescence measures were performed

in a 96 well plate with a Tecan Infinite 200 Pro microplate reader.

To validate our standard curve, we compared predicted OD_{600} of *P. agricolaris*

and *K. pneumoniae* to colony forming unit (CFU) counts from the same samples. Linear

regression revealed that predicted OD_{600} measurements explained most of the variation in

CFUs, showing that our assay is reliable (Figure S1). We also checked our standard curve

for significant quadratic terms, which can cause measurement errors when combining

OD_{600} and fluorescence measures at high densities (Meyers *et al.* 2018), but our curve did

not have a significant quadratic term.

Host spore production

To measure host spore production, we estimated spore concentration in the

supernatants from washed plates using a hemocytometer. We then calculated the total

number of spores per plate by multiplying by the volume of wash solution.

Spore production simulations

To test whether *Paraburkholderia* symbionts induce a bet-hedging phenotype in

hosts, we simulated repeated bouts of growth and dispersal across soil patches with

different probabilities of having food bacteria. Bet hedging occurs when the best

phenotype in terms of arithmetic mean fitness is not the same as the best phenotype in

terms of geometric mean fitness (Seger & Brockmann 1987). We separately modeled

three host phenotypes: (1) unifected (2) infected with *P. agricolaris*, or (3) infected with

P. hayleyella. Infections with both species at once are possible, but are rare in nature

(Haselkorn *et al.* 2019) so we exclude coinfections from our analysis.

We assumed that environments consisted of *x* discrete soil patches. In each round

of the simulation, patches are randomly assigned as food-rich with probability *p* or food-

poor with probabity 1-*p*. Thus, in simulations with one patch, populations experience

variation in food abundance that is completely temporal. With additional patches,

variation can also occur across space. This will reduce temporal variation because fitness

at a given time will be summed over multiple patches.

Initially all patches are colonized. Each patch produces a number of spores,

drawn from the distribution of our empirical spore production values, according to

whether it is a food-rich or food-poor patch. Infected hosts have been found to have

reduced spore production in food-rich environments that depends on infection density of

Paraburkholderia endosymbionts (Brock *et al.* 2011; DiSalvo *et al.* 2015; Miller *et al.*

2020). We did not detect this cost in our data, perhaps because we infected with fewer

Paraburkholderia. Since these costs appear to vary and could be an important factor

affecting fitness in the soil environment when food abundance varies, we included them

as a variable. To model costs, we penalized host spore production in food rich

environments by multiplying spore production values by $1 - c$, where c is the proportion

of spores lost because of the cost of infection. When c is 0, we model the scenario

observed in this study, with no infection cost. These cost-adjusted spore production

values are summed to get the total spore production across all patches. This is divided by

2×10^5, a rough estimate of the number of spores in a typical sorus, to get the total number

of sori, which we are assuming to be the dispersal unit.

The global pool of sori is used to seed the next round. Patches are assumed to be

empty and are randomly reassigned as food-poor or food-rich. Each sorus is randomly

assigned to a patch and it successfully germinates with probability g (we assume g = 0.5

in the main text; other values (0.1 and 0.9) result in similar outcomes (Figure S2&S3).

We assumed that patches colonized by multiple sori were the same as singly colonized

patches for the purposes of determining their subsequent spore production. Some patches

may remain unfilled.

We let cost of infection c and the frequency of food-poor patches p vary from 0 to

1 in increments of 0.1 and simulated environments with one, 10, and 100 soil patches.

We simulated dispersal to new patches for 100 rounds of growth and dispersal using 500

replicates for each combination of probability of p and c for each phenotype. At the end

of the 100 rounds, we calculated the total spore production per round and calculated

geometric mean spore production from these values. To calculate arithmetic mean spore

production, we sampled our spore production values from food-rich and food-poor plates

according to the number of patches and the value of p. We used t tests to compare

uninfected hosts' spore production to that of either *P. hayleyella* or *P. agricolaris*

infected hosts for a given parameter combination. If there was no significant difference

between infected and uninfected, we categorized results as "no best phenotype". Because

we used many parameter combinations, we corrected p-values for multiple comparisons

using false discovery rate at the level of different numbers of patches and different

infection statuses (at the level of individual grids in Figure 4).

Statistical Methods

We performed statistics in R version 3.6.3 (R Core Team 2020). To compare

bacteria density and spore production, we used linear mixed models (LMM) with the *lme*

function in the nlme package (Pinheiro & Bates 2006). To account for random variation

from replicate clones and effects of dates when experiments were performed, we included

clone and the date the experiment was performed — along with each variable on its own

— as random effects. To select the best model of random effects, we used AICc, a

sample-size-corrected measure of model fit that balances predictive ability and model

complexity (Burnham & Anderson 2004). Many of our models showed different

variances between treatments. To account for these differences in variance, we weighted

models with the *varIdent* function in nlme (Pinheiro & Bates 2006). We used the

emmeans package (Lenth *et al.* 2018) to perform contrasts.

To understand how *Paraburkholderia* density affects host spore production across

food conditions, we fit a LMM using only infected hosts that included symbiont density

leftover on plates and whether the plate was food-rich or food-poor, along with the

interaction between these variables. We included random effects for clone, date, and both

crossed effects and selected the best random effect structure with AICc. We determined

whether the interaction was important by comparing AICc of the model including the

interaction with models including the other variables but lacking the interaction.

Results:

Paraburkholderia dispersed by Dictyostelium sori have lower growth when host food bacteria are abundant

The context of a food-poor environment is known to be important for *D. discoideum* hosts. It is not known how *Paraburkholderia* are affected by this same context, but reduced competition with food bacteria seems likely. We tested this by growing infected sorus contents on food-poor and food-rich nutrient plates and measuring the density of *Paraburkholderia* after *D. discoideum* fruiting body formation (Figure 1). After infected hosts formed fruiting bodies, *Paraburkholderia* densities were lowest in food-rich conditions (Figure 2A), as expected if they compete with food bacteria. There was around five times more *P. agricolaris* on food-poor than food-rich plates (LMM, $p < 0.001$). *P. hayleyella* growth was higher in food-poor conditions than food-rich, but this difference was not significant after 6 days (LMM, $p = 0.416$). Because *P. hayleyella* grows slowly, we performed two more experiments with *P. hayleyella* with 8 and 12-day growth periods (Figure 2B). Allowing for longer incubations did not result in significantly higher density of *P. hayleyella* (LMM, $p = 0.633$), suggesting that *P. hayleyella* reach their maximum density at or before 6 days, but including these additional experiments gave us enough power to find a significant increase in *P. hayleyella* density in food-poor conditions relative to food-rich (LMM, $p = 0.027$). These results show that symbiont density is context-dependent.

Higher symbiont density harms hosts, but less so in food-poor contexts

The host-food context may affect the relationship between symbiont density and

host spore production and therefore the degree of conflict or cooperation between them.

To investigate this, we also measured total host spore production from plates where we

measured the growth of *Paraburkholderia* symbionts (Figure 1). We used uninfected

hosts as a baseline for fitness without symbionts. We confirmed prior studies (Brock *et*

al. 2011; DiSalvo *et al.* 2015) showing that infected hosts could carry food bacteria and

proliferate on food-poor plates, while uninfected host could not (Figure 3A).

Surprisingly, we did not observe a cost of being infected in food-rich conditions ($p > 0.5$

for both species) which has been seen in previous studies (Brock *et al.* 2011; DiSalvo *et*

al. 2015; Shu *et al.* 2018a). This is likely a result of our lower infection dosage of 0.1%.

While having some symbionts is essential for hosts to be able to carry food and

survive in food-poor conditions, higher symbiont densities may nevertheless harm hosts,

perhaps in ways that depend on food context. We found that larger populations of

symbionts as measured by OD_{600} were associated with lower host spore production, but

this harm was reduced in food-poor conditions. Lower host spore production was

associated with being in a food-poor environment ($\beta_{food-poor} = -3.283$, se = 0.853) and

symbiont density ($\beta_{density} = -10.317$, se = 6.364), but the interaction between food scarcity

and symbiont density showed that the harmful effect of higher symbiont densities was

lessened on food-poor plates ($\beta_{food-poor*density} = 8.078$, se = 6.381; Figure 3B). These

results indicate that symbiont density may come at the expense of host spore production,

but that this cost decreases in food-poor environments.

Symbionts can induce a bet-hedging phenotype in hosts

Because *Paraburkholderia* helps hosts in food-poor contexts, we hypothesized

that they could give hosts a bet-hedging advantage that increases geometric mean fitness

at the expense of arithmetic mean fitness across spatial and temporal variation in food

abundance in the soil. We modeled this by using our empirical spore production values to

simulate 100 rounds of growth and dispersal across environments with one, 10, and 100

soil patches. We modeled three symbiosis phenotypes: (1) uninfected, (2) infected with

P. agricolaris, and (3) infected with *P. hayleyella* (Figure 4A, Animations S1-3) in their

own separate environments and then compared their geometric and arithmetic mean spore

productions.

Soil patches were either food-rich or food-poor and changed randomly in each

round. By changing the number of patches, we are able to understand how host fitness is

affected when variation in food abundance is more temporal (few soil patches) or spatial

(many soil patches). We varied the frequency of food-poor soil patches and calculated a

phenotype's success as the geometric or arithmetic mean spore production. We included

the cost of infection observed in other studies (DiSalvo *et al.* 2015; Shu *et al.* 2018a;

Haselkorn *et al.* 2019; Miller *et al.* 2020) by subtracting some proportion of spores from

our empirical measures for infected hosts in food-rich conditions. We classified a

phenotype as a bet-hedger if the best phenotype in terms of geometric mean fitness

differed from the best phenotype in terms of arithmetic mean fitness (Figure 4B).

We found that uninfected hosts were favored by both geometric and arithmetic

means when costs of infection were high, food was abundant, and food abundance

variation was spatial (Figure 4C&D; pink). Associating with *Paraburkholderia*,

especially *P. hayleyella*, provided an advantage in terms of arithmetic mean fitness across

a wide range of parameters, even with fairly large costs (Figure 4C&D; blue). Symbionts

were most beneficial when variation in food abundance was temporal (one soil patch). As

the number of patches in the environment increased, the benefits of having symbionts

decreased, with *P. agricolaris* being beneficial only when the probability of food was

high. This lowered benefit of *P. agricolaris* is especially apparent when fewer potential

dispersers manage to colonize new patches (Figure S2).

We found that bet-hedging is likely when variation in food abundance is more

temporal (one patch), when food-poor patches are common, and when costs of infection

are high (Figure 4C&D; green). This bet-hedging advantage in more temporally varying

environments is because uninfected populations go extinct when generations without

food occur (as illustrated in Figure 4A with a single soil patch, Animation S1). Bet-

hedging becomes rarer as the number of patches increases (middle and rightmost panels

in Figure 4C&D) and the worst consequences of food-poor patches (extinction) become

rare due to the presence of some patches with food. Bet-hedging occurs across a broader

range of parameters in *P. agricolaris* simulations (Figure 4C). These results show that

symbionts can provide an outright benefit for hosts in many conditions and can induce a

bet-hedging phenotype in some conditions.

Discussion

Our results show how the context of host food abundance affects the

Dictyostelium-Paraburkholderia symbiosis beyond the previously demonstrated

advantage to hosts when food is rare (Brock *et al.* 2011). We found evidence that both

Paraburkholderia species benefit from reduced competition when they are carried to

food-poor environments (Figure 2). Symbiont density negatively affected host spore

production, but symbionts harmed hosts less in food-poor conditions (Figure 3B).

Symbionts gave hosts an outright or a bet-hedging advantage across a range of

parameters in patchy soils (Figure 4).

Our finding that symbionts benefit from reduced competition when dispersed to

food-poor environments fits in with other findings of competitive benefits for symbionts

(Iwai 2019). Competition between symbionts and food bacteria may also be the cause of

context-dependent effects of symbiont density on host spore production. Our spore

production results showed that higher symbiont densities resulted in lower host spore

production, indicating that symbionts are harmful to hosts. However, higher symbiont

densities are less harmful in food-poor conditions when competition is lower (Figure 3B).

The reduced harm for hosts could be the result of less antagonism between bacteria,

which results in less collateral damage to amoebae through secreted toxins or other

competitive interactions between food bacteria and symbionts. The generality of our

results is limited somewhat by only using one species of food bacteria. While using a

single food bacterium is more experimentally tractable, amoebae encounter multiple

bacteria species in their natural environments (Brock *et al.* 2018). Different species, or

combinations of species, could change competition with symbionts and affect host spore

production in different ways.

Paraburkholderia symbionts benefit hosts by giving hosts the ability to carry food

to food-poor contexts (Brock *et al.* 2011; DiSalvo *et al.* 2015). Using simulations, we

found that this benefit was prevalent across a range of costs, food probabilities, and

number of soil patches according to measures of arithmetic mean spore production

(Figure 4). Hosts could also benefit across contexts through bet-hedging, where

geometric mean fitness trades off with arithmetic mean fitness (Seger & Brockmann

1987). Bet-hedging occurred more often in our simulations when costs imposed by the

symbiont were relatively high and food abundance varied temporally (few soil patches).

Temporal variation disadvantaged uninfected amoebae because they went extinct in food-

poor patches while infected amoebae weathered these harsh contexts. These results are

interesting when paired with the finding that infected hosts constitute about 25% of the

population (Haselkorn *et al.* 2019). This may mean that symbiosis is not universally

favored — if so, *Paraburkholderia* infection would be fixed in the population — but

instead depends on ecological conditions.

Our simulations were necessarily simplified to make them tractable. Two notable

limitations from this are that soil patches are discrete and equally sized and that host

spore production as measured in the lab may not accurately reflect nature. Though not

unique to our model (Starrfelt & Kokko 2012), our finding that bet hedging is most likely

in temporally varying environments also limits the scope of our results. However, such

temporal variation may result naturally in the soil from rainfall variation and has been

shown to impact soil bacteria abundance (Barnard *et al.* 2013). Though it remains

uncertain how much *Paraburkholderia* induces host bet-hedging in the wild, our model

illustrates the potential for a symbiotic partner to induce bet-hedging.

Our simulation results suggest that bet-hedging may be an important, but

underappreciated, consequence of some symbioses. Bet-hedging has been suggested in

some symbioses (Lekberg & Koide 2014), but has not been tested in detail. Though our

simulations were tailored to the symbiosis between *D. discoideum* and *Paraburkholderia*,

they should apply to other cases of context-dependent symbiosis with costs and temporal

variation. Since context-dependence is common in symbioses (Bronstein 1994), we

expect that bet-hedging that involves symbiosis may be relatively common. Accounting

for geometric mean fitness in symbiotic interactions will be crucial to determining the

true extent of bet-hedging in symbioses.

Bet-hedging may be especially relevant in symbioses where hosts gain novel

phenotypes from symbionts (Moran 2007; Oliver *et al.* 2010) because they can allow

symbionts to survive in new, previously harsh, contexts (Montllor *et al.* 2002; Tsuchida

2004). However, it is important to note that bet-hedging will not evolve unless these

harsh contexts occur probabilistically. If the new harsh environment is constant,

symbiosis will be favored without any bet-hedging. If the harsh environment is too

predictable, plasticity will be favored instead of bet-hedging (Botero *et al.* 2015). Thus an

understanding of the ecological contexts will also be important for identifying symbiont-

induced bet-hedging.

Acknowledgements: This material is based upon work funded by the National Science

Foundation under grants DEB-1753743 and IOS-1656756. We thank Tyler Larsen and

James Medina for comments on the manuscript. We would also like to thank the

Strassmann/Queller lab group, especially Debbie Brock, for feedback during the

development and execution of this project.

Authorship statement: TS, DQ, and JS designed the study and wrote the manuscript. TS

performed the experiments and simulations and analyzed the data.

Data Statement: Data will be archived at Dryad upon acceptance. Data and code for

simulations and analysis will be available at www.gitlab.com/treyjscott/farmerBH.

References

Barnard, R.L., Osborne, C.A. & Firestone, M.K. (2013). Responses of soil bacterial and
fungal communities to extreme desiccation and rewetting. *ISME J.*, 7, 2229–
2241.
Bonner, J.T. (1982). Evolutionary Strategies and Developmental Constraints in the
Cellular Slime Molds. *Am. Nat.*, 119, 530–552.
Botero, C.A., Weissing, F.J., Wright, J. & Rubenstein, D.R. (2015). Evolutionary tipping
points in the capacity to adapt to environmental change. *Proc. Natl. Acad. Sci.*,
112, 184–189.
Brock, D.A., Douglas, T.E., Queller, D.C. & Strassmann, J.E. (2011). Primitive
agriculture in a social amoeba. *Nature*, 469, 393–396.
Brock, D.A., Haselkorn, T.S., Garcia, J.R., Bashir, U., Douglas, T.E., Galloway, J., *et al.*
(2018). Diversity of Free-Living Environmental Bacteria and Their
Interactions With a Bactivorous Amoeba. *Front. Cell. Infect. Microbiol.*, 8.
Brock, D.A., Jones, K., Queller, D.C. & Strassmann, J.E. (2016). Which phenotypic
traits of Dictyostelium discoideum farmers are conferred by their bacterial
symbionts? *Symbiosis*, 68, 39–48.
Brock, D.A., Noh, S., Hubert, A.N.M., Haselkorn, T.S., DiSalvo, S., Suess, M.K., *et al.*
(2020). Endosymbiotic adaptations in three new bacterial species associated
with *Dictyostelium discoideum* : *Paraburkholderia agricolaris* sp. nov.,
Paraburkholderia hayleyella sp. nov., and *Paraburkholderia bonniea* sp. nov.
PeerJ, 8, e9151.
Bronstein, J.L. (1994). Conditional outcomes in mutualistic interactions. *Trends Ecol.*
Evol., 9, 214–217.
Burnham, K.P. & Anderson, D.R. (2004). Multimodel Inference: Understanding AIC
and BIC in Model Selection. *Sociol. Methods Res.*, 33, 261–304.
Chamberlain, S.A., Bronstein, J.L. & Rudgers, J.A. (2014). How context dependent are
species interactions? *Ecol. Lett.*, 17, 881–890.
DiSalvo, S., Haselkorn, T.S., Bashir, U., Jimenez, D., Brock, D.A., Queller, D.C., *et al.*
(2015). *Burkholderia* bacteria infectiously induce the proto-farming
symbiosis of *Dictyostelium* amoebae and food bacteria. *Proc. Natl. Acad. Sci.*,
112, E5029–E5037.
Fey, P., Dodson, R.J., Basu, S. & Chisholm, R.L. (2013). One stop shop for everything
Dictyostelium: dictyBase and the Dicty Stock Center in 2012. In:
Dictyostelium Discoideum Protocols. Springer, pp. 59–92.
Garcia, J.R. & Gerardo, N.M. (2014). The symbiont side of symbiosis: do microbes
really benefit? *Front. Microbiol.*, 5.

Garcia, J.R., Larsen, T.J., Queller, D.C. & Strassmann, J.E. (2019). Fitness costs and
benefits vary for two facultative *Burkholderia* symbionts of the social
amoeba, *Dictyostelium discoideum*. *Ecol. Evol.*, ece3.5529.
Haselkorn, T.S., DiSalvo, S., Miller, J.W., Bashir, U., Brock, D.A., Queller, D.C., *et al.*
(2019). The specificity of *Burkholderia* symbionts in the social amoeba
farming symbiosis: Prevalence, species, genetic and phenotypic diversity.
Mol. Ecol., 28, 847–862.
Haselkorn, T.S., Jimenez, D., Bashir, U., Sallinger, E., Queller, D.C., Strassmann, J.E., *et*
al. (2021). Novel Chlamydiae and *Amoebophilus* endosymbionts are prevalent
in wild isolates of the model social amoeba *Dictyostelium discoideum*. *Environ.*
Microbiol. Rep., 1758-2229.12985.
Iwai, S. (2019). Photosynthetic Endosymbionts Benefit from Host's Phagotrophy,
Including Predation on Potential Competitors. *Curr. Biol.*, 29, 3114–3119.
Keeling, P.J. & McCutcheon, J.P. (2017). Endosymbiosis: The feeling is not mutual. *J.*
Theor. Biol., 434, 75–79.
Kessin, R.H. (2001). *Dictyostelium: evolution, cell biology, and the development of*
multicellularity. Cambridge University Press.
Khojandi, N., Haselkorn, T.S., Eschbach, M.N., Naser, R.A. & DiSalvo, S. (2019).
Intracellular Burkholderia Symbionts induce extracellular secondary
infections; driving diverse host outcomes that vary by genotype and
environment. *ISME J.*
Kuserk, F.T. (1980). The Relationship Between Cellular Slime Molds and Bacteria in
Forest Soil. *Ecology*, 61, 1474–1485.
Lekberg, Y. & Koide, R.T. (2014). Integrating physiological, community, and
evolutionary perspectives on the arbuscular mycorrhizal symbiosis. *Botany*,
92, 241–251.
Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. (2018). Emmeans:
Estimated marginal means, aka least-squares means. *R Package Version*, 1, 3.
Lowe, C.D., Minter, E.J., Cameron, D.D. & Brockhurst, M.A. (2016). Shining a Light on
Exploitative Host Control in a Photosynthetic Endosymbiosis. *Curr. Biol.*, 26,
207–211.
Meyers, A., Furtmann, C. & Jose, J. (2018). Direct optical density determination of
bacterial cultures in microplates for high-throughput screening applications.
Enzyme Microb. Technol., 118, 1–5.
Miller, J.W., Bocke, C.R., Tresslar, A.R., Schniepp, E.M. & DiSalvo, S. (2020).
Paraburkholderia Symbionts Display Variable Infection Patterns That Are
Not Predictive of Amoeba Host Outcomes. *Genes*, 11, 674.
Montllor, C.B., Maxmen, A. & Purcell, A.H. (2002). Facultative bacterial
endosymbionts benefit pea aphids Acyrthosiphon pisum under heat stress.
Ecol. Entomol., 27, 189–195.
Moran, N.A. (2007). Symbiosis as an adaptive process and source of phenotypic
complexity. *Proc. Natl. Acad. Sci.*, 104, 8627–8633.
Oliver, K.M., Degnan, P.H., Burke, G.R. & Moran, N.A. (2010). Facultative Symbionts in
Aphids and the Horizontal Transfer of Ecologically Important Traits. *Annu.*
Rev. Entomol., 55, 247–266.

Philippi, T. & Seger, J. (1989). Hedging one's evolutionary bets, revisited. *Trends*
567   *Ecol. Evol.*, 4, 41–44.
568   Pinheiro, J. & Bates, D. (2006). *Mixed-effects models in S and S-PLUS*. Springer Science
569   & Business Media.
570   R Core Team. (2020). *R: A language and environment for statistical computing*. R
571   Foundation for Statistical Computing, Vienna, Austria.
572   Raper, K.B. (1937). Growth and development of Dictyostelium discoideum with
573   different bacterial associates. *J. Agric. Res.*, 55, 289–316.
574   Rubenstein, D.R. (2011). Spatiotemporal environmental variation, risk aversion, and
575   the evolution of cooperative breeding as a bet-hedging strategy. *Proc. Natl.*
576   *Acad. Sci.*, 108, 10816–10822.
577   Seger, J. & Brockmann, H. (1987). What is bet-hedging? In: *Oxford surveys in*
578   *evolutionary biology*. Oxford University Press.
579   Shu, L., Brock, D.A., Geist, K.S., Miller, J.W., Queller, D.C., Strassmann, J.E., *et al.*
580   (2018a). Symbiont location, host fitness, and possible coadaptation in a
581   symbiosis between social amoebae and bacteria. *eLife*, 7, 25.
582   Shu, L., Zhang, B., Queller, D.C. & Strassmann, J.E. (2018b). Burkholderia bacteria use
583   chemotaxis to find social amoeba Dictyostelium discoideum hosts. *ISME J.*, 12,
584   1977–1993.
585   Simons, A.M. & Johnston, M.O. (2003). Suboptimal timing of reproduction in Lobelia
586   inflata may be a conservative bet-hedging strategy. *J EVOL BIOL*, 11.
587   Slatkin, M. (1974). Hedging one's evolutionary bets. *Nature*, 250, 704–705.
588   smith, jeff, Queller, D.C. & Strassmann, J.E. (2014). Fruiting bodies of the social
589   amoeba Dictyostelium discoideum increase spore transport by Drosophila.
590   *BMC Evol. Biol.*, 14, 105.
591   Starrfelt, J. & Kokko, H. (2012). Bet-hedging-a triple trade-off between means,
592   variances and correlations. *Biol. Rev.*, 87, 742–755.
593   Thompson, J.N. (1994). *The coevolutionary process*. University of Chicago Press.
594   Tsuchida, T. (2004). Host Plant Specialization Governed by Facultative Symbiont.
595   *Science*, 303, 1989–1989.
596   Vos, M., Wolf, A.B., Jennings, S.J. & Kowalchuk, G.A. (2013). Micro-scale determinants
597   of bacterial diversity in soil. *FEMS Microbiol. Rev.*, 37, 936–954.
598   Young, I.M. (2004). Interactions and Self-Organization in the Soil-Microbe Complex.
599   *Science*, 304, 1634–1637.

Figure Legends

Figure 1: Schematic of experimental design. (A) Uninfected and infected *D. discoideum*

fruiting bodies are collected and plated on food-rich and food-poor plates (after one

passage on GFP-expressing *K. pneumoniae* food bacteria). These plates are grown for six

days and then washed for bacterial measurement and spore counting (B). Bacteria are

measured by calculating GFP fluorescence and optical density (see Methods).

Figure 2: More *Paraburkholderia* were recovered from plates after fruiting body

formation from food-poor plates (those that had not received additional *K. pneumoniae*).

(A) *Paraburkholderia* density after 6 days. (B) *P. hayleyella* density after 8 and 12 days.

Point shapes show individual clones 4 of each (see Figure 1).

Figure 3: Effects of *Paraburkholderia* infection and density on host spore production.

(A) Spore production of hosts from food-rich and food-poor plates for uninfected, *P. agricolaris* infected, and *P. hayleyella* infected hosts. (B) Interaction between measured *Paraburkholderia* density (OD_{600}) and food environment on host spore production. This interaction model explained 95% of the variance in spore production. Inset shows food-rich results on smaller scale. Point shapes show individual clones (see Figure 1).

Figure 4: Benefits of associating with *Paraburkholderia* symbionts depends on variation

in food availability, fitness costs, and number of soil patches. (A) Toy example of

simulations with five rounds of growth and dispersal where spores are dispersed to a

single soil patch each generation. Lack of food could lead to the extinction of uninfected

clones. (B) Illustration of bet-hedging trade-off between geometric and arithmetic mean

spore production in a single soil patch. Bet-hedging occurs when the food-poor

probability is between 0.1 and 0.7 (note how uninfected hosts have the highest arithmetic

mean while infected hosts have the highest geometric mean). Infected hosts suffered a

50% reduction in spore production in these simulations. Performance of *P. agricolaris*

(C) and *P. hayleyella* (D) relative to uninfected for 1, 10, and 100 soil patches. Pink

shows when uninfected hosts have higher arithmetic and geometric mean spore

production; blue shows when infected hosts have higher arithmetic and geometric mean

spore production; green shows when arithmetic fitness is reduced for higher geometric

mean fitness (bet-hedging).

 Supplemental Material

Figure S1: Linear regression results relating estimated optical density (predicted from

standard curve) to colony counts from serial dilutions of the same *P. agricolaris* (A) and

K. pneumoniae (B) samples.

670 **Figure S2:** Plot of the best phenotype for a given probability of being in a no food

671 environment and cost of having symbionts. These are the same as Figure 4C and D with

672 10% of sori successfully germinating.

Figure S3: Plot of the best phenotype for a given probability of being in a no food environment and cost of having symbionts. These are the same as Figure 4C-H with 90% of sori successfully germinating.

Animation S1: Example simulation with 100 rounds of growth and dispersal with one soil patch, a 50% cost of infection, and a food-poor frequency of 0.3.

Animation S2: Example simulation with 100 rounds of growth and dispersal with 10 soil patches, a 50% cost of infection, and a food-poor frequency of 0.3.

Animation S3: Example simulation with 100 rounds of growth and dispersal with 100 soil patches, a 50% cost of infection, and a food-poor frequency of 0.3.