Genetically diverse mouse models of SARS-CoV-2 infection model clinical variation and

cytokine responses in COVID-19.

Shelly J Robertson[1], Olivia Bedard[2], Kristin L. McNally[1], Matthew Lewis[1], Chad Clancy[3], Carl

Shaia[3], Rebecca M. Broeckel[1], Abhilash I. Chiramel[1], Gail L. Sturdevant[1], Elvira Forte[2],

Christoph Preuss[2], Candice N. Baker[2], Daniel E. Sturdevant[4], Craig Martens[4], Steven M.

Holland[5], Nadia A. Rosenthal[2,6]* and Sonja M. Best[1]*

[1]Laboratory of Virology, Rocky Mountain Laboratories, National Institute of Allergy and

Infectious Diseases, NIH, Hamilton MT USA

[2]The Jackson Laboratory, Bar Harbor Maine USA

[3]Rocky Mountain Veterinary Branch, Rocky Mountain Laboratories, National Institute of

Allergy and Infectious Diseases, NIH, Hamilton MT USA

[4]Research Technologies Branch, Rocky Mountain Laboratories, National Institute of Allergy and

Infectious Diseases, NIH, Hamilton MT USA

[5]Division of Intramural Research, National Institute of Allergy and Infectious Diseases, NIH,

Bethesda MD USA

[6]National Heart and Lung Institute, Imperial College London

*co-corresponding authors

Abstract

Host genetics are a significant determinant of coronavirus disease 2019 (COVID-19)[1]. Animal

models that reflect genetic diversity and a range of clinical outcomes observed in human

populations are needed to understand mechanisms of severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) infection dynamics and disease[2]. Here, we report a mouse panel

comprising the diverse genetic backgrounds of the Collaborative Cross (CC) founder strains

crossed to C57BL/6J mice expressing the K18-hACE2 transgene[3] that enables infection by

SARS-CoV-2. Infection of CCxK18-hACE2 F1 progeny resulted in a spectrum of weight loss,

survival, viral replication kinetics, histopathology, and cytokine profiles, some of which were

sex-specific. Importantly, survival was closely associated with early type I interferon expression

and a phased proinflammatory response distinct from mice with severe disease. Thus, dynamics

of inflammatory responses observed in COVID-19 can be modeled in diverse mice that provide a

genetically tractable platform for understanding antiviral immunity and evaluating

countermeasures.

Main

Mice humanized for the main cellular receptor for SARS-CoV-2 entry, angiotensin converting

enzyme 2 (ACE2), are valuable pre-clinical infection models[2]. The most widely used model,

K18-hACE2, supports high early virus replication in lung epithelial cells, resulting in

inflammatory cell infiltration, interstitial edema and focal consolidation of the lung with rapid

lethality[4-9]. However, this model on a standard C57BL/6J genetic background does not reflect the

wide variety of COVID-19 outcomes in humans, from asymptomatic to severe[10], that are

dependent on genetic polymorphisms, age, sex and the presence of underlying conditions[11-13].

Multiple studies have demonstrated the value of using genetically diverse mice to assess the

impact of host genetics on infectious disease outcomes [14-18]. Indeed, a single genetic background

likely will not capture the range of responses and can fail to model key features of disease[19].

To determine if incorporating genetic diversity could recapitulate the broad phenotypic variation

observed in human COVID-19, we crossed the K18-hACE2 strain to the genetically distinct

inbred strains A/J, 129S1J, NOD, NZO, PWK, CAST, and WSB that along with C57BL/6J

comprise the founders of the Collaborative Cross (CC). This reference population represents

90% of the genetic diversity in *M. musculus* populations[20,21] and are used to understand genetic

regulation of complex immune responses. Notably, NOD and NZO mice are commonly used in

studies of diabetes and metabolic dysfunction that are factors contributing to COVID-19

severity[22]. In addition to the 8 CC founder strains, BALB/c and DBA strains were included

because BALB/c is widely used in infectious disease modelling, and DBA is a founder strain for

BXD strains also used in genetic mapping[23].

CCxK18-hACE2 F1 mice were infected via intranasal inoculation with 10^3 pfu SARS-CoV-2

strain nCoV-WA1-2020 (MN985325.1) and mice were monitored for 21 days. This virus dose is

sub-uniformly lethal in the prototype K18-hACE2 strain (C57BL/6J background), resulting in

80-90% of mice reaching end-point criteria by 7 days post infection (dpi) (Fig.1). Infection was

confirmed by seroconversion to SARS-CoV-2 nucleoprotein in all survivors. Comparative

analysis of the eight CCxK18-hACE2 F1 cohorts documented a broad spectrum of weight loss

and survival curves characteristic to each CC founder genotype, some of which were sex-specific

(Fig. 1). Highly sensitive mice strains included C57BL/6J and A/JxK18-hACE2 that lost

approximately 20% of their starting weight between 4 and 7 dpi with no clear sex bias. In

contrast, CCxK18-hACE2 F1 progeny from PWK, NZO, 129S1/J (Fig. 1), BALB/c and DBA

(Fig. S1) were resistant to clinical disease, generally losing 5-10% starting weight with ~80% of

mice surviving infection. Finally, CCxK18-hACE2 F1 progeny of three strains (CAST, NOD,

WSB) had marked sexual dimorphism in response, with CAST and NOD F1 males susceptible to

lethal disease, although differences in weight loss between sexes were not apparent. Yet another

phenotype was obtained in SARS-CoV-2 infected WSB F1 progeny, where infection was

uniformly lethal in females. Weight loss in WSB F1 males began later than other groups at 6 dpi

and surviving males exhibited sustained weight loss until the end of the observation period in

contrast to the rapid weight gain associated with recovery of other CCxK18-hACE2 F1 cohorts.

In K18-hACE2 mice, SARS-CoV-2 titers peak in the lung at 2-3 dpi and high titers of virus can

be isolated from the CNS which may contribute to lethality in this model[24,25]. Thus, virus

replication kinetics were determined at 3 and 6 dpi in lung and brain of CCxK18-hACE2 F1

cohorts (Fig. 2A, 2B). F1 progeny of sensitive founder strains C57BL/6J and A/J showed high

levels of infectious SARS-CoV-2 at 3dpi in lung homogenates that was reduced by 1-2 $\log_{10}$ by 6

dpi with no differences between sexes. Infectious virus was generally not recovered from the

CNS at 3dpi, but 50% of C57BL/6J and A/J F1 progeny had high virus burden in the CNS by

6dpi (10^6-10^8 PFU/g tissue). In contrast, peak lung virus titer in F1 progeny of the most resistant

founder strain, PWK, was 150-fold lower than the sensitive strains at 3 dpi and was below the

limit of detection at 6dpi. Control of virus replication in PWK F1s was also evident in the CNS

where detection of infectious virus was sporadic. Interestingly, F1 progeny of additional CC

founder strains classed as resistant had equivalent peak viral titers in the lung to the sensitive

K18-hACE2 and A/J F1 progeny, but they tended to control replication by 6 dpi to levels $1\log_{10}$

lower than sensitive mice and had less virus burden in the CNS. Finally, lung titers in (CAST,

NOD and WSB)xK18-hACE2 F1 progeny were not different between males and females despite

differences in clinical severity associated with sex. Thus, F1 progeny of CCxK18-hACE2 mouse

strains can be further stratified into a) sensitive F1 progeny with high sustained virus replication

in lung and CNS (C57BL/6J, A/J), b) resistant F1 progeny associated with lower peak virus titer

and earlier control of replication in the lung with no or low dissemination to other organs (PWK,

NZO, 129S1/J), and c) F1 progeny with sex bias where resistance is independent of virus titer in

the lung suggesting a sex-based difference in host response (CAST, NOD, WSB). Relative

expression of the K18-hACE2 transgene in lung suggested some variability, but expression was

not associated with clinical phenotypes or peak viral burden (Fig. S2). Together, these data

suggest that the range of clinical severity in mice is only partially associated with control of virus

replication (Summarized in Table 1).

Pathological changes in SARS-CoV-2-infected K18-hACE2 mice were similar to those

previously described[7-9,26]. Inflammatory infiltrates evident by 3dpi included perivascular

lymphocytes with alveolar septa thickened by neutrophils, macrophages and edema. At 6 dpi,

pulmonary pathology was multifocal, and consistent with interstitial pneumonia including type II

pneumocyte hyperplasia, septal, alveolar and perivascular inflammation comprised of

lymphocytes, macrophages and neutrophils, with alveolar fibrin and edema evident. Bronchiolar

pathology was not observed in these mice. Pathology was classified as none, mild (rare scattered

inflammatory foci), moderate (coalescing inflammatory foci) or severe (widespread, large

inflammatory foci) as exemplified in Fig. 2c. The classification of 'severe' is a comparative term

for these mice, as we generally did not observe pathology equivalent to that of the more severe

Syrian hamster model of SARS-CoV-2 infection[27-29]. Surprisingly, in comparison to F1 progeny

of sensitive founder strains (K18-hACE2 and A/JxK18-hACE2), F1 progeny of resistant PWK,

female NZO and 129S/J mice, as well as those of NOD and WSB tended to have higher

pathology scores in lungs (Fig. 2d). Lung distribution of viral RNA by RNAscope was limited to

type I and II pneumocytes in all strains (Fig. 2b). Pathology in the CNS was scored as present or

absent, as it generally consisted of subtle inflammatory foci, including perivascular cuffing and

increased gliosis associated with necrotic cells observed in K18-hACE2 and DBAxK18-hACE2

F1 males (Fig. S3A-C, G). However, pathology in the CNS of CASTxK18-hACE2 F1 progeny

was striking in that microthrombi were evident in capillaries with extensive hemorrhage in the

absence of encephalitis, and was associated with viral RNA distribution in the same area (Fig.

S3D-F).

In-depth longitudinal analyses of immune responses in patients with COVID-19 have defined

key immunological correlates of disease outcome[30-43]. Moderate COVID-19 is associated with

increased plasma levels of a core inflammatory signature of pro-inflammatory cytokines,

chemokines and growth factors (e.g., IL-1α, IL-1β, IFNα, IL-12 p70, and IL-17A) that are

effectively resolved[43]. In contrast, severe COVID-19 includes sustained expression of these

markers with additional signatures including IL-6, IL-10, IL-18, IL-23, TNF-α, and eotaxin

among others suggesting that specific timing and failure to resolve inflammatory responses are

important factors in disease progression[43]. In SARS-CoV-2-infected CCxK18-hACE2 F1 mice,

cytokines and chemokines were quantified in the serum and bronchoalveolar lavage fluid (BAL)

by multiplex analysis at 3 and 6 dpi. In general, cytokine levels were much higher in BAL fluid

than in serum (Fig. S4A, B; individual data points are provided in Figs. S5-8). Strikingly in the

BAL, comparison of sensitive (K18-hACE2 and A/JxK18-hACE2) versus resistant (PWK, NZO

and 129S1)xK18-hACE2 F1 progeny revealed that high IFNα expression at 3dpi was associated

with survival in both males and females (Fig. 3A, B). Resistant mice also expressed

proinflammatory cytokine IL-6, Th1 cytokines (IL-12p70, IL-27) and chemokines (CXCL10,

Gro-α/KC, CCL2, CCL3, CCL4, CCL5, CCL7 and CXCL2) (designated as Group A) in BAL at

this early timepoint (Fig. 3B, Cluster 1). At 6 dpi, many Group A cytokines were resolving (Fig.

3C) while IFN-γ, TNF-α, IL-18, IL-1β, IL-2, IL-4, IL-5, and IL-13 (designated as Group B)

were added to the cytokine signature in resistant mice (Fig. 3B, Cluster 2). Interestingly,

resistant PWK and NZO x K18-hACE2 mice exhibited a unique profile characterized by

tempered levels of Group A and Group B along with high production of IFNγ, Gro-α/KC and

eotaxin at 6dpi, a timepoint coincident with the most efficient clearance of virus from the lung

(Cluster 4). In contrast, sensitive strains generally failed to produce Group A cytokines at 3 dpi

(Cluster 3) and increases in production of Group A and Group B mediators in BAL occurred by

6 dpi (Fig. 3B, Cluster 2). Cytokine signatures were less clear in F1 progeny with sex bias

(CAST, NOD and WSB) likely due to intermediate survival phenotypes observed in one or both

sexes, although responses in BAL were generally higher in males (Fig. S4C). Together, these

genetically diverse mice model major dynamic phenotypes observed in human COVID-19.

Specifically, reduced disease severity in mice is associated with early type I IFN expression and

a phased, controlled inflammatory response. In contrast, a delayed and unorchestrated innate

response is associated with lethality.

Taken together, these observations demonstrate that use of host genetic variation in mice can

model different outcomes of SARS-CoV-2 infection, addressing a major deficiency in the toolkit

required to combat COVID-19. The phenotypes in diverse mice support a role for the innate

immune system in determining COVID-19 severity[32,44-47] and can be used to address key

knowledge gaps, including mechanisms of innate immune control of virus replication, defining

events needed for a well-orchestrated inflammatory response independent of early viral burden,

molecular mechanisms of sex-dependent disease severity, and longer-term implications for tissue

repair[48] and lung function.

Acknowledgement

This work was funded by the Division of Intramural Research, National Institutes of Health,

National Institute of Allergy and Infectious Diseases. Thank you to the animal caretaker staff at

campuses of both RML and The Jackson Laboratory for their work in animal husbandry.

Declaration of Interests:

The authors have no interests to declare.

METHODS

Ethics statement

Animal study protocols were reviewed and approved by the Institutional Animal Care and Use Committee

(IACUC) at Rocky Mountain Laboratories (RML), NIAID, NIH in accordance with the recommendations in

the Guide for the Care and Use of Laboratory Animals of the NIH. All animal experiments were performed

in an animal biosafety level 3 (ABSL3) research facility at RML. Standard operating procedures for work

with infectious SARS-CoV-2 and protocols for virus inactivation were approved by the Institutional

Biosafety Committee (IBC) and performed under BSL3 conditions.

Virus preparation

SARS-CoV-2 (USA_WA1/2020) from University of Texas Medical Branch (Vero passage 4) was

propagated on Vero cells cultured in DMEM supplemented with 10% fetal bovine serum, 1 mM L-

glutamine, 50U/ml penicillin and 50ug/ml streptomycin. Culture supernatants were collected at 72 hpi,

aliquoted and stored at -80ºC.

Mice and virus infection

CC founder x C57BL/6J -K18-hACE2 F1 were provided by The Jackson Laboratories and include the

following: B6.Cg-Tg (K18-ACE2)2Prlmn/J), 034860; (A/J x B6.Cg-Tg(K18-ACE2)2Prlmn/J)F1/J, 035940;

(PWK/PhJ x B6.Cg-Tg(K18-ACE2)2Prlmn/J)F1/J, 035938; (NZO/HlLtJ x B6.Cg-Tg(K18-

ACE2)2Prlmn/J)F1/J, 035936; (129S1/SvlmJ x B6.Cg-Tg(K18-ACE2)2Prlmn/J)F1/J, 035934; (CAST/EiJ x

B6.Cg-Tg(K18-ACE2)2Prlmn/J)F1/J, 035937; (NOD/ShiLtJ x B6.Cg-Tg(K18-ACE2)2Prlmn/J)F1/J,

035935; (WSB/EiJ x B6.Cg-Tg(K18-ACE2)2Prlmn/J)F1/J, 035939; (BALB/cJ x B6.Cg-Tg(K18-

ACE2)2Prlmn/J)F1/J, 035941; (BALB/cJ x B6.Cg-Tg(K18-ACE2)2Prlmn/J)F1/J 035943. Six to 12 week-

old male and female mice were inoculated by the intranasal route with 10^3 pfu of SARS-CoV-2 in a

volume of 50ul PBS (Gibco). Prior to inoculation mice were anesthetized by inhalation of isoflurane.

Mice were monitored daily for clinical signs of disease and weight loss.

Virus titration

Infectious virus in lung and brain tissue was quantified by plaque assay. Tissues were collected in 0.5ml

of DMEM containing 2% FBS, 50U/ml penicillin and 50ug/ml streptomycin and immediately frozen.

Tissue samples were weighed, and then homogenized using 5mm steel beads and TissueLyzer II high-

speed shaker (Qiagen). Ten-fold serial dilutions of homogenates were prepared in duplicate and used to

inoculate Vero cells grown in 48-well tissue culture plates. Following 1 hour incubation at 37ºC, the cells

were overlayed with 1.5% carboxymethyl cellulose (CMC) in MEM and incubated at 37ºC for 3-4 days.

Cells were then fixed in 10%formalin and plaques were visualized by staining with 1% crystal violet

diluted in 10% ethanol.

Measurement of SARS-CoV-2-specific IgG

Sera were collected at 21 dpi from mice that survived SARS-CoV-2 infection. SARS-CoV-2 spike protein-

specific IgG was measured using SARS-CoV2 spike protein serological IgG ELISA kit (Cell Signaling

Technology) per manufacturer's instructions.

Multiplex cytokine/chemokine analysis

Sera were collected from SARS-CoV-2-infected mice at 3 and 6 dpi by centrifugation of whole blood in

GelZ serum separation tubes (Sarstedt). BAL samples were recovered by insufflation of lungs with 1 ml

sterile PBS followed by aspiration to collect ~0.5ml volume of fluid. SARS-CoV-2 in sera and BAL fluid

was inactivated by using γ- irradiation (2 MRad) and removed from the BSL3 laboratory. Cytokine

concentrations in serum and BAL were measured using a 26-Procartaplex mouse cytokine/chemokine

panel (ThermoFisher EPX260-26088-901) combined with a Simplex IFNα assay (ThermoFisher EPX01A-

26027-901). Samples were run on a Luminex Bio-Plex 200 system with BioPlex Manager 6.1.1 software.

Histopathology and *in situ* hybridization

Tissues were fixed in 10% neutral buffered formalin for a minimum of 7 days with 2 changes according to

IBC-approved standard operating procedure. Tissues were processed with a Sakura VIP-6 Tissue Tek,

on a 12-hour automated schedule, using a graded series of ethanol, xylene, and PureAffin. Embedded

tissues were sectioned at 5 µm and dried overnight at 42°C prior to staining with hematoxylin and

eosin. Chromogenic detection of SARS-CoV-2 viral RNA was performed using the RNAscope VS

Universal AP assay (Advanced Cell Diagnostics Inc.) on the Ventana Discovery ULTRA stainer using a

SARS-CoV-2 specific probe (Advanced Cell Diagnostics Inc. cat#848569). *In situ* hybridization was

performed according to manufacturer's instructions.

Quantitative real-time PCR

Total lung RNA was extracted from homogenized lysates using the *mir*Vana miRNA Isolation Kit

(Invitrogen cat. AM1560) following the manufacturer's protocol for total RNA isolation. Total RNA samples

were quantified using a Nanodrop spectrophotometer (Thermo cat. ND-2000C), and reverse transcribed

to cDNA using the SuperScript IV VILO Master Mix with ezDNase enzyme (Invitrogen cat. 11766050)

following the manufacturer's instructions. Expression of mouse *Ace2*, mouse *Gapdh*, and human ACE2

were quantified by qPCR using the ViiA7 Real-Time PCR System (Thermo cat. 44535545). TaqMan

probes targeting mouse *Ace2* (Thermo probe ID Mm01159006_m1), mouse *Gapdh* (Thermo probe ID

Mm99999915_g1), and human ACE2 (Thermo probe ID Hs01085333_m1) were used in conjunction with

the TaqMan Fast Advanced Master Mix (Thermo cat. 4444556) and the ViiA7 Real-Time PCR System

with QuantStudio Software (Thermo cat. 44535545). Samples were run in technical triplicate in 10ul

reaction volumes on a 384-well plate, and m*Ace2* and hACE2 transcript abundance values were

normalized to m*Gapdh* in each sample and relative expression was calculated using a common reference

lung sample (C57BL/6J male) and the delta delta-Ct method.

Statistical analysis

Comparison of survival curves in males versus females for each strain was performed using the log-rank

(Mantel-Cox) test. One-way analysis of variance (ANOVA) with Tukey's multiple comparison posttest was

used to compare cytokine and chemokine levels of BAL and serum. Differences between groups were

considered significant at a *P* value of <0.05. All statistical analyses were performed with graphPad Prism

8.0 (GraphPad Software) or Qlucore Omics Explorer Version 3.7 (Qlucore AB).

Figure Legends:

Figure 1: CCxK18-hACE2 F1 progeny mice demonstrate a range of clinical severity

following infection with SARS-CoV-2. Mice were infected intranasally with 10^3PFU of SARS-

CoV-2 and monitored for survival and weight loss. F1 founder strains were classified as: **a,**

sensitive with 50% or greater mortality and no difference between males and females, **b,** resistant

with 35% or less mortality, or **c,** sensitive with a sex bias as having statistically significant

difference between sensitivity of male and female mice. Mice without evidence of

seroconversion at the end of the study were removed from analysis (N=6-8/per sex).

Figure 2: Virus replication kinetics and pathology in CCxK18-hACE2 F1 progeny. a, Virus

titers (PFU/g tissue) in lung and brain of male and female mice were measured at 3 and 6 dpi. **b,**

RNAscope of SARS-CoV-2 genome predominantly localizing to pneumocytes in areas of lung

inflammation. Serial sections demonstrate pathology (upper) and viral RNA distribution (lower;

Red). Scale bars represent 100μm. **c,** Examples of pathology scoring used in this study. Scale

bars represent 500μm. **d,** Lung pathology scores for CC Founder F1 males and females at 3 and

6 dpi. N = 6/time point/sex. ND, not determined.

Figure 3: Timing of type I IFN and inflammatory cytokine responses in the lung is

associated with clinical severity in CCxK18-hACE2 F1 progeny. a, IFNα protein in BAL

from resistant and sensitive mice. **b,** cytokine and chemokine responses in BAL at 3 and 6 dpi. **c,**

Resolving group A cytokines in mice resistant to lethality, but not sensitive mice.

Supplemental Figure 1: Survival, weight loss and SARS-CoV-2 replication kinetics in

BALB/cxK18-hACE2 or DBAxK18-hACE2 F1 progeny. a, Survival and weight loss in

BALB/cxK18-hACE2 F1. **b,** Survival and weight loss in DBA x K18-hACE2 F1. **c,** Virus titers

(PFU/g tissue) in lung and brain of male and female mice were measured at 3 and 6 dpi. **d,**

Cytokine expression in BAL.

Supplemental Figure 2: Relative expression of human *ACE2* in CCxK18-hACE2 F1

progeny. Gene expression was measured by real-time RT-PCR and expressed relative to that in

K18-hACE2 mice.

Supplemental Figure 3: Examples of pathology in the brain of SARS-CoV-2-infected

CCxK18-hACE2 F1 progeny. a, Small focal areas of inflammation were most commonly

observed (circled). Scale bar = 1000μm. **b,** increased magnification of a. demonstrating

perivascular cuffing, gliosis, and evidence of necrotic cells in areas of focal inflammation. Scale

bar = 200μm. **c,** Example of a pronounced lymphocytic inflammatory response with perivascular

cuffing and increased cellularity (arrowheads). Scale bar = 100μm. **d,** Microthrombus occluding

cerebral capillaries and hemorrhage (arrow) observed in CASTxK18-hACE2. Scale bar =

100μm. **e,** Example of extensive hemorrhage in the brain of CASTxK18-hACE2 F1 progeny

associated with extensive virus replication as demonstrated in f. Scale bar = 1000μm. **f,** serial

section relative to e. showing extensive SARS-CoV-2 genome by RNAscope. Scale bar =

1000μm. **g,** Incidence of encephalitis for CC Founder F1 males and females at 3 and 6 dpi. ND,

not determined.

Supplemental Figure 4: Interferon, cytokine and chemokine expression in the serum

compared to the BAL of SARS-CoV-2-infected female CCxK18-hACE2 F1 progeny.

Cytokines were measured at 3 and 6 dpi by bioplex. **a,** 3 dpi, and **b,** 6dpi for sensitive and

resistant strains with no sex-bias. **c,** 3 and 6 dpi from strains with a sex-bias.

Supplemental Figure 5: Interferon, cytokine and chemokine expression in the

bronchoalveolar lavage from SARS-CoV-2-infected male CCxK18-hACE2 F1 progeny.

Cytokines were measured at 3 and 6 dpi by bioplex. The average cytokine expression in BAL of

naïve mice is indicated by a grey dashed line.

Supplemental Figure 6: Interferon, cytokine and chemokine expression in the serum of

SARS-CoV-2-infected male CCxK18-hACE2 F1 progeny. Cytokines were measured at 3 and

6 dpi by bioplex. The average cytokine expression in serum of naïve mice is indicated by a grey

dashed line.

Supplemental Figure 7: Interferon, cytokine and chemokine expression in the

bronchoalveolar lavage from SARS-CoV-2-infected female CCxK18-hACE2 F1 progeny.

Cytokines were measured at 3 and 6 dpi by bioplex. The average cytokine expression in BAL of

naïve mice is indicated by a grey dashed line.

Supplemental Figure 8: Interferon, cytokine and chemokine expression in the serum of

SARS-CoV-2-infected female CCxK18-hACE2 F1 progeny. Cytokines were measured at 3

and 6 dpi by bioplex. The average cytokine expression in serum of naïve mice is indicated by a

grey dashed line.

References:

1   Severe Covid, G. G. *et al.* Genomewide Association Study of Severe Covid-19 with
349   Respiratory Failure. *N Engl J Med* **383**, 1522-1534, doi:10.1056/NEJMoa2020283 (2020).
350   2   Munoz-Fontela, C. *et al.* Animal models for COVID-19. *Nature* **586**, 509-515,
351   doi:10.1038/s41586-020-2787-6 (2020).
352   3   McCray, P. B., Jr. *et al.* Lethal infection of K18-hACE2 mice infected with severe acute
353   respiratory syndrome coronavirus. *J Virol* **81**, 813-821, doi:10.1128/JVI.02012-06 (2007).
354   4   Chen, R. E. *et al.* In vivo monoclonal antibody efficacy against SARS-CoV-2 variant
355   strains. *Nature* **596**, 103-108, doi:10.1038/s41586-021-03720-y (2021).
356   5   Johnson, B. A. *et al.* Loss of furin cleavage site attenuates SARS-CoV-2 pathogenesis.
357   *Nature* **591**, 293-299, doi:10.1038/s41586-021-03237-4 (2021).
358   6   VanBlargan, L. *et al.* A potently neutralizing anti-SARS-CoV-2 antibody inhibits variants of
359   concern by binding a highly conserved epitope. *bioRxiv*, doi:10.1101/2021.04.26.441501
360   (2021).
361   7   Winkler, E. S. *et al.* SARS-CoV-2 infection of human ACE2-transgenic mice causes severe
362   lung inflammation and impaired function. *Nat Immunol* **21**, 1327-1335,
363   doi:10.1038/s41590-020-0778-2 (2020).
364   8   Yinda, C. K. *et al.* K18-hACE2 mice develop respiratory disease resembling severe COVID-
365   19. *PLoS Pathog* **17**, e1009195, doi:10.1371/journal.ppat.1009195 (2021).
366   9   Zheng, J. *et al.* COVID-19 treatments and pathogenesis including anosmia in K18-hACE2
367   mice. *Nature* **589**, 603-607, doi:10.1038/s41586-020-2943-z (2021).
368   10   Long, Q. X. *et al.* Clinical and immunological assessment of asymptomatic SARS-CoV-2
369   infections. *Nat Med* **26**, 1200-1204, doi:10.1038/s41591-020-0965-6 (2020).
370   11   Docherty, A. B. *et al.* Features of 20 133 UK patients in hospital with covid-19 using the
371   ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study.
372   *BMJ* **369**, m1985, doi:10.1136/bmj.m1985 (2020).

12   Richardson, S. *et al.* Presenting Characteristics, Comorbidities, and Outcomes Among
5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* **323**, 2052-
2059, doi:10.1001/jama.2020.6775 (2020).
13   Pairo-Castineira, E. *et al.* Genetic mechanisms of critical illness in COVID-19. *Nature* **591**,
92-98, doi:10.1038/s41586-020-03065-y (2021).
14   Graham, J. B. *et al.* Genetic diversity in the collaborative cross model recapitulates
human West Nile virus disease outcomes. *mBio* **6**, e00493-00415,
doi:10.1128/mBio.00493-15 (2015).
15   Gralinski, L. E. *et al.* Genome Wide Identification of SARS-CoV Susceptibility Loci Using
the Collaborative Cross. *PLoS Genet* **11**, e1005504, doi:10.1371/journal.pgen.1005504
(2015).
16   Martin, M. D., Sompallae, R., Winborn, C. S., Harty, J. T. & Badovinac, V. P. Diverse CD8 T
Cell Responses to Viral Infection Revealed by the Collaborative Cross. *Cell Rep* **31**,
107508, doi:10.1016/j.celrep.2020.03.072 (2020).
17   Noll, K. E. *et al.* Complex Genetic Architecture Underlies Regulation of Influenza-A-Virus-
Specific Antibody Responses in the Collaborative Cross. *Cell Rep* **31**, 107587,
doi:10.1016/j.celrep.2020.107587 (2020).
18   Price, A. *et al.* Transcriptional Correlates of Tolerance and Lethality in Mice Predict Ebola
Virus Disease Patient Outcomes. *Cell Rep* **30**, 1702-1713 e1706,
doi:10.1016/j.celrep.2020.01.026 (2020).
19   Sarkar, S. & Heise, M. T. Mouse Models as Resources for Studying Infectious Diseases.
Clin Ther **41**, 1912-1922, doi:10.1016/j.clinthera.2019.08.010 (2019).
20   Roberts, A., Pardo-Manuel de Villena, F., Wang, W., McMillan, L. & Threadgill, D. W. The
polymorphism architecture of mouse genetic resources elucidated using genome-wide
resequencing data: implications for QTL discovery and systems genetics. *Mamm*
Genome **18**, 473-481, doi:10.1007/s00335-007-9045-1 (2007).
21   Srivastava, A. *et al.* Genomes of the Mouse Collaborative Cross. *Genetics* **206**, 537-556,
doi:10.1534/genetics.116.198838 (2017).
22   Williamson, E. J. *et al.* Factors associated with COVID-19-related death using
OpenSAFELY. *Nature* **584**, 430-436, doi:10.1038/s41586-020-2521-4 (2020).
23   Ashbrook, D. G. *et al.* A platform for experimental precision medicine: The extended
BXD mouse family. *Cell Syst* **12**, 235-247 e239, doi:10.1016/j.cels.2020.12.002 (2021).
24   Gan, E. S. *et al.* A mouse model of lethal respiratory dysfunction for SARS-CoV-2
infection. *Antiviral Res* **193**, 105138, doi:10.1016/j.antiviral.2021.105138 (2021).
25   Kumari, P. *et al.* Neuroinvasion and Encephalitis Following Intranasal Inoculation of
SARS-CoV-2 in K18-hACE2 Mice. *Viruses* **13**, doi:10.3390/v13010132 (2021).
26   Oladunni, F. S. *et al.* Lethality of SARS-CoV-2 infection in K18 human angiotensin-
converting enzyme 2 transgenic mice. *Nat Commun* **11**, 6122, doi:10.1038/s41467-020-
19891-7 (2020).
27   Francis, M. E. *et al.* SARS-CoV-2 infection in the Syrian hamster model causes
inflammation as well as type I interferon dysregulation in both respiratory and non-
respiratory tissues including the heart and kidney. *PLoS Pathog* **17**, e1009705,
doi:10.1371/journal.ppat.1009705 (2021).

28   Imai, M. *et al.* Syrian hamsters as a small animal model for SARS-CoV-2 infection and
countermeasure development. *Proc Natl Acad Sci U S A* **117**, 16587-16595,
doi:10.1073/pnas.2009799117 (2020).
29   Tostanoski, L. H. *et al.* Ad26 vaccine protects against SARS-CoV-2 severe clinical disease
in hamsters. *Nat Med* **26**, 1694-1700, doi:10.1038/s41591-020-1070-6 (2020).
30   Bernardes, J. P. *et al.* Longitudinal Multi-omics Analyses Identify Responses of
Megakaryocytes, Erythroid Cells, and Plasmablasts as Hallmarks of Severe COVID-19.
Immunity **53**, 1296-1314 e1299, doi:10.1016/j.immuni.2020.11.017 (2020).
31   Schulte-Schrepping, J. *et al.* Severe COVID-19 Is Marked by a Dysregulated Myeloid Cell
Compartment. *Cell* **182**, 1419-1440 e1423, doi:10.1016/j.cell.2020.08.001 (2020).
32   Schultze, J. L. & Aschenbrenner, A. C. COVID-19 and the human innate immune system.
Cell **184**, 1671-1692, doi:10.1016/j.cell.2021.02.029 (2021).
33   Arunachalam, P. S. *et al.* Systems biological assessment of immunity to mild versus
severe COVID-19 infection in humans. *Science* **369**, 1210-1220,
doi:10.1126/science.abc6261 (2020).
34   Bastard, P. *et al.* Autoantibodies against type I IFNs in patients with life-threatening
COVID-19. *Science* **370**, doi:10.1126/science.abd4585 (2020).
35   Bost, P. *et al.* Host-Viral Infection Maps Reveal Signatures of Severe COVID-19 Patients.
Cell **181**, 1475-1488 e1412, doi:10.1016/j.cell.2020.05.006 (2020).
36   Chakraborty, S. *et al.* Proinflammatory IgG Fc structures in patients with severe COVID-
19. *Nat Immunol* **22**, 67-73, doi:10.1038/s41590-020-00828-7 (2021).
37   Consiglio, C. R. *et al.* The Immunology of Multisystem Inflammatory Syndrome in
Children with COVID-19. *Cell* **183**, 968-981 e967, doi:10.1016/j.cell.2020.09.016 (2020).
38   Mathew, D. *et al.* Deep immune profiling of COVID-19 patients reveals distinct
immunotypes with therapeutic implications. *Science* **369**, doi:10.1126/science.abc8511
(2020).
39   Shen, B. *et al.* Proteomic and Metabolomic Characterization of COVID-19 Patient Sera.
Cell **182**, 59-72 e15, doi:10.1016/j.cell.2020.05.032 (2020).
40   Silvin, A. *et al.* Elevated Calprotectin and Abnormal Myeloid Cell Subsets Discriminate
Severe from Mild COVID-19. *Cell* **182**, 1401-1418 e1418, doi:10.1016/j.cell.2020.08.002
(2020).
41   Takahashi, T. *et al.* Sex differences in immune responses that underlie COVID-19 disease
outcomes. *Nature* **588**, 315-320, doi:10.1038/s41586-020-2700-3 (2020).
42   Wilk, A. J. *et al.* A single-cell atlas of the peripheral immune response in patients with
severe COVID-19. *Nat Med* **26**, 1070-1076, doi:10.1038/s41591-020-0944-y (2020).
43   Lucas, C. *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19.
Nature **584**, 463-469, doi:10.1038/s41586-020-2588-y (2020).
44   Zhang, Q. *et al.* Inborn errors of type I IFN immunity in patients with life-threatening
COVID-19. *Science* **370**, doi:10.1126/science.abd4570 (2020).
45   Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives Development of
COVID-19. *Cell* **181**, 1036-1045 e1039, doi:10.1016/j.cell.2020.04.026 (2020).
46   Ziegler, C. G. K. *et al.* Impaired local intrinsic immunity to SARS-CoV-2 infection in severe
COVID-19. *Cell* **184**, 4713-4733 e4722, doi:10.1016/j.cell.2021.07.023 (2021).

47　Zanoni, I. Interfering with SARS-CoV-2: are interferons friends or foes in COVID-19? *Curr*
Opin Virol **50**, 119-127, doi:10.1016/j.coviro.2021.08.004 (2021).
48　Delorey, T. M. *et al.* COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular
targets. *Nature* **595**, 107-113, doi:10.1038/s41586-021-03570-8 (2021).
463

464　**Table 1:** Summary of percent survival, weight loss, virus burden and BAL cytokine profiles in

465　K18-hACE2 x CC Founder F1 progeny.

				Peak weight loss		Average Virus Titers in Lung		IFNα in BAL	Cytokine profile BAL	
			Mean time to death			3 dpi	6 dpi			
Strain	Sex	%Survival	(days)	%	DPI	Mean (Std Dev)	Mean (Std Dev)	3dpi	3 dpi	6 dpi
K18-hACE2	M	14	6	16.7	7	1.5e6 (1.9e6)	7.6e4 (1.3e5)	Low	No/Low	High Group A and Group B
	F	29	7	12.0	7	1.5e6 (1.4e6)	8.2e3 (5.7e3)	Low		
A/J x K18-hACE2 F1	M	33	8	15.0	7	2.8e6 (2.2e6)	4.65e4 (6.9e4)	Low	No/Low	High Group A and Group B
	F	66	UD	6.1	7	3.3e6 (2.4e6)	3.0e4 (2.9e4)	Low		
PWK x K18-hACE2 F1	M	83	UD	-	-	1.6e5 (2.3e5)	BLD	High	High Group A	Moderate Group A and Group B
	F	83	UD	7.0	7	1.5e4 (2.3e4)	BLD	Low		
NZO x K18-hACE2 F1	M	86	UD	13.9	9	2.6e6 (2.9e6)	3.8e3 (3.7e3)	High	High Group A	High Group A and B
	F	71	UD	21.9	9	2.0e6 (9.4e5)	1.3e3 (1.0e3)	High		
129S1J x K18-ACE2 F1	M	86	UD	11.8	7	1.1e6 (8.3e5)	3.0e3 (2.8e3)	High	High Group A	High Group A and Group B
	F	71	UD	12.0	7	8.1e5 (6.4e4)	2.4e4 (4.0e4)	Moderate		
CAST x K18-ACE2 F1	M	0	6	17.2	7	3.4e6 (3.2e6)	4.8e5(3.7e5)	Low	Low Group A	Low Group A Males = Females
	F	50	15.5	35.8	8	1.75e6(1.5e6)	3.4e4 (2.9e4)	Low		
NOD x K18-ACE2 F1	M	33	6.5	6.0	6	1.7e6 (3.7e6)	4.8e3 (5.3e3)	Low	No/Low	Group A and Group B Males > Females
	F	83	UD	-	-	4.3e5 (7.4e5)	6.4e3 (8.1e3)	Low		
WSB x K18-ACE2 F1	M	66	UD	32.0	11	4.2e6 (1.8e6)	3.1e4 (1.6e4)	High	Moderate Group A	Group A and B Males > Females
	F	0	8	25.1	8	3.7e6 (4.1e6)	2.5e4 (1.2e4)	High		
BALB/c x K18-ACE2 F1	M	100	UD	3.0	5	6.0e5 (5.5e5)	1.6e4 (1.3e4)	Moderate	No/Low	Group A and B Females > Males
	F	66	UD	8.3	6	4.4e5 (4.3e5)	6.0e2 (1.2e3)	High	High Group A	
DBA/2 x K18-ACE2 F1	M	57	UD	16.0	10	1.5e6 (1.7e6)	1.7e4 (2.2e4)	Moderate	No/Low	Group A and B Females > Males
	F	71	UD	3.6	6	2.0e6 (1.0e6)	1.2e3 (1.0e3)	Moderate		

466　UD: undetermined

467　BLD: below limit of detection

a Sensitive

C57Bl/6JxK18-hACE2

A/JxK18-hACE2

b Resistant

PWKxK18-hACE2   NZOxK18-hACE2

129S1xK18-hACE2

c Sensitive with Sex Bias

CASTxK18-hACE2   NODxK18-hACE2

WSBxK18-hACE2

Fig. 1

Figure 2

Figure 3

Fig. S3

Fig. S4

Male BAL cytokine panel

Male serum cytokine panel

Fig. S6

Female BAL cytokine panel

Fig. S7

Female serum cytokine panel

Fig. S8