**Exploring the structural basis to develop efficient multi-epitope vaccines displaying interaction with HLA and TAP and TLR3 molecules to prevent NIPAH infection, a global threat to human health**

Sukrit Srivastava*[a,b], Ajay Kumar Saxena[b], Michael Kolbe*[c,d],

a     Infection Biology Group, Indian Foundation for Fundamental Research, RaeBareli 229001, India
b     Molecular Medicine Lab., School of Life Science, Jawaharlal Nehru University, 110067 New Delhi, India.
c     Department for Structural Infection Biology, Centre for Structural Systems Biology (CSSB) & Helmholtz-Centre for Infection Research, Notkestraße 85, 22607 Hamburg, Germany.
d     Faculty of Mathematics, Informatics and Natural Sciences, University of Hamburg, Rothenbaumchaussee 19, 20148 Hamburg, Germany.

*Correspondence should be addressed to S.S. (srivastava.sukrit@iffr.in; srivastav.sukrit@gmail.com); and M.K. (Michael.Kolbe@helmholtz-hzi.de)

## ABSTRACT

Nipah virus (NiV) is an emerging zoonotic virus responsible to cause several serious outbreaks in South Asian region with high mortality rate of 40 to 90% since 2001. NiV infection causes lethal encephalitis and respiratory disease with the symptom of endothelial cell-cell fusion. No specific vaccine has yet been reported against NiV infection. Recently, some Multi-Epitope Vaccines (MEV) has been proposed but they involve limited number of epitopes which further limits the potential of vaccine. To address the urgent need for a specific and effective vaccine against NiV infection, in the present study, we have design two multi-epitope vaccines (MEVs) composed of 33 Cytotoxic T lymphocyte (CTL) epitopes and 38 Helper T lymphocyte (HTL) epitopes. Both the MEVs carry

potential B cell linear epitope overlapping regions, B cell discontinuous epitopes as well as IFN-γ inducing epitopes. Hence the designed MEVs carry potential to elicit cell-mediated as well as humoral immune response. Selected CTL and HTL epitopes were validated for their stable molecular interactions with HLA class I and II alleles as well as in case of CTL epitopes, with human transporter associated with antigen processing (TAP). Human β-defensin 2 and β-defensin 3 were used as adjuvants to enhance the immune response of both the MEVs. The molecular dynamics simulation study of MEVs-TLR3(ECD) (Toll-Like Receptor 3 Ectodomain) complex indicated stable molecular interaction. Further, the codon optimized cDNA of both the MEVs has shown high expression potential in the mammalian host cell line (Human). Hence for further studies, both the design of CTL and HTL MEVs could be cloned, expressed and tried for *in-vivo* validations (animal trails) as potential vaccine candidates against NiV infection.

**Key words:** Nipah virus (NiV), Human Transporter associated with antigen processing (TAP), Toll-Like Receptor 3 (TLR-3), Epitope, Immunoinformatics, Molecular Docking, Molecular Dynamics (MD) simulation, Multi-epitope Vaccine (MEV)

**INTRODUCTION**

Nipah virus (NiV) is an emerging zoonotic virus of the genus Henipavirus of the Paramyxoviridae family [1]. NiV infection causes fatal encephalitis and respiratory disease with a particular symptom of endothelial cell-cell fusion [2].

The first NiV infection to human was first reported in Malaysia in 1998. Later NiV

outbreak was reported from Meherpur, Bangladesh in 2001. In the Malaysia NiV

infection, the transmission was primarily due being in contact of pigs, whereas in

later outbreaks of Bangladesh and India the transmission was associated with

contaminated date palm sap and human-to-human contact [3]. Bats are identified

as the main reservoir for the NiV and they are responsible for the transmission of

the infection to both humans and animals [4]. After 2001 NiV outbreak has been

reported from different district of Bangladesh almost every year (2003-05, 2007-

12). Till March 31, 2012, a total of 209 confirmed cases of NiV infections were

reported out of which 161 people died resulting in the mortality rate as high as

77%. After several outbreaks in Bangladesh, in total three NiV outbreaks have

also been reported from India. Two of them occurred in the state of West Bengal

in 2001 and 2007 [5]. The most recent NiV outbreak was reported from the

Kerala state of India during the period of May to June-2018. The Kerala outbreak

claimed 17 lives leaving only two survivors out of 19 confirmed cases [6]. Till

present, there has been no specific vaccine reported against NiV infection, and

the pathogenesis mechanism of NiV to human cells is largely unknown. Hence,

an immune-informatics approach investigating the potential of different NiV

proteins for vaccine design would be an important and essential step forward for

vaccine development.

NiV infection of human cells involves several protein-protein interactions

and protein cluster formation on the host cell surface. Essential proteins involved

in NiV pathogenesis include C protein, Fusion glycoproteins (F), Glycoproteins

(G), Matrix proteins (M), Nucleocapsid protein, Phosphoprotein, Polymerase, V protein and the W protein [7-21]. The C protein regulates the early host pro-inflammatory response as well as the pathogen virulence thus providing a conducive environment for a successful NiV infection [7]. The attachment glycoprotein (G), the fusion protein (F) and the matrix protein (M) together form a cluster on the human cell membrane facilitating virus particle assembly and pathogenesis [8-12]. The G and F proteins of NiV have been shown to be immunogenic by inducing protective immune responses in hamsters [22, 23]. The NiV matrix protein is observed to play a central role in virus particle formation and is essentially required for viral budding from the infected human cells [13-15]. The NiV Polymerase is responsible for the initiation of RNA synthesis, primer extension, and transition to elongation mode and hence the enzyme facilitates viral pathogenesis and survival in host cells [16]. The phosphoprotein and the glycoprotein of NiV are crucially involved in the regulation of viral replication [17, 18] while the V protein of NiV is responsible for the host interferon (IFN) signalling evasion during pathogenesis [19, 20]. Interestingly, the identical N-terminal region of the pathogen's V and W protein is sufficient to exert the IFN-antagonist activity [21]. Hence, all the nine above mentioned NiV proteins are crucial in different ways for viral pathogenesis and are important drug and vaccine candidates.

The Nipah virus is a zoonotic RNA virus and it infects human respiratory epithelium cells as well as differentiated neurons (in the brain and spinal cord). Thus, as understood by previous animal model studies, recovery from viral

infection and the clearance of viral RNA requires the presence of virus-specific

antibodies and interferon gamma (IFN-γ) secretion from T cells [24-26]. Along

with the B cell, the T cell also play a critical role in immune response against NiV

infection. In recent studies a number of B cell and T cell epitopes from the NIPAH

proteome have been reported [27-38]. Most of these epitopes show strong

interaction tendency with their respective HLA molecule binders. Further different

approaches were proposed for the design of multi epitope vaccines [39, 40]. The

proposed vaccines utilized a limited number (6 to 8) of T and B cell epitopes. The

use of limited number of epitopes could be challenging for the successful

presentation of the exogenous vaccine candidates in view of the proteolytic

cleavage by Antigen Presenting Cells (APC).

In present study we have screened out the most potential Cytotoxic T

lymphocyte (CTL) epitopes, Helper T lymphocyte (HTL) and B cell epitopes from

the NiV proteome. We have shortlisted and priotized  the most potential and

highest scoring 33 CTL, 38 HTL and 16 B cell epitopes. We further studied

several critical properties (like IC(50), Immunogenicity, Conservancy, Non-toxic

etc) to identify the most potential epitopes against Nipah virus. The shortlisted

epitopes were utilized for the design of CTL and HTL multi-epitope vaccines

against Nipah virus. The designed vaccines were further studied for their stable

interaction with immunological receptor the Toll-Like Receptor 3 (TLR3). The

analysis of cDNA of the designed multi-epitope vaccine has predicted to be

highly favorable for expression in mammalian cell line. Overall in the present

study we have designed and proposed potential multi-epitope vaccines against

Nipah virus infection.

**METHODOLOGY**

In the present study, we have designed two multi-epitope vaccines

(MEVs) composed of thoroughly screened most potential Cytotoxic T lymphocyte

(CTL) epitopes and Helper T lymphocyte (HTL) epitopes derived from the nine

NiV proteins (glycoprotein (gi-253559848); C protein (gi-1859635642); fusion

protein (gi-13559813); matrix protein (gi-13559811); nucleoprotein (gi-

1679387250); phosphoprotein (gi-1802790259); V protein (gi-1802790260); RNA

polymerase (gi-15487370); W protein (gi-374256971)). CTL and HTL epitopes

would be the most potential vaccine candidates since they are responsible for

cell-mediated immune response by their presentation on the surface of antigen

presenting cells (APCs) by their respective Class I and II human leukocyte

antigen (HLA) allele binders. Both the CTL and HTL epitopes chosen for MEV

design also carry overlapping regions of linear B cell epitopes. Moreover, both

the MEVs also carry potential discontinuous B cell epitopes as well as IFN-γ

inducing epitopes in their tertiary structure model. Hence both the designed

MEVs carry potential to elicit cell-mediated as well as humoral immune response.

Furthermore, both the MEVs were designed with human β Defensin 2 and human

β Defensin 3 as adjuvant at their N and C termini to enhance the immunogenic

response [41-42]. The β-defensins have considerable immunological adjuvant

activity. The β-defensins 2 & 3 have been shown in previous studies to generate

potent humoral immune responses when fused with B-cell lymphoma epitopes in

mouse model [43-46]. Since, the pro-inflammatory mediators enhance the expression of β-defensins 2 & 3 in airway epithelial cells we chose β-defensins in our study. The selected CTL and HTL epitopes were validated for their stable molecular interactions with their respective HLA alleles binders; for CTL epitopes, their molecular interaction with human Transporter associated with antigen processing (TAP) was also analyzed [47, 48]. This analysis validated the CTL epitopes that get transported through the TAP cavity for their presentation on cell surface or not. The human TAP selectively pumps cytosolic peptides into the lumen of the endoplasmic reticulum in an ATP-dependent manner [49]. The tertiary structure models of both the MEVs were generated, refined, and further docked with the human Toll-Like Receptor 3 (TLR3), which is an essential immunoreceptor in this pathway [50, 51]. The nuclear localization of Nipah virus W protein inhibits the signaling pathway of TLR3 upon pathogenesis to suppress the TLR3 induced activation of the IFN response to eventually prevent relay of the warning signals to uninfected cells. TLR3 is preferentially expressed by human astrocytes of the central nervous system (CNS) upon infection.  The NiV infection involves invasion of the neurons of CNS and hence causes infection in CNS. These studies indicate the importance of the TLR3 responses in immune response and hence TLR3 has been chosen to be studied for its stable binding with the designed multi-epitope vaccines [52-55]. The complexes of CTL and HTL MEVs formed with the human TLR3 were further analysed for their stable molecular interaction by a molecular dynamics simulation study. The cDNA of the designed MEVs were generated and analysed for their high expression tendency

in the mammalian host cell line (Human). Overall, from the present *in-silico* study,

we may put forward the design of two MEVs, which qualify all the significant

criterions for being a potential vaccine candidate against NiV infection. The

corresponding workflow is shown in Supplementary figure S1.

*NiV proteins selected for potential epitope screening.* In the present study, nine

NiV proteins were used for epitope screening. They include C protein, Fusion

glycoproteins (F), Glycoproteins (G), Matrix proteins (M), Nucleocapsid protein,

Phosphoprotein, Polymerase, V protein and the W protein. The full-length protein

sequences of NiV proteins were retrieved from the NCBI database (National

Center for Biotechnology Information (https://www.ncbi.nlm.nih.gov/protein). A

total of 161 protein sequences available at NCBI, belonging to different strains

and origins of NiV, were retrieved. For structural based epitope screenings

available tertiary structures of NiV proteins were retrieved from Protein Data

Bank (PDB) (http://www.rcsb.org/pdb/home/home.do). For the NiV proteins with

no structure available, homology models were generated by Swiss-model,

(http://swissmodel.expasy.org/) [56] (Supplementary table S1).

**Screening of Potential Epitopes**

***T cell Epitope Prediction***

*Screening of Cytotoxic T lymphocyte (CTL) Epitope.* The screening of Cytotoxic

T lymphocyte epitopes was performed by the IEDB (Immune Epitope Database)

tool "Proteasomal cleavage/TAP transport/MHC class I combined predictor"

(http://tools.iedb.org/processing/) [57-59]. Proteasome cleavage score depend on

the total amount of cleavage site in the protein. TAP score estimates an effective

log -(IC50) values (half maximal inhibitory concentration (IC50)) for binding to

TAP of a peptide or its N-terminal prolonged precursors. The MHC binding

prediction score is -log(IC50) values for binding to MHC of a peptide [60]. The

tool gives a "Total Score" which is a combined score of the proteasome, MHC,

TAP (N-terminal interaction), and processing analysis scores. The total score is

generated by using the combination of six different methods viz. Consensus, NN-

align, SMM-align, Combinatorial library, Sturniolo and NetMHCIIpan. The IC(50)

(nM) value for each epitope and MHC allele binding pairs were also obtained by

this IEDB tool. Epitopes having high, intermediate, and least affinity of binding to

their HLA allele binders have IC50 values < 50 nM, < 500 nM and < 5000 nM

respectively. Immunogenicity of all the screened CTL epitopes was also obtained

by    using    "MHC    I    Immunogenicity"    tool    of    IEDB

(http://tools.iedb.org/immunogenicity/) with all the parameters set to default

analyzing 1st, 2nd, and C-terminus amino acids of the given screened epitope

[60]. The tool predicts the immunogenicity of a given peptide-MHC (pMHC)

complex on the basis of the physiochemical properties of constituting amino acid

and their position within the peptide sequence.

*Screening of Helper T lymphocyte (HTL) Epitopes.* To identify the Helper T

lymphocyte epitopes from NiV proteins, the IEDB tool "MHC-II Binding

Predictions" (http://tools.iedb.org/mhcii/) was used. Peptides with IC50 values

<50 nM are considered high affinity, <500 nM intermediate affinity and <5000 nM

low affinity [61-64]. The tool generates percentile rank for each potential peptide.

This percentile rank is generated by the combination of three different methods

viz. combinatorial library, SMM_align and Sturniolo and by comparing the score

of the peptide against the scores of other random five million 15-mer peptides of

SWISSPROT database [61-64]. The rank from the consensus of all three

methods was generated by the median percentile rank of the three methods.

Lower the value of percentile, higher would be the rank.

*Population Coverage by CTL and HTL epitopes.* The "Population Coverage" tool

of IEDB (http://tools.iedb.org/population/) was used to elucidate the world human

population coverage by the shortlisted 33 CTL and 38 HTL epitopes derived from

nine NiV proteins [65]. The T cells recognize complex between a specific major

MHC molecule and a particular pathogen-derived epitope. The given epitope will

elicit a response only in an individual that express an MHC molecule, which is

capable of binding that particular epitope. This denominated MHC restriction of T

cell responses and the MHC polymorphism provide basis for population coverage

study. The MHC types are expressed at dramatically different frequencies in

different ethnicities. Hence a vaccine with larger population coverage could be of

greater importance [65]. Clinical administration of multiple-epitopes involving both

the CTL and the HTL epitopes are predicted here to have a greater probability of

larger human population coverage worldwide.

***B Cell Epitope Prediction***

*Sequence-based B Cell epitope prediction.* Protein sequence-based six different

methods were utilized to screen linear B cell epitopes from nine different NiV

proteins. These methods are available at "B Cell Epitope Prediction Tools" of

IEDB server (http://tools.iedb.org/bcell/). In this screening the parameters such

as hydrophilicity, flexibility, accessibility, turns, exposed surface, polarity and

antigenic propensity of the polypeptides are correlated with their localization in

the protein. This allows the search for continuous epitopes prediction from

protein sequence. The prediction is based on the propensity scales for each of

the 20 amino acids. For a window size n, the i - (n-1)/2 neighboring residues on

each side of residue i are used to compute the score for the residue i [66-71].

These methods utilize the propensity scale method as well as the physiochemical

properties of the given antigenic sequence to screen potential epitopes using

"Bepipred Linear Epitope Prediction", "Chou & Fasman Beta-Turn Prediction",

"Emini Surface Accessibility Prediction", "Karplus & Schulz Flexibility Prediction",

"Kolaskar & Tongaonkar Antigenicity" and "Parker Hydrophilicity Prediction" tools

[66-71].

*Structure-based B cell epitope prediction.* The Ellipro (ElliPro: Antibody Epitope

Prediction tool; http://tools.iedb.org/ellipro/) and the DiscoTope2.0 (DiscoTope:

Structure-based    Antibody    Prediction    tool;    http://tools.iedb.org/discotope/)

methods available at IEDB, were used to screen the linear and the discontinuous

B cell epitopes [72, 73]. The ElliPro method analyses on the basis of the location

of residue in the protein's 3D structure. The residues lying outside of an ellipsoid

covering 90% of the inner core protein residues score highest Protrusion Index

(PI) of 0.9; and so on. The discontinuous epitopes predicted by the ElliPro tool

are clustered on the basis of the distance "R" in Å between two residue's centers

of mass lying outside of the largest possible ellipsoid. The larger value of R

indicates larger distant residues (residue discontinuity) are screened in the

epitopes. The Discotope 2.0 method is based on the "contact number" of the

residue's Cα carbon atom as well as on the propensity of a given residue to be a

part of an epitope [72, 73]. The residue "contact number" is the number of Cα

atoms in the antigen within a distance of 10 Å of the particular residue's Cα atom.

A low contact number would indicate the residue being close to the surface or in

protruding regions of the antigen's structures.

**Characterisation of potential epitopes**

*Epitope conservation analysis.* The shortlisted CTL, HTL and B cell epitopes

screened from nine NiV proteins were analysed for the conservancy of their

amino    acid    sequence    by    "Epitope    Conservancy    Analysis"    tool

(http://tools.iedb.org/conservancy/) of IEDB. The epitope conservancy is the

fraction of protein sequences that contain that particular epitope. The analysis

was done against their entire respective source protein sequences of NiV

proteins retrieved from the NCBI protein database [74].

*Epitope          Toxicity          prediction.*          The          tool          ToxinPred

(http://crdd.osdd.net/raghava/toxinpred/multi_submit.php) was used to analyse

the toxicity of shortlisted CTL, HTL and B cell epitopes. The tool allows to

identifying highly toxic or non-toxic short peptides. The toxicity check analysis

was done by the "SVM (Swiss-Prot) based" (support vector machine) method

utilizing dataset of 1805 sequences as positive, 3593 negative sequences from

Swissprot as well as an alternative dataset comprises the same 1805 positive

sequences and 12541 negative sequences from TrEMBLE [75].

*Overlapping residue analysis.* The overlapping residue analysis for the shortlisted

CTL, HTL and the B cell linear epitopes was performed by the Multiple Sequence

Alignment (MSA) analysis by Clustal Omega tool

(https://www.ebi.ac.uk/Tools/msa/clustalo/) of EBI (European Bioinformatics

Institute) [76]. The Clustal Omega multiple sequence alignment tool virtually

aligns any number of protein sequences and delivers an accurate alignment.

*Epitope selected for molecular interaction study with HLA allele and TAP*

*transporter.* On the basis of the overlapping residue analysis of shortlisted CTL,

HTL and linear B cell epitopes few numbers of CTL and HTL epitopes were

chosen for further analysis involving stable interaction with their respective HLA

allele binders and TAP cavity interaction (Supplementary table S3 & S4, Figure

3). These epitopes were chosen on the basis of them having partial or full

overlapping sequence region amongst all three types of epitopes (CTL, HTL and

B Cell), or having full sequence overlap amongst any of the two types of

epitopes, or having the highest number of the HLA allele binders.

**Molecular interaction analysis of selected epitopes with HLA allele and TAP**

**transporter.**

*Tertiary structure modelling of HLA alleles and selected T cell epitopes.* The

Swiss-model was used for homology modelling of the HLA class I and II allele

binders of shortlisted epitopes [56]. The amino acid sequence of the HLA allele

binders were retrieved from Immuno Polymorphism Database (IPD-IMGT/HLA)

(https://www.ebi.ac.uk/ipd/imgt/hla/allele.html).    Templates    for    homology

modelling were chosen on the basis of highest amino acid sequence similarity.

All the HLA allele models were further validated by their QMEAN value. The

QMEAN value gives a composite quality estimate involving both global as well as

local analysis of the model [77]. Generated Model having acceptable QMEAN

value (cutoff -4.0) were chosen for further studies (Supplementary table S2).

The "Natural Peptides Module for Beginners" module of PEPstrMOD

(http://osddlinux.osdd.net/raghava/pepstrmod/nat_ss.php)    was    utilized    to

generate tertiary structures for the selected CTL and HTL epitopes [78]. The time

window for simulation was set to 100 picoseconds (ps) in a vacuum environment.

*Molecular interaction analysis of selected CTL and HTL epitopes with HLA*

*alleles.* The AutoDock 4.2 (ADT) and the AutoDock Vina were used for *in-silico*

molecular docking study of the selected CTL and HTL epitopes with their

respective HLA class I and II allele binders [79, 80]. The generated docked

complexes were studied for their stable nature by molecular dynamics simulation.

MD simulation was performed by the Gromacs 5.1.4 using the Optimized

Potential for Liquid Simulations - all-atom force field (OPLS-AA) [81, 82].

*Molecular interaction analysis of selected CTL epitopes with TAP transporter.*

TAP transporter plays an important role in the presentation of CTL epitope. From

the cytosol after proteasome processing, the fragmented peptide of foreign

protein gets transported to endoplasmic reticulum (ER) through the TAP

transporter, from the ER these short peptides reach to the Golgi bodies and then

get presented on the cell surface [83]. Molecular interaction study of the

shortlisted CTL epitopes with the TAP transporter cavity was performed by

molecular docking using AutoDock Vina [79, 80]. As structural model the cryo-

EM structure of TAP transporter (PDB ID: 5u1d) after removing the antigen from

TAP cavity of the original structure [48] was used for epitope-TAP interaction

study.

**Design, characterisation and molecular interaction analysis of Multi-**

**Epitope Vaccines with immunological receptor.**

*Design of Multi-Epitope Vaccines.* All the screened CTL and HTL epitopes were

utilized to design CTL and HTL Multi-Epitope vaccines. Short peptides EAAAK

and GGGGS were used as rigid and flexible linkers respectively (Figure 1). The

GGGGS linker provides proper conformational flexibility to the vaccine tertiary

structure and hence facilitates stable conformation to the vaccine. The EAAAK

linker facilitates in domain formation and hence facilitates the vaccine to obtain

its final stable structure [84-86]. The human beta-defensin 2 (hBD-2) (PDB ID:

1FD3, Sequence: GIGDPVTCLKSGAICHPVFCPRRYKQIGTCGLPGTKCCKKP)

and the human beta-defensin 3 (hBD-3) (PDB ID: 1KJ6, Sequence:

GIINTLQKYYCRVRGGRCAVLSCLPKEEQIGKCSTRGRKCCRRKK) were used

as adjuvants in the design of both the MEVs at N and C terminals respectively

[41-46, 87,]. Human Beta-defensins (hBD) have an important role in the

chemotactic activity memory T cells, immature dendritic cells and monocytes.

Beta-defensins are also involved in degranulation of the mast cells. Due to the

important role of hBDs in immune response enhancement, hBDs have been

chosen and utilized as adjuvants for the MEV designs.

**Figure 1. Design of Multi-Epitope Vaccine (MEVs). (A)** CTL and **(B)** HTL

epitopes were linked by the short peptide linker 'GGGGS'. Human β Defensin 2

and β Defensin 3 were used as an adjuvant at the N and C terminals

respectively. The short peptide EAAAK was used to link the β Defensin 2 and β

Defensin 3. Epitopes from different proteins were coloured in different colours. C

terminal 6xHis is designed as His tag. * Epitopes common to Phosphoprotein, V

Protein and W protein.

*Characterisation of designed Multi-Epitope Vaccines*

*Interferon gamma inducing epitope prediction.* From the designed amino acid

sequence of both the MEVs potential interferon gamma (IFN-γ) epitopes were

screened by the "IFNepitope" server

(http://crdd.osdd.net/raghava/ifnepitope/scan.php) using "Motif and SVM hybrid",

(MERCI: Motif-EmeRging and with Classes-Identification, and SVM: support

vector machine) method. The tool predicts peptides from protein sequences

having the capacity to induce IFN-gamma release from CD4+T cells. This

module generates overlapping IFN-gamma inducing peptides from query

sequence. . For the screening, IEDB database with 3705 IFN-gamma inducing

and 6728 non-inducing MHC class II binders is utilized [88, 89].

*MEVs allergenicity and antigenicity prediction.* The designed MEVs were further

analysed for allergenicity and antigenicity prediction by utilizing the AlgPred

(http://crdd.osdd.net/raghava/algpred/submission.html) and the Vaxigen

(http://www.ddg-pharmfac.net/vaxijen/VaxiJen/VaxiJen.html) tools respectively

[90, 91]. The AlgPred prediction is based on the similarity of already known

epitope with any region of the submitted protein. For the screening of

allergenicity, the Swiss-prot dataset consisting of 101725 non-allergens and 323

allergens is utilized [90]. The VaxiJen utilizes an alignment-free approach, solely

based on the physicochemical properties of the query amino acid sequence. For

prediction of antigenicity, the Bacterial, viral and the tumour protein datasets

were used to derive models for the prediction of whole protein antigenicity. Every

set consisted of known 100 antigens and 100 non-antigens [91].

*Physicochemical property analysis of designed MEVs.* The ProtParam

(https://web.expasy.org/protparam/) tool was utilized to analyse the

physiochemical properties of the designed CTL and HTL MEVs [92]. The

ProtParam analysis performs an empirical investigation for the given query amino

acid sequence. ProtParam computes various physicochemical properties derived

from a given protein sequence.

*Tertiary structure modelling and refinement of MEVs.* The tertiary structure of the

designed CTL and HTL MEVs were calculated by homology modelling utilizing

the RaptorX structure prediction tool

(http://raptorx.uchicago.edu/StructurePrediction/predict/) [93]. RaptorX predicts

template-based secondary and tertiary structures, contacts, solvent accessibility,

disordered regions and binding sites for given protein sequence, with or with out

having close homologs in the Protein Data Bank (PDB). RaptorX also assigns

confidence scores to indicate the quality of a predicted 3D model [94-96]. Quality

assessment for both the generated homology models of CTL and HTL MEVs was

performed by their respective P-values. The P-value for a predicted homology

model is a probability score of the generated model being worse than the best.

Hence the P-value indicates a relative quality of the generated model in terms of

modelling error, combining the global distance test (GDT) and the un-normalized

global distance test (uGDT) indicating the error involved at each residue. The

smaller the P-value the greater the quality of a predicted model.

The refinement of both the generated MEV models was performed by

ModRefiner (https://zhanglab.ccmb.med.umich.edu/ModRefiner/) and

GalaxyRefine    tool    (http://galaxy.seoklab.org/cgi-bin/submit.cgi?type=REFINE)

[97, 98]. Modrefiner improves the physical realism and structural accuracy of the

model using a Two-step Atomic-level Energy Minimization. ModRefiner is an

algorithm    for    the    atomic-level,    high-resolution    protein    tertiary    structure

refinement. Both the side-chain and the backbone atoms of protein structure are

completely    flexible    during    the    structure    refinement    simulations.    The

conformational search is guided by a composite of physics and knowledge based

force field. The tool generates significant improvement in the physical quality of

the local structures [97]. TM-score generated by ModRefiner indicates the

structural similarity of the refined model with the original input model. Closer the

TM-Score to 1, higher would be the similarity of original and the refined model.

The GalaxyRefine tool refines the query tertiary structure by repeated structure

perturbation as well as by utilizing the subsequent structural relaxation by the

molecular dynamics simulation. The tool GalaxyRefine generates reliable core

structures from multiple templates and then re-builds unreliable loops or termini

by using an optimization-based refinement method [96, 99]. To avoid any breaks

in the 3D model GalaxyRefine uses triaxial loop closure method. The MolProbity

score    generated    for    a    given    refined    model    indicates    the    log-weighted

combination of the clash score (the number of atomic clashes per 1000 atoms),

the Ramachandran favored backbone torsion angles and the percentage of bad

side-chain rotamers (the percentages of rotamer outliers). The 'GDT-HA' (Global

Distance Test-High Accuracy) generated by the tool indicates the backbone

structure accuracy; 'RMSD' (Root mean Square Deviation) indicates the overall

structural deviation in refined model from the initial model and the 'Rama favored'

indicates percentage of Ramachandran favored residues.

*Validation of CTL and HTL MEVs refined models.* Both the refined CTL and HTL

MEV 3D models were further validated by RAMPAGE analysis tool

(http://mordred.bioc.cam.ac.uk/~rapper/rampage.php) [100, 101]. The generated

Ramachandran plots for the MEV models show the sterically allowed and

disallowed residues along with their dihedral psi ($\psi$) and phi ($\phi$) angles.

*Discontinuous B-cell epitope prediction from MEVs.* Both the generated tertiary

models of designed CTL and HTL MEVs were subjected to discontinuous B cell

epitopes prediction. The structure-based discontinuous B cell epitopes were

screened from both the MEV models by utilizing the ElliPro method as described

earlier [72].

**Molecular interaction analysis of MEVs with TLR-3**

Molecular interaction analysis of both the designed MEVs with Toll-Like receptor-

3 (TLR-3), was performed by molecular docking and molecular dynamics

simulation. Molecular docking was performed by PatchDock server

(http://bioinfo3d.cs.tau.ac.il/PatchDock/) [102-104]. PatchDock utilizes algorithm

for unbound (real life) docking of molecules for protein-protein. The algorithm

carries out the rigid docking, with the surface variability/flexibility implicitly

addressed through liberal intermolecular penetration. The algorithm focuses on

the (i) initial molecular surface fitting on localized, curvature based surface

patches (ii) use of Geometric Hashing and Pose Clustering for initial

transformation detection (iii) computation of shape complementarity utilizing the

Distance Transform (iv) efficient steric clash detection and geometric fit scoring

based on a multi-resolution shape representation and (v) utilization of biological

information by focusing on hot spot rich surface patches [102-104]. For molecular

docking, the 3D structure of human TLR-3 ectodomain (ECD) was retrieved from

the PDB databank (PDB ID: 2A0Z). Further, the molecular dynamics simulation

study of the MEVs-TLR-3 complexes were performed by Gromacs 5.1.4, utilizing

the Optimized Potential for Liquid Simulations - all-atom force field (OPLS-AA)

[81, 82]. MD simulation was performed to understand the properties of the MEVs-

TLR3 complexes in terms of their structure and the microscopic interactions in

the complex. The study provides dynamical properties of the designed system

with MEVs-TLR3 complexes with a guess at the interactions between the

molecules, and also it gives 'exact' predictions of bulk properties. MD Simulations

act as a bridge between theory and experiment [81, 82].

***In-silico* analysis of MEVs for cloning and expression**

*Analysis of cDNA of the MEVs for cloning and expression in mammalian cell line.*

Complementary DNA of both the MEVs, codon optimized for expression in

Mammalian cell line (Human) were generated by Java Codon Adaptation Tool

(http://www.jcat.de/). The generated cDNA were further analysed by GenScript

Rare Codon Analysis Tool (https://www.genscript.com/tools/rare-codon-analysis).

The tool analyses the GC content, Codon Adaptation Index (CAI) and the

Tandem rare codon frequency for a given cDNA [105-107]. The CAI indicates the

possibility of cDNA expression in a chosen expression system. The tandem rare

codon frequency indicates the presence of low-frequency codons in the given

cDNA.

**RESULTS & DISCUSSION**

**Screening of potential epitopes**

***T cell Epitope Prediction***

*Screening of Cytotoxic T lymphocyte (CTL) Epitope.* Cytotoxic T lymphocyte

(CTL) epitopes screened were shortlisted according to the highest "Total Score",

low IC(50) (nM) value for epitope-HLA class I allele complexes, and epitopes with

the larger number of the HLA class I allele binders. The immunogenicity of the

shortlisted CTL epitopes was also determined; the higher immunogenicity score

indicates the greater immunogenic potential of the given epitope (Supplementary

table S3, S7). A total of 33 CD8+ T cell epitopes were finally chosen. Out of the

33 CTL epitopes reported here 10 epitopes (Fusion Protein: FALSNGVLF;

Glycoprotein: TVYHCSAVY; Nucleocapsid: YPALALNEF; Phosphoprotein:

VSDAKMLSY; Polymerase: YPECNNILF, FPVMGNRIY,

AEFFSFFRTF, IPFLFLSAY, ETDDYNGIY, SQNLLVTSY) show a match with

previous studies [35-38], indicating consensus of epitope screening by different

approaches and methods (Supplementary table S3).

*Screening of Helper T lymphocyte (HTL) epitopes.* The screening of helper T

lymphocyte (HTL) epitopes from nine different proteins of NiV was performed on

the basis of "Percentile rank". The smaller the value of percentile rank the higher

would be the affinity of the peptide with its respective HLA allele binders. In our

initial screening, we got several potential CD4+ T cell epitopes with high scoring.

38 epitopes out of initial screening were shortlisted on the basis that they had

highest percentile rank and highest number of HLA class II allele binders

(Supplementary table S4, S8).

*Population Coverage by CTL and HTL epitopes.* The population coverage by the

shortlisted epitopes was also studied, in particular involving countries of South

Asia, East Asia, Northeast Asia and the Middle East. From this study, we may

conclude that the combined use of all the shortlisted CTL and HTL epitopes

would have an average worldwide population coverage as high as 97.88%, with

a standard deviation of 21.97 (Supplementary table S5).

**B Cell epitope prediction**

*Sequence-based B Cell epitope prediction.* In our initial study, we screened a

total of 116 B Cell epitope from nine different NiV proteins, with the epitope

length of at least four amino acids utilizing the Bepipred Linear Epitope Prediction

method. B cell epitopes predicted by another five different methods based on

different physiochemical properties were found to have significant consensus

with the epitope amino acid sequences predicted by Bepipred Linear Epitope

Prediction. Here, 16 out of the 116 epitopes were shortlisted having a length of 4

to 19 amino acids (Supplementary table S6, Figure 2). One of these 16 B cell

epitopes (Matrix Protein: SIPREFMIY) matches with a previous study [39],

indicating epitope screening consensus using different approaches and methods

(Supplementary table S6).

**Figure 2. Overlapping regions amongst the linear B cell epitopes predicted**

**by the BepiPred method and other seven different B cell epitopes**

**prediction methods.** B cell epitopes predicted by the BepiPred method and

other different protein sequence based (Chou.., Emini.., Karplus.., Kolaskar..,

and Parker..) and protein structure based (DiscoTope and ElliPro) prediction

methods were found to have significant consensus. Consensus overlapping

regions of BepiPred epitopes are underlined by the different colour,

corresponding to respective prediction method.

*Structure-based B cell epitope prediction.* Structure-based discontinuous and

linear epitopes predicted by the DiscoTope 2.0 and the Ellipro methods have

shown a significant consensus of overlapping amino acid sequence with the

linear epitopes predicted by Bepipred linear epitopes method (Supplementary

table S6, Figure 2). This result further confirms that the shortlisted B Cell

Bepipred Linear Epitopes have a high chance of being highly immunogenic

epitopes.

**Characterisation of potential epitopes**

*Epitope conservation analysis.* Sequence conservation analysis of the shortlisted 33 CTL, 38 HTL and 16 B cell epitopes showed that the '100% amino acid sequence' conservancy of CTL, HTL and B cell epitopes amongst all retrieved NiV protein sequences is mostly 100%, as shown in Supplementary table S3, S4, S6. This result indicates the high conservancy nature of the amino acid sequence of the shortlisted CTL, HTL and B cell epitopes.

*Epitope toxicity prediction.* Toxicity analysis of all the shortlisted CTL, HTL and B Cell epitopes was also performed. The ToxinPred study indicated the non-toxic nature of all the shortlisted epitopes (Supplementary table S3, S4, S6).

*Overlapping residue analysis.* Amino acid sequence overlap analysis amongst the shortlisted CTL, HTL and B cell epitopes from nine NiV proteins was performed by using the Multiple Sequence Alignment (MSA) analysis tool Clustal Omega. Our analysis showed that several epitopes of CTL, HTL and B cell have overlapping amino acid sequence. The CTL, HTL and B cell epitopes having two or more amino acids overlap are shown in Figure 3.

**Figure 3. Overlapping CTL, HTL and B cell epitopes**. Multiple sequence alignment performed by Clustal Omega at EBI to identify the consensus overlapping regions of CTL (red), HTL (blue) and B cell epitopes (green) amongst shortlisted epitopes. Epitopes with overlapping region amongst all the

three types of epitopes (CTL, HTL and B Cell epitopes), epitopes with full

sequence overlap and epitopes with the highest number of HLA allele binders

were chosen for further studies (encircled).

*Epitope selected for molecular interaction study with HLA allele and TAP*

*transporter.* Amongst all the shortlisted epitope peptides seven CTL and

seventeen HTL epitope peptides have partial or full overlapping sequences or

have the highest number of HLA allele binders were shortlisted for further studies

(Figure 3, Supplementary table S3 & S4).

**Molecular interaction analysis of selected epitopes with HLA allele and TAP**

**transporter.**

*Molecular interaction analysis of selected CTL and HTL epitopes with HLA*

*alleles.* The molecular docking study of the shortlisted CTL and HTL epitopes

with their respective HLA class I and II allele binders was performed using

AutoDock Vina. Docking studies revealed for all epitopes significant molecular

interaction with their HLA allele binders having low binding energies and multiple

hydrogen bonds formated (Figure 4A & 4B). The stability of the obtained docking

complexes was further tested by  molecular dynamics (MD) simulation studies.

MD simulations were performed over a time interval of 0.5-1 ns at the invariable

temperature of ~ 300 K and at invariable pressure of ~ 1 bar. All the complexes

showed reasonably invariant root mean square deviation (RMSD) value

(between ~ 0.2 to 0.4 nm) indicating the stable nature of the tested epitope-HLA

allele complexes (Figure 5A & 5B). Moreover, the reasonably invariant Rg (radius of gyration) of the complexes, throughout the MD simulation (Supplementary figure S2), and the root mean square fluctuation (RMSF) for all the atoms of the complexes (Supplementary figure S3) again indicate the stable nature of the epitopes and HLA allele complexes. Furthermore, the B-factor analysis of all the epitope-HLA allele complexes indicated most of the complex regions to be stable (blue) with a very small region being acceptably fluctuating (yellow and orange) (VIBGYOR colour presentation) Supplementary figure S4.

**Figure 4. (A) Molecular Docking analysis of CTL epitopes and HLA alleles.** Molecular docking of selected CTL epitopes (cyan sticks) with their respective HLA class I allele binders (gray sticks). The study shows the docked complexes to have significantly negative binding energy along with hydrogen bonds (green dots) formation in the complex interface. **(B) Molecular Docking analysis of HTL epitopes and HLA alleles.** Molecular docking of selected HTL epitopes (cyan sticks) with their respective HLA class II allele binders (gray sticks). The study shows the docked complexes to have significantly negative binding energy along with hydrogen bonds (green dots) formation in the complex interface.

**Figure 5. (A) Molecular Dynamics simulation analysis of CTL epitopes and HLA allele complexes.** Molecular Dynamics simulation study reveals a stable nature of the CTL-HLA allele complexes throughout 0.5-1 ns time window with reasonably invariable RMSD. **(B) Molecular Dynamics simulation analysis of**

**CTL epitopes and HLA allele complexes.** Molecular Dynamics simulation study

reveals a stable nature of the HTL-HLA allele complexes throughout 0.5-1 ns

time window with reasonably invariable RMSD.

*Molecular interaction analysis of selected CTL epitopes with TAP transporter.*

The molecular docking interaction analysis of the chosen CTL epitopes with the

TAP transporter cavity showed a significantly strong molecular interaction with

low binding energy and several hydrogen bonds formed at different sites of the

TAP transporter cavity. Two sites of interaction were of particular interest, one

located near the cytoplasmic end and the other in the vicinity of the ER lumen

(Figure 6). Our study confirms the transportation feasibility of the chosen CTL

epitopes from the cytoplasm into the ER lumen which is essential for the

representation of peptides by the HLA allele molecules on the surface of antigen

presenting cells.

**Figure 6. Molecular docking analysis of CTL epitopes within the TAP**

**transporter cavity.** Molecular interaction of CTL epitopes (cyan sticks) within the

TAP cavity (gray ribbon/sticks) is shown in detail. For every panel of epitope-TAP

complex, **(A)** shows the binding of epitope at two different sites within TAP cavity,

**(B)** and **(C)** show detailed molecular interaction between epitopes and TAP

cavity; (a, b) show chain A and B of TAP transporter. H-bonds are shown in

yellow dots. (*) Binding energy, shown in kcal/mol.

**Characterisation and molecular interaction analysis of designed Multi-Epitope Vaccines with immunological receptor**

*Characterisation of designed Multi-Epitope Vaccines*

*Interferon-gamma inducing epitope prediction.* Interferon-gamma (IFN-γ) inducing epitopes are involved in both the adaptive and the innate immune response. The IFN-γ inducing 15-mer peptide epitopes were screened from the CTL and HTL MEVs by utilizing the IFNepitope server. A total of 33 CTL MEV and 43 HTL MEV INF-γ inducing POSITIVE epitopes with a score of 1 or more than 1 were shortlisted (Supplementary table S9, Figure 7D & 7I).

**Figure 7. Tertiary structure modelling of CTL and HTL Multi-Epitope Vaccines. (A) & (F)**: Tertiary structural models of CTL and HTL MEVs showing epitopes in different colours corresponds to as in Figure 1. **(B) & (G)**: Show the different domains of CTL and HTL MEVs. **(C) & (H)**: The overlapping linear B cell epitope region present in CTL and HTL MEVs, shown by spheres. **(D) & (I):** From the CTL and HTL MEVs, the INF-γ inducing epitopes are shown in cyan, discontinuous B Cell epitopes are shown in magenta and the region common amongst INF-γ and discontinuous B Cell epitopes are shown in wheat colour. **(E) & (J)**: RAMPAGE analysis of the refined CTL and HTL MEV models.

*MEVs allergenicity and antigenicity prediction.* Both the CTL and HTL MEVs were analyzed to be NON-ALLERGEN by the AlgPred analysis scoring -0.61243421 and -0.93493027 respectively while default threshold value being -

0.4. CTL and HTL MEVs were also analyzed by VaxiJen to be probable

ANTIGENS with the prediction score of 0.4447 and 0.4836 respectively, while the

default threshold value for viral proteins being 0.4. Hence, with the mentioned

analysis tools both the CTL and HTL MEVs are predicted to be non-allergic as

well as antigenic in nature.

*Physicochemical property analysis of designed MEVs.* ProtParam analysis were

performed for both the CTL and HTL MEVs to analyse their physiochemical

properties. The CTL MEV is composed of 576 amino acids, has a molecular

weight of 58.57 kDa and a  theoretical pI of 8.19. The expected half-life of the

CTL MEV in *E.coli*, yeast and mammalian reticulocytes were predicted with 10 h,

20 min, and 30 h respectively; the aliphatic index of CTL MEV was found to be

58.42, and grand average of hydropathicity (GRAVY) of CTL MEV was found to

be -0.010, both indicating globular and hydrophilic nature of the CTL MEV. The

instability index score of the CTL MEV was 48.03 indicating its stable folding

under native conditions.

Further, the ProtParam analysis of the HTL MEV showed for the 857

amino acids, a molecular weight of 87.62 kDa and a  theoretical pI of 5.99. The

expected half-life of HTL MEV in *E.coli*, yeast and mammalian reticulocytes was

predicted to be 10 h, 20 min and 30 h, respectively. The aliphatic index of HTL

MEV was calculated as 82.99, and the grand average of hydropathicity (GRAVY)

of the HTL MEV was found to be 0.188, both indicating that HTL MEV has a

globular and hydrophilic nature. The instability index of the HTL MEV was 45.66

indicating its stable nature.

*Tertiary structure modelling and refinement of MEVs.* 3D homology models were

generated for both the CTL and HTL MEVs by utilizing the RaptorX modelling

tool (Figure 7A, 7F). The model obtained for CTL MEV has 6% helix, 27% β-

sheet, 66% coil content and structural elements are 23% exposed, 39% medium

and 36% buried. The structural model has three domains ranging from amino

acid 1 to 46 (1$^{st}$ domain, template-1fd3:A), 47 to 520 (2$^{nd}$ domain, templates-

1yrzA, 1y7bA, 5jozA, 3zxjA, 5z5dA) and 521 to 576 (3$^{rd}$ domain , template-

1kj6:A) (Figure 7B). Similarly, the 3D model calculated with RaptorX for the HTL

MEV has 21% helix, 22% β-sheet, 56% coil content with 24% of the amino acids

exposed, 39% medium and 36% buried. The structural model has three domains

ranging from amino acid 1 to 46 (1$^{st}$ domain, template-1fd3:A), 47 to 801 (2$^{nd}$

domain, templates-5m5zA, 3eqnA), 802 to 857 (3$^{rd}$ domain, template-1kj6:A)

(Figure 7G). The P-Value for the best template based CTL and HTL MEV

homology models were 2.79e-04 and 5.99e-03 respectively. Good quality, mostly

alpha proteins have a P-value of less than $10^{-3}$ and that of mostly beta proteins

has a P-value of less than $10^{-4}$. Hence both the homology models of CTL and

HTL MEVs are predicted to be of good quality. Since for the CTL and HTL MEV

design, the CTL and HTL epitopes used also show overlapping common regions

with the linear B cell epitopes (Figure 2), both the generated CTL and HTL MEV

models also carry the overlapping regions of linear B Cell epitopes (Figure 7C,

7H).

The generated CTL and HTL 3D models were further refined using

ModRefiner to fix structural gaps followed by GalaxyRefine refinement.

Refinement with ModRefiner showed a TM-score of 0.9703 and 0.8934 for the

CTL and HTL models respectively, both being close to 1 indicating the initial and

the refined models were structurally similar. For the CTL MEV model refinement,

the sore of models 1 for different parameters were, Rama favored was 90.8%,

GDT-HA was 0.9596, RMSD was 0.385, MolProbity was 2.673, Clash score was

29.9, and Poor rotamers was 1.8. Likewise for HTL MEV model refinement, the

sore of models 1 for different parameters were, Rama favoured was 88.5%,

GDT-HA was 0.9463, RMSD was 0.419, MolProbity was 2.811, Clash score was

38.4, and Poor rotamers was 1.6. Here, MolProbity shows the log-weighted

combination of the clash score, percentage Ramachandran not favoured and the

percentage bad side-chain rotamers. After refinement, all the mentioned

parameters were significantly improved in comparison to the initial CTL and HTL

MEV models (Supplementary table S10).

*Validation of CTL and HTL MEVs refined models.* Both the CTL and HTL model

were analysed with the RAMPAGE analysis tool after refinement. The refined

CTL MEV model has 91.5% residues in favored regions, 6.8% residues in

allowed regions, and only 1.7% residues in the outlier region; while the refined

HTL MEV model was found to have 89.5% of residues in favored region, 8.7%

residues in allowed region, and 1.9% residues in the outlier region (Figure 7E,

7J).

*Discontinuous B-cell epitope prediction from MEVs.* Discontinuous B-cell

epitopes were further predicted from the final refined 3D models of CTL and HTL

MEVs utilizing the ElliPro tool on IEDB server. The screening revealed that the

CTL MEV carries 3 and the HTL MEV has 2 potential discontinuous epitopes.

The PI (Protrusion Index) score of the CTL MEV discontinuous B cell epitopes

ranges from 0.682 to 0.747 and that of HTL MEV it ranges from 0.687 to 0.745

(Supplementary table S11, Figure 7D & 7I). The higher PI score indicates a

greater potential of the discontinuous B cell epitope.

**Molecular interaction analysis of MEVs with TLR-3 .**

The refined models of CTL and HTL MEVs were further studied for their

molecular interaction with the ectodomain (ECD) of human TLR-3. Therefore,

molecular docking of CTL and HTL MEVs model with the TLR-3 crystal structure

model (PDB ID: 2A0Z) was performed utilizing the PatchDock tool. Generated

docking conformation with highest scores of 22382 and 18264 for CTL and HTL

MEVs, respectively were selected for further studies. The highest docking score

predicted with the PatchDock tool indicates the best geometric shape

complementarity fitting conformation of MEVs and the TLR-3 receptor. Both, the

CTL and HTL MEVs were fitting into the ectodomain region of TLR-3 after

docking (Figue 8A & 8E). The CTL and HTL MEVs have shown to form multiple

hydrogen bonds within the ectodomain cavity region of TLR-3. Further, the

molecular dynamics simulation study was also performed for the docked

complexes of both the MEVs and TLR-3. In MD simulations both the complexes

have shown reasonably stable RMSD value between ~ 0.2 to 0.5 nm for a given

time window of 10 ns at invariable pressure (~ 1 bar) and temperature (~ 300 K)

(Figure 8B & 8F). The reasonably invariant radius of gyration (Rg) of both MEVs-

TLR-3 complexes (Figure 8C & 8G), and RMS fluctuation (RMSF) for all the

atoms in both the complexes (Figure 8D & 8H) indicate that the MEVs-TLR3

complexes are stable. The B-factor analysis of MEVs-TLR3 complexes was also

performed. The B-factor indicates the displacement of the atomic positions from

an average (mean) value i.e. the more flexible an atom is the larger the

displacement from the mean position will be (mean-squares displacement)

(Figure 8A and 8E). The areas with high B-factors are colored red (hot), while low

B-factors are colored blue (cold) (VIBGYOR presentation). The B-factor of most

of the regions of MEVs-TLR3 complexes indicates the stable nature of the

complexes while a very small region is found to be fluctuating. The results

suggest a stable complex formation tendency for both the CTL and HTL MEVs

with the ectodomain of the human TLR-3 receptor.

**Figure 8. Molecular Docking and dynamics simulation study of CTL and**

**HTL MEVs with TLR-3. (A)** CTL and **(E)** HTL MEVs (VIBGYOR) docked

complex with TLR-3 (gray). Both the complexes are forming several hydrogen

bonds in the MEV and TLR-3 interface, as shown by green dots. B-Factor of the

docked MEVs is shown by a rainbow (VIBGYOR) presentation. The regions in

blue being indicated stable and the region in red indicate unstable. In the above

complexes, most of the region of docked MEVs is in blue and with the very small

region is green, yellow or orange, hence the complexes are predicted to be very

stable. **(B)** and **(F),** RMSD as generated by the molecular Dynamics simulation

study of CTL, HTL MEVs and TLR-3 complexes. **(C)** & **(G)** Rg (radius of gyration)

across the time window of 10 nanosecond. **(D)** & **(H)**, RMS fluctuation for all the

atoms of the CTL, HTL MEVs and TLR-3 complexes.

***In-silico* analysis of MEVs for cloning and expression**

*Analysis of cDNA of both the MEVs for cloning and expression in mammalian cell*

*line.* Complementary DNA codon optimized for CTL and HTL expression in

mammalian host cell lines (Human) was generated with the Java Codon

Adaptation Tool. The generated optimized cDNA's for both the MEVs were also

analysed by utilizing the GenScript Rare Codon Analysis Tool revealing aGC

content of optimized CTL-MEV cDNA of 69.79% and a CAI (Codon Adaptation

Index) score of 1.00 with 0% tandem rare codons. Likewise, the GC content of

the optimized HTL-MEV cDNA was 70.69%, CAI score was 1.00 with 0% tandem

rare codons. Since for higher possibility for cDNA expression in human

expression system, the GC content of a cDNA should be within the range of 30%

to 70%, the CAI score should be between 0.8-1.0, and the tandem rare codon

frequency that indicates the presence of low-frequency codons, should be <30%,

the cDNA constructs of CTL and HTL MPVs are expected to have high potential

for expression in human expression system. The tandem rare codons may hinder

the proper expression of the cDNA or even interrupt the translational machinery

of the chosen expression system. According to the GenScript Rare Codon

Analysis the cDNA of both the MEVs satisfy all the mentioned parameters and

are predicted to have high expression in the mammalian host cell line (Human).

**CONCLUSION**

In the present study, we have designed and validated two multi-epitope

vaccines derived from CTL and HTL epitopes. The selected peptides show

significant sequence overlap with screened linear B cell epitopes. Both the CTL

and HTL MEVs tertiary models carry potential discontinuous B cell epitopes and

INF-γ epitopes. Consequently, the designed MEVs might have the potential to

elicit profound humoral and cellular immune responses. Human β-Defensin 2 and

3 fused to the N and C terminal ends of both the MEVs serving as adjuvants to

enhance the immune response. The identified epitopes for the CTL and HTL

MEVs were also validated by molecular docking and MD simulation studies to

test the interaction with their respective HLA allele binders. Molecular interaction

of the selected CTL epitopes within the TAP transporter cavity was also

evaluated indicating a favorable transport of epitopes from cytoplasm to lumen of

Endoplasmic Reticulum for further presentation on cell surface by Golgi bodies.

Our analysis of the shortlisted CTL and HTL epitopes combined revealed

coverage of 97.88% world human population. The molecular interaction analysis

of both the CTL and HTL MEVs with the immunoreceptor TLR3 showed

structural fit of the MEVs into the ectodomain of TLR3 receptor cavity and the MD

simulations indicate a very stable complex formation. Since both the CTL and

HTL MEVs carry CTL, HTL as well as discontinuous B cell epitopes, the

combined administration of both the MEVs, is predicted to elicit both the humoral

as well as cell-mediated immune response. The cDNA for both the MEVs were

designed considering codon-bias for the expression in mammalian host cell lines

(Human). The cDNAs were optimized with respect to their GC content and zero

tandem rare codons  to gain high expression in mammalian host cell lines

(Human). In future experimental studies the designed CTL and HTL MEVs

should be cloned, expressed and validated *in-vivo* and in animal trails as

potential vaccine candidates against NiV infection.

**Supplementary figure S1.** Workflow chart.

847
848   **Supplementary figure S2. (A)** Rg (radius of gyration) for the CTL epitope – HLA
849   class I allele complexes, across the time window of 1 nano second. **(B)** Rg for
850   the HTL epitope – HLA class II allele complexes, across the time window of 1
851   nano second.

852
853   **Supplementary figure S3. (A)** RMS fluctuation in nanometers for all the atoms
854   of the CTL epitope – HLA class I allele complexes. **(B)** RMS fluctuation in
855   nanometers for all the atoms of the HTL epitope – HLA class II allele complexes.

856
857   **Supplementary figure S4. (A)** B-Factor of CTL epitope – HLA class I allele
858   complexes **(B)** B-Factor of HTL epitope – HLA class II allele complexes. Epitopes
859   are shown in sticks and HLA alleles are shown in gray cartoons. B-factor is
860   indicated by rainbow (VIBGYOR) colour, blue for stable region and red for most
861   unstable region.

862
863   **Supplementary table S1. Protein sequence retrieval, tertiary structures**
864   **retrieval and homology modeling of nine Nipah proteins.** Nipah  protein
865   sequences were retrieved from NCBI. Available structure files (pdb) for Nipah
866   proteins were retrieved from RCSB PDB. Nipah proteins with no tertiary structure
867   available were subjected to homology modeling by Swissmodel.

868
869   **Supplementary table S2. Homology modeling for HLA alleles.** Tertiary
870   structure of HLA alleles were modeled by homology modeling using SwissModel

871  server. Templates were chosen with highest sequence identity. Generated
872  models with acceptable QMEAN value were chosen for further studies.
873

**Supplementary table S3. Shortlisted high scoring CTL epitopes.** Selected
high scoring CTL epitopes and their respective HLA alleles binders are listed. *In-*
*silico* analysis have shown all the selected epitopes to be non-toxic (Non-Toxin)
as well as they show significant conservancy. ToxinPred analysis is based on the
ToxinPred main dataset used by "ToxinPred" algorithm to predict toxicity of any
unknown peptide. # Epitope match with previous studies indicating consensus in
epitope screening by different approaches and methods.
881

882  **Supplementary table S4. Shortlisted high scoring HTL epitopes.** Selected
883  high scoring HTL epitopes and their respective HLA alleles binders are listed
884  above. *In-silico* analysis have shown all the selected epitopes to be non-toxic
885  (Non-Toxin) as well as they show significant conservancy.
886

887  **Supplementary table S5. World population coverage by the shortlisted CTL**
888  **and HTL epitopes combined.** With a standard deviation of 21.97 on an average
889  97.88% of world population could be covered by the joint administration of
890  selected CTL and HTL epitopes as vaccine candidate.
891

**Supplementary table S6. Shortlisted B Cell epitopes.** BepiPred Linear B Cell
epitopes showing sequence overlap with CTL and HTL epitopes are shortlisted
above. *In-silico* analysis have shown all the selected epitopes to be non-toxic
(Non-Toxin) as well as they show significant amino acid sequence conservancy.
# Epitope match with previous studies indicating consensus in epitope screening by
different approaches and methods.
898

899  **Supplementary table S7. CTL epitope prediction.** Detailed scoring of all
900  screened CTL epitopes and their respective HLA class I allele binders. CTL
901  epitopes were chosen on the basis of high "Total score" and higher number of
902  HLA allele binders. Total score is a combined score of TAP score, MHC score,
903  Proteasome score and Processing score.
904

905  **Supplementary table S8. HTL epitope prediction.** Percentile rank of HTL
906  epitopes and their respective HLA class II allele binders. HTL epitopes were
907  screened on the basis of percentile rank (lower the percentile number, higher the
908  rank) and larger number of HLA allele binders. Last column show the method
909  used for epitope screening.
910

911  **Supplementary table S9. INF-γ epitopes from CTL and HTL MEVs.** INF-γ
912  inducing (POSITIVE) epitopes from CTL and HTL MEVs were screened by using
913  "Motif and SVM hybrid" (MERCI & SVM) approaches.
914

915 **Supplementary table S10. Refinement models of CTL and HTL MEVs.** CTL
916 and HTL MEVs models were refined by GalaxyWEB server and used for further
917 studies. After refinement in particular Rama favored residues increased
918 significantly.
919
920 **Supplementary table S11. B Cell discontineous epitopes of CTL & HTL**
921 **MEVs.** Discontinuous B Cell epitopes predicted by ElliPro (IEDB) from CTL &
922 HTL MEVs.
923

**ACKNOWLEDGEMENTS:**

We acknowledge the Advance Instrumentation Research Facility (AIRF) at
Jawaharlal Nehru University, New Delhi for providing the advance computational
facility to conduct experiments. We also acknowledge Indian Foundation for
Fundamental Research (IFFR) for providing resources and funding

**FUNDING:**

Indian Foundation for Fundamental Research (IFFR)

**AUTHOR CONTRIBUTION:**

Idea conceived, methodology designed and performed by S.S., critical data
analysis and scientific writing was done by S.S. and M.K., facility of MD
simulation was provided by A.K.S, draft was finalized by S.S., M.K., A.K.S.

**ADDITIONAL INFORMATION:**

Authors declare to have no competing interests.

951 **FIGURES:**

**A. CTL epitopes**
**B. HTL epitopes**

ß Defensin 2: GIGDPVTCLKSGAICHPVFCPRRYKQIGTCGLPGTKCCKKP

| B. HTL epitopes | A. CTL epitopes |
|---|---|
| EAAAK | EAAAK |
| **C Protein** | |
| VQMTYNWTQWLQTLY | MMASILLTLF |
| GGGGS | GGGGS |
| DLALSKYLSDLLFVF | AQITAGVALY |
| GGGGS | GGGGS |
| LALSKYLSDLLFVFG | FALSNGVLF |
| GGGGS | GGGGS |
| ALSKYLSDLLFVFGP | KYLSDLLFVF |
| GGGGS | GGGGS |
| LSKYLSDLLFVFGPN | MTIQAISQAF |
| GGGGS | GGGGS |
| SKYLSDLLFVFGPNL | AENPVFTVF |
| GGGGS | GGGGS |
| KYLSDLLFVFGPNLQ | AVYNNEFYY |
| GGGGS | GGGGS |
| YLSDLLFVFGPNLQD | LAMDEGYFAY |
| GGGGS | GGGGS |
| **Fusion Protein** | |
| ASFSWDTMIKFGDVL | TVYHCSAVY |
| GGGGS | GGGGS |
| FSWDTMIKFGDVLTV | NYMYLICYGF |
| GGGGS | GGGGS |
| GVYNDAFLIDRINWI | YMIPRTMLEF |
| GGGGS | GGGGS |
| NDAFLIDRINWISAG | EIISDIGNY |
| GGGGS | GGGGS |
| DAFLIDRINWISAGV | TPFVDSRAY |
| GGGGS | GGGGS |
| AFLIDRINWISAGVF | YPALALNEF |
| GGGGS | GGGGS |
| FLIDRINWISAGVFL | LDPVVTDVVY* |
| GGGGS | GGGGS |
| **Glycoprotein** | |
| IPREFMIYDDVFIDN | LVSDAKMLSY* |
| GGGGS | GGGGS |
| FMIYDDVFIDNTGRI | MPSDDFSNTF |
| GGGGS | GGGGS |
| **Matrix Protein** | |
| LSSDQVAELAAAVQE | VSDAKMLSY* |
| GGGGS | GGGGS |
| SSDQVAELAAAVQET | AEFFSFFRTF |
| GGGGS | GGGGS |
| SDQVAELAAAVQETS | ETDDYNGIY |
| GGGGS | GGGGS |
| DQVAELAAAVQETSA | FPISRLFNMY |
| GGGGS | GGGGS |
| QVAELAAAVQETSAG | FPVMGNRIY |
| GGGGS | GGGGS |
| **Nucleocapsid** | |
| *NNGNVCLVSDAKMLS | IATVYTWAY |
| GGGGS | GGGGS |
| *NGNVCLVSDAKMLSY | IMKKSFKAY |
| GGGGS | GGGGS |
| *GNVCLVSDAKMLSYA | IPFLFLSAY |
| GGGGS | GGGGS |
| *NVCLVSDAKMLSYAP | KWYECFLFWF |
| GGGGS | GGGGS |
| *VCLVSDAKMLSYAPE | KYYQIDQPFF |
| GGGGS | GGGGS |
| **Phosphoprotein** | |
| NIDNIHLLAEFFSFF | LETDDYNGIY |
| GGGGS | GGGGS |
| IDNIHLLAEFFSFFR | RLFNMYRSY |
| GGGGS | GGGGS |
| DNIHLLAEFFSFFRT | SQNLLVTSY |
| GGGGS | GGGGS |
| NIHLLAEFFSFFRTF | SYFGLVLVCF |
| GGGGS | GGGGS |
| IHLLAEFFSFFRTFG | TSDLDFVIFY |
| GGGGS | GGGGS |
| LELASFLMDRRVILP | YPECNNILF |
| GGGGS | GGGGS |
| ELASFLMDRRVILPR | |
| GGGGS | |
| LASFLMDRRVILPRA | |
| GGGGS | |
| ASFLMDRRVILPRAA | |
| GGGGS | |
| LDFVIFYASLTYLRR | |
| GGGGS | |
| FVIFYASLTYLRRGI | |
| EAAAK | EAAAK |

**Polymerase**

ß Defensin 3: GIINTLQKYYCRVRGGRCAVLSCLPKEEQIGKCSTRGRKCCRRKK
HHHHHH        HHHHHH

**Figure 1. Design of Multi-Epitope Vaccine (MEVs). (A)** CTL and **(B)** HTL epitopes were linked by the short peptide linker 'GGGGS'. Human β Defensin 2 and β Defensin 3 were used as an adjuvant at the N and C terminals respectively. The short peptide EAAAK was used to link the β Defensin 2 and β Defensin 3. Epitopes from different proteins were coloured in different colours. C terminal 6xHis is designed as His tag. * Epitopes common to Phosphoprotein, V Protein and W protein.

**Overlapping regions of predicted B cell Bepipred linear Epitope and epitopes predicted by other protein sequence and structure based methods**

| Protein | Bepipred epitope | Overlapping regions of sequence based B cell linear epitope prediction | | | | | Overlapping regions of structure based B cell epitope prediction | | |
|---|---|---|---|---|---|---|---|---|---|
| | | Chou & Fasman Beta-Turn | Emini Surface Accessibility | Karplus & Schulz Flexibility | Kolaskar & Tongaonkar Antigenicity | Parker Hydrophilicity | DiscoTope: discontinuous epitopes | ElliPro Linear | ElliPro Discontinuous |
| Fusion Protein | 215 GPNLQDPVSNSM 226 | 215-226 | | 215-226 | 215-216 | 217-226 | | | |
| Glycoprotein | 271 WTPPNPNT 278 | 271-278 | 271-278 | 272-278 | | 272-278 | | | 271-278 |
| Glycoprotein | 459 SWDTMI 464 | 459-460 | | | | | | | 463-464 |
| Glycoprotein | 529 NQTAE 533 | 529-532 | 529-533 | 529-533 | 529-533 | 529-533 | | | |
| Glycoprotein | 184 SGIYM 188 | 184-188 | | 184-188 | 184-188 | 184-188 | | | |
| Matrix Protein | 330 SIPREFMIY 338 | 330-335 | | 330-336 | | 331-334 | 332-333 | | 333-338 |
| Matrix Protein | 340 DVFIDNTGRI 349 | 341-349 | | 340-349 | | 340-344 | | | 340-349 |
| Matrix Protein | 125 GYGFTSSPERGWSDYTSGA 143 | 125-143 | 130-136 | 128-143 | | 127-143 | | | |
| Phosphoprotein | 163 IAVSKEDR 170 | 166-170 | 166-170 | 163-170 | 163-166 | 165-170 | | | |
| Phosphoprotein | 346 NIDN 349 | 348-349 | | | | | | | |
| Polymerase | 364 GHPILE 369 | 364-367 | | | 365-369 | | | | 366-369 |
| Polymerase | 1095 DKSFDEDLEL 1104 | 1096-1103 | 1096-1101 | 1095-1104 | | 1096-1103 | | | 1095-1104 |
| Polymerase | 1108 LMDR 1111 | | | | | | | | 1108-1111 |
| Polymerase | 1409 LRLETDDYNG 1418 | 1412-1418 | 1410-1416 | 1412-1418 | | 1411-1418 | 1409-1417 | | 1409-1417 |
| Polymerase | 1982 GFPIS 1986 | 1982-1986 | | 1982-1986 | 1982-1986 | | | | 1182-1986 |
| Polymerase | 2004 PVYSNPD 2010 | 2004-2010 | 2006-2010 | 2007-2010 | 2004-2008 | 2007-2010 | | | 2004-2010 |

**Figure 2. Overlapping regions amongst the linear B cell epitopes predicted**

**by the BepiPred method and other seven different B cell epitopes**

**prediction methods.** B cell epitopes predicted by the BepiPred method and

other different protein sequence based (Chou.., Emini.., Karplus.., Kolaskar..,

and Parker..) and protein structure based (DiscoTope and ElliPro) prediction

methods were found to have significant consensus. Consensus overlapping

regions of BepiPred epitopes are underlined by the different colour,

corresponding to respective prediction method.

Clusters of overlapping CTL, HTL & B cell epitopes

**Figure 3. Overlapping CTL, HTL and B cell epitopes**. Multiple sequence

alignment performed by Clustal Omega at EBI to identify the consensus

overlapping regions of CTL (red), HTL (blue) and B cell epitopes (green)

amongst shortlisted epitopes. Epitopes with overlapping region amongst all the

three types of epitopes (CTL, HTL and B Cell epitopes), epitopes with full

sequence overlap and epitopes with the highest number of HLA allele binders

were chosen for further studies (encircled).

A. CTL epitopes - HLA Class I allele complexes

B. HTL epitopes - HLA Class II allele complexes

**Figure 4 (A). Molecular Docking analysis of CTL epitopes and HLA alleles.**

Molecular docking of selected CTL epitopes (cyan sticks) with their respective HLA class I allele binders (gray sticks). The study shows the docked complexes to have significantly negative binding energy along with hydrogen bonds (green dots) formation in the complex interface. **(B) Molecular Docking analysis of HTL epitopes and HLA alleles.** Molecular docking of selected HTL epitopes (cyan sticks) with their respective HLA class II allele binders (gray sticks). The study shows the docked complexes to have significantly negative binding energy along with hydrogen bonds (green dots) formation in the complex interface.

**Figure 5 (A). Molecular Dynamics simulation analysis of CTL epitopes and**

**HLA allele complexes.** Molecular Dynamics simulation study reveals a stable

nature of the CTL-HLA allele complexes throughout 0.5-1 ns time window with

reasonably invariable RMSD. **(B) Molecular Dynamics simulation analysis of**

**CTL epitopes and HLA allele complexes.** Molecular Dynamics simulation study

reveals a stable nature of the HTL-HLA allele complexes throughout 0.5-1 ns

time window with reasonably invariable RMSD.

**Figure 6. Molecular docking analysis of CTL epitopes within the TAP transporter cavity.** Molecular interaction of CTL epitopes (cyan sticks) within the TAP cavity (gray ribbon/sticks) is shown in detail. For every panel of epitope-TAP complex, **(A)** shows the binding of epitope at two different sites within TAP cavity, **(B)** and **(C)** show detailed molecular interaction between epitopes and TAP cavity; (a, b) show chain A and B of TAP transporter. H-bonds are shown in yellow dots. (*) Binding energy, shown in kcal/mol.

**CTL multi-epitope vaccine:**

**HTL multi-epitope vaccine:**

**Figure 7. Tertiary structure modelling of CTL and HTL Multi-Epitope**

**Vaccines. (A) & (F)**: Tertiary structural models of CTL and HTL MEVs showing

epitopes in different colours corresponds to as in Fig. 1. **(B) & (G)**: Show the

different domains of CTL and HTL MEVs. **(C) & (H)**: The overlapping linear B cell

epitope region present in CTL and HTL MEVs, shown by spheres. **(D) & (I):** From

the CTL and HTL MEVs, the INF-γ inducing epitopes are shown in cyan,

discontinuous B Cell epitopes are shown in magenta and the region common

amongst INF-γ and discontinuous B Cell epitopes are shown in wheat colour. **(E)**

**& (J)**: RAMPAGE analysis of the refined CTL and HTL MEV models.

**Figure 8. Molecular Docking and dynamics simulation study of CTL and**

**HTL MEVs with TLR-3. (A)** CTL and **(E)** HTL MEVs (VIBGYOR) docked

complex with TLR-3 (gray). Both the complexes are forming several hydrogen

bonds in the MEV and TLR-3 interface, as shown by green dots. B-Factor of the

docked MEVs is shown by a rainbow (VIBGYOR) presentation. The regions in

blue being indicated stable and the region in red indicate unstable. In the above

complexes, most of the region of docked MEVs is in blue and with the very small

region is green, yellow or orange, hence the complexes are predicted to be very

stable. **(B)** and **(F),** RMSD as generated by the molecular Dynamics simulation

study of CTL, HTL MEVs and TLR-3 complexes. **(C)** & **(G)** Rg (radius of gyration)

across the time window of 10 nanosecond. **(D)** & **(H)**, RMS fluctuation for all the

atoms of the CTL, HTL MEVs and TLR-3 complexes.

1532

1533

1534

1535

1536

1537

1538

1539

1540

1541

1542

1543

1544

1545

1546

1547

**REFERENCES:**

1. Angeletti, S., Presti, A.L., Cella, E. and Ciccozzi, M., 2016. Molecular epidemiology and phylogeny of nipah virus infection: a mini review. Asian Pacific journal of tropical medicine, 9(7), pp.630-634.
2. Aguilar, H.C., Henderson, B.A., Zamora, J.L. and Johnston, G.P., 2016. Paramyxovirus glycoproteins and the membrane fusion process. Current clinical microbiology reports, 3(3), pp.142-154.
3. Ang, B.S., Lim, T.C. and Wang, L., 2018. Nipah Virus Infection. Journal of clinical microbiology, pp.JCM-01875.
4. WHO Report, Surveillance and outbreak alert, Nipah virus; https://www.who.int/health-topics/nipah-virus-infection#tab=tab_1
5. Plowright RK, Becker DJ, Crowley DE, Washburne AD, Huang T, Nameer PO, Gurley ES, Han BA. Prioritizing surveillance of Nipah virus in India. PLoS Negl Trop Dis. 2019 Jun 27;13(6):e0007393. doi: 10.1371/journal.pntd.0007393. PMID: 31246966; PMCID: PMC6597033.
6. Thomas, B., Chandran, P., Lilabi, M. P., George, B., Sivakumar, C. P., Jayadev, V. K., Bindu, V., Rajasi, R. S., Vijayan, B., Mohandas, A., & Hafeez, N. (2019). Nipah Virus Infection in Kozhikode, Kerala, South India, in 2018: Epidemiology of an Outbreak of an Emerging Disease. Indian journal of community medicine : official publication of Indian Association of Preventive & Social Medicine, 44(4), 383–387. https://doi.org/10.4103/ijcm.IJCM_198_19
7. Mathieu, C., Guillaume, V., Volchkova, V.A., Pohl, C., Jacquot, F., Looi, R.Y., Wong, K.T., Legras-Lachuer, C., Volchkov, V.E., Lachuer, J. and Horvat, B., 2012. Nonstructural Nipah virus C protein regulates both the early host proinflammatory response and viral virulence. Journal of virology, pp.JVI-01203.
8. Liu, Q., Chen, L., Aguilar, H.C. and Chou, K.C., 2018. A stochastic assembly model for Nipah virus revealed by super-resolution microscopy. Nature communications, 9(1), p.3050.
9. Johnston, G.P., Contreras, E.M., Dabundo, J., Henderson, B.A., Matz, K.M., Ortega, V., Ramirez, A., Park, A. and Aguilar, H.C., 2017. Cytoplasmic motifs in the nipah virus fusion protein modulate virus particle assembly and egress. Journal of virology, pp.JVI-02150.
10. Satterfield, B.A., Cross, R.W., Fenton, K.A., Borisevich, V., Agans, K.N., Deer, D.J., Graber, J., Basler, C.F., Geisbert, T.W. and Mire, C.E., 2016. The Nipah virus C and W proteins contribute to respiratory disease in ferrets. Journal of virology, pp.JVI-00215.
11. Lamp, B., Dietzel, E., Kolesnikova, L., Sauerhering, L., Erbar, S., Weingartl, H. and Maisner, A., 2013. Nipah virus entry and egress from polarized epithelial cells. Journal of virology, pp.JVI-02696.
12. Weise, C., Erbar, S., Lamp, B., Vogt, C., Diederich, S. and Maisner, A., 2010. Tyrosine residues in the cytoplasmic domains affect sorting and fusion activity of the Nipah virus glycoproteins in polarized epithelial cells. Journal of virology, 84(15), pp.7634-7641.
13. Ciancanelli, M.J. and Basler, C.F., 2006. Mutation of YMYL in the Nipah virus

matrix protein abrogates budding and alters subcellular localization. Journal of virology, 80(24), pp.12070-12078.

14. Patch, J.R., Crameri, G., Wang, L.F., Eaton, B.T. and Broder, C.C., 2007. Quantitative analysis of Nipah virus proteins released as virus-like particles reveals central role for the matrix protein. Virology journal, 4(1), p.1.

15. Patch, J.R., Han, Z., McCarthy, S.E., Yan, L., Wang, L.F., Harty, R.N. and Broder, C.C., 2008. The YPLGVG sequence of the Nipah virus matrix protein is required for budding. Virology journal, 5(1), p.137.

16. Jordan, P.C., Liu, C., Raynaud, P., Lo, M.K., Spiropoulou, C.F., Symons, J.A., Beigelman, L. and Deval, J., 2018. Initiation, extension, and termination of RNA synthesis by a paramyxovirus polymerase. PLoS pathogens, 14(2), p.e1006889.

17. Ranadheera, C., Proulx, R., Chaiyakul, M., Jones, S., Grolla, A., Leung, A., Rutherford, J., Kobasa, D., Carpenter, M. and Czub, M., 2018. The interaction between the Nipah virus nucleocapsid protein and phosphoprotein regulates virus replication. Scientific reports, 8(1), p.15994.

18. Baronti, L., Erales, J., Habchi, J., Felli, I.C., Pierattelli, R. and Longhi, S., 2015. Dynamics of the intrinsically disordered C-terminal domain of the Nipah virus nucleoprotein and interaction with the X domain of the phosphoprotein as unveiled by NMR spectroscopy. ChemBioChem, 16(2), pp.268-276.

19. Uchida, S., Horie, R., Sato, H., Kai, C. and Yoneda, M., 2018. Possible role of the Nipah virus V protein in the regulation of the interferon beta induction by interacting with UBX domain-containing protein1. Scientific reports, 8(1), p.7682.

20. Ludlow, L.E., Lo, M.K., Rodriguez, J.J., Rota, P.A. and Horvath, C.M., 2008. Henipavirus V protein association with Polo-like kinase reveals functional overlap with STAT1 binding and interferon evasion. Journal of virology, 82(13), pp.6259-6271.

21. Park, M.S., Shaw, M.L., Munoz-Jordan, J., Cros, J.F., Nakaya, T., Bouvier, N., Palese, P., García-Sastre, A. and Basler, C.F., 2003. Newcastle disease virus (NDV)-based assay demonstrates interferon-antagonist activity for the NDV V protein and the Nipah virus V, W, and C proteins. Journal of virology, 77(2), pp.1501-1511.

22. Sakib, M.S., Islam, M., Hasan, A.K.M. and Nabi, A.H.M., 2014. Prediction of epitope-based peptides for the utility of vaccine development from fusion and glycoprotein of nipah virus using in silico approach. Advances in bioinformatics, 2014.

23. Guillaume, V., Contamin, H., Loth, P., Georges-Courbot, M.C., Lefeuvre, A., Marianneau, P., Chua, K.B., Lam, S.K., Buckland, R., Deubel, V. and Wild, T.F., 2004. Nipah virus: vaccination and passive protection studies in a hamster model. Journal of virology, 78(2), pp.834-840.

24. Koyuncu, O.O., Hogue, I.B. and Enquist, L.W., 2013. Virus infections in the nervous system. Cell host & microbe, 13(4), pp.379-393.

25. Griffin, D.E. and Metcalf, T., 2011. Clearance of virus infection from the CNS. Current opinion in virology, 1(3), pp.216-221.

26. Kong, D., Wen, Z., Su, H., Ge, J., Chen, W., Wang, X., Wu, C., Yang, C., Chen, H. and Bu, Z., 2012. Newcastle disease virus-vectored Nipah encephalitis vaccines induce B and T cell responses in mice and long-lasting neutralizing antibodies in pigs. Virology, 432(2), pp.327-335.

27. Kamthania, M. and Sharma, D.K., 2015. Screening and structure-based modeling of T-cell epitopes of Nipah virus proteome: an immunoinformatic approach for designing peptide-based vaccine. *3 Biotech*, *5*(6), pp.877-882.

28. Kamthania, M. and Sharma, D.K., 2016. Epitope-based peptides prediction from proteome of nipah virus. *International Journal of Peptide Research and Therapeutics*, *22*(4), pp.465-470.

29. Ali, M.T., Morshed, M.M. and Hassan, F., 2015. A computational approach for designing a universal epitope-based peptide vaccine against Nipah virus. *Interdisciplinary Sciences: Computational Life Sciences*, *7*(2), pp.177-185.

30. kumar Sharma, S., Srivastava, S., Kumar, A. and Srivastava, V., 2021. Anticipation of Antigenic Sites for the Goal of Vaccine Designing Against Nipah Virus: An Immunoinformatics Inquisitive Quest. International Journal of Peptide Research and Therapeutics, pp.1-13.

31. Dey, S., Roy, P., Dutta, T., Nandy, A. and Basak, S.C., 2018. Rational Design of Peptide Vaccines for the Highly Lethal Nipah and Hendra Viruses. *bioRxiv*, p.425819.

32. Krishnamoorthy, P.K., Subasree, S., Arthi, U., Mobashir, M., Gowda, C. and Revanasiddappa, P.D., 2020. T-cell Epitope-based Vaccine Design for Nipah Virus by Reverse Vaccinology Approach. *Combinatorial chemistry & high throughput screening*, *23*(8), pp.788-796.

33. Sakib, M.S., Islam, M., Hasan, A.K.M. and Nabi, A.H.M., 2014. Prediction of epitope-based peptides for the utility of vaccine development from fusion and glycoprotein of nipah virus using in silico approach. *Advances in bioinformatics*, *2014*.

34. Eshaghi, M., Tan, W.S. and Yusoff, K., 2005. Identification of epitopes in the nucleocapsid protein of Nipah virus using a linear phage-displayed random peptide library. *Journal of medical virology*, *75*(1), pp.147-152.

35. Mohammed, A.A., Shantier, S.W., Mustafa, M.I., Osman, H.K., Elmansi, H.E., Osman, I.A.A., Mohammed, R.A., Abdelrhman, F.A., Elnnewery, M.E., Yousif, E.M. and Mustafa, M.M., 2020. Epitope-based peptide vaccine against glycoprotein G of Nipah henipavirus using immunoinformatics approaches. *Journal of immunology research*, *2020*.

36. Gupta, A.K., Kumar, A., Rajput, A., Kaur, K., Dar, S.A., Thakur, A., Megha, K. and Kumar, M., 2020. NipahVR: a resource of multi-targeted putative therapeutics and epitopes for the Nipah virus. *Database*, *2020*.

37. Habib, P.T., 2021. Learning from COVID-19 Pandemic: In Silico Vaccine and Cloning Design Against Nipah Virus by Studying and Analyzing the Whole Nipah Virus Proteome.

38. Singh, R.K., Dhama, K., Chakraborty, S., Tiwari, R., Natesan, S., Khandia, R., Munjal, A., Vora, K.S., Latheef, S.K., Karthik, K. and Singh Malik, Y., 2019. Nipah virus: epidemiology, pathology, immunobiology and advances in diagnosis, vaccine designing and control strategies–a comprehensive review. *Veterinary Quarterly*, *39*(1), pp.26-55.

39. Majee, P., Jain, N. and Kumar, A., 2021. Designing of a multi-epitope vaccine candidate against Nipah virus by in silico approach: a putative prophylactic solution for the deadly virus. *Journal of Biomolecular Structure and Dynamics*, *39*(4), pp.1461-1480.

40. Ojha, R., Pareek, A., Pandey, R.K., Prusty, D. and Prajapati, V.K., 2019. Strategic development of a next-generation multi-epitope vaccine to prevent Nipah virus zoonotic infection. *ACS omega*, *4*(8), pp.13069-13079.

41. Wilson, S.S., Wiens, M.E. and Smith, J.G., 2013. Antiviral mechanisms of human defensins. Journal of molecular biology, 425(24), pp.4965-4980.

42. Duits, L.A., Nibbering, P.H., Strijen, E., Vos, J.B., Mannesse-Lazeroms, S.P., Sterkenburg, M.A. and Hiemstra, P.S., 2003. Rhinovirus increases human β-defensin-2 and-3 mRNA expression in cultured bronchial epithelial cells. Pathogens and Disease, 38(1), pp.59-64.

43. Yang, D., Biragyn, A., Kwak, L.W. and Oppenheim, J.J., 2002. Mammalian defensins in immunity: more than just microbicidal. Trends in immunology, 23(6), pp.291-296.

44. Biragyn, A., Surenhu, M., Yang, D., Ruffini, P.A., Haines, B.A., Klyushnenkova, E., Oppenheim, J.J. and Kwak, L.W., 2001. Mediators of innate immunity that target immature, but not mature, dendritic cells induce antitumor immunity when genetically fused with nonimmunogenic tumor antigens. The Journal of Immunology, 167(11), pp.6644-6653.

45. Duits, L.A., Nibbering, P.H., van Strijen, E., Vos, J.B., Mannesse-Lazeroms, S.P., van Sterkenburg, M.A. and Hiemstra, P.S., 2003. Rhinovirus increases human β-defensin-2 and-3 mRNA expression in cultured bronchial epithelial cells. FEMS Immunology & Medical Microbiology, 38(1), pp.59-64.

46. Kohlgraf, K.G., Pingel, L.C., Dietrich, D.E. and Brogden, K.A., 2010. Defensins as anti-inflammatory compounds and mucosal adjuvants. Future microbiology, 5(1), pp.99-113.

47. Antoniou, A.N., Powis, S.J. and Elliott, T., 2003. Assembly and export of MHC class I peptide ligands. Current opinion in immunology, 15(1), pp.75-81.

48. Oldham, M.L., Grigorieff, N. and Chen, J., 2016. Structure of the Transporter associated with antigen processing trapped by herpes simplex virus. eLife, 5, p.e21829.

49. Meena, S.R., Gangwar, S.P. and Saxena, A.K., 2012. Purification, crystallization and preliminary X-ray crystallographic analysis of the ATPase domain of human TAP in nucleotide-free and ADP-, vanadate- and azide-complexed forms. Acta Crystallographica Section F: Structural Biology and Crystallization Communications, 68(6), pp.655-658.

50.  Delneste, Y., Beauvillain, C. and Jeannin, P., 2007. Innate immunity:
structure and function of TLRs. Medecine sciences: M/S, 23(1), pp.67-73.
51. Totura, A.L., Whitmore, A., Agnihothram, S., Schäfer, A., Katze, M.G., Heise,
M.T. and Baric, R.S., 2015. Toll-like receptor 3 signaling via TRIF
contributes to a protective innate immune response to severe acute
respiratory syndrome coronavirus infection. MBio, 6(3), pp.e00638-15.
52.  Shaw, M.L., Cardenas, W.B., Zamarin, D., Palese, P. and Basler, C.F.,
2005. Nuclear localization of the Nipah virus W protein allows for inhibition
of both virus-and toll-like receptor 3-triggered signaling pathways. Journal
of virology, 79(10), pp.6078-6088.
53.  Seto, J., Qiao, L., Guenzel, C.A., Xiao, S., Shaw, M.L., Hayot, F. and
Sealfon, S.C., 2010. Novel Nipah virus immune-antagonism strategy
revealed by experimental and computational study. Journal of virology,
84(21), pp.10965-10973.
54. Farina, C., Krumbholz, M., Giese, T., Hartmann, G., Aloisi, F. and Meinl, E.,
2005. Preferential expression and function of Toll-like receptor 3 in human
astrocytes. Journal of neuroimmunology, 159(1-2), pp.12-19.
55. Weingartl, H., Czub, S., Copps, J., Berhane, Y., Middleton, D., Marszal, P.,
Gren, J., Smith, G., Ganske, S., Manning, L. and Czub, M., 2005. Invasion
of the central nervous system in a porcine host by Nipah virus. Journal of
virology, 79(12), pp.7528-7534.
56. Arnold K, Bordoli L, Kopp J, and Schwede T (2006). The SWISS-MODEL
Workspace: A web-based environment for protein structure homology
modelling. Bioinformatics.,22,195-201.
57. Tenzer S, Peters B, Bulik S, Schoor O, Lemmel C, Schatz MM, Kloetzel PM,
Rammensee HG, Schild H, Holzhutter HG. 2005. Modeling the MHC class
I pathway by combining predictions of proteasomal cleavage, TAP
transport and MHC class I binding. Cell Mol Life Sci 62:1025-1037.
58. Peters B, Bulik S, Tampe R, Van Endert PM, Holzhutter HG. 2003. Identifying
MHC class I epitopes by predicting the TAP transport efficiency of epitope
precursors. J Immunol171:1741-1749.
59.  Hoof, I., Peters, B., Sidney, J., Pedersen, L.E., Sette, A., Lund, O., Buus, S.
and Nielsen, M., 2009. NetMHCpan, a method for MHC class I binding
prediction beyond humans. Immunogenetics, 61(1), p.1.
60. Calis JJA, Maybeno M, Greenbaum JA, Weiskopf D, De Silva AD, Sette A,
Kesmir C, Peters B. 2013. Properties of MHC class I presented peptides
that enhance immunogenicity. PloS Comp. Biol. 8(1):361.
61.  Wang, P., Sidney, J., Kim, Y., Sette, A., Lund, O., Nielsen, M. and Peters,
B., 2010. Peptide binding predictions for HLA DR, DP and DQ molecules.
BMC bioinformatics, 11(1), p.568.
62.  Sidney, J., Assarsson, E., Moore, C., Ngo, S., Pinilla, C., Sette, A. and
Peters, B., 2008. Quantitative peptide binding motifs for 19 human and
mouse MHC class I molecules derived using positional scanning
combinatorial peptide libraries. Immunome research, 4(1), p.2.
63. Nielsen, M., Lundegaard, C. and Lund, O., 2007. Prediction of MHC class II
binding affinity using SMM-align, a novel stabilization matrix alignment

1778    method. BMC bioinformatics, 8(1), p.238.

1779 64. Sturniolo, T., Bono, E., Ding, J., Raddrizzani, L., Tuereci, O., Sahin, U.,
1780    Braxenthaler, M., Gallazzi, F., Protti, M.P., Sinigaglia, F. and Hammer, J.,
1781    1999. Generation of tissue-specific and promiscuous HLA ligand
1782    databases using DNA microarrays and virtual HLA class II matrices.
1783    Nature biotechnology, 17(6), p.555.

1784 65. Bui H. H, Sidney J, Dinh K, Southwood S, Newman M. J, Sette A. 2006.
1785    Predicting population coverage of T-cell epitope-based diagnostics and
1786    vaccines. BMC Bioinformatics 17:153.

1787 66. Larsen JE, Lund O, Nielsen M. 2006. Improved method for predicting linear
1788    B-cell epitopes. Immunome Res 2:2.

1789 67. Chou PY, Fasman GD. 1978. Prediction of the secondary structure of
1790    proteins from their amino acid sequence. Adv Enzymol Relat Areas Mol
1791    Biol 47:45-148.

1792 68. Emini EA, Hughes JV, Perlow DS, Boger J. 1985. Induction of hepatitis A
1793    virus-neutralizing antibody by a virus-specific synthetic peptide. J Virol
1794    55:836-839.

1795 69. Karplus PA, Schulz GE. 1985. Prediction of chain flexibility in proteins.
1796    Naturwissenschaften 72:212-213.

1797 70. Kolaskar AS, Tongaonkar PC. 1990. A semi-empirical method for prediction
1798    of antigenic determinants on protein antigens. FEBS Lett276:172-174.

1799 71. Parker JM, Guo D, Hodges RS. 1986. New hydrophilicity scale derived from
1800    high-performance liquid chromatography peptide retention data:
1801    correlation of predicted surface residues with antigenicity and X-ray-
1802    derived accessible sites. Biochemistry 25:5425-5432.

1803 72. J. V. Kringelum, C. Lundegaard, O. Lund, M. Nielsen. 2012. Reliable B cell
1804    epitope predictions: impacts of method development and improved
1805    benchmarking. PLoS Comput Biol. 8:(12):e1002829.

1806 73. Ponomarenko JV, Bui H, Li W, Fusseder N, Bourne PE, Sette A, Peters B.
1807    2008. ElliPro: a new structure-based tool for the prediction of antibody
1808    epitopes. BMC Bioinformatics 9:514.

1809 74. Bui HH, Sidney J, Li W, Fusseder N, Sette A. 2007. Development of an
1810    epitope conservancy analysis tool to facilitate the design of epitope-based
1811    diagnostics and vaccines. BMC Bioinformatics 8:361.

1812 75. Gupta, S., Kapoor, P., Chaudhary, K., Gautam, A., Kumar, R., Raghava,
1813    G.P. and Open Source Drug Discovery Consortium, 2013. In silico
1814    approach for predicting toxicity of peptides and proteins. PLoS One, 8(9),
1815    p.e73957.

1816 76. Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R.,
1817    McWilliam, H., Remmert, M., Söding, J. and Thompson, J.D., and Higgins
1818    D.G., 2011. Fast, scalable generation of high-quality protein multiple
1819    sequence alignments using Clustal Omega. Molecular systems biology,
1820    7(1), p.539.

1821 77. Benkert, P., Tosatto, S.C. and Schomburg, D., 2008. QMEAN: A
1822    comprehensive scoring function for model quality assessment. Proteins:
1823    Structure, Function, and Bioinformatics, 71(1), pp.261-277.

  60

78. Singh, S., Singh, H., Tuknait, A., Chaudhary, K., Singh, B., Kumaran, S. and
Raghava, G.P.S. (2015) PEPstrMOD: structure prediction of peptides
containing natural, non-natural and modified residues. Biology Direct
10:73.
79. Morris, G.M., Huey, R., Lindstrom, W., Sanner, M.F., Belew, R.K., Goodsell,
D.S. and Olson, A.J., 2009. AutoDock4 and AutoDockTools4: Automated
docking with selective receptor flexibility. Journal of computational
chemistry, 30(16), pp.2785-2791.
80. O. Trott, A. J. Olson, AutoDock Vina: improving the speed and accuracy of
docking with a new scoring function, efficient optimization and
multithreading, Journal of Computational Chemistry 31 (2010) 455-461
81. Abraham, M. J., Murtola, T., Schulz, R., Pa′ll, S., Smith, J. C., Hess, B.,
Lindahl, E. GROMACS: High performance molecular simulations through
multi-level parallelism from lap- tops to supercomputers. SoftwareX 1–
2:19–25, 2015.
82. Jorgensen, W.L., Maxwell, D.S. and Tirado-Rives, J., 1996. Development
and testing of the OPLS all-atom force field on conformational energetics
and properties of organic liquids. Journal of the American Chemical
Society, 118(45), pp.11225-11236.
83. Abele, R. and Tampé, R., 2004. The ABCs of immunology: structure and
function of TAP, the transporter associated with antigen processing.
Physiology, 19(4), pp.216-224.
84. Hu W, Li F, Yang X, Li Z, Xia H, Li G, Wang Y, Zhang Z. A flexible peptide
linker enhances the immunoreactivity of two copies HBsAg preS1 (21-47)
fusion protein. J Biotechnol. 2004;107:83–90.
85. Hajighahramani, N., Nezafat, N., Eslami, M., Negahdaripour, M.,
Rahmatabadi, S.S. and Ghasemi, Y., 2017. Immunoinformatics analysis
and in silico designing of a novel multi-epitope peptide vaccine against
Staphylococcus aureus. Infection, Genetics and Evolution, 48, pp.83-94.
86. Chen, X., Zaro, J.L. and Shen, W.C., 2013. Fusion protein linkers: property,
design and functionality. Advanced drug delivery reviews, 65(10),
pp.1357-1369.
87. Hoover, D.M., Rajashankar, K.R., Blumenthal, R., Puri, A., Oppenheim, J.J.,
Chertov, O. and Lubkowski, J., 2000. The structure of human β-defensin-2
shows evidence of higher order oligomerization. Journal of Biological
Chemistry, 275(42), pp.32911-32918. PDB ID: 1FD3.
88. Nagpal, G., Gupta, S., Chaudhary, K., Dhanda, S.K., Prakash, S. and
Raghava, G.P., 2015. VaccineDA: Prediction, design and genome-wide
screening of oligodeoxynucleotide-based vaccine adjuvants. Scientific
reports, 5, p.12478.
89. Dhanda, S. K., Vir, P. & Raghava, G. P. Designing of interferon-gamma
inducing MHC class-II binders. Biol. Direct. 8, 30 (2013).
90. Saha, S. & Raghava, G. AlgPred: prediction of allergenic proteins and
mapping of IgE epitopes. Nucleic. Acids. Res. 34, W202–W209 (2006).
91. Irini A Doytchinova and Darren R Flower. VaxiJen: a server for prediction of
protective antigens, tumour antigens and subunit vaccines. BMC

Bioinformatics. 2007 8:4.

92. Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S.E., Wilkins, M.R., Appel, R.D. and Bairoch, A., 2005. Protein identification and analysis tools on the ExPASy server (pp. 571-607). Humana Press.

93. Morten Källberg, Haipeng Wang, Sheng Wang, Jian Peng, Zhiyong Wang, Hui Lu, and Jinbo Xu. Template-based protein structure modeling using the RaptorX web server. Nature Protocols 7, 1511-1522, 2012.

94. Ma, J., Wang, S., Zhao, F. and Xu, J., 2013. Protein threading using context-specific alignment potential. Bioinformatics, 29(13), pp.i257-i265.

95. Wang, Z., Zhao, F., Peng, J. and Xu, J., 2010, December. Protein 8-class secondary structure prediction using conditional neural fields. In 2010 IEEE International Conference on Bioinformatics and Biomedicine (BIBM) (pp. 109-114). IEEE.

96. Wang, Z. and Xu, J., 2013. Predicting protein contact map using evolutionary and physical constraints by integer programming. Bioinformatics, 29(13), pp.i266-i273.

97. Dong Xu and Yang Zhang. Improving the Physical Realism and Structural Accuracy of Protein Models by a Two-step Atomic-level Energy Minimization. Biophysical Journal, vol 101, 2525-2534 (2011).

98. J. Ko, H. Park, L. Heo, and C. Seok, GalaxyWEB server for protein structure prediction and refinement, Nucleic Acids Res. 40 (W1), W294-W297 (2012).

99. Shin, W.H., Lee, G.R., Heo, L., Lee, H. and Seok, C., 2014. Prediction of protein structure and interaction by GALAXY protein modeling programs. Bio Design, 2(1), pp.1-11.

100. Ramakrishnan, C. and Ramachandran, G.N., 1965. Stereochemical criteria for polypeptide and protein chain conformations: II. Allowed conformations for a pair of peptide units. Biophysical journal, 5(6), pp.909-933.

101. S.C. Lovell, I.W. Davis, W.B. Arendall III, P.I.W. de Bakker, J.M. Word, M.G. Prisant, J.S. Richardson and D.C. Richardson (2002) Structure validation by Calpha geometry: phi,psi and Cbeta deviation. Proteins: Structure, Function & Genetics. 50: 437-450.

102. Bell, J.K., Botos, I., Hall, P.R., Askins, J., Shiloach, J., Segal, D.M. and Davies, D.R., 2005. The molecular structure of the Toll-like receptor 3 ligand-binding domain. Proceedings of the National Academy of Sciences of the United States of America, 102(31), pp.10976-10980.

103. Duhovny D, Nussinov R, Wolfson HJ. Efficient Unbound Docking of Rigid Molecules. In Gusfield et al., Ed. Proceedings of the 2'nd Workshop on Algorithms in Bioinformatics(WABI) Rome, Italy, Lecture Notes in Computer Science 2452, pp. 185-200, Springer Verlag, 2002

104. Schneidman-Duhovny D, Inbar Y, Nussinov R, Wolfson HJ. PatchDock and SymmDock: servers for rigid and symmetric docking. Nucl. Acids. Res. 33: W363-367, 2005.

105. Nezafat, N., Eslami, M., Negahdaripour, M., Rahbar, M.R. and Ghasemi, Y., 2017. Designing an efficient multi-epitope oral vaccine against Helicobacter pylori using immunoinformatics and structural vaccinology

1916     approaches. Molecular BioSystems, 13(4), pp.699-713.
1917 106. Morla, S., Makhija, A. & Kumar, S. Synonymous codon usage pattern in
1918     glycoprotein gene of rabies virus. Gene. 584, 1–6 (2016).
1919 107. Wu, X., Wu, S., Li, D., Zhang, J., Hou, L., Ma, J., Liu, W., Ren, D., Zhu, Y.
1920     and He, F., 2010. Computational identification of rare codons of
1921     Escherichia coli based on codon pairs preference. Bmc Bioinformatics,
1922     11(1), p.61.
1923
1924

**A. CTL epitopes**     **B. HTL epitopes**

**ß Defensin 2:** GIGDPVTCLKSGAICHPVFCPRRYKQIGTCGLPGTKCCKKP

| B. HTL epitopes | linker | A. CTL epitopes |
|---|---|---|
| | EAAAK | EAAAK |
| **C Protein** | | |
| VQMTYNWTQWLQTLY | GGGGS | MMASILLTLF |
| **Fusion Protein** | | |
| DLALSKYLSDLLFVF | GGGGS | AQITAGVALY |
| LALSKYLSDLLFVFG | GGGGS | FALSNGVLF |
| ALSKYLSDLLFVFGP | GGGGS | KYLSDLLFVF |
| LSKYLSDLLFVFGPN | GGGGS | MTIQAISQAF |
| SKYLSDLLFVFGPNL | GGGGS | AENPVFTVF |
| KYLSDLLFVFGPNLQ | GGGGS | AVYNNEFYY |
| YLSDLLFVFGPNLQD | GGGGS | LAMDEGYFAY |
| **Glycoprotein** | | |
| ASFSWDTMIKFGDVL | GGGGS | TVYHCSAVY |
| FSWDTMIKFGDVLTV | GGGGS | NYMYLICYGF |
| GVYNDAFLIDRINWI | GGGGS | YMIPRTMLEF |
| NDAFLIDRINWISAG | GGGGS | EIISDIGNY |
| DAFLIDRINWISAGV | GGGGS | TPFVDSRAY |
| AFLIDRINWISAGVF | GGGGS | YPALALNEF |
| FLIDRINWISAGVFL | GGGGS | LDPVVTDVVY* |
| **Matrix Protein** | | |
| IPREFMIYDDVFIDN | GGGGS | LVSDAKMLSY* |
| FMIYDDVFIDNTGRI | GGGGS | MPSDDFSNTF |
| **Nucleocapsid** | | |
| LSSDQVAELAAAVQE | GGGGS | VSDAKMLSY* |
| SSDQVAELAAAVQET | GGGGS | AEFFSFFRTF |
| SDQVAELAAAVQETS | GGGGS | ETDDYNGIY |
| DQVAELAAAVQETSA | GGGGS | FPISRLFNMY |
| QVAELAAAVQETSAG | GGGGS | FPVMGNRIY |
| **Phosphoprotein** | | |
| *NNGNVCLVSDAKMLS | GGGGS | IATVYTWAY |
| *NGNVCLVSDAKMLSY | GGGGS | IMKKSFKAY |
| *GNVCLVSDAKMLSYA | GGGGS | IPFLFLSAY |
| *NVCLVSDAKMLSYAP | GGGGS | KWYECFLFWF |
| *VCLVSDAKMLSYAPE | GGGGS | KYYQIDQPFF |
| NIDNIHLLAEFFSFF | GGGGS | LETDDYNGIY |
| IDNIHLLAEFFSFFR | GGGGS | RLFNMYRSY |
| **Polymerase** | | |
| DNIHLLAEFFSFFRT | GGGGS | SQNLLVTSY |
| NIHLLAEFFSFFRTF | GGGGS | SYFGLVLVCF |
| IHLLAEFFSFFRTFG | GGGGS | TSDLDFVIFY |
| LELASFLMDRRVILP | GGGGS | YPECNNILF |
| ELASFLMDRRVILPR | GGGGS | |
| LASFLMDRRVILPRA | GGGGS | |
| ASFLMDRRVILPRAA | GGGGS | |
| LDFVIFYASLTYLRR | GGGGS | |
| FVIFYASLTYLRRGI | EAAAK | EAAAK |

**ß Defensin 3:** GIINTLQKYYCRVRGGRCAVLSCLPKEEQIGKCSTRGRKCCRRKK

HHHHHH     HHHHHH

| Overlapping regions of predicted B cell Bepipred linear Epitope and epitopes predicted by other protein sequence and structure based methods | Overlapping regions of sequence based B cell linear epitope prediction | | | | | Overlapping regions of structure based B cell epitope prediction | | |
|---|---|---|---|---|---|---|---|---|
| | Chou & Fasman Beta-Turn | Emini Surface Accessibility | Karplus & Schulz Flexibility | Kolaskar & Tongaonkar Antigenicity | Parker Hydrophilicity | DiscoTope: discontinuous epitopes | ElliPro Linear | ElliPro Discontinuous |
| Fusion Protein — 215 GPNLQDPVSNSM 226 | 215-226 | | 215-226 | 215-216 | 217-226 | | | |
| Glycoprotein — 271 WTPPNPNT 278 | 271-278 | 271-278 | 272-278 | | 272-278 | | | 271-278 |
| Glycoprotein — 459 SWDTMI 464 | 459-460 | | | | | | | 463-464 |
| Glycoprotein — 529 NQTAE 533 | 529-532 | 529-533 | 529-533 | | 529-533 | | | |
| Matrix Protein — 184 SGIYM 188 | 184-188 | | 184-188 | | 184-188 | | | |
| Matrix Protein — 330 SIPREFMIY 338 | 330-335 | | 330-336 | | 331-334 | 332-333 | 332-333 | 333-338 |
| Matrix Protein — 340 DVFIDNTGRI 349 | 341-349 | | 340-349 | | 340-344 | | | 340-349 |
| Phosphoprotein — 125 GYGFTSSPERGWSDYTSGA 143 | 125-143 | 130-136 | 128-143 | | 127-143 | | | |
| Phosphoprotein — 163 IAVSKEDR 170 | 166-170 | 166-170 | 163-170 | 163-166 | 165-170 | | | |
| Polymerase — 346 NIDN 349 | 348-349 | | | | | | | |
| Polymerase — 364 GHPILE 369 | 364-367 | | | 365-369 | | | | 366-369 |
| Polymerase — 1095 DKSFDEDLEL 1104 | 1096-1103 | 1096-1101 | 1095-1104 | | 1096-1103 | | | 1095-1104 |
| Polymerase — 1108 LMDR 1111 | | | | | | | | 1108-1111 |
| Polymerase — 1409 LRLETDDYNG 1418 | 1412-1418 | 1410-1416 | 1412-1418 | | 1411-1418 | 1409-1417 | 1409-1417 | |
| Polymerase — 1982 GFPIS 1986 | 1982-1986 | | 1982-1986 | 1982-1986 | | | | 1182-1986 |
| Polymerase — 2004 PVYSNPD 2010 | 2004-2010 | 2006-2010 | 2007-2010 | 2004-2008 | 2007-2010 | | | 2004-2010 |

# Clusters of overlapping CTL, HTL & B cell epitopes

| | |
|---|---|
| **Fusion Protein** | `------------------------MTIQAISQAF-`<br>`----KYLSDLLFVF-----------------`<br>`------YLSDLLFVFGPNLQD---------`<br>`----KYLSDLLFVFGPNLQ----------`<br>`-----SKYLSDLLFVFGPNL---------`<br>`----LSKYLSDLLFVFGPN----------`<br>`---ALSKYLSDLLFVFGP-----------`<br>`--LALSKYLSDLLFVFG------------`<br>`-DLALSKYLSDLLFVF-------------`<br>`----------------GPNLQDPVSNSM----------`<br>199-236 LDLALSKYLSDLLFVFGPNLQDPVSNSMTIQAISQAFG |
| **Glycoprotein** | `--------------------------------------AENPVFTVF-`<br>`----------------FLIDRINWISAGVFL---------------`<br>`-------------AFLIDRINWISAGVF----------------`<br>`-------------DAFLIDRINWISAGV----------------`<br>`--------TVYHCSAVY-------`  `----FSWDTMIKFGDVLTV--`  `------NDAFLIDRINWISAG---------------`<br>`-------------AVYNNEFYY-`  `--ASFSWDTMIKFGDVL----`  `-GVYNDAFLIDRINWI------------------`<br>`-WTPPPNPNT---------------`  `-----SWDTMI------`  `-------------------------NQTAE-------------`<br>270-293 VWTPPNPNTVYHCSAVYNNEFYYV   454-474 YQASFSWDTMIKFGDVLTVNP   505-541 EGVYNDAFLIDRINWISAGVFLDSNQTAENPVFTVFK |
| **Matrix Protein** | `-----FMIYDDVFIDNTGRI-`<br>`-IPREFMIYDDVFIDN-----`<br>`----YMIPRTMLEF-`  `-----------DVFIDNTGRI-`<br>`-SGIYM---------`  `SIPREFMIY------------`<br>183-197 DSGIYMIPRTMLEFR   330-350 SIPREFMIYDDVFIDNTGRIL |
| **Nucleocapsid** | `------QVAELAAAVQETSAG--`<br>`-----DQVAELAAAVQETSA-`<br>`----SDQVAELAAAVQETS--`<br>`---SSDQVAELAAAVQET---`<br>`--LSSDQVAELAAAVQE------`<br>382-404 LGLSSDQVAELAAAVQETSAGRQ |
| **Phosphoprotein** | `------------LVSDAKMLSY-----------`<br>`-------------VSDAKMLSY-----------`<br>`-------VCLVSDAKMLSYAPE----------`<br>`--------NVCLVSDAKMLSYAP--------`<br>`-------GNVCLVSDAKMLSYA--------`<br>`------NGNVCLVSDAKMLSY---------`<br>`-----NNGNVCLVSDAKMLS----------`<br>`--GYGFTSSPERGWSDYTSGA---------`  `-----IAVSKEDR--`<br>123-172 CTGYGFTSSPERGWSDYTSGANNGNVCLVSDAKMLSYAPEIAVSKEDRET |
| **Polymerase** | `---------AEFFSFFRTF-------`<br>`-----IHLLAEFFSFFRTFG------`<br>`----NIHLLAEFFSFFRT--------`  `------ASFLMDRRVILPRAA-----`<br>`---DNIHLLAEFFSFFRT--------`  `-------LASFLMDRRVILPRA-----`<br>`--IDNIHLLAEFFSFFR--------`  `------ELASFLMDRRVILPR------`<br>`-NIDNIHLLAEFFSFF---------`  `-----LELASFLMDRRVILP------`  `----ETDDYNGIY--`<br>`-NIDN-----------------`  `------------LMDR----------`  `----LETDDYNGIY--`<br>`--------------GHPILE-`  `--DKSFDEDLEL---------------`  `-LRLETDDYNG----`<br>345-370 LNIDNIHLLAEFFSFFRTFGHPILEA   1093-1124 FHDKSFDEDLELASFLMDRRVILPRAAHEILD   1408-1422 RLRLETDDYNGIYHL |
| **Polymerase** | `------------SYFGLVLVCF----------`<br>`-----------RLFNMYRSY-----------`<br>`-TSDLDFVIFY-----------`  `--FPISRLFNMY------------------`<br>`----VIFYASLTYLRRGI-`  `-------------------PVYSNPD--`<br>`----LDFVIFYASLTYLRR---`  `-GFPIS----------------------`<br>1715-1736 NTSDLDFVIFYASLTYLRRGII   1981-2012 PGFPISRLFNMYRSYFGLVLVCFPVYSNPDST |

**A. CTL epitopes - HLA Class I allele complexes**

1. AEFFSFFRTF-B4403
2. AENPVFTVF-B4402
3. AENPVFTVF-B4403
4. AVYNNEFYY-A1101
5. AVYNNEFYY-A3002
6. KYLSDLLFVF-A2301
7. KYLSDLLFVF-A2402
8. LDPVVTDVVY-B3501
9. RLFNMYRSY-A3002
10. RLFNMYRSY-A3201
11. RLFNMYRSY-B1501
12. VSDAKMLSY-A0101

5. GNVCLVSDAKMLSYA-DRB1-0301
6. IDNIHLLAEFFSFFR-DPA1-0103
7. IDNIHLLAEFFSFFR-DPB1-0201
8. IHLLAEFFSFFRTFG-DPA1-0103
9. IHLLAEFFSFFRTFG-DPB1-0201
10. KYLSDLLFVFGPNLQ-DRB3-0101
11. LALSKYLSDLLFVFG-DRB3-0101
12. LSKYLSDLLFVFGPN-DRB3-0101
13. NGNVCLVSDAKMLSY-DRB1-0301
14. NIDNIHLLAEFFSFF-DPA1-0103
15. NIDNIHLLAEFFSFF-DPB1-0201
16. NIHLLAEFFSFFRTF-DPA1-0103
17. NIHLLAEFFSFFRTF-DPB1-0201
18. NNGNVCLVSDAKMLS-DRB1-0301
19. NVCLVSDAKMLSYAP-DRB1-0301
20. SKYLSDLLFVFGPNL-DRB3-0101

**B. HTL epitopes - HLA Class II allele complexes**

1. ALSKYLSDLLFVFGP-DRB3-0101
2. DLALSKYLSDLLFVF-DRB3-0101
3. DNIHLLAEFFSFFRT-DPA1-0103
4. DNIHLLAEFFSFFRT-DPB1-0201
21. VCLVSDAKMLSYAPE-DRB1-0301
22. YLSDLLFVFGPNLQD-DRB3-0101

**A**

1. AEFFSFFRTF-B4403
2. AENPVFTVF-B4402
3. AENPVFTVF-B4403
4. AVYNNEFYY-A1101
5. AVYNNEFYY-A3002
6. KYLSDLLFVF-A2301
7. KYLSDLLFVF-A2402
8. LDPVVTDVVY-B3501
9. RLFNMYRSY-A3002
10. RLFNMYRSY-A3201
11. RLFNMYRSY-B1501
12. VSDAKMLSY-A0101

RMSD (nm)

TIME (ns)

**B**

1. ALSKYLSDLLFVFGP-DRB3-0101
2. DLALSKYLSDLLFVF-DRB3-0101
3. DNIHLLAEFFSFFRT-DPA1-0103
4. DNIHLLAEFFSFFRT-DPB1-0201
5. GNVCLVSDAKMLSYA-DRB1-0301
6. IDNIHLLAEFFSFFR-DPA1-0103
7. IDNIHLLAEFFSFFR-DPB1-0201
8. IHLLAEFFSFFRTFG-DPA1-0103
9. IHLLAEFFSFFRTFG-DPB1-0201
10. KYLSDLLFVFGPNLQ-DRB3-0101
11. LALSKYLSDLLFVFG-DRB3-0101
12. LSKYLSDLLFVFGPN-DRB3-0101
13. NGNVCLVSDAKMLSY-DRB1-0301
14. NIDNIHLLAEFFSFF-DPA1-0103
15. NIDNIHLLAEFFSFF-DPB1-0201
16. NIHLLAEFFSFFRTF-DPA1-0103
17. NIHLLAEFFSFFRTF-DPB1-0201
18. NNGNVCLVSDAKMLS-DRB1-0301
19. NVCLVSDAKMLSYAP-DRB1-0301
20. SKYLSDLLFVFGPNL-DRB3-0101
21. VCLVSDAKMLSYAPE-DRB1-0301
22. YLSDLLFVFGPNLQD-DRB3-0101

RMSD (nm)

TIME (ns)

1. Epitope AEFFSFFRTF through TAP cavity

2. Epitope AENPVFTVF through TAP cavity

3. Epitope AVYNNEFYY through TAP cavity

4. Epitope KYLSDLLFVF through TAP cavity

5. Epitope LDPVVTDVVY through TAP cavity

6. Epitope RLFNMYRSY through TAP cavity

7. Epitope VSDAKMLSY through TAP cavity

# CTL multi-epitope vaccine:

# HTL multi-epitope vaccine:

General/Pre-Pro/Proline Favoured    General/Pre-Pro/Proline Allowed
Glycine Favoured    Glycine Allowed

**Nine NIPAH proteins were chosen**
*Protein sequences and available 3D structures
retrieved from NCBI & RCSB PDB or modelled by SwissModel*

**Screening of potential CTL, HTL and B Cell epitopes**

Screening of CTL epitope
*(IEDB "Proteasomal cleavage/
TAP transport/MHC class I combined predictor")*

Screening of HTL epitopes
*(IEDB "MHC-II Binding Predictions")*

Screening of protein sequence &
structure based B Cell epitope *(IEDB "B Cell
Epitope Prediction Tools", DiscoTope2.0 & Ellipro)*

Population coverage by screened
CTL and HTL epitopes *(IEDB "Population Coverage")*

**Characterization of screened potential epitopes**

Epitope conservation analysis
*(IEDB "Epitope Conservancy Analysis")*

Epitope toxicity prediction
*(ToxinPred)*

Overlapping residue analysis for
CTL, HTL & B Cell epitopes *(Clustal Omega)*

**Molecular interaction study of selected
overlapping CTL & HTL epitopes with HLA allele and TAP transporter**

3D structure modelling of HLA alleles
and selected CTL & HTL epitopes *(Swiss-model & PEPstrMOD)*

Molecular interaction analysis of
selected overlapping epitopes with HLA allele
*(AutoDock Tool 4.2, AutoDock Vina, & Gromacs 5.1.4)*

Molecular interaction analysis of selected
CTL epitopes with TAP transporter *(AutoDock Vina)*

**Design and characterisation of CTL & HTL Multi-Epitope Vaccines**

Interferon gama inducing epitope
prediction from designed MEVs *("IFNepitope")*

MEVs allergenicity and
antigenicity prediction *(AlgPred & VaxiJen)*

Physicochemical
analysis of MEVs *(ProtParam)*

**3D structure modelling and refinement
of CTL & HTL MEVs *(RaptorX, GalaxyWEB)***

*In-silico* validation of refined CTL
& HTL MEV models *(RAMPAGE)*

Discontinuous B-cell epitope prediction
from CTL & HTL MEVs *(IEDB "ElliPro")*

**Molecular interaction and complex stability study of
CTL & HTL MEVs with TLR2 & TLR4**
*(PatchDock & Gromacs 5.1.4)*

***In-silico* analysis of cDNA of CTL & HTL MEVs for
cloning and expression in mammalian cell line (Human)**
*(Codon Usage Wrangler Tool & GenScript Rare Codon Analysis Tool)*

**A. B-factor of CTL epitope & HLA I allele complexes**

1. AEFFSFFRTF-B4403
2. AENPVFTVF-B4402
3. AENPVFTVF-B4403
4. AVYNNEFYY-A1101
5. AVYNNEFYY-A3002
6. KYLSDLLFVF-A2301
7. KYLSDLLFVF-A2402
8. LDPVVTDVVY-B3501
9. RLFNMYRSY-A3002
10. RLFNMYRSY-A3201
11. RLFNMYRSY-B1501
12. VSDAKMLSY-A0101

**B. B-factor of HTL epitope & HLA II allele complexes**

1. ALSKYLSDLLFVFGP-DRB3-0101
2. DLALSKYLSDLLFVF-DRB3-0101
3. DNIHLLAEFFSFFRT-DPA1-0103
4. DNIHLLAEFFSFFRT-DPB1-0201

5. GNVCLVSDAKMLSYA-DRB1-0301
6. IDNIHLLAEFFSFFR-DPA1-0103
7. IDNIHLLAEFFSFFR-DPB1-0201
8. IHLLAEFFSFFRTFG-DPA1-0103
9. IHLLAEFFSFFRTFG-DPB1-0201
10. KYLSDLLFVFGPNLQ-DRB3-0101
11. LALSKYLSDLLFVFG-DRB3-0101
12. LSKYLSDLLFVFGPN-DRB3-0101
13. NGNVCLVSDAKMLSY-DRB1-0301
14. NIDNIHLLAEFFSFF-DPA1-0103
15. NIDNIHLLAEFFSFF-DPB1-0201
16. NIHLLAEFFSFFRTF-DPA1-0103
17. NIHLLAEFFSFFRTF-DPB1-0201
18. NNGNVCLVSDAKMLS-DRB1-0301
19. NVCLVSDAKMLSYAP-DRB1-0301
20. SKYLSDLLFVFGPNL-DRB3-0101
21. VCLVSDAKMLSYAPE-DRB1-0301
22. YLSDLLFVFGPNLQD-DRB3-0101